

A NEW CLASS OF OPTICAL SOLITONS

Sasanka Ghosh

Physics Department, Indian Institute of Technology, Guwahati

Panbazar, Guwahati - 781 001, India

Phone: + 91-361-521-915, Fax: + 91-361-521-916

E-mail: sasanka@iitg.ernet.in

and

Sudipta Nandy

Physics Department, Indian Institute of Technology, Guwahati

Panbazar, Guwahati - 781 001, India

Phone: + 91-361-521-915, Fax: + 91-361-521-916

E-mail: sudipta@iitg.ernet.in

ABSTRACT

Existence of a new class of soliton solutions is shown for higher order nonlinear Schrodinger equation, describing third order dispersion, Kerr effect and stimulated Raman scattering. These new solutions have been obtained by invoking a group of nonlinear transformations acting on localised stable solutions. Stability of the solutions has been studied for different values of the arbitrary coefficient, involved in the recursion relation and consequently, different values of coefficient lead to different transmission rate for almost same input power. Another series solution containing even powers of localised stable solution is shown to exist for higher order nonlinear schrodinger equation.

Key words : Nonlinear dynamics, Optical solitons, Raman scattering, Kerr effect, Group of nonlinear transformation.

Theoretical prediction of Hasegawa and Tappert that an optical pulse in a dielectric fibre forms an envelope soliton [1] and subsequent experimental verification by Mollenauer [2] have made a significant achievement in ultra-high speed telecommunication. Contrary to conventional optical pulses, solitonic optical pulses are dispersionless, which makes them so important for long distance communication. Optical solitons are currently under intense study not only for providing distortionless signal transmission, but also for many nonlinear optical applications in fibre.

Nonlinear Schrodinger equation (NLS) and their various higher order gener-

alizations play a pivotal role in describing the dynamics of optical solitons. The successful propagation of NLS solitons for long distance transmission eventually turns out to be limited to a few kilometers [2]. This is due to the fact that solitons, which follow the dynamics of NLS cannot take care of propagation loss, as it propagates along the optical fibre and consequently the pulse needs to be reshaped time to time to compensate propagation loss. As a remedy, at present, a Raman pump light is used to send a signal simultaneously along the fibre to adjust the propagation loss [3, 4]. This is achieved by utilizing the effect of stimulated Raman scattering. It is seen later [5, 6, 7] that the effect of stimulated Raman scattering already exists within the spectra of soliton solutions, if we include a few more terms in the equation of motion over NLS. No additional pump light is needed in the later case to adjust the propagation loss. The equation of motion, thus obtained, is called higher order nonlinear Schrodinger equation (HNLS):

$$\partial_z E = i[\beta_1 \partial_t E + \beta_2 \partial_t^2 E] + [\beta_3 \partial_t^3 E + \beta_4 \partial_t (\partial_t^2 E) + \beta_5 \partial_t (\partial_t \partial_t^2 E)] \quad (1)$$

where, E is envelop of electric field propagating in z direction at a time t . Notice that in absence of last three terms, (1) reduces to NLS. The coefficients of the higher order terms, namely β_3 , β_4 and β_5 respectively represent third order dispersion (TOD), self-steepening (SS) related to Kerr effect and the self frequency shifting via stimulated Raman scattering. It is the last term, which plays an important role in the propagation of distortionless optical pulses over long distance. On the other hand, for ultra-short pulses, the effects of last three

terms cannot be neglected.

Several attempts are made to find the soliton solutions of HNLS (1) [8-12] by various methods like Hirota's bilinear method, Painleve analysis, Backlund transformation, inverse scattering technique. Recently, HNLS is solved for soliton solutions by inverse scattering technique method for an arbitrary value of the coefficients [13], where it is convenient to solve a gauge equivalent equation,

$$\partial_t q(x;t) = \partial_{xxx} q(x;t) + \alpha_1 q(x;t) \partial_x q(x;t) + \alpha_2 \partial_x (q(x;t) \partial_x q(x;t)): \quad (2)$$

The equations (1) and (2) are related explicitly through the following gauge transformation:

$$E(z; \lambda) = \exp[-i p x + i p^3 t] q(x;t) \quad (3a)$$

$$x = \lambda + \alpha_1 p z \quad (3b)$$

$$t = \alpha_3 z \quad (3c)$$

with $p = \alpha_1 = (3 - \alpha_3) = \alpha_2 = (\alpha_4)$, $\alpha_1 = \alpha_4 = \alpha_3$, and $\alpha_2 = \alpha_5 = \alpha_3$.

One soliton solution, found by inverse scattering method, may be given by [13]

$$E(z; \lambda) = \left(\frac{p}{\lambda} \right)^{\frac{1}{2}} (n - 1) \operatorname{sech} \tilde{A}(z; \lambda) \exp(i \tilde{B}(z; \lambda)) \quad (4)$$

with

$$\begin{aligned} \tilde{A}(z; \lambda) &= \lambda + \left[\alpha_3 \left(\lambda^3 - \alpha_3 \lambda^2 \right) + \frac{\alpha_1^2}{3 \alpha_3} \right] z - \frac{1}{2} \ln(n - 1) \\ \tilde{B}(z; \lambda) &= \left[-\frac{1}{3 \alpha_3} \right] + \left[\alpha_3 \left(\lambda^2 - \alpha_3 \lambda^2 - \frac{\alpha_1^2}{3 \alpha_3^2} \right) - \frac{2 \alpha_1^3}{27 \alpha_3^2} \right] z + \end{aligned}$$

where, n is dimension of the Lax operator and the location of the simple pole is chosen at $\lambda = \frac{1}{2}(\alpha + i\beta)$ in the complex spectral parameter plane, α and β being real parameters. Simple sech type structure of one soliton in (4) is quite interesting, since it may be obtained from the output of a mode locked laser. N soliton solutions is obtained for a more a general dynamical equation in [14].

In this paper, we discuss a new class of solution of (1). For this, let us assume a trial solution of (1) in the form

$$E(z; \lambda) = E_0 e^{i\lambda z + im} y(\lambda) \quad (5)$$

where λ and m are real parameters and $y(\lambda)$ is chosen as a real function. Substituting (5) in (1), we obtain a set of two differential equations in y :

$$(\lambda^2 - \lambda m + \alpha m^3)y(\lambda) + (\lambda - \alpha - \beta m)y''(\lambda) + (\lambda^2 - \alpha m)E_0 y^3(\lambda) = 0 \quad (6a)$$

$$(2 - \lambda m - \alpha - \beta m^2)y'(\lambda) + \beta y'''(\lambda) + \frac{1}{3}(\lambda^2 + 2\alpha - \beta)(y^3)'(\lambda) = 0 \quad (6b)$$

Consistency conditions of (6a) and (6b) demands that

$$\lambda^2 = (\lambda + \beta m)^2 \quad (7)$$

$$\lambda = \alpha - m(1 + m)^2 \quad (8)$$

$$\lambda^2 + 2\alpha - \beta = \frac{3E_0^2}{3} [\lambda^2 - m - \alpha] \quad (9)$$

Notice that (6) leads a solution of the form

$$y = \text{sech}(E_0 \lambda) \quad (10)$$

provided the following conditions are satisfied

$$m = \frac{1 - \alpha^2}{3 - \alpha^2} \quad (11)$$

$$\begin{aligned}
1 &= \frac{2(\alpha_1 - \alpha_3)(\alpha_1 + 2\alpha_3)^2}{3^2 \alpha_3^2} \\
E_0^2 &= \frac{(\alpha_1 - 3\alpha_3)(\alpha_1 + \alpha_3)}{3^2 \alpha_3^2} \\
2 &= 2\alpha_3 - \alpha_4 + \frac{1}{3} \frac{\alpha_4^2}{\alpha_3} \\
5 &= \frac{(3\alpha_3 - \alpha_1)(\alpha_1 + \alpha_3)}{2\alpha_3} - \frac{4}{2}
\end{aligned}$$

such type solution of (6) is a desired one, as it has been obtained earlier [13] by inverse scattering method (4). In this paper, however, we look for a more general solution of the form

$$y = \sum_{j=0}^{\infty} C_j y^j(x) \quad (12)$$

Substituting (12) into (6), we obtain

$$C_0 = 0; \quad \left(\frac{1 + m^2 \alpha_1 - m^3 \alpha_3}{(\alpha_2 - m^2 \alpha_1) E_0^2} \right)$$

Unfortunately, nontrivial values of C_0 lead only to trivial solutions of (6) consistent with the conditions (12). Thus $C_0 = 0$ is an acceptable solution and consequently the recursion relation among the coefficients C_j may be given by

$$\begin{aligned}
& [(\alpha_1 - m^2 \alpha_1 + m^3 \alpha_3) + \alpha_0^2 (\alpha_1 - 3m\alpha_3)(j+2)^2] C_{j+2} \\
& = \alpha_0^2 (\alpha_1 - 3m\alpha_3)(j^2 + j) C_j - \alpha_0^2 (\alpha_2 - m\alpha_4) \sum_{m=0}^{j-2} C_{j+2-m} C_{m-1} C_1
\end{aligned} \quad (13)$$

for $j = 0; 1; 2; 3; \dots$. If we now choose arbitrary, all the even coefficients become trivial

$$C_{2j} = 0 \quad (14)$$

whereas odd coefficients satisfy a recursion relation

$$C_{2j+1} = \frac{1}{2j(j+1)} [j(2j-1)C_{2j-1} - \sum_{m=0}^{2j-1} C_{2j+1-m} C_{m-1} C_1] \quad (15)$$

provided the coefficients satisfy the conditions (12), which is not surprising, as we will see $y = \text{sech}(E_0)$ itself is a special solution of (12). It is interesting to observe from the recursion relation (15) that if the arbitrary coefficient C_1 is chosen to be unity all higher order coefficients starting from C_3 become zero and the series leaves with one term, $y = \text{sech}(E_0)$. Thus the solution (10) already exists as a special case in (12) for nontrivial odd coefficients. Notice that a similar solution also exists for NLS [15].

Let us now introduce an operatorial notation for the series solution (12) as

$$H_{C_1} Y = \sum_{j=1}^{\infty} C_{2j+1} Y^{2j+1} \quad (16)$$

for arbitrary C_1 such that

$$H_1 Y = Y$$

i.e. the coefficients satisfy the recursion relations (14) and (15). It emerges from (16) and the recursion relations that

$$H_{C_1} H_{C_1^0} = H_{C_1 C_1^0} \quad (17)$$

implying that $\{H_{C_1}; C_1 \in \mathbb{R}\}$ is the commutative multiplicative group of nonlinear transformation (GNT). Thus, like NLS, the commutative group of nonlinear transformations for HNLS also leads to a new solution by operating on stationary localised solutions. The advantage of GNT method is that it enables us to

determine the stability of the new solution by the following relation

$$\lim_{\epsilon \rightarrow 0} H_{1+\epsilon} Y = Y \quad (18)$$

The main motivation in obtaining new type of solutions by GNT method is to increase the transmission rate. It is observed in NLS that higher values of the coefficient C_1 gives rise to higher transmission rate for almost the same input power [15]. Interestingly, the new form of solutions (14) and (15) also assert the same conclusion for HNLS with an added advantage that HNLS takes care of propagation loss, unlike NLS.

Let us present another interesting solution for HNLS, which emerges from the recursion relation (14). If we choose C_2 arbitrary and impose the condition that

$$2(\epsilon_1 m^2 + \epsilon_3 m^3) = \epsilon_0^2 (\epsilon_1 - 3m^3) \quad (19)$$

all odd coefficients will become trivial, i.e.

$$C_{2j+1} = 0 \quad (20a)$$

whereas even coefficients will satisfy a recursion relation, given by

$$C_{2j} = \frac{2}{(8j^2 + 1)} [2(j-1)(2j-1)C_{2j-2} - \frac{2}{1-3m^3} \sum_{m=0}^{j-1} X^{2j} X^m C_{2j-m} C_{m-1} C_1] \quad (20b)$$

To conclude, a new class of soliton solutions are obtained for HNLS. This is found by invoking a group of nonlinear transformations acting on a stable

localised solution. It turns out that the new solution reduces to the most stable one for $C_1 = 1$. For higher values of C_1 , however, it leads to a faster rate of propagation for an almost same input power. This, indeed, is a considerable improvement in the field of all optical communication through fibre. The existence of another solution of HNLS, where all odd coefficients are trivial, also earns a lot of interest and needs to be investigated more thoroughly. It is quite useful to see the consequences of the even series solution in the context of fibre communication.

S.N. would like to thank CSIR, Govt. of India for fellowship and financial support.

References

- [1] A. Hasegawa and F.D. Tappert, Appl. Phys. Lett. 23, 142 (1973).
- [2] L.F. Mollenauer, R.H. Stolen and J.P. Gordon, Phys. Rev. Lett 45, 1095 (1980).
- [3] A. Hasegawa, Appl. Opt. 23, 3302 (1984); Optical Solitons in Fibres, (Springer, Heidelberg, 1989).
- [4] L.F. Mollenauer, K. Smith, Opt. Lett. 13, 675 (1988).
- [5] J.P. Gordon, Opt. Lett. 11, 662 (1986).

- [6] F.M. Mitschke and L.F. Mollenauer, *Opt. Lett.* 11, 657 (1986).
- [7] Y. Kodama and A. Hasegawa, *IEEE J. Quantum Electron.* QE-23, 5610 (1987).
- [8] N. Sasa and J. Satsuma, *J. Phys. Soc. Jpn.* 60, 409 (1991).
- [9] K. Porsezian and K. Nakkeeran, *Phys. Rev. Lett.* 76, 3955 (1996).
- [10] M. Gedalin, T.C. Scott and Y.B. Band, *Phys. Rev. Lett.* 78, 448 (1997).
- [11] K. Porsezian, M. Daniel and M. Lakshmanan, *Proc. of the Int. Conf. on Nonlinear Evolution Equations and Dynamical Systems*, Eds. V.G. Makhankov, I. Puzynin and O. Pashaev 436 (World Sc., Singapore).
- [12] S. Ghosh, A. Kundu and S. Nandy, *Soliton Solutions, Liouville Integrability and Gauge Equivalence of Sasa Satsuma Equation*, *J. Math. Phys.* 40, 1993 (1999).
- [13] S. Ghosh and S. Nandy, *Optical Solitons in Higher Order Nonlinear Schrodinger Equation*, *solv-int* 9904019.
- [14] S. Ghosh and S. Nandy, *Inverse Scattering Method and Vector Higher Order Nonlinear Schrodinger Equation*, *solv-int* 9904021.
- [15] S. Lekic, S. Glamic and Z. Rajlic, *Optical Fibre Communication: Group of Nonlinear Transformations*, *solv-int* 9812012.