

Stability of wormholes in DGP gravity

Martín G. Richarte^{1, a}

¹ *Departamento de Física, Facultad de Ciencias Exactas y Naturales,
Universidad de Buenos Aires, Ciudad Universitaria, Pabellón I, 1428 Buenos Aires, Argentina*

Spherically symmetric thin-shell wormholes in DGP theory are constructed using the generalized junction conditions. We concentrate on the mechanical stability of wormholes under radial perturbation preserving the original spherical symmetry. In order to do that, we consider linearized perturbations around static solutions. We obtain that for certain values of the mass μ and crossover scale r_c stable wormholes exist with very small values of squared speed sound. Unlike the case of Einstein's gravity, this type of wormholes fulfil the energy conditions. Besides, we show that the gravitational field associated to these wormhole configurations is attractive for $\mu > 0$.

PACS numbers:

Keywords:

I. INTRODUCTION

In the last years a new type of gravitational model was widely studied in the context of cosmology as well as particle physics, the so called Dvali, Gabadadze and Porrati (DGP) theory. It predicts deviations from the standard 4D gravity over large distances. The transition between four and higher-dimensional gravitational potentials in the DGP model arises because of the presence of both the brane and bulk Hilbert–Einstein(H–E) terms in the action [1]. Cosmological considerations of the DGP model were first discussed in [2] where it was shown that in a Minkowski bulk spacetime we can obtain self-accelerating solutions.

Besides, a lot of efforts have been devoted to investigate wormhole configurations beyond Einstein framework because they seem to possess some interesting properties regarding the kind of matter which could support them [3],[4],[5]. To be more concrete, it was shown that static thin-shell wormholes could be supported by ordinary matter fulfilling the energy conditions in the Einstein–Gauss–Bonnet theory [3], in Brans–Dicke gravity [5], and in DGP gravity [6]. Interestingly, the former configurations turned out to be stable under radial perturbations[7].

In general, alternative gravity theories provide generalized junction conditions for the thin-shell wormholes which modify considerably the projected field equations on the joining surface, adding new types of geometrical objects besides the usual extrinsic curvature tensor. As we said, the contributions from the curvature tensor, theoretically, seem to allow the existence of wormholes supported by ordinary matter [3],[5],[6]. However, traversable wormhole geometries result of physical interest mainly as long as they are stable, at least under a simple kind of perturbation. Thus, beyond the mere characterization of static wormhole solutions, their stability under perturbations should always be explored.

Within the Einstein framework a straightforward approach for analyzing the stability of thin-shell wormhole solutions was proposed by Poisson and Visser in Ref. [8]. Hence, starting with a known static thin-shell wormhole solution of the Lanczos equations [9] the authors consider linearized spherically symmetric perturbation around the static solution. This choice leads to a single dynamical master equation which completely determines the motion of the wormhole throat[8]. Such a procedure has been subsequently followed to study the stability of more general spherically symmetric configurations (see, for example, Refs. [10]).

Another consequence of the non-linearity introduced by the DGP theory is related with the way in which the stability analysis is carried out for the dynamic case, that is, in this context it is not completely clear how to undertake the seek of the allowed stability zones.

Our goal is to perform a study of the linear stability of thin-shell wormholes within the DGP gravity preserving the original symmetry. We focus on configurations supported by ordinary matter which satisfy the energy conditions. Moreover, we shall show that there exist stable wormholes with squared speed sound within the range $0 \leq v_s^2 \leq 0.1$ indicating that the matter located at the throat of the wormhole could be non-relativistic.

II. THE DGP GRAVITY

We start from the action for the DGP theory in a five-dimensional manifold \mathcal{M}_5 with four-dimensional boundary Σ (cf. [11]),

$$S = 2M_5^3 \int_{\mathcal{M}_5} d^5x \sqrt{-g} R(g_{\mu\nu}) + \int_{\Sigma} d^4x \sqrt{-\gamma} 2M_4^2 \mathcal{R}(\gamma_{ab}) + \int_{\Sigma} d^4x \sqrt{-\gamma} \left(-4M_5^3 \mathcal{K}(\gamma_{ab}) + \mathcal{L}_m \right),$$

where $g_{\mu\nu}$ is the five-dimensional metric, γ_{ab} is the four-dimensional induced metric on the boundary Σ , and \mathcal{K} is the trace of extrinsic curvature. Here, the matter

^aElectronic address: martin@df.uba.ar

fields in \mathcal{L}_m are confined to a 4-dimensional boundary. The extra term in the boundary introduces a mass scale $m_c = 2M_5^3/M_4^2 = r_c^{-1}$ which determines a scale that separates two different regimes of the theory. For distances much smaller than m_c^{-1} one would expect the solutions to be well approximated by General Relativity and the modifications to appear at larger distances. This is indeed the case for distributions of matter and radiation which are homogeneous and isotropic at scales $\gtrsim r_c$. Typically, $m_c \sim 10.42\text{GeV}$, so it sets the distance/time scale $r_c = m_c^{-1}$ at which the Newtonian potential significantly deviates from the conventional one. In the bulk the DGP equations are the Einstein ones in vacuum : $G_{\mu\nu}^{(5)} = 0$. Then, the Birkhoff's theorem forces the bulk metric to be static, and of the Schwarzschild form:

$$ds^2 = -f(r)dt^2 + \frac{1}{f(r)}dr^2 + r^2d\Omega_3^2, \quad (1)$$

$$f(r) = 1 - \frac{\mu}{r^2} \quad (2)$$

where the parameter μ is related to the five dimensional Arnowitt–Deser–Misner (ADM) mass, $M_{ADM} = 3\pi^2\mu M_5^3$. The above spacetime has only one horizon placed at $r_+ = \sqrt{\mu}$ with $\mu > 0$. Besides, when $\mu < 0$ the manifold only presents a naked singularity at the origin $r = 0$.

In the next sections, without loss of generality, we shall take $8M_5^3 = 1$ so the crossover scale reads as $r_c = 4M_4^2$.

III. THIN-SHELL CONSTRUCTION IN DGP THEORY

Employing the metric Eqs.(1-2) we build a spherically symmetric thin-shell wormhole in DGP theory. We take two copies of the spacetime and remove from each manifold the five-dimensional regions described by

$$\mathcal{M}_\pm = \{x/r_\pm \leq a, a > r_h\}. \quad (3)$$

where a is chosen to include possible singularities or horizon within the region \mathcal{M}_\pm . The resulting manifolds have boundaries given by the timelike hypersurfaces

$$\Sigma_\pm = \{x/r_\pm = a, a > r_h\}. \quad (4)$$

Then we identify these two timelike hypersurfaces to obtain a geodesically complete new manifold $\mathcal{M} = \mathcal{M}^+ \cup \mathcal{M}^-$ with a matter shell at the surface $r = a$ where the throat of the wormhole is located. This manifold is constituted by two regions which are asymptotically flat. To study this type of wormholes we apply the Darmois-Israel formalism generalized [11],[12],[13] to the case of DGP theory. We can introduce the coordinates $\xi^a = (\tau, \chi, \theta, \phi)$ in Σ , with τ the proper time on the throat. Though we shall first focus in static configurations, in the subsequent analysis we could allow the radius of the throat be a function of the proper time for

studying the dynamics evolution of the wormholes, then in general we have that the boundary hypersurface reads:

$$\Sigma : \mathcal{H}(r, \tau) = r - a(\tau) = 0. \quad (5)$$

Because the Birkhoff theorem holds for the original manifold then the geometry remains static outside the throat, regardless the radius $a(\tau)$ can vary with time, so no gravitational waves are present.

The field equations projected on the shell Σ are the generalized junction (or Darmois–Israel) conditions [11, 12]

$$r_c \left(\mathcal{R}_{ab} - \frac{1}{2} \gamma_{ab} \mathcal{R} \right) - 2 \left(\langle \mathcal{K}_{ab} - \mathcal{K} \gamma_{ab} \rangle \right) = \mathcal{S}_{ab}, \quad (6)$$

where the bracket $\langle \cdot \rangle$ stands for the jump of a given quantity across the hypersurface Σ and γ_{ab} is the induced metric on Σ . The extrinsic curvature \mathcal{K}_{ab} is defined as follows:

$$\mathcal{K}_{ab}^\pm = -n_\kappa^\pm \left(\frac{\partial^2 X^\kappa}{\partial \xi^a \partial \xi^b} + \Gamma_{\mu\nu}^\kappa \frac{\partial X^\mu}{\partial \xi^a} \frac{\partial X^\nu}{\partial \xi^b} \right)_{r=a}, \quad (7)$$

where n_κ^\pm are the unit normals ($n_\kappa n^\kappa = 1$) to the surface Σ .

Notice that the first term in (6) is not enclosed within the brackets because this contribution comes from the four dimensional E-H term in the DGP action which already lives in the boundary, so it does not need to be projected on Σ . By taking the limit $r_c \rightarrow 0$ we recover the standard Darmois–Israel junction condition found in [9]. For a review of the junction conditions within the DGP theory see [13],[14].

After some algebraic manipulation, we obtain that the energy density ($\sigma = -\mathcal{S}_\tau^\tau$) and the pressure ($p = \mathcal{S}_i^i$) can be recast as

$$\sigma = 3r_c \left(\frac{\dot{a}^2}{a^2} + \frac{1}{a^2} \right) - \frac{6}{a} \ell, \quad (8)$$

$$p = -r_c \left(\frac{\dot{a}^2}{a^2} + \frac{1}{a^2} + \frac{2\ddot{a}}{a} \right) + \left(2\ddot{a} + f' \right) \ell^{-1} + \frac{4\ell}{a}. \quad (9)$$

where $\ell := \sqrt{f(a) + \dot{a}^2}$, the prime indicates the derivatives with respect to a while the dot stands for a derivative with respect to τ . The DGP contributions are encoded in the r_c factor of the above equations. If we take $r_c \rightarrow 0$ in both equations (8) and (9) we recover the expression for the energy density σ and the pressure p found in [3], ignoring the Gauss-Bonnet contribution.

It is easy to see from σ and p that the energy conservation equation is fulfilled:

$$\frac{d(a^3\sigma)}{d\tau} + p \frac{da^3}{d\tau} = 0, \quad (10)$$

The first term in Eq. (10) represents the internal energy change of the shell and the second the work by internal forces of the shell. The dynamical evolution of the wormhole throat is governed by the generalized Lanczos equations and to close the system we must supply an equation of state $p = p(\sigma)$ that relates p and σ .

IV. THE STABILITY ANALYSIS

In general to obtain the dynamic picture of the wormholes within the DGP gravity is a very complicated task. As it can be seen from the Eqs. (8-9) the nonlinear character of these expressions makes the idea of obtaining exact solutions very hard to implement. However, we can follow another route and study the stability of static solutions by linearizing the field equation. A physically interesting wormhole geometry should last enough so that its traversability makes sense. Thus the stability of a given wormhole configuration becomes a central aspect of its study. Here we shall analyze the stability under small perturbations preserving the spherical symmetry of the configuration; for this we shall proceed as [8], [15]. As we said, the dynamical evolution is determined by Eqs. (8) and (9), or by any of them and Eq. (10), and to complete the system we must add an equation of state that relates p with σ .

Our first move to address the stability issue is to recast Eq. (8) in such a way that it allows us to get $\dot{a} = \mathcal{F}(a, \sigma(a))$. Then, by squaring appropriately the energy density, we obtain a quadratic polynomial in the variable $X = a^{-2}(1 + \dot{a}^2)$ as it reads below:

$$r_c^2 X^2 - 2X(r_c \bar{\sigma} + 2) + \bar{\sigma}^2 - 4G(a) = 0 \quad (11)$$

with $\bar{\sigma} = \sigma/3$, and $G(a) = -\mu/a^4$. From the master equation (11) we get a single dynamical equation which completely determines the motion of the wormhole throat after the energy density is selected:

$$\dot{a}^2 = -V(a) \quad (12)$$

$$V(a) = 1 - \frac{a^2}{r_c} \left[\bar{\sigma} + \frac{2}{r_c} (1 + \epsilon Y(a)) \right] \quad (13)$$

$$Y(a) = \left(1 + r_c^2 G + r_c \bar{\sigma} \right)^{\frac{1}{2}} \quad (14)$$

where ϵ denotes either $+1$ or -1 . For $\mu = 0$ Eq.(12) is similar to the Friedmann one found by Maeda et al in the context of brane world cosmology with induced gravity [13]. In order to keep the potential defined on a real domain the following reality condition must hold

$$a^4 \left(1 + r_c \bar{\sigma} \right) - r_c^2 \mu \geq 0 \quad (15)$$

Now, making a Taylor expansion to second order of the potential V around the static solution yields

$$V(a) = V(a_0) + V'(a_0)(a - a_0) + \frac{1}{2}V''(a_0)(a - a_0)^2 + \mathcal{O}[(a - a_0)^3]$$

From Eq.(13) we get that the first derivative of V is

$$V' = -\frac{2a}{r_c} \left[\bar{\sigma} + \frac{2}{r_c} (1 + \epsilon Y(a)) \right] - \frac{a^2}{r_c} \left[\bar{\sigma}' + \frac{2\epsilon}{r_c} Y'(a) \right]$$

whereas the second derivate is given by

$$V'' = -\frac{2}{r_c} \left(\bar{\sigma} + \frac{2}{r_c} (1 + \epsilon Y(a)) \right) - \frac{4a}{r_c} \left(\bar{\sigma}' + \frac{2\epsilon}{r_c} Y'(a) \right) - \frac{a^2}{r_c} \left(\bar{\sigma}'' + \frac{2\epsilon}{r_c} Y''(a) \right)$$

Now, it is useful to rewrite the energy conservation as $a\bar{\sigma}'(a) = -3(\bar{p} + \bar{\sigma})$, where $\bar{p} = p/3$. Using the latter identity we can obtain the second derivative of the energy density

$$\bar{\sigma}''(a) = \frac{3}{a^2} (4 + 3\eta^2) (\bar{\sigma} + \bar{p}) \quad (16)$$

where the parameter η is defined by the relation

$$\eta(\sigma) = \frac{\partial p}{\partial \sigma} \quad (17)$$

which for ordinary matter would represent the squared speed sound: $v_s^2 = \eta$. Here, however, we simply consider η as a parameter entering the equations of state. Besides, we shall be interested in the stability of wormholes supported by ordinary matter like those found in [6]. To study the stability of the static solutions under perturbations preserving the spherical symmetry we linearize the equation of state around the static solution as follows

$$p - p_0 = \eta_0(\sigma - \sigma_0) \quad (18)$$

where the surface energy density σ_0 and the transverse pressure p_0 for a static configuration ($a = a_0$, $\dot{a} = 0$, $\ddot{a} = 0$) are given by

$$\sigma_0 = \frac{3r_c}{a_0^2} - \frac{6}{a_0} \sqrt{f(a_0)}, \quad (19)$$

$$p_0 = -\frac{r_c}{a_0^2} + \left(\frac{4f(a_0) + a_0 f'(a_0)}{a_0 \sqrt{f(a_0)}} \right) \quad (20)$$

When evaluated the potential at the static solution $a = a_0$ it is easy to see that $V(a_0) = V'(a_0) = 0$, so the potential is

$$V(a) = \frac{1}{2}V''(a_0)(a - a_0)^2 + \mathcal{O}[(a - a_0)^3] \quad (21)$$

where the second derivatives has three parts as it can be seen below

$$V''(a_0) = V''_I(a_0) + V''_{II}(a_0) + V''_{III}(a_0), \quad (22)$$

with

$$V''_I = -\frac{2}{r_c} \left(\bar{\sigma}_0 + \frac{2}{r_c} (1 + \epsilon Y(a_0)) \right), \quad (23)$$

$$V''_{II} = -\frac{4a_0}{r_c} \left(-\frac{3}{a_0} (\bar{\sigma}_0 + \bar{p}_0) + \frac{2\epsilon}{r_c} Y'(a_0) \right), \quad (24)$$

$$V''_{III} = -\frac{a_0^2}{r_c} \left(\frac{3}{a_0^2} (4 + 3\eta_0^2) (\bar{\sigma}_0 + \bar{p}_0) + \frac{2\epsilon}{r_c} Y''(a_0) \right). \quad (25)$$

Before to proceed with the stability issue we must verify that the following constraints hold at the same time

$$\text{I} \quad \sigma_0 > 0, \quad (26)$$

$$\text{II} \quad \sigma_0 + p_0 > 0, \quad (27)$$

$$\text{III} \quad a^4 \left(1 + r_c \frac{\sigma_0}{3} \right) - r_c^2 \mu \geq 0. \quad (28)$$

These conditions must be supplemented with IV: $a_0 > \sqrt{\mu}$ to ensure the existence of the wormholes manifold. Notice that I-II refer to the energy conditions (see [17]) that is, we are only interested in wormholes supported by ordinary matter, while inequality III is completely necessary to keep the potential defined on a real domain. From Fig.1 we obtain that the space spanned by the above restrictions is quite representative, in the sense that it covers a vast region. So the wormholes are physically relevant and do not correspond to a finely-tuned value of mass μ or crossover scale r_c . Of course, there are others types of solutions but not so interesting like these. To




FIG. 1: We show moduli space where the constrains I-IV are satisfy for $\mu = 1, 100$.

sum up, wormholes are stable if and only if $V''(a_0) > 0$, while for $V''(a_0) < 0$ perturbations can grow, at least until the nonlinear regime is reached.

From now on, without loss of generality, we consider the stability zones with $\mu = 1$. Let us investigate how the regions of stability change with the parameters r_c and ϵ . In the case of $\epsilon = -1$ we get that there are stable wormholes with $\eta_0 \geq 0$ for $r_c = 10$ and $r_c = 100$ (see Fig.2); moreover as it be seen that $\eta_0 \geq 1$, which would correspond to superluminal sound velocity in the wormhole throat. However, for $\epsilon = +1$ and $r_c = 10$ the model exhibits stable wormholes with $\eta_0 \geq 0$ only for small radii close to $a_0 \gtrsim 1$, whereas wormholes with large radius only are possible with $\eta_0 < 0$. In addition, for $r_c = 100$ we obtain that wormhole stable configurations have considerably large radius and $0 \leq \eta_0 \leq 0.1$, indicating that η_0 could represent the speed of sound of non-relativistic matter (see Fig. 2). Interestingly, we get that stable wormholes are achieved with values of r_c parameter far away from the General Relativity limit ($r_c \rightarrow 0$). A novel characteristic introduced by the induced gravity theory

called DGP is that these wormholes not only seem to be stable for many different choices of μ, r_c, ϵ but also they fulfil energy conditions(see Fig.3), being these conditions completely independent of ϵ . Notice that wormholes fulfilling energy conditions are not possible within Einstein's gravity [16].




FIG. 2: We show stability zones with $\mu = 1, r_c=10,100$ and for the branches $\epsilon = \pm 1$




FIG. 3: We show the energy conditions in terms of a_0 for $\mu = 1, r_c = 10, 100$.

As a final comment, let us consider the attractive or repulsive character of the wormhole geometry. The wormholes studied could be both attractive or repulsive. To

characterize this aspect of the configurations we analyze the force on a test particle at rest in the geometry described above. For this, we evaluate the radial acceleration given by

$$\mathcal{A}^r = -\Gamma_{tt}^r \left(\frac{dt}{d\tau} \right)^2 \quad (29)$$

The sign of the acceleration of a particle initially at rest is then given by minus the sign of the component Γ_{tt}^r of the connection, which for the metric considered is equal to $f'f/2$. Thus we have an attractive gravitational field for $f' > 0$ and a repulsive field for $f' < 0$ (of course we consider only the possibility $f > 0$). In our case, we find that $f'(r) = 2\mu/r^3$ so the gravitational field turns to be attractive, indicating that the wormholes are always attractive as long as $\mu > 0$. However, when the original manifold has a naked singularity ($\mu < 0$), the situation is reversed and we obtain that the field associated to the wormhole configurations is repulsive.

V. SUMMARY

We have built spherically symmetric wormholes within the induced gravity theory proposed by Dvali–

Gabadadze–Porrati (DGP). We mainly focus in the mechanical stability of wormhole configurations under radial perturbations preserving the spherical symmetry. We found stable wormholes supported by ordinary matter, that is, configurations which verify the energy conditions for many different choices of the parameter space spanned by μ, r_c, ϵ . Further, the stability analysis shows that in a scenario with a crossover scale $r_c \sim \mathcal{O}(10^2)$ (far away from the General Relativity limit) stable wormholes with very small squared speed sound $v_s^2 \in [0, 0.1]$ are obtained. Besides, studying the radial acceleration experimented by a test particle on the gravitational field we obtained that the latter one turns to be attractive, indicating that the wormholes are always attractive as long as $\mu > 0$.

Acknowledgments

MGR thanks the University of Buenos Aires for partial support under project X044. MGR is also supported by the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET).

-
- [1] G.R. Dvali, G. Gabadadze, and M. Porrati, Phys.Lett.B **485** 208-214 (2000).; G. R. Dvali and G. Gabadadze, Phys. Rev. D **63**, 065007 (2001) (e-Print: arXiv:hep-th/0008054).
 - [2] C. Deffayet, Phys. Lett. B **502**, 199 (2001); C. Deffayet, G. R. Dvali and G. Gabadadze, Phys. Rev. D **65**, 044023 (2002) (e-Print: astro-ph/0105068).
 - [3] M. G. Richarte and C. Simeone Phys. Rev. D **76**, 087502 (2007); Erratum-ibid **77**, 089903 (2008) (e-Print: arXiv:0710.2041 [gr-qc]);
 - [4] Hideki Maeda, Masato Nozawa, Phys.Rev.D **78** 024005 (2008).
 - [5] E. F. Eiroa, M. G. Richarte, and C. Simeone, Phys.Lett.A **373** 1-4 (2008), Erratum-ibid.373:2399-2400,2009, (e-Print: arXiv:0809.1623 [gr-qc]).
 - [6] M.G.Richarte, (e-Print:arXiv:1001.4034 v1).
 - [7] S.Habib Mazharimousavi, M. Halilsoy and Z. Amirabi, (e-Print:arXiv:1001.4384).
 - [8] E. Poisson and M. Visser, Phys. Rev. D **52**, 7318 (1995).
 - [9] G. Darmois, Mémorial des Sciences Mathématiques, Fascicule XXV ch V (Gauthier-Villars, Paris, 1927) ; W. Israel, Nuovo Cimento **44B**, 1 (1966); **48B**, 463(E) (1967).
 - [10] E. F. Eiroa and G. E. Romero, Gen. Relativ. Gravit. **36**, 651 (2004); F. S. N. Lobo and P. Crawford, Class. Quantum Grav. **21**, 391 (2004); M. Thibeault, C. Simeone and E. F. Eiroa, Gen. Relativ. Gravit. **38**, 1593 (2006); E. F. Eiroa and C. Simeone, Phys.Rev.D **71** 127501 (2005).
 - [11] M. A. Luty, M. Porrati, and R. Rattazzi, JHEP **0309** 029 (2003), (e-Print: hep-th/0303116);
 - [12] R. Gregory, N. Kaloper, R. Myers, and A. Padilla, JHEP **0710** 069 (2007), (e-Print: arXiv:0707.2666 [hep-th]);
 - [13] K. Maeda, S. Mizuno and T. Torii, Phys.Rev. D**68** (2003) 024033.
 - [14] A. Davidson and I. Gurwich, Phys.Rev. D**74** (2006) 044023; A.O. Barvinsky, C. Deffayet and A.Yu. Kamenshchik, (e-Print: arXiv:0912.4604).
 - [15] E. F. Eiroa, Phys. Rev. D **78**, 024018, (2008).
 - [16] M. Visser, *Lorentzian Wormholes* (AIP Press, New York, 1996); M. S. Morris and K. S. Thorne, Am. J. Phys. **56**, 395 (1988); M. Visser, Phys. Rev. D **39**, 3182 (1989); M. Visser, Nucl. Phys. **B328**, 203 (1989).
 - [17] The *weak energy condition* (WEC) states that for any timelike vector U^ξ it must be $T_{\xi\eta}U^\xi U^\eta \geq 0$; the WEC also implies, by continuity, the *null energy condition* (NEC), which means that for any null vector k^ξ it must be $T_{\xi\eta}k^\xi k^\eta \geq 0$ [3]. In an orthonormal basis the WEC reads $\rho \geq 0$, $\rho + p_l \geq 0 \forall l$ while the NEC takes the form $\rho + p_l \geq 0 \forall l$. Besides, the *strong energy condition* states that $\rho + p_l \geq 0 \forall l$, and $\rho + 3p_l \geq 0 \forall l$.