

ON THE TRACE-FREE EINSTEIN EQUATIONS AS A VIABLE ALTERNATIVE TO GENERAL RELATIVITY

GEORGE F. R. ELLIS^{1,*}, HENK VAN ELST^{1,2,†}, JEFF MURUGAN^{1,4,‡}, and JEAN-PHILIPPE UZAN^{1,3,4,§}

¹*Astrophysics, Cosmology and Gravitation Centre*

*Department of Mathematics and Applied Mathematics, University of Cape Town
Rondebosch 7701, South Africa*

²*Fakultät I: Betriebswirtschaft und Management, Karlshochschule International University
Karlstraße 36–38, 76133 Karlsruhe, Germany*

³*Institut d'Astrophysique de Paris, UMR-7095 du CNRS, Université Paris VI Pierre et Marie Curie
98 bis bd Arago, 75014 Paris, France*

⁴*National Institute for Theoretical Physics (NITheP), Stellenbosch 7600, South Africa*

April 2, 2011

Abstract

The quantum field theoretic prediction for the vacuum energy density leads to a value for the effective cosmological constant that is incorrect by between 60 to 120 orders of magnitude. We review an old proposal of replacing Einstein's Field Equations by their trace-free part (the Trace-Free Einstein Equations), together with an independent assumption of energy–momentum conservation by matter fields. While this does not solve the fundamental issue of why the cosmological constant has the value that is observed cosmologically, it is indeed a viable theory that resolves the problem of the discrepancy between the vacuum energy density and the observed value of the cosmological constant. However, one has to check that, as well as preserving the standard cosmological equations, this does not destroy other predictions, such as the junction conditions that underlie the use of standard stellar models. We confirm that no problems arise here: hence, the Trace-Free Einstein Equations are indeed viable for cosmological and astrophysical applications.

1 Introduction

The interpretation of dark energy is a major puzzle [19]. The gravitational effect of the quantum vacuum is expected to be equivalent to an effective cosmological constant [21], which according to the standard view will cause an accelerated expansion of the universe. However, simple estimates of its expected magnitude are very large, exceeding the observed value by between 60 and 120 orders of magnitude [20, 4, 11]; a blatant contradiction with observations. This indicates a profound discrepancy between General Relativity (GR) and Quantum Field Theory (QFT) — a major problem for theoretical physics [17, 20]. This will presumably be resolved when a full-blown theory of quantum gravity is finalized and accepted. However, the field theory view of gravity as a massless spin-2 field, and with the quantum vacuum contributing to the cosmological constant, is a half-way house between GR and a full quantum gravity theory (which will need to have both GR and spin-2 QFT as appropriate limits). We need to resolve that discrepancy, no matter what final quantum gravity theory is adopted.

*E-mail: George.Ellis@uct.ac.za

†E-mail: hvanelst@karlshochschule.de

‡E-mail: jeff@nassp.uct.ac.za

§E-mail: uzan@iap.fr

The key issue is: “*What are the gravitational effects of the vacuum energy?*” One of the proposals for solving the problem, summarized by Weinberg [20], is use of Trace-Free Einstein Equations (TFE) instead of the standard Einstein’s Field Equations [7, 8]. If these trace-free equations are adopted, the vacuum energy has no gravitational effect. This does not determine a unique value for the effective cosmological constant, but it does solve the huge discrepancy between theory and observation. However, that resolution is dependent on this form of the field equations giving the same results as the standard equations, not only for the solar system and black holes (which will follow because the vacuum field equations are unchanged), but also on the one hand for cosmological models, and on the other for astrophysical objects such as stellar models, where one has standard junction conditions between the interior and exterior solutions. The former must work out as expected (because of Weinberg’s results), but how this happens is not immediately obvious. In the latter case, it is not *a priori* clear that the usual results should remain true. Indeed, one might even expect a discrepancy between the stellar mass and the mass of the exterior Schwarzschild solution.

In this article, we confirm that the trace-free Einstein equations are compatible with either case, and so are indeed a viable alternative to the GR field equations — with the improvement that, unlike Einstein’s Field Equations, they do not suffer from the crucial discrepancy with the standard results of quantum field theory, which imply a very large value for the vacuum energy: so large that even solar system results would be affected. Pauli in the 1930’s (as quoted by Straumann: [17], pp.6-7) estimated the zero-point energy of the radiation field, assuming that the cut-off of the theory was the electron mass. He then estimated that the radius of the de Sitter universe ($\Lambda \simeq 1/R^2$) would be smaller than the Earth–Moon distance; actually, one can easily compute that $R \simeq 31$ km [17]. So this value drastically affects the Solar system, since there would be no solar system. It seems prudent to look for a way out.

2 Basic problem

Classical gravitational dynamics is encoded in the Einstein Field Equations (EFE) [7, 8]

$$G_{ab} + \Lambda g_{ab} = R_{ab} - \frac{1}{2} R g_{ab} + \Lambda g_{ab} = \frac{8\pi G}{c^4} T_{ab} , \quad (1)$$

where Λ is the cosmological constant. These are subject to the conservation equations

$$\nabla_b G^{ab} = 0 \quad \Rightarrow \quad \nabla_b T^{ab} = 0 , \quad (2)$$

for the total energy–momentum tensor, which guarantee the consistency of the time development of the EFE.

In the standard cosmological application, the metric tensor is assumed to take the spatially homogeneous and isotropic form

$$ds^2 = -c^2 dt^2 + a^2(t) d\sigma^2 , \quad u^a = \delta_0^a , \quad (3)$$

with $a(t)$ a universal time-dependent scale factor, $d\sigma^2$ the metric of a 3-space of constant curvature k , and u^a the normalised matter 4-velocity field ($u_a u^a = -1$). Because of the symmetries of the metric, the energy–momentum tensor of the matter necessarily takes a perfect fluid form:

$$T_{ab} = \left(\rho + \frac{p}{c^2} \right) c^2 u_a u_b + p g_{ab} , \quad (4)$$

where the matter mass density ρ and the matter isotropic pressure p are related through an equation of state $p = p(\rho)$. Whenever the energy–momentum tensor takes the form (4), the conservation equations (2) give the mass–energy conservation relation

$$\dot{\rho} + 3 \frac{\dot{a}}{a} \left(\rho + \frac{p}{c^2} \right) = 0 , \quad (5)$$

where a dot denotes a covariant derivative with respect to proper time, d/dt . For the case of the Robertson–Walker metric (3), the EFE (1) reduce to two non-trivial equations: the Raychaudhuri equation

$$3 \frac{\ddot{a}}{a} + 4\pi G \left(\rho + 3 \frac{p}{c^2} \right) - \Lambda c^2 = 0 , \quad (6)$$

and the Friedmann equation

$$3 \left(\frac{\dot{a}}{a} \right)^2 = - \frac{3kc^2}{a^2} + 8\pi G \rho + \Lambda c^2 . \quad (7)$$

The latter equation is a first integral of the other two; indeed any two of (5) to (7) imply the third. When they are satisfied, all 10 EFE (1) are satisfied.

The key equation as regards gravitational attraction is the Raychaudhuri equation (6), which shows that $(\rho + 3p/c^2)$ is the active gravitational mass density. The equation of state for a vacuum

$$p_{\text{vac}} = - \rho_{\text{vac}} c^2 \quad (8)$$

shows this is negative:

$$\left(\rho_{\text{vac}} + 3 \frac{p_{\text{vac}}}{c^2} \right) = - 2\rho_{\text{vac}} , \quad (9)$$

where we can represent the effect of the vacuum in (6) either as a fluid with equation of state (8), or as an effective cosmological constant,

$$\Lambda_{\text{vac}} = \frac{8\pi G}{c^2} \rho_{\text{vac}} . \quad (10)$$

The problem is that the QFT view of the vacuum as an infinite set of oscillators, each with zero-point oscillatory energy $\frac{1}{2}\hbar\omega_k$, gives a diverging value for the vacuum energy E_{vac} . With a suitable high-energy cut-off, the vacuum energy density is estimated by Weinberg [20] to be of the order

$$\langle \rho_{\text{vac}} \rangle \simeq 2 \times 10^{71} \text{ GeV}^4 , \quad (11)$$

whereas the effective value of the cosmological constant as determined by astronomical observations is of the order

$$\langle \rho_{\text{obs}} \rangle \simeq 10^{-47} \text{ GeV}^4 . \quad (12)$$

Because the value (11) is a constant, it has no effect on local dynamics, and so can be subtracted off the total energy density as far as local physics is concerned, but on the standard view, because of (9), it will have a gravitational effect hugely bigger than the observed value (12).

There are three ways out of this problem.

1. Other fields may contribute negative energy densities that will cancel the positive terms, and leave a very small residue as observed; or maybe there is a symmetry implying $\rho_{\text{vac}} + \Lambda_{\text{bare}} = 0$, and one has to have extra fields ('quintessence') to give the observed acceleration. This is indeed possible in principle; for example, each mode of a Dirac field gives a negative contribution to the vacuum energy (hence, this is the option that would be realised via supersymmetry, were supersymmetry not broken in the real universe). But it is very hard to make this work in practice: many fields contribute to the vacuum energy density, and it is highly unlikely they would just happen to cancel the value (11) accurately to 120 decimal places, but not exactly, so as to give (12).
2. The value in (11) is only an expectation value: maybe we live in a multiverse where ρ_{vac} takes all sorts of values, and anthropic selection effects determine we live in a universe domain where its value (12) is very far from the expectation value. This is a possible solution, but many regard it as an act of desperation, abandoning the idea of Λ as a fundamental constant in favour of regarding its value as being contingent.

3. Maybe the EFE (1) are not the true effective equations of gravitational interactions: a variant to the EFE arises from the underlying quantum gravity theory, and negates the gravitational effect of the vacuum.

We explore aspects of the latter option in this article.

3 Trace-free Einstein gravity

Einstein continually worried about what should go on the right-hand side of the relativistic gravitational field equations. An interesting proposal is to take the trace-free part of the EFE to get the Trace-Free Einstein Equations (TFE), which is a subset of the EFE that can give back the full EFE with an integration constant. This is an old proposal, essentially initiated by Einstein [9] himself, and developed by many others since, see [20, pp. 11–13] and references given there. More recently, it has been developed under the name of “unimodular gravity”, see [1, 18, 10, 15] and references therein.

We use a hat to denote the trace-free part of a symmetric tensor: so

$$\hat{G}_{ab} := R_{ab} - \frac{1}{4} R g_{ab}, \quad \hat{T}_{ab} := T_{ab} - \frac{1}{4} T g_{ab} \quad \Rightarrow \quad \hat{G}_a{}^a = 0, \quad \hat{T}_a{}^a = 0.$$

On taking its trace-free part, the EFE (1) implies the TFE

$$\hat{G}_{ab} = \frac{8\pi G}{c^4} \hat{T}_{ab} \quad \Leftrightarrow \quad R_{ab} - \frac{1}{4} R g_{ab} = \frac{8\pi G}{c^4} \left(T_{ab} - \frac{1}{4} T g_{ab} \right). \quad (13)$$

We adopt these as the gravitational field equations, instead of (1). The twice-contracted second Bianchi identity for the Einstein tensor G_{ab} still holds:

$$\nabla_b G^{ab} = 0. \quad (14)$$

However, now the corresponding divergence relation for the energy–momentum tensor T_{ab} ,

$$\nabla_b T^{ab} = 0, \quad (15)$$

is no longer a consequence of the geometrical identity (14), as in (2): it is a *separate assumption*.

The gravity theory based on the TFE recovers all the vacuum solutions of the EFE unchanged, so e.g. results from the Schwarzschild and Kerr black hole solutions are still valid. However, it no longer has a cosmological constant problem, as Λ does not affect spacetime curvature. Indeed, for a perfect fluid the matter source term is the manifestly trace-free energy–momentum tensor

$$\hat{T}_{ab} = \left(\rho + \frac{p}{c^2} \right) c^2 \left(u_a u_b + \frac{1}{4} g_{ab} \right); \quad (16)$$

hence, matter enters the field equations only in terms of the inertial mass density $(\rho + p/c^2)$, which vanishes for a vacuum. However, as discussed in [20], the theory acquires a new *integrability condition*: differentiating (13), and using (14) and (15), gives

$$\nabla_b \left(R^{ab} - \frac{1}{4} R g^{ab} \right) = \frac{8\pi G}{c^4} \nabla_b \left(T^{ab} - \frac{1}{4} T g^{ab} \right) \quad \Rightarrow \quad \nabla_a R = - \frac{8\pi G}{c^4} \nabla_a T. \quad (17)$$

Integrating, $\left(R + \frac{8\pi G}{c^4} T \right)$ is a constant:

$$R + \frac{8\pi G}{c^4} T =: 4\hat{\Lambda}. \quad (18)$$

Substituting into (13) to eliminate T gives back (1), but with a new effective cosmological constant:

$$R_{ab} - \frac{1}{2} R g_{ab} + \hat{\Lambda} g_{ab} = \frac{8\pi G}{c^4} T_{ab} . \quad (19)$$

So the way it works is as follows: we *assume* both the TFE (13) and the matter conservation equations (15). The integrability condition (17) follows from these equations. Integrating gives (19). This is not surprising in that we have assumed the validity of the trace-free part of the EFE (1); and (18) is just the trace of those equations. The same consistency results hold as for the full EFE: by (14) the time evolution equations amongst the TFE preserve the constraints. This result can be implemented as a dynamical theory where the spacetime volume density $\sqrt{-g}$ is *not* a dynamical variable [20].

Hence, we have a remarkable result [20]: *the TFE (13) together with the differential relations (15) are functionally equivalent to the EFE (19), with the cosmological constant an arbitrary integration constant $\hat{\Lambda}$ unrelated to the vacuum energy Λ_{vac} .* Note this is not the same as either

$$\hat{G}_{ab} = \frac{8\pi G}{c^4} T_{ab} \quad \Rightarrow \quad T = 0, \quad R = \text{constant} \quad (20)$$

(as proposed by Einstein in 1919, see [9]), or

$$G_{ab} = \frac{8\pi G}{c^4} \hat{T}_{ab} \quad \Rightarrow \quad R = 0, \quad T = \text{constant} \quad (21)$$

(which also decouples spacetime curvature from vacuum energy). Each of these proposals has 10 field equations, and allows only very restricted forms of matter to occur. The proposal that makes physical sense is (13), where both sides of the equation have been specifically constructed so as to have the same symmetry. Then we have 9 field equations and the energy–momentum tensor of the matter can have a non-zero trace, but only the trace-free part gravitates; this can accommodate generic inhomogeneous matter.

What about experiments? *The experimental predictions of the two theories are the same, so no experiment can tell the difference between them, except for one fundamental feature: the EFE (confirmed in the solar system and by binary pulsar measurements to high accuracy) together with the QFT prediction for the vacuum energy density (confirmed by Casimir force measurements) give the wrong answer by many orders of magnitude; the TFE does not suffer this problem.* In this respect, the TFE are strongly preferred by experiment.

This trace-free Einstein gravity theory then should give cosmology without the vacuum energy problem. We explore this in the next section, but we will not deal with the relation of this theory to commonplace variational principle approaches, which is adequately covered elsewhere (see [1, 20, 15] and references there).

4 FLRW cosmologies

This works out in the spatially homogeneous and isotropic case as a special case of the above general theory (as, of course, it has to). That is, starting from (5), (6) and (7), we first determine a trace-free dynamic equation by eliminating Λ between (6) and (7), and then derive an integrability condition for this equation. Thereafter, we show how this latter equation can be integrated to recover (6) and (7) with a new effective cosmological constant $\hat{\Lambda}$ as an integration constant, independent of any dynamical values we might assign to the vacuum energy. Thus, this solves the major problem of a vacuum energy many orders of magnitude larger than measured by cosmological observations.

In detail: the trace-free equation is

$$\frac{\ddot{a}}{a} - \left(\frac{\dot{a}}{a}\right)^2 - \frac{kc^2}{a^2} = -4\pi G \left(\rho + \frac{p}{c^2}\right) . \quad (22)$$

The spacetime Ricci curvature scalar R is given by

$$Rc^2 = -\frac{8\pi G}{c^2}T + 4\Lambda c^2 = 8\pi G\left(\rho - 3\frac{p}{c^2}\right) + 4\Lambda c^2 = 6\left[\frac{\ddot{a}}{a} + \left(\frac{\dot{a}}{a}\right)^2 + \frac{kc^2}{a^2}\right]. \quad (23)$$

The time derivative of this equation is the needed integrability condition for (22), encoding the fact that Λ is a constant (see (17)). It has the form

$$0 = 6\frac{d}{dt}\left[\frac{\ddot{a}}{a} + \left(\frac{\dot{a}}{a}\right)^2 + \frac{kc^2}{a^2} - \frac{4\pi G}{3}\left(\rho - 3\frac{p}{c^2}\right)\right]. \quad (24)$$

Now (24), being a vanishing total time derivative, can be easily integrated to yield an integration constant $\frac{2}{3}\hat{\Lambda}$. Eliminating \ddot{a}/a between (22) and the integral of (24), and including the (now assumed) mass–energy conservation relation (5), one recovers the original dynamic equations (6) and (7), but with a renormalized Λ . That is, we obtain in this way the Raychaudhuri equation

$$3\frac{\ddot{a}}{a} + 4\pi G\left(\rho + 3\frac{p}{c^2}\right) + \hat{\Lambda}c^2 = 0, \quad (25)$$

and the Friedmann equation

$$3\left(\frac{\dot{a}}{a}\right)^2 = -\frac{3kc^2}{a^2} + 8\pi G\rho + \hat{\Lambda}c^2, \quad (26)$$

with effective cosmological constant $\hat{\Lambda}$ — an arbitrary integration constant, unrelated to Λ_{vac} . This solves the basic discrepancy between QFT estimates of the energy density of the quantum vacuum, and the disastrous result if we assume this is a source term in the Raychaudhuri equation in an obvious way. Thus, we arrive at

Hypothesis: The EFE are not the true effective equations of gravity: rather — whatever the underlying quantum theory of gravity — the effective theory is trace-free Einstein gravity, as described above.

In that case the basic equations of cosmology are (22) and (5), supplemented by an equation of state determining the pressure from the mass density, and (25) and (26) are consequences. The value of the vacuum term Λ_{vac} does not affect the cosmic expansion. Note the following remarkable feature: *the active gravitational mass density $\rho_{\text{grav}} = (\rho + 3p/c^2)$ does not occur in either of the equations (22) or (5) that we take as the basis of the dynamics: only the inertial mass density $\rho_{\text{inert}} = (\rho + p/c^2)$ occurs there. Nevertheless, in the end ρ_{grav} turns out to be the effective gravitational mass density, see (25).* What happens is that while (5) determines the time evolution of the energy density, the assumed equation of state determines the time evolution of the pressure, and does so in such a way that (25) results. It is interesting to note that (22) is in the form of the dynamical equations often used for studies of inflationary universe models.¹

Let us also mention an important point concerning scalar field cosmology. The transformation $T_{ab} \rightarrow T_{ab} + \Lambda g_{ab}$ leaves \hat{T}_{ab} and the conservation equation (15) unchanged only if Λ is a constant. In the case of a self-interacting scalar field φ evolving in a potential $V(\varphi)$, it is easily checked for the energy–momentum tensor that

$$T_{ab} = \partial_a\varphi\partial_b\varphi - \frac{1}{2}[\partial_c\varphi\partial^c\varphi + 2V(\varphi)]g_{ab} \quad \Rightarrow \quad \hat{T}_{ab} = \partial_a\varphi\partial_b\varphi - \frac{1}{4}(\partial_c\varphi\partial^c\varphi)g_{ab} \quad (27)$$

whatever the potential, the latter of which, therefore, does not appear in the TFE (17), and so has no gravitational effect. Nevertheless, the potential is, of course, of dynamical relevance, since it cannot be

¹We thank Roy Maartens for this comment.

eliminated from the conservation equation as long as $\dot{\varphi} \neq 0$. For an FLRW scalar field cosmology, (15) yields the equation of motion

$$\dot{\varphi} \left(\ddot{\varphi} + 3H\dot{\varphi} + \frac{dV}{d\varphi} \right) = 0. \quad (28)$$

The dynamics of a universe containing a scalar field (e.g. during inflation) will then be the same as in standard GR. Note, however, that we now have the freedom to shift the minimum of the potential at will, since $V(\varphi) \rightarrow V(\varphi) + V_0$ leaves the equation of motion unchanged.

5 Junction conditions

To complete the picture of the trace-free Einstein gravity proposal, we need to verify that in the case of a stellar model the interior and exterior solutions match, so that the mass measured outside is the same as the mass of the interior solution; if this was not true, it would be a disaster for these equations, as they would not give standard results for stellar structure models. In the GR context, this issue is settled by the Darmois–Israel junction conditions [6, 12], which arise from the local embedding relations of Gauß and Codazzi for a timelike or spacelike 3-surface in a 4-dimensional spacetime manifold.

5.1 Covariant 3 + 1 viewpoint

Consider a non-null 3-surface $\Sigma : \{\phi = \text{constant}\}$ (of codimension 1) embedded in a spacetime manifold, with unit normal n_a given by

$$n_a := \nabla_a \phi, \quad (29)$$

such that

$$n_a n^a = \varepsilon = \begin{cases} +1, & \text{when } \Sigma \text{ is timelike} \\ -1, & \text{when } \Sigma \text{ is spacelike} \end{cases} \quad (30)$$

applies. Then

$$\delta_a^b := \varepsilon n_a n^b + h_a^b \quad (31)$$

defines tensors $\perp_a^b := \varepsilon n_a n^b$ and h_a^b which project orthogonally and tangentially to Σ , respectively, with $0 = h_a^b n_b$. The covariant derivative of n_a is given by

$$\nabla_a n_b = \varepsilon n_a \dot{n}_b - K_{ab}, \quad (32)$$

which defines the non-geodesicity of n_a by $\dot{n}_a := n^b \nabla_b n^a$, and the extrinsic curvature tensor of Σ by $K_{ab} := -h_a^c h_b^d \nabla_c n_d =: -D_{(a} n_{b)}$, where D_a is the induced connection of Σ . Note that $0 = \dot{n}_a n^a = K_{ab} n^b$ applies.

Certain orthogonal and tangential projections (by means of \perp_a^b and h_a^b) of the 4-dimensional Ricci identity generate for Σ the well-known local embedding equations of Gauß and Codazzi (see e.g. [16]),

$${}^3R_{abcd} - \varepsilon(K_{ac}K_{bd} - K_{ad}K_{bc}) = h_a^e h_b^f h_c^g h_d^h R_{efgh} \quad (33)$$

$$D_a K_{bc} - D_b K_{ac} = -h_a^d h_b^e h_c^f n^g R_{defg}, \quad (34)$$

with ${}^3R_{abcd}$ the intrinsic 3-Riemann curvature tensor of Σ . Linear combinations of contractions of these relations then lead to (see [13, 16])

$$n^a n^b G_{ab} = \frac{1}{2} \left(K^2 - K_{ab} K^{ab} - \varepsilon {}^3R \right) \quad (35)$$

$$-n^b h_a^c G_{bc} = D_b K_a^b - D_a K, \quad (36)$$

where $K := K_a^a$. In the absence of matter surface layers, Darmois' junction conditions [6] require that across Σ the induced 3-metric and the extrinsic curvature be continuous, i.e.,

$$0 = [h_{ab}] \Rightarrow 0 = [D_a], \quad 0 = [{}^3R_{abcd}], \quad (37)$$

$$0 = [K_{ab}] \Rightarrow 0 = [D_a K_{bc}], \quad (38)$$

employing $[Q] := Q_+ - Q_-$ to denote the change of any quantity Q across Σ . With (35) and (36), this leads to Israel's junction conditions [12]

$$0 = [n_a n_b G^{ab}], \quad 0 = [n_b h_c^a G^{bc}]. \quad (39)$$

As this is a purely geometrical result, it does not make a qualitative difference whether we use (i) the EFE (1), or (ii) the TFE (13) together with (18), to replace the Einstein tensor G_{ab} in these relations.

5.2 Including a matter surface layer

The previous consideration of the Darmois–Israel junction conditions assumes that there is no matter surface layer on the 3-surface Σ between the two parts of spacetime to be matched. An alternative approach to the derivation of the junction conditions relies on the use of a local coordinate approach. Using the same conventions as in the previous paragraph, in a parametric local representation of Σ given by $x^a = X^a(\sigma^i)$ (where $a, b, \dots = 0, \dots, 3$ and $i, j, \dots = 1, \dots, 3$) the coordinate components of the induced 3-metric are expressed by

$$\gamma_{ij} := \frac{\partial X^a}{\partial \sigma^i} \frac{\partial X^b}{\partial \sigma^j} g_{ab}. \quad (40)$$

Viewed from the 4-dimensional perspective, this is the first fundamental form $h^{ab} = \frac{\partial X^a}{\partial \sigma^i} \frac{\partial X^b}{\partial \sigma^j} \gamma^{ij} = g^{ab} - \varepsilon n^a n^b$. If we assume that n_a is smoothly continued into the 4-dimensional spacetime along a set of geodesics, i.e. such that $\dot{n}_a = 0$, the extrinsic curvature reduces to $K_{ab} = -\nabla_a n_b$ [see (32)]. While g_{ab} is everywhere continuous, its derivative along n_a may not be continuous across Σ , so that K_{ab} may also be discontinuous.

Now consider a neighbourhood of Σ represented in terms of a Gaussian normal coordinate system, where the coordinate s varying in the interval $s_- < s < s_+$ measures the distance to Σ ; we assume Σ to be located at $s = 0$. This implies that $n_a = \nabla_a s$. We then denote by $Q' = \partial_s Q = n^a \partial_a Q$ the partial derivative of any quantity Q along the normal direction to Σ . The change of Q across Σ is given by

$$[Q] = Q_+ - Q_- = \overline{Q}' := \int_{s_-}^{s_+} Q' ds. \quad (41)$$

This definition is such that $Q' \rightarrow \overline{Q}' \delta(s)$ in the limit of an infinitesimally thin 3-surface. We use angular brackets to denote the mean value,

$$\langle Q \rangle := \frac{1}{2} (Q_+ + Q_-). \quad (42)$$

In Gaussian normal coordinates, equations (33) and (34) are complemented by the spatial projection [13]

$$h_a^c h_d^b G_c^d = {}^3G_a^b + \varepsilon n^c \partial_c (K_a^b - K h_a^b) - K K_a^b + \frac{1}{2} (K^2 + K_{cd} K^{cd}) h_a^b. \quad (43)$$

If there is matter localised on Σ , then its energy–momentum tensor can be computed as

$$S_a^b = \overline{T}_a^b = \lim_{s \rightarrow 0} \int_{-s}^s T_a^b ds, \quad (44)$$

which satisfies $0 = S_{ab}n^b$. In the infinitesimally thin limit, the 4-dimensional energy–momentum tensor on Σ is of the form $T_{ab} = S_{ab}\delta(s)$. For the geometry of Σ to be well-defined, both g_{ab} and h_{ab} should be continuous across Σ and thus contain neither a $\delta(s)$ factor nor a discontinuity. The first implies that ${}^3R_{abcd}$ and thus ${}^3R_{ab}$ have no $\delta(s)$ factor, while the second implies that K_{ab} has no $\delta(s)$ factor. Integration of (43) then implies that

$$h_a{}^c h_d{}^b \bar{G}_c{}^d = \varepsilon \left[K_a{}^b - K h_a{}^b \right], \quad (45)$$

since only the terms containing a derivative along s contribute to the integral, i.e., $G_{ab} \simeq \varepsilon(K'_{ab} - K' h_{ab})$. Using that $G_{ab} = (G_{cd}n^c n^d)n_a n_b + \varepsilon G_{cd}n^c h_a{}^d n_b + G_{cd}h_a{}^c h_b{}^d$, and the fact that the integrals along n^a of expressions (33) and (34) vanish even when matter is localized on Σ , we conclude that

$$\bar{G}_a{}^b = \varepsilon \left[K_a{}^b - K h_a{}^b \right]. \quad (46)$$

With the standard EFE (1), this would imply that $\left[K_a{}^b - K h_a{}^b \right] = \frac{8\pi G}{c^4} S_a{}^b$, a generalisation of (38).

For the TFE (13), we first need to compute the jump in the trace-free Einstein tensor \hat{G}_{ab} by using that $G \simeq -2\varepsilon K'$, so that

$$\bar{\hat{G}}_{ab} = \varepsilon \left[K_{ab} - K \left(h_{ab} - \frac{1}{2} g_{ab} \right) \right] = \frac{8\pi G}{c^4} \left(S_{ab} - \frac{1}{4} S g_{ab} \right). \quad (47)$$

While the junction condition for the trace K cannot be extracted from this equation, one has to rely on (17). We decompose the total energy–momentum tensor according to $T_{ab} = T_{ab}^{\text{mat}} + T_{ab}^{\Sigma}$, with $T_{ab}^{\Sigma} = S_{ab}\delta(s)$. The hypothesis (15) implies that

$$\nabla_b T_{\Sigma}^{ab} = f^a = -\nabla_b T_{\text{mat}}^{ab}. \quad (48)$$

We assume for simplicity that Σ does not interact with the matter fields (see e.g. [3] for the treatment of such a case, and [5] for an example in which Σ is coupled to a form field allowing a jump in the cosmological constant). Since $\nabla_b T_{\text{mat}}^{ab} = 0$, this implies that $n_b (T_{\text{mat}}^{ab})' = 0$, and thus $[T_{\text{mat}}^{ab}] \perp_{ab} = 0$. Then, (17) implies $n^a \nabla_a \left(R + \frac{8\pi G}{c^4} T \right) = 0$, so that $R + \frac{8\pi G}{c^4} T$ remains constant across Σ (i.e., $\bar{\Lambda} = 0$). Now, since $R \simeq \varepsilon 2K'$, we deduce that $\bar{R} = \varepsilon 2[K] = -\frac{8\pi G}{c^4} S$, where the second equality arises from the constancy of $R + \frac{8\pi G}{c^4} T$, which implies $\bar{R} = -\frac{8\pi G}{c^4} S$. It follows that $\frac{1}{2} \varepsilon [K] g_{ab} = -\frac{8\pi G}{c^4} \frac{1}{4} S g_{ab}$, so that the junction condition (47) reduces to

$$\varepsilon [K_{ab} - K h_{ab}] = \frac{8\pi G}{c^4} S_{ab}, \quad (49)$$

i.e., to the same relation as in classical GR.

In conclusion, the standard GR stellar structure models, with an interior solution matched to an exterior solution across a suitable 3-surface Σ , will remain valid in the case of trace-free Einstein gravity. There will not be a problem of the interior and exterior masses not matching. Similar issues arise for the junction conditions in Swiss-Cheese models. Again, the TFE will be acceptable: the usual mass matching condition will be fulfilled, so they do not lead to anomalies here either.

6 Viability of the TFE equations

We have revisited the possibility that the true effective gravitational field equations are given by the TFE, implying that only the trace-free part of the energy–momentum tensor T_{ab} of matter is gravitating.

Then the effective cosmological constant $\hat{\Lambda}$ is a constant of integration that is arbitrarily disposable (as in classical GR), and, hence, is independent of any fundamental value assigned to Λ (cf. [9, p. 196]). We do not require a fine tuning $\rho_{\text{vac}} + \Lambda \simeq \Lambda_{\text{obs}}$, because ρ_{vac} is not gravitating.

Thus, employing the TFE in place of the EFE appears to be a good theoretical assumption to make: any huge Λ_{vac} is powerless to affect cosmology, or indeed the solar system, as the zero point energy will not affect spacetime geometry. The EFE will be as usual, but with $\hat{\Lambda}$ an integration constant that may be small, or may be zero. As observations indicate, this constant corresponds to a particular cosmological length scale ($\hat{\Lambda} \simeq H_0^2$) that should be determined from initial conditions for our universe; see (17). In that sense, the vacuum energy problem vanishes in the trace-free Einstein gravity proposal, while the almost equality between $\hat{\Lambda}$ and the Hubble constant, i.e., the coincidence problem, remains.

We have checked here that these equations are compatible with usual cosmological models and also with standard junction conditions, without and with a matter surface layer. In particular, this ensures that stellar structure models will be the same as in GR: there will not be any mass anomaly between the interior and exterior solutions. Hence, the TFE work both for cosmology and for astrophysics.

Overall, this proposal does not solve the issue of why the cosmological constant has the value it has today, but it does resolve the issue of why it does not have the huge value implied by the obvious use of the QFT prediction for the vacuum energy in conjunction with the EFE. The patently incorrect result obtained in this way is a major crisis for theoretical physics, because it suggests a profound contradiction between two of our most successful theories, namely QFT and GR. Use of the TFE instead of the EFE solves that problem.

Acknowledgments

We thank Roy Maartens for helpful comments on an earlier draft of this article.

References

- [1] James L. Anderson and David Finkelstein (1971) “Cosmological constant and fundamental length” *Am. J. Phys.* **39** 901–904.
- [2] Richard A. Battye and Brandon Carter (2001) “Generic junction conditions in brane-world scenarios” *Phys. Lett. B* **509** 331–336 [arXiv:hep-th/0101061 v1].
- [3] Richard A. Battye, Brandon Carter, Andrew Mennim and Jean-Philippe Uzan (2001) “Einstein equations for an asymmetric brane-world” *Phys. Rev. D* **64** 124007 [arXiv:hep-th/0105091 v1].
- [4] Sean M. Carroll (2001) “The cosmological constant” *Living Rev. Rel.* **4** 1–56 [Online Article]: cited [Aug 3, 2010], <http://www.livingreviews.org/lrr-2001-1> [arXiv:astro-ph/0004075 v2].
- [5] Brandon Carter and Jean-Philippe Uzan (2001) “Reflection symmetry breaking scenarios with minimal gauge form coupling in brane world cosmology,” *Nucl. Phys. B* **606** 45–58 [arXiv:gr-qc/0101010 v2].
- [6] Georges Darmois (1927) *Les équations de la gravitation Einsteinienne* (Chapitre V) Mémorial de Sciences Mathématiques, Fascicule XXV (Paris: Gauthier–Villars).
- [7] Albert Einstein (1915) “Die Feldgleichungen der Gravitation” *Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.)* **1915** 844–847.
- [8] Albert Einstein (1917) “Kosmologische Betrachtungen zur allgemeinen Relativitätstheorie” *Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.)* **1917** 142–152.

- English translation: (1923) “Cosmological considerations on the general theory of relativity” In *The Principle of Relativity* (London: Methuen). Reprinted: (1952) (Mineola, NY: Dover Publications) 175–188.
- [9] Albert Einstein (1919) “Spielen Gravitationsfelder im Aufbau der materiellen Elementarteilchen eine wesentliche Rolle?” *Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.)* **1919** 349–356. English translation: (1923) “Do gravitational fields play an essential part in the structure of the elementary particles of matter?” In *The Principle of Relativity* (London: Methuen). Reprinted: (1952) (Mineola, NY: Dover Publications) 189–198.
- [10] David R. Finkelstein, Andrei A. Galiatdinov and James E. Baugh (2001) “Unimodular relativity and cosmological constant” *J. Math. Phys.* **42** 340–346 [arXiv:gr-qc/0009099v1].
- [11] Joshua A. Frieman, Michael S. Turner and Dragan Huterer (2008) “Dark energy and the accelerating universe” *Ann. Rev. Astron. Astrophys.* **46** 385–432 [arXiv:0803.0982v1 [astro-ph]].
- [12] Werner Israel (1966) “Singular hypersurfaces and thin shells in general relativity” *Nuovo Cimento B* **44** 1–14.
Erratum: (1967) *Nuovo Cimento B* **48** 463.
- [13] Charles W. Misner, Kip S. Thorne and John A. Wheeler (1973) *Gravitation* (San Francisco: Freeman).
- [14] Patrick Peter and Jean-Philippe Uzan (2009) *Primordial cosmology* (Oxford: Oxford University Press).
- [15] Lee Smolin (2009) “Quantization of unimodular gravity and the cosmological constant problems” *Phys. Rev. D* **80** 084003 [arXiv:0904.4841v1 [hep-th]].
- [16] Hans Stephani (1991) *Allgemeine Relativitätstheorie* 4th Edition (Berlin: Dt. Verlag d. Wissenschaften).
- [17] Norbert Straumann (2009) “Wolfgang Pauli and modern Physics” *Space Sci. Rev.* **148** 25–36 [arXiv:0810.2213v1 [physics.hist-ph]].
- [18] William G. Unruh (1989) “Unimodular theory of canonical quantum gravity” *Phys. Rev. D* **40** 1048–1052.
- [19] Jean-Philippe Uzan (2010) “Dark energy, gravitation and the Copernican principle”. In *Dark Energy: observational and theoretical approaches*, Edited by P. Ruiz-Lapuente (Cambridge: Cambridge University Press) [arXiv:0912.5452v1 [gr-qc]].
- [20] Steven Weinberg (1989) “The cosmological constant problem” *Rev. Mod. Phys.* **61** 1–23.
- [21] Yakov B. Zel’dovich (1967) “Cosmological Constant and Elementary Particles” *JETP Lett.* **6** 316– [Pisma Zh. Eksp. Teor. Fiz. **6** (1967) 883–].