

Phase transitions far away from stationary points of the energy landscape

Michael Kastner*

*National Institute for Theoretical Physics (NITheP), Stellenbosch 7600, South Africa and
Institute of Theoretical Physics, University of Stellenbosch, Stellenbosch 7600, South Africa*

Dhagash Mehta†

Department of Physics, Syracuse University, Syracuse, NY 13244, USA

(Dated: April 29, 2022)

The stationary points of the potential energy function of the ϕ^4 model on a two-dimensional square lattice with nearest-neighbor interactions are studied. We employ two numerical methods and corroborate the results by an analytical upper bound on the potential energy at stationary points. Our findings, supported by all three methods, have striking consequences regarding phase transitions and their relation to stationary points (or, equivalently, changes of configuration space topology): A theorem by Franzosi and Pettini [Phys. Rev. Lett. **92**, 060601 (2004)], purporting a rigorous relation between stationary points of the potential energy function and phase transitions, is shown to be incorrect. A numerical method by Franzosi *et al.* [Phys. Rev. Lett. **84**, 2774 (2000)], yielding results in support of that theorem, is found to be unreliable. These findings are supplemented by a more detailed study of a certain region of the potential energy landscape.

PACS numbers: 05.50.+q, 64.60.A-, 05.70.Fh

The stationary points of the potential energy function or other classical energy functions can be employed to calculate or estimate certain physical quantities of interest. Well-known examples include transition state theory or Kramers’s reaction rate theory for the thermally activated escape from metastable states, where the barrier height (corresponding to the difference between potential energies at certain stationary points of the potential energy function) plays an essential role. More recently, a large variety of related techniques has become popular under the name of “energy landscape methods” [1], allowing for applications to many-body systems as diverse as metallic clusters, biomolecules and their folding transitions, or glass formers undergoing a glass transition.

In the late 1990s it was observed that properties of stationary points of the potential energy function V , i.e. configuration space points q^s satisfying $dV(q^s) = 0$, reflect in dynamical and statistical physical quantities simultaneously, showing pronounced signatures near a phase transition [2]. This observation sparked quite some research activity, reviewed in [3], including a theorem asserting that stationary points with $V(q^s)/N = v_c$ are indispensable for the occurrence of an equilibrium phase transition at potential energy v_c [4]. Subsequently, it was noticed that the Hessian determinant of the potential energy function, evaluated at the stationary points, is crucial for discriminating whether or not the stationary points can induce a phase transition in the thermodynamic limit [5]. For some models, even in the absence of an exact solution, this insight may facilitate the exact analytic computation of transition energies [6].

In contrast to the applications of energy landscapes reviewed in [1], the study of phase transitions within this framework was pursued almost exclusively by analytical methods. An exception is the numerical work by Franzosi

et al. [7] that uses a rather indirect approach to study the phase transition of the ϕ^4 model on a two-dimensional square lattice with nearest-neighbor interactions.

In this Letter, we employ three different methods to extract information about the stationary points of the potential energy function of this model. The two-dimensional nearest-neighbor ϕ^4 model is known to have a continuous phase transition which is in the universality class of the two-dimensional Ising model but, in contrast to that model, is amenable to energy landscape methods by virtue of its continuous configuration space. Our first numerical method, previously used for analyzing gauge fixing functionals in lattice gauge theory [8], uses algebraic-geometrical techniques to obtain *all* stationary points of a given function of many variables, but is restricted to fairly small lattice sizes. The second method makes use of a globally convergent version of the Newton-Raphson algorithm for searching the zeros of a real-valued function (see Sec. 9.7 of [9]). The numerical results are corroborated by an analytical upper bound on the potential energy per lattice site $v^s = V(q^s)/N$ evaluated at a stationary point q^s . All three methods evidence that $v^s \leq 0$ for all q^s .

The implications of this finding are significant: First, it follows that the only previously published numerical method in the field [7] fails to approximate the exact result. Second, the above mentioned theorem by Franzosi and Pettini [4] is shown to be flawed: For suitable parameter values, the phase transition of the two-dimensional nearest-neighbor ϕ^4 model can take place at a critical potential energy v_c much larger than zero. Since the stationary values v^s are never larger than zero, this finding falsifies the theorem.

Two-dimensional nearest-neighbor ϕ^4 model.—On a finite square lattice $\Lambda \subset \mathbb{Z}^2$ consisting of $N = L^2$ sites, a

real degree of freedom ϕ_i is assigned to each lattice site $i \in \Lambda$. By $\mathcal{N}(i)$ we denote the subset of Λ consisting of the four nearest-neighboring sites of i on the lattice under the assumption of periodic boundary conditions. The potential energy function of this model is given by

$$V(q) = \sum_{i \in \Lambda} \left[\frac{\lambda}{4!} q_i^4 - \frac{\mu^2}{2} q_i^2 + \frac{J}{4} \sum_{j \in \mathcal{N}(i)} (q_i - q_j)^2 \right], \quad (1)$$

where $q = (q_1, \dots, q_N)$ denotes a point in configuration space $\Gamma = \mathbb{R}^N$ [10]. $J > 0$ determines the coupling strength between nearest-neighboring sites and the parameters $\lambda, \mu > 0$ characterize a local double-well potential each degree of freedom is experiencing.

In the thermodynamic limit $N \rightarrow \infty$ this model is known to undergo, at some critical temperature T_c , a continuous phase transition, in the sense that the configurational canonical free energy $f(T)$ is nonanalytic at $T = T_c$. The transition is from a “ferromagnetic” phase with nonzero average particle displacement to a “paramagnetic” phase with vanishing average displacement (see [11] for more details as well as for Monte Carlo results).

Instead of T_c , it is more adequate for our purposes to compare to the critical potential energy per lattice site, v_c , of the transition. Both quantities are unambiguously related to each other in the thermodynamic limit via the caloric curve $v(T)$, independently of the statistical ensemble used. The critical potential energy v_c is less frequently studied, in fact the only data we could find in the literature are from Monte Carlo simulations of fairly small system sizes $N = 20 \times 20$ in [12], with parameter values $\lambda = 3/5$, $\mu^2 = 2$, and $J = 1$. We mostly use the same parameter values in the following, and since the value of v_c is a crucial benchmark when relating our stationary point analysis to the phase transition of the ϕ^4 model, we have performed standard Metropolis Monte Carlo simulations for somewhat larger system sizes up to 128×128 and 10^7 lattice sweeps. We have sampled, among other observables, the canonical average $\langle v \rangle$ of the potential energy per particle, and plots of some of the Monte Carlo results are shown in Fig. 1. Although our simulations are not of high-end quality, the data are sufficiently good to extract a fair estimate of $v_c \approx 2.2$, establishing a critical potential energy well above zero.

Numerical continuation method.—For $J = 0$, i.e., in the absence of coupling, the stationary points q^s of the potential V in (1) can be calculated analytically without difficulty, obtaining the 3^N solutions $q^s = (q_1^s, \dots, q_N^s)$ with $q_j^s \in \{0, \pm\sqrt{6\mu^2/\lambda}\}$. Knowledge of these solutions permits us to continue them to $J > 0$ by numerical continuation; see [13] for a description of the homotopy continuation method we have actually been using. Under certain conditions on the initial (decoupled) and final (coupled) potentials, this method is known to yield all stationary points of V . However, since the number of stationary points for $J = 0$ grows exponentially with the




FIG. 1. (Color online) Monte Carlo results for the two-dimensional ϕ^4 model (1) with $\lambda = 3/5$, $\mu^2 = 2$, and $J = 1$. System sizes $N = L \times L$ are plotted with L ranging from 32 to 128. Left panel: the canonical average $\langle v \rangle$ of the potential energy per lattice site as a function of the temperature T . Right panel: The canonical specific heat $c = (\langle V^2 \rangle - \langle V \rangle^2)/(NT^2)$, plotted as a function of $\langle v \rangle$. The lines between the data points serve as a guide to the eye.

number of lattice sites, such an analysis is restricted to fairly small systems.

We have used the homotopy continuation method to compute all stationary points of V for various values of J and system sizes up to 4×4 , and the results show the following features. First, upon increasing the coupling constant J , the number $\#(q^s)$ of stationary points decreases gradually from 3^N for $J = 0$ to only 3 stationary points for some sufficiently large J . This behavior is illustrated for $N = 3 \times 3$ in the left panel of Fig. 2. Second, the stationary values $v^s = V(q^s)/N$, i.e., the potential energy per lattice site evaluated at a stationary point, are found to be never larger than zero. This is illustrated for $N = 4 \times 4$ in the right panel of Fig. 2. Before discussing the implications of these observations, we want to confirm and extend them by other methods.

Newton-Raphson method.—Variants of this method are frequently used in energy landscape studies [1]. From a given initial point in phase space, Newton-Raphson finds a stationary point that is in some sense close by [9]. From a large and well-chosen set of initial points (generated for example by Monte Carlo dynamics), a large




FIG. 2. (Color online) Numerical results from the homotopy continuation method. Left panel: The number of stationary points of V for $N = 3 \times 3$, plotted logarithmically as a function of the coupling J . Right panel: For all 10,288,973 stationary points q^s of a 4×4 lattice with $J = 0.2$, the scaled Hessian determinant $D = |\det \mathcal{H}_V(q^s)|^{1/N}$ is shown vs. the stationary value $v^s = V(q^s)/N$, demonstrating that $v^s \leq 0$ for all q^s .




FIG. 3. Numerical results from the Newton-Raphson method. For system sizes $N = L \times L$ with $L = 3, 4, 6, 8,$ and 16 , the scaled Hessian determinant $D = |\det \mathcal{H}_V(q^s)|^{1/N}$ is shown versus the stationary value $v^s = V(q^s)/N$. Up to 10^6 different stationary points q^s per system size have been computed for $J = 0.1$ (left panel) and $J = 0.2$ (right panel).

number of stationary points can be calculated. In contrast to the homotopy continuation method, there is in general no way of knowing whether all stationary points have been found. An advantage, however, is that the Newton-Raphson method can be applied to system sizes much larger than 4×4 .

We have used the routine `newt` from [9], a globally convergent version of the Newton-Raphson method, to compute stationary points of ϕ^4 lattices of sizes up to 32×32 . At least for small J , the number of stationary points of such larger systems is huge and only a tiny fraction of them can be tracked down. Similarly to the homotopy continuation results, the number of stationary points is found to decrease dramatically with increasing coupling J . For sufficiently large J , only three such points are found: the two global minima $q^s = (q_1^s, \dots, q_N^s)$ where all $q_j^s = \sqrt{6\mu^2/\lambda}$, respectively $-\sqrt{6\mu^2/\lambda}$, and a saddle of index 1 when $q_j^s = 0$ for all j . Moreover, we again observe that the stationary values obey $v^s \leq 0$ for all stationary points q^s found by the routine (see Fig. 3). Since in particular the latter observation has serious implications, we will corroborate the numerics by an analytic calculation.

Upper bound on the stationary values.—The stationary points of the potential V are the solutions of the N coupled nonlinear equations

$$\frac{\partial V(q)}{\partial q_k} = \frac{\lambda}{3!} q_k^3 - \mu^2 q_k + J \sum_{j \in \mathcal{N}(k)} (q_k - q_j) = 0. \quad (2)$$

By elementary algebraic manipulations this set of equations can be written in the form

$$J \sum_{j \in \mathcal{N}(k)} q_j = \frac{\lambda}{3!} q_k^3 + (4J - \mu^2) q_k. \quad (3)$$

Although it is not feasible to solve these equations explicitly, Eq. (3) can be used to determine the potential energy at a stationary point. To this aim, we rewrite the

potential (1) in the form

$$V(q) = \sum_{i \in \Lambda} q_i \left[\frac{\lambda}{4!} q_i^3 + \left(\frac{4J - \mu^2}{2} \right) q_i - \frac{J}{2} \sum_{j \in \mathcal{N}(i)} q_j \right]. \quad (4)$$

Substituting (3) into (4), we obtain the potential energy at a stationary point $q^s = (q_1^s, \dots, q_N^s)$,

$$V(q^s) = -\frac{\lambda}{4!} \sum_{i \in \Lambda} q_i^4. \quad (5)$$

Hence, for $\lambda \geq 0$, the potential energy per lattice site at any stationary point is bounded above by zero,

$$v^s = V(q^s)/N \leq 0. \quad (6)$$

In order to relate these findings to results in the literature, we next need to translate them into topology changes in configuration space.

Stationary points and configuration space topology.—Morse theory is a branch of mathematics relating the topology of some manifold M to the stationary points of a smooth function defined on M . For our purposes, we consider submanifolds of the configuration space Γ ,

$$M_v = \{q \in \Gamma \mid V(q) \leq Nv\}, \quad (7)$$

and relate their topology to the stationary points of the potential V . One of the basic theorems of Morse theory then amounts to the following statement [14]: If V has no stationary values $V(q^s)/N$ in the interval $[a, b]$, then M_a and M_b are topologically equivalent, i.e., there exists a homeomorphism mapping M_a onto M_b .

Franzosi and Pettini announced, and allegedly proved, a theorem stating that, under some conditions on the potential V , the following holds [4]: If there exists an interval $[a, b]$ such that, for any N larger than some constant N_0 , the manifolds M_v with $v \in [a, b]$ are homeomorphic, then in the thermodynamic limit the canonical configurational free energy $f(\beta)$ is two times continuously differentiable in the interval $(\beta(a), \beta(b))$, where $\beta(a)$ and $\beta(b)$ are the values of the inverse temperature corresponding to the potential energies a and b , respectively.

Translating the homeomorphicity condition into properties of stationary points, the theorem asserts that neither a first-order nor a second-order phase transition can occur at a critical potential energy v_c unless there are stationary values v^s converging to v_c in the thermodynamic limit. The nearest-neighbor ϕ^4 potential (1) satisfies all requirements on V demanded by this theorem. Accordingly, taking into account that all stationary values are nonpositive (6), the theorem asserts that the critical potential energy of the second-order phase transition of the model cannot be positive. This prediction is in contradiction to the thermodynamic properties of this model and the theorem in [4] is disproved.

Previous numerical results.—The only previous numerical study of the relation between configuration space topology and phase transitions was published by Franzosi *et al.* in 2000 [7]. Their method is very different from ours and rather intricate: Monte Carlo sampling is used to determine an estimator of the Gauss-Kronecker curvature K_G of the constant potential energy manifolds

$$\Sigma_v = \{q \in \Gamma \mid V(q) = Nv\} \quad (8)$$

in phase space. Then, K_G is related to the Euler characteristic $\chi(\Sigma_v)$ of these manifolds by means of the Gauss-Bonnet-Hopf theorem. Applying this method to the two-dimensional ϕ^4 model, the numerical results for $\chi(\Sigma_v)$ vary as a function of v and display a kink-like feature in the vicinity of the phase transition (see Fig. 3 of [7]).

The upper bound (6) on the stationary values v^s allows us to analytically determine the Euler characteristic $\chi(\Sigma_v)$: From the absence of stationary points with $v^s > 0$ we can infer that, for all $v > 0$, the manifold M_v as defined in (7) is homeomorphic to an N -dimensional ball. The constant potential energy manifold Σ_v is the boundary of M_v and is therefore homeomorphic to an $(N - 1)$ -sphere, implying that the Euler characteristic $\chi(\Sigma_v)$, being a topological invariant, is constant for all $v > 0$. The kink of $\chi(\Sigma_v)$ observed in [7] is therefore an artifact of the numerical method used.

More on the shape of the energy landscape.—Contrary to the claims in [4, 7], we have seen that, for large enough couplings J , the energy landscape in the vicinity of the critical potential energy v_c of a phase transition can be locally trivial, i.e., the configuration space submanifolds M_v can be homeomorphic for v slightly above and below v_c (and the same is true for the submanifolds Σ_v). For $v < 0$ and large system sizes, however, when stationary points abound, it is more difficult to explore the shape of the energy landscape. As a first step towards this aim, we study the properties of the stationary point $q^s = (0, \dots, 0)$. This point corresponds to the largest stationary value $v^s = 0$ at which M_v becomes homeomorphic to an N -dimensional ball. Its index I , i.e., the number of negative eigenvalues of the Hessian matrix of V evaluated at q^s , characterizes the topology change of M_v at $v = 0$. With increasing system size $N = L \times L$, we observe that the relative index $i = I/N$ converges to a finite value, and this value $i_\infty = \lim_{N \rightarrow \infty} i/N$ depends on the coupling J (see Fig. 4). The existence of such a limiting value might be taken as a starting point for discussing the topology of the potential energy landscape in the infinite system limit.

Conclusions.—By analytical and numerical methods, we have probed certain features of the potential energy landscape of the two-dimensional ϕ^4 lattice model. For large coupling J , the model's phase transition was found to occur at energies well separated from the stationary values of the potential V (or, equivalently, from topology changes in configuration space). These findings con-




FIG. 4. (Color online) Left panel: The relative index i of the stationary point $q^s = (0, \dots, 0)$, plotted as a function of the linear system size L for various couplings J . From top to bottom: $J = 0.1, 0.36, 0.5, 0.64, 1, 2, 10$. Right panel: The large-system limit i_∞ of the relative index, plotted as a function of the coupling constant J .

tradict, and thereby disprove, a theorem put forward in [4]. Since our results imply that the configuration space submanifolds M_v are simply connected for $v \geq 0$, the symmetry breaking phase transition appears to be driven only by a concentration-of-measure effect, but not by the connectivity of the underlying finite-system manifolds.

D.M. acknowledges support by the U.S. Department of Energy under contract DE-FG02-85ER40237 and by the Science Foundation Ireland grant 08/RFP/PHY1462. M.K. acknowledges support by the *Incentive Funding for Rated Researchers* programme of the National Research Foundation of South Africa.

* kastner@sun.ac.za

† dbmehta@syr.edu

- [1] D. J. Wales, *Energy Landscapes* (Cambridge University Press, Cambridge, 2004).
- [2] L. Casetti, M. Pettini, and E. G. D. Cohen, *Phys. Rep.*, **337**, 237 (2000).
- [3] M. Kastner, *Rev. Mod. Phys.*, **80**, 167 (2008).
- [4] R. Franzosi and M. Pettini, *Phys. Rev. Lett.*, **92**, 060601 (2004); R. Franzosi, M. Pettini, and L. Spinelli, *Nuclear Phys. B*, **782**, 189 (2007).
- [5] M. Kastner, S. Schreiber, and O. Schnetz, *Phys. Rev. Lett.*, **99**, 050601 (2007); M. Kastner and O. Schnetz, *ibid.*, **100**, 160601 (2008); M. Kastner, O. Schnetz, and S. Schreiber, *J. Stat. Mech. Theory Exp.*, **2008**, P04025 (2008).
- [6] C. Nardini and L. Casetti, *Phys. Rev. E*, **80**, 060103(R) (2009); M. Kastner, *ibid.*, **83**, 031114 (2011).
- [7] R. Franzosi, M. Pettini, and L. Spinelli, *Phys. Rev. Lett.*, **84**, 2774 (2000).
- [8] D. Mehta, A. Sternbeck, L. von Smekal, and A. G. Williams, *PoS, QCD-TNT09*, 025 (2009).
- [9] W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, *Numerical Recipes in C: The Art of Scientific Computing*, 2nd ed. (Cambridge University Press, Cambridge, 1992).
- [10] Our definition of V coincides with the one in [12], but differs from [7] by a factor 1/6 in the quartic term. Judging from the critical temperatures and energies reported in the latter, as well as from their reference to [12], we

assume that there is a misprint in [7]. Note that for the main conclusions of the present Letter the precise values of any of the constants do not matter.

- [11] A. Milchev, D. W. Heermann, and K. Binder, *J. Stat. Phys.*, **44**, 749 (1986).
- [12] R. Franzosi, L. Casetti, L. Spinelli, and M. Pettini, *Phys. Rev. E*, **60**, R5009 (1999).
- [13] D. Mehta, “Finding all the stationary points of a potential energy landscape via numerical polynomial homotopy continuation method,” arXiv:1104.5497.
- [14] Theorem 3.1 in Y. Matsumoto, *An Introduction to Morse Theory*, Translations of Mathematical Monographs, Vol. 208 (American Mathematical Society, 2002).