

Possibility of Skyrmion Superconductivity in Doped Antiferromagnet $K_2Fe_4Se_5$

G. Baskaran

Quantum Science Centre, The Institute of Mathematical Sciences, C I T Campus, Chennai 600 113, India
Perimeter Institute for Theoretical Physics, 31 Caroline St N, Waterloo, Ontario, Canada N2L 2Y5

Intercalated Fe chalcogenide $K_2Fe_4Se_5$ family exhibits high T_c (~ 30 K) superconductivity and spin-8 high T_N (~ 560 K) antiferromagnetism (AFM). We present a model Hamiltonian and suggest *Skyrmion superconductivity*. A doped electron creates an orbitally non degenerate $S=7\frac{1}{2}$ state in a Fe_4Se_9 cluster and moves in a single correlated band, exchange coupled to a robust $S=8$ AFM order. Skyrmion, a topological excitation of 2 dimensional AFM order acquires an *induced charge* $-2e$ through a quantum anomaly and becomes a Cooper pair. Superconductivity emerges for a range of doping. Fluctuating superconductivity, arising from preformed Cooper pairs (stable Skyrmions), around room temperatures is likely. There is hope for higher T_c 's in other high spin Mott insulators.

PACS numbers:

Discovery of superconductivity in Fe based pnictides [1, 2] and chalcogenides [3] have created another wave of excitement in condensed matter physics. Even though T_c has not matched the scale of 100 K seen in cuprates, Fe systems continue to offer surprises. A recent surprise is the discovery [4] of superconductivity in intercalated Fe chalcogenide family, where a robust antiferromagnetic order with a sublattice moment close to $13 \mu_B$ and a $T_N \approx 560$ K seems to coexist with a high T_c superconductivity of $T_c \approx 30$ K. From materials science point of view the vacancy order present in $K_2Fe_4Se_5$ has enabled a rich physics. Ordered vacancies is a new route to novel states of quantum matter [5].

In the present article we provide a minimal model and identify a mechanism of electron pairing. We use some general considerations and present certain consequences that are compelling. We show that a single band model of strongly correlated electrons, in a square lattice, close to half filling exchange coupled to a square lattice of large spin- $7\frac{1}{2}$ core moments of Fe_4 plaquettes captures the essential low energy physics. In our theory topological excitations of an ordered 2 dimensional Heisenberg antiferromagnet, namely Skyrmion [6], plays a fundamental role. Relevance of topological excitations [7, 8] to lightly doped quantum antiferromagnets began in 1987 with the advent of RVB theory of high T_c superconductivity. This study continues [10, 11].

Briefly, kinetic energy frustration of doped electrons in the presence of a rigid AFM order is relieved by a Skyrmion which help delocalize 2 electrons, using its spin twist. Skyrmion, through a quantum anomaly, acquires an *induces* charge $-2e$ and becomes a Cooper pair. For very low doping we get an ordered Skyrmion crystal. Beyond a range of doping Skyrmion crystal quantum melts, leading to *Skyrmion superconductivity*. For even higher dopings ferromagnetism is likely to intervene, with a possible island of p-wave superconductivity (figure 4).

Following the discovery of superconductivity in FeSe [3], a new family, A_xFe_ySe , with intercalated alkali atoms ($A = K, Cs, Rb$) was synthesized [4]. Stoichiometric




FIG. 1: $\sqrt{5} \times \sqrt{5}$ a) vacancy (o) order and $S=8$ ($=2 \times 4$) AFM order of Fe_4 plaquettes in FeSe layer of $K_2Fe_4Se_5$. Only Fe atoms (\bullet) are shown. Fe^{2+} is in $S = 2$, the maximal spin state. b) Se atoms above (red) and below (pink) Fe layers are shown. Dashed square in the middle marks a Fe_4Se_9 unit.

$K_2Fe_4Se_5$ has ordered Fe vacancies and $\sqrt{5} \times \sqrt{5}$ periodicity. Experiments indicate [12, 13] that $K_2Fe_4Se_5$ is a Mott insulator. Neutron scattering [12] shows that the Fe_4 plaquettes attain a large cluster moment and reaches a $S=8$ spin state. Further the plaquettes develop a (π, π) long range AFM order (figure 1) with a $T_N \sim 560$ K.

Superconducting T_c is around 30 K, in various members of the family. Many experiments show coexistence of magnetism and superconductivity [12, 13, 15]. However, NMR and STM results [14] seem to challenge coexistence.

There are some theoretical studies on this system [19]. We present a new approach and develop a minimal microscopic model to understand magnetic and superconducting properties. We first propose that Fe_4 *plaquette is a basic electronic building block. Electronic integrity of Fe_4 plaquette is revealed by its robust total spin-8 state arising from strongly exchange coupled four Fe atoms that carry spin-2 each. The $S=8$ plaquette spins have a (π, π) AFM order in the 2D square lattice of plaquettes. An important question is where do doped carriers go? This question is similar to Zhang-Rice singlet formation in cuprates. In view of available experiments we will focus on electron doping.*

Simple quantum chemical and phenomenological con-




FIG. 2: a) Schematic energy levels of 16 singly occupied and spin polarized plaquette orbitals of the Fe_4Se_9 cluster. b) The strongly hybridized core spin- $7\frac{1}{2}$ moment and the valence spin- $\frac{1}{2}$ moment distributed in the Fe_4 plaquette is schematically shown.

siderations suggest that a doped electron enters a Fe_4 plaquette and create an *orbitally non degenerate* spin- $7\frac{1}{2}$, *hexadectic state* in the Fe_4 plaquette [16]. Briefly, Fe atoms of Fe_4 plaquette are tetrahedrally coordinated by Se atoms and form a Fe_4Se_9 unit. In view of strong hybridization of 3d orbitals of Fe and 4s/4p orbitals of Se atoms, the 16 electrons that makeup $S=8$ spin state of Fe_4 plaquette reside in 16 *plaquette orbitals* in a strongly exchange coupled parallel spin state (figure 1b). A doped electron enters the top most singly occupied non degenerate orbital. We call this as *valence orbital* of Fe_4 plaquette that carries a *spin- $\frac{1}{2}$ valence moment*.

Anticipating addition of an electron (doping) we conveniently separate $S=8$ spin of Fe_4 into $S=7\frac{1}{2}$ core spin moment \mathbf{S}_i , ferromagnetically coupled by a *plaquette Hund coupling* J_H to a $S=\frac{1}{2}$ valence moment \mathbf{s}_i (figure 1b). Neighboring Fe_4 plaquette spins are antiferromagnetically coupled by an effective superexchange J . Thus the spin Hamiltonian of the Mott insulator is:

$$H_{\text{AFM}} = -J_H \sum_i \mathbf{S}_i \cdot \mathbf{s}_i + J \sum_{\langle ij \rangle} (\mathbf{S}_i + \mathbf{s}_i) \cdot (\mathbf{S}_j + \mathbf{s}_j) \quad (1)$$

$$= J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j - J_H \sum_i \mathbf{S}_i \cdot \mathbf{s}_i + \frac{J}{2} \sum_{\langle ij \rangle} (\mathbf{S}_i \cdot \mathbf{s}_j + \mathbf{S}_j \cdot \mathbf{s}_i)$$

Here $\mathbf{s}_i \equiv c_{i\alpha}^\dagger \vec{\sigma}_{\alpha\beta} c_{i\beta}$, is the valence spin- $\frac{1}{2}$ operator written in terms of electron operators ($c_{i\sigma}^\dagger, c_{i\sigma}$) of the Fe_4 plaquette valence orbital. For simplicity we ignore the weak c-axis exchange coupling and any anisotropic spin interaction terms.

In the limit $J_H \gg J$, equation 1 becomes a Spin-8 Heisenberg AFM Hamiltonian on a square lattice. The effective exchange coupling J can be estimated from the *acoustic spin wave* band width from inelastic neutron scattering results[12]. The estimate is $JS = 8J \approx 17$ meV. Location of the first *optic spin wave* branch give a value for $J_H \approx 110$ meV.

The Hamiltonian of our electron doped system is a single band model of plaquette valence spin electrons *close*




FIG. 3: Spin configuration of a Skyrmion. A circle in the middle has only xy component of sublattice spins and defines an XY-vortex. An anti Skyrmion will have an antivortex.

to *half filling*, coupled to $S = 7\frac{1}{2}$ core spins of Fe_4 plaquettes:

$$H = - \sum_{ij} t_{ij} c_{i\sigma}^\dagger c_{j\sigma} + \text{h.c.} + J \sum_{\langle ij \rangle} \mathbf{s}_i \cdot \mathbf{s}_j + J \sum_{\langle ij \rangle} \mathbf{S}_i \cdot \mathbf{S}_j$$

$$- J_H \sum_i \mathbf{S}_i \cdot \mathbf{s}_i + \frac{J}{2} \sum_{\langle ij \rangle} (\mathbf{S}_i \cdot \mathbf{s}_j + \mathbf{S}_j \cdot \mathbf{s}_i) \quad (2)$$

with a local constraint $\sum_\sigma c_{i\sigma}^\dagger c_{i\sigma} \neq 0$ (for electron doping). Here t is the inter plaquette hopping matrix element. From known band structure results, after folding the BZ due to vacancy ordering, we estimate $t \sim 50$ meV. The band width $W \sim 400$ meV is large compared to the charging energy $U_{\text{eff}} \approx \frac{e^2}{\epsilon_0 R} \sim 500$ meV of the $(\text{Fe}_2\text{Se}_5)^{2+}$ plaquette. Here R is the size of the Fe_4 plaquette. Our doping fraction x is with reference to the Fe_4 plaquette orbital band. Thus in the formula $\text{K}_{2+y}\text{Fe}_4\text{Se}_5$, our doping $x = \frac{y}{4}$.

The low energy effective Hamiltonian of the interacting core spin Heisenberg antiferromagnet is a $2 + 1$ dimensional $O(3)$ nonlinear sigma model with action:

$$S \sim \int dx dy d\tau [(\partial_\mu \mathbf{n}(\mathbf{r}))^2 - \frac{1}{v_s^2} (\partial_t \mathbf{n}(\mathbf{r}))^2] \quad (3)$$

Here $\mu = x, y$; v_s is the spin wave velocity and $\mathbf{n}(\mathbf{r})$ is the normalized, $\mathbf{n}(\mathbf{r}) \cdot \mathbf{n}(\mathbf{r}) = 1$, sublattice magnetization.

The above system supports small amplitude gapless spin wave (Goldstone mode) excitations and finite energy topological excitations namely Skyrmons. Skyrmons are analogue of a pair of domain walls of 1D Heisenberg antiferromagnet. In 1D system a single domain wall is a π -twist that interpolates up and down spin vacua in a continuous fashion. In the process it effectively decouples a single spin (a *spinon* localized around the domain wall) from the antiferromagnetic mean field. In the language of spin density wave (similar to polyacetylene) it creates a midgap state containing an unpaired spin.

A Skyrmion interpolates the down spin vacuum in its central region and up spin vacuum at the boundary, in a radially symmetric fashion (figure 3). In the process it effectively decouples two spin- $\frac{1}{2}$ *spinons* [8]. This can be

seen in several ways[7–9]. A simple way is in the SDW band picture, as studied by John and collaborators[9]. A skyrmion or 2 half skyrmions (2 merons) configuration traps two unpaired spins in two mid gap states.

Skyrmion has topological properties. It carries chirality or Pontryagin index Q ($= 0, \pm 1, \pm 2, \dots$). Its sign measures the sense, and magnitude number of windings, when we map order parameter $\mathbf{n}(\mathbf{r})$ field from the 2D plane onto S^2 , the surface of an ordinary sphere:

$$Q \equiv \frac{1}{8\pi} \int dx dy \mathbf{n} \cdot (\partial_x \mathbf{n} \times \partial_y \mathbf{n}) \quad (4)$$

In the language of homotopy theory, $\Pi_2(S^2) = \mathbb{Z}$. Skyrmion has a zero mode associated with a residual U(1) symmetry about the z-axis of the spin configuration. Quantized zero mode represents spin current fluctuation [17] and quantum Skyrmion is a *chiral doublet*. This is a 2D analogue of Affleck's dyon[18] in 1D Heisenberg antiferromagnets.

Valence electrons living in a single band in our model are exchange coupled to an antiferromagnetic mean field provided by the core spin moments from Fe_4 plaquettes. An electron in a rigid AFM background can not delocalize. We have kinetic energy frustration. Skyrmion ($Q = \pm 1$) relieves the frustration by delocalizing 2 electrons using spin current (spin twist) under its wings.

Taking care of the local constraints and exchange interactions in a mean field fashion we get the mean field Hamiltonian of the valence electrons system as:

$$H_{\text{mF}} \approx \sum_{ij} \tilde{t}_{ij} c_{i\sigma}^\dagger c_{j\sigma} + \text{h.c} - \tilde{J}_H S \sum_i \mathbf{n}_i \cdot c_{i\alpha}^\dagger \vec{\sigma}_{\alpha\beta} c_{i\beta} \quad (5)$$

Here \tilde{t}_{ij} is a renormalized interplaquette hopping matrix element, (we choose π flux mean field solution); $\tilde{J}_H = J_H + 4J$. The (π, π) antiferromagnetic order introduces a gap in the two Dirac nodes. To calculate induced electric charge, in the presence of a slowly varying order parameter $\mathbf{n}(\mathbf{r})$ and U(1) gauge field A_μ^c , we use the language of 2D Dirac fermions and follow Wigmann and Abanov[7]. The effective action of the valence electron quasi particles coupled to slowly varying background AFM field is:

$$S \sim \int d^3x \bar{\psi} [-i\gamma_\mu (\partial_\mu + A_\mu^c) + im \vec{\sigma} \cdot \mathbf{n}(\mathbf{r})] \psi \quad (6)$$

Here $\mu = x, y$ and $\tau, \bar{\psi} = i\psi^\dagger \gamma^0$, and $\gamma^0, \gamma^x, \gamma^y \equiv \sigma^z, \sigma^y, -\sigma^x$. We have rescaled the Fermi velocity of electron to 1 for brevity. Further $m \approx SJ_H$ is a *mass gap* provided by the strong exchange field. Integrating out the fermions and in the large mass approximation one gets [7]:

$$S = S_R + 2iA_\mu^c J_\mu^T \quad (7)$$

Here $J_\mu^T(\mathbf{r}) = \frac{1}{8\pi} \epsilon^{\mu\nu\lambda} \mathbf{n} \cdot \partial_\nu \mathbf{n} \times \partial_\lambda \mathbf{n}$ is the topological spin current density, whose time component $J_0^T(\mathbf{r}) \equiv$

$\frac{1}{8\pi} \mathbf{n} \cdot (\partial_x \mathbf{n} \times \partial_y \mathbf{n})$ is the Skyrmion density (equation 4). The real part of the induced interaction renormalizes the NLS action (equation 3). The imaginary part is the key U(1) gauge field A_μ^c - topological spin current J_μ^T coupling. The time component of the above, $2iA_0^c J_0^T$ is the coupling of dopant chemical potential to Skyrmion density (equation 4). Hence we get an induced U(1) (electric) charge Q_{Skyrmion}^c given by the expression[7]:

$$Q_{\text{Skyrmion}}^c = \pm 2e |Q| \quad (8)$$

where Q is the Pontryagin index or Skyrmion number.

To calculate the spin of the $-2e$ Skyrmion, we start with an even number of electrons in an ordered antiferromagnet and create one Skyrmion. This generates two singly occupied mid gap states. Addition of two electrons to these midgap states completely fill the two states and create spin zero midgap states. Thus nominally a 2D Skyrmion carrying charge $-2e$ is a spin singlet.

$2e$ Skyrmion, a charged chiral doublet, as discussed by Lee [11] for the case of a spin liquid, represents a $L_z = \pm 2$ angular momentum, $d \pm id$ state of a Cooper pair. Skyrmions that are real combination of the above doublet will lead to a *real* d-wave superconducting state. Otherwise we will get a $d \pm id$ superconducting state that breaks PT symmetry. This will be a chiral superconductor *à la* Laughlin.

In an ideal O(3) non linear sigma model in 2D, Skyrmion energy is finite and size independent. In practice, coulomb repulsion among 2 electrons, spin anisotropy and interchain coupling determine the size. For our case, the mean size of Skyrmion is determined by the inter Skyrmion distance. Cooper pair size or coherence length is selfconsistently limited by mean inter Cooper pair distance.

Skyrmion delocalization and kinetic energy gain is determined by their mutual overlap, proportional to x . As a first approximation, Skyrmion band width is a fraction x of the fermi energy Wx of free dopants. That is, the Skyrmion band width $W_{\text{Skyrmion}} \sim Wx^2$. Since $W \sim 400$ meV, $Wx^2 \sim 4$ meV, for a doping of $x \sim 0.1$. As $-2e$ Skyrmions are Bosons this is also Bose condensation energy scale. This energy is in the right ball park of the observed $T_c \approx 30$ K. Our superconductor is best viewed as a 2D Kosterlitz-Thouless condensed state of interacting bosons. Weak Josephson coupling between layers will make it a 3D superconductor.

Skyrmions are stable, in principle upto Neel temperature, as long as AFM order survives. In this sense $2e$ Skyrmions are *performed Cooper pairs* in the background of an AFM order. As a consequence we could see Cooper pairs well above T_c , through for example anomalous Nernst effect. We predict that $\text{K}_2\text{Fe}_4\text{Se}_5$ family offers a wonderful and some what unique opportunity to explore and utilise fluctuating superconductivity around room temperature scales.




FIG. 4: Schematic phase diagram for doped $K_2Fe_4Se_5$ family.

In our theory the large spin AFM order and superconductivity coexist for a range of doping. Dopant delocalization will eventually destroy the long range AFM order. In the low doping regime unscreened coulomb interaction will localize Skyrmions resulting in Skyrmion crystal or Skyrmion glass. Berciú and John [9] have discussed this for cuprates. A universal resistivity maxima seen in $K_2Fe_4Se_5$ superconductor could indicate onset of Skyrmion crystallization frustrated by growing quantum fluctuations (dashed line in figure 4 inside AFM dome).

It is known that large spin doped antiferromagnets leads to ferromagnetism through *double exchange*, for example in doped manganites, a spin-2 Mott antiferromagnet. As shown by de Gennes, small doping leads to spin twist (canting), before ferromagnetic order sets in at higher dopings. In a 2 dimensional system such as ours, the system naturally organize spin twists in the form of topologically stable Skyrmion eigen excitations and help delocalization. What is exciting is that *superconductivity is an alternative to ferromagnetism in 2 dimensions*. So we have a new avenue for high T_c search, namely high spin antiferromagnets in lower dimensions.

It also follows from double exchange physics that heavily doped $K_2Fe_4Se_5$ will eventually become a ferromagnetic metal. If we take clue from Sr_2RuO_4 , another 2D even electron ($Ru^{2+} = 4d^4$) system, Hund coupling and a growing ferromagnetic fluctuation could stabilise an island of spin triplet p-wave superconductivity.

A robust AFM magnetic order of a large spin Mott antiferromagnet offers exciting possibilities, as shown schematically in figure 4. It will be useful to study doped $K_2Fe_4Se_5$ family experimentally with controlled doping.

I thank D. Abanin, P.W. Anderson, T. Imai, D.H. Lee, N.P. Ong and T. Xiang for discussion. This research was supported by Perimeter Institute for Theoretical Physics.

[1] I.Y. Kamihara et al., J. Am. Chem. Soc. **130**, 3296 (2008)
 [2] Cao Wang et al., Europhys. Lett., **83** 67006 (2008); Han et al., Phys. Rev. **B 80** 024506 (2009)
 [3] F. C. Hsu et al., PNAS (USA), **105** 14262 (2008)
 [4] J. Guo et al., Phys. Rev. **B 82** 180520(R) (2010); I thank T. Xiang for bringing this & related work to my attention.

[5] M. Alaei, S. Akbar Jafari and H. Akbarzadeh, J. Phys. Chem. Sol. **69** 3283 (2008)
 [6] A.A. Belavin and A.M. Polyakov, JETP Lett., **22** 245 (75); W.J. Zakrzewski, Low Dimensional Sigma Models, Adam Hilger (NY, 1990); F. Wilczek and A. Zee, Phys. Rev. Lett., **51** 2250 (83); D.J. Gross, Nucl. Phys. **B 132** 439 (1978); F. D. M. Haldane, Phys. Rev. Lett. **61** 1029 (1988);
 [7] I. Dzyaloshinskii, A. Polyakov and P. Wiegmann, Phys. Lett., **127** 112 (1988); P.B. Wiegmann, Phys. Rev. **B 59** 15705 (1999); A.G. Abanov and P.B. Wiegmann, Phys. Rev. Lett. **86** 1319 (2001)
 [8] P.W. Anderson, S. John, G. Baskaran, B. Doucot and S. Liang, Princeton Preprint (1988); G. Baskaran, Phys. Rev. **B 68** 21409 (2003)
 [9] S. John and A. Golubnetsev, Phys. Rev. Lett., **71** 3343 (1993); M. Berciú and S. John, Phys. Rev. **B 59** 15443 (1999); *ibid*, **B 61** 16454 (2000)
 [10] X.G. Wen and A. Zee, Nucl. Phys. **B 15** 135 (1990); P. Voruganti and S. Doniach, Phys. Rev. **B 41** 9358 (1990); B.I. Shraiman and E.D. Siggia, Phys. Rev. **B 42** 2485 (1990); J. A. Vergas et al., Phys. Rev. **B 43**, 6099 1991; R. J. Gooding, Phys. Rev. Lett., **66** 2266 (1991); E. C. Marino and M. B. Silva Neto, Phys. Rev. **B 64** 092511 (2001); T. Morinari, Phys. Rev., **B 73** 064504 (2006); D. H. Lee and C. L. Kane, Phys. Rev. Lett. **64** 1313 (1990); S. L. Sondhi et al., Phys. Rev. **B 47** 16419 (1993); T. Grover and T. Senthil, Phys. Rev. Lett. **100** 156804 (2008); D. A. Abanin, A. Parameswaran, and S. L. Sondhi, Phys. Rev. Lett. **103** 076802 (2009)
 [11] D.H. Lee, Phys. Rev. **B 60**, 12429 (1999); mid gap states in a spin liquid state and *pairing via index theorem*, discussed in this reference is deeply related to 2e Skyrmion in an ordered AFM. I thank D.H. Lee for a reminder.
 [12] Wei Bao et al., Chinese Phys. Lett. **28**, 086104 (2011); M. Wang et al., arXiv:1105.4675; J.T. Park et al., arXiv:1107.1703
 [13] M. Fang et al., EPL, **94** (2011) 27009; Z.G. Chen et al., Phys. Rev. **B83**, 220507(R) (2011)
 [14] D.A. Torchetti et al., Phys. Rev., **B83** 104508 (2011); L. Ma, Phys. Rev. **B 83** 174510 (2011); W.Q. Yu et al., Phys. Rev. Lett. **106** 197001 (2011); Wei Li et al., arXiv:1108.0069; Peng Cai et al., arXiv:1108.2798
 [15] R.H. Liu et al., EPL, **94**, 27008 (2011); Z. Shermadini et al., Phys. Rev. Lett. **106** 117602 (2011); A.M. Zhang et al., arXiv:1106.2706; D.H. Ryan, Phys. Rev. **B 83** 104526 (2011); Z. Li et al., arXiv:1103.0098; F. Ye et al., arXiv:1102.2882; D. Mou et al., Phys. Rev. Lett., **106** 107001 (2011)
 [16] P.W. Anderson suggests (private communications) that on hole doping, a highly stable $S = \frac{5}{2}$ state of Fe^{3+} could stabilize a $S = 8 \frac{1}{2}$ (*octadectic*) Fe_4 plaquette spin.
 [17] G. Baskaran, unpublished
 [18] I. Affleck, Phys. Rev. Lett., **57** 1048 (1986)
 [19] R. Yu, Jian-Xin Zhu and Q. Si, Phys. Rev. Lett., **106** 186401 (2011); G.M. Zhang, Z.Y. Lu and T. Xiang, arXiv:1102.4575; W. Lv et al., arXiv:1105.0432; Y. Zhou, arXiv:1101.4462; C. Cao and J. Dai, arXiv:1102.1344; Fa Wang et al., Euro. Phys. Lett., **93** 57003; (2011); Y. Zhou et al., arXiv:1101.4462; ; T. Das and A.V. Balatsky, arXiv:1106.3289; T.A. Maier, Phys. Rev., **B 83** 100515(R) (2011); Yi-Zhuang Wu et al. arXiv:1103.1902; C. Platt et al., arXiv:1106.5964