

Ordinarization Transform of a Numerical Semigroup and Semigroups with a Large Number of Intervals

Maria Bras-Amorós

February 19, 2022

Abstract

A numerical semigroup is said to be ordinary if it has all its gaps in a row. Indeed, it contains zero and all integers from a given positive one. One can define a simple operation on a non-ordinary semigroup, which we call here the ordinarization transform, by removing its smallest non-zero non-gap (the multiplicity) and adding its largest gap (the Frobenius number). This gives another numerical semigroup and by repeating this transform several times we end up with an ordinary semigroup. The genus, that is, the number of gaps, is kept constant in all the transforms.

This procedure allows the construction of a tree for each given genus containing all semigroups of that genus and rooted in the unique ordinary semigroup of that genus. We study here the regularity of these trees and the number of semigroups at each depth. For some depths it is proved that the number of semigroups increases with the genus and it is conjectured that this happens at all given depths. This may give some lights to a former conjecture saying that the number of semigroups of a given genus increases with the genus.

We finally give an identification between semigroups at a given depth in the ordinarization tree and semigroups with a given (large) number of gap intervals and we give an explicit characterization of those semigroups.

Keywords: Numerical semigroup, semigroup tree, Freĭman's theorem, sums.

1 Introduction

Let \mathbb{N}_0 denote the set of non-negative integers. A numerical semigroup is a subset of \mathbb{N}_0 which is closed under addition, contains 0, and its complement in \mathbb{N}_0 is finite. One main reference for numerical semigroups is [14]. For a numerical semigroup Λ the elements in $\mathbb{N}_0 \setminus \Lambda$ are called *gaps* and the number of gaps is the *genus* of the semigroup. The largest gap is called the *Frobenius*

number. The *multiplicity* m of a numerical semigroup is its first non-zero non-gap. A numerical semigroup different than \mathbb{N}_0 is said to be *ordinary* if its gaps are all in a row.

It was conjectured in [2] that the number n_g of numerical semigroups of genus g asymptotically behaves like the Fibonacci numbers. More precisely, it was conjectured that $n_g \geq n_{g-1} + n_{g-2}$, that the limit of the ratio $\frac{n_g}{n_{g-1} + n_{g-2}}$ is 1 and so that the limit of the ratio $\frac{n_g}{n_{g-1}}$ is the golden ratio $\phi = \frac{1+\sqrt{5}}{2}$. Many other papers deal with the sequence n_g [10, 11, 3, 5, 4, 6, 16, 1, 9] and recently Alex Zhai gave a proof for the asymptotic Fibonacci-like behavior of n_g [15]. However, it has still not been proved that n_g is increasing. In the present work we approach this problem.

All numerical semigroups can be organized in a tree \mathcal{T} whose root is the semigroup \mathbb{N}_0 and in which the parent of a numerical semigroup Λ of genus g is the numerical semigroup obtained by adjoining to Λ its Frobenius number. So, the parent of a numerical semigroup of genus g has genus $g - 1$ and all numerical semigroups are in \mathcal{T} , at a depth equal to its genus. In particular, n_g is the number of nodes of \mathcal{T} at depth g . This construction was already considered in [13].

Here we will see that all numerical semigroups of a given genus g can be organized in a tree \mathcal{T}_g rooted at the unique ordinary semigroup of genus g . One significant difference between \mathcal{T}_g and \mathcal{T} is that the first one has only a finite number of nodes, indeed, it has n_g nodes, while \mathcal{T} is an infinite tree. We will see some relations between the trees \mathcal{T}_g and \mathcal{T} . We conjecture that the number of numerical semigroups in \mathcal{T}_g at a given depth is at most the number of numerical semigroups in \mathcal{T}_{g+1} at the same depth. This conjecture would prove that $n_{g+1} \geq n_g$. Here we show this result for the lowest and largest depths.

We finally study semigroups with a large number of intervals. We prove that if $\lfloor \frac{n}{2} \rfloor \geq \frac{g+2}{3}$, then the set of numerical semigroups of genus g and n intervals of gaps is empty if n and g have different parity and it is exactly the set of numerical semigroups of genus g and depth $\lfloor \frac{n}{2} \rfloor$ in \mathcal{T}_g otherwise. Furthermore we give an explicit description of the form of these semigroups.

2 A tree with all semigroups of a given genus

2.1 Ordinarization transform and ordinarization number of a numerical semigroup

The *ordinarization transform* of a non-ordinary semigroup Λ with Frobenius number F and multiplicity m is the set $\Lambda' = \Lambda \setminus \{m\} \cup \{F\}$. The ordinarization transform of an ordinary semigroup is itself. For instance, the ordinarization transform of the semigroup $\Lambda = \{0, 4, 5, 8, 9, 10, 12, \dots\}$ is the semigroup $\Lambda' = \{0, 5, 8, 9, 10, 11, 12, \dots\}$. Note that the genus of the ordinarization transform of a semigroup is the genus of the semigroup.

We can iterate the ordinarization transform and set $\Lambda'' = (\Lambda')'$ and in gen-

eral $\Lambda^{(i)} = \Lambda^{(i-1)'}$. It is easy to check that if Λ has genus g then there exists an integer i such that $\Lambda^{(i)} = \{0, g+1, g+2, \dots\}$. If Λ is non-ordinary then there exists a unique i with this property and satisfying $\Lambda^{(i-1)} \neq \{0, g+1, g+2, \dots\}$. We call this i the *ordinarization number* of Λ . By extension, the ordinarization number of an ordinary semigroup is set to be 0. For instance, if $\Lambda = \{0, 4, 5, 8, 9, 10, 12, \dots\}$ then $\Lambda' = \{0, 5, 8, 9, 10, 11, 12, \dots\}$ which is not the ordinary semigroup while $\Lambda'' = \{0, 7, 8, 9, 10, 11, 12, \dots\}$ which is the ordinary semigroup of genus 6. Thus the ordinarization number of Λ is 2.

Lemma 1. *The ordinarization number of a numerical semigroup of genus g is the number of its non-zero non-gaps which are smaller than or equal to g .*

Proof. A numerical semigroup of genus g is non-ordinary if and only if its multiplicity is at most g . So, we can transform a numerical semigroup while its multiplicity is at most g . The number of times that we can transform a semigroup before getting the ordinary semigroup is thus the number of its non-zero non-gaps which are smaller than or equal to the genus. \square

Given a numerical semigroup Λ it will be convenient to consider its enumeration λ as the unique increasing bijective map between \mathbb{N}_0 and Λ . We will use λ_i for $\lambda(i)$. By the previous lemma, if the ordinarization number of a semigroup is r then the non-gaps which are at most g are $\lambda_0 = 0, \lambda_1, \dots, \lambda_r$.

Lemma 2. *The maximum ordinarization number of a semigroup of genus g is $\lfloor \frac{g}{2} \rfloor$.*

Proof. Suppose that the ordinarization number of a semigroup is r . On one hand, since the Frobenius number F is at most $2g-1$, the total number of gaps from 1 to $2g-1$ is g and so the number of non-gaps from 1 to $2g$ is g . The number of those non-gaps which are larger than the genus is $g-r$. On the other hand $\lambda_r + \lambda_1, \lambda_r + \lambda_2, \dots, 2\lambda_r$ are different non-gaps between $g+1$ and $2g$. So, the number of non-gaps between $g+1$ and $2g$ is at least r . Putting this altogether we get that $g-r \geq r$ and so $r \leq \frac{g}{2}$.

On the other hand, the bound is attained by the semigroup

$$\{0, 2, 4, \dots, 2 \lfloor \frac{g}{2} \rfloor, 2 \left(\lfloor \frac{g}{2} \rfloor + 1 \right), \dots, 2g, 2g+1, 2g+2, \dots\}. \quad (1)$$

\square

We want to see that the maximum ordinarization number as stated in the previous lemma is attained only by the semigroup in (1). For this purpose we need the next lemma. Its proof has been omitted but can easily be obtained.

Lemma 3. *Consider a finite subset $A = \{a_1 < \dots < a_n\} \subseteq \mathbb{N}_0$.*

1. *The set $A + A$ contains at least $2n - 1$ elements*
2. *The set $A + A$ contains exactly $2n - 1$ elements if and only if $a_i = a_1 + \alpha i$ for all i and for a given positive integer α .*

Lemma 4. *Let $g > 0$. The unique numerical semigroup of genus g and ordinarization number $\lfloor \frac{g}{2} \rfloor$ is $\{0, 2, 4, \dots, 2g, 2g+1, 2g+2, \dots\}$.*

Proof. Suppose that the ordinarization number of Λ is $\lfloor \frac{g}{2} \rfloor$. Since $\lambda_{\lfloor \frac{g}{2} \rfloor} \leq g$, we know that the set of all non-gaps between 0 and $2g$ must contain all the sums $\{\lambda_i + \lambda_j : 0 \leq i, j \leq \lfloor \frac{g}{2} \rfloor\}$. But the number of non-gaps between 0 and $2g$ is $g+1$, while by Lemma 3 the set of sums above has at least $2\lfloor \frac{g}{2} \rfloor + 1$ elements.

If g is even then by the second item of Lemma 3 we get that $\lambda_i = i\lambda_1$ for $i \leq \frac{g}{2}$. Now $\lambda_{\frac{g}{2}} = \frac{g}{2}\lambda_1 \leq g$ means that $\lambda_1 \leq 2$. If $\lambda_1 = 1$ this contradicts $g > 0$. So, $\lambda_i = 2i$ for $0 \leq i \leq \frac{g}{2}$ and the remaining non-gaps between $g+1$ and $2g$ are necessarily $\lambda_i = 2i$ for $i = \frac{g}{2} + 1$ to $i = g$.

If g is odd and $g \notin \Lambda$ then we know that the set of all non-gaps between 0 and $2g-1$ must contain all the sums $\{\lambda_i + \lambda_j : 0 \leq i, j \leq \lfloor \frac{g}{2} \rfloor\}$. But the number of non-gaps between 0 and $2g-1$ is g , while by Lemma 3 the set of sums above has at least $2\lfloor \frac{g}{2} \rfloor + 1 = g$ elements and we can argue as before. \square

2.2 The tree \mathcal{T}_g

The definition of the ordinarization transform of a numerical semigroup allows the construction of a tree \mathcal{T}_g on the set of all numerical semigroups of a given genus rooted at the unique ordinary semigroup of this genus, where the parent of a semigroup is its ordinarization transform and the descendants of a semigroup are the semigroups obtained by taking away a generator larger than the Frobenius number and adding a new non-gap smaller than the multiplicity in a licit place. To illustrate this construction with an example in Figure 1 we depicted \mathcal{T}_6 . The depth of a numerical semigroup of genus g in \mathcal{T}_g is then its ordinarization number.

The next two lemmas show some relations between \mathcal{T} and \mathcal{T}_g .

Lemma 5. *If Λ_1 is a descendant of Λ_2 in \mathcal{T} then Λ'_1 is a descendant of Λ'_2 in \mathcal{T} .*

Proof. This is obvious when Λ_1 (and so Λ_2) is an ordinary semigroup. Suppose that Λ_1 is not ordinary. It is easy to check that in this case Λ_1 and Λ_2 have the same multiplicity m and that $\Lambda_1 \setminus \{m\}$ is a descendant of $\Lambda_2 \setminus \{m\}$ in \mathcal{T} . Now the lemma is a consequence of the fact that if we adjoin to a semigroup (in our case $\Lambda_1 \setminus \{m\}$) its Frobenius number we obtain its parent in \mathcal{T} (in our case $\Lambda_2 \setminus \{m\} = \Lambda'_1$) and if we repeat the same procedure (in our case obtaining Λ'_2) we obtain the parent of the parent in \mathcal{T} (in our case the parent of Λ'_1 in \mathcal{T}). \square

Lemma 6. *If two non-ordinary semigroups Λ_1 and Λ_2 with the same genus g have the same parent in \mathcal{T} then they also have the same parent in \mathcal{T}_g .*

Proof. Indeed, the parent of Λ_1 and Λ_2 in \mathcal{T}_g is the parent they have in \mathcal{T} without its multiplicity. \square




Figure 1: \mathcal{T}_6

2.3 Conjecture

Let $n_{g,r}$ be the number of numerical semigroups of genus g and ordinarization number r . For each genus $g \leq 49$ we computed $n_{g,r}$ for each ordinarization number r from 0 up to $\lfloor \frac{g}{2} \rfloor$. The results are given in Table 1. One can observe that for each ordinarization number the number of semigroups of this ordinarization number increases with the genus or stays the same. By extending the definition of $n_{g,r}$ for $r > \lfloor \frac{g}{2} \rfloor$ by setting $n_{g,r} = 0$ in this case, this leads to the next conjecture.

Conjecture 7. *For each genus $g \in \mathbb{N}_0$ and each ordinarization number $r \in \mathbb{N}_0$, $n_{g,r} \leq n_{g+1,r}$.*

This is equivalent to the number of numerical semigroups at a given depth of \mathcal{T}_g being at most the number of numerical semigroups at the same depth of \mathcal{T}_{g+1} . If the conjecture were true then the total number of nodes in \mathcal{T}_g would be at most the total number of nodes in \mathcal{T}_{g+1} proving that n_g increases with g .

3 Partial proofs of the conjecture

We will prove the conjecture for particular values of the pair g, r . We wrote these values in bold face in Table 1. It is obvious that for $r = 0$ we always have $n_{g,r} = 1$ since for any genus the ordinary semigroup is the unique numerical semigroup of ordinarization number 0. In the next subsections we will prove the conjecture for $n_{g,1}$ and any g and for $n_{g,r}$ and any g , whenever $r \geq \max(\frac{g}{3} + 1, \lfloor \frac{g+1}{2} \rfloor - 14)$.

3.1 Ordinarization number 1

Lemma 8. *Let $g \in \mathbb{N}_0$. The number of semigroups of genus g and ordinarization number 1 is*

$$n_{g,1} = \left\lceil \frac{g-1}{2} \right\rceil \left\lfloor \frac{g+1}{2} \right\rfloor + \frac{\lfloor \frac{g-1}{2} \rfloor \lfloor \frac{g+1}{2} \rfloor}{2} = \begin{cases} \frac{3}{8}g^2 - \frac{1}{4}g & \text{if } g \text{ is even,} \\ \frac{3}{8}g^2 - \frac{3}{8} & \text{if } g \text{ is odd.} \end{cases}$$

Proof. The semigroups of ordinarization number one are obtained from the ordinary semigroup $\{0, g+1, g+2, \dots\}$ by taking out one non-gap a and adding a non-zero non-gap b smaller than $g+1$. Since any element in the ordinary semigroup larger than $2g+1$ is a sum of two non-zero non-gaps it can not be taken out. So,

$$g+1 \leq a \leq 2g+1.$$

Fix a in the previous range. For b we have four necessary conditions which together become sufficient:

1. $b \leq g$ since b must be a gap of the ordinary semigroup;
2. $a-b \leq g$ since otherwise $a = b + (a-b)$ and so a must be in the new semigroup;

3. $2b \geq g + 1$ because otherwise $b, 2b$ are two different non-gaps which are at most g contradicting that the ordinaryization number of the new semigroup is 1;
4. $2b \neq a$ because otherwise a must be a non-gap.

From the first three conditions we deduce

$$\max\left(a - g, \left\lceil \frac{g+1}{2} \right\rceil\right) \leq b \leq g.$$

Now, taking also the fourth condition into consideration we get that the number of options for the pair a, b (and so $n_{g,1}$) is

$$\begin{aligned} n_{g,1} &= \sum_{a=g+1}^{2g+1} (g+1 - \max(a-g, \left\lceil \frac{g+1}{2} \right\rceil)) - \#\{\text{even integers in } \{g+1, \dots, 2g+1\}\} \\ &= \sum_{a=g+1}^{\left\lceil \frac{g+1}{2} \right\rceil + g} (g+1 - \left\lceil \frac{g+1}{2} \right\rceil) + \sum_{\left\lceil \frac{g+1}{2} \right\rceil + g+1}^{2g+1} (g+1 - a + g) - \left\lfloor \frac{g+1}{2} \right\rfloor \\ &= \sum_{a=g+1}^{\left\lceil \frac{g+1}{2} \right\rceil + g} \left\lfloor \frac{g+1}{2} \right\rfloor + \sum_{\left\lceil \frac{g+1}{2} \right\rceil + g+1}^{2g+1} (2g+1 - a) - \left\lfloor \frac{g+1}{2} \right\rfloor \\ &= \left\lfloor \frac{g+1}{2} \right\rfloor \left\lfloor \frac{g+1}{2} \right\rfloor + \sum_{i=0}^{\left\lfloor \frac{g-1}{2} \right\rfloor} i - \left\lfloor \frac{g+1}{2} \right\rfloor \\ &= \left\lfloor \frac{g-1}{2} \right\rfloor \left\lfloor \frac{g+1}{2} \right\rfloor + \frac{\left\lfloor \frac{g-1}{2} \right\rfloor \left\lfloor \frac{g+1}{2} \right\rfloor}{2}. \end{aligned}$$

□

Corollary 9. For each genus $g \in \mathbb{N}_0$, $n_{g,1} \leq n_{g+1,1}$.

Proof. If g is even and $g+1$ is odd then

$$\begin{aligned} n_{g,1} &= \frac{3}{8}g^2 - \frac{1}{4}g \\ n_{g+1,1} &= \frac{3}{8}(g+1)^2 - \frac{3}{8} = \frac{3}{8}g^2 + \frac{3}{4}g \end{aligned}$$

So, $n_{g+1,1} = n_{g,1} + g \geq n_{g,1}$.

On the other hand, if g is odd and $g+1$ is even then

$$\begin{aligned} n_{g,1} &= \frac{3}{8}g^2 - \frac{3}{8} \\ n_{g+1,1} &= \frac{3}{8}(g+1)^2 - \frac{1}{4}(g+1) = \frac{3}{8}g^2 + \frac{g}{2} + \frac{1}{8} \end{aligned}$$

So, $n_{g+1,1} = n_{g,1} + \frac{g+1}{2} \geq n_{g,1}$.

□

3.2 High ordinarization numbers

Next we will need Freïman's Theorem [7, 8] as formulated in [12].

Theorem 10 (Freïman). *Let A be a set of integers such that $\#A = k \geq 3$. If $\#(A + A) \leq 3k - 4$, then A is a subset of an arithmetic progression of length $\#(A + A) - k + 1 \leq 2k - 3$.*

By means of Freïman's Theorem we can prove the next lemma which tells that the semigroups of large ordinarization number have the first non-gaps even.

Lemma 11. *If a semigroup Λ of genus g has ordinarization number r with $\frac{g+2}{3} \leq r \leq \lfloor \frac{g}{2} \rfloor$ then all its non-gaps which are less than or equal to g are even.*

Proof. Suppose that Λ is a semigroup of genus g and ordinarization number $r \geq \frac{g+2}{3}$. This means in particular that $\lambda_0 = 0, \lambda_1, \dots, \lambda_r \leq g$ and $\lambda_{r+1} \geq g + 1$.

Let $A = \Lambda \cap [0, g] = \{\lambda_0, \lambda_1, \dots, \lambda_r\}$. We have that the elements in $A + A$ are upper bounded by $2g$ and so $A + A \subseteq \Lambda \cap [0, 2g]$. Then $\#(A + A) \leq \#(\Lambda \cap [0, 2g])$. Since the Frobenius number of Λ is at most $2g - 1$, $\#(\Lambda \cap [0, 2g]) = 2g + 1 - g = g + 1$. So, $\#A = r + 1$ while $\#(A + A) \leq g + 1$. Now, since $r \geq \frac{g+2}{3}$ we have $g + 1 \leq 3r - 1 = 3(r + 1) - 4$ and we can apply Theorem 10 with $k = r + 1$. Thus we have that A is a subset of an arithmetic progression of length at most $g + 1 - k + 1 = g - r + 1$. Let $d(A)$ be the common difference of this arithmetic progression.

Now, $d(A)$ can not be larger than or equal to three since otherwise $\lambda_r \geq r \cdot d(A) \geq 3r \geq 3 \frac{g+2}{3} > g$, a contradiction with r being the ordinarization number.

If $d(A) = 1$ then $A \subseteq [0, g - r]$ and we claim that in this case $A \subseteq \{0\} \cup [\lceil \frac{g+1}{2} \rceil, g - r]$. Indeed, suppose that $x \in A$. Then $2x$ satisfies either $2x \leq g - r$ or $2x \geq g + 1$. If the second inequality is satisfied then it is obvious that $x \in \{0\} \cup [\lceil \frac{g+1}{2} \rceil, g - r]$. If the first inequality is satisfied then we will prove that $mx \leq g - r$ for all $m \geq 2$ by induction on m and this leads to $x = 0$. Indeed, if $mx \leq g - r$ then $x \leq \frac{g-r}{m} \leq \frac{g-\frac{g+2}{3}}{m} = \frac{2g-2}{3m} < \frac{2g}{3m}$. Now $(m+1)x < \frac{2g(m+1)}{3m} = \frac{(2m+2)g}{3m}$ and since $m \geq 2$, we have $(m+1)x < \frac{(2m+m)g}{3m} = g$. Since $(m+1)x$ is in $\Lambda \cap [0, g] = A \subseteq [0, g - r]$ this means that $(m+1)x \leq g - r$ and this proves the claim.

Now, $A \subseteq \{0\} \cup [\lceil \frac{g+1}{2} \rceil, g - r]$ implies that $r \leq g - r - \lceil \frac{g+1}{2} \rceil + 1 = \lfloor \frac{g+1}{2} \rfloor - r \leq \frac{g+1}{2} - \frac{g+2}{3} = \frac{g-1}{6} < r$, a contradiction.

So, we deduce that $d(A) = 2$, leading to the proof of the lemma. □

Lemma 12. *Suppose that a numerical semigroup Λ has ω gaps between 1 and $n - 1$ and $n \geq 2\omega + 2$ then*

1. $n \in \Lambda$,
2. the Frobenius number of Λ is smaller than n ,
3. the genus of Λ is ω .

Proof. 1. The number of pairs s, t with $1 \leq s \leq t \leq n-1$ such that $s+t = n$ is $\lfloor \frac{n}{2} \rfloor \geq \lfloor \frac{2\omega+2}{2} \rfloor = \omega + 1$. Since there are ω gaps between 1 and $n-1$ this means that in at least one of these pairs both s and t are non-gaps and so n is a sum of two non-gaps and so a non-gap.

2. Using the same argument one can show by induction that $m \in \Lambda$ for all $m \geq n$. Thus, the Frobenius number must be smaller than n .

3. It is a consequence of the previous statements. □

A consequence of this Lemma 12 is the well known fact that the Frobenius number of a numerical semigroup of genus g is at most $2g - 1$. Indeed, otherwise take $\omega = g - 1$ and n the Frobenius number of the semigroup and get a contradiction.

Let Λ be a numerical semigroup. We say that a set $B \subset \mathbb{N}_0$ is Λ -closed if for any $b \in B$ and any λ in Λ , the sum $b+\lambda$ is either in B or it is larger than $\max(B)$. If B is Λ -closed so is $B - \min(B)$. Indeed, $b - \min(B) + \lambda = (b + \lambda) - \min(B)$ is either in $B - \min(B)$ or it is larger than $\max(B) - \min(B) = \max(B - \min(B))$. The new Λ -closed set contains 0. We will denote by $C(\Lambda, i)$ the Λ -closed sets of size i that contain 0.

Theorem 13. *Let $g \in \mathbb{N}_0$ and let r be an integer with $\frac{g+2}{3} \leq r \leq \lfloor \frac{g}{2} \rfloor$. Define $\omega = \lfloor \frac{g}{2} \rfloor - r$*

1. *If Ω is a numerical semigroup of genus ω and B is a Ω -closed set of size $\omega + 1$ and first element equal to 0 then*

$$\{2j : j \in \Omega\} \cup \{2j - 2 \max(B) + 2g + 1 : j \in B\} \cup (2g + \mathbb{N}_0)$$

is a numerical semigroup of genus g and ordinarization number r .

2. *All numerical semigroups of genus g and ordinarization number r can be uniquely written as*

$$\{2j : j \in \Omega\} \cup \{2j - 2 \max(B) + 2g + 1 : j \in B\} \cup (2g + \mathbb{N}_0)$$

for a unique numerical semigroup Ω of genus ω and a unique Ω -closed set B of size $\omega + 1$ and first element equal to 0.

3. *The number of numerical semigroups of genus g and ordinarization number r depends only on ω . It is exactly*

$$\sum_{\text{Semigroups } \Omega \text{ of genus } \omega} \#C(\Omega, \omega + 1).$$

Proof. 1. Suppose that Ω is a numerical semigroup of genus ω and B is a Ω -closed set of size $\omega + 1$ and first element equal to 0. Let $X = \{2j : j \in \Omega\}$, $Y = \{2j - 2 \max(B) + 2g + 1 : j \in B\}$, $Z = (2g + \mathbb{N}_0)$.

First of all let us see that the complement $\mathbb{N}_0 \setminus (X \cup Y \cup Z)$ has g elements. Notice that all elements in X are even while all elements in Y are odd. So, X and Y do not intersect. Also the unique element in $Y \cap Z$ is $2g+1$. The number of elements in the complement will be

$$\begin{aligned} \#\mathbb{N}_0 \setminus (X \cup Y \cup Z) &= 2g - \#\{x \in X : x < 2g\} - \#Y + 1 \\ &= 2g - \#\{s \in \Omega : s < g\} - \#B + 1 \\ &= 2g - \omega - \#\{s \in \Omega : s < g\}. \end{aligned}$$

We know that all gaps of Ω are at most $2\omega - 1 < g$. So, $\#\{s \in \Omega : s < g\} = g - \omega$ and we conclude that $\#\mathbb{N}_0 \setminus (X \cup Y \cup Z) = g$.

Before proving that $X \cup Y \cup Z$ is a numerical semigroup, let us prove that the number of non-zero elements in $X \cup Y \cup Z$ which are smaller than or equal to g is r . Once we prove that $X \cup Y \cup Z$ is a numerical semigroup, this will mean, by Lemma 1, that it has ordinarization number r . On one hand, all elements in Y are larger than g . Indeed, if λ is the enumeration of Ω then $\max(B) \leq \lambda_\omega \leq 2\omega = 2\lfloor \frac{g}{2} \rfloor - 2r \leq g - 2\frac{g+2}{3} < \frac{g}{3}$. Now, for any $j \in B$, $2j - 2\max(B) + 2g + 1 > 2g - 2\max(B) > g$. On the other hand, all gaps of Ω are at most $2\omega - 1 = 2\lfloor \frac{g}{2} \rfloor - 2r - 1 \leq g - \frac{2(g+2)}{3} - 1 < \frac{g}{3} - 1$ and so all the even integers not belonging to X are less than g . So, the number of non-zero non-gaps of $X \cup Y \cup Z$ smaller than or equal to g is $\lfloor \frac{g}{2} \rfloor - \omega = r$.

To see that $X \cup Y \cup Z$ is a numerical semigroup we only need to see that it is closed under addition. It is obvious that $X + Z \subseteq Z$, $Y + Z \subseteq Z$, $Z + Z \subseteq Z$. It is also obvious that $X + X \subseteq X$ since Ω is a numerical semigroup and that $Y + Y \subseteq Z$ since, as we proved before, all elements in Y are larger than g .

It remains to see that $X + Y \subseteq X \cup Y \cup Z$. Suppose that $x \in X$ and $y \in Y$. Then $x = 2i$ for some $i \in \Omega$ and $y = 2j - 2\max(B) + 2g + 1$ for some $j \in B$. Then $x + y = 2(i + j) - 2\max(B) + 2g + 1$. Since B is Ω -closed, we have that either $i + j \in B$ and so $x + y \in Y$ or $i + j > \max(B)$. In this case $x + y \in Z$. So, $X + Y \subseteq Y \cup Z$.

2. First of all notice that, since the Frobenius number of a semigroup Λ of genus g is smaller than $2g$, it holds

$$\Lambda \cap (2g + \mathbb{N}_0) = (2g + \mathbb{N}_0).$$

For any numerical semigroup the set $\Omega = \{\frac{j}{2} : j \in \Lambda \cap (2\mathbb{N}_0)\}$ is a numerical semigroup. If Λ has ordinarization number $r \geq \frac{g+2}{3}$ then, by Lemma 11,

$$\Lambda \cap [0, g] = (2\Omega) \cap [0, g].$$

The semigroup Ω has exactly $r + 1$ non-gaps between 0 and $\lfloor \frac{g}{2} \rfloor$ and $\omega = \lfloor \frac{g}{2} \rfloor - r$ gaps between 0 and $\lfloor \frac{g}{2} \rfloor$. We can use Lemma 12 with $n = \lfloor \frac{g}{2} \rfloor + 1$ since

$$2\omega + 2 = 2\left\lfloor \frac{g}{2} \right\rfloor - 2r + 2 \leq g - \frac{2(g+2)}{3} + 2 = \frac{g+2}{3},$$

which implies $2\omega + 2 \leq \lfloor \frac{g+2}{3} \rfloor \leq \lfloor \frac{g+2}{2} \rfloor = n$. Then the genus of Ω is ω and the Frobenius number of Ω is at most $\lfloor \frac{g}{2} \rfloor$. This means in particular that all even integers larger than g belong to Λ .

Define $D = (\Lambda \cap [0, 2g]) \setminus 2\Omega$. That is, D is the set of odd non-gaps of Λ smaller than $2g$. We claim that $\bar{B} = \{\frac{j-1}{2} : j \in D \cup \{2g+1\}\}$ is a Ω -closed set of size $\omega + 1$. The size follows from the fact that the number of non-gaps of Λ between $g+1$ and $2g$ is $g-r$ and that the number of even integers in the same interval is $\lceil \frac{g}{2} \rceil$. Suppose that $\lambda \in \Omega$ and $b \in \bar{B}$. Then $b = \frac{j-1}{2}$ for some j in $D \cup \{2g+1\}$ and $b + \lambda = \frac{(j+2\lambda)-1}{2}$. If $\frac{(j+2\lambda)-1}{2} \geq \max(\bar{B}) = \frac{(2g+1)-1}{2}$ we are done. Otherwise we have $j + 2\lambda \leq 2g$. Since Λ is a numerical semigroup and both $j, 2\lambda \in \Lambda$, it holds $j + 2\lambda \in \Lambda \cap [0, 2g]$. Furthermore, $j + 2\lambda$ is odd since so is j . So, $b + \lambda$ is either larger than $\max(\bar{B})$ or it is in \bar{B} . Then $B = \bar{B} - \min(\bar{B})$ is a Λ -closed set of size $\omega + 1$ and first element zero.

3. It is a consequence of the previous point. □

Define the sequence f_ω by $f_\omega = \sum_{\text{Semigroups } \Omega \text{ of genus } \omega} \#C(\Omega, \omega + 1)$. The first elements in the sequence, from f_0 to f_{14} are

ω	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
f_ω	1	2	7	23	68	200	615	1764	5060	14626	41785	117573	332475	933891	2609832

From these first elements and Theorem 13 we can deduce $n_{g,r}$ for any g , whenever $r \geq \max(\frac{g+2}{3}, \lfloor \frac{g}{2} \rfloor - 14)$. This is illustrated in Table 2. Since the sequence f_ω is increasing for ω between 0 and 14 we deduce the next corollary.

Corollary 14. *For any $g \in \mathbb{N}$ and any $r \geq \max(\frac{g}{3} + 1, \lfloor \frac{g+1}{2} \rfloor - 14)$, it holds $n_{g,r} \geq n_{g+1,r}$.*

If we proved that $f_\omega \leq f_{\omega+1}$ for any ω , this would imply $n_{g,r} \leq n_{g+1,r}$ for any $r > \frac{g}{3}$.

4 On numerical semigroups with a large number of gap intervals

In this final section we present a bijection between semigroups at a given depth in the ordinarization tree and semigroups with a given (large) number of gap intervals.

Lemma 15. *Suppose that a numerical semigroup Λ has genus g and ordinarization number $r \geq \frac{g+2}{3}$. Then it has $2r$ intervals of gaps if g is even and $2r + 1$ intervals of gaps if g is odd.*

Proof. By Theorem 13 we deduce that, defining $\omega = \lfloor \frac{g}{2} \rfloor - r$, Λ should be

$$\{2j : j \in \Omega\} \cup \{2j - 2 \max(B) + 2g + 1 : j \in B\} \cup (2g + \mathbb{N}_0)$$

for a unique numerical semigroup Ω of genus ω and a unique Ω -closed set B of size $\omega + 1$ and first element equal to 0. But this semigroup has $g - 2\omega$ intervals of gaps. This number is equal to $g - 2\lfloor \frac{g}{2} \rfloor + 2r$ which equals $2r$ if g is even and $2r + 1$ if g is odd. \square

Lemma 16. *A numerical semigroup with n intervals of gaps has ordinarization number at least $\lfloor \frac{n}{2} \rfloor$.*

Proof. The maximum number of gaps between 1 and g is obtained for the semigroup (should it be a semigroup) that has $g - n + 1$ non-gaps in a row right after 0 and then $n - 1$ sequences of a non-gap and a gap and then no more gaps. In this case there would be $\lceil \frac{n-1}{2} \rceil = \lfloor \frac{n}{2} \rfloor$ non-zero non-gaps in the same interval. \square

Theorem 17. *Suppose that a numerical semigroup Λ has genus g and n intervals of gaps with $\lfloor \frac{n}{2} \rfloor \geq \frac{g+2}{3}$. Then g and n have the same parity and Λ has ordinarization number equal to $\lfloor \frac{n}{2} \rfloor$.*

Proof. By Lemma 16, $r \geq \frac{g+2}{3}$, and Lemma 15 gives the result. \square

Concluding, if $\lfloor \frac{n}{2} \rfloor \geq \frac{g+2}{3}$, then the set of numerical semigroups of genus g and n intervals of gaps is empty if n and g have different parity and it is exactly the set of numerical semigroups of genus g and ordinarization number $\lfloor \frac{n}{2} \rfloor$ otherwise.

Acknowledgments

The author is grateful to Anna de Mier for her suggestions on this work. This work was partly supported by the Spanish Government through projects TIN2009-11689 ‘‘RIPUP’’ and CONSOLIDER INGENIO 2010 CSD2007-00004 ‘‘ARES’’, and by the Government of Catalonia under grant 2009 SGR 1135.

References

- [1] Victor Blanco, Pedro A. Garcıa-Sanchez, and Justo Puerto. Counting numerical semigroups with short generating functions. *International Journal of Algebra and Computation*, 21(7):1217–1235, 2011.
- [2] Maria Bras-Amoros. Fibonacci-like behavior of the number of numerical semigroups of a given genus. *Semigroup Forum*, 76(2):379–384, 2008.
- [3] Maria Bras-Amoros. Bounds on the number of numerical semigroups of a given genus. *J. Pure Appl. Algebra*, 213(6):997–1001, 2009.
- [4] Maria Bras-Amoros and Stanislav Bulygin. Towards a better understanding of the semigroup tree. *Semigroup Forum*, 79(3):561–574, 2009.
- [5] Maria Bras-Amoros and Anna de Mier. Representation of numerical semigroups by Dyck paths. *Semigroup Forum*, 75(3):677–682, 2007.

Table 1: Number of numerical semigroups of each ordinarization number for each genus $g \leq 49$.

$r \setminus g$	g=0	g=1	g=2	g=3	g=4	g=5	g=6	g=7	g=8	g=9	g=10	g=11	g=12	g=13	g=14	g=15	g=16	g=17	g=18	g=19	g=20	g=21	
r=0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
r=1						5	9	12	18	22	30	35	45	51	63	70	84	92	108	117	135	145	165
r=2				3	1	2	9	19	39	70	118	196	281	432	586	838	1080	1490	1835	2449	2956	3804	
r=3							1	1	4	16	47	97	228	442	844	1462	2447	4017	6127	9516	13693	20152	
r=4									1	1	2	3	28	60	180	442	1083	2202	4611	8579	15830	27493	
r=5											1	1	2	9	27	83	215	721	1685	4417	9653	19633	
r=6													1	1	2	2	7	9	45	89	319	889	
r=7															1	1	2	2	7	7	25	47	
r=8																	1	1	2	2	7	7	
r=9																			1	1	2	2	
r=10																					1	1	

$r \setminus g$	g=22	g=23	g=24	g=25	g=26	g=27	g=28	g=29	g=30	g=31	g=32	g=33	g=34	g=35	g=36	g=37
r=0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
r=1	176	198	210	234	247	273	287	315	330	360	376	408	425	459	477	513
r=2	4498	5090	6582	8162	9352	11370	12879	15480	17317	20569	22877	26812	29610	34454	37739	45338
r=3	27768	30726	52312	72494	93341	125600	157738	208370	255661	331626	401389	510031	608332	764927	899285	1114817
r=4	46615	76616	120795	189550	285103	429618	618555	905721	1255646	1790138	2418323	3354611	4425179	6031518	7767784	10392189
r=5	21378	41912	83951	153896	281388	487211	831654	1374366	2218771	3524257	5445975	8352388	12435320	18555615	26699519	38853706
r=6	2635	6446	17582	39214	90574	188067	394521	759910	1469758	2662254	4823902	8344482	14314198	23747986	38898550	62372773
r=7	142	340	1266	3483	10171	26489	69692	161111	382713	816457	1763299	3533977	7088495	13371197	25321828	45509829
r=8	23	24	96	157	553	1570	5281	14835	43790	113548	294908	701946	1652408	3632809	7973030	16368101
r=9	7	7	23	23	69	95	301	627	2457	7168	23475	68223	194677	512838	1323375	3178140
r=10	2	2	7	7	23	23	68	70	228	309	1142	2994	10901	33846	109619	318308
r=11	1	1	2	2	7	7	23	23	68	68	202	232	740	1249	4843	14332
r=12			1	1	2	2	7	7	23	23	68	68	200	201	649	759
r=13					1	1	1	1	2	2	7	23	200	68	68	200
r=14								1	1	2	7	7	23	23	68	68
r=15									1	1	2	7	7	23	23	23
r=16											1	1	2	2	7	7
r=17													1	1	2	2
r=18															1	1

$r \setminus g$	g=38	g=39	g=40	g=41	g=42	g=43	g=44	g=45	g=46	g=47	g=48	g=49
r=0	1	1	1	1	1	1	1	1	1	1	1	1
r=1	532	570	590	630	651	693	715	759	782	828	852	900
r=2	4710	5430	58886	67072	72419	81855	88142	98946	106170	118716	128844	141164
r=3	1299978	1590237	1836517	2226669	2545983	3059220	3477286	4134725	4699073	5518427	6185260	7256830
r=4	13180451	17322789	21616641	28040199	34458068	44142389	53663689	67788397	81530366	102094609	121404838	15047267
r=5	54507523	77486888	106904921	148091995	198378083	272201928	358476988	483240666	626315811	833944191	1063739070	1397557241
r=6	98298482	152816803	232801667	352797809	521753229	772496765	1114485292	1614321267	227596011	3242295418	4478817624	6268430457
r=7	81612546	140878791	241699680	402445801	664483703	1072569052	1711738040	2688862529	4168280931	6388426599	9636305171	14462411903
r=8	33550240	65385970	129699443	235541563	436401532	774742260	1380117648	2375549463	4064063006	6774823275	11221522599	18209647612
r=9	7487630	16760501	36890000	77385799	160762381	319966692	631894288	1203245544	2273796763	4158338885	7567139870	13367227712
r=10	899807	2383461	6101724	14810797	34997273	79159902	175168573	373545010	782283651	1585487022	3171168252	6150909456
r=11	51063	164512	519339	1509557	4237829	11221868	28679326	70097864	169662233	379419480	845334246	1824288237
r=12	2527	5652	21994	71261	252707	803934	2492982	7226212	20114114	53281902	136131501	334153690
r=13	616	649	1925	2679	9947	27432	106780	361575	1245778	3945659	12053243	34718395
r=14	200	200	615	617	1800	1939	6144	11138	43824	140489	537134	1835716
r=15	68	68	200	200	615	615	1766	1804	5254	6320	22087	52194
r=16	23	23	68	68	200	200	615	1764	1765	5102	5278	5278
r=17	7	7	23	23	68	68	200	615	615	1764	1764	1764
r=18	2	2	7	7	23	23	68	68	200	200	615	615
r=19	1	1	2	2	7	7	23	23	68	68	200	200
r=20			1	1	2	2	7	7	23	23	68	68
r=21					1	1	2	2	7	7	23	23
r=22							1	1	2	2	7	7
r=23									1	1	2	2
r=24											1	1

- [6] Sergi Elizalde. Improved bounds on the number of numerical semigroups of a given genus. *J. Pure Appl. Algebra*, 214(10):1862–1873, 2010.
- [7] G. A. Freĭman. On the addition of finite sets. *Dokl. Akad. Nauk SSSR*, 158:1038–1041, 1964.
- [8] G. A. Freĭman. *Nachala strukturnoi teorii slozheniya mnozhestv*. Kazan. Gosudarstv. Ped. Inst, 1966.
- [9] Nathan Kaplan. Counting numerical semigroups by genus and some cases of a question of Wilf. *Journal of Pure and Applied Algebra*, 216(5):1016–1032, 2012.
- [10] Jiryō Komeda. On non-Weierstrass gap sequences. Technical report, Kanagawa Institute of Technology B-13, 1989.
- [11] Jiryō Komeda. Non-Weierstrass numerical semigroups. *Semigroup Forum*, 57(2):157–185, 1998.
- [12] Melvyn B. Nathanson. *Additive number theory*, volume 165 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1996. Inverse problems and the geometry of sumsets.
- [13] J. C. Rosales, P. A. García-Sánchez, J. I. García-García, and J. A. Jiménez Madrid. The oversemigroups of a numerical semigroup. *Semigroup Forum*, 67(1):145–158, 2003.
- [14] José C. Rosales and Pedro A. García-Sánchez. *Numerical semigroups*, volume 20 of *Developments in Mathematics*. Springer, New York, 2009.
- [15] Alex Zhai. Fibonacci-like growth of numerical semigroups of a given genus. arXiv:1111.3142v1, 2011.
- [16] Yufei Zhao. Constructing numerical semigroups of a given genus. *Semigroup Forum*, 80(2):242–254, 2010.