

INEQUALITIES FOR EIGENVALUES OF THE WEIGHTED HODGE LAPLACIAN

DAGUANG CHEN* AND YINGYING ZHANG

ABSTRACT. In this paper, we obtain "universal" inequalities for eigenvalues of the weighted Hodge Laplacian on a compact self-shrinker of Euclidean space. These inequalities generalize the Yang-type and Levitin-Parnovski inequalities for eigenvalues of the Laplacian and Laplacian. From the recursion formula of Cheng and Yang [12], the Yang-type inequality for eigenvalues of the weighted Hodge Laplacian are optimal in the sense of the order of eigenvalues.

1. INTRODUCTION

Let M^m be an m -dimensional complete Riemannian manifold and Ω be a bounded domain in M^m . The Dirichlet eigenvalue problem of Laplacian is given by

$$\begin{cases} \Delta u = -\lambda u, & \text{in } \Omega \\ u = 0, & \text{on } \partial\Omega. \end{cases} \quad (1.1)$$

It is well known that the spectrum of this problem is real and discrete:

$$0 < \lambda_1 < \lambda_2 \leq \lambda_3 \leq \cdots \nearrow \infty,$$

where each λ_i has finite multiplicity which is repeated according to its multiplicity.

The main developments were obtained by Payne, Pólya and Weinberger [32], Hile and Protter [24] and Yang [36]. In 1956, Payne, Pólya and Weinberger [32] proved that

$$\lambda_{k+1} - \lambda_k \leq \frac{4}{mk} \sum_{i=1}^k \lambda_i. \quad (1.2)$$

In 1980, Hile and Protter [24] improved (1.2) to

$$\sum_{i=1}^k \frac{\lambda_i}{\lambda_{k+1} - \lambda_i} \geq \frac{mk}{4}. \quad (1.3)$$

2000 *Mathematics Subject Classification.* 35P15; 58J50; 58C40; 58A10.

Key words and phrases. Eigenvalues, Weighted Hodge Laplacian, Universal inequalities, Self-shrinker.

* This work of the first named author was partially supported by NSFC grant No. 11101234.

In 1991, Yang (see [36] and more recently [11]) obtained a very sharp inequality

$$\sum_{i=1}^k (\lambda_{k+1} - \lambda_i) (\lambda_{k+1} - (1 + \frac{4}{m}) \lambda_i) \leq 0. \quad (1.4)$$

There has been much work dedicated to extending and strengthening the classical inequalities of Payne-Pólya-Weinberger, Hile-Protter and Yang. When M^m is an m -dimensional compact manifold, there are similar results about the eigenvalue estimates for the Laplacian (see, e.g. [10, 11, 17, 28, 31, 37]). For the compact Riemannian manifolds isometrically immersed in an Euclidean space or a sphere, J. M. Lee [27] proved Hile-Protter type bounds for eigenvalues for Hodge Laplacian on p -forms. In 2002, B. Colbois [15] derived a Payne-Pólya-Weinberger type inequality for the rough Laplacian. In [25], S. Ilias and O. Makhoul obtained inequalities for the eigenvalues of the Hodge Laplacian.

In 1991, N. Anghel [1] obtained the analogous estimate of (1.2) for the Dirac operator. In 2009, the Yang-type inequality (1.4) was extended to the eigenvalues of Dirac operator by the first author in [8].

In [19], Harrell gave an abstract algebraic argument involving operators, their commutators and traces, which generalize the original PPW arguments. These algebraic ideas were developed in different contexts to produce many new universal eigenvalues inequalities (see [5, 20–23, 30]).

In present paper, making use of a theorem of Ashbaugh and Hermi [5], we obtain the Yang-type inequality for higher order eigenvalues of the weighted Hodge Laplacian for submanifolds in Euclidean space.

Theorem 1.1. *Let $x : (M^m, g) \rightarrow (\mathbb{R}^n, \text{can})$ be a compact self-shrinker, $\Delta_{p,x} = \Delta_H + \frac{1}{2} \mathcal{L}_{\nabla|x|^2}$ (see below (2.9)) be the weighted Hodge Laplacian acting on p -forms over M^m . Assume that $\{\lambda_i^{(p)}\}_{i=1}^{\infty}$ are the eigenvalues of $\Delta_{p,x}$ and $\{\varphi_i\}_{i=1}^{\infty}$ is a corresponding orthonormal basis of p -eigenforms. We have, for any $p \in \{0, 1, \dots, m\}$,*

$$\begin{aligned} m \sum_{i=1}^k \left(\lambda_{k+1}^{(p)} - \lambda_i^{(p)} \right)^2 &\leq \sum_{i=1}^k \left(\lambda_{k+1}^{(p)} - \lambda_i^{(p)} \right) \left(4\lambda_i^{(p)} + 2m \right. \\ &\quad - \int_{M^m} |x|^2 |\varphi_i|^2 e^{-\frac{|x|^2}{2}} d\text{vol} \\ &\quad - 4 \int_{M^m} \langle \mathfrak{Ric} \varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} d\text{vol} \\ &\quad \left. + 4 \int_{M^m} \langle \nabla(\nabla \frac{|x|^2}{2}) \varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} d\text{vol} \right) \end{aligned} \quad (1.5)$$

where $\{e_i\}_{i=1}^m$ is a local orthonormal basis of TM^m with respect to the induced metric g and $\mathfrak{Ric} = -\omega^i \wedge \iota(e_j) R(e_i, e_j)$ (see also 2.8) is the curvature operator acting on p -forms.

Remark 1.1. When $p = 0$, i.e., $\lambda_i := \lambda_i^{(0)}$ are the eigenvalues of the operator $\mathfrak{L} := \Delta_{0,x} = \Delta + \langle x, \cdot \rangle$ acting on scalar functions, we have

$$\begin{aligned} m \sum_{i=1}^k (\lambda_{k+1} - \lambda_i)^2 &\leq \sum_{i=1}^k (\lambda_{k+1} - \lambda_i) \left(4\lambda_i + 2m - \int_{M^m} |x|^2 |\varphi_i|^2 e^{-\frac{|x|^2}{2}} dvol \right) \\ &\leq \sum_{i=1}^k (\lambda_{k+1} - \lambda_i) \left(4\lambda_i + 2m - \min_{M^m} |x|^2 \right), \end{aligned} \quad (1.6)$$

which is Theorem 1.1 in [13]. Therefore, Theorem 1.1 generalizes eigenvalue estimates from the operator \mathfrak{L} to the weighted Hodge Laplacian $\Delta_{p,x}$.

Remark 1.2. If $|x| = c$, ($c > 0$), the manifold M^m is a submanifold of sphere $\mathbb{S}^{n-1}(\frac{1}{c})$ in Euclidean space \mathbb{R}^n . Furthermore, the weighted Hodge Laplacian $\Delta_{p,x}$ is reduced to the ordinary one.

For a compact self-shrinker (see (2.4)) in Euclidean space, we have

Corollary 1.1. Let $x : (M^m, g) \rightarrow (\mathbb{R}^n, \text{can})$ be a compact self-shrinker, H, h be the second fundamental form and the mean curvature of the immersion x , respectively. We have, $p \in \{1, \dots, m\}$,

$$\begin{aligned} &\sum_{i=1}^k (\lambda_{k+1}^{(p)} - \lambda_i^{(p)})^2 \\ &\leq \frac{4}{m} \sum_{i=1}^k (\lambda_{k+1}^{(p)} - \lambda_i^{(p)}) \left[\lambda_i^{(p)} + \frac{m}{2} + 1 \right. \\ &\quad \left. + \int_{M^m} \left(p|H||h| - \Phi(H, h) - \frac{1}{4}|x|^2 \right) |\varphi_i|^2 e^{-\frac{|x|^2}{2}} dvol \right] \quad (1.7) \\ &\leq \frac{4}{m} \sum_{i=1}^k (\lambda_{k+1}^{(p)} - \lambda_i^{(p)}) \left[\lambda_i^{(p)} + \frac{m}{2} + 1 \right. \\ &\quad \left. + \max_{M^m} \left(p|H||h| - \Phi(H, h) - \frac{1}{4}|x|^2 \right) \right], \end{aligned}$$

where $\Phi(H, h)$ is a function depending on the second fundamental form h and the mean curvature H defined in (3.10).

From Theorem 1.1, we can obtain the spectral gaps of the consecutive eigenvalues of the weighted Hodge Laplacian $\Delta_{p,x}$.

Corollary 1.2. *Under the same assumption in Corollary 1.1, we have*

$$\lambda_{k+1}^{(p)} - \lambda_k^{(p)} \leq 2 \left[\left(\frac{2}{m} \frac{1}{k} \sum_{i=1}^k \lambda_i^{(p)} + \frac{2}{m} + \frac{2}{m} \max_{M^m} \left(p|H||h| - \Phi(H, h) - \frac{1}{4}|x|^2 \right) \right)^2 - \left(1 + \frac{4}{m} \right) \frac{1}{k} \sum_{j=1}^k \left(\lambda_j^{(p)} - \frac{1}{k} \sum_{i=1}^k \lambda_i^{(p)} \right)^2 \right]^{\frac{1}{2}}$$

For the lower order eigenvalues of (1.1), in 1956, Payne, Pólya and Weinberger [32] proved that for $\Omega \subset \mathbb{R}^2$,

$$\lambda_2 + \lambda_3 \leq 6\lambda_1,$$

which was extended to domains $\Omega \subset \mathbb{R}^m$ in [35] (or see Section 3.2 of [2])

$$\sum_{i=1}^m (\lambda_{i+1} - \lambda_1) \leq 4\lambda_1.$$

There are also a variety of extensions of results of this type, for examples, see [2, 7–11, 34]. Recently, S. Ilias and O. Makhoul [26] obtained the universal inequality for eigenvalues of the Hodge Laplacian.

In the second part of this paper, by using an algebraic identity deduced by Levitin and Parnovski [30], we can obtain

Theorem 1.2. *Let $x : (M^m, g) \rightarrow (\mathbb{R}^n, \text{can})$ be a compact self-shrinker and $\Delta_{p,x}$ be the weighted Hodge Laplacian defined acting on p -forms over M^m . Assume that $\{\lambda_i^{(p)}\}_{i=1}^{\infty}$ are the eigenvalues of $\Delta_{p,x}$ and $\{\varphi_i\}_{i=1}^{\infty}$ is a corresponding orthonormal basis of p -eigenforms. We have, for any $p \in \{0, 1, \dots, m\}$,*

$$\begin{aligned} \sum_{l=1}^m (\lambda_{i+l}^{(p)} - \lambda_i^{(p)}) &\leq 4\lambda_i^{(p)} + 2m - \int_{M^m} |x|^2 |\varphi_i|^2 e^{-\frac{|x|^2}{2}} d\text{vol} \\ &\quad - 4 \int_{M^m} \langle \mathfrak{Ric} \varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} d\text{vol} \\ &\quad + 4 \int_{M^m} \langle \nabla(\nabla \frac{|x|^2}{2}) \varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} d\text{vol}. \end{aligned} \quad (1.8)$$

Remark 1.3. *When $p = 0$, i.e., $\lambda_i = \lambda_i^{(0)}$ is the eigenvalues of the operator $\mathfrak{L} = \Delta_{0,x} = \Delta + \langle x, \cdot \rangle$ acting on scalar functions, we have*

$$\begin{aligned} \sum_{l=1}^m (\lambda_{i+l} - \lambda_i) &\leq 4\lambda_i + 2m - \int_{M^m} |x|^2 |\varphi_i|^2 e^{-\frac{|x|^2}{2}} d\text{vol} \\ &\leq 4\lambda_i + 2m - \min_{M^m} |x|^2. \end{aligned} \quad (1.9)$$

Since i is arbitrary, (1.9) is more general than Proposition 4.1 in [13].

Corollary 1.3. *Let $x : (M^m, g) \rightarrow (\mathbb{R}^n, \text{can})$ be a self-shrinker, H, h be the second fundamental form and the mean curvature of the immersion x , respectively. Assume that $\{\lambda_i^{(p)}\}_{i=1}^\infty$ are the eigenvalues of $\Delta_{p,x}$ and $\{\varphi_i\}_{i=1}^\infty$ is a corresponding orthonormal basis of p -eigenforms. We obtain, for $p \in \{1, \dots, m\}$,*

$$\begin{aligned} \sum_{l=1}^m \left(\lambda_{i+l}^{(p)} - \lambda_i^{(p)} \right) &\leq 4\lambda_i^{(p)} + 2m + 4 \\ &\quad + 4 \int_{M^m} \left(p|H||h| - \Phi(H, h) - \frac{1}{4}|x|^2 \right) |\varphi_i|^2 e^{-\frac{|x|^2}{2}} d\text{vol} \\ &\leq 4\lambda_i^{(p)} + 2m + 4 + \max_{M^m} \left(p|H||h| - \Phi(H, h) - \frac{1}{4}|x|^2 \right). \end{aligned} \tag{1.10}$$

Furthermore, from the recursion formula of Cheng and Yang [12], we can obtain an upper bound for eigenvalue $\lambda_k^{(p)}$:

Corollary 1.4. *Let M^m be an m -dimensional compact self-shrinker in \mathbb{R}^n . Then, eigenvalues of the weighted Hodge Laplacian $\Delta_{p,x}$ satisfy, for any $k \geq 1$,*

$$\mu_{k+1} \leq C_0(m) k^{\frac{2}{m}} \mu_1$$

where $C_0(m) \leq 1 + \frac{4}{m}$ is a constant and $\mu_i = \lambda_i^{(p)} + \frac{m}{2} + 1 + \max_{M^m} \left(p|H||h| - \Phi(H, h) - \frac{1}{4}|x|^2 \right)$.

This paper is organized as follows: In Section 2, we present some formulas for submanifolds in Euclidean space, the definitions of the weighted Hodge Laplacian. In Section 3, in order to prove main theorems, we derive several lemmas for differential forms. In Section 4 and Section 5, we give the proofs of Theorem 1.1 and 1.2.

Acknowledgments. The authors wish to express their gratitude to Professors Huaidong Cao and Xiaofeng Sun for their suggestions and useful discussions. This work of the first named author was done while the author visited Department of Mathematics, Lehigh University, USA. He also would like to thank the institute for its hospitality.

2. PRELIMINARIES

2.1. Submanifold in Euclidean space and self-shrinker. Let $x : M^m \rightarrow \mathbb{R}^n$ be an m -dimensional submanifold of n -dimensional Euclidean space \mathbb{R}^n . Let $\{e_1, \dots, e_m\}$ be a local orthonormal basis of TM^m with respect to the induced metric, and $\{\omega^1, \dots, \omega^m\}$ be their dual 1-forms. Let $\{e_{m+1}, \dots, e_n\}$ be the local orthonormal unit normal vector fields. In this paper we make the following conventions on the range of indices:

$$1 \leq i, j, k \leq m; \quad m+1 \leq \alpha, \beta, \gamma \leq n.$$

Then we have the following structure equations (see [8, 13])

$$\begin{aligned} dx &= \omega^i e_i, & \omega^\alpha &= 0, \\ de_i &= \omega_i^j e_j + \omega_i^\alpha e_\alpha, & \omega_i^\alpha &= h_{ij}^\alpha \omega^j, \\ de_\alpha &= \omega_\alpha^j e_j + \omega_\alpha^\beta e_\beta, \end{aligned} \quad (2.1)$$

where h_{ij}^α denote the the components of the second fundamental form of M^m . We denote by $|h|^2 = \sum_{\alpha, i, j} (h_{ij}^\alpha)^2$, the norm square of the second fundamental form, $H = \sum_\alpha H^\alpha e_\alpha = \sum_\alpha (\sum_i h_{ii}^\alpha) e_\alpha$ the mean curvature vector field over M^m .

One can deduce that, pointwise on M^m ,

$$\sum_{A=1}^n |\nabla x^A|^2 = m, \quad (2.2)$$

and

$$\frac{1}{2} |x|_{,ij}^2 = \frac{1}{2} \left(\sum_{A=1}^n (x^A)^2 \right)_{,ij} = \langle h_{ij}^\alpha e_\alpha, x \rangle + \delta_{ij}. \quad (2.3)$$

The submanifold M^m is called a self-shrinker [16] if it satisfies the quasi-linear elliptic system:

$$H = -x^\perp, \quad (2.4)$$

where H denotes the mean curvature vector field of the immersion and \perp is the projection onto the normal bundle of M^m .

2.2. Differential forms and the weighted Hodge Laplacian. Let (M^m, g) be an m -dimensional compact Riemannian manifold. For any two p -forms φ and ψ , we let $\varphi_{i_1 \dots i_p} = \varphi(e_{i_1}, \dots, e_{i_p})$ and $\psi_{i_1 \dots i_p} = \psi(e_{i_1}, \dots, e_{i_p})$ denote the components of φ and ψ , with respect to a local orthonormal frame $\{e_i\}_{i=1}^m$. Their pointwise inner product with respect to Riemannian metric g is given by

$$\begin{aligned} \langle \varphi, \psi \rangle &= \sum_{1 \leq i_1 < \dots < i_p \leq m} \varphi_{i_1 \dots i_p} \psi_{i_1 \dots i_p} \\ &= \frac{1}{p!} \sum_{1 \leq i_1, \dots, i_p \leq m} \varphi_{i_1 \dots i_p} \psi_{i_1 \dots i_p}. \end{aligned}$$

We denote by Δ_p the Hodge Laplacian acting on p -forms

$$\Delta_p = (d\delta + \delta d), \quad (2.5)$$

where d is the exterior derivative acting on p -forms and δ is the adjoint of d with respect to Riemannian measure $dvol$.

In [6, 33], the operator (2.5) is generalized to the weighted Hodge Laplacian acting on differential forms. Let $f \in C^\infty(M^m, \mathbb{R})$ be a smooth function

defined on M^m . When the Riemannian measure is changed from being $dvol$ to $e^{-f}dvol$, it is natural to define the weighted Hodge Laplacian by

$$\Delta_{p,f} = d\delta' + \delta'd \quad (2.6)$$

where $\delta' = e^f\delta e^{-f}$, which is the adjoint operator of the exterior derivative d with respect to Riemannian measure $e^{-f}dvol$.

For the weighted Hodge Laplacian, we have the following Bochner-Weitzenböck type formula [33]

$$\begin{aligned} \Delta_{p,f} &= \Delta_p + \mathcal{L}_{\nabla f} \\ &= \nabla^*\nabla - \omega^i \wedge \iota(e_j)R(e_i, e_j) + \mathcal{L}_{\nabla f} \\ &= \nabla_f^*\nabla - \omega^i \wedge \iota(e_j)R(e_i, e_j) - \nabla(\nabla f) \\ &= \nabla_f^*\nabla + \mathfrak{Ric} - \nabla(\nabla f) \end{aligned} \quad (2.7)$$

where \mathcal{L} is the Lie derivative, $\iota(X)$ for $X \in \Gamma(TM^m)$ is inner product acting on forms, ∇X acting on form φ is given by

$$\nabla X\varphi = X^j{}_{,l}\omega^l \wedge \iota(e_j)\varphi$$

and

$$\mathfrak{Ric} = -\omega^i \wedge \iota(e_j)R(e_i, e_j). \quad (2.8)$$

With respect to the measure $e^{-f}dvol$, the spectrum of $\Delta_{p,f}$ consists of a nondecreasing, unbounded sequence of eigenvalues with finite multiplicities

$$\text{Spec}(\Delta_{p,f}) = \{0 \leq \lambda_1^{(p)} \leq \lambda_2^{(p)} \leq \lambda_3^{(p)} \leq \dots \leq \lambda_k^{(p)} \leq \dots\}.$$

Let $x = (x^1, \dots, x^n) : M^m \rightarrow \mathbb{R}^n$ be an m -dimensional submanifold of \mathbb{R}^n . In this article, we will consider the operator (2.6) over M^m , for $f = e^{\frac{1}{2}|x|^2}$,

$$\Delta_{p,x} = d\delta' + \delta'd. \quad (2.9)$$

For $\Delta_{p,x}$ acting on the scalar functions, the operator (2.9) [16]¹ is given by

$$\mathfrak{L} = \Delta + \langle x, \cdot \rangle = -e^{\frac{|x|^2}{2}} \text{div} \left(e^{-\frac{|x|^2}{2}} d \right) = \delta'd = \Delta_{0,x}. \quad (2.10)$$

where Δ is the positive operator. If M^m is a self-shrinker, we have

$$\mathfrak{L}x^A = x^A, \quad A = 1, \dots, n. \quad (2.11)$$

3. SOME LEMMAS

In order to prove our main theorems, we will derive some lemmas in this section.

By the direct calculations, we have

¹The Laplacian operator is different in [16] with a minus sign.

Lemma 3.1. For $f, u \in C^\infty(M, \mathbb{R})$ and $\varphi \in \bigwedge^p(T^*M^m)$, we have

$$\mathcal{L}_{\nabla f}(u\varphi) = g(\nabla f, \nabla u)\varphi + u\mathcal{L}_{\nabla f}\varphi. \quad (3.1)$$

$$[\Delta_{p,f}, u]\varphi = [\Delta_p, u]\varphi + [\mathcal{L}_{\nabla f}, u]\varphi \quad (3.2)$$

$$\delta_f(u\varphi) = -\iota(\nabla u)\varphi + u\delta_f\varphi \quad (3.3)$$

where $[\Delta_{p,f}, u]\varphi = \Delta_{p,f}(u\varphi) - u\Delta_{p,f}\varphi$.

Lemma 3.2. Assuming that T_{ij} is a symmetric 2-tensor, we have, for any p -form φ ,

$$\begin{aligned} T_{ij}\omega^i \wedge \iota(e_j)\varphi &= \frac{1}{p!} \sum_{i_1, \dots, i_p} (T\varphi)_{i_1 \dots i_p} \omega^{i_1} \wedge \dots \wedge \omega^{i_p} \\ &= \frac{1}{(p-1)!} \sum_{i_1, \dots, i_p} \left(\sum_j T_{ji_1} \varphi_{ji_2 \dots i_p} \right) \omega^{i_1} \wedge \dots \wedge \omega^{i_p}, \end{aligned} \quad (3.4)$$

and

$$\left\langle \sum_{i,j=1}^m T_{ij}\omega^i \wedge \iota(e_j)\varphi, \varphi \right\rangle = \frac{1}{(p-1)!} \sum_{j, i_1, \dots, i_p} T_{ji_1} \varphi_{ji_2 \dots i_p} \varphi_{i_1 \dots i_p} \quad (3.5)$$

where $(T\varphi)_{i_1 \dots i_p} = \sum_{j=1}^m \sum_{k=1}^p T_{jk} \varphi_{i_1 \dots j \dots i_p}$.

Proof. Assuming that $\varphi = \frac{1}{p!} \sum_{i_1, \dots, i_p} \varphi_{i_1 \dots i_p} \omega^{i_1} \wedge \dots \wedge \omega^{i_p}$, then we get

$$\begin{aligned} & T_{ij}\omega^i \wedge \iota(e_j)\varphi \\ &= \frac{1}{p!} \sum_{i,j=1}^m T_{ij}\omega^i \wedge \iota(e_j) \left(\sum_{i_1, \dots, i_p} \varphi_{i_1 \dots i_p} \omega^{i_1} \wedge \dots \wedge \omega^{i_p} \right) \\ &= \frac{1}{p!} \sum_{i,j, i_1, \dots, i_p} \sum_{k=1}^p (-1)^{k-1} T_{ij} \delta_j^{i_k} \varphi_{i_1 \dots i_p} \omega^i \wedge \omega^{i_1} \wedge \dots \wedge \widehat{\omega^{i_k}} \wedge \dots \wedge \omega^{i_p} \\ &= \frac{1}{p!} \sum_{i, i_1, \dots, i_k, \dots, i_p} \sum_{k=1}^p (-1)^{k-1} T_{ij} \varphi_{i_1 \dots j \dots i_p} \omega^i \wedge \omega^{i_1} \wedge \dots \wedge \widehat{\omega^{i_k}} \wedge \dots \wedge \omega^{i_p} \\ &= \frac{1}{p!} \sum_{i_1, \dots, i_p} (T\varphi)_{i_1 \dots i_p} \omega^{i_1} \wedge \dots \wedge \omega^{i_p} \\ &= \frac{1}{(p-1)!} \sum_{i_1, \dots, i_p} T_{ji_1} \varphi_{ji_2 \dots i_p} \omega^{i_1} \wedge \dots \wedge \omega^{i_p}. \end{aligned}$$

Therefore, we obtain

$$\begin{aligned} \left\langle \sum_{i,j=1}^m T_{ij} \omega^i \wedge \iota(e_j) \varphi, \varphi \right\rangle &= \frac{1}{p!} \sum_{i_1, \dots, i_p} (T\varphi)_{i_1 \dots i_p} \varphi_{i_1 \dots i_p} \\ &= \frac{1}{(p-1)!} \sum_{j, i_1, \dots, i_p} T_{ji_1} \varphi_{ji_2 \dots i_p} \varphi_{i_1 \dots i_p}. \end{aligned}$$

□

Lemma 3.3. *Under the same assumptions in Lemma 3.2, then we have*

$$\left| \sum_{i,j,i_2, \dots, i_p} T_{ij} \varphi_{ii_2 \dots i_p} \varphi_{ji_2 \dots i_p} \right| \leq p! |T| |\varphi|^2 \quad (3.6)$$

and

$$\left\langle \sum_{i,j=1}^m T_{ij} \omega^i \wedge \iota(e_j) \varphi, \varphi \right\rangle \leq p! |T| |\varphi|^2 \quad (3.7)$$

where $|T| = \left(\sum_{i,j} T_{ij}^2 \right)^{\frac{1}{2}}$.

Proof.

$$\begin{aligned} \left| \sum_{i,j,i_2, \dots, i_p} T_{ij} \varphi_{ii_2 \dots i_p} \varphi_{ji_2 \dots i_p} \right| &= \left| \sum_{i_2, \dots, i_p} \sum_j \left(\sum_i T_{ij} \varphi_{ii_2 \dots i_p} \right) (\varphi_{ji_2 \dots i_p}) \right| \\ &\leq \sum_{i_2, \dots, i_p} \left(\sum_j \left(\sum_i T_{ij} \varphi_{ii_2 \dots i_p} \right)^2 \sum_k \varphi_{ki_2 \dots i_p}^2 \right)^{\frac{1}{2}} \\ &\leq \sum_{i_2, \dots, i_p} \left(\sum_j \sum_i T_{ij}^2 \sum_l \varphi_{li_2 \dots i_p}^2 \sum_k \varphi_{ki_2 \dots i_p}^2 \right)^{\frac{1}{2}} \\ &= \left(\sum_{i,j} T_{ij}^2 \right)^{\frac{1}{2}} \sum_{k, i_2, \dots, i_p} \varphi_{ki_2 \dots i_p}^2 \\ &= |T| \sum_{i, i_2, \dots, i_p} \varphi_{ii_2 \dots i_p}^2 \\ &= p! |T| |\varphi|^2. \end{aligned}$$

□

Lemma 3.4. *Assume that $x : M^m \rightarrow \mathbb{R}^n$ is a compact self-shrinker, H, h are the second fundamental form and the mean curvature of the immersion*

x , respectively. We have, for any p -form φ , $p \in \{1, \dots, m\}$

$$\begin{aligned} & \int_{M^m} \langle \nabla(\nabla \frac{|x|^2}{2}) \varphi, \varphi \rangle e^{-\frac{|x|^2}{2}} d\text{vol} \\ & \leq \int_{M^m} |\varphi|^2 e^{-\frac{|x|^2}{2}} d\text{vol} + p \int_{M^m} |H||h||\varphi|^2 e^{-\frac{|x|^2}{2}} d\text{vol} \\ & \leq \int_{M^m} |\varphi|^2 e^{-\frac{|x|^2}{2}} d\text{vol} + p \max_{M^m} |H||h|. \end{aligned} \quad (3.8)$$

Proof. From (2.3), and taking $T_{ij} = \langle h_{ij}^\alpha e_\alpha, x \rangle$ in (3.7), we obtain

$$\begin{aligned} & \int_{M^m} \langle \nabla(\nabla \frac{|x|^2}{2}) \varphi, \varphi \rangle e^{-\frac{|x|^2}{2}} d\text{vol} \\ & = \int_{M^m} \langle \langle h_{ij}^\alpha e_\alpha, x \rangle \omega^i \wedge \iota(e_j) \varphi, \varphi \rangle e^{-\frac{|x|^2}{2}} d\text{vol} + \int_{M^m} |\varphi|^2 e^{-\frac{|x|^2}{2}} d\text{vol} \\ & \leq \int_{M^m} |\varphi|^2 e^{-\frac{|x|^2}{2}} d\text{vol} + p \int_{M^m} \left(\sum_{i,j} \langle h_{ij}^\alpha e_\alpha, x \rangle^2 \right)^{\frac{1}{2}} |\varphi|^2 e^{-\frac{|x|^2}{2}} d\text{vol} \\ & = \int_{M^m} |\varphi|^2 e^{-\frac{|x|^2}{2}} d\text{vol} + p \int_{M^m} \left(\sum_{i,j} (H^\alpha h_{ij}^\alpha)^2 \right)^{\frac{1}{2}} |\varphi|^2 e^{-\frac{|x|^2}{2}} d\text{vol} \\ & \leq \int_{M^m} |\varphi|^2 e^{-\frac{|x|^2}{2}} d\text{vol} + p \int_{M^m} |H||h||\varphi|^2 e^{-\frac{|x|^2}{2}} d\text{vol} \\ & \leq \int_{M^m} |\varphi|^2 e^{-\frac{|x|^2}{2}} d\text{vol} + p \max_{M^m} |H||h|. \end{aligned}$$

□

Combining Proposition 4.1 in [18] and Theorem 1.1 in [29], we obtain the estimate of \mathfrak{Ric} (2.8) acting on p -forms. (c.f. Theorem 3.2 of [25])

Lemma 3.5.

$$\langle \mathfrak{Ric}(\varphi), \varphi \rangle \geq \Phi(h, H) |\varphi|^2, \quad \varphi \in \wedge^p(T^*M^m), \quad (3.9)$$

where

$$\begin{aligned} \Phi(h, H) = & \left\{ -p^2 \left[\left(\frac{m-5}{4} \right) |H|^2 + |h|^2 - \frac{1}{4m^2} \left(\sqrt{m-1}(m-2)|H| \right. \right. \right. \\ & \left. \left. \left. - 2\sqrt{m|h|^2 - |H|^2} \right)^2 \right] - \frac{1}{2} \sqrt{p(p-1)} (|H|^2 + |h|^2) \right\}. \end{aligned} \quad (3.10)$$

4. INEQUALITIES FOR EIGENVALUES

In order to obtain the extrinsic bounds of higher order eigenvalues of the weighted Hodge Laplacian, we firstly introduce the abstract formula derived by Ashbaugh and Hermi [5].

Let \mathfrak{H} be a complex Hilbert space with inner product (\cdot, \cdot) , $\mathcal{A} : \mathcal{D} \subset \mathfrak{H} \rightarrow \mathfrak{H}$ a self-adjoint operator defined on a dense domain \mathcal{D} that is bounded below

and has a discrete spectrum

$$\lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \cdots .$$

Let $\{\mathcal{B}_k : \mathcal{A}(\mathcal{D}) \rightarrow \mathfrak{H}\}_{k=1}^N$ be a collection of symmetric operators leaving \mathcal{D} invariant and $\{\varphi_i, \lambda_i\}_{i=1}^\infty$ be the spectral resolution of \mathcal{A} . Moreover, $\{\varphi_i\}_{i=1}^\infty$ consisting of the orthonormal basis w.r.t. inner product (\cdot, \cdot) for \mathfrak{H} is assumed. Define the commutator $[\mathcal{A}, \mathcal{B}]$ and the norm $\|\varphi\|$ by, respectively

$$[\mathcal{A}, \mathcal{B}] = \mathcal{A}\mathcal{B} - \mathcal{B}\mathcal{A}, \quad \|\varphi\|^2 = (\varphi, \varphi).$$

Based on commutator algebra and the Rayleigh-Ritz principle, M.S. Ashbaugh and L. Hermi [5] obtained

Theorem 4.1. *The eigenvalues λ_i of the operator \mathcal{A} satisfy the Yang-type inequality*

$$\sum_{i=1}^k (\lambda_{k+1} - \lambda_i)^2 \rho_i \leq \sum_{i=1}^k (\lambda_{k+1} - \lambda_i) \Lambda_i \quad (4.1)$$

where ρ_i, Λ are defined by, respectively,

$$\begin{aligned} \rho_i &= \sum_{k=1}^N \langle [\mathcal{A}, \mathcal{B}_k] \varphi_i, \mathcal{B}_k \varphi_i \rangle \\ \Lambda_i &= \sum_{k=1}^N \|[\mathcal{A}, \mathcal{B}_k] \varphi_i\|^2. \end{aligned}$$

Applying Theorem 4.1 to the weighted Hodge Laplacian $\Delta_{p,f}$, we have

Lemma 4.1. *Let (M^m, g) be an m -dimensional Riemannian manifold with Riemannian measure $e^{-f} dvol$ and u be a smooth function defined on M^m . For the eigenvalues $\{\lambda_i^{(p)}\}_{i=1}^\infty$ of the weighted Hodge Laplacian $\Delta_{p,f}$ (2.6) acting on p -forms, we have, $p \in \{0, 1, \dots, m\}$,*

$$\begin{aligned} & \sum_{i=1}^k (\lambda_{k+1}^{(p)} - \lambda_i^{(p)})^2 \int_{M^m} |\nabla u|^2 |\varphi_i|^2 e^{-f} dvol \\ & \leq \sum_{i=1}^k (\lambda_{k+1}^{(p)} - \lambda_i^{(p)}) \int_{M^m} \left((\Delta_{0,f} u)^2 |\varphi_i|^2 + 4 |\nabla_{\nabla u} \varphi_i|^2 \right. \\ & \quad \left. - 4 \langle \Delta_{0,f} u \varphi_i, \nabla_{\nabla u} \varphi_i \rangle \right) e^{-f} dvol \end{aligned} \quad (4.2)$$

where $\{\varphi_i\}_{i=1}^\infty$ is a corresponding orthonormal basis of p -eigenforms, i.e.

$$\int_{M^m} \langle \varphi_i, \varphi_j \rangle e^{-f} dvol = \delta_{ij}.$$

Proof. It is easy to check that $\mathcal{A} = \Delta_{p,f}$ and $\mathcal{B} = u \in C^\infty(M^m, \mathbb{R})$ satisfy the conditions in Theorem 4.1. Therefore, by the estimate of (4.1), we have

$$\begin{aligned} \sum_{i=1}^k \left(\lambda_{k+1}^{(p)} - \lambda_i^{(p)} \right)^2 \int_M \langle [\Delta_{p,f}, u] \varphi_i, u \varphi_i \rangle e^{-f} \, \text{dvol} \\ \leq \sum_{i=1}^k \left(\lambda_{k+1}^{(p)} - \lambda_i^{(p)} \right) \| [\Delta_{p,f}, u] \varphi_i \|^2, \end{aligned} \quad (4.3)$$

where

$$\| [\Delta_{p,f}, u] \varphi_i \|^2 = \int_{M^m} \langle [\Delta_{p,f}, u] \varphi_i, [\Delta_{p,f}, u] \varphi_i \rangle e^{-f} \, \text{dvol}.$$

By direct calculations, we have

$$\begin{aligned} [\Delta_{p,f}, u] \varphi_i &= [\Delta_p + \mathcal{L}_{\nabla f}, u] \varphi_i \\ &= [\Delta_p, u] \varphi_i + [\mathcal{L}_{\nabla f}, u] \varphi_i. \end{aligned} \quad (4.4)$$

From (3.1), we obtain

$$[\mathcal{L}_{\nabla f}, u] \varphi_i = g(\nabla f, \nabla u) \varphi_i. \quad (4.5)$$

By (2.7), we have

$$\begin{aligned} [\Delta_p, u] \varphi_i &= [\nabla^* \nabla, u] \varphi_i \\ &= \Delta u \varphi_i - 2 \nabla_{\nabla u} \varphi_i. \end{aligned} \quad (4.6)$$

Therefore, from (4.4) to (4.6) we get

$$[\Delta_{p,f}, u] \varphi_i = \Delta_{0,f} u \varphi_i - 2 \nabla_{\nabla u} \varphi_i \quad (4.7)$$

From (4.7), we have

$$\begin{aligned} \int_{M^m} \langle [\Delta_{p,f}, u] \varphi_i, u \varphi_i \rangle e^{-f} \, \text{dvol} \\ = \int_{M^m} \langle \Delta_{0,f} u \varphi_i - 2 \nabla_{\nabla u} \varphi_i, u \varphi_i \rangle e^{-f} \, \text{dvol}. \end{aligned}$$

By integration by parts, we have

$$\begin{aligned} 2 \int_{M^m} \langle \nabla_{\nabla u} \varphi_i, u \varphi_i \rangle e^{-f} \, \text{dvol} &= \frac{1}{2} \int_{M^m} \langle \nabla |\varphi_i|^2, \nabla u^2 \rangle e^{-f} \, \text{dvol} \\ &= \int_{M^m} (u \Delta_{0,f} u - |\nabla u|^2) |\varphi_i|^2 e^{-f} \, \text{dvol}. \end{aligned}$$

Finally, we obtain

$$\int_{M^m} \langle [\Delta_{p,f}, u] \varphi_i, u \varphi_i \rangle e^{-f} \, \text{dvol} = \int_{M^m} |\nabla u|^2 |\varphi_i|^2 e^{-f} \, \text{dvol}. \quad (4.8)$$

On the other hand, using (4.7), we get

$$\begin{aligned} \| [\Delta_{p,f}, u] \varphi_i \|^2 &= \int_{M^m} \left((\Delta_{0,f} u)^2 |\varphi_i|^2 + 4 |\nabla_{\nabla u} \varphi_i|^2 \right. \\ &\quad \left. - 4 \langle \Delta_{0,f} u \varphi_i, \nabla_{\nabla u} \varphi_i \rangle \right) e^{-f} \, \text{dvol}. \end{aligned} \quad (4.9)$$

Inserting (4.8) and (4.9) into (4.3), we obtain (4.2). \square

Proof of Theorem 1.1. Letting $f = \frac{1}{2}|x|^2$ and therefore $\Delta_{p,x} = \Delta_{p,f}$, substituting $u = x^A, A = 1, \dots, n$, the p^{th} component of the isometric immersion $x = (x^1, \dots, x^n) : M^m \rightarrow \mathbb{R}^n$ in (4.2), and taking summation on p from 1 to n , we have

$$\begin{aligned} & \sum_{A=1}^n \sum_{i=1}^k (\lambda_{k+1}^{(p)} - \lambda_i^{(p)})^2 \int_{M^m} |\nabla x^A|^2 |\varphi_i|^2 e^{-\frac{|x|^2}{2}} \text{dvol} \\ & \leq \sum_{i=1}^k \left(\lambda_{k+1}^{(p)} - \lambda_i^{(p)} \right) \int_{M^m} \sum_{A=1}^n \left((\mathfrak{L}x^A)^2 |\varphi_i|^2 + 4|\nabla_{\nabla x^A} \varphi_i|^2 \right. \\ & \quad \left. - 4\langle \mathfrak{L}x^A \varphi_i, \nabla_{\nabla x^A} \varphi_i \rangle \right) e^{-\frac{|x|^2}{2}} \text{dvol}, \end{aligned} \quad (4.10)$$

where \mathfrak{L} is the weighted Hodge Laplacian acting on functions given by (2.10). From (2.2), we obtain

$$\begin{aligned} \int_{M^m} \sum_{p=1}^n |\nabla x^A|^2 |\varphi_i|^2 e^{-\frac{|x|^2}{2}} \text{dvol} &= m \int_{M^m} |\varphi_i|^2 e^{-\frac{|x|^2}{2}} \text{dvol} \\ &= m. \end{aligned} \quad (4.11)$$

From (2.11) and (4.9), we get

$$\begin{aligned} & \sum_{A=1}^n \|\Delta_{p,x} x^A \varphi_i\|^2 \\ &= \sum_{A=1}^n \int_{M^m} \left((\mathfrak{L}x^A)^2 |\varphi_i|^2 + 4|\nabla_{\nabla x^A} \varphi_i|^2 - 4\langle \mathfrak{L}x^A \varphi_i, \nabla_{\nabla x^A} \varphi_i \rangle \right) e^{-\frac{|x|^2}{2}} \text{dvol} \\ &= \sum_{A=1}^n \int_{M^m} \left((x^A)^2 |\varphi_i|^2 + 4|\nabla_{\nabla x^A} \varphi_i|^2 - 4\langle x^A \varphi_i, \nabla_{\nabla x^A} \varphi_i \rangle \right) e^{-\frac{|x|^2}{2}} \text{dvol} \end{aligned} \quad (4.12)$$

Since M^m is a compact self-shrinker, by integration by parts and (2.3), we have

$$4 \sum_{A=1}^n \int_{M^m} \langle x^A \varphi_i, \nabla_{\nabla x^A} \varphi_i \rangle e^{-\frac{|x|^2}{2}} \text{dvol} = - \int_{M^m} 2(m - |x|^2) |\varphi_i|^2 e^{-\frac{|x|^2}{2}} \text{dvol}$$

Since $\sum_{A=1}^n |\nabla_{\nabla x^A} \varphi_i|^2 = |\nabla \varphi_i|^2$, we have

$$\begin{aligned} \sum_{A=1}^n \|\Delta_{p,x} x^A \varphi_i\|^2 &= 2m - \int_{M^m} |x|^2 |\varphi_i|^2 e^{-\frac{|x|^2}{2}} \text{dvol} \\ &\quad + 4 \int_{M^m} |\nabla \varphi_i|^2 e^{-\frac{|x|^2}{2}} \text{dvol}. \end{aligned} \quad (4.13)$$

By integration by parts, from (2.7), (3.8) and (3.9) , we have

$$\begin{aligned}
\int_{M^m} |\nabla \varphi_i|^2 e^{-\frac{|x|^2}{2}} \, d\text{vol} &= \int_{M^m} \langle \nabla'^* \nabla \varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} \, d\text{vol} \\
&= \int_{M^m} \langle (\Delta_{p,x} - \mathfrak{Ric} + \nabla(\nabla \frac{|x|^2}{2})) \varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} \, d\text{vol} \\
&= \lambda_i^{(p)} - \int_{M^m} \langle \mathfrak{Ric} \varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} \, d\text{vol} \\
&\quad + \int_{M^m} \langle \nabla(\nabla \frac{|x|^2}{2}) \varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} \, d\text{vol},
\end{aligned} \tag{4.14}$$

where ∇'^* is the adjoint operator of ∇ with respect to the Riemannian measure $e^{-\frac{|x|^2}{2}} \, d\text{vol}$. Therefore, we obtain

$$\begin{aligned}
\sum_{A=1}^n \|\Delta_{p,x} \varphi_i\|^2 &= 4\lambda_i^{(p)} + 2m \\
&\quad + 4 \int_{M^m} \langle \nabla(\nabla \frac{|x|^2}{2}) \varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} \, d\text{vol} \\
&\quad - 4 \int_{M^m} \langle \mathfrak{Ric} \varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} \, d\text{vol} \\
&\quad - \int_{M^m} |x|^2 |\varphi_i|^2 e^{-\frac{|x|^2}{2}} \, d\text{vol}.
\end{aligned} \tag{4.15}$$

From (4.3) and (4.15), we get

$$\begin{aligned}
m \sum_{i=1}^k (\lambda_{k+1}^{(p)} - \lambda_i^{(p)})^2 &\leq \sum_{i=1}^k \left(\lambda_{k+1}^{(p)} - \lambda_i^{(p)} \right) \left(4\lambda_i^{(p)} + 2m \right. \\
&\quad + 4 \int_{M^m} \langle \nabla(\nabla \frac{|x|^2}{2}) \varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} \, d\text{vol} \\
&\quad - 4 \int_{M^m} \langle \mathfrak{Ric} \varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} \, d\text{vol} \\
&\quad \left. - \int_{M^m} |x|^2 |\varphi_i|^2 e^{-\frac{|x|^2}{2}} \, d\text{vol} \right),
\end{aligned}$$

which completes the proof of Theorem 1.1. \square

Proof of Corollary 1.1. From (4.10), (3.8) and (3.9), we have

$$\begin{aligned}
 & m \sum_{i=1}^k (\lambda_{k+1}^{(p)} - \lambda_i^{(p)})^2 \\
 & \leq \sum_{i=1}^k \left(\lambda_{k+1}^{(p)} - \lambda_i^{(p)} \right) \left(4\lambda_i^{(p)} + 2m + 4 \right. \\
 & \quad + 4p \int_{M^m} |H||h|\varphi_i|^2 e^{-\frac{|x|^2}{2}} \text{dvol} \\
 & \quad \left. - \int_{M^m} |x|^2 |\varphi_i|^2 e^{-\frac{|x|^2}{2}} \text{dvol} - 4 \int_{M^m} \langle \Re i c \varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} \text{dvol} \right) \\
 & \leq 4 \sum_{i=1}^k \left(\lambda_{k+1}^{(p)} - \lambda_i^{(p)} \right) \left[\lambda_i^{(p)} + \frac{m}{2} + 1 \right. \\
 & \quad \left. + \int_{M^m} \left(p|H||h| - \Phi(H, h) - \frac{1}{4}|x|^2 \right) |\varphi_i|^2 e^{-\frac{|x|^2}{2}} \text{dvol} \right] \\
 & \leq 4 \sum_{i=1}^k \left(\lambda_{k+1}^{(p)} - \lambda_i^{(p)} \right) \left[\lambda_i^{(p)} + \frac{m}{2} + 1 \right. \\
 & \quad \left. + \max_{M^m} \left(p|H||h| - \Phi(H, h) - \frac{1}{4}|x|^2 \right) \right].
 \end{aligned}$$

□

5. GENERALIZATION OF THE LEVITIN-PARNOVSKI INEQUALITY

In this section, we will give the proof of Theorem 1.2 by similar argument in [26]. Firstly, we recall the following algebraic identity obtained by Levitin and Parnovski (see identity 2.2 of Theorem 2.2 in [30]).

Lemma 5.1. *Let \mathcal{L} and \mathcal{G} be two self-adjoint operators with domains $D_{\mathcal{L}}$ and $D_{\mathcal{G}}$ contained in a same Hilbert space and such that $G(D_{\mathcal{L}}) \subseteq D_{\mathcal{L}} \subseteq D_{\mathcal{G}}$. Let λ_j and u_j be the eigenvalues and orthonormal eigenvectors of \mathcal{L} . Then, for each j ,*

$$\sum_{k=1}^{\infty} \frac{|\langle [\mathcal{L}, \mathcal{G}] u_j, u_k \rangle|_{L^2}^2}{\lambda_k - \lambda_j} = -\frac{1}{2} \langle [[\mathcal{L}, \mathcal{G}], \mathcal{G}] u_j, u_j \rangle_{L^2} \quad (5.1)$$

(The summation is over all k and is correctly defined even when $\lambda_k = \lambda_j$ because in this case $\langle [\mathcal{L}, \mathcal{G}] u_j, u_k \rangle = 0$).

Proof of Theorem 1.2. By applying Lemma 5.1 with $\mathcal{L} = \Delta_{p,x}$ and $\mathcal{G} = x^A$, where x^A is one of the components of the isometric immersion x , we have

$$\sum_{k=1}^{\infty} \frac{|\langle [\Delta_{p,x}, x^A] u_j, u_k \rangle|_{L^2}^2}{\lambda_k - \lambda_j} = -\frac{1}{2} \langle [[\Delta_{p,x}, x^A], x^A] u_j, u_j \rangle_{L^2}. \quad (5.2)$$

From (4.7), we have

$$\begin{aligned}
[[\Delta_{p,x}, x^A], x^A]\varphi_i &= [\Delta_{p,x}, x^A](x^A\varphi_i) - x^A([\Delta_{p,x}, x^A]\varphi_i) \\
&= \mathfrak{L}(x^A)x^A\varphi_i - 2\nabla_{\nabla x^A}(x^A\varphi_i) - x^A(\mathfrak{L}x^A\varphi_i - 2\nabla_{\nabla x^A}\varphi_i) \\
&= -2\nabla_{\nabla x^A}(x^A\varphi_i) + 2x^A\nabla_{\nabla x^A}\varphi_i \\
&= -2|\nabla x^A|^2\varphi_i,
\end{aligned}$$

hence

$$-\frac{1}{2} \int_{M^m} \langle [[\Delta_{p,x}, x^A], x^A]\varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} \text{dvol} = \int_{M^m} |\nabla x^A|^2 |\varphi_i|^2 e^{-\frac{|x|^2}{2}} \text{dvol}.$$

From (5.2), we have

$$\begin{aligned}
&\int_{M^m} |\nabla x^A|^2 |\varphi_i|^2 e^{-\frac{|x|^2}{2}} \text{dvol} \\
&= \sum_{k=1}^{\infty} \frac{1}{\lambda_k^{(p)} - \lambda_i^{(p)}} \left(\int_{M^m} \langle [\Delta_{p,x}, x^A]\varphi_i, \varphi_k \rangle e^{-\frac{|x|^2}{2}} \text{dvol} \right)^2. \tag{5.3}
\end{aligned}$$

For a fixed i , from the Gram-Schmidt orthogonalization, we can find the coordinate system $\{x^A\}_{A=1}^n$ in Euclidean space \mathbb{R}^n such that the matrix

$$\left(\int_{M^m} \langle [\Delta_{p,x}, x^A]\varphi_i, \varphi_{i+k} \rangle e^{-\frac{|x|^2}{2}} \text{dvol} \right)_{1 \leq k, A \leq n}$$

is a real upper triangular matrix. That is,

$$\int_{M^m} \langle [\Delta_{p,x}, x^A]\varphi_i, \varphi_{i+k} \rangle e^{-\frac{|x|^2}{2}} \text{dvol} = 0, \quad 1 \leq k < A \leq n. \tag{5.4}$$

By (5.4), we can estimate the right hand side of (5.3) in the following

$$\begin{aligned}
&\sum_{k=1}^{\infty} \frac{1}{\lambda_k^{(p)} - \lambda_i^{(p)}} \left(\int_{M^m} \langle [\Delta_{p,x}, x^A]\varphi_i, \varphi_k \rangle e^{-\frac{|x|^2}{2}} \text{dvol} \right)^2 \\
&= \sum_{k=1}^{i-1} \frac{1}{\lambda_k^{(p)} - \lambda_i^{(p)}} \left(\int_{M^m} \langle [\Delta_{p,x}, x^A]\varphi_i, \varphi_k \rangle e^{-\frac{|x|^2}{2}} \text{dvol} \right)^2 \\
&\quad + \sum_{k=i+A}^{\infty} \frac{1}{\lambda_k^{(p)} - \lambda_i^{(p)}} \left(\int_{M^m} \langle [\Delta_{p,x}, x^A]\varphi_i, \varphi_k \rangle e^{-\frac{|x|^2}{2}} \text{dvol} \right)^2 \\
&\leq \sum_{k=i+A}^{\infty} \frac{1}{\lambda_k^{(p)} - \lambda_i^{(p)}} \left(\int_{M^m} \langle [\Delta_{p,x}, x^A]\varphi_i, \varphi_k \rangle e^{-\frac{|x|^2}{2}} \text{dvol} \right)^2 \\
&\leq \frac{1}{\lambda_{i+A}^{(p)} - \lambda_i^{(p)}} \sum_{k=1}^{\infty} \left(\int_{M^m} \langle [\Delta_{p,x}, x^A]\varphi_i, \varphi_k \rangle e^{-\frac{|x|^2}{2}} \text{dvol} \right)^2 \\
&= \frac{1}{\lambda_{i+A}^{(p)} - \lambda_i^{(p)}} \int_{M^m} |[\Delta_{p,x}, x^A]\varphi_i|^2 e^{-\frac{|x|^2}{2}} \text{dvol}
\end{aligned}$$

where Parseval's identity is used in the last equality.

Taking summation on A from 1 to n , from (5.3), (4.13), (4.14) and (2.7), we have

$$\begin{aligned}
 & \sum_{A=1}^n (\lambda_{i+A}^{(p)} - \lambda_i^{(p)}) \int_{M^m} |\nabla x^A|^2 |\varphi_i|^2 e^{-\frac{|x|^2}{2}} \, \text{dvol} \\
 & \leq \sum_{A=1}^n \int_{M^m} |[\Delta_{p,x}, x^A] \varphi_i|^2 e^{-\frac{|x|^2}{2}} \, \text{dvol} \\
 & = 2m - \int_{M^m} |x|^2 |\varphi_i|^2 e^{-\frac{|x|^2}{2}} \, \text{dvol} + 4 \int_{M^m} |\nabla \varphi_i|^2 e^{-\frac{|x|^2}{2}} \, \text{dvol} \\
 & = 4\lambda_i^{(p)} + 2m - \int_{M^m} |x|^2 |\varphi_i|^2 e^{-\frac{|x|^2}{2}} \, \text{dvol} \\
 & \quad - 4 \int_{M^m} \langle \mathfrak{Ric} \varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} \, \text{dvol} \\
 & \quad + 4 \int_{M^m} \langle \nabla(\nabla \frac{|x|^2}{2}) \varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} \, \text{dvol}.
 \end{aligned} \tag{5.5}$$

Since M^n is isometrically immersed in \mathbb{R}^n , it is easy to check

$$\sum_{A=1}^n \lambda_{i+A}^{(p)} |\nabla x^A|^2 \geq \sum_{l=1}^m \lambda_{i+l}^{(p)}. \tag{5.6}$$

Therefore, we have

$$\begin{aligned}
 \sum_{l=1}^m (\lambda_{i+l}^{(p)} - \lambda_i^{(p)}) & \leq 4\lambda_i^{(p)} + 2m - \int_{M^m} |x|^2 |\varphi_i|^2 e^{-\frac{|x|^2}{2}} \, \text{dvol} \\
 & \quad - 4 \int_{M^m} \langle \mathfrak{Ric} \varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} \, \text{dvol} \\
 & \quad + 4 \int_{M^m} \langle \nabla(\nabla \frac{|x|^2}{2}) \varphi_i, \varphi_i \rangle e^{-\frac{|x|^2}{2}} \, \text{dvol}.
 \end{aligned}$$

□

Proof of Corollary 1.3. The proof of Corollary 1.3 follows directly from (3.8) and (3.9). □

REFERENCES

- [1] N. Anghel, *Extrinsic upper bounds for eigenvalues of Dirac-type operators*, Proc. Amer. Math. Soc., 117(2):501–509, 1993.
- [2] M. S. Ashbaugh, *Isoperimetric and universal inequalities for eigenvalues*, in Spectral theory and geometry (Edinburgh, 1998), volume 273, London Math. Soc. Lecture Note Ser., pages 95–139. Cambridge Univ. Press, Cambridge, 1999.
- [3] M. S. Ashbaugh, *The universal eigenvalue bounds of Payne-Pólya-Weinberger, Hile-Protter, and H. C. Yang*, Proc. Indian Acad. Sci. Math. Sci., 112(1):3–30, 2002.
- [4] M. S. Ashbaugh, R. D. Benguria, *More bounds on eigenvalue ratios for Dirichlet Laplacians in n dimensions*, SIAM J. Math. Anal. **24** (1993), 1622–1651.
- [5] M.S. Ashbaugh and L. Hermi, *A unified approach to universal inequalities for eigenvalues of elliptic operators*, Pacific J. Math., 217(2):201–219, 2004.

- [6] E. Bueler, *The heat kernel weighted Hodge Laplacian on non compact manifolds*, Trans. A.M.S. 351, 2 (1999), 683-713.
- [7] J. J. A. M. Brands, *Bounds for the ratios of the first three membrane eigenvalues*, Arch. Rational Mech. Anal. **16** (1964), 265-268.
- [8] D.G. Chen, *Extrinsic estimates for eigenvalues of the Dirac operator*, Math. Z. 262 (2009), no. 2, 349-361.
- [9] D.G. Chen and T. Zheng, *Bounds for ratios of the membrane eigenvalues*, J. Differential Equations 250 (2011) 1575-1590.
- [10] D.G. Chen and Q.-M. Cheng, *Extrinsic estimates for eigenvalues of the Laplace operator*, J. Math. Soc. Japan 602 (2008), 325-339.
- [11] Q.-M. Cheng and H.C. Yang, *Estimates on eigenvalues of laplacian*, Math. Ann., 331:445–460, 2005.
- [12] Q. -M. Cheng and H. C. Yang, *Bounds on eigenvalues of Dirichlet Laplacian*, Math. Ann., **337** (2007), 159-175.
- [13] Q.-M. Cheng and Y. Peng, *Estimates for eigenvalues of \mathfrak{L} operator on self-shrinkers*, arXiv:1112.5938, 2011.
- [14] S.-Y. Cheng, *Eigenfunctions and eigenvalues of the Laplacian*, Part II, Amer. Math. Soc. Proc. Symp. Pure. Math., 27:185–193, 1975.
- [15] B. Colbois, *Une inégalité du type Payne-Polya-Weinberger pour le laplacien brut*, Proc. Amer. Math. Soc., 131(12):3937–3944, 2003.
- [16] T. H. Colding and W. P. Minicozzi II, *Generic mean curvature flow I; generic singularities*, Annals of Mathematics 175 (2012), 755-833.
- [17] A. El Soufi, E.M. Harrell, II, and S. Ilias. *Universal inequalities for the eigenvalues of Laplace and Schrödinger operators on Submanifolds*, Trans. Amer. Math. Soc., 361(5):2337–2350, 2009.
- [18] J. F. Grosjean, *Minimal submanifolds with a parallel or a harmonic p -form*, J. Geom. Phys. 51(2), 211-228 (2004).
- [19] E.M. Harrell II, *General bounds for the eigenvalues of Schrödinger operators*, Maximum principles and eigenvalue problems in partial differential equations (Knoxville, TN, 1987), volume 175, Pitman Res. Notes Math. Ser., pages 146–166. Longman Sci. Tech., Harlow, 1988.
- [20] E. M. Harrell, II, *Commutators, eigenvalue gaps, and mean curvature in the theory of Schrödinger operators*, Comm. Partial Differential Equations 32:3 (2007), 401-413.
- [21] E. M. Harrell, II and P. L. Michel, *Commutator bounds for eigenvalues, with applications to spectral geometry*, Comm. Partial Differential Equations 19:11-12 (1994), 2037-2055.
- [22] E.M. Harrell II and J. Stubbe, *On trace identities and universal eigenvalue estimates for some partial differential operators*, Trans. Amer. Math. Soc., 349(5):1797–1809, 1997.
- [23] E.M. Harrell II and J. Stubbe, *Universal bounds and semiclassical estimates for eigenvalues of abstract schrödinger operators*, Siam J. Math. Anal., Vol. 42, No. 5, pp. 2261-2274, 2010.
- [24] G. N. Hile and M. H. Protter, *Inequalities for eigenvalues of the Laplacian*, Indiana Univ. Math. J., 29(4):523–538, 1980.
- [25] S. Ilias and O. Makhoul, *"Universal" inequalities for the eigenvalues of the Hodge de Rham Laplacian*, Ann. Glob. Anal. Geom. (2009) 36:191-204.
- [26] S. Ilias and O. Makhoul, *A Generalization of a Levitin and Parnovski Universal Inequality for Eigenvalues*, J. Geom Anal (2012) 22:206-222.
- [27] J. M. Lee, *The gaps in the spectrum of the Laplace-Beltrami operator*, Houston J. Math., 17(1):1–24, 1991.
- [28] P.-F. Leung, *On the consecutive eigenvalues of the Laplacian of a compact minimal submanifold in a sphere*, J. Austral. Math. Soc. Ser. A, 50(3):409–416, 1991.

- [29] P.-F. Leung, *An estimate on the Ricci curvature of a submanifold and some applications*, Proc. Amer. Math. Soc. 114(4), 1051-1061 (1992).
- [30] M. Levitin and L. Parnovski, *Commutators, spectral trace identities, and universal estimates for eigenvalues*, J. Funct. Anal., 192(2):425-445, 2002.
- [31] P. Li, *Eigenvalue estimates on homogeneous manifolds*, Comment. Math. Helv., 55(3):347-363, 1980.
- [32] L. E. Payne, G. Pólya, and H. F. Weinberger, *On the ratio of consecutive eigenvalues*, J. Math. Phys., 35:289-298, 1956.
- [33] P. Petersen, *Demystifying the Weitzenböck curvature operator*, <http://www.math.ucla.edu/~petersen/BLWformulas.pdf>.
- [34] H. Sun, Q.M. Cheng, and H.C Yang, *Lower order eigenvalues of Dirichlet Laplacian*, Manuscripta Math., 125:139-156, 2008.
- [35] C. J. Thompson, *On the ratio of consecutive eigenvalues in n -dimensions*, Stud. Appl. Math., 48 (1969), 281-283.
- [36] H.C. Yang, *An estimate of the difference between consecutive eigenvalues*, preprint IC/91/60 of the Intl. Center for Theoretical Physics, Trieste, 1991(revised preprint, Academia Sinica, 1995).
- [37] P.C. Yang and S.T. Yau. *Eigenvalues of the Laplacian of compact Riemann surfaces and minimal submanifolds*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 7(1):55-63, 1980.

E-mail address: `dgchen@math.tsinghua.edu.cn`

DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING 100084,
P.R. CHINA.

E-mail address: `yiz308@lehigh.edu`

DEPARTMENT OF MATHEMATICS, LEHIGH UNIVERSITY, BETHLEHEM, PA USA 18015.