

DIFFERENTIABLE CLASSIFICATION OF 4-MANIFOLDS WITH SINGULAR RIEMANNIAN FOLIATIONS

JIANQUAN GE AND MARCO RADESCHI

ABSTRACT. In this paper, we first prove that any closed simply connected 4-manifold that admits a decomposition into two disk bundles of rank greater than 1 is diffeomorphic to one of the standard elliptic 4-manifolds: \mathbb{S}^4 , $\mathbb{C}\mathbb{P}^2$, $\mathbb{S}^2 \times \mathbb{S}^2$, or $\mathbb{C}\mathbb{P}^2 \# \pm \mathbb{C}\mathbb{P}^2$. As an application we prove that any closed simply connected 4-manifold admitting a nontrivial singular Riemannian foliation is diffeomorphic to a connected sum of copies of standard \mathbb{S}^4 , $\pm \mathbb{C}\mathbb{P}^2$ and $\mathbb{S}^2 \times \mathbb{S}^2$. A classification of singular Riemannian foliations of codimension 1 on all closed simply connected 4-manifolds is obtained as a byproduct. In particular, there are exactly 3 non-homogeneous singular Riemannian foliations of codimension 1, complementing the list of cohomogeneity one 4-manifolds.

1. INTRODUCTION

Four-dimensional manifolds form an extremely rich and interesting class of manifolds¹. This is the lowest dimension in which exotic smooth structures arise, e.g., the noncompact 4-space \mathbb{R}^4 [8, 14, 23, 49] and compact $m\mathbb{C}\mathbb{P}^2 \# n\overline{\mathbb{C}\mathbb{P}^2}$ for many pairs of $(m \geq 1, n \geq 2)$ [2, 9, 13]. Moreover, exotic smooth structures abound in this dimension and it is not known whether there is a 4-manifold with only one (standard) smooth structure, even for the simplest 4-manifold \mathbb{S}^4 , the affirmative side of which is called the *smooth Poincaré conjecture in dimension 4*.

It is thus natural to try to classify subclasses of 4-manifolds with additional structures. The study of 4-manifolds admitting smooth group actions received a lot of attention, and there is rich literature on the subject. Joining several different independent results (cf. [11, 12, 42, 43, 44]) we know that any closed simply connected 4-manifold admitting a smooth action by a compact Lie group is diffeomorphic to a connected sum of copies of standard \mathbb{S}^4 , $\pm \mathbb{C}\mathbb{P}^2$ and $\mathbb{S}^2 \times \mathbb{S}^2$. When admitting a cohomogeneity one action, the closed simply connected 4-manifold splits as a union of two disk bundles, glued along their common boundary. In this case the classification was

2000 *Mathematics Subject Classification.* 53C24, 57R30, 57R55, 57R60.

Key words and phrases. singular Riemannian foliation, isoparametric foliation, diffeomorphism, isotopy.

The first author is partially supported by the NSFC (No. 11001016 and No. 11331002), the SRFDP (No. 20100003120003), the Fundamental Research Funds for the Central Universities, and a research fellowship from the Alexander von Humboldt Foundation.

¹Throughout this paper, all manifolds considered are connected and smooth.

carried out by Parker [44] (see also [25, 26, 32]), who proved that such manifolds must be diffeomorphic to S^4 , $\mathbb{C}\mathbb{P}^2$, $S^2 \times S^2$, or $\mathbb{C}\mathbb{P}^2 \# -\mathbb{C}\mathbb{P}^2$.

The first part of this paper is concerned with a classification of closed simply connected 4-manifolds which admit a splitting structure into disk bundles but without requiring any group action.

Theorem 1.1. *Let N be a closed simply connected 4-manifold obtained by gluing two disk bundles over closed submanifolds of codimension greater than 1. Then N is diffeomorphic to one of the standard S^4 , $\mathbb{C}\mathbb{P}^2$, $S^2 \times S^2$, or $\mathbb{C}\mathbb{P}^2 \# \pm \mathbb{C}\mathbb{P}^2$.*

In [18], the first named author and Tang proved that if a homotopy 4-sphere admits some properly transnormal function, or equivalently it has a splitting structure as in Theorem 1.1, then it must be diffeomorphic to the standard S^4 . Thus, Theorem 1.1 was known when N is a homotopy (topological) 4-sphere. As an immediate application, we see that there exist no properly transnormal (isoparametric) function on any closed simply connected 4-manifold other than the five standard elliptic 4-manifolds. This should be compared to the interesting existence result of Qian and Tang [45] that every homotopy n -sphere ($n > 4$) carries a properly isoparametric function.

Recall that a singular Riemannian foliation on a Riemannian manifold is, roughly speaking, a partition of M into connected complete, injectively immersed submanifolds which stay at a constant distance from each other, and it provides a generalization of smooth actions of Lie groups. If the foliation has codimension 1, it gives rise to a splitting structure as in Theorem 1.1. Therefore, Theorem 1.1 provides a generalization of Parker's result on cohomogeneity one 4-manifolds. Moreover, we obtain

Corollary 1.2. *There are exactly 3 foliated diffeomorphism classes of non-homogeneous singular Riemannian foliations of codimension 1 in closed simply connected 4-manifolds.*

In fact, in Corollary 3.4 we will recover all cohomogeneity one actions, together with these three non-homogeneous singular Riemannian foliations of codimension 1, one of which on $\mathbb{C}\mathbb{P}^2 \# -\mathbb{C}\mathbb{P}^2$ and the other two on $\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2$ (see some further description in Subsection 3.4). Notice that $\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2$ does not admit cohomogeneity one actions, though it indeed admits metrics of non-negative curvatures as the other four cohomogeneity one 4-manifolds. This suggests the following

Conjecture 1.3. *Any closed simply connected non-negatively curved Riemannian manifold admits singular Riemannian foliations of codimension 1, under the same metric or a different bundle-like metric.*

Existence of such foliations ensures that topologically the manifold admits a splitting structure into two disk bundles as in Theorem 1.1. Hence this conjecture is essentially attributed to Karsten Grove who conjectured that every compact non-negatively

curved manifold splits as a union of two disk bundles. A positive answer to this conjecture will in particular solve affirmatively the long-standing conjecture: a closed simply connected non-negatively curved 4-manifold is diffeomorphic to one of the standard \mathbb{S}^4 , $\mathbb{C}\mathbb{P}^2$, $\mathbb{S}^2 \times \mathbb{S}^2$, or $\mathbb{C}\mathbb{P}^2 \# \pm \mathbb{C}\mathbb{P}^2$. The latter conjecture is still open even for homeomorphism.

In the second part of this paper we turn to consider singular Riemannian foliations of general codimension. At last we are able to recover and generalize the differentiable classification obtained for group actions.

Theorem 1.4. *Let N be a closed simply connected 4-manifold admitting a nontrivial singular Riemannian foliation. Then it is diffeomorphic to a connected sum of copies of standard \mathbb{S}^4 , $\pm \mathbb{C}\mathbb{P}^2$ and $\mathbb{S}^2 \times \mathbb{S}^2$.*

When the singular Riemannian foliation is closed and of dimension 1, this result has been proven by Galaz-Garcia and the second named author [16, Cor. 8.6], by showing that such a foliation comes from a smooth effective circle action which then derives the conclusion from the classification of circle actions on 4-manifolds by Fintushel [11, 12]. When the foliation is of dimension 3, Theorem 1.4 reduces to Theorem 1.1.

2. PRELIMINARIES

In this section we collect some background materials on singular Riemannian foliations, most part of which is based on the Preliminary section of [16]. We also refer the reader to [40, 4] for further results on the theory.

2.1. Singular Riemannian foliations. A *transnormal system* \mathcal{F} on a complete Riemannian manifold M is a decomposition of M into complete, injectively immersed connected submanifolds, called *leaves*, such that every geodesic emanating perpendicularly to one leaf remains perpendicular to all leaves. A *singular Riemannian foliation* is a transnormal system \mathcal{F} which is also a *singular foliation*, i.e., such that there are smooth vector fields X_i on M that span the tangent space $T_p L_p$ to the leaf L_p through each point $p \in M$. If furthermore \mathcal{F} is regular, i.e., the leaves have the same dimension, then \mathcal{F} is called a *Riemannian foliation*. If M is a smooth manifold and (M, \mathcal{F}) is a singular foliation, a metric g on M is called *bundle-like* for \mathcal{F} if (M, g, \mathcal{F}) becomes a singular Riemannian foliation. Slightly abusing notation, the pair (M, \mathcal{F}) will also denote a singular Riemannian foliation \mathcal{F} on a complete Riemannian manifold (M, g) .

We will call the quotient space M/\mathcal{F} the *leaf space*, and will also denote it by M^* . We will let $\pi : M \rightarrow M/\mathcal{F}$ be the leaf projection map. A singular Riemannian foliation \mathcal{F} will be called *closed* if all its leaves are closed in M . If \mathcal{F} is closed, then the leaf space M/\mathcal{F} is a Hausdorff metric space.

A leaf of maximal dimension is called a *regular leaf*, and its dimension is defined to be the *dimension* of \mathcal{F} , denoted by $\dim \mathcal{F}$. Leaves of lower dimensions are

called *singular* leaves. The *codimension* of \mathcal{F} is defined to be the codimension of a regular leaf. A singular Riemannian foliation (M, \mathcal{F}) of codimension one is called an *isoparametric* foliation if the regular leaves have constant mean curvature, and a *totally isoparametric* foliation if the regular leaves have constant principal curvatures in M . In fact, when (M, \mathcal{F}) gives a splitting structure as (3.1), it can become isoparametric by modifying the bundle-like metric on M (cf. [45]). A codimension one closed singular Riemannian (resp. isoparametric, totally isoparametric) foliation on a closed Riemannian manifold would be given by level sets of a transnormal (resp. isoparametric, totally isoparametric) function, which we do not introduce here but refer the reader to [18, 20, 24, 45, 50, 51].

2.2. Stratification. Let (M, \mathcal{F}) be a singular Riemannian foliation. For any point $p \in M$, we denote by L_p the leaf of \mathcal{F} through p . For $k \leq \dim \mathcal{F}$, define

$$\Sigma_k = \{ p \in M : \dim L_p = k \}.$$

Every connected component C of the set Σ_k , called a *stratum*, is an embedded (possibly non-complete) submanifold of M and the restriction of \mathcal{F} to C is a Riemannian foliation. Moreover, any horizontal geodesic (perpendicular to the leaves) tangent to Σ_k , stays in the closure of Σ_k for all time. The subset $\Sigma_{\dim \mathcal{F}}$ of regular leaves is open, dense and connected in M ; it is called the *regular stratum* of M , and it will be denoted by M_0 . All other strata have codimension at least 2 in M and are called *singular strata*.

The quotient M/\mathcal{F} inherits a stratification from M , where the strata are the projections Σ/\mathcal{F} of the strata Σ of \mathcal{F} . Any such stratum Σ/\mathcal{F} is an orbifold and in particular the regular stratum M_0/\mathcal{F} is an orbifold which is open and dense in M/\mathcal{F} .

2.3. Holonomy map. The fundamental group $\pi_1(L_p)$ acts on the normal space $\nu_p L_p$ of a regular leaf L_p , in such a way that if $x, y \in \nu_p L_p$ belong to the same $\pi_1(L_p)$ -orbit, then $\exp_p(tx), \exp_p(ty)$ belong to the same leaf, for all t . Such an action is called *holonomy map*. Fixed an ϵ -tubular neighborhood U of L_p for some small ϵ , the universal cover \tilde{U} admits a foliation $\tilde{\mathcal{F}}$ which is the lift of (U, \mathcal{F}) . One checks that \tilde{U} splits as a product $\tilde{L}_p \times D$, where D is an ϵ -disk in $\nu_p L_p$ around the origin. Therefore $U = \tilde{L}_p \times_{\pi_1(L_p)} D$, where $\pi_1(L_p)$ acts by deck transformations on the first factor, and by the holonomy map on the second. In particular, the normal bundle is orientable if and only if the holonomy map acts preserving the orientation of $\nu_p L_p$. Moreover, the holonomy group is the local group of the orbifold M_0/Σ at the point $\pi(p)$ (cf. [40, Section 3.6]).

A regular leaf L is called *principal* if the holonomy map acts trivially on the normal space $\nu_p L$ (the definition is independent on the point $p \in L$) and *exceptional* otherwise. From the local description of \mathcal{F} around L , a regular leaf is principal if and only if it projects to a manifold point of M_0/\mathcal{F} .

2.4. Infinitesimal singular Riemannian foliations. Given a point $p \in M$, let \mathbb{S}_p be the unit sphere in the normal space $\nu_p L_p$ of the leaf through p . On \mathbb{S}_p we define a foliation \mathcal{F}_p by saying that $x, y \in \mathbb{S}_p$ belong to the same leaf in \mathcal{F}_p if $\exp_p(\epsilon x)$ and $\exp_p(\epsilon y)$ belong to the same leaf of \mathcal{F} , for every $\epsilon > 0$ small enough. If \mathbb{S}_p is equipped with the round metric, the foliation $(\mathbb{S}_p, \mathcal{F}_p)$ is a singular Riemannian foliation, and it is called the *infinitesimal foliation* of \mathcal{F} at p . If p is a regular point, \mathcal{F}_p is the trivial foliation whose leaves are points.

Infinitesimal foliations are useful to understand the relation between leaves of different types, as follows. Consider a point $p \in M$, a vector $x \in \mathbb{S}_p$, and let $q = \exp_p \epsilon x$. If ϵ is small enough, there is a well defined, smooth closest-point projection $\mathfrak{p} : L_q \rightarrow L_p$, that is a locally trivial fibration. Moreover, the connected component of the fiber of \mathfrak{p} through q can be identified with the leaf $\mathcal{L}_x \in \mathcal{F}_p$ through x . There is a cover $\bar{L}_p \rightarrow L_p$ of L_p such that \mathfrak{p} lifts to a fibration $\bar{\mathfrak{p}} : L_q \rightarrow \bar{L}_p$ with connected fiber \mathcal{L}_x (just take $\bar{L}_p = \tilde{L}_p / \mathfrak{p}_*(\pi_1(L_q))$ where \tilde{L}_p is the universal cover of L_p):

$$(2.1) \quad \mathcal{L}_x \longrightarrow L_q \longrightarrow \bar{L}_p.$$

3. SINGULAR RIEMANNIAN FOLIATIONS OF CODIMENSION ONE

In this section we first prove Theorem 1.1 which is essentially the codimension 1 case of Theorem 1.4, and then give a classification of singular Riemannian foliations of codimension 1 on closed simply connected 4-manifolds in Corollary 3.4. Throughout this paper we denote diffeomorphisms and homeomorphisms by “ \cong ” and “ \simeq ” respectively.

Let N be a closed simply connected 4-manifold obtained by gluing two disk bundles over closed submanifolds M_\pm of codimension m_\pm greater than one, i.e.,

$$(3.1) \quad N = D(M_+) \bigcup_f D(M_-),$$

where $f : \partial D(M_+) \rightarrow \partial D(M_-)$ is a diffeomorphism between the boundaries of the disk bundles $D(M_\pm)$ of rank m_\pm over M_\pm . We denote the common boundary by $M := \partial D(M_+) \cong \partial D(M_-)$ and it follows from the proof of [39, Cor. 11.4 and Thm. 11.3] that M is an orientable hypersurface of N . Without loss of generality, we assume $2 \leq m_+ \leq m_- \leq 4$. As remarked in the introduction, the case when N is a 4-sphere has been solved in [18] and henceforth we assume $\dim H^2(N, \mathbb{Z}_2) = b_2 \geq 1$ for simplicity, although this case could also be derived by the same arguments.

For a splitting structure as (3.1), we have the following exact cohomology sequences ([10, 30]):

$$(3.2) \quad \cdots \rightarrow H^{i-1}(M_\pm, \mathbb{Z}_2) \xrightarrow{\theta_\pm} H^{i-m_\mp}(M_\mp, \mathbb{Z}_2) \xrightarrow{\alpha_\pm} H^i(N, \mathbb{Z}_2) \xrightarrow{i_\pm^*} H^i(M_\pm, \mathbb{Z}_2) \rightarrow \cdots$$

Since N is simply connected, by Poincaré duality we have $H^1(N, \mathbb{Z}_2) = H^3(N, \mathbb{Z}_2) = 0$.

To prove Theorem 1.1 we analyze case by case according to the value of $m_- \in \{2, 3, 4\}$ in the following subsections.

3.1. $m_- = 4$. Then $M_- = \{pt\}$ is a point in N and $M \cong \mathbb{S}^3$. By (3.2) we have

$$0 \longrightarrow H^{2-m_+}(M_+, \mathbb{Z}_2) \xrightarrow{\alpha_-} H^2(N, \mathbb{Z}_2) \xrightarrow{i_-^*} 0 \xrightarrow{\theta_-} H^{3-m_+}(M_+, \mathbb{Z}_2) \longrightarrow 0,$$

which, due to the assumption $b_2 \geq 1$, implies $m_+ = 2$, $H^1(M_+, \mathbb{Z}_2) = 0$ and $H^2(N, \mathbb{Z}_2) = H^0(M_+, \mathbb{Z}_2) = \mathbb{Z}_2$. Because 2-dimensional manifolds are determined by their cohomology structures and closed simply connected 4-manifolds are determined up to homeomorphism by the second Betti number when it is less than 2 (cf. [15]), the equalities above lead to $M_+ \cong \mathbb{S}^2 \cong \mathbb{C}\mathbb{P}^1$, $N \simeq \mathbb{C}\mathbb{P}^2$ and

$$N = D(\mathbb{C}\mathbb{P}^1) \bigcup_f D^4,$$

where $f \in \text{Diff}(\mathbb{S}^3)$ is a diffeomorphism of the common boundary \mathbb{S}^3 of the 4-disk D^4 and the 2-disk bundle $D(\mathbb{C}\mathbb{P}^1)$. Meanwhile it is known that one of the two isoparametric foliations in $\mathbb{C}\mathbb{P}^2$ (Both are homogeneous! cf. [20], etc.) splits $\mathbb{C}\mathbb{P}^2$ as

$$\mathbb{C}\mathbb{P}^2 = D(\mathbb{C}\mathbb{P}^1) \bigcup_{id} D^4,$$

where $\mathbb{C}\mathbb{P}^1 \subset \mathbb{C}\mathbb{P}^2$ is the canonical inclusion and id is the identity map of the distance sphere around the focal point of $\mathbb{C}\mathbb{P}^1$ in $\mathbb{C}\mathbb{P}^2$. Note that one still gets $\mathbb{C}\mathbb{P}^2$ if the gluing map id were replaced by an orientation reversing isometry (e.g. a reflection) since it can be radially extended to the 4-disk D^4 . Now by Cerf [7] and Hatcher [29], $f \in \text{Diff}(\mathbb{S}^3)$ is isotopic to id up to an orientation reversing isometry, and therefore using standard argument with the isotopy extension theorem (cf. [31, Thm. 2.3]) we obtain $N \cong \mathbb{C}\mathbb{P}^2$.

3.2. $m_- = 3$. Then $M_- \cong \mathbb{S}^1$ and $M \cong \mathbb{S}^1 \times \mathbb{S}^2$ since the only nontrivial 2-sphere bundle over \mathbb{S}^1 is non-orientable and thus impossible as discussed before. Using (3.2) again we get the short exact sequence

$$0 \longrightarrow \mathbb{Z}_2 \xrightarrow{\theta_-} H^{2-m_+}(M_+, \mathbb{Z}_2) \xrightarrow{\alpha_-} H^2(N, \mathbb{Z}_2) \longrightarrow 0,$$

which implies $m_+ = 2$ and $\mathbb{Z}_2 = H^0(M_+, \mathbb{Z}_2) = \mathbb{Z}_2 \oplus H^2(N, \mathbb{Z}_2)$, contradicting with the assumption $b_2 \geq 1$. This case occurs only if $N \cong \mathbb{S}^4$.

3.3. $m_- = 2$. Then $m_+ = 2$. By (3.2) we have the following exact sequence

$$0 \longrightarrow H^1(M_-, \mathbb{Z}_2) \xrightarrow{\theta_-} \mathbb{Z}_2 \xrightarrow{\alpha_-} H^2(N, \mathbb{Z}_2) \xrightarrow{i_-^*} \mathbb{Z}_2 \xrightarrow{\theta_-} H^1(M_+, \mathbb{Z}_2) \longrightarrow 0,$$

which gives

$$(3.3) \quad \begin{aligned} \mathbb{Z}_2 &= H^1(M_-, \mathbb{Z}_2) \oplus \text{Image}(\alpha_-), & \mathbb{Z}_2 &= H^1(M_+, \mathbb{Z}_2) \oplus \text{Image}(i_-^*), \\ H^2(N, \mathbb{Z}_2) &= \text{Image}(\alpha_-) \oplus \text{Image}(i_-^*). \end{aligned}$$

These equations show that $\dim H^2(N, \mathbb{Z}_2) = b_2 \leq 2$ and hence $b_2 = 1$ or 2 .

When $b_2 = 1$, $N \simeq \mathbb{C}\mathbb{P}^2$ (cf. [15]) and either $\text{Image}(\alpha_-) = 0$, $\text{Image}(i_-^*) = \mathbb{Z}_2$ or $\text{Image}(\alpha_-) = \mathbb{Z}_2$, $\text{Image}(i_-^*) = 0$, correspondingly either $H^1(M_-, \mathbb{Z}_2) = \mathbb{Z}_2$, $H^1(M_+, \mathbb{Z}_2) = 0$ or $H^1(M_-, \mathbb{Z}_2) = 0$, $H^1(M_+, \mathbb{Z}_2) = \mathbb{Z}_2$. Therefore M_\pm are \mathbb{S}^2 and $\mathbb{R}\mathbb{P}^2$. It is well known (cf. [47]) that any orientable circle bundle with Euler number m over \mathbb{S}^2 is a lens space $L(m, 1)^2$ which is a quotient of \mathbb{S}^3 by some \mathbb{Z}_m -action. Recall (cf. [41]) that the orientable³ circle bundle over $\mathbb{R}\mathbb{P}^2$ with Euler number e , denoted by $(On1 | e)$, is two-fold covered by the orientable circle bundle over \mathbb{S}^2 with Euler number $2e$, denoted by $(Oo0 | 2e)$. Moreover, $(On1 | 1) \cong (Oo0 | 4) \cong L(4, 1)$. Thus $(On1 | e)$ is the quotient of \mathbb{S}^3 (viewed as the unit quaternions) by the subgroup Q_{4e} generated by $\omega = \cos(\pi/|e|) + i \sin(\pi/|e|)$ and j (cf. [34]). Note that Q_{4e} is a binary dihedral group which is abelian only if $e = 1$. It follows that the common boundary M , as circle bundles over both \mathbb{S}^2 and $\mathbb{R}\mathbb{P}^2$, can only be $\mathbb{S}^3/Q_4 = L(4, 1)$.

In conclusion, we have proved

$$N = D(\mathbb{S}^2) \bigcup_f D(\mathbb{R}\mathbb{P}^2),$$

where $f \in \text{Diff}(L(4, 1))$ is a diffeomorphism of the common boundary $M \cong L(4, 1)$. On the other hand, it is known that the other (homogeneous) isoparametric foliation on $\mathbb{C}\mathbb{P}^2$ decomposes it as ([20], [48])

$$\mathbb{C}\mathbb{P}^2 = D(\mathbb{Q}^1) \bigcup_{id} D(\mathbb{R}\mathbb{P}^2),$$

where $\mathbb{Q}^1 \cong \mathbb{S}^2$ is the standard complex quadric in $\mathbb{C}\mathbb{P}^2$ and id is the identity on $M = \partial D(\mathbb{Q}^1) = \partial D(\mathbb{R}\mathbb{P}^2) = L(4, 1)$. Due to the celebrated work on the (generalized) Smale Conjecture of Hong, et al. [33], the inclusion of the isometry group $\text{Isom}(L(m, q)) \rightarrow \text{Diff}(L(m, q))$ of any lens space $L(m, q)$ ($m \geq 3$) is a homotopy equivalence. As a result, $f \in \text{Diff}(L(4, 1))$ is isotopic to the identity id or an orientation-preserving isometry $f_0 \in \text{Isom}(L(4, 1)) = O(2) \times SO(3)$ which preserves the fibration of $L(4, 1)$ over \mathbb{S}^2 and hence is radially extendable to an orientation-preserving diffeomorphism \tilde{f}_0 of the corresponding disk bundle $D(\mathbb{S}^2)$. This derives

$$(3.4) \quad N = D(\mathbb{S}^2) \bigcup_f D(\mathbb{R}\mathbb{P}^2) \cong D(\mathbb{S}^2) \bigcup_{id \text{ or } f_0} D(\mathbb{R}\mathbb{P}^2) \cong D(\mathbb{Q}^1) \bigcup_{id} D(\mathbb{R}\mathbb{P}^2) = \mathbb{C}\mathbb{P}^2,$$

where the second diffeomorphism with respect to f_0 comes from gluing $\tilde{f}_0 : D(\mathbb{S}^2) \rightarrow D(\mathbb{Q}^1)$ and the identity $id : D(\mathbb{R}\mathbb{P}^2) \rightarrow D(\mathbb{R}\mathbb{P}^2)$ (cf. [31, Thm. 2.2]).

Remark 3.1. *In fact, here and later on, it is sufficient to use only the “ π_0 -part” of the Smale conjecture: $\pi_0(\text{Isom}(M)) \rightarrow \pi_0(\text{Diff}(M))$ is an isomorphism induced from the natural inclusion of $\text{Isom}(M)$ in $\text{Diff}(M)$. This part has been confirmed for every elliptic 3-manifold (cf. [7], [38] and references therein). In particular, $\pi_0(\text{Diff}(L(m, 1))) \cong$*

²Here we ignore the specific orientations and identify $L(\pm m, 1)$.

³Here orientability refers to that of the total space.

$\pi_0(\text{Isom}(L(m,1))) \cong C_2$ is a cyclic group of order 2 for each $m \geq 1$. Moreover, it consists of (path components of) the identity and an orientation-reversing isometry f_1 if $m = 1$ or 2, and an orientation-preserving isometry f_0 , which preserves the fibration of $L(m,1)$ over \mathbb{S}^2 and hence is radially extendable to an orientation-preserving diffeomorphism \tilde{f}_0 of the corresponding disk bundle $D(\mathbb{S}^2)$, if $m \geq 3$.

Now we turn to deal with the case when $b_2 = 2$, i.e., $N \simeq \mathbb{S}^2 \times \mathbb{S}^2$, or $\mathbb{C}\mathbb{P}^2 \# \pm \mathbb{C}\mathbb{P}^2$ (cf. [15]). By (3.3) we have $H^1(M_\pm, \mathbb{Z}_2) = 0$ and thus $M_+ \cong M_- \cong \mathbb{S}^2$, M is a lens space $L(m,1)$ for some $m \geq 0$. These give a splitting structure on N as

$$(3.5) \quad N = D(\mathbb{S}^2) \bigcup_f D(\mathbb{S}^2),$$

where $f \in \text{Diff}(L(m,1))$ is a gluing diffeomorphism on the common boundary $M \cong L(m,1)$ for some $m \geq 0$. Note that m is just the self intersection number of either \mathbb{S}^2 in N or equivalently the Euler number of the circle bundles $L(m,1) \rightarrow \mathbb{S}^2$. The proof of this case (and hence of the total Theorem 1.1) will be completed by the following.

Proposition 3.2. *Let N be a closed (simply connected) 4-manifold admitting a splitting structure as (3.5). Then N is diffeomorphic to $\mathbb{S}^2 \times \mathbb{S}^2$, or $\mathbb{C}\mathbb{P}^2 \# \pm \mathbb{C}\mathbb{P}^2$.*

Remark 3.3. *The classification can be explicitly described as follows. Without loss of generality, we suppose the splitting structure (3.5) is given by gluing two copies of an oriented disk bundle $D_m(\mathbb{S}^2)$ of opposite orientations over \mathbb{S}^2 through a diffeomorphism $f : L(m,1) \rightarrow L(m,1)$ of the boundary with induced orientation from $D_m(\mathbb{S}^2)$ for some integer $m \geq 0$. Then N is diffeomorphic to*

- (1) $\mathbb{S}^2 \times \mathbb{S}^2$ if $m > 2$ is even, or $m = 2$ and $[f] = [id] \in \pi_0(\text{Diff}(\mathbb{R}\mathbb{P}^3)) \cong C_2$ (orientation-preserving), or $m = 0$ and $[f] \in \pi_0(\text{Isom}(\mathbb{S}^1 \times \mathbb{S}^2)) \cong C_2 \times C_2 \subset \pi_0(\text{Diff}(\mathbb{S}^1 \times \mathbb{S}^2)) \cong C_2 \times C_2 \times C_2$ (isotopic to an isometry);
- (2) $\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2$ if $m = 2$ and $[id] \neq [f] \in \pi_0(\text{Diff}(\mathbb{R}\mathbb{P}^3)) \cong C_2$ (orientation-reversing), or $m = 1$ and $[id] \neq [f] \in \pi_0(\text{Diff}(\mathbb{S}^3)) \cong C_2$ (orientation-reversing);
- (3) $\mathbb{C}\mathbb{P}^2 \# -\mathbb{C}\mathbb{P}^2$ if $m > 2$ is odd, or $m = 1$ and $[f] = [id] \in \pi_0(\text{Diff}(\mathbb{S}^3)) \cong C_2$ (orientation-preserving), or $m = 0$ and $[f] \in \pi_0(\text{Diff}(\mathbb{S}^1 \times \mathbb{S}^2)) - \pi_0(\text{Isom}(\mathbb{S}^1 \times \mathbb{S}^2))$ (not isotopic to an isometry).

Proof. It follows from van Kampen theorem that N is simply connected and thus orientable, implying the splitting structure (3.5) has the form as described in the remark above. For $m > 2$, $f \in \text{Diff}(L(m,1))$ is isotopic to the identity id or a fiber-preserving orientation-preserving isometry f_0 which can be extended to an orientation-preserving diffeomorphism \tilde{f}_0 of the disk bundle $D_m(\mathbb{S}^2)$ (see Remark 3.1), both giving rise to the same double manifold $N \cong D_m(\mathbb{S}^2) \bigcup_{id} -D_m(\mathbb{S}^2)$ as in (3.4). It is easily seen from [47] that this double manifold is an oriented \mathbb{S}^2 bundle over \mathbb{S}^2 and $m \in \mathbb{Z} \cong \pi_1 SO(2)$ determines a reduction of the structural group $SO(3)$ to $SO(2)$ of the 2-sphere bundle.

Note that $\pi_1 SO(3) \cong \mathbb{Z}_2$, thus each even m corresponds to the trivial bundle and each odd m corresponds to the only nontrivial bundle, the total space of which is shown to be diffeomorphic to $\mathbb{C}\mathbb{P}^2 \# -\mathbb{C}\mathbb{P}^2$ in [47]. Hence, N is $\mathbb{S}^2 \times \mathbb{S}^2$ if m is even, $\mathbb{C}\mathbb{P}^2 \# -\mathbb{C}\mathbb{P}^2$ if m is odd. The same argument holds for the cases when $m = 1, 2$ and f is orientation-preserving, or when $m = 0$ and f is isotopic to an isometry, since each isometry in this case can be radially extended to the trivial disk bundle $\mathbb{S}^2 \times D^2$ (see the π_0 -part or the homotopy type of $\text{Diff}(\mathbb{S}^1 \times \mathbb{S}^2)$ in [22], [28]).

When $m = 2$ (resp. 1) and f is isotopic to an orientation-reversing isometry (all in one path component) in $\text{Isom}(\mathbb{R}\mathbb{P}^3)$ (resp. $\text{Isom}(\mathbb{S}^3)$), we would first get a homeomorphism $N \simeq \mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2$ by checking the second Betti number $b_2 = 2$ and the signature $\sigma = 2$ using the Novikov's additivity theorem. As there is only one path component for the gluing diffeomorphism f , i.e., all gluing diffeomorphisms are isotopic to each other, using the standard gluing argument as in (3.4) would deduce a diffeomorphism once we establish the same splitting structure on $\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2$ for $m = 2, 1$ respectively. Therefore it is sufficient to check the existence of the following splitting structures

$$\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2 = D_2(\mathbb{S}^2) \bigcup_{id} D_2(\mathbb{S}^2) = D_1(\mathbb{S}^2) \bigcup_{id} D_1(\mathbb{S}^2),$$

where id denotes the identity on $\partial D_2(\mathbb{S}^2) = \mathbb{R}\mathbb{P}^3$ and $\partial D_1(\mathbb{S}^2) = \mathbb{S}^3$ respectively. The existence can be confirmed as follows. Consider the two embeddings of \mathbb{S}^2 in $\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2$ as two connected sums $\mathbb{C}\mathbb{P}^1 \# \mathbb{C}\mathbb{P}^1$ via different embeddings of $\mathbb{C}\mathbb{P}^1 \subset \mathbb{C}\mathbb{P}^2$, e.g., defined by the vanishing of different coordinate functions. The neighborhoods of these two embeddings will give the splitting structure for the case $m = 2$. Considering the standard embeddings of $\mathbb{C}\mathbb{P}^1$ in the two copies of $\mathbb{C}\mathbb{P}^2$ in $\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2$ will give rise to the splitting structure for the case $m = 1$.

The last case left to analyze is when $m = 0$ and f is not isotopic to an isometry, i.e., it belongs to the four "rotation" path components of $\text{Diff}(\mathbb{S}^1 \times \mathbb{S}^2)$ given by compositions of isometries of $\mathbb{S}^1 \times \mathbb{S}^2$ with a "rotation" (generator) diffeomorphism (cf. [22])

$$(3.6) \quad \tau(z, w) = (z, \phi(z)w)$$

where $\phi(z)w = (z \cdot (w_1, w_2), w_3)$ is a rotation of $w = (w_1, w_2, w_3) \in \mathbb{S}^2$ around the w_3 -axis through the angle of $z \in \mathbb{S}^1$. We can finally produce only one 4-manifold from these four isotopy classes of gluing diffeomorphisms since as mentioned before, all isometries of $\mathbb{S}^1 \times \mathbb{S}^2$ can be radially extended to diffeomorphisms of the trivial disk bundle $D_0(\mathbb{S}^2) = \mathbb{S}^2 \times D^2$ and consequently their compositions with τ as gluing maps give rise to the same 4-manifold as that by τ itself. It thus suffices to prove

$$(3.7) \quad N = \mathbb{S}^2 \times D^2 \bigcup_{\tau} -(\mathbb{S}^2 \times D^2) \cong \mathbb{C}\mathbb{P}^2 \# -\mathbb{C}\mathbb{P}^2.$$

Now we regard N as given by first cutting out the sphere-factors along their equators, then gluing the disk-factors pointwise along the north and south hemispheres, at last regluing these two hemispheres back using the transformation map $\tau_0 : \mathbb{S}^1 \rightarrow SO(2)$

induced from τ . Then it turns out to be an oriented \mathbb{S}^2 bundle over \mathbb{S}^2 with the reduced structural group $SO(2) \subset SO(3)$ corresponding to the element $[\tau_0] = 1 \in \mathbb{Z} \cong \pi_1 SO(2)$. Hence $N \cong \mathbb{C}\mathbb{P}^2 \# -\mathbb{C}\mathbb{P}^2$ as before (cf. [47]).

The proof is now complete. \square

As a corollary, we conclude the following differential classification of singular Riemannian foliations of codimension one on closed simply connected 4-manifolds. Note that by the result of [45] introduced in subsection 2.1, these singular Riemannian foliations can become isoparametric with only the bundle-like Riemannian metrics on the 4-manifolds modified.

Corollary 3.4. *Let N be a closed simply connected 4-manifold admitting a singular Riemannian foliation \mathcal{F} of codimension one with regular leaf M and two singular leaves M_{\pm} . Then the following Table 1 gives a foliated diffeomorphism classification of (N, \mathcal{F}) with \mathcal{F} in terms of (M, M_{\pm}) :*

TABLE 1. SRF of codim 1 on closed simply connected 4-manifolds

N	\mathcal{F}		<i>Properties</i>		
	M	M_{\pm}	<i>Homog</i>	<i>T-Isopar</i>	<i>Isopar</i>
\mathbb{S}^4	$L(1, 1)$	pt, pt	<i>Yes</i>	<i>Yes</i>	<i>Yes</i>
	$L(0, 1)$	$\mathbb{S}^1, \mathbb{S}^2$			
	$SO(3)/(\mathbb{Z}_2 \oplus \mathbb{Z}_2)$	$\mathbb{R}\mathbb{P}^2, \mathbb{R}\mathbb{P}^2$			
$\mathbb{C}\mathbb{P}^2$	$L(1, 1)$	pt, \mathbb{S}^2	<i>No</i>	<i>Yes</i>	<i>Yes</i>
	$L(4, 1)$	$\mathbb{R}\mathbb{P}^2, \mathbb{S}^2$			
$\mathbb{S}^2 \times \mathbb{S}^2$	$L(2m, 1), m \geq 0$	$\mathbb{S}^2, \mathbb{S}^2$	<i>Yes</i>	<i>Unknown</i>	
$\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2$	$L(1, 1)$	$\mathbb{S}^2, \mathbb{S}^2$	<i>No</i>	<i>Unknown</i>	
	$L(2, 1)$	$\mathbb{S}^2, \mathbb{S}^2$			
$\mathbb{C}\mathbb{P}^2 \# -\mathbb{C}\mathbb{P}^2$	$L(2m + 1, 1), m \geq 0$	$\mathbb{S}^2, \mathbb{S}^2$	<i>Yes</i>	<i>Unknown</i>	
	$L(0, 1)$	$\mathbb{S}^2, \mathbb{S}^2$	<i>No</i>		

where the column “Homog” (resp. “T-Isopar”, “Isopar”) means whether there exist a homogeneous (resp. totally isoparametric, isoparametric) representative in the foliated diffeomorphism class.

Proof. The classification is clear by the proof of Theorem 1.1 and that of [18, Thm. 1.1] for \mathbb{S}^4 , where the equivalence of the foliations with the same (M, M_{\pm}) follows from the consideration on isotopy classes of the gluing diffeomorphisms and the observation that all possible extensions of the gluing diffeomorphisms can be made radially (so to preserve the leaves). As for the homogeneity, one can compare with the classification of cohomogeneity one 4-manifolds in [44]⁴. Naturally, the 7 homogeneous foliations are

⁴In [44] the second case in $\mathbb{C}\mathbb{P}^2$ was missing, as remarked also in [26, 32]. Note also that it should be $\mathbb{C}\mathbb{P}^2 \# -\mathbb{C}\mathbb{P}^2$ other than $\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2$, as the covering space of the manifolds 43 and 49 in Parker’s list.

totally isoparametric. The 2 nonhomogeneous foliation classes on $\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2$ can also be represented by totally isoparametric foliations as follows. Fix an invariant metric \tilde{g}_1 for the homogeneous foliation $(\mathbb{C}\mathbb{P}^2 \# - \mathbb{C}\mathbb{P}^2, \mathcal{F}' = (L(1, 1), \mathbb{S}^2, \mathbb{S}^2))$. This induces a singular Riemannian foliation representing $(\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2, \mathcal{F}_1 = (L(1, 1), \mathbb{S}^2, \mathbb{S}^2))$, by gluing the homogeneous foliations of the two disk bundles $D_1(\mathbb{S}^2)$ in $(\mathbb{C}\mathbb{P}^2 \# - \mathbb{C}\mathbb{P}^2, \mathcal{F}')$ through an orientation-reversing isometry $f_1 \in \text{Isom}(\mathbb{S}^3)$ (with respect to the metric induced from \tilde{g}_1), with the bundle-like metric g_1 on $(\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2, \mathcal{F}_1)$ also glued from the restrictions of \tilde{g}_1 to the two disk bundles $D_1(\mathbb{S}^2)$ by the isometry f_1 . The metric g_1 is smooth and well-defined because the gluing map f_1 is an isometry with respect to $\tilde{g}_1|_{\partial D_1(\mathbb{S}^2)}$. It follows that each regular leaf $L(1, 1)$ of \mathcal{F}_1 has constant principal curvatures and thus $(\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2, g_1, \mathcal{F}_1)$ is totally isoparametric. The same argument is applicable to the case $(\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2, \mathcal{F}_2 = (L(2, 1), \mathbb{S}^2, \mathbb{S}^2))$ with the homogeneous foliation $(\mathbb{S}^2 \times \mathbb{S}^2, \mathcal{F}'' = (L(2, 1), \mathbb{S}^2, \mathbb{S}^2))$ taking place of $(\mathbb{C}\mathbb{P}^2 \# - \mathbb{C}\mathbb{P}^2, \mathcal{F}')$. The last nonhomogeneous foliation class $(\mathbb{C}\mathbb{P}^2 \# - \mathbb{C}\mathbb{P}^2, \mathcal{F} = (L(0, 1), \mathbb{S}^2, \mathbb{S}^2))$ can be represented by an isoparametric foliation, either by the result of [45] cited above, or simply using the isoparametric foliation obtained from the pull-back of the standard isoparametric foliation on \mathbb{S}^2 through a Riemannian submersion $\mathbb{C}\mathbb{P}^2 \# - \mathbb{C}\mathbb{P}^2 \rightarrow \mathbb{S}^2$ with totally geodesic \mathbb{S}^2 -fibres (cf. [18]). We remark that one can not use directly the argument above for $(\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2, \mathcal{F}_1)$ to obtain a totally isoparametric representative in this case, since the gluing map now does not belong to the isometry group of $\mathbb{S}^1 \times \mathbb{S}^2$. \square

Note that we have found two nonhomogeneous examples of totally isoparametric foliations on $\mathbb{C}\mathbb{P}^2 \# \mathbb{C}\mathbb{P}^2$ while such foliations were guessed to be homogeneous (cf. [20]). However, we do not know whether there exists any totally isoparametric representative in the last nonhomogeneous foliation class $(\mathbb{C}\mathbb{P}^2 \# - \mathbb{C}\mathbb{P}^2, \mathcal{F} = (L(0, 1), \mathbb{S}^2, \mathbb{S}^2))$. Observing that now \mathbb{S}^2 represents the homology class $\pm(1, -1) \in H_2(\mathbb{C}\mathbb{P}^2 \# - \mathbb{C}\mathbb{P}^2)$, we see that this question relates with the *homology representation (minimal genus problem)* (cf. [35, 36]) with further geometric restrictions: the complement of the embedding sphere is an open disk bundle over another embedding sphere, and every tubular hypersurface of either embedding sphere has constant principal curvatures under some Riemannian metric on $\mathbb{C}\mathbb{P}^2 \# - \mathbb{C}\mathbb{P}^2$. Note moreover that the latter curvature condition forces both embedding spheres to have constant principal curvatures which are independent of the choice of unit normal vectors (cf. [19]). Nevertheless, the answer to this question seems to be negative in view of the following observation motivated by the pull-back construction in the proof above.

Proposition 3.5. *There exists no totally isoparametric foliation $\mathcal{F} = (\mathbb{S}^1 \times \mathbb{S}^2, \mathbb{S}^2, \mathbb{S}^2)$ on $\mathbb{C}\mathbb{P}^2 \# - \mathbb{C}\mathbb{P}^2$ that can be projected to a singular Riemannian foliation on \mathbb{S}^2 via a Riemannian submersion $\pi : \mathbb{C}\mathbb{P}^2 \# - \mathbb{C}\mathbb{P}^2 \rightarrow \mathbb{S}^2$ with totally geodesic fibres.*

Proof. Let $\pi : (N = \mathbb{C}\mathbb{P}^2 \# - \mathbb{C}\mathbb{P}^2, g) \rightarrow (\mathbb{S}^2, g')$ be a Riemannian submersion with totally geodesic fibers, and let $\mathcal{F}' = (\mathbb{S}^1, pt, pt)$ be a codimension 1 singular Riemannian

foliation on (\mathbb{S}^2, g') . The pull-back foliation $\mathcal{F} = \pi^{-1}(\mathcal{F}')$, obtained by the preimages of the leaves in \mathcal{F}' , is a singular Riemannian foliation on (N, g) , and suppose that any regular leaf $L = \pi^{-1}(\mathbb{S}^1) \cong \mathbb{S}^1 \times \mathbb{S}^2$ of \mathcal{F} has constant principal curvatures in N .

As mentioned in Subsection 2.1, the foliations $\mathcal{F}, \mathcal{F}'$ consist of level sets of some (non-unique) functions F, f on N, \mathbb{S}^2 , respectively, where $F = f \circ \pi$ is totally isoparametric on (N, g) and f is transnormal on (\mathbb{S}^2, g') . In fact, for our purpose it suffices to let F, f be the distance functions to one of the singular leaves, which are well-defined and smooth on the regular parts $N_0 = N - (\mathbb{S}^2 \sqcup \mathbb{S}^2) = \pi^{-1}(N'_0)$, $N'_0 = \mathbb{S}^2 - (pt \sqcup pt)$ of the foliations $\mathcal{F}, \mathcal{F}'$. By the relation between the shape operator S on L with respect to the unit normal vector field $\xi := \nabla F/|\nabla F|$ (normalized gradient) and the Hessian H_F of F on (N, g) , namely

$$g(S(X), Y) = -H_F(X, Y)/|\nabla F|, \quad \text{for } X, Y \in TL,$$

\mathcal{F} is totally isoparametric if and only if F is transnormal, i.e., $|\nabla F|^2 = b(F)$ for some function $b: \mathbb{R} \rightarrow \mathbb{R}$, and $H_F|_{TL}$ has constant eigenvalues on any regular leaf L .

Let $L' \cong \mathbb{S}^1$ be any regular leaf in $(\mathbb{S}^2, g', \mathcal{F}')$ and $L = \pi^{-1}(L')$. Let e be the basic horizontal unit vector field on L that projects to the (oriented) unit vector field on L' , and let $\{u, v\}$ be a local vertical orthonormal frame on L , such that $\{e, u, v\}$ is a local orthonormal frame on L . Notice that the basic horizontal vector fields e, ξ are well-defined on the whole of the regular part N_0 . Recall that the O'Neill's integrability tensor A acts skew-symmetrically on the horizontal orthonormal frame $\{e, \xi\}$ as

$$A_e \xi = g(\nabla_e \xi, u)u + g(\nabla_e \xi, v)v,$$

where ∇ denotes the covariant derivative on (N, g) . This tensor measures the obstruction to integrability of the horizontal distribution. When the fibres are totally geodesic, $A \equiv 0$ if and only if the total space N is locally a Riemannian product of the base manifold and the fiber (see more details and properties of the A -tensor in, e.g., [17]). In particular, in our case the A -tensor cannot vanish identically on N since otherwise N would have to split isometrically as $\mathbb{S}^2 \times \mathbb{S}^2$.

Straightforward calculations show that under the orthonormal frame $\{e, u, v\}$,

$$H_F|_{TL} = \begin{pmatrix} H_F(e, e) & |\nabla F|g(\nabla_e \xi, u) & |\nabla F|g(\nabla_e \xi, v) \\ |\nabla F|g(\nabla_e \xi, u) & 0 & 0 \\ |\nabla F|g(\nabla_e \xi, v) & 0 & 0 \end{pmatrix}.$$

Therefore, $H_F|_{TL}$ has constant eigenvalues if and only if $H_F(e, e) = \Delta f \circ \pi$ is constant (i.e., f is isoparametric) and $|\nabla F|^2 |A_e \xi|^2 = |\nabla F|^2 (g(\nabla_e \xi, u)^2 + g(\nabla_e \xi, v)^2)$ is constant on L . Then $|A_e \xi|^2$ is constant on L because $|\nabla F|^2 = b(F) \neq 0$ is constant on L . Noticing that $A_e \xi$ is now a global tangent vector field of constant length along each \mathbb{S}^2 -fibre in L , we conclude that $A \equiv 0$ on L and hence on N_0 . By continuity $A \equiv 0$ on $N = \overline{N_0}$ which gives the contradiction as described in the preceding paragraph. \square

3.4. Further remark on the 3 non-homogeneous foliations. To conclude this section, we would like to remark further on the exactly 3 non-homogeneous singular Riemannian foliations of codimension one in Table 1 of Corollary 3.4. In fact, these non-homogeneous foliations all arise by somewhat “twisted” gluing of two copies of a homogeneous foliation on a disk bundle.

Explicitly, $(\mathbb{C}\mathbb{P}^2\# - \mathbb{C}\mathbb{P}^2, \mathcal{F} = (L(0, 1) = \mathbb{S}^1 \times \mathbb{S}^2, \mathbb{S}^2, \mathbb{S}^2))$ can be regarded as the pull-back of the homogeneous foliation of \mathbb{S}^2 by concentric circles, via the \mathbb{S}^2 -bundle projection $\mathbb{C}\mathbb{P}^2\# - \mathbb{C}\mathbb{P}^2 \rightarrow \mathbb{S}^2$ as in Proposition 3.5. This can thus also be seen as a twisted gluing of the homogeneous foliations on the two trivial \mathbb{S}^2 -bundles over the south and north hemispheres, i.e., on the trivial disk bundle $\mathbb{S}^2 \times D^2$. The twisted gluing diffeomorphism is nothing but the non-isometric “rotation” τ in (3.6) (under appropriate choices of orientations of the trivial disk bundles and possibly some isotopy), which proves directly the diffeomorphism (3.7) and moreover, implies the “failure” (Proposition 3.5) of gluing two copies of the homogeneous foliation into a totally isoparametric foliation since τ is not isometric. This also gives an interesting example of non-homogeneous foliation (M, \mathcal{F}) that projects, via a foliated map $M \rightarrow N$, to a homogeneous foliation on N . Conversely, examples of homogeneous foliations that project to non-homogeneous foliations were rather well-known on the Hopf fibrations $\mathbb{S}^{2n+1} \rightarrow \mathbb{C}\mathbb{P}^n$ (see for example [20]).

This “twisted” gluing phenomenon occurs similarly for the other two non-homogeneous foliations on $\mathbb{C}\mathbb{P}^2\#\mathbb{C}\mathbb{P}^2$, but with isometric twisted (orientation-reversing) gluing diffeomorphisms instead, which ensures the success of the gluing into totally isoparametric foliations. One is glued from two copies of the (unique) homogeneous foliation on $D_1(\mathbb{S}^2)$, the disk bundle over \mathbb{S}^2 with Euler number 1 along the boundary $L(1, 1) = \mathbb{S}^3$. The other is glued from two copies of the (unique) homogeneous foliation on $D_2(\mathbb{S}^2)$, the disk bundle over \mathbb{S}^2 with Euler number 2 along the boundary $L(2, 1) = \mathbb{R}\mathbb{P}^3$.

Notice that there are neither non-isometric, nor orientation-reversing isotopy classes of diffeomorphisms on the lens spaces $L(m, 1)$ for $m > 2$. This explains somewhat why there are only these 3 “twisted” gluing cases and hence non-homogeneous classes.

4. SINGULAR RIEMANNIAN FOLIATIONS OF GENERAL CODIMENSION

In this section we first prove a conjecture of Molino for 4-manifolds which reduces the objects to closed singular Riemannian foliations. Then we prove Theorem 1.4 by verifying the only remaining case of closed 2-dimensional singular Riemannian foliations.

Let (M, \mathcal{F}) be a singular Riemannian foliation. Recall that $\overline{\mathcal{F}}$ is defined as the partition of M given by the closures of the leaves in \mathcal{F} . It is a transnormal system, i.e. the leaves are locally at a constant distance from each other. Moreover, Molino [40] proved that in the regular part of \mathcal{F} , $\overline{\mathcal{F}}$ is a singular Riemannian foliation, and he

conjectured that $\overline{\mathcal{F}}$ is actually a singular Riemannian foliation on the whole of M . In the following proposition, we show that Molino's conjecture holds for 4-manifolds, and therefore $\overline{\mathcal{F}}$ is a closed singular Riemannian foliation.

Proposition 4.1. *Molino's conjecture holds for 4-manifolds.*

Proof. If $\dim \mathcal{F} = 2, 3$, it is easily seen that for every point $p \in M$, the infinitesimal foliation $(\mathbb{S}_p, \mathcal{F}_p)$ either consists of points (if p is regular), or it consists of one leaf, or it is a foliation of codimension one. All these foliations are polar, i.e., the leaf space $\mathbb{S}_p/\mathcal{F}_p$ is isometric to a Riemannian orbifold, and this makes \mathcal{F} *infinitesimally polar*. Molino's conjecture is known to hold for such foliations [5] and thus one only needs to prove it for 1-dimensional singular Riemannian foliations.

We need to prove that there is a family of smooth vector fields $\{X_i\}$ such that, for each point $p \in M$, the tangent space of the leaf $\overline{L_p}$ of $\overline{\mathcal{F}}$ through p is the span of the vectors $\{X_i(p)\}$. Notice that this is a local condition. Moreover, Molino himself proved that this condition is satisfied around regular points of \mathcal{F} , so we only have to prove that the condition holds around the singular leaves of \mathcal{F} .

Since $\dim \mathcal{F} = 1$, the singular leaves are just points and in particular they are closed. Moreover, a metric ball around each singular leaf is foliated diffeomorphic to the orbit decomposition of a representation $\mathbb{R} \rightarrow O(4)$. The closure of such actions is well known to be homogeneous, and more precisely given by the action of a torus $T^2 \rightarrow O(4)$. In particular, the closure of \mathcal{F} around the singular leaves of \mathcal{F} is a singular foliation, which is what we wanted to prove. \square

From the proposition above, if a 4-manifold M admits a singular Riemannian foliation \mathcal{F} then it also admits a closed singular Riemannian foliation $\overline{\mathcal{F}}$. Moreover, if M is simply connected and closed, its Euler characteristic is positive and by [21] there is a compact leaf. In particular, the leaves of \mathcal{F} cannot be dense in M , and $\overline{\mathcal{F}}$ is a nontrivial closed foliation.

In what follows, we will assume that (M, \mathcal{F}) is a closed, nontrivial singular Riemannian foliation on a closed simply connected 4-manifold. As introduced in Section 1, Theorem 1.4 has been proven by Galaz-Garcia and the second named author [18, Cor. 8.6] for closed foliations of dimension 1, by showing that such a foliation comes from a smooth effective circle action and then applying Fintushel's classification of 4-manifolds with a circle action [13, 14]. For foliations of dimension 3 Theorem 1.4 reduces to Theorem 1.1, and therefore we are left to study the case when $\dim \mathcal{F} = 2$. Now the codimension of \mathcal{F} is 2, and by [37] the leaf space M^* is a simply connected orbifold, which has no boundary if and only if \mathcal{F} is a regular foliation. According to this criterion we prove the theorem separately in the following two subsections.

4.1. If the foliation is regular, the leaf space M^* is an orbifold without boundary. On the one hand we know that the projection $\pi : M \rightarrow M^*$ induces a surjection

$\pi_1(M) \rightarrow \pi_1(M^*)$, and therefore M^* must be topologically a 2-sphere. On the other hand, by [37, Cor. 5.3] the orbifold fundamental group of M^* is also trivial, and therefore M^* is an orbifold 2-sphere $\mathbb{S}^2(p, q)$ with at most 2 orbifold points of coprime order p, q . In particular, it is possible to write M^* as a union of 2 disks around the orbifold points, and the preimages of these disks decompose M into a union of two disk bundles along possibly exceptional leaves, which proves the result by Theorem 1.1. In this case it is actually possible to know more about which manifolds come up as follows.

Proposition 4.2. *If (M, \mathcal{F}) is a (regular) closed Riemannian foliation of dimension 2 on a compact simply connected 4-manifold M , then the leaves are the fibers of a \mathbb{S}^2 -bundle over \mathbb{S}^2 . In particular, M is diffeomorphic to $\mathbb{S}^2 \times \mathbb{S}^2$ or $\mathbb{C}\mathbb{P}^2 \# -\mathbb{C}\mathbb{P}^2$.*

Proof. Let M^* be the leaf space of (M, \mathcal{F}) . The frame bundle $Fr(M^*)$ is a smooth manifold with an almost free $O(2)$ -action (cf. [1, Thm. 1.23]), and the projection $P : Fr(M^*) \rightarrow M^*$ is a smooth map in the orbifold sense. If $EO(2)$ denotes the universal $O(2)$ -bundle, we construct the *Haefliger classifying space* $B = Fr(M^*) \times_{O(2)} EO(2)$ of M^* . The map $B \rightarrow M^*$ taking $[x, g]$ to $P(x)$ induces an isomorphism in rational cohomology and thus on rational homotopy (cf. [27]), and therefore $\pi_2(B) \otimes \mathbb{Q} = \mathbb{Q}$.

Up to homotopy, there is a fibration (see for example [16])

$$(4.1) \quad L \rightarrow \hat{M} \rightarrow B$$

where \hat{M} is homotopic to M and $L \rightarrow \hat{M}$ is homotopic to the inclusion of a regular leaf. From the long exact sequence in homotopy of the fibration (4.1) it follows that $\pi_1(L) \otimes \mathbb{Q} = \mathbb{Q}$ or 0 . Since L is a compact (orientable) surface, it follows that $L = \mathbb{S}^2$. The only possible exceptional leaf would be $\mathbb{R}\mathbb{P}^2$. In such case the holonomy of \mathcal{F} (cf. Subsection 2.3) would act on $\nu_p(\mathbb{R}\mathbb{P}^2) = \mathbb{R}^2$ without fixed points except the origin, and therefore it would act by an orientation-preserving map. This implies that the normal bundle of $\mathbb{R}\mathbb{P}^2$ is orientable, but this would give a contradiction since M is orientable and $\mathbb{R}\mathbb{P}^2$ is not. In particular there cannot be exceptional leaves, M^* is diffeomorphic to \mathbb{S}^2 , and M is an $L = \mathbb{S}^2$ -bundle over $M^* = \mathbb{S}^2$. Therefore M is foliated diffeomorphic to $(\mathbb{S}^2 \times \mathbb{S}^2, \mathbb{S}^2 \times \{pt\})$, or $\mathbb{C}\mathbb{P}^2 \# -\mathbb{C}\mathbb{P}^2$ foliated by the fibers of the 2-sphere bundle $\mathbb{C}\mathbb{P}^2 \# -\mathbb{C}\mathbb{P}^2 \rightarrow \mathbb{S}^2$. \square

4.2. If the foliation is singular, the leaf space M^* is homeomorphic to a disk D^2 , where the boundary points correspond exactly to the singular leaves of \mathcal{F} . In this case the leaves are orientable surfaces, and they admit a fibration over the singular leaves as described in (2.1). In particular, unless the the regular leaves are tori the only possible singular leaves are points. In such a case, let p be a singular point and let $(\mathbb{S}_p, \mathcal{F}_p)$ be the infinitesimal foliation at p . Since $\mathbb{S}_p = \mathbb{S}^3$ in this case, by [46] the regular leaf \mathcal{L} of \mathcal{F}_p must be diffeomorphic to either \mathbb{S}^2 or T^2 . The fibration (2.1) in this case becomes $\mathcal{L} \rightarrow L \rightarrow p$, where L is a regular leaf of \mathcal{F} , and thus L must also be diffeomorphic to either \mathbb{S}^2 or T^2 .

If the regular leaves of \mathcal{F} are diffeomorphic to \mathbb{S}^2 then the singular leaves must be points, and the singular stratum Σ_0 is the whole boundary of M^* , which is diffeomorphic to \mathbb{S}^1 . In this case, M^* can be written as a union of a ball of a point in the interior, and a tubular neighborhood of the boundary. The preimage of this decomposition under π gives M a splitting structure as (3.1), and the result follows from Theorem 1.1.

The only remaining case to consider is the one where the regular leaf L of \mathcal{F} is diffeomorphic to T^2 . In this case we have the following structure [16]:

- The leaf space M^* is homeomorphic to a disk D^2 with boundary and corners. There are at least 2 corners.
- The leaves corresponding to the interior points of M^* are diffeomorphic to T^2 . The leaves corresponding to points in the boundary of M^* (not corner points) are diffeomorphic to \mathbb{S}^1 . The leaves corresponding to the corners are points.

The preimage of each component of ∂M^* under π is a singular stratum of \mathcal{F} . Fix a regular leaf $L_0 \cong T^2$, and for each edge E_i of ∂M^* fix a leaf $L_i \cong \mathbb{S}^1$ in the preimage of the corresponding edge, and a fibration $\mathbf{p}_i : L_0 \rightarrow L_i$. Let $(m_i, n_i) \in \pi_1(L_0) = \mathbb{Z} \oplus \mathbb{Z}$, $\gcd(m_i, n_i) = 1$, be a primitive generator of the kernel of $\mathbf{p}_{i*} : \pi_1(L_0) \rightarrow \pi_1(L_i)$. Notice that $\{\pm(m_i, n_i)\}$ does not depend on the choice of L_i and \mathbf{p}_i . Call (m_i, n_i) the *weight* of the edge E_i . These correspond precisely to the weights defined in [42]. In particular, the following properties hold:

- Proposition 4.3.** (i) *For every $\gamma : [0, 1] \rightarrow M^*$ such that $\gamma(0) \in E_i$, $\gamma(1) \in E_j$, and $\gamma'(0) \perp E_i$, $\gamma'(1) \perp E_j$, the preimage of $\text{Im}(\gamma)$ in M is a lens space, with fundamental group $\mathbb{Z}/k\mathbb{Z}$, where $k = \begin{vmatrix} m_i & m_j \\ n_i & n_j \end{vmatrix}$. Call any such curve a (E_i, E_j) -curve.*
- (ii) *If ∂M^* consists of at most 4 edges, then M can be divided as a union of two disk bundles of rank greater than 1.*
- (iii) *If ∂M^* consists of more than 4 edges, then there exist edges E_i, E_j , $j \neq i \pm 1$, such that the preimage of every (E_i, E_j) -curve is diffeomorphic to \mathbb{S}^3 .*
- (iv) *If ∂M^* consists of more than 4 edges, then (M, \mathcal{F}) can be written as a foliated connected sum $(\widetilde{M}_1, \mathcal{F}_1) \# (\widetilde{M}_2, \mathcal{F}_2)$, where $\mathcal{F}_1, \mathcal{F}_2$ are codimension 2 singular Riemannian foliations by tori, whose leaf spaces $\widetilde{M}_1^*, \widetilde{M}_2^*$ have boundaries with a number of edges strictly lower than that of ∂M^* .*

This proves Theorem 1.4 for 2-dimensional foliations by induction and use of Theorem 1.1. Hence, the proof of the total Theorem 1.4 will be complete.

Proof. i) Given a (E_i, E_j) -curve γ , let S_{ij} be the preimage of γ under π . It is clear that $\pi^{-1}(\gamma|_{(0,1)})$ is a manifold. Moreover, since $\gamma'(0)$ is perpendicular to E_i , the preimage of $\gamma|_{(0,\epsilon)}$, for $\epsilon > 0$ small enough, is diffeomorphic to a vector bundle over $\pi^{-1}(\gamma(0)) = \mathbb{S}^1$,

and therefore $\pi^{-1}(\gamma|_{[0,\epsilon]})$ is a manifold. By symmetry, $\pi^{-1}(\gamma|_{(1-\epsilon,1]})$ is a manifold, and therefore S_{ij} is a manifold as well.

Without loss of generality, we can suppose that γ passes through $\pi(L_0)$. Consider the open cover $U_0 = \pi^{-1}(\gamma[0,1))$, $U_1 = \pi^{-1}(\gamma(0,1])$ of S_{ij} . U_0 and U_1 retract to $\pi^{-1}(\gamma(0))$ and $\pi^{-1}(\gamma(1))$ respectively, which are both diffeomorphic to \mathbb{S}^1 , while $U_0 \cap U_1$ retracts to L_0 . Moreover, the inclusion maps $\iota_0 : U_0 \cap U_1 \rightarrow U_0$ and $\iota_1 : U_0 \cap U_1 \rightarrow U_1$ are homotopic to the projections $\mathfrak{p}_i : L_0 \rightarrow L_i$ and $\mathfrak{p}_j : L_0 \rightarrow L_j$, respectively. The maps $\mathfrak{p}_{i*}, \mathfrak{p}_{j*} : \mathbb{Z}^2 \rightarrow \mathbb{Z}$ induced between the fundamental groups, are

$$\mathfrak{p}_{i*}(x, y) = n_i x - m_i y, \quad \mathfrak{p}_{j*}(x, y) = n_j x - m_j y.$$

By van Kampen theorem, the fundamental group of S_{ij} has a presentation

$$\langle g_1, g_2 \mid g_1^{n_i} = g_2^{n_j}, g_1^{m_i} = g_2^{m_j} \rangle.$$

Given integers p, q such that $m_i p + n_i q = 1$, we obtain $g_1 = g_1^{m_i p + n_i q} = g_2^{m_j p + n_j q}$ and therefore $\pi_1(S_{ij})$ is generated by g_2 only. Moreover, $1 = g_1^{m_i n_i - n_i m_i} = g_2^{m_i n_j - n_i m_j}$ and thus $\pi_1(S_{ij})$ is cyclic of order $m_i n_j - n_i m_j$, as we wanted to prove.

ii) If ∂M^* has at most 2 edges, M^* can be decomposed as a union of two balls around 2 points in ∂M^* , containing the vertices. If ∂M^* has three vertices, M^* can be decomposed as the union of a disk around one vertex, and the disk around the opposite edge. Finally, if ∂M^* has four vertices, it is diffeomorphic (as an orbifold) to a rectangle, and M^* can be decomposed as a union of the disks around two opposite edges. Therefore, in each case the preimage of the decomposition in M^* divides M as a union of disk bundles, where the preimage of an edge is \mathbb{S}^2 .

iii) This point was proved in [42, Thm. 5.7]. For the sake of completeness, we exhibit the proof here. Let $r \geq 5$ be the number of edges of M^* . Notice that the preimage of a (E_i, E_{i+1}) -curve is diffeomorphic to the unit sphere \mathbb{S}^3 around the (0-dimensional) leaf that is the preimage of the corner between E_i and E_{i+1} . From part i) it follows that

$$(4.2) \quad \begin{vmatrix} m_i & m_{i+1} \\ n_i & n_{i+1} \end{vmatrix} = \pm 1, \quad i = 1, \dots, r,$$

where one should read $r+1 = 1$. We now choose the generators of $\pi_1(L_0) = \mathbb{Z}^2$ so that the edges E_1 has weight $(0, 1)$, and E_2 has weight $(1, 0)$. From (4.2), the weights of E_3 and E_r are, respectively, $(m_3, \pm 1)$, $(\pm 1, n_r)$. Notice that if $m_3 = 0$ (*resp.* $m_3 = \pm 1$), then the preimage of a (E_r, E_3) -curve (*resp.* a (E_1, E_3) -curve) is diffeomorphic to \mathbb{S}^3 . In the same way, if $n_r = 0$, (*resp.* $n_r = \pm 1$) then the preimage of a (E_r, E_3) -curve (*resp.* a (E_r, E_2) -curve) is diffeomorphic to \mathbb{S}^3 . Moreover, for any i if we have $|m_i| = |n_i|$, then $m_i, n_i = \pm 1$ since m_i, n_i are coprime, and thus the preimage of any (E_1, E_i) -curve or (E_2, E_i) -curve is \mathbb{S}^3 .

Therefore, we can now restrict to the case $|m_3| > 1 = |n_3|$, $|n_r| > 1 = |m_r|$, and $|m_i| \neq |n_i|$ for every $i = 3, \dots, r$. Let $3 \leq i < r$ be the first index such that $|m_i| \geq |n_i| + 1$, $|n_{i+1}| \geq |m_{i+1}| + 1$. The existence of such an i is assured by (4.2). Then

$$|m_i n_{i+1}| \geq (|n_i| + 1)(|m_{i+1}| + 1) = |m_{i+1} n_i| + |m_{i+1}| + |n_i| + 1$$

and therefore

$$1 = |m_i n_{i+1} - m_{i+1} n_i| \geq |m_i n_{i+1}| - |m_{i+1} n_i| \geq |m_{i+1}| + |n_i| + 1.$$

In particular $m_{i+1} = n_i = 0$, which implies $|n_{i+1}| = |m_i| = 1$, and applying the result in i) again, we obtain that the preimages of both (E_1, E_i) -curves and (E_2, E_{i+1}) -curves are diffeomorphic to \mathbb{S}^3 .

iv) By iii), the preimage $S \cong \mathbb{S}^3$ of some (E_i, E_j) -curve decomposes M as a union of two connected 4-manifolds M_1, M_2 along S . In the following we will focus on M_1 , but everything will hold for M_2 as well. Now $(M_1, \mathcal{F}|_{M_1})$ is a foliated 4-manifold with boundary S , and the restriction of \mathcal{F} to S gives a codimension 1 B-foliation $(S, \mathcal{F}|_S)$. By [16] it is foliated diffeomorphic to $(\mathbb{S}^3, \mathcal{F}')$, where \mathcal{F}' is given by the orbit decomposition of the (unique up to conjugation) isometric T^2 -action on the round \mathbb{S}^3 . This action can be extended to an isometric action on the 4-disk \mathbb{D}^4 , and this action induces a singular Riemannian foliation $\mathcal{F}_{\mathbb{D}^4}$. The manifold

$$\widetilde{M}_1 := M_1 \cup_{S \cong \mathbb{S}^3} \mathbb{D}^4$$

is canonically foliated by the singular foliation \mathcal{F}_1 that restricts to $\mathcal{F}|_{M_1}$ on M_1 and $\mathcal{F}_{\mathbb{D}^4}$ on \mathbb{D}^4 . In the next section we will prove the following, rather technical, lemma:

Lemma 4.4 (Gluing Lemma). *There is a metric \tilde{g} on \widetilde{M}_1 that is bundle-like for \mathcal{F}_1 , i.e., such that $(\widetilde{M}_1, \tilde{g}, \mathcal{F}_1)$ is a singular Riemannian foliation.*

This proves, in particular, that (M, \mathcal{F}) is foliated diffeomorphic to a foliated connected sum $(\widetilde{M}_1, \mathcal{F}_1) \# (\widetilde{M}_2, \mathcal{F}_2)$ of two codimension 2 singular Riemannian foliations by tori. Moreover, it is easy to see that the leaf spaces $\widetilde{M}_i / \mathcal{F}_i$ are obtained by gluing a corner $\mathbb{D}^4 / \mathcal{F}_{\mathbb{D}^4} \simeq \{(x, y) \in \mathbb{D}^2 \mid x \geq 0, y \geq 0\}$ to $M_i / \mathcal{F}|_{M_i}$ along the common boundary $\mathbb{S}^3 / \mathcal{F}' \simeq S / \mathcal{F}|_S$, and in particular have strictly less boundary edges than M / \mathcal{F} by iii).

The proof of the proposition is now complete. \square

5. THE GLUING LEMMA

The goal of this section is to prove Lemma 4.4 above. Since $(S, \mathcal{F}|_S)$ is foliated diffeomorphic to $(\mathbb{S}^3, \mathcal{F}')$ we will from now on identify these two spaces, and suppose that the (foliated) boundary of M_1 is $(\mathbb{S}^3, g_0, \mathcal{F}')$ with some metric g_0 .

In what follows, we write \mathbb{S}^3 as a union of disk bundles

$$(5.1) \quad \mathbb{S}^3 = \mathbb{D}^2 \times \mathbb{S}^1 \bigcup_{\phi_0} \mathbb{D}^2 \times \mathbb{S}^1,$$

where $\phi_0 \in \text{Diff}(T^2 = \partial(\mathbb{D}^2 \times \mathbb{S}^1))$ is the map interchanging the two factors, such that the leaves of \mathcal{F}' are the concentric tori with respect to the canonical metric in $\mathbb{D}^2 \times \mathbb{S}^1$.

First of all, we observe the following.

Lemma 5.1. *There is a neighborhood U of \mathbb{S}^3 in M_1 that is foliated diffeomorphic to $(\mathbb{S}^3 \times [-\epsilon, 0], \mathcal{F}' \times \{pt\})$. Moreover, the metric g on M_1 induces a metric on $\mathbb{S}^3 \times [-\epsilon, 0]$ of the form $g_t + dt^2$, where $\mathfrak{F}_- = \{g_t\}_{t \in [-\epsilon, 0]}$ is a family of bundle-like metrics on $(\mathbb{S}^3, \mathcal{F}')$.*

Proof. Consider the unique unit length normal vector field X on $\mathbb{S}^3 \subseteq M_1$, pointing in the *outward* direction of M_1 . Since X is uniquely defined, its restriction to every leaf of \mathcal{F}' is basic with respect to the projection $M_1 \rightarrow M_1/\mathcal{F}|_{M_1}$. Since \mathbb{S}^3 is compact, there is a neighborhood U of \mathbb{S}^3 in M_1 such that the map $(p, t) \mapsto \exp_p tX(p)$ defines a diffeomorphism $\exp^\perp : \mathbb{S}^3 \times [-\epsilon, 0] \rightarrow U$. On the other hand, by the equifocality of singular Riemannian foliations [6, Thm. 1.5] \exp^\perp is also a foliated map because X is basic. Moreover, X is perpendicular to the family of submanifolds $S_t = \exp^\perp(\mathbb{S}^3 \times \{t\})$, all diffeomorphic to \mathbb{S}^3 via \exp^\perp , and therefore the tangent space of U splits orthogonally as $TS_t \oplus \langle X \rangle$. If dt denotes the (exact) 1-form dual to X , then the metric g splits as $g|_{TS_t} + dt^2$. By setting $g_t = (\exp^\perp)^*(g|_{TS_t})$ we obtain the result. \square

In the same way, there is a neighborhood of $(\mathbb{S}^3 = \partial\mathbb{D}^4, g_1, \mathcal{F}')$ in $(\mathbb{D}^4, \mathcal{F}_{\mathbb{D}^4})$ that is foliated diffeomorphic to $\mathbb{S}^3 \times [1, 1 + \epsilon]$, where the metric has the form $g_t + dt^2$ for some family of bundle-like metrics $\mathfrak{F}_+ = \{g_t\}_{t \in [1, 1 + \epsilon]}$ on \mathbb{S}^3 .

Suppose now that the metrics g_0 and g_1 on \mathbb{S}^3 can be connected by a family $\mathfrak{F} = \{g_t\}_{t \in [0, 1]}$ of smoothly varying bundle-like metrics for \mathcal{F}' , i.e., such that $(\mathbb{S}^3, g_t, \mathcal{F}')$ is a singular Riemannian foliation for all $t \in [0, 1]$. In this case one can extend \mathfrak{F} to a smooth family $\{g_t\}_{t \in [-\epsilon, 1 + \epsilon]}$ using \mathfrak{F}_\pm . In particular $(\mathbb{S}^3 \times [0, 1], g_t + dt^2, \mathcal{F}' \times \{pt\})$ is a singular Riemannian foliation that can be glued smoothly with $(M_1, \mathcal{F}|_{M_1})$ and $(\mathbb{D}^4, \mathcal{F}_{\mathbb{D}^4})$, giving a singular Riemannian foliation on

$$M_1 \bigcup (\mathbb{S}^3 \times [0, 1]) \bigcup \mathbb{D}^4 \cong \widetilde{M}_1.$$

We are thus left to produce a smooth family \mathfrak{F} of bundle-like metrics connecting g_0 to g_1 . Notice that it is enough to produce a piecewise smooth family. The construction will proceed along the following steps:

- (i) Find an orientation-preserving foliated diffeomorphism $\overline{F} \in \text{Diff}(\mathbb{S}^3, \mathcal{F}')$ such that $\overline{F}^* g_0$ and g_1 have the same horizontal spaces near the singular leaves.
- (ii) Prove that \overline{F} is foliated isotopic to the identity. If \overline{F}_t is such an isotopy, then $\overline{F}_t^* g_0$ is a one parameter family \mathfrak{F}_1 between g_0 and $\overline{F}^* g_0$.
- (iii) Find a one parameter family \mathfrak{F}_2 between $\overline{F}^* g_0$ to a metric g' that has the same horizontal spaces as g_1 .
- (iv) Find a one parameter family \mathfrak{F}_3 between g' and g_1 .

The one parameter family we need is obtained by composing $\mathfrak{F}_1 * \mathfrak{F}_2 * \mathfrak{F}_3$. We prove each step in a separate lemma.

Lemma 5.2 (Step i). *There exist orientation-preserving foliated diffeomorphisms $F, G \in \text{Diff}(\mathbb{D}^2 \times \mathbb{S}^1)$ with $F|_{\mathbb{S}^1} = id = G|_{\mathbb{S}^1}$, which glue to a diffeomorphism $\overline{F} := (F, G) \in \text{Diff}(\mathbb{S}^3, \mathcal{F}')$ such that \overline{F}^*g_0 and g_1 have the same horizontal spaces around the singular leaves.*

Proof. Let $r : \mathbb{D}^2 \times \mathbb{S}^1 \rightarrow [0, 1]$ denote the radial function on $\mathbb{D}^2 \times \mathbb{S}^1$. Consider the homothetic transformation with respect to g_1 (cf. [40]):

$$h_\lambda^1 : \mathbb{D}^2 \times \mathbb{S}^1 \rightarrow \mathbb{D}^2 \times \mathbb{S}^1, \quad \lambda \in (0, 1],$$

defined in such a way that if $q = \exp_p(x)$ for some $p \in \mathbb{S}^1$ and some g_1 -horizontal vector x , then $h_\lambda^1(q) = \exp_p(\lambda x)$. In the same way, define the homothetic transformation h_λ^0 with respect to g_0 . Notice that for each $\lambda \in (0, 1]$, $f_\lambda = (h_\lambda^1)^{-1} \circ h_\lambda^0$ is a foliated diffeomorphism of $\mathbb{D}^2 \times \mathbb{S}^1$ that restricts to the identity on \mathbb{S}^1 , $f_1 = id$, and f_λ converges smoothly to $\exp_{g_1}^\perp \circ (\exp_{g_0}^\perp)^{-1}$ as $\lambda \rightarrow 0$, where $\exp_{g_1}^\perp, \exp_{g_0}^\perp$ denote the normal exponential maps of $\{0\} \times \mathbb{S}^1 \subseteq \mathbb{D}^2 \times \mathbb{S}^1$, and the normal bundles of the singular leaf L^+ are identified via $\nu L^+ \simeq T\mathbb{S}^3/T\mathbb{S}^1$. In particular, if we define $f_0 = \exp_{g_1}^\perp \circ (\exp_{g_0}^\perp)^{-1}$, then $f_0|_{\mathbb{S}^1} = id$ and f_0 takes g_0 -horizontal geodesics to g_1 -horizontal geodesics (and thus $f_0^*g_0$ and g_1 have the same horizontal spaces). Let

$$\begin{aligned} f : \mathbb{D}^2 \times \mathbb{S}^1 \times [0, 1] &\longrightarrow \mathbb{D}^2 \times \mathbb{S}^1 \\ (p, t) &\longmapsto f_t(p) \end{aligned}$$

Now consider a function $\varphi : [0, 1] \rightarrow [0, 1]$ such that $\varphi(t) = 0$ in $[0, \epsilon)$, and $\varphi(t) = 1$ in $(1 - \epsilon, 1]$, and define an embedding

$$\begin{aligned} \iota : \mathbb{D}^2 \times \mathbb{S}^1 &\longrightarrow \mathbb{D}^2 \times \mathbb{S}^1 \times [0, 1] \\ p &\longmapsto (p, \varphi(r(p))) \end{aligned}$$

The composition $F := f \circ \iota$ gives an orientation-preserving foliated diffeomorphism of $\mathbb{D}^2 \times \mathbb{S}^1$ that coincides with f_0 in a neighborhood of \mathbb{S}^1 , and is the identity next to the boundary. If we denote by G the same map on the other copy of $\mathbb{D}^2 \times \mathbb{S}^1$, we can glue the diffeomorphisms by ϕ_0 in (5.1) and obtain the desired $\overline{F} = (F, G) \in \text{Diff}(\mathbb{S}^3, \mathcal{F}')$. \square

Lemma 5.3 (Step ii). *The map \overline{F} in Lemma 5.2 is foliated isotopic to the identity.*

Proof. Consider tubular neighborhoods U^\pm of small radius ϵ around the singular leaves L^\pm . First, noticing that $\overline{F} = id$ at the singular leaves, we can regard $\overline{F}|_{U^\pm}$ alternatively as a new tubular neighborhood map $\overline{F}|_{U^\pm} : \nu L^\pm \rightarrow \mathbb{S}^3$, where we have identified U^\pm and the normal bundles νL^\pm . Since tubular neighborhood maps are foliated isotopic to each other (cf. Theorems 5.3 and 6.5 in Chapter 4 in [31]), $\overline{F}|_{U^\pm}$ is foliated isotopic to the identity inclusion $id : \nu L^\pm \rightarrow \mathbb{S}^3$. Then by the isotopy extension theorem (Theorem

1.3 in Chapter 8 in [31]), \overline{F} is foliated isotopic to a diffeomorphism that restricts to the identity on U^\pm .

Suppose now that \overline{F} fixes $U^+ \cup U^-$. Let $r = \text{dist}_{L^+}$, where the distance is taken with respect to either metric, and let $L_t = r^{-1}(t)$. We can assume, up to rescaling the metric, that the two singular leaves correspond to $L^+ = r^{-1}(0)$ and $L^- = r^{-1}(1)$. Set $T = L_\epsilon$. Using for example the normal holonomy with respect to g_0 , we can identify any L_r , $r \in [\epsilon, 1 - \epsilon]$ with T . Since \overline{F} sends any leaf L_r to itself, and it fixes every leaf L_t with $t \in [0, \epsilon] \cup [1 - \epsilon, 1]$, we can think of \overline{F} as a closed loop $\gamma : t \in [\epsilon, 1 - \epsilon] \mapsto \overline{F}|_{L_t} \in \text{Diff}(L_t) = \text{Diff}(T)$ based at the identity. In particular, the path lies in the identity component $\text{Diff}_0(T)$. Let $(\mathbb{D}^2 \times \mathbb{S}^1, T)$ denote the homogeneous foliation where $T = T^2$ (seen as a Lie group) acts linearly on $\mathbb{D}^2 \times \mathbb{S}^1$. By fixing a foliated diffeomorphism $U^+ \simeq \mathbb{D}^2 \times \mathbb{S}^1$, we obtain an action of T on each L_t , $t \leq \epsilon$, and in particular on T itself. We thus obtain a map $\iota : T \rightarrow \text{Diff}_0(T)$ that is known to be a deformation retract of $\text{Diff}_0(T)$. In particular, we can homotope the loop $\gamma(t)$ to a loop in $\iota(T)$, and this gives an isotopy between \overline{F} and a diffeomorphism \hat{F} such that $\hat{F}|_{L_t} \in \iota(T)$ for all t .

The map $\hat{\gamma} : t \in [\epsilon, 1 - \epsilon] \mapsto \hat{F}|_{L_t} \in \iota(T)$ is a loop based at the identity. However, since every diffeomorphism in $\iota(T)$ can be completed to a diffeomorphism of U^+ - and this completion can be made canonical using the foliated diffeomorphism $U^+ \simeq \mathbb{D}^2 \times \mathbb{S}^1$ - we can homotope the curve $\hat{\gamma}$ by only fixing the end point, and letting the initial point free to move within $\iota(T)$. In this way we obtain an isotopy through foliated diffeomorphisms that still fix U^- but may not in general fix U^+ . It is clear then, that by allowing such freedom in $\hat{\gamma}$ we can homotope it to the constant map $t \mapsto \text{id} \in \text{Diff}_0(T)$, which corresponds to isotoping \hat{F} (and hence \overline{F}) to the identity. \square

Lemma 5.4 (Step iii). *If g, \tilde{g} are two bundle-like metrics of $(\mathbb{S}^3, \mathcal{F}')$ with the same horizontal spaces near the singular leaves, there is a one-parameter family of bundle-like metrics g_t from \tilde{g} to a metric g' with the same horizontal spaces as g .*

Proof. Denote by \mathbb{S}_{reg}^3 the complement in \mathbb{S}^3 of the singular leaves. There are two horizontal distributions $\Delta, \tilde{\Delta}$ given by the g - and \tilde{g} -orthogonal spaces to \mathcal{F}' , which are both of dimension one in \mathbb{S}_{reg}^3 . Notice moreover that, by assumption, $\Delta = \tilde{\Delta}$ in a neighbourhood of the singular leaves. Choose a variation of distributions Δ_t from $\tilde{\Delta}$ to Δ , in such a way that Δ_t is always transverse to \mathcal{F} , and $\Delta_t(p)$ is constant wherever $\Delta(p) = \tilde{\Delta}(p)$. For each point $p \in \mathbb{S}^3$, define Δ_t^\perp the \tilde{g} -orthogonal distribution to Δ_t , and the \tilde{g} -orthogonal projections $\pi_t : T_p \mathbb{S}^3 \rightarrow \Delta_t^\perp$, $\pi_h : T_p \mathbb{S}^3 \rightarrow \tilde{\Delta}$. Finally, define

$$h_t(x, y) = \tilde{g}(\pi_t x, \pi_t y) + \tilde{g}(\pi_h x, \pi_h y)$$

Notice that h_t is a metric, and it satisfies the following properties:

- If Δ_t varies smoothly, h_t varies smoothly since it is defined in terms of functions that depend smoothly on Δ_t .

- The h_t -orthogonal space to \mathcal{F}' is Δ_t . In fact, if $v \in T_p L_p$ and $x \in \Delta_t$, then $\pi_t(x) = 0$, $\pi_h(v) = 0$ and thus $h_t(v, x) = 0$.
- Wherever $\Delta_t(p) = \tilde{\Delta}(p)$, we also have $h_t(p) = \tilde{g}(p)$.
- The transverse metric h_t^T equals the transverse metric \tilde{g}^T .

In particular, h_t is a bundle-like metric (cf. [3]), and thus defining $g' = h_1$ completes the proof of the lemma. \square

Lemma 5.5 (Step iv). *Let (M, \mathcal{F}) be any singular Riemannian foliation, and let g, g' be two bundle-like metrics with the same horizontal distribution. Then for any t , the metric $g_t = tg + (1 - t)g'$ is a bundle-like metric.*

Proof. Of course g_t is a metric, and the g_t -horizontal distribution is the same as the g - and g' -horizontal distributions. In particular, $g_t^T = t \cdot g^T + (1 - t) \cdot g'^T$. On each stratum Σ , we can take a vertical vector field $X \in \mathfrak{X}(\mathcal{F})$, and since g, g' are bundle-like metrics of $(\Sigma, \mathcal{F}|_\Sigma)$, we have $L_X g^T = L_X g'^T = 0$. Therefore

$$L_X g_t^T = t \cdot L_X g^T + (1 - t) \cdot L_X g'^T = 0$$

and therefore g_t is a bundle-like metric on each stratum. By [3], this is enough to ensure that g_t is a bundle-like metric for \mathcal{F} . \square

Acknowledgements . *The authors are very grateful to Alexander Lytchak for kindly introducing each other, very nice suggestions and useful comments. The authors also thank Zizhou Tang, Gudlaugur Thorbergsson, Burkhard Wilking and Wolfgang Ziller for their support and valuable discussion. Many thanks also to Marcos M. Alexandrino and Fernando Galaz-Garcia for their interest and helpful conversation. The first author would like to thank the Alexander von Humboldt Foundation and the University of Cologne for their support and hospitality during his Humboldt postdoctoral position in 2012–2014, under the supervision of professor Gudlaugur Thorbergsson.*

REFERENCES

- [1] A. Adem, J. Leida and Y. Ruan, *Orbifolds and stringy topology*. Cambridge Tracts in Mathematics, **171**. Cambridge University Press, Cambridge, (2007).
- [2] A. Akhmedov and B. Doug Park, *Exotic smooth structures on small 4-manifolds with odd signatures*, Invent. Math. **181** (2010), no. 3, 483–492.
- [3] M.M. Alexandrino, *Desingularization of singular Riemannian foliation*, Geom. Dedicata **149** (2010), 397–416.
- [4] M.M. Alexandrino, R. Briquet, and D. Töben, *Progress in the theory of singular Riemannian foliations* Differential Geom. Appl. **31** (2013), no. 2, 248–267.
- [5] M.M. Alexandrino and A. Lytchak, *On smoothness of isometries between orbit spaces*, Riemannian geometry and applications—Proceedings RIGA 2011, 17–28, Ed. Univ. Bucureşti, Bucharest, 2011.
- [6] M.M. Alexandrino and D. Töben, *Equifocality of a singular riemannian foliation*, Proc. Amer. Math. Soc. **136** (2008), 3271–3280.
- [7] J. Cerf, *Sur les difféomorphismes de la sphère de dimension trois ($\Gamma_4 = 0$)*, Lecture Notes in Mathematics, 53, Springer-Verlag, Berlin-New York, (1968).

- [8] S.K. Donaldson, *An application of gauge theory to four-dimensional topology*, J. Diff. Geom. **18** (1983), 279–315.
- [9] S.K. Donaldson, *Irrationality and the h-cobordism conjecture*, J. Diff. Geom. **26** (1987), 141–168.
- [10] H.B. Duan, *A generalization of the Wang sequence*, Homol. Homotopy Appl. **13** (2011), 37–42.
- [11] R. Fintushel, *Circle actions on simply connected 4-manifolds*, Trans. Amer. Math. Soc. **230** (1977), 147–171.
- [12] R. Fintushel, *Classification of circle actions on 4-manifolds*, Trans. Amer. Math. Soc., **242** (1978), 377–390.
- [13] R. Fintushel and R.J. Stern, *Double node neighborhoods and families of simply connected 4-manifolds with $b^+ = 1$* , J. Amer. Math. Soc. **19** (2005), 171–180.
- [14] M.H. Freedman, *The topology of four dimensional manifolds*, J. Diff. Geom. **17** (1982), no. 3, 357–453.
- [15] M. Freedman and F. Quinn, *Topology of 4-manifolds*, Princeton Mathematical Series 39, Princeton University Press, Princeton, N.J., (1990).
- [16] F. Galaz-Garcia and M. Radeschi, *Singular Riemannian foliations and applications to positive and nonnegative curvature*, preprint, 2013, arXiv:1302.4593.
- [17] J.Q. Ge, *DDVV-type inequality for skew-symmetric matrices and Simons-type inequality for Riemannian submersions*, Advances in Math. **251** (2014), 62–86.
- [18] J.Q. Ge and Z.Z. Tang, *Isoparametric functions and exotic spheres*, J. reine angew. Math. **683** (2013), 161–180.
- [19] J.Q. Ge and Z.Z. Tang, *Geometry of isoparametric hypersurfaces in Riemannian manifolds*, Asian J. Math. **18** (2014), 117–126.
- [20] J.Q. Ge, Z.Z. Tang and W.J. Yan, *A Filtration for Isoparametric Hypersurfaces in Riemannian Manifolds*, to appear in J. Math. Soc. Jpn., arXiv:1102.1126.
- [21] E. Ghys, *Feuilletages riemanniens sur les variétés simplement connexes*, Ann. Inst. Fourier (Grenoble) **34** (1984), 203–223. Theorem A
- [22] H. Gluck, *The embedding of two-spheres in the four-sphere*, Trans. Amer. Math. Soc. **104** (1962), 308–333.
- [23] R. Gompf, *Three exotic R^4 's and other anomalies*, J. Diff. Geom. **18** (1983), 317–328.
- [24] D. Gromoll and G. Walschap, *Metric foliations and curvature*, Progress in Mathematics, **268**, Birkhäuser Verlag, Basel, 2009.
- [25] K. Grove and S. Halperin, *Dupin hypersurfaces, group actions and the double mapping cylinder*, J. Diff. Geom. **26** (1987), 429–459.
- [26] K. Grove and W. Ziller, *Lifting group actions and nonnegative curvature*, Trans. Amer. Math. Soc. **363** (2011), 2865–2890.
- [27] A. Haefliger, *Holonomy Groupoïdes d'holonomie et classifiants*, Transversal structure of foliations (Toulouse, 1982). Astérisque **116** (1984), 70–97.
- [28] A.E. Hatcher, *On the diffeomorphism group of $\mathbb{S}^1 \times \mathbb{S}^2$* , Proc. Amer. Math. Soc. **83** (1981), 427–430.
- [29] A.E. Hatcher, *A proof of the Smale conjecture, $\text{Diff}(\mathbb{S}^3) \simeq O(4)$* , Ann. of Math. **117** (1983), 553–607.
- [30] J.J. Hebda, *The regular focal locus*, J. Diff. Geom. **16** (1981), 421–429.
- [31] M.W. Hirsch, *Differential Topology* (GTM. **33**), Springer-Verlag, New York etc., 1976.
- [32] C.A. Hoelscher, *Classification of cohomogeneity one manifolds in low dimensions*, Pac. J. Math. **246** (2010), 129–185.
- [33] S. Hong, J. Kalliongis, D. McCullough and J. H. Rubinstein, *Diffeomorphisms of elliptic 3-manifolds*, Lecture Notes in Mathematics **2055**, Springer, Heidelberg, 2012.

- [34] T. Lawson, *Normal bundles for an embedded $\mathbb{R}P^2$ in a positive definite 4-manifold*, J. Diff. Geom. **22** (1985), 215–231.
- [35] T. Lawson, *The minimal genus problem*, Exposition. Math. **15** (1997), 385–431.
- [36] B.H. Li and T.J. Li, *Minimal genus smooth embeddings in $S^2 \times S^2$ and $\mathbb{C}P^2 \# n\overline{\mathbb{C}P^2}$ with $n \leq 8$* , Topology **37** (1998), 575–594.
- [37] A. Lytchak, *Geometric resolution of singular Riemannian foliations*, Geometriae Dedicata **149** (2010), 379–395.
- [38] D. McCullough, *Isometries of elliptic 3-manifolds*, J. London Math. Soc. **65** (2002), 167–182.
- [39] J.W. Milnor and J.D. Stasheff, *Characteristic classes*, Ann. Math. Studies No.76, Princeton University, (1974).
- [40] P. Molino, *Riemannian foliations*, Translated from the French by Grant Cairns. With appendices by G. Cairns, Y. Carrière, E. Ghys, E. Salem, V. Sergiescu. Progress in Math. **73**. Birkhäuser Boston, Inc., Boston, MA, 1988.
- [41] J.M. Montesinos, *Classical tessellations and three manifolds*, Universitext. Springer-Verlag, Berlin, 1987.
- [42] P. Orlik and F. Raymond, *Actions of the torus on 4-manifolds I*, Trans. Amer. Math. Soc., **152** (1970), 531–559.
- [43] P. Orlik and F. Raymond, *Actions of the torus on 4-manifolds. II*, Topology **13** (1974), 89–112.
- [44] J. Parker, *4-dimensional G-manifolds with 3-dimensional orbits*, Pac. J. Math. **125** (1986), 187–204.
- [45] C. Qian and Z.Z. Tang, *Isoparametric functions on exotic spheres*, Advances in Math. **272** (2015), 611–629.
- [46] M. Radeschi, *Low dimensional singular Riemannian foliations on spheres*, Ph.D. thesis, University of Pennsylvania (2012).
- [47] N. Steenrod, *The Topology of Fibre Bundles*, PMS-14, Princeton University Press, Princeton, N.J., 1951.
- [48] R. Takagi, *On homogeneous real hypersurfaces in a complex projective space*, Osaka J. Math. **10** (1973), 495–506.
- [49] C.H. Taubes, *Gauge theory on asymptotically periodic 4-manifolds*, J. Diff. Geom. **25** (1987), 363–430.
- [50] G. Thorbergsson, *Singular Riemannian foliations and isoparametric submanifolds*, Milan J. Math. **78** (2010), 355–370.
- [51] Q.M. Wang, *Isoparametric Functions on Riemannian Manifolds. I*, Math. Ann. **277** (1987), 639–646.

SCHOOL OF MATHEMATICAL SCIENCES, LABORATORY OF MATHEMATICS AND COMPLEX SYSTEMS,
BEIJING NORMAL UNIVERSITY, BEIJING 100875, P.R. CHINA.

E-mail address: jqge@bnu.edu.cn

MATHEMATISCHES INSTITUT, WWU MÜNSTER, EINSTEINSTR. 62, MÜNSTER, GERMANY.

E-mail address: mrade_02@uni-muenster.de