

YANG-MILLS THEORY AND JUMPING CURVES

YASHA SAVELYEV

To Anna

ABSTRACT. We study a smooth analogue of jumping curves of a holomorphic vector bundle, and use Yang-Mills theory over S^2 to show that any non-trivial, smooth Hermitian vector bundle E over a smooth simply connected manifold, must have such curves. This is used to give new examples complex manifolds for which a non-trivial holomorphic vector bundle must have jumping curves in the classical sense, (when $c_1(E)$ is zero). We also use this to give a new proof a theorem of Gromov on the norm of curvature of unitary connections, and make the theorem slightly sharper. Lastly we define a sequence of new non-trivial integer invariants of smooth manifolds, connected to this theory of smooth jumping curves, and make some computations of these invariants. Our methods include an application of the recently developed Morse-Bott chain complex for the Yang-Mills functional over S^2 .

1. INTRODUCTION

To study a holomorphic vector bundle E over a complex manifold X , one strategy is to extract information about E from information obtained by restrictions of E to (families) of complex genus 0 curves in X . By Birkhoff-Grothendieck theorem, a complex rank r holomorphic vector bundles over $\mathbb{C}\mathbb{P}^1$ is holomorphically isomorphic to

$$\sum_{1 \leq i \leq r} \mathcal{O}(d_i),$$

with weights d_i uniquely determined, (up to order) which makes their theory particularly transparent. We shall say that a holomorphic curve $u : \mathbb{C}\mathbb{P}^1 \rightarrow X$ is a *jumping curve* if the pullback of E to $\mathbb{C}\mathbb{P}^1$ by u is holomorphically non-trivial. When $c_1(E) \neq 0$ the classical definition is different but for our purpose we stick to the above for convenience. It is not actually very hard to extend our arguments to the case of classical jumping curves (when $c_1(E) \neq 0$), but at the cost of much added complexity. The strategy then consists of studying splitting types of jumping curves. For $X = \mathbb{C}\mathbb{P}^n$, one may take this approach very far, see for example [14], in particular the authors prove that a holomorphic vector bundle over $\mathbb{C}\mathbb{P}^n$ is non trivial if and only if it has jumping lines.

Jumping curves can be defined for a smooth Hermitian vector bundle with a connection, as a direct generalization of the holomorphic case. Let X be a smooth manifold, E a Hermitian vector bundle over X , and A a Hermitian, i.e. unitary connection on E . Given a smooth map $u : S^2 \rightarrow X$, the pull-back bundle u^*E , with its pull back connection u^*A , has a natural induced holomorphic structure. It is

1991 *Mathematics Subject Classification.* 53C07, 37D15, 14D20.

Key words and phrases. YangMills flow; jumping curves; smooth invariants; cascade complex.

the holomorphic structure whose Dolbeault operator is induced by the anti-complex linear part of the covariant derivative operator corresponding to A , this operator will be denoted by $\bar{\partial}_A$. This Dolbeault operator is necessarily integrable, see for instance [1]. We shall say that u is a *smooth jumping curve* if the $\bar{\partial}_{u^*A}$ holomorphic structure on u^*E is non-trivial. We will also just say jumping curves where there is no possibility of confusion with holomorphic case.

Theorem 1.1. *Let X be a simply connected smooth manifold, E a non-trivial, smooth Hermitian vector bundle over X , then for any Hermitian connection A on E , there are smooth jumping curves in any class $[u] \in \pi_2(X)$.*

This is proved by studying the map $\Omega^2 X \rightarrow \Omega^2 BU(r)$ induced by the classifying map for E , via Yang-Mills flow, where $\Omega^2 X$ denotes the spherical mapping space. More precisely we use the energy flow for the Yang-Mills functional, and at some points we use its Morse-Bott complex, which for the base S^2 has recently been nicely developed, [17].

Let us explain the relationship of the above to the holomorphic setting. Suppose that $E \rightarrow X$ is a holomorphic vector bundle. Pick a Hermitian structure h on E . Then there is a unique compatible Hermitian connection A_h on E , inducing the given holomorphic structure, [7]. From this it readily follows that a holomorphic curve $u : \mathbb{C}\mathbb{P}^1 \rightarrow X$, is a jumping curve if and only if it is a smooth jumping curve with respect to A_h . Let $Hol^{[u]}(\mathbb{C}\mathbb{P}^1, X)$ denote the space of based holomorphic maps in class $[u]$. Then an argument for theorem above, together with the discussion above give the following.

Theorem 1.2. *Suppose that E is a non-trivial (topologically as a complex vector bundle) holomorphic vector bundle over X , with $\pi_1(X) = 0$ and suppose that for every $k > 0$ there is a class $[u_k]$ such that the inclusion $Hol^{[u_k]}(\mathbb{C}\mathbb{P}^1, X) \rightarrow \Omega^{2, [u_k]} X$ induces an isomorphism of homotopy groups, in range $[0, k]$. Then E has (holomorphic) jumping curves. In particular this holds for $X = \mathbb{C}\mathbb{P}^n$, a generalized flag manifold, a toric manifold, and $X = \Omega G$, with G compact Lie group.*

Note that we can no longer control the class of jumping curves without stronger assumptions. The hypothesis are known to be satisfied for $X = \mathbb{C}\mathbb{P}^n$ [16], flag manifolds for example [9], [12] toric manifolds [10], loop groups ΩG , with G compact Lie group, for example [5]. Thus 1.2 also reproduces a slightly weakened version of the result in [14] mentioned above, weaker because in principle we obtain only jumping curves rather than lines, and we must ask for topological non-triviality rather than holomorphic. For the other cases the above existence result on jumping curves seems to be new.

1.1. Some new integer invariants of smooth vector bundles and smooth manifolds. For a smooth complex rank r vector bundle E over X , and a unitary connection A , if $u : S^2 \rightarrow X$ is a smooth curve define

$$(1.1) \quad |u|_A = \sum_i |d_i|^2,$$

$$(1.2) \quad |u|_{A, \infty} = \max_i |d_i|,$$

with d_i the weights in the Birkhoff-Grothendieck splitting of $(u^*E, \bar{\partial}_{u^*A})$. Note that $|u|_A$ is the energy of the homomorphism $S^1 \rightarrow U(n)$, with weights d_i , after

appropriate choice of normalization of the bi-invariant metric on $U(n)$. This will come into the proof of 1.5.

Define:

$$\zeta_{YM}(E, n) = \inf_{A, [f], f' \in [f]} \sup_{s \in S^n} |f'(s)|_A \in \mathbb{N},$$

where the infimum is over all $[f] \in \pi_n(\Omega^2 X)$, s.t. the map $\Omega^2 X \rightarrow \Omega^2 BU(r)$ induced by the classifying map of E is non-vanishing on $[f]$, if there is no such $[f]$ set $\zeta_{YM}(E, n) = 0$. Here $\Omega^2 X$ denotes the component of the spherical mapping space in the constant class, (which is why the invariant is \mathbb{N} valued). Also define:

$$\zeta_{YM}(E) = \inf_n \zeta_{YM}(E, n) \in \mathbb{N},$$

$$K_{YM}(X, r) = \inf_E \zeta_{YM}(E) \in \mathbb{N},$$

$$K_{YM}(X) = \inf_{r > 0} K_{YM}(X, r) \in \mathbb{N}.$$

where the second infimum is over all rank $r > 0$ complex vector bundles E with some non-vanishing Chern number.

Theorem 1.3. *If E is a non-trivial complex vector bundle over a simply connected X , then the sequence of positive integers $\{\zeta_{YM}(E, n)\}$ cannot be identically 0 and hence $\zeta_{YM}(E) \neq 0$, $K_{YM}(X, r) \neq 0$, and $K_{YM}(X) \neq 0$.*

Conjecture 1.4. *For $f : X \rightarrow Y$ a smooth homotopy equivalence, and E a complex vector bundle over Y :*

$$\zeta_{YM}(E) = \zeta_{YM}(f^* E),$$

in particular the invariants $K_{YM}(X, r)$, $K_{YM}(X)$ of a smooth manifold X are homotopy invariants.

This conjecture is extremely likely. It would follow immediately if the perturbation property needed in the proof of Theorem 1.5, held for every X as opposed to just $X = S^m$. With the perturbation property, it also becomes much easier to compute (at least in principle) the above invariants.

We note that unlike Pontryagin and Chern numbers, the integers $\zeta_{YM}(E, n)$ are not stable. That is, for a direct sum $E' = E \oplus \epsilon^k$ of vector bundles with ϵ^k the trivial \mathbb{C}^k vector bundle, we do not in general have $\zeta_{YM}(E, n) = \zeta_{YM}(E', n)$. In fact we have:

Theorem 1.5. *For any rank r complex vector bundle $E \rightarrow S^m$, and any $k > 0$, so that $n \leq 2(r + k) - 2$ we have*

$$\zeta_{YM}(E \oplus \epsilon^k, m - 2) \leq r + k.$$

On the other hand for $l = 2$, or any l of the form $l = \sum_{1 \leq i \leq r} |d_i|^2$, with $d_i \in \mathbb{Z}$, $\sum_i d_i = 0$, there is a complex rank r vector bundle E over S^{m_l} , with $\zeta_{YM}(E, m_l - 2) = l$.

1.2. Gromov norm. Given a unitary connection A on a rank r complex vector bundle $E \rightarrow X$, a metric g on X , and a fixed bi-invariant metric on $\mathfrak{u}(r)$, we can define the norm of its curvature R_A by

$$|R_A|_g \equiv \sup_{|v \wedge w|_g = 1} |R_A(v, w)|,$$

where R_A is considered as a $\mathfrak{u}(r)$ valued 2-form. Then 1.1 gives another very different in nature proof of the following theorem of Gromov.

Theorem 1.6 ([8]). *For a simply connected Riemannian manifold X, g , and E a non-trivial Hermitian vector bundle over X , we have*

$$\inf_A |R_A|_g > 0.$$

We can sharpen it as follows.

Theorem 1.7. *For X, E, g as above:*

$$\inf_A |R_A|_g \geq \inf_A \sup_{n, [f], f' \in [f], s \in S^n} \frac{|f'(s)|_{A, \infty}}{\text{area}_g(f'(s))} > 0,$$

with the supremum over all $[f] \in \pi_n(\Omega^2 X)$, s.t. the map $\Omega^2 X \rightarrow \Omega^2 BU(r)$ is non-vanishing on $[f]$, and with area_g denoting the area functional with respect to g .

This is proved using Chern-Weil theory, in connection with our approach to jumping curves via Yang-Mills theory.

2. ACKNOWLEDGEMENTS

I am grateful to Fran Presas for related discussions, Dietmar Salamon for providing some references, and Egor Shelukhin for supportive comments. This work was partially completed during my stay at CRM Barcelona and ICMAT Madrid, under funding of Viktor Ginzburg laboratory and Severo Ochoa project, I am very grateful for the support.

3. PRELIMINARIES

We briefly collect here some principal points of Yang-Mills theory, and set up our conventions and notation. Our notation will mostly follow [3] and [1].

The Yang-Mills functional denoted by YM is defined on the space \mathcal{A} of G connections on a principal G bundle P over a Riemann manifold (X, g) by:

$$YM(A) = \int_X \|F_A\|^2 d\text{vol}_g,$$

for A a G connection, and F_A the curvature 2-form on X , with values in $\text{Ad } P = P \times_{\text{Ad } G} \mathfrak{g}$. The norm $\|\cdot\|$ above is the natural norm induced by the metric g on X and a fixed bi-invariant inner product on \mathfrak{g} , inducing a metric on $P \times_{\text{Ad } G} \mathfrak{g}$.

In this note G will be $U(r)$, sometimes $SU(r)$, and $X = S^2$. The value $YM(A)$, will be called *Yang-Mills energy* of a connection and critical points will be called *Yang-Mills connections*. For $X = S^2$ or more generally a Riemann surface, the negative gradient flow for YM exists for all time on the space of smooth connections, and moreover gradient lines converge to Yang-Mills connections. We shall call the negative gradient flow for YM on \mathcal{A} , with respect to the natural L^2 metric, the *Yang-Mills flow*. The functional YM and the metric are equivariant with respect to the action of the gauge group of P , i.e. the group of its G bundle automorphisms, and the flow is defined on the space of gauge equivalence classes. A crucial property of the Yang-Mills flow is that it preserves the orbit of A under the complexified gauge group. In our case P is associated to some complex vector bundle E , and this means that Yang-Mills flow preserves the holomorphic isomorphism type of (E, ∂_A) .

For the proof of 1.5 we also need some kind of Morse homology for YM . Much additional work is required to set up Morse-Smale-Witten complex on the space of gauge equivalence classes of connections. For $X = S^2$ this is started in [3]. Note

however that unlike [3] it is critical for us to work with the reduced gauge group which we call $\mathcal{G}_0(P)$ and which consists of bundle automorphisms fixing the fiber over the base point $0 \in S^2$, as this group acts freely. When we say *gauge equivalence class*, we mean the equivalence class under the action of \mathcal{G}_0 , as opposed to the full gauge group. In this case *YM* on $\mathcal{A}/\mathcal{G}_0$ only gives rise to a Morse-Bott complex, full details of this appear in [17].

4. PROOFS

We need a few preliminary steps. Fix a rank r Hermitian vector bundle E over $\mathbb{C}\mathbb{P}^1$. Let $\mathcal{G}_0(E)$ denote the group of bundle automorphisms of E , fixing the fiber over 0. Denote by $\mathcal{A}(E)$ the space of unitary connection on E . It is easy to check that the group $\mathcal{G}_0(E)$ acts freely on $\mathcal{A}(E)$. Elements of $\mathcal{A}(E)/\mathcal{G}_0(E)$ will be called *gauge equivalence classes* of connections. Let also \mathcal{E} denote the universal rank r Hermitian vector bundle over $BU(r)$, and let $\Omega^{2,[u]}BU(r)$ denote the smooth spherical mapping space in the component of a map $u : S^2 \rightarrow BU(r)$. To make sense of smooth here as well as of smooth connections on the universal bundle to be used later, we only need to observe that $BU(r)$ has a homotopy model that is a direct limit of smooth manifolds, $BU(r) \simeq \lim_n Gr_{\mathbb{C}}(r, \mathbb{C}^n)$, with maps in the directed system also smooth. This will be sufficient to make sense of what follows. For $v \in \Omega^{2,[u]}BU(r)$ set $v^*\mathcal{E} = \mathcal{E}_v$.

Lemma 4.1. *Given a smooth unitary connection A on the universal bundle \mathcal{E} over $BU(r)$, there is a natural induced map $W_{A,u} : \Omega^{2,[u]}BU(r) \rightarrow \mathcal{A}(\mathcal{E}_u)/\mathcal{G}_0(\mathcal{E}_u)$.*

Proof. For $v \in \Omega^{2,[u]}BU(r)$ the bundles \mathcal{E}_u and \mathcal{E}_v are isomorphic via a bundle map fixing the fiber over 0. We may get such an isomorphism by fixing a path m from u to v and fixing an auxiliary connection on the induced bundle $\mathcal{E}_m \rightarrow \mathbb{C}\mathbb{P}^1 \times [0, 1]$, trivial over $\{0\} \times [0, 1]$.

So we may define

$$W_{A,u}(v) \equiv iso^*A_v \in \mathcal{A}(\mathcal{E}_u)/\mathcal{G}_0(\mathcal{E}_u),$$

where A_v is the connection v^*A on \mathcal{E}_v , for iso any fixed isomorphism as above. We need to check that this is well defined, i.e. independent of the choice of iso . But this is immediate for if iso' is another such isomorphism, the connections iso^*A_v , iso'^*A are obviously gauge equivalent. \square

Proposition 4.2. *$W_{A,u}$ is a homotopy equivalence for any A .*

Proof. The spaces $\Omega^{2,[u]}BU(r)$ and $\mathcal{A}(\mathcal{E}_u)/\mathcal{G}_0(\mathcal{E}_u)$ are homotopy equivalent and have the homotopy types of CW complexes, [1, Proposition 2.4], and we show that that our particular map $W_{A,u}$ is an isomorphism on homotopy groups. We show injectivity. Thus, suppose that $f'_1 = W_{A,u} \circ f_1$, $f'_2 = W_{A,u} \circ f_2$, for $f_i : S^k \rightarrow \Omega^{2,[u]}BU(r)$, are homotopic. As $\mathcal{A}(\mathcal{E}_u)/\mathcal{G}_0(\mathcal{E}_u) \simeq B\mathcal{G}_0(\mathcal{E}_u)$, to a given $f : S^k \rightarrow \mathcal{A}(\mathcal{E}_u)/\mathcal{G}_0(\mathcal{E}_u)$ we have associated principal $\mathcal{G}_0(\mathcal{E}_u)$ -bundles \mathcal{P}_f . Let \mathcal{P}_i denote the principal $\mathcal{G}_0(\mathcal{E}_u)$ bundles associated to f'_i , and let $pr : \mathcal{P}_i \rightarrow S^k$ denote the associated \mathcal{E}_u bundles, (a bundle with fiber modeled on the complex vector bundle E_u over $\mathbb{C}\mathbb{P}^1$). We may also think of \mathcal{P}_i as total spaces of rank r Hermitian vector bundles over $S^k \times S^2$. For distinction let us call them E_i . It is elementary to verify that E_i are classified by the maps $cl_i : S^k \times S^2 \rightarrow BU(r)$, induced by f_i . If $H : S^k \times [0, 1] \rightarrow \mathcal{A}(\mathcal{E}_u)/\mathcal{G}_0(\mathcal{E}_u)$ is a homotopy between f'_1, f'_2 , we get

again an induced vector bundle E_H over $S^k \times S^2 \times [0, 1]$. By observation above it's classifying map to $BU(r)$ induces a homotopy between \tilde{f}_i , and consequently a homotopy between f_i .

Note that the homotopy class of $W_{A,u}$ is clearly independent of A , consequently to show surjectivity it suffices to construct for each smooth map $f : S^k \rightarrow \mathcal{A}(\mathcal{E}_u)/\mathcal{G}_0(\mathcal{E}_u)$, a smooth connection A_f on \mathcal{E} and $\tilde{f} : S^k \rightarrow \Omega^{2,[u]}BU(r)$, s.t. $W_{A_f,u} \circ \tilde{f}$ is homotopy equivalent to f . Let $pr : P_f \rightarrow S^k$ denote the associated \mathcal{E}_u bundle, constructed as before (a bundle with fiber modeled on the Hermitian vector bundle E_u over $\mathbb{C}\mathbb{P}^1$). We now construct a smooth family of unitary connections $\{A_s\}$, $s \in S^k$, with A_s a connection on the fiber $pr^{-1}(s)$, with the property that $A_s = f(s) \in \mathcal{A}(\mathcal{E}_u)/\mathcal{G}_0(\mathcal{E}_u)$ for every s . Smooth here is in the sense that the family $\{A_s\}$ is smooth in smooth trivializations of P_f . Cover S^k by a pair of disks $\{D_i^k\}$, with intersection diffeomorphic to S^{k-1} and pick any pair of smooth sections $\{A_s\}$ over D_i^k of $f^*(\mathcal{A}(E_u))$. This sections may not coincide but they are related by a transition map

$$g : U_1 \cap U_2 \simeq S^{k-1} \rightarrow \mathcal{G}_0(\mathcal{E}_u).$$

In particular they induce as required a well defined smooth family $\{A_s\}$ on

$$P_f \simeq \mathcal{E}_u \times D^k \sqcup_g \mathcal{E}_u \times D^k,$$

with the latter being the bundle obtained by using g as a clutching map. Let $E_f \rightarrow S^k \times S^2$ be the associated Hermitian vector bundles. The family $\{A_s\}$ may be extended to a unitary connection A_f on E_f . Let $cl_f : S^k \times S^2 \rightarrow BU(r)$ be an embedding classifying E_f . We may then take any connection A on the universal bundle so that its pullback by cl_f is A_f , and we take the map \tilde{f} to be the map induced by cl_f . \square

Proof of 1.1. Suppose that E has complex rank r . Clearly we may suppose that $c_1(E) = 0$, so that $cl_E : X \rightarrow BU(r)$ vanishes on π_2 . As E is non-trivial, the classifying map $cl_E : X \rightarrow BU(r)$, must be non-vanishing on some π_k , with $k > 2$. Consequently, since X is simply connected the induced map $\tilde{cl}_E : \Omega^{2,[const]}X \rightarrow \Omega^{2,jconst}BU(r)$, is non-vanishing on some $[f] \in \pi_k(\Omega^{2,[const]}X)$ with $k > 0$. Approximate the classifying map by a smooth embedding cl_E , and push-forward the connection A to a connection on the universal bundle \mathcal{E} over the image of f_E , then extend to a connection \tilde{A} on \mathcal{E} over $BU(r)$. (There are no obstructions to existence of such an extension, it can be constructed via partitions of unity for example). Now suppose that there are no jumping curves in X for E, A in class $[u]$, then we claim that the map $\Omega^{2,[u]}X \rightarrow \Omega^{2,[const]}BU(r)$ induced by the classifying map vanishes on all homotopy groups, and consequently $\Omega^{2,[const]}X \rightarrow \Omega^{2,[const]}BU(r)$ also vanishes on all homotopy groups, which is a contradiction. Indeed let $f : S^k \rightarrow \Omega^{2,[u]}X$, $k > 1$ be a continuous map. We show that the map

$$\tilde{f} = W_{\tilde{A},[const]} \circ \tilde{cl}_E \circ f : S^k \rightarrow \mathcal{A}(\epsilon^r)/\mathcal{G}_0(\epsilon^r),$$

is null-homotopic, where $const$ denotes the constant based map $const : S^2 \rightarrow BU(r)$, and ϵ^r is shorthand for the trivial \mathbb{C}^r vector bundle \mathcal{E}_{const} . This will be a contradiction and hence conclude the proof.

Indeed let $\phi_{YM}(t)$ denote the time t flow map of the Yang-Mills flow on $\mathcal{A}(\epsilon^r)/\mathcal{G}_0(\epsilon^r)$. We need a pair of facts: the isomorphism class of the induced holomorphic structure on ϵ^r by elements $|A| \in \mathcal{A}(\epsilon^r)/\mathcal{G}_0(\epsilon^r)$ is constant along negative gradient flow lines of the Yang-Mills functional, which follows by [1, 8.12], and a critical $|A|$ for

the Yang-Mills functional (a.k.a a Yang-Mills connection) determines a non-trivial holomorphic structure on ϵ^r , so long as it is not the gauge equivalence class of a trivial connection. Consequently, as the holomorphic structure induced by each $\tilde{f}(s)$, $s \in S^k$ is trivial by assumption, the negative gradient lines $t \mapsto \phi_{YM}(t)(\tilde{f}(s))$ must converge to the gauge equivalence class of a trivial connection. Reparametrizing Yang-Mills flow, (by Yang-Mills energy), we obtain a null-homotopy of \tilde{f} . \square

Proof of 1.3. This follows readily by the proof of 1.1. \square

Proof of 1.2. As previously observed the map $\tilde{cl}_E : \Omega^2 X \rightarrow \Omega^2 BU(r)$ induced by the classifying map for E , must be non-vanishing on some $[f] \in \pi_k(\Omega^{2,[const]} X)$, with $k > 0$. Let $[u_k]$ be as in the hypothesis and suppose that there are no class $[u_k]$, holomorphic jumping curves. Let $\tilde{f} : S^k \rightarrow \Omega^{2,[u_k]}(X)$, be a representative for $iso \circ f$, for $iso : \Omega^{2,[const]} X \rightarrow \Omega^{2,[u_k]} X$ the canonical homotopy equivalence, so that \tilde{f} is in the image of the inclusion $Hol^{[u_k]}(\mathbb{CP}^1, X) \rightarrow \Omega^{2,[u_k]} X$. Consequently there are no smooth jumping curves in the image of \tilde{f} . Thus running the argument in 1.1 we get that $\tilde{cl}_E \circ \tilde{f}$ is null-homotopic, but then \tilde{cl}_E must be vanishing on the class $[f]$ as well, a contradiction. \square

Proof of 1.7. This follows by the proof of 1.1 once we note:

Lemma 4.3. *Let E, X, A, g be as in the hypothesis. Suppose $u : \mathbb{CP}^1 \rightarrow X$ is a smooth jumping curve. Then the norm of the curvature of u^*E, u^*A , $|R(u^*A)|_g$ is at least $|u|_{A,\infty}/area_g(u)$.*

Proof. Indeed for the holomorphic structure on u^*E , induced by u^*A , u^*A preserves the Grothendieck splitting of u^*E . Consequently the norm of the curvature of u^*A on the maximal weight subspace \mathcal{O}_{max} of u^*E cannot exceed the norm of curvature of u^*A on u^*E , see [7, Page 79]. This follows from “positivity” of curvature on a holomorphic vector bundle.

As $c_1(\mathcal{O}_{max}) = |u|_{A,\infty}$, by assumption, Chern-Weil theory gives a ready estimate

$$|R(u^*A)|_{\mathcal{O}_{max}}|_{u^*g} \cdot area_g(u) \geq |u|_{A,\infty}.$$

\square

\square

Proof of 1.5. As we are working in the component of $\Omega^2 X$ in the constant class we may restrict our arguments to $SU(r)$ connections on E , and $\mathcal{A}(\epsilon^r)$ will denote the space of smooth $SU(r)$ connections on ϵ^r .

Morse theory for the Yang-Mills functional YM on $\mathcal{A}^r \equiv \mathcal{A}(\epsilon^r)/\mathcal{G}_0(\epsilon^r)$ is known to be essentially equivalent to Morse theory on $\Omega SU(r)$ for the energy functional, [6], [18]. Although we run the argument with Yang-Mills functional, it will be helpful to refer to the correspondence in a few instances. Moreover the energy flow picture, as developed in great detail in [15], is probably more accessible, and may help the reader to get intuition. The correspondence is given via radial trivialization. We briefly sketch this. Given an element $|A| \in \mathcal{A}^r$ we get a loop $Rad(|A|) \in \Omega SU(r)$, by fixing an S^1 -family of rays $\{r_\theta\}$ from 0 to $\infty \in \mathbb{CP}^1$, identifying the fibers over $0, \infty$ via A -parallel transport along r_0 , and then getting a loop in $SU(r)$ via A -parallel transport along rays $\{r_\theta\}$, for $A \in |A|$, which is well defined with respect to equivalence under action of $\mathcal{G}_0(\epsilon^r)$. The functions and their flow are then shown to

behave well with respect to this correspondence. In particular critical points and their indices correspond. Thus radial trivialization of a Yang-Mills connection is an S^1 subgroup of $SU(r)$.

We shall use the construction of Morse-Bott-Smale-Witten complex for YM , in [17]. The specific complex that we use is the so called *cascade complex* originally appearing in Fraunfelder [4], in a Floer theory setting. For another discussion of the cascade complex also comparing various different approaches to Morse-Bott complex, see [2]. Very briefly given a Morse-Bott function h on a manifold M , g a metric on M , so that (h, g) is a Morse-Bott-Smale pair and $\{aux_i\}$ auxiliary Morse-Smale (with respect to g) functions on the critical manifolds $\{C_i\}$, one constructs a complex $C(h, \{aux_i\}, g)$ which is generated by critical points of $\{aux_i\}$, graded by

$$\text{index}(p) = \text{index}_{MB}(C_i) + \text{index}_{aux_i} p,$$

for $p \in \text{crit}(C_i, aux_i)$, index_{MB} the Morse-Bott index and $\text{index}_{aux_i} p$ the Morse index with respect to aux_i . The differential is obtained by count of \mathbb{R} reparametrized, Fredholm index 1, *cascade flow lines* abutting to generators. The (unbroken) cascade flow lines are formal concatenations of negative gradient trajectories

$$(\gamma_{aux_{i_1}}, \gamma_h, \gamma_{aux_{i_0}}),$$

where $\gamma_h : (-\infty, \infty) \rightarrow M$ is a negative gradient trajectory for h , $\gamma_{aux_{i_1}} : [0, \infty) \rightarrow M$ is a negative gradient trajectory for aux_{i_1} , and likewise $\gamma_{aux_{i_0}} : (-\infty, 0]$ is a negative gradient trajectory for aux_{i_0} . It is shown in [2] in finite dimensional context how to use the functions aux_i to perturb h to a Morse-Smale function h' , so that there is an index preserving correspondence between the generators of the cascade complex and the critical points of h' , and a certain correspondence between the cascade flow lines and the classical flow-lines for (h', g) . Passing to our infinite dimensional setting in principle requires a bit of care, but we leave this out as it is not very interesting. We shall say in this context that the flow of h is *cascade perturbed*.

The critical sets of YM on \mathcal{A}^r split into disjoint unions of submanifolds $\Lambda_{\mathbf{d}}$, composed of (Yang-Mills) connections, inducing the holomorphic structure of type \mathbf{d} , where \mathbf{d} by Birkhoff-Grothendieck can be taken to mean an unordered collection of r integers. The manifolds $\Lambda_{\mathbf{d}}$, which are isomorphic to the manifolds of S^1 subgroups of $SU(r)$, conjugate to the diagonal S^1 subgroup with weights \mathbf{d} , are certain complex flag manifolds and admit perfect Morse functions. Consequently, the homology of \mathcal{A}^r is computed by the cascade Morse-Bott-Smale-Witten complex, with vanishing differentials, as the Morse-Bott index in this case is always even, (see elaboration below). We will not mention the auxiliary Morse-Smale functions on the critical manifolds explicitly, and just write $\mathcal{C}_*(YM)$, for this complex.

Up to degree $2r - 2$ all the generators of $\mathcal{C}_*(YM)$, come from critical manifolds $\Lambda_{\mathbf{d}}$, with \mathbf{d} having all weights 1, -1 or 0. This readily follows upon computing the Morse-Bott index along $\Lambda_{\mathbf{d}}$, (i.e. the normal component of Morse index). As the Morse index for YM of a critical $|A| \in \mathcal{A}^r$, coincides with the Morse index of $\text{rad}(|A|) \in \Omega SU(r)$, we may use the classical calculation of Morse index of geodesics in $SU(r)$, see for example [13]. This tells us that for a homomorphism

$\gamma : S^1 \rightarrow SU(r)$, conjugate to

$$\gamma = \begin{pmatrix} e^{2\pi i a_1} & & \\ & e^{2\pi i a_2} & \\ & & \dots \end{pmatrix}$$

with weights a_1, \dots, a_r , organized so that $a_1 \geq a_2 \geq \dots \geq a_r$ the Morse index is given by

$$\sum_{i < j} 2|a_i - a_j| - 2.$$

It is then easy to verify that if $\mathbf{d} = \{a_1, \dots, a_r\}$ does not satisfy that all a_i are either 1, -1 or 0, then the Morse index is strictly greater than $2r - 2$, see [13, page 132] for a similar calculation.

We may then use unstable manifolds of the cascade perturbed Yang-Mills flow, to induce a cellular decomposition of \mathcal{A}^r , with all cells of even dimension. Likewise there is a dual stratification of \mathcal{A}^r by finite co-dimension strata, intersecting the cascade perturbed unstable manifolds transversally, corresponding to cascade perturbed stable manifolds for YM . It follows that given a map $f : S^k \rightarrow \mathcal{A}^r$ with $k \leq 2r - 2$, after a small perturbation of f the (reparametrized by energy) (cascade perturbed) Yang-Mills flow will give a homotopy of f into the $2r - 2$ skeleton of \mathcal{A}^r . That is, we just have to take a perturbation of f so that we are transverse to the finite codimension strata. This perturbation exists by general differential topology considerations. Let's now apply this.

Following the proof of 1.1, we have a map

$$\Omega^2 S^m \rightarrow \mathcal{A}^{r+k}$$

induced by the classifying map for $E \oplus \epsilon^k$. Take k so that $n \leq 2(r + k) - 2$. It follows by discussion above that the composition

$$\tilde{f} = W_{\tilde{A}, [\text{const}]} \circ \tilde{cl}_{E \oplus \epsilon^k} \circ f : S^n \rightarrow \mathcal{A}^{r+k}$$

can be perturbed so that the energy flow (after reparametrization) takes \tilde{f} into the $2(r + k) - 2$ skeleton of \mathcal{A}^{r+k} .

Moreover we have:

Lemma 4.4. *Any abstract perturbation of \tilde{f} , can be obtained via perturbation of the connection A on E .*

If we accept this for the moment then the first part of the theorem follows, as the Birkhoff-Grothendieck splitting type of elements in the $2(r + k) - 2$ skeleton, by preliminary discussion above, is such that all weights are either 1, -1 or 0. Consequently, (after perturbation of A) $|f(s)|_A \leq r + k$ for each $s \in S^n$.

Proof of Lemma 4.4. Let us then verify the claim about the perturbation. This is where the assumption $X = S^n$ comes into the proof. Take a representative $f : S^n \rightarrow \Omega^2 S^m$, for $[f]$ so that the induced map $g : S^n \times S^2 \rightarrow S^m$ is an embedding outside the submanifolds $S^n \times \{0\}$, $\{0\} \times S^2$ which are collapsed to a point. Given an abstract perturbation \tilde{f}' of \tilde{f} , with $\tilde{F} : S^n \times I \rightarrow \mathcal{A}^{r+k}$ a homotopy between \tilde{f} , and \tilde{f}' , we get an associated structure group $\mathcal{G}_0(\epsilon^{r+k})$ bundle

$$\pi_{\tilde{F}} : P_{\tilde{F}} \rightarrow S^n \times [0, 1]$$

with fiber $\mathbb{C}^{r+k} \times S^2$. We also get a bundle $G_{\tilde{F}} \rightarrow S^n \times [0, 1]$, with fiber over (s, t) the space of connections on $\pi_{\tilde{F}}^{-1}(s, t)$, in the gauge equivalence class of $\tilde{F}(s, t)$. Note that this is in general not isomorphic to the principal $\mathcal{G}_0(\epsilon^{r+k})$ bundle pulled back by \tilde{F} . By construction $G_{\tilde{F}}$ has a section over $S^n \times \{0\}$, use homotopy lifting property to obtain a section $\mathcal{S}_{\tilde{F}}$ over $S^n \times \{1\}$. $\mathcal{S}_{\tilde{F}}$ can be thought of as smooth family of connections $\{A_s\}$ over $\{\{s\} \times S^2 \subset S^n \times S^2\}$, for the \mathbb{C}^{r+k} bundle $E_{\tilde{F}} \rightarrow S^n \times S^2$ canonically associated to $P_{\tilde{F}}$. Note that $E_{\tilde{F}}$ is canonically trivial over $S^n \times \{0\}$, $\{0\} \times S^2$, consequently we may extend $\{A_s\}$ to a connection A'' on $E_{\tilde{F}} \rightarrow S^n \times S^2$, trivial over $S^n \times \{0\}$, $\{0\} \times S^2$. (The extension may be obtained via classical partition of unity argument.) It follows that we may push forward A'' via g to a connection A' on E , with exactly the right property. \square

We now verify the second part of the theorem. We will say that a skeleton of \mathcal{A}^r (in the previous sense) has energy l , if all the generators of $\mathcal{C}_*(YM)$ in it have YM energy less then or equal to l . As $\mathcal{C}_*(YM)$ is perfect, skeleta with distinct energy have distinct homology. From this it follows that a skeleton \mathcal{S} with energy l admits a non null-homotopic based map $f : S^{m_l-2} \rightarrow \mathcal{S}$, which cannot be pushed below the energy level l . As otherwise \mathcal{S} could be compressed into a lower energy skeleton, which would contradict the above. (The compression argument is just Whitehead's theorem, see for example [11]). As the map $W_{A, const}$ is a homotopy equivalence for every A , we may find a $g : S^{m_l-2} \rightarrow \Omega^2 BU(r)$, s.t. $W_{A, const_*}[g] = [f]$. Following the proof of Lemma 4.4 we get a family $\{A'_s\}$ of connections on $\{E'_f|_{\{s\} \times S^2}\}$ where E'_f is the \mathbb{C}^r vector bundle over $S^{m_l-2} \times S^2$ associated to f as in the proof of 4.4. Extend $\{A'_s\}$ to a connection A'_f on E_f , trivial over $S^{m_l-2} \times \{0\}$, where the bundle is canonically trivialized. By construction A'_f is also the trivial connection over $\{0\} \times S^2$. It follows that E'_f induces a vector bundle over

$$S^{m_l} = S^{m_l-2} \vee S^2,$$

and A'_f induces a connection A_f on E_f . By construction the infimum in $\zeta_{YM}(E_f, m_l)$ is attained on A_f .

The proof is finished as the normalization for the bi-invariant metric on $SU(r)$ that we take, is such that an S^1 geodesic with weights $\{d_i\}$, has energy given by (1.1). \square

REFERENCES

- [1] M. F. ATIYAH AND R. BOTT, *The Yang-Mills equations over Riemann surfaces.*, Philos. Trans. R. Soc. Lond., A, 308 (1983), pp. 523–615.
- [2] A. BANYAGA AND D. E. HURTUBISE, *Cascades and perturbed Morse-Bott functions.*, Algebr. Geom. Topol., 13 (2013), pp. 237–275.
- [3] T. J. DAVIES, *The Yang-Mills functional over Riemann surfaces and the loop group*, thesis, (1996).
- [4] U. FRAUENFELDER, *The Arnold-Givental conjecture and moment Floer homology.*, Int. Math. Res. Not., 2004 (2004), pp. 2179–2269.
- [5] J. GRAVESEN, *On the topology of spaces of holomorphic maps*, Acta Math., 162 (1989), pp. 247–286.
- [6] J. GRAVESEN, *Loop groups and Yang-Mills theory in dimension two.*, Commun. Math. Phys., 127 (1990), pp. 597–605.
- [7] P. GRIFFITHS AND J. HARRIS, *Principles of algebraic geometry*, Wiley Classics Library, John Wiley & Sons Inc., New York, 1994. Reprint of the 1978 original.

- [8] M. GROMOV, *Positive curvature, macroscopic dimension, spectral gaps and higher signatures*. Gindikin, Simon (ed.) et al., Functional analysis on the eve of the 21st century. Volume II. In honor of the eightieth birthday of I. M. Gelfand. Proceedings of a conference, held at Rutgers University, New Brunswick, NJ, USA, October 24-27, 1993. Boston, MA: Birkhäuser. Prog. Math. 132, 1-213 (1996).
- [9] M. A. GUEST, *Topology of the space of absolute minima of the energy functional*, Amer. J. Math., 106 (1984), pp. 21–42.
- [10] M. A. GUEST, *The topology of the space of rational curves on a toric variety*, Acta Math., 174 (1995), pp. 119–145.
- [11] A. HATCHER, *Algebraic topology.*, Cambridge: Cambridge University Press, 2002.
- [12] F. KIRWAN, *On spaces of maps from Riemann surfaces to Grassmannians and applications to the cohomology of moduli of vector bundles.*, Ark. Mat., 24 (1986), pp. 221–275.
- [13] J. MILNOR, *Morse theory*, Based on lecture notes by M. Spivak and R. Wells. Annals of Mathematics Studies, No. 51, Princeton University Press, Princeton, N.J., 1963.
- [14] C. OKONEK, M. SCHNEIDER, AND H. SPINDLER, *Vector bundles on complex projective spaces. With an appendix by S. I. Gelfand. Corrected reprint of the 1988 edition.*
- [15] A. PRESSLEY AND G. SEGAL, *Loop groups*, Oxford Mathematical Monographs, (1986).
- [16] G. SEGAL, *The topology of spaces of rational functions*, Acta Math., 143 (1979), pp. 39–72.
- [17] J. SWOBODA, *Morse Homology for the Yang-Mills Gradient Flow*, [arXiv:1103.0845](#), (2011).
- [18] ———, *The Yang-Mills gradient flow and loop spaces of compact lie groups*, [arXiv:1104.5514](#), (2012).

CALLE NICOLAS CABRERA 13 N 13-15 ICMAT, 28049 MADRID
E-mail address: yasha.savelyev@gmail.com