

ON THE EQUIVALENCE BETWEEN LOW RANK MATRIX COMPLETION AND TENSOR RANK

HARM DERKSEN

ABSTRACT. The Rank Minimization Problem asks to find a matrix of lowest rank inside a linear variety of the space of $n \times m$ matrices. The Low Rank Matrix Completion problem asks to complete a partially filled matrix such that the resulting matrix has smallest possible rank. The Tensor Rank Problem asks to determine the rank of a tensor. We show that these three problems are equivalent: each one of the problems can be reduced to the other two.

1. INTRODUCTION

Suppose that \mathbb{F} is a field. We will consider the following computational problems:

Rank Minimization (RM). Given matrices $A, B_1, \dots, B_s \in \text{Mat}_{n,m}(\mathbb{F})$, find $x_1, \dots, x_s \in \mathbb{F}$ for which $\text{rank}(A + x_1 B_1 + \dots + x_s B_s)$ is minimal.

Special Rank Minimization (1–RM). This problem is the same as the Rank Minimization Problem, but we assume that the matrices B_1, \dots, B_s have rank 1.

Low Rank Matrix Completion (LRMC). Given an $n \times m$ matrix A which is partially filled with entries of \mathbb{F} , fill in the remaining entries such that the resulting matrix has minimal rank.

Tensor Rank (TR). Given a tensor $T \in \mathbb{F}^{n_1} \otimes \mathbb{F}^{n_2} \otimes \dots \otimes \mathbb{F}^{n_d}$ Find pure tensors $v_j = v_j^{(1)} \otimes \dots \otimes v_j^{(d)}$ for $j = 1, 2, \dots, r$ such that $T = v_1 + \dots + v_r$ and r is minimal. (This minimal r is called the *tensor rank* of T , see [17].)

Order 3 Tensor Rank (3–TR). This is the Tensor Rank problem where the order d is equal to 3.

The LRMC problem has many applications. One such application is *collaborative filtering*. A typical example is the *Netflix problem*. Suppose that the rows of a matrix correspond to users, and the columns correspond to movies. Each user has only rated some of the movies. This leads to a partially filled matrix. To predict whether users would like a certain movie that they have not seen yet, one would like to complete the matrix. Because preferences only depend on a few parameters, it is reasonable to assume that the rank of the completed matrix should be small. The LRMC problem is known to be NP complete (see [26, Theorem 3.1]). For the field $\mathbb{F} = \mathbb{R}$, some algorithms have been proposed for the LRMC problem using convex relaxation, for example [10, 6, 4, 6].


If the matrices B_1, \dots, B_s in the Rank Minimization problems are matrices that have only one nonzero entry, then the RM problem becomes an LRMC problem. So the RM problem is

The author was partially supported by NSF grant DMS 1302032.

more general. Over finite fields, RM has applications to coding theory ([1, 28]). For the real numbers, applications are in control ([11, 12, 13]), systems identification ([24]) and Euclidean embedding problems ([23, 18]). Some algorithms have appeared in [11, 27, 24, 4, 25].

The TR problem is NP-complete if the field \mathbb{F} is finite ([15, 16]) and NP-hard if the field is \mathbb{Q} ([15, 16]), or if it contains \mathbb{Q} ([19]). Approximation of a tensor by another tensors of low rank is known as the PARAFAC ([14]) or CANDECOP ([5]) model. There are various heuristic approaches for finding low-rank approximations. See [22, 29] for a discussion. The problems of finding the rank of a tensor, and to approximate tensors with tensors of low rank has many applications, such as the complexity of matrix multiplication, fluorescence spectroscopy, statistics, psychometrics, geophysics and magnetic resonance imaging. In general, the rank of higher tensors more ill-behaved than the rank of a matrix (see [9]).

We will show that these 5 problems are equivalent to each other by giving explicit reductions. Some reductions are obvious: 1–RM is a special case of RM, 3–TR is a special case of TR and LRMC is a special case of 1–RM. The reduction from LRMC to 3–TR was done in [8]. Other reductions are explicitly described in this paper and these reductions are denoted by double arrows. By composition of reduction steps, we can reduce each of the 5 problems to any other problem. In [2] the authors give an algorithm for tensor decompositions and more generally, multi-homogeneous tensors (see also [3] for symmetric tensor decompositions). A crucial step (Algorithm 3.1, step (1)) in their algorithm asks for a low rank completion of a Hankel matrix (Theorem 3.3.), so their algorithm can also be viewed as a reduction from TR to RM.



2. REDUCTION FROM RM TO 1–RM

We first reduce the Rank Minimization Problem with $s = 1$ to the Special Rank Minimization Problem. Suppose that $A, B \in \text{Mat}_{n,m}(\mathbb{F})$ and we would like to minimize $\text{rank}(A + xB)$ for $x \in \mathbb{F}$. Let $r = \text{rank } B$. We can write

$$B = \sum_{i=1}^r v_i w_i^t$$

where $v_1, \dots, v_r \in \mathbb{F}^n$ and $w_1, \dots, w_r \in \mathbb{F}^m$. We introduce new variables:

$$\begin{aligned} x_i, & \quad 1 \leq i \leq r \\ y_{i,j}, & \quad 1 \leq i, j \leq r \text{ and } i \neq j \\ z_{i,j}, & \quad 1 \leq i < j \leq r. \end{aligned}$$

If $i > j$, then we define $z_{i,j} = z_{j,i}$. We will not have such a convention for the y 's, so $y_{i,j}$ and $y_{j,i}$ are distinct variables. Define

$$\begin{aligned}\mathbf{x} &= (x_1, \dots, x_r) \\ \mathbf{y} &= (y_{i,j} \mid i \neq j, 1 \leq i, j \leq r) \\ \mathbf{z} &= (z_{i,j} \mid 1 \leq i < j \leq r).\end{aligned}$$

For $1 \leq i < j \leq r$ define the $n \times 2$ matrices

$$C_{i,j}(\mathbf{x}, \mathbf{y}) := ((y_{i,j} - x_i)v_i \quad (y_{j,i} - x_j)v_j)$$

and the 1×2 matrices

$$D_{i,j}(\mathbf{y}, \mathbf{z}) := (y_{i,j} - z_{i,j} \quad y_{j,i} - z_{j,i}).$$

Also, define an $n \times m$ matrix

$$E(\mathbf{x}) := A + \sum_{i=1}^r x_i v_i w_i^t.$$

Let J be the matrix

$$J(\mathbf{x}, \mathbf{y}, \mathbf{z}) := \left(\begin{array}{c|cccc} E(\mathbf{x}) & C_{1,2}(\mathbf{x}, \mathbf{y}) & C_{1,3}(\mathbf{x}, \mathbf{y}) & \cdots & C_{r-1,r}(\mathbf{x}, \mathbf{y}) \\ \hline 0 & D_{1,2}(\mathbf{y}, \mathbf{z}) & 0 & \cdots & 0 \\ 0 & 0 & D_{1,3}(\mathbf{y}, \mathbf{z}) & & 0 \\ \vdots & \vdots & & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & D_{r-1,r}(\mathbf{y}, \mathbf{z}) \end{array} \right)$$

The size of J is $(n + \frac{1}{2}r(r-1)) \times (m + r(r-1))$.

We will use the convention $C_{i,j} := C_{j,i}$ and $D_{i,j} := D_{j,i}$ if $i > j$.

Theorem 1. *For every $\mathbf{x} = (x_1, \dots, x_r) \in \mathbb{F}^r$, $\mathbf{y} \in \mathbb{F}^{r(r-1)}$, $\mathbf{z} \in \mathbb{F}^{r(r-1)/2}$ one can choose $x \in \{x_1, \dots, x_r\}$ such that*

$$\text{rank } J(\mathbf{x}, \mathbf{y}, \mathbf{z}) \geq \text{rank}(A + xB).$$

In particular,

$$\min_{\mathbf{x}, \mathbf{y}, \mathbf{z}} \text{rank } J(\mathbf{x}, \mathbf{y}, \mathbf{z}) = \min_x \text{rank}(A + xB).$$

Proof. Let

$$Z := \{i \mid y_{i,j} \neq z_{i,j} \text{ or } x_i \neq y_{i,j} \text{ for some } j\}.$$

Choose a $j \in \{1, \dots, r\}$ such that either $j \notin Z$ or $Z = \{1, \dots, r\}$. Define $x := x_j$. Suppose that $x_i \neq x$. If $Z = \{1, \dots, r\}$ then we have $i \in Z$. Otherwise, $j \notin Z$ and $x_i \neq x_j = y_{j,i} = z_{j,i} = z_{i,j}$. So $x_i \neq y_{i,j}$ or $y_{i,j} \neq z_{i,j}$ and it follows that $i \in Z$. We conclude that $x_i \neq x$ implies $i \in Z$.

If $y_{i,j} \neq x_i$, then $C_{i,j}$ contains column that is nonzero multiple v_i . If $y_{i,j} \neq z_{i,j}$ then we wipe $C_{i,j}$ with $D_{i,j}$ using elementary row operations to obtain a matrix $C'_{i,j}$ which has a column that is a nonzero multiple of v_i . So after some elementary row operations, we obtain a matrix

$$J' = \left(\begin{array}{c|cccc} E & C'_{1,2} & C'_{1,3} & \cdots & C'_{r-1,r} \\ \hline 0 & D_{1,2} & 0 & \cdots & 0 \\ 0 & 0 & D_{1,3} & & 0 \\ \vdots & \vdots & & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & D_{r-1,r} \end{array} \right) = \left(\begin{array}{c|c} E & C' \\ \hline 0 & D \end{array} \right)$$

such that for every $i \in Z$, C' has a column that is a nonzero multiple of the column vector v_i .

Using elementary column operations, we can replace the submatrix

$$E(\mathbf{x}) = E(x_1, \dots, x_r) = A + \sum_{i=1}^r x_i v_i w_i^t$$

of J' by

$$E(x, x, \dots, x) = A + x \sum_{i=1}^r v_i w_i^t = A + xB = E(\mathbf{x}) + \sum_{i \in Z} (x - x_i) v_i w_i^t.$$

So after elementary row and column operations, $A + xB$ becomes a submatrix of $J(\mathbf{x}, \mathbf{y}, \mathbf{z})$. This proves that

$$\text{rank } J(\mathbf{x}, \mathbf{y}, \mathbf{z}) \geq \text{rank}(A + xB).$$

If we set $x_i = y_{i,j} = z_{i,j} = x$ for all i, j , then we get

$$J = \begin{pmatrix} A + xB & 0 \\ 0 & 0 \end{pmatrix}.$$

This proves that

$$\min_{\mathbf{x}, \mathbf{y}, \mathbf{z}} \text{rank } J(\mathbf{x}, \mathbf{y}, \mathbf{z}) = \min_x \text{rank}(A + xB).$$

□

We can write J in the form

$$J(\mathbf{x}, \mathbf{y}, \mathbf{z}) = \begin{pmatrix} A & 0 \\ 0 & 0 \end{pmatrix} + \sum_{i=1}^r x_i F_i + \sum_{i \neq j} y_{i,j} G_{i,j} + \sum_{i < j} z_{i,j} H_{i,j}.$$

The crucial point here is that $F_i, G_{i,j}, H_{i,j}$ all have rank 1. In fact, $G_{i,j}$ has only one nonzero column, and $H_{i,j}$ has only one nonzero row.

Example 2. Suppose that

$$A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Let

$$v_1 = w_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad v_2 = w_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

so that

$$B = v_1 w_1^t + v_2 w_2^t = \begin{pmatrix} 1 \\ 0 \end{pmatrix} (1 \ 0) + \begin{pmatrix} 0 \\ 1 \end{pmatrix} (0 \ 1).$$

Define

$$J := \left(\begin{array}{c|c} E & C_{1,2} \\ \hline 0 & D_{1,2} \end{array} \right)$$

where

$$E := A + x_1 v_1 w_1^t + x_2 v_2 w_2^t = \begin{pmatrix} x_1 & 1 \\ -1 & x_2 \end{pmatrix},$$

$$C_{1,2} := ((y_{1,2} - x_1)v_1 \ (y_{2,1} - x_2)v_2) = \begin{pmatrix} y_{1,2} - x_1 & 0 \\ 0 & y_{2,1} - x_2 \end{pmatrix}$$

and

$$D_{1,2} := (y_{1,2} - z_{1,2} \quad y_{2,1} - z_{1,2})$$

So we have

$$J = \left(\begin{array}{cc|cc} x_1 & 1 & y_{1,2} - x_1 & 0 \\ -1 & x_2 & 0 & y_{2,1} - x_2 \\ \hline 0 & 0 & y_{1,2} - z_{1,2} & y_{2,1} - z_{1,2} \end{array} \right).$$

We can write

$$\begin{aligned} J = & \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} + x_1 \begin{pmatrix} 1 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} + x_2 \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 0 \end{pmatrix} + y_{1,2} \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} + \\ & + y_{2,1} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix} + z_{1,2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & -1 \end{pmatrix}. \end{aligned}$$

Example 3. Suppose that

$$A = \begin{pmatrix} 0 & 3 & 4 \\ 0 & 0 & 5 \\ 0 & 0 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 2 & 1 \end{pmatrix}$$

We can write

$$B = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix} \begin{pmatrix} 0 & 1 & 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \begin{pmatrix} 0 & 0 & 1 \end{pmatrix}$$

We have

$$E = \begin{pmatrix} x_1 & 3 & 4 \\ 2x_1 & x_2 & 5 \\ 3x_1 & 2x_2 & x_3 \end{pmatrix},$$

$$\begin{aligned} C = (C_{1,2} \quad C_{1,3} \quad C_{2,3}) = \\ = \begin{pmatrix} y_{1,2} - x_1 & 0 & y_{1,3} - x_1 & 0 & 0 & 0 \\ 2y_{1,2} - 2x_1 & y_{2,1} - x_2 & 2y_{1,3} - 2x_1 & 0 & y_{2,3} - x_2 & 0 \\ 3y_{1,2} - 3x_1 & 2y_{2,1} - 2x_2 & 3y_{1,3} - 3x_1 & y_{3,1} - x_3 & 2y_{2,3} - 2x_2 & y_{3,2} - x_3 \end{pmatrix} \end{aligned}$$

and

$$\begin{aligned} D = \begin{pmatrix} D_{1,2} & 0 & 0 \\ 0 & D_{1,3} & 0 \\ 0 & 0 & D_{2,3} \end{pmatrix} = \\ = \begin{pmatrix} y_{1,2} - z_{1,2} & y_{2,1} - z_{1,2} & 0 & 0 & 0 & 0 \\ 0 & 0 & y_{1,3} - z_{1,3} & y_{3,1} - z_{1,3} & 0 & 0 \\ 0 & 0 & 0 & 0 & y_{2,3} - z_{2,3} & y_{3,2} - z_{2,3} \end{pmatrix} \end{aligned}$$

So we have

$$J = \left(\begin{array}{ccc|cccccc} x_1 & 3 & 4 & y_{1,2} - x_1 & 0 & y_{1,3} - x_1 & 0 & 0 & 0 \\ 2x_1 & x_2 & 5 & 2y_{1,2} - 2x_1 & y_{2,1} - x_2 & 2y_{1,3} - 2x_1 & 0 & y_{2,3} - x_2 & 0 \\ 3x_1 & 2x_2 & x_3 & 3y_{1,2} - 3x_1 & 2y_{2,1} - 2x_2 & 3y_{1,3} - 3x_1 & y_{3,1} - x_3 & 2y_{2,3} - 2x_2 & y_{3,2} - x_3 \\ \hline 0 & 0 & 0 & y_{1,2} - z_{1,2} & y_{2,1} - z_{1,2} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & y_{1,3} - z_{1,3} & y_{3,1} - z_{1,3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & y_{2,3} - z_{2,3} & y_{3,2} - z_{2,3} \end{array} \right)$$

We will formulate the general rank minimization problem with upper indices, so that we can reserve lower indices for other uses. Suppose $A, B^{(1)}, \dots, B^{(s)} \in \text{Mat}_{m,n}(\mathbb{F})$ and we want to minimize

$$\text{rank}(A + x^{(1)}B^{(1)} + \dots + x^{(s)}B^{(s)}).$$

over all $x^{(1)}, \dots, x^{(s)} \in \mathbb{F}$. Inductively we construct a matrices by

$$J_0(A) = A.$$

and

$$J_i(A, B^{(1)}, \dots, B^{(i)}, \mathbf{x}^{(1)}, \dots, \mathbf{x}^{(i)}, \mathbf{y}^{(1)}, \dots, \mathbf{y}^{(i)}, \mathbf{z}^{(1)}, \dots, \mathbf{z}^{(i)}) := J(J_{i-1}(A, B^{(1)}, \dots, B^{(i-1)}, \mathbf{x}^{(1)}, \dots, \mathbf{x}^{(i-1)}, \mathbf{y}^{(1)}, \dots, \mathbf{y}^{(i-1)}, \mathbf{z}^{(1)}, \dots, \mathbf{z}^{(i-1)}), B^{(i)}, \mathbf{x}^{(i)}, \mathbf{y}^{(i)}, \mathbf{z}^{(i)}).$$

for $i = 1, 2, \dots, s$. Inductively, we can find $x^{(s)}, x^{(s-1)}, \dots, x^{(1)}$ (in that order) such that

$$\begin{aligned} \text{rank } J_i(A + x^{(i+1)}B^{(i+1)} + \dots + x^{(s)}B^{(s)}, B^{(1)}, \dots, B^{(i)}, \mathbf{x}^{(1)}, \dots, \mathbf{x}^{(i)}, \mathbf{y}^{(1)}, \dots, \mathbf{y}^{(i)}, \mathbf{z}^{(1)}, \dots, \mathbf{z}^{(i)}) &\geq \\ \geq \text{rank } J_{i-1}(A + x^{(i)}B^{(i)} + \dots + x^{(s)}B^{(s)}, \mathbf{x}^{(1)}, \dots, \mathbf{x}^{(i-1)}, \mathbf{y}^{(1)}, \dots, \mathbf{y}^{(i-1)}, \mathbf{z}^{(1)}, \dots, \mathbf{z}^{(i-1)}). \end{aligned}$$

for all i . Combining all the inequalities together gives

$$\begin{aligned} \text{rank } J_s(A, B^{(1)}, \dots, B^{(s)}, \mathbf{x}^{(1)}, \dots, \mathbf{x}^{(s)}, \mathbf{y}^{(1)}, \dots, \mathbf{y}^{(s)}, \mathbf{z}^{(1)}, \dots, \mathbf{z}^{(s)}) &\geq \\ \text{rank } J_0(A + x^{(1)}B^{(1)} + \dots + x^{(s)}B^{(s)}) = \text{rank}(A + x^{(1)}B^{(1)} + \dots + x^{(s)}B^{(s)}). \end{aligned}$$

If we set all the entries of $\mathbf{x}^{(i)}, \mathbf{y}^{(i)}, \mathbf{z}^{(i)}$ equal to $x^{(i)}$ for all i , then we have equality.

Example 4. Suppose that

$$A = \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, B^{(1)} = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, B^{(2)} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, B^{(3)} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}.$$

so that

$$A + x^{(1)}B^{(1)} + x^{(2)}B^{(2)} + x^{(3)}B^{(3)} = \begin{pmatrix} -1 & x^{(1)} & x^{(2)} \\ -x^{(1)} & -1 & x^{(3)} \\ -x^{(2)} & -x^{(3)} & -1 \end{pmatrix}.$$

We write

$$\begin{aligned}
B^{(1)} &= \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} (0 \ 1 \ 0) + \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} (-1 \ 0 \ 0), \\
B^{(2)} &= \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} (0 \ 0 \ 1) + \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} (-1 \ 0 \ 0), \\
B^{(3)} &= \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} (0 \ 0 \ 1) + \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} (0 \ -1 \ 0).
\end{aligned}$$

We get

$$J_3 = \left(\begin{array}{ccc|cc|cc|cc}
-1 & x_1^{(1)} & x_1^{(2)} & y_{1,2}^{(1)} - x_1^{(1)} & 0 & y_{1,2}^{(2)} - x_1^{(2)} & 0 & 0 & 0 \\
-x_2^{(1)} & -1 & x_1^{(3)} & 0 & y_{2,1}^{(1)} - x_2^{(1)} & 0 & 0 & y_{1,2}^{(3)} - x_1^{(3)} & 0 \\
-x_2^{(2)} & -x_2^{(3)} & -1 & 0 & 0 & 0 & y_{2,1}^{(2)} - x_2^{(2)} & 0 & y_{2,1}^{(3)} - x_2^{(3)} \\
\hline
0 & 0 & 0 & y_{1,2}^{(1)} - z_{1,2}^{(1)} & y_{2,1}^{(1)} - z_{1,2}^{(1)} & 0 & 0 & 0 & 0 \\
\hline
0 & 0 & 0 & 0 & 0 & y_{1,2}^{(2)} - z_{1,2}^{(2)} & y_{2,1}^{(2)} - z_{1,2}^{(2)} & 0 & 0 \\
\hline
0 & 0 & 0 & 0 & 0 & 0 & 0 & y_{1,2}^{(3)} - z_{1,2}^{(3)} & y_{2,1}^{(3)} - z_{1,2}^{(3)}
\end{array} \right).$$

3. REDUCTION FROM 1-RM TO LRMC

We first deal with the Special Rank Minimization Problem with $s = 1$. Suppose that A, B are $n \times m$ matrices, and that B has rank 1. We can write $B = vw^t$ with $v \in \mathbb{F}^n$ and $w \in \mathbb{F}^m$.

From

$$\begin{pmatrix} A + xvw^t & 0 \\ w^t & -1 \end{pmatrix} = \begin{pmatrix} I & xv \\ 0 & 1 \end{pmatrix} \begin{pmatrix} A & xv \\ w^t & -1 \end{pmatrix}$$

follows that

$$\begin{aligned}
\text{rank}(A + xB) + 1 &= \text{rank}(A + xvw^t) + 1 = \text{rank} \begin{pmatrix} A + xvw^t & 0 \\ w^t & -1 \end{pmatrix} = \\
&= \text{rank} \begin{pmatrix} A & xv \\ w^t & -1 \end{pmatrix} = \text{rank}(A' + xB'),
\end{aligned}$$

where

$$A' = \begin{pmatrix} A & 0 \\ w^t & -1 \end{pmatrix}, \quad B' = \begin{pmatrix} 0 & v \\ 0 & 0 \end{pmatrix}.$$

Note that B' has at most 1 nonzero column. If B has at most 1 nonzero row, then v will have only one nonzero entry and B' will only have one nonzero entry. In that case we have reduced to the case where B has only one nonzero entry. So we have reduced a special rank minimization problem to a matrix completion problem in this case.

Example 5. Suppose that

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 0 \\ 2 & 3 \end{pmatrix}.$$

We have

$$A' = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & 3 & -1 \end{pmatrix}, \quad B' = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

So we have reduced the problem of minimizing the rank of

$$A + xB = \begin{pmatrix} 1 & 0 \\ 2x & 1 + 3x \end{pmatrix}$$

to the matrix completion problem

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & ? \\ 2 & 3 & -1 \end{pmatrix}.$$

The minimal rank of $A + xB$ is 1, namely for $x = -\frac{1}{3}$. The smallest rank for the matrix completion problem is 2. This is the case when the missing entry is $-\frac{1}{3}$.

Suppose again that $A, B \in \text{Mat}_{n,m}(\mathbb{F})$ and $B = vw^t$ has rank 1. By symmetry, we also have

$$\begin{aligned} \text{rank}(A + xB) + 1 &= \text{rank}(A + xvw^t) + 1 = \text{rank} \begin{pmatrix} A + xvw^t & v \\ 0 & -1 \end{pmatrix} = \\ &= \text{rank} \begin{pmatrix} A & v \\ xw^t & -1 \end{pmatrix} = \text{rank}(A'' + xB''), \end{aligned}$$

where

$$A'' = \begin{pmatrix} A & v \\ 0 & -1 \end{pmatrix}, \quad B'' = \begin{pmatrix} 0 & 0 \\ w^t & 0 \end{pmatrix}.$$

Note that B' has at most 1 nonzero row. If B has at most 1 nonzero column, then w^t will have only one nonzero entry and B'' will only have one nonzero entry. In that case we have reduced the special matrix minimization problem to a matrix completion problem.

In the general case, where $B = vw^t$ has more than 1 nonzero row and more than 1 nonzero column, we can use the first construction to obtain matrices A', B' such that

$$\text{rank}(A + xB) + 1 = \text{rank}(A' + xB')$$

and B' has at most one nonzero column. Then we can use the second construction to obtain matrices A'', B'' such that

$$\text{rank}(A' + xB') + 1 = \text{rank}(A'' + xB'')$$

such that B'' has only one nonzero entry. Note that

$$\text{rank}(A + xB) + 2 = \text{rank}(A'' + xB'').$$

So we have reduced the problem of minimizing the rank of $A + xB$ to the matrix completion problem.

Example 6. Suppose that

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 2 \\ 3 & 6 \end{pmatrix} = \begin{pmatrix} 1 \\ 3 \end{pmatrix} \begin{pmatrix} 1 & 2 \end{pmatrix}.$$

If we define

$$A' = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 2 & -1 \end{pmatrix}, \quad B' = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 3 \\ 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 3 \\ 0 \end{pmatrix} \begin{pmatrix} 0 & 0 & 1 \end{pmatrix}$$

then

$$\text{rank}(A + xB) + 1 = \text{rank}(A' + xB').$$

Now, define

$$A'' = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 3 \\ 1 & 2 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad B'' = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

Then we have

$$\text{rank}(A + xB) + 2 = \text{rank}(A'' + xB'').$$

So we have reduced the problem of minimizing the rank of $A + xB$ to the low rank matrix completion problem

$$\begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 3 \\ 1 & 2 & -1 & 0 \\ 0 & 0 & ? & -1 \end{pmatrix}.$$

The smallest possible rank of this matrix is 3, when the missing entry is $-\frac{1}{7}$.

Suppose that $A \in \text{Mat}_{n,m}(\mathbb{F})$ and that $B_1, \dots, B_s \in \text{Mat}_{n,m}(\mathbb{F})$ have rank 1. By repeated use of the construction above, we can reduce the problem of minimizing

$$\text{rank}(A + x_1B_1 + \dots + x_sB_s)$$

to the Low Rank Matrix Completion Problem.

Example 7. Consider again Example 2. We defined

$$A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

and we reduced the problem of minimizing

$$\text{rank}(A + xB)$$

to the problem of minimizing the rank of

$$J = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} + x_1 \begin{pmatrix} 1 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} + x_2 \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 0 \end{pmatrix} + y_{1,2} \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} + \\ + y_{2,1} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix} + z_{1,2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & -1 \end{pmatrix}.$$

This can be reduced to the Low Rank Matrix Completion Problem:

$$\left(\begin{array}{cccc|cccc} 0 & 1 & 0 & 0 & ? & 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & ? & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & ? \\ \hline 1 & 0 & -1 & 0 & -1 & 0 & 0 & 0 & 0 \\ \hline 0 & 1 & 0 & -1 & 0 & -1 & 0 & 0 & 0 \\ \hline 0 & 0 & ? & 0 & 0 & 0 & -1 & 0 & 0 \\ \hline 0 & 0 & 0 & ? & 0 & 0 & 0 & -1 & 0 \\ \hline 0 & 0 & -1 & -1 & 0 & 0 & 0 & 0 & -1 \end{array} \right).$$

Over the field \mathbb{R} , the smallest possible rank of this matrix is $2 + 5 = 7$, for example when all the missing entries are 0. But over \mathbb{C} , the minimal rank of this matrix is $1 + 5 = 6$. This is the case when all the missing entries are equal to the imaginary number i .

4. REDUCTION FROM 1-RM TO 3-TR

Suppose that $A = (a_{i,j}) \in \text{Mat}_{n,m}(\mathbb{F})$ and that $B_1, \dots, B_s \in \text{Mat}_{n,m}(\mathbb{F})$ have rank 1 and are linearly independent. We can write $B_i = v_i w_i^t$ where $v_i \in \mathbb{F}^n$ and $w_i \in \mathbb{F}^m$ for all i . We will identify $\text{Mat}_{n,m}(\mathbb{F})$ with $\mathbb{F}^n \otimes \mathbb{F}^m$. Then B_i will be identified with $v_i \otimes w_i \in \mathbb{F}^n \otimes \mathbb{F}^m$. We define a third order tensor $T \in \mathbb{F}^n \otimes \mathbb{F}^m \otimes \mathbb{F}^{s+1}$ by

$$(1) \quad T = A \otimes e_{s+1} + \sum_{k=1}^s B_k \otimes e_k = \sum_{i=1}^n \sum_{j=1}^m a_{i,j} (e_i \otimes e_j \otimes e_{s+1}) + \sum_{k=1}^s v_k \otimes w_k \otimes e_k.$$

Let $l = \text{rank } T$. We can write

$$T = \sum_{j=1}^l a_j \otimes b_j \otimes c_j.$$

Lemma 8. *The vectors c_1, \dots, c_l, e_{s+1} span \mathbb{F}^{s+1} .*

Proof. For every i , choose a linear function $h_i : \mathbb{F}^n \otimes \mathbb{F}^m \rightarrow \mathbb{F}$ such that $h_i(B_i) = 1$ and $h_i(B_j) = 0$ if $j \neq i$. Applying $h_i \otimes \text{id} : \mathbb{F}^n \otimes \mathbb{F}^m \otimes \mathbb{F}^{s+1} \rightarrow \mathbb{F}^{s+1}$ to the tensor T gives

$$\sum_{j=1}^l h_i(a_j \otimes b_j) c_j = (h_i \otimes \text{id})(T) = e_i + h_i(A) e_{s+1}.$$

This shows that e_i lies in the span of c_1, \dots, c_l, e_{s+1} for all i . So c_1, \dots, c_l, e_{s+1} span the vector space \mathbb{F}^{s+1} . \square

After rearranging c_1, \dots, c_l , we may assume that $c_1, c_2, \dots, c_s, e_{s+1}$ is a basis of \mathbb{F}^{s+1} . There exists a linear function $f : \mathbb{F}^{s+1} \rightarrow \mathbb{F}$ such that $f(c_1) = \dots = f(c_s) = 0$ and $f(e_{s+1}) = 1$. We apply $\text{id} \otimes \text{id} \otimes f : \mathbb{F}^n \otimes \mathbb{F}^m \otimes \mathbb{F}^{s+1} \rightarrow \mathbb{F}^n \otimes \mathbb{F}^m$ to T :

$$A + \sum_{k=1}^s f(e_k) B_k = (\text{id} \otimes \text{id} \otimes f)(T) = \sum_{j=1}^l f(c_j) a_j \otimes b_j = \sum_{j=s+1}^l f(c_j) a_j \otimes b_j.$$

The resulting matrix clearly has rank at most $l - s$.

Theorem 9. *The smallest possible rank of*

$$A + x_1 B_1 + \cdots + x_s B_s$$

over all $(x_1, \dots, x_s) \in \mathbb{F}^s$ is $l - s$. This minimum is attained when $x_i = f(e_i)$ for $i = 1, 2, \dots, s$.

Proof. Suppose that $x_1, \dots, x_s \in \mathbb{F}$ and $\text{rank}(A + x_1 B_1 + \cdots + x_s B_s) = r$. We have to show that $r \geq l - s$. We can write

$$A + x_1 B_1 + \cdots + x_s B_s = C_1 + \cdots + C_r,$$

where C_1, \dots, C_r are matrices of rank 1. We have

$$\begin{aligned} T &= A \otimes e_{s+1} + \sum_{k=1}^s B_k \otimes e_k = (A + \sum_{k=1}^s x_k B_k) \otimes e_{s+1} - \sum_{k=1}^s x_k B_k \otimes e_{s+1} + \sum_{k=1}^s B_k \otimes e_k = \\ &= \sum_{j=1}^r C_j \otimes e_{s+1} + \sum_{k=1}^s B_k \otimes (e_k - x_k e_{s+1}) \end{aligned}$$

We have written T as a sum of $r + s$ pure tensors. So $l = \text{rank } T \leq r + s$. □

Remark 10. The reduction in this section also easily generalizes to the higher order analog of matrix completion: *tensor completion*. Let $V = \mathbb{F}^{n_1} \otimes \cdots \otimes \mathbb{F}^{n_d}$ and suppose that $A \in V$, and B_1, \dots, B_s are pure tensors that are linearly dependent. Define

$$T = A \otimes e_{s+1} + \sum_{k=1}^s B_k \otimes e_k \in V \otimes \mathbb{F}^{s+1} = \mathbb{F}^{n_1} \otimes \cdots \otimes \mathbb{F}^{n_d} \otimes \mathbb{F}^{s+1}.$$

Suppose that T has rank l . Then we can write

$$T = \sum_{j=1}^l D_j \otimes c_j$$

where $D_1, \dots, D_l \in V$ are pure tensors. The vectors c_1, \dots, c_l, e_{s+1} span \mathbb{F}^{s+1} . After reordering, we may assume that c_1, \dots, c_s, e_{s+1} is a basis. Define a linear map $f: \mathbb{F}^{s+1} \rightarrow \mathbb{F}$ by $f(c_1) = \cdots = f(c_s) = 0$ and $f(e_{s+1}) = 1$. Now the smallest possible rank of the tensor

$$A + \sum_{i=1}^s x_i B_i$$

over all $x_1, \dots, x_s \in \mathbb{F}$ is $l - s$. Equality is attained when we take $x_i = f(e_i)$ for $i = 1, 2, \dots, s$, and we have a decomposition

$$A + \sum_{i=1}^s f(e_i) B_i = \sum_{j=s+1}^l f(c_j) D_j.$$

Example 11. Consider the matrix completion problem over \mathbb{C} for the matrix

$$\begin{pmatrix} 1 & \varepsilon \\ ? & 1 \end{pmatrix}$$

where $\varepsilon \in \mathbb{C}$ is some constant. If $\varepsilon \neq 0$, then the matrix can be completed to a rank 1 matrix:

$$\begin{pmatrix} 1 & \varepsilon \\ \varepsilon^{-1} & 1 \end{pmatrix}.$$

But if $\varepsilon = 0$, then the rank of any completion will be 2. We can view this as a rank minimization problem where

$$A = \begin{pmatrix} 1 & \varepsilon \\ 0 & 1 \end{pmatrix} = e_1 \otimes (e_1 + \varepsilon e_2) + e_2 \otimes e_2 \in \mathbb{C}^2 \otimes \mathbb{C}^2$$

and

$$B = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} = e_2 \otimes e_1 \in \mathbb{C}^2 \otimes \mathbb{C}^2,$$

and we want to minimize the rank of $A + xB$. Let

$$T = e_1 \otimes (e_1 + \varepsilon e_2) \otimes e_2 + e_2 \otimes e_2 \otimes e_2 + e_2 \otimes e_1 \otimes e_1.$$

For $\varepsilon \neq 0$, this tensor has rank 2:

$$T = (e_1 + \varepsilon^{-1}e_2) \otimes (e_1 + \varepsilon e_2) \otimes e_2 + e_2 \otimes e_1 \otimes (e_1 - \varepsilon^{-1}e_2).$$

But for $\varepsilon = 0$ the tensor has rank 3. This is the *border rank phenomenon*: a rank 3 tensor can be the limit of rank 2 tensors. Of course, for matrices (order 2 tensors), this is not possible.

5. REDUCTION FROM 3-TR TO RM

Suppose that $T \in \mathbb{F}^p \otimes \mathbb{F}^q \otimes \mathbb{F}^r$. We would like to find pure tensors v_1, \dots, v_k such that $T = v_1 + \dots + v_k$ and k is minimal. In this section we reduce this problem to rank minimization. First we need some theoretical results.

Lemma 12. *We can write $T = \sum_{i=1}^r S_i \otimes e_i$ where $S_1, \dots, S_r \in \mathbb{F}^p \otimes \mathbb{F}^q \cong \text{Mat}_{p,q}(\mathbb{F})$. If $\text{rank } T = l$ then there exist rank 1 matrices $U_1, \dots, U_l \in \text{Mat}_{p,q}(\mathbb{F})$ such that S_1, \dots, S_r lie in the span of U_1, \dots, U_l .*

Proof. We can write $T = \sum_{i=1}^l U_i \otimes f_i$. Let $\pi_i : \mathbb{F}^r \rightarrow \mathbb{F}$ be the projection on the i -th coordinate, and consider the projection

$$\text{id} \otimes \text{id} \otimes \pi_i : \mathbb{F}^p \otimes \mathbb{F}^q \otimes \mathbb{F}^r \rightarrow \mathbb{F}^p \otimes \mathbb{F}^q.$$

We have $(\text{id} \otimes \text{id} \otimes \pi_i)(T) = S_i = \sum_{j=1}^l \pi_i(f_j)U_j$. □

Lemma 13. *Suppose that $a_1, \dots, a_k \in \mathbb{F}^p$, $b_1, \dots, b_k \in \mathbb{F}^q$ and $\lambda_1, \dots, \lambda_k \in \mathbb{F}$ and let*

$$A = (a_1 \ \cdots \ a_k), \quad B = (b_1 \ \cdots \ b_k), \quad \Lambda = \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_k \end{pmatrix}, \quad C = \begin{pmatrix} S & A & 0 \\ 0 & I_k & -\Lambda \\ B^t & 0 & I_k \end{pmatrix}.$$

Then we have

$$\text{rank } C = 2k + \text{rank} \left(S - \sum_{j=1}^k \lambda_j a_j b_j^t \right).$$

Proof. We perform elementary row operations on the matrix C :

$$\begin{pmatrix} I_p & 0 & 0 \\ 0 & I_k & \Lambda \\ 0 & 0 & I_k \end{pmatrix} C = \begin{pmatrix} I_p & 0 & 0 \\ 0 & I_k & \Lambda \\ 0 & 0 & I_k \end{pmatrix} \begin{pmatrix} S & A & 0 \\ 0 & I_k & -\Lambda \\ B^t & 0 & I_k \end{pmatrix} = \begin{pmatrix} S & A & 0 \\ \Lambda B^t & I_k & 0 \\ B^t & 0 & I_k \end{pmatrix}$$

$$\begin{pmatrix} I_p & -A & 0 \\ 0 & I_k & 0 \\ 0 & 0 & I_k \end{pmatrix} \begin{pmatrix} S & A & 0 \\ \Lambda B^t & I_k & 0 \\ B^t & 0 & I_k \end{pmatrix} = \begin{pmatrix} S - A\Lambda B^t & 0 & 0 \\ \Lambda B^t & I_k & 0 \\ B^t & 0 & I_k \end{pmatrix}$$

So we have

$$\text{rank } C = \text{rank} \begin{pmatrix} S - A\Lambda B^t & 0 & 0 \\ \Lambda B^t & I_k & 0 \\ B^t & 0 & I_k \end{pmatrix} = 2k + \text{rank}(S - A\Lambda B^t) = 2k + \text{rank} \left(S - \sum_{j=1}^k \lambda_j a_j b_j^t \right).$$

□

From now on, assume that $k \geq \text{rank } T$. Let $A \in \text{Mat}_{p,k}(\mathbb{F})$ and $B \in \text{Mat}_{q,k}(\mathbb{F})$ be matrices with indeterminate entries. For every i with $1 \leq i \leq r$ define

$$\Lambda_i = \begin{pmatrix} \lambda_{i,1} & & \\ & \ddots & \\ & & \lambda_{i,k} \end{pmatrix},$$

where $\lambda_{i,1}, \dots, \lambda_{i,k}$ are indeterminates. Let \mathbf{a} be a list of the entries of A , \mathbf{b} a list of the entries of B and λ a list of all $\lambda_{i,j}$. Define

$$C_i(\mathbf{a}, \mathbf{b}, \lambda) = \begin{pmatrix} S_i & A & 0 \\ 0 & I_k & -\Lambda_i \\ B^t & 0 & I_k \end{pmatrix}$$

and

$$U_i(\mathbf{a}, \mathbf{b}, \lambda) = S_i - \sum_{j=1}^k \lambda_{i,j} a_j b_j^t$$

where a_1, \dots, a_k are the columns of A and b_1, \dots, b_k are the columns of B . We have

$$\text{rank } C_i(\mathbf{a}, \mathbf{b}, \lambda) = 2k + \text{rank } U_i(\mathbf{a}, \mathbf{b}, \lambda).$$

We can write

$$\begin{aligned} T &= \sum_{i=1}^r S_i \otimes e_i = \sum_{i=1}^r U_i(\mathbf{a}, \mathbf{b}, \lambda) \otimes e_i + \sum_{i=1}^r \sum_{j=1}^k \lambda_{i,j} a_j b_j^t \otimes e_i = \\ &= \sum_{i=1}^r U_i(\mathbf{a}, \mathbf{b}, \lambda) \otimes e_i + \sum_{j=1}^k a_j b_j^t \otimes \left(\sum_{i=1}^r \lambda_{i,j} e_i \right). \end{aligned}$$

It follows that

$$(2) \quad \text{rank } T \leq \sum_{i=1}^r \text{rank } U_i(\mathbf{a}, \mathbf{b}, \lambda) + \sum_{j=1}^k \text{rank}(a_j b_j^t).$$

Define

$$D_j(\mathbf{a}, \mathbf{b}) = \begin{pmatrix} a_j \\ b_j \end{pmatrix}.$$

Then we have

$$\text{rank } D_j(\mathbf{a}, \mathbf{b}) \geq \text{rank}(a_j b_j^t)$$

with equality if a_i and b_j are both zero or both nonzero. Define

$$E(\mathbf{a}, \mathbf{b}, \lambda) = \begin{pmatrix} C_1(\mathbf{a}, \mathbf{b}, \lambda) & & & & & & \\ & \ddots & & & & & \\ & & C_r(\mathbf{a}, \mathbf{b}, \lambda) & & & & \\ & & & D_1(\mathbf{a}, \mathbf{b}) & & & \\ & & & & \ddots & & \\ & & & & & & D_k(\mathbf{a}, \mathbf{b}) \end{pmatrix}.$$

Theorem 14. *We have*

$$\min_{\mathbf{a}, \mathbf{b}, \lambda} \text{rank } E(\mathbf{a}, \mathbf{b}, \lambda) = 2kr + \text{rank } T.$$

Proof. From (2) follows that

$$\begin{aligned} \text{rank } E(\mathbf{a}, \mathbf{b}, \lambda) &= \sum_{i=1}^r \text{rank}(C_i(\mathbf{a}, \mathbf{b}, \lambda)) + \sum_{j=1}^k \text{rank } D_j(\mathbf{a}, \mathbf{b}) \geq \\ &\geq 2kr + \sum_{i=1}^r \text{rank } U_i(\mathbf{a}, \mathbf{b}, \lambda) + \sum_{j=1}^k \text{rank}(a_j b_j^t) \geq 2kr + \text{rank } T. \end{aligned}$$

We have to show now that we can have equality for some choice of $\mathbf{a}, \mathbf{b}, \lambda$. Suppose that $\text{rank } T = l$. Then we can write $T = \sum_{j=1}^l a_j \otimes b_j \otimes c_j$ with $l \leq k$. Define $a_j = 0$ and $b_j = 0$ for $j = l+1, \dots, k$. If we apply the map

$$\text{id} \otimes \text{id} \otimes \pi_i : \mathbb{F}^p \otimes \mathbb{F}^q \otimes \mathbb{F}^r \rightarrow \mathbb{F}^p \otimes \mathbb{F}^q.$$

to T , we get

$$S_i = \text{id} \otimes \text{id} \otimes \pi_i(T) = \sum_{j=1}^l \pi_i(c_j) a_j b_j^t$$

Define $\lambda_{i,j} = \pi_i(c_j)$. Then we have $U_i(\mathbf{a}, \mathbf{b}, \lambda) = 0$ for $i = 1, 2, \dots, r$. It follows that

$$\begin{aligned} \text{rank } E(\mathbf{a}, \mathbf{b}, \lambda) &= \sum_{i=1}^r \text{rank } C_i(\mathbf{a}, \mathbf{b}, \lambda) + \sum_{j=1}^k \text{rank } D_k(\mathbf{a}, \mathbf{b}) = \\ &= \left(\sum_{i=1}^r 2k \right) + l = rk + l = 2rk + \text{rank } T. \end{aligned}$$

□

Suppose that $\text{rank } E(\mathbf{a}, \mathbf{b}, \lambda)$ is minimal over all choices for $\mathbf{a}, \mathbf{b}, \lambda$. We can write

$$\begin{aligned} T &= \sum_{i=1}^r S_i \otimes e_i = \sum_{i=1}^r U_i(\mathbf{a}, \mathbf{b}, \lambda) \otimes e_i + \sum_{i=1}^r \left(\sum_{j=1}^k \lambda_{i,j} a_j b_j^t \right) \otimes e_i = \\ &= \sum_{i=1}^r U_i(\mathbf{a}, \mathbf{b}, \lambda) \otimes e_i + \sum_{j=1}^k a_j b_j^t \otimes \left(\sum_{i=1}^r \lambda_{i,j} e_i \right). \end{aligned}$$

If $d_i = \text{rank } U_i(\mathbf{a}, \mathbf{b}, \lambda)$ then we can write $U_i(\mathbf{a}, \mathbf{b}, \lambda)$ as sum of d_i rank 1 matrices. Since

$$\sum_{i=1}^r \text{rank } U_i(\mathbf{a}, \mathbf{b}, \lambda) + \sum_{i=1}^k \text{rank}(a_j b_j^t) \leq \sum_{i=1}^r \text{rank } C_i(\mathbf{a}, \mathbf{b}, \lambda) + \sum_{j=1}^k \text{rank } D_j(\mathbf{a}, \mathbf{b}) - 2kr = l,$$

we have written T as a sum of $l = \text{rank } T$ pure tensors.

The matrix $E(\mathbf{a}, \mathbf{b}, \lambda)$ has $(p + 2k)r + (p + q)k$ rows, and $(q + 2k)r + k$ columns. There are $(p + q + k)r$ variables. We can write

$$E(\mathbf{a}, \mathbf{b}, \lambda) = A + \sum_{i,j} a_{i,j} F_{i,j} + \sum_{i,j} b_{i,j} G_{i,j} + \sum_{i,j} \lambda_{i,j} H_{i,j}$$

Here $F_{i,j}$ and $G_{i,j}$ have rank $\leq r + 1$ because the variables $a_{i,j}$ and $b_{i,j}$ appear $r + 1$ times. The matrices $H_{i,j}$ have rank 1 because they only have 1 nonzero entry.

Example 15. Suppose that $T = \sum_{i,j,k} a_{i,j,k} e_i \otimes e_j \otimes e_k$ is a tensor in $\mathbb{F}^2 \otimes \mathbb{F}^2 \otimes \mathbb{F}^2$. Identifying $\mathbb{F}^2 \otimes \mathbb{F}^2 \otimes \mathbb{F}^2$ with $\text{Mat}_{2,2}(\mathbb{F}) \otimes \mathbb{F}^2$ gives us

$$T = \begin{pmatrix} t_{1,1,1} & t_{1,2,1} \\ t_{2,1,1} & t_{2,2,1} \end{pmatrix} \otimes e_1 + \begin{pmatrix} t_{1,1,2} & t_{1,2,2} \\ t_{2,1,2} & t_{2,2,2} \end{pmatrix} \otimes e_2 = S_1 \otimes e_1 + S_2 \otimes e_2.$$

The largest possible rank of a tensor T in $\mathbb{F}^2 \otimes \mathbb{F}^2 \otimes \mathbb{F}^2$ is 3. So let us take $k = 3$. Define

$$A = \begin{pmatrix} a_{1,1} & a_{1,2} & a_{1,3} \\ a_{2,1} & a_{2,2} & a_{2,3} \end{pmatrix}, B = \begin{pmatrix} b_{1,1} & b_{1,2} & b_{1,3} \\ b_{1,2} & b_{2,2} & b_{2,3} \end{pmatrix}.$$

For $i = 1, 2$, define

$$\Lambda_i = \begin{pmatrix} \lambda_{i,1} & 0 & 0 \\ 0 & \lambda_{i,2} & 0 \\ 0 & 0 & \lambda_{i,3} \end{pmatrix},$$

$$C_i = \begin{pmatrix} S_i & A & 0 \\ 0 & I_3 & \Lambda_i \\ B^t & 0 & I_3 \end{pmatrix} = \left(\begin{array}{cc|ccc|ccc} t_{1,1,i} & t_{1,2,i} & a_{1,1} & a_{1,2} & a_{1,3} & 0 & 0 & 0 \\ t_{2,1,i} & t_{2,2,i} & a_{2,1} & a_{2,2} & a_{2,3} & 0 & 0 & 0 \\ \hline 0 & 0 & 1 & 0 & 0 & \lambda_{i,1} & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & \lambda_{i,2} & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & \lambda_{i,3} \\ \hline b_{1,1} & b_{2,1} & 0 & 0 & 0 & 1 & 0 & 0 \\ b_{1,2} & b_{2,2} & 0 & 0 & 0 & 0 & 1 & 0 \\ b_{1,3} & b_{2,3} & 0 & 0 & 0 & 0 & 0 & 1 \end{array} \right)$$

$$D_j = \begin{pmatrix} a_{1,j} \\ a_{2,j} \\ b_{1,j} \\ b_{2,j} \end{pmatrix}$$

So finally, we have

$$E(\mathbf{a}, \mathbf{b}, \lambda) = \begin{pmatrix} C_1 & 0 & 0 & 0 & 0 \\ 0 & C_2 & 0 & 0 & 0 \\ 0 & 0 & D_1 & 0 & 0 \\ 0 & 0 & 0 & D_2 & 0 \\ 0 & 0 & 0 & 0 & D_3 \end{pmatrix} =$$

$$\left(\begin{array}{cccccc|cccc|ccc|ccc} t_{1,1,1} & t_{1,2,1} & a_{1,1} & a_{1,2} & a_{1,3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ t_{2,1,1} & t_{2,2,1} & a_{2,1} & a_{2,2} & a_{2,3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & \lambda_{1,1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & \lambda_{1,2} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & \lambda_{1,3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ b_{1,1} & b_{2,1} & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ b_{1,2} & b_{2,2} & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ b_{1,3} & b_{2,3} & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & t_{1,1,2} & t_{1,2,2} & a_{1,1} & a_{1,2} & a_{1,3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & t_{2,1,2} & t_{2,2,2} & a_{2,1} & a_{2,2} & a_{2,3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & \lambda_{2,1} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & \lambda_{2,2} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & \lambda_{2,3} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & b_{1,1} & b_{2,1} & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & b_{1,2} & b_{2,2} & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & b_{1,3} & b_{2,3} & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & a_{1,1} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & a_{2,1} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & b_{1,1} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & b_{2,1} & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & a_{1,2} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & a_{2,2} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & b_{1,2} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & b_{2,2} & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & a_{1,3} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & a_{2,3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & b_{1,3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & b_{2,3} & 0 & 0 \end{array} \right).$$

We have

$$\min_{\mathbf{a}, \mathbf{b}, \lambda} E(\mathbf{a}, \mathbf{b}, \lambda) = 12 + \text{rank } T.$$

6. REDUCTION FROM TR TO RM

For a matrix $A \in \text{Mat}_{m,n}(\mathbb{F})$, we define its *padding* as the partially filled matrix

$$\text{pad}(A) = \begin{pmatrix} A & ? \\ ? & 1 \end{pmatrix}$$

Lemma 16. *We have $\text{mrank pad}(A) = \max\{1, \text{rank } A\}$.*

Proof. It is clear that $\text{rank pad}(A) \geq \max\{1, \text{rank } A\}$, and we have equality if $A = 0$. Suppose that A is nonzero and let v be a nonzero column v of A . Choose a vector w so that $(w^t \ 1)$ is a multiple of a row of the matrix $(A \ v)$. We have

$$\text{rank} \begin{pmatrix} A & v \\ w^t & 1 \end{pmatrix} = \text{rank} (A \ v) = \text{rank}(A) = \max\{1, \text{rank}(A)\}.$$

□

Suppose that $T \in \mathbb{F}^{n_1} \otimes \cdots \otimes \mathbb{F}^{n_d}$. Inductively, we will define partially filled matrices $A(T)$ and $B(T)$. If $d = 2$ then T can be viewed as a $n_1 \times n_2$ matrix and we define $A(T) = \text{pad}(T)$ and $B(T) = T$.

We can write

$$T = \sum_{i=1}^{n_d} S_i \otimes e_i.$$

where $S_i \in \mathbb{F}^{n_1} \otimes \cdots \otimes \mathbb{F}^{n_{d-1}}$. We choose a linear isomorphism

$$\phi : \mathbb{F}^{n_1} \otimes \cdots \otimes \mathbb{F}^{n_{d-1}} \rightarrow \mathbb{F}^{n_1 n_2 \cdots n_{d-1}},$$

where $\phi(S)$ will be viewed as a column vector. Define

$$N(T) = (\phi(S_1) \ \cdots \ \phi(S_{n_d}))$$

and

$$A(T) = \begin{pmatrix} \text{pad}(N(T)) & & & \\ & A(S_1) & & \\ & & \ddots & \\ & & & A(S_{n_d}) \end{pmatrix}$$

Lemma 17. *We have the inequality*

$$\text{mrank } A(T) \geq \max\{1, \text{rank } T\} + K(n_1, \dots, n_d),$$

where

$$K(n_1, \dots, n_d) = n_3 n_4 \cdots n_d + n_4 n_5 \cdots n_d + \cdots + n_d.$$

If T has rank 0 or 1 then we have equality.

Proof. We prove the inequality by induction on d . If $d = 2$ then $A(T) = \text{pad}(T)$ and $K(n_1, n_2) = 0$. So we have

$$\text{mrank } A(T) = \text{mrank pad}(T) = \max\{\text{rank } T, 1\} = \max\{\text{rank } T, 1\} + K(n_1, n_2).$$

Suppose that $d \geq 3$. We prove the inequality by induction on n_d . If $n_d = 1$ then we have $T = S_1 \otimes e_1$, $\text{rank pad}(N(T)) = 1$, and

$$K(n_1, \dots, n_d) = K(n_1, \dots, n_{d-1}, 1) = K(n_1, \dots, n_{d-1}) + 1.$$

It follows that

$$\begin{aligned} \text{mrank } A(T) &= \text{mrank pad}(N(T)) + \text{mrank } A(S_1) \geq \\ &\geq 1 + \max\{1, \text{rank } S_1\} + K(n_1, \dots, n_{d-1}) = \max\{1, \text{rank } T\} + K(n_1, \dots, n_d). \end{aligned}$$

If T has rank ≤ 1 , then $\text{rank } S_1 \leq 1$ and we have equality.

Suppose that $n_d > 1$. Let $T' = \sum_{i=1}^{n_d-1} S_i \otimes e_i$.

First, assume that S_1, \dots, S_{n_d} are linearly independent. Then T and T' are nonzero. We have

$$\begin{aligned} \text{mrank } A(T) &= \text{mrank pad}(N(T)) + \sum_{i=1}^{n_d} \text{mrank } A(S_i) = \text{rank } N(T) + \sum_{i=1}^{n_d} \text{mrank } A(S_i) = \\ &= \text{rank } N(T') + 1 + \sum_{i=1}^{n_d} \text{mrank } A(S_i) = \text{mrank } A(T') + 1 + \text{mrank } A(S_{n_d}) \geq \\ &\geq \text{rank } T' + K(n_1, \dots, n_{d-1}, n_d - 1) + 1 + \text{rank } S_{n_d} + K(n_1, \dots, n_{d-1}) = \\ &= \text{rank } T' + \text{rank } S_{n_d} + K(n_1, \dots, n_d) \geq \text{rank } T + K(n_1, \dots, n_d). \end{aligned}$$

If S_1, \dots, S_{n_d} are linearly dependent, we may assume without loss of generality that S_{n_d} lies in the span of S_1, \dots, S_{n_d-1} . Then we have $\text{rank } T = \text{rank } T'$, and

$$\text{mrank pad}(N(T)) = \text{mrank pad}(N(T')).$$

It follows that

$$\begin{aligned} \text{mrank } A(T) &= \text{mrank pad}(N(T)) + \sum_{i=1}^{n_d} \text{mrank } A(S_i) = \\ &= \text{mrank pad}(N(T')) + \sum_{i=1}^{n_d} \text{mrank } A(S_i) = \text{mrank } A(T') + \text{mrank } A(S_{n_d}) \geq \\ &\geq \max\{1, \text{rank } T'\} + K(n_1, \dots, n_{d-1}, n_d - 1) + \max\{1, \text{rank } S_{n_d}\} + K(n_1, \dots, n_{d-1}) \geq \\ &\geq \max\{1, \text{rank } T\} + K(n_1, \dots, n_{d-1}, n_d - 1) + K(n_1, \dots, n_{d-1}) + 1 = \\ &= \max\{1, \text{rank } T\} + K(n_1, \dots, n_d). \end{aligned}$$

If $\text{rank } T \leq 1$ then $\text{rank } S_i \leq 1$ for all i , and we have

$$\begin{aligned} \text{mrank } A(T) &= \text{mrank pad}(N(T)) + \sum_{i=1}^{n_d} \text{rank } A(S_i) = \\ &= 1 + n_d(1 + K(n_1, \dots, n_{d-1})) = 1 + K(n_1, \dots, n_d) = \max\{\text{rank } T, 1\} + K(n_1, \dots, n_d). \end{aligned}$$

□

Let us define

$$B(T) = \begin{pmatrix} N(T) & & & \\ & A(S_1) & & \\ & & \ddots & \\ & & & A(S_{n_d}) \end{pmatrix}.$$

Lemma 18. *We have*

$$\text{mrank } B(T) \geq \text{rank } T + K(n_1, \dots, n_d)$$

with equality if $\text{rank}(T) \leq 1$.

Proof. If T is nonzero, then we have $\text{rank } N(T) = \text{mrank pad}(N(T))$, so

$$\text{mrank } B(T) = \text{mrank } A(T) \geq \text{rank } T + K(n_1, \dots, n_d).$$

with equality if $\text{rank } T = 1$.

If $T = 0$, then $\text{rank } N(T) = 0$ and $\text{rank pad}(N(T)) = 1$. It follows that

$$\text{mrnk } B(T) = \text{mrnk } A(T) - 1 \geq K(n_1, \dots, n_d) = \text{rank } T + K(n_1, \dots, n_d).$$

□

For tensors U_1, \dots, U_k we define

$$C(U_1, \dots, U_k) = \begin{pmatrix} B(U_1 - U_2) & & & \\ & \ddots & & \\ & & B(U_{k-2} - U_{k-1}) & \\ & & & B(U_{k-1} - U_k) \end{pmatrix}$$

Lemma 19. *If $\text{rank } T \leq k$, then we have*

$$\text{rank } T + kK(n_1, \dots, n_d) = \min_{U_1, \dots, U_{k-1}} \text{mrnk } C(T, U_1, \dots, U_{k-1}).$$

Proof. For all U_1, \dots, U_{k-1} we have

$$\begin{aligned} \text{rank } C(T, U_1, \dots, U_{k-1}) &= \text{rank } B(T - U_1) + \sum_{i=2}^k \text{rank } B(U_i - U_{i-1}) \geq \\ &\geq \text{rank}(T - U_1) + \sum_{i=1}^k \text{rank}(U_i - U_{i-1}) + kK(n_1, \dots, n_d) \geq \text{rank } T + kK(n_1, \dots, n_d). \end{aligned}$$

We can choose U_1, \dots, U_{k-1} such that $T - U_1, U_1 - U_2, \dots, U_{k-1} - U_k$ are pure tensors. In that case we have equality. □

Acknowledgment. The author would like to thank Bernard Mourrain for pointing out some references.

REFERENCES

- [1] Z. Bar-Yossef, Y. Birk, T. S. Jayram and T. Kol, *Index coding with side information*, IEEE Trans. on Inf. Th. **57** (2011), no. 3, 1479–1494.
- [2] A. Bernardi, J. Brachat, P. Comon and B. Mourrain, *General tensor decomposition, moment matrices and applications*, J. Symbolic Computation **52** (2013), 51–71.
- [3] J. Brachat, P. Comon, B. Mourrain, E. P. Tsigaridas, *Symmetric tensor decomposition*, Linear Algebra and its Applications **433** (2010), no. 1, 1851–1872.
- [4] J.-F. Cai, E. Candès and Z. Shen, *A singular value thresholding algorithm for matrix completion*, SIAM J. Optim. **20** (2010), no. 4, 1956–1982.
- [5] J. D. Carroll and J. Chang, *Analysis of individual differences in multidimensional scaling via an n -way generalization of “Eckart-Young” decomposition*, Psychometrika **35** (1970), 283–319.
- [6] E.J. Candès and B. Recht, *Exact matrix completion via convex optimization*, Foundations of Computational Mathematics **9**, 2009, 717–772.
- [7] E. J. Candès and T. Tao, *The power of convex relaxation: Near-optimal matrix completion*, IEEE Trans. Inform. Theory **56** (5), 2053–2080.
- [8] H. Derksen, *Matrix completion and tensor rank*, preprint, arXiv:1302.2639.
- [9] V. De Silva and L.-H. Lim, *Tensor rank and the ill-posedness of the best low-rank approximation problem*, SIAM J. Matrix Analysis Appl. **30** (2008), no. 3, 1084–1127.
- [10] M. Fazel, *Matrix Rank Minimization with Applications*, Ph.D. thesis, Stanford University, (2002).
- [11] M. Fazel, H. Hindi and S. Boyd, *A rank minimization heuristic with application to minimum order system approximation*, in: Proceedings of the American Control Conference **6** (2001), 4734–4739.

- [12] M. Fazel, H. Hindi and S. Boyd, *Rank minimization and applications in system theory*, in: American Control Conference, 2004, 3273–3278.
- [13] L. E. Ghaoui, P. Gahinet, *Rank minimization under LMI constraints: A framework for output feedback problems*, in: Proceedings of the European Control Conference, 1993.
- [14] R. A. Harshman, *Foundations of the parafac procedure: models and conditions for an “explanatory” multimodal factor analysis*, UCLA working papers in Phonetics **16** (1970), 1–84.
- [15] J. Håstad, *Tensor rank is NP-complete*, J. Algorithms **11** (1990), no. 4, 644–654.
- [16] J. Håstad, *Tensor rank is NP-complete*, Automata, languages and programming (Stresa, 1989), Lecture Notes in Comput. Sci. **372**, Springer, Berlin, 1989, 451–460.
- [17] F. L. Hitchcock, *Multiple invariants and generalized rank of a p -way matrix or tensor*, J. Math. and Phys. **7** (1927), no. 1, 40–79.
- [18] M. Laurent, *A connection between positive semidefinite and Euclidean distance matrix completion problems*, Linear Algebra Appl. **273** (1998), 9–22.
- [19] C. Hillar and L.-H. Lim, *Most tensor problems are NP-hard*, Journal of the ACM (2013), to appear.
- [20] C. R. Johnson, *Matrix completion problems: a survey*, in: Matrix Theory and Applications, AMS, Providence, RI, 1990, 171–198.
- [21] R. H. Keshavan, A. Montanari and S. Oh, *Matrix completion from a few entries*, IEEE Trans. on Information Theory **56** (2010), 2980–2998.
- [22] T. G. Kolda, B. W. Bader, *Tensor Decompositions and Applications*, SIAM Rev. **51** (3), 455–500.
- [23] N. Linial, E. London and Y. Rabinovich, *The geometry of graphs and some of its algorithmic applications*, Combinatorica **15** (1995), 215–245.
- [24] Z. Liu, L. Vandenberghe, *Interior-point method for nuclear norm approximation with application to system identification*, SIAM J. Matrix Anal. Appl. **31** (2009), no. 3, 1235–1256.
- [25] S. Ma, D. Goldfarb, L. Chen, *Fixed point and Bregman iterative methods for matrix rank minimization*, Math. Program. **128** (2011), 321–353.
- [26] R. Peeters, *Orthogonal representations over finite fields and the chromatic number of graphs*, Combinatorica **16** (1996), 417–431.
- [27] B. Recht, M. Fazel and P. Parillo, *Guaranteed minimum rank solutions of matrix equations via nuclear norm minimization*, SIAM Review **52** (2010), no. 3, 471–501.
- [28] V. Y. F. Tan, L. Balzano, S. C. Draper, *Rank Minimization over finite fields: fundamental limits and coding-theoretic interpretations*, IEEE transactions on Inf. Th. **58** (2012), no. 4, 2018–2039.
- [29] G. Tomasi and R. Bro, *A comparison of algorithms for fitting the parafac model*, Computational Statistics and Data Analysis **50**, 7, 1700–1734.