

Optimal investment with time-varying stochastic endowments

Christoph Belak^a, An Chen^b, Carla Mereu^c, and Robert Stelzer^d

^aTU Berlin, Institute of Mathematics, Straße des 17. Juni 136, 10623 Berlin, Germany.
belak@math.tu-berlin.de

^bUlm University, Institute of Insurance Science, Helmholtzstrasse 20, 89081 Ulm, Germany.
an.chen@uni-ulm.de

^cd-fine GmbH, An der Hauptwache 7, 60313 Frankfurt, Germany. ccarlamereu@gmail.com

^dUlm University, Institute of Mathematical Finance, Helmholtzstrasse 18, 89081 Ulm, Germany.
robert.stelzer@uni-ulm.de

October 19, 2021

This paper considers a utility maximization and optimal asset allocation problem in the presence of a stochastic endowment that cannot be fully hedged through trading in the financial market. After studying continuity properties of the value function for general utility functions, we rely on the dynamic programming approach to solve the optimization problem for power utility investors including the empirically relevant and mathematically challenging case of relative risk aversion larger than one. For this, we argue that the value function is the unique viscosity solution of the Hamilton-Jacobi-Bellman (HJB) equation. The homogeneity of the value function is then used to reduce the HJB equation by one dimension, which allows us to prove that the value function is even a classical solution thereof. Using this, an optimal strategy is derived and its asymptotic behavior in the large wealth regime is discussed.

Keywords. Utility maximization; Hamilton-Jacobi-Bellman equation; stochastic endowment; viscosity solution.

In the present paper, we analyze the utility maximization or optimal investment problem of an economic agent who receives stochastic endowments over a finite time period. We deal with an incomplete market setting as the endowment risk is not perfectly correlated with the traded assets in the market. A specific example to motivate our model would be an economic agent who receives random salaries during her working life and invests a fixed proportion of the salaries in financial assets, optimizing the utility of her wealth at the retirement age. Another example could be a pension fund which receives random contributions from the members of the pension fund and invests to generate cash flows. In fact, the original idea of the paper arises from the study of defined contribution (DC) pension plans. This kind of pension scheme has become very popular over the last years and is substituting defined benefit pension plans (see e.g. [4]).

The problem of maximizing the expected utility of an economic agent by investment and/or consumption dates back to [26] and [27] and is further studied e.g. in [8, 12, 14, 23], just to quote

a few. Such problems have originally been solved using the dynamic programming approach which requires the assumption of Markovianity on the state process and leads to a Hamilton-Jacobi-Bellman (HJB) equation. In the literature, this approach is called primal approach. In the 1980s, researchers developed an alternative approach, the so called dual approach where the assumption of Markovian asset prices can be relaxed to solve the optimal investment problem. In a complete market setting the dual method has been studied e.g. in [6, 32] and in an incomplete market setting e.g. by [18] and [19].

In the present paper we consider the optimal asset allocation problem on a finite time horizon for an investor receiving an exogenous stochastic endowment modeled as a time-inhomogeneous geometric Brownian motion. We follow the primal HJB approach and argue that, in the case of power utility and under suitable regularity assumptions on the market parameters, the value function is a classical solution of the HJB equation. To obtain these results, we use a viscosity solution approach combined with classical existence results on regular solutions for nonlinear parabolic partial differential equations. More precisely, we first argue that the value function is the unique continuous viscosity solution of the HJB equation. Using the fact that the value function is homogeneous, we reduce the dimension of the HJB equation by one and obtain a fully nonlinear second-order parabolic equation which is uniformly elliptic in the spatial variable. From this, we are able to argue that the value function is even a classical solution of the HJB equation.

Proceeding from there, we then construct an optimal trading strategy. Since we restrict to strategies taking values in a compact-convex set, we shall see that it is a priori not clear if the stochastic differential equation (SDE) for the wealth process under the candidate optimal strategy (given in feedback form through the maximizer of the Hamiltonian in the HJB equation) admits a strong solution due to a lack of global Lipschitz properties of the drift and diffusion coefficients of the SDE. We are therefore forced to follow a non-classical approach to the construction of an optimal strategy: We first argue that an optimal strategy exists and then show that it can be represented in feedback form. In particular, this implies that the optimal strategy is unique. Given the optimal strategy, we finally analyze its asymptotic properties for large wealth-to-endowment ratios. We find that, asymptotically, the optimal strategy converges to the famous Merton ratio, which is the optimal strategy in the absence of random endowments.

The idea of incorporating a stochastic endowment in the classical utility maximization problem is indeed not new. Both dual and primal approaches (sometimes combined with Backward Stochastic Differential Equation (BSDE) techniques) are adopted to solve this optimization problem. Most of the papers in this field restrict themselves however to the analytically more tractable case of exponential utility functions (among several others [10] and [21]). Some authors deal also with the problem of maximizing expected power utility of terminal wealth in the presence of exogenous endowments. In [9], the problem is solved via duality for a broad class of utility functions under the assumption that the random endowment is bounded. The existence and uniqueness of an optimal control are proven, but an explicit representation of the optimal strategy is subordinated to the decomposition of the elements of $(L^\infty)^*$ into a regular and a singular part, which is hard to characterize explicitly. The authors of [22] overcome the problem and relax the hypothesis of boundedness by introducing the number of random endowments as a new control variable. [28]

aims to examine under which conditions on the market model and on the utility stochastic field the maximizer of the optimal consumption problem exists. In addition, [28] looks into the properties of the value function and the corresponding dual problem. [29] studies existence and uniqueness of optimal investment and consumption strategies in a general incomplete semimartingale setting and provides a dual characterization in terms of an optional strong supermartingale deflator and a decreasing process. [37] focuses on examining the stability of the utility maximization problem with random endowment, i.e. when there is model and/or preference misspecification for the optimal investment problem. [20] extends the approach of [21] to the case of power utility functions, based on the martingale optimality principle combined with BSDE methods. They reduce the problem to the solution of a fully-coupled forward-backward stochastic differential equation, which is still not easy to solve.

Most closely related to our paper are [3] and [12]. The setting of [3] is similar to ours and considers a utility maximization problem with stochastic income modelled by geometric Brownian motion. However, they find a near-optimal consumption and investment strategy which leads to a small wealth-equivalent loss compared to the unknown optimal strategy. In [12], the expected HARA utility from consumption is optimized over an infinite-time horizon and it yields an elliptic HJB equation, not depending on time. We adopt and extend their methods, developed in the elliptic case (infinite-time horizon), to the parabolic case (finite-time horizon). We further extend their model to a slightly broader class of random endowments, i.e., we allow for time-varying coefficients. Most importantly, while [12] deals exclusively with the case of relative risk aversion less than one, we additionally extend the model to the empirically more relevant case of power utilities with relative risk aversion larger than one. Especially the extension to the latter class of utility functions poses significant mathematical challenges in the viscosity characterization of the value function. In fact, we are able to establish a strong comparison principle for the (unreduced) HJB equation which covers the large relative risk aversion case as well, which is often a significant challenge in utility maximization problems; see, e.g., [33, 2].

The remainder of the paper is organized as follows. Section 1 describes the model setup and particularly the endowment process. In Section 2, we introduce the optimization problem and study some of its basic properties such as well-posedness, continuity, concavity, and homogeneity. In Section 3 we restrict ourselves to power utility and study properties of the associated HJB equation. Here, we show that the value is the unique viscosity solution of the HJB equation and prove that it is even a classical solution if the market parameters are sufficiently regular. In Section 4, we construct an optimal strategy in the regular case and analyze its asymptotic behavior as the wealth-to-endowment ratio approaches infinity. Finally, Section 5 concludes the paper.

1 The model

On a fixed filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0, T]}, \mathbb{P})$ satisfying the usual hypotheses, consider a financial market consisting of a riskless and a risky asset. From now on let $T > 0$ be a fixed finite time point. Let S^0 and S^1 denote respectively the savings account and the risky asset. We

assume that the two assets follow a Black-Scholes model:

$$dS_t^0 = rS_t^0 dt, \quad dS_t^1 = \mu S_t^1 dt + \sigma S_t^1 dW_t^1,$$

where $\mu, r \in \mathbb{R}$, $\sigma > 0$ and W^1 is a Brownian motion on our filtered probability space.

The endowment process c is assumed to have stochastic dynamics driven by another Brownian motion W^C defined on the same probability space, which is assumed to be correlated with W^1 with a correlation coefficient¹ $\rho \in (-1, 1)$. In other words, we assume that there is another Brownian motion W^2 independent of W^1 such that $W^C = \rho W^1 + \sqrt{1 - \rho^2} W^2$, which leads to the random endowment process c given by

$$dc_t = \mu_C(t)c_t dt + \sigma_C(t)c_t dW_t^C, \quad t \in [0, T], \quad c_0 = y,$$

where $y > 0$ and $\mu_C : [0, T] \rightarrow \mathbb{R}$, $\sigma_C : [0, T] \rightarrow (0, \infty)$ are deterministic continuous functions.

Remark 1.1. Time-homogeneous geometric Brownian motions are sometimes chosen in the literature to model a stochastic income (see [34]). Thinking of the example of a DC pension scheme, we can interpret the process c as a diffusion income, or rather a proportion of it possibly changing over time, which is paid continuously into the pension fund. From an analytical point of view, the fundamental feature of this family of random endowments is that, in the power utility case, the value function of the optimization problem turns out to be homogeneous in the spatial variables. This is the crucial property to achieve a reduction in the dimension of the problem.

We assume that an agent with an initial wealth $x > 0$ invests at any time t a proportion π_t of the wealth in the stock S^1 and $1 - \pi_t$ in the risk-free asset S^0 with interest rate r . In addition, the random income is paid continuously to the account at rate c_t . The wealth process corresponding to the strategy π , denoted by A^π , is assumed to have the following dynamics:

$$dA_t^\pi = \frac{\pi_t A_t^\pi}{S_t^1} dS_t^1 + \frac{(1 - \pi_t) A_t^\pi}{S_t^0} dS_t^0 + c_t dt.$$

Defining $\theta := (\mu - r)/\sigma$, this can be written as

$$(1.1) \quad dA_t^\pi = [(r + \sigma\theta\pi_t)A_t^\pi + c_t] dt + \sigma\pi_t A_t^\pi dW_t^1, \quad t \in [0, T], \quad A_0^\pi = x.$$

This definition reflects the fact that the only additional cash injections are due to the continuous payments at rate c_t .

Note that in our set-up a strategy π encodes the fraction of the overall wealth invested in the risky asset. We assume that π is a progressively measurable process taking values in a compact-convex set $A := [\underline{\pi}, \bar{\pi}]$ for real numbers $\underline{\pi} < \bar{\pi}$. This includes the case of short-selling of the risky asset and borrowing of cash being restricted. We write \mathcal{A} for the set of all such trading strategies.

¹We exclude the perfect correlation cases $\rho = 1$ and $\rho = -1$. In these extreme cases, both the risky asset and the random income are fully driven by W^1 . The optimization problem becomes simpler and differs from what we will present in the remaining text. For instance, in a retirement context, [13] study the optimal consumption and retirement problem with stochastic labor income, where the labor income is perfectly correlated with the financial market risk.

To compare and discuss optimal trading in the presence and the absence of random endowments, we subsequently assume $\pi_M \in A$, where π_M denotes the Merton fraction

$$\pi_M := \frac{\theta}{(1-\gamma)\sigma}.$$

It is a classical result that π_M is the optimal strategy in the absence of random endowments, i.e. in case of $c \equiv 0$, and power utility with $1 - \gamma$ being the relative risk aversion. Let us highlight that our compactness assumption does not imply that the money invested in the risky asset or the cash position is bounded. Instead it means that there is a relative bound on the extent to which short-selling of the risky asset and taking credit on the bank account is allowed. As an example, choosing $A = [0, 1]$ amounts to assuming that short-selling of the risky asset and borrowing of cash are prohibited. Mathematically, the compactness assumption makes the study of the value function of the optimal investment problem significantly easier than in the unrestricted case. In fact, we shall make ample use of this assumption in our results in Section 2. In contrast, this trading constraint leads to significant challenges in the construction of an optimal strategy. In fact, even though we are able to show that the value function is a classical solution of the HJB equation and obtain a candidate optimal strategy in feedback form, it is unclear if the wealth process associated with the candidate optimal strategy exists; see the discussion in the beginning of Section 4 for details. We overcome this issue by first establishing an abstract existence result for optimal strategies and subsequently verifying that any optimal strategy coincides with our candidate optimal control.

Remark 1.2. Evidently, as A is compact, any $\pi \in \mathcal{A}$ is bounded. In particular, this ensures the existence and uniqueness of a solution for Equation (1.1).

Remark 1.3. In the definition of admissibility, one usually has to ensure that the wealth process never becomes negative. Under our assumptions, the wealth process stays positive without any extra requirement on the admissible strategies. This can be seen from the fact that the explicit solution of (1.1) is given by

$$A_t^\pi = \left(x + \int_0^t \frac{c_s}{Z_s^\pi} ds \right) Z_t^\pi, \quad t \in [0, T],$$

where Z^π is a stochastic exponential factor given as

$$Z_t^\pi := \exp \left\{ \int_0^t \left[\pi_s \sigma \theta + r - \frac{1}{2} (\pi_s \sigma)^2 \right] ds + \int_0^t \sigma \pi_s dW_s^1 \right\}, \quad t \in [0, T].$$

2 The optimization problem: statement and properties

Recall that the Brownian motion W^C represents the uncertainty in the income, which is supposed not to be traded in the market. This makes the market incomplete and we are hence facing the problem of maximizing expected utility of an individual in an incomplete market. More precisely,

we are looking for an optimal investment strategy $\pi^* \in \mathcal{A}$ such that

$$\mathbb{E}[U(A_T^{\pi^*})] = \sup_{\pi \in \mathcal{A}} \mathbb{E}[U(A_T^\pi)],$$

where $U : (0, \infty) \rightarrow \mathbb{R}$ is a utility function which is twice continuously differentiable, increasing, concave, and satisfies the Inada conditions $\lim_{x \uparrow +\infty} U'(x) = 0$ and $\lim_{x \downarrow 0} U'(x) = \infty$.

Later on, we will restrict to the special case of U being a constant relative risk aversion (CRRA) power utility function, i.e. for a risk aversion parameter $\gamma < 1$, $\gamma \neq 0$, we shall assume that $U = U_\gamma$ with

$$U_\gamma(x) = \frac{1}{\gamma} x^\gamma, \quad x > 0.$$

The power utility function is abundantly used in both theoretical and empirical research because of its analytical tractability. Most importantly, the use of power utility is also well-motivated economically, since the long-run behavior of the economy suggests that the long run risk aversion does not strongly depend on wealth, see [5].

The agent is assumed to maximize expected utility of terminal wealth at the final time T , and thus the value function of the utility maximization problem is given by²

$$(2.1) \quad \mathcal{V} : [0, T] \times \mathcal{O} \rightarrow \mathbb{R} \cup \{-\infty\}, \quad (t, x, y) \mapsto \mathcal{V}(t, x, y) := \sup_{\pi \in \mathcal{A}} \mathbb{E}[U(A_T^{\pi, t, x, y})],$$

where $\mathcal{O} := (0, \infty) \times (0, \infty)$. In the definition of the value function, the state process is taken to be the pair $(A^{\pi, t, x, y}, c^{t, y})$ and the notation $A^{\pi, t, x, y}$ and $c^{t, y}$ is used for the wealth and endowment process started at time t in x and y , respectively.

2.1 A priori estimates, local boundedness, and continuity

We first provide simple sufficient conditions under which the value function is finite and continuous. A key ingredient is the following collection of a priori estimates on the state processes.

Lemma 2.1 (A priori estimates). *Let $\alpha, \beta, \eta > 0$. Then there exist $C_\alpha, C_\beta, C_\eta > 0$ such that*

$$(2.2) \quad \mathbb{E} \left[\sup_{s \in [t, T]} |c_s^{t, y}|^\alpha \right] \leq C_\alpha |y|^\alpha, \quad (t, y) \in [0, T] \times (0, \infty),$$

$$(2.3) \quad \sup_{\pi \in \mathcal{A}} \mathbb{E} \left[\sup_{s \in [t, T]} |A_s^{\pi, t, x, y}|^{-\beta} \right] \leq C_\beta |x|^{-\beta}, \quad (t, x, y) \in [0, T] \times \mathcal{O},$$

$$(2.4) \quad \sup_{\pi \in \mathcal{A}} \mathbb{E} \left[\sup_{s \in [t, T]} |A_s^{\pi, t, x, y}|^\eta \right] \leq C_\eta |(x, y)|^\eta, \quad (t, x, y) \in [0, T] \times \mathcal{O}.$$

Moreover, there exists a constant $C > 0$ such that

$$(2.5) \quad \sup_{\pi \in \mathcal{A}} \mathbb{E} \left[|A_T^{\pi, t, x, y} - A_T^{\pi, t', x', y'}|^2 \right] \leq C^2 \left(|(x, y) - (x', y')|^2 + |(x, y)|^2 |t - t'| \right)$$

²A priori, it is of course not evident that \mathcal{V} is finite or even well-defined. We shall, however, see in Proposition 2.2 below that the positive part of $U(A_T^{\pi, t, x, y})$ is integrable with a bound not depending on the trading strategy $\pi \in \mathcal{A}$, in which case \mathcal{V} is well-defined as an $\mathbb{R} \cup \{-\infty\}$ -valued function.

for all $(t, x, y), (t', x', y') \in [0, T] \times \mathcal{O}$.

Proof. The estimate (2.5) follows from a direct application of Proposition 3.22 in [30] (applying Eq. (3.78) with $p = 2$, $q = \infty$, and $\lambda > 1$ arbitrarily in the notation of [30]). To obtain (2.2), we first observe that an application of Itô's lemma shows that the process $\tilde{c} := |c^{t,y}|^{\alpha/2}$ satisfies the (linear) Lipschitz SDE

$$d\tilde{c}_s = \tilde{c}_s \frac{\alpha}{2} \left[\left(\mu_C(s) + \frac{1}{4}(\alpha - 2)\sigma_C(s)^2 \right) ds + \sigma_C(s) dW_s^C \right], \quad s \in [t, T], \quad \tilde{c}_t = |y|^{\alpha/2}.$$

Classical a priori estimates for solutions of Lipschitz SDEs such as Corollary 2.10 in [25] applied to \tilde{c} hence imply the existence of a constant $C_\alpha > 0$ such that

$$\mathbb{E} \left[\sup_{s \in [t, T]} |c_s^{t,y}|^\alpha \right] = \mathbb{E} \left[\sup_{s \in [t, T]} |\tilde{c}_s|^2 \right] \leq C_\alpha |\tilde{c}_t|^2 = C_\alpha |y|^\alpha.$$

Turning to (2.3), we use that $c^{t,y} \geq 0$ and $-\beta < 0$ to estimate

$$(2.6) \quad |A_s^{\pi,t,x,y}|^{-\beta} = \left(x + \int_t^s \frac{c_u^{t,y}}{Z_u^\pi} du \right)^{-\beta} |Z_s^\pi|^{-\beta} \leq |x|^{-\beta} |Z_s^\pi|^{-\beta} = |x|^{-\beta} |\tilde{Z}_s^\pi|^2,$$

where, again by Itô's lemma, $\tilde{Z}^\pi := |Z^\pi|^{-\beta/2}$ solves the (linear) Lipschitz SDE

$$d\tilde{Z}_s^\pi = -\frac{\beta}{2} \tilde{Z}_s^\pi \left[\left(r + \sigma\theta\pi_s - \frac{1}{4}(\beta + 2)\sigma^2\pi_s^2 \right) ds + \sigma\pi_s dW_s^1 \right], \quad s \in [t, T], \quad \tilde{Z}_t = 1.$$

As before, we conclude that there exists a constant $C_\beta > 0$ (independent of π as the Lipschitz constant in the SDE for \tilde{Z}^π can be chosen independent of π since the set A is bounded) such that

$$\mathbb{E} \left[\sup_{s \in [t, T]} |A_s^{\pi,t,x,y}|^{-\beta} \right] \leq |x|^{-\beta} \mathbb{E} \left[\sup_{s \in [t, T]} |\tilde{Z}_s^\pi|^2 \right] \leq C_\beta |x|^{-\beta} |\tilde{Z}_t^\pi|^2 = C_\beta |x|^{-\beta}$$

and hence (2.3) is obtained. Finally, regarding (2.4), we note that

$$\mathbb{E} \left[\sup_{s \in [t, T]} |A_s^{\pi,t,x,y}|^\eta \right] \leq \mathbb{E} \left[\sup_{s \in [t, T]} |(A_s^{\pi,t,x,y}, c_s^{t,y})|^\eta \right] \leq \mathbb{E} \left[\sup_{s \in [t, T]} |(A_s^{\pi,t,x,y}, c_s^{t,y})|^{2\vee\eta} \right]^{(\eta/2) \wedge 1},$$

where we have to use Jensen's inequality in the last step if $\eta < 2$. Applying again the a priori estimate for Lipschitz SDEs yields the existence of $\tilde{C}_\eta > 0$ independent of π such that

$$\mathbb{E} \left[\sup_{s \in [t, T]} |(A_s^{\pi,t,x,y}, c_s^{t,y})|^{2\vee\eta} \right] \leq \tilde{C}_\eta |(x, y)|^{2\vee\eta},$$

which implies (2.4) with $C_\eta := |\tilde{C}_\eta|^{(\eta/2) \wedge 1}$. \square

The a priori estimates can be used to provide simple sufficient conditions under which the value function is finite and $\mathbb{E}[U(A_T^{\pi,t,x,y})]$ is bounded uniformly from below for all admissible strategies. As usual we define the positive and negative part of a function $U^+(x) := \max\{U(x), 0\}$ and $U^-(x) := \max\{-U(x), 0\}$. Moreover, we remark that the Inada condition at infinity, i.e.

$\lim_{x \uparrow \infty} U'(x) = 0$, implies that there exist constants $\eta \in (0, 1]$ and $K_+ \geq 0$ such that

$$(2.7) \quad U^+(x) \leq K_+(1 + |x|^\eta), \quad x > 0.$$

E.g., in the case of power utility $U = U_\gamma$, this estimate holds for $(K_+, \eta) = (1/\gamma, \gamma)$ if $\gamma \in (0, 1)$ and $K_+ = 0$ if $\gamma < 0$.

Proposition 2.2 (Local boundedness). *(i) Let $\eta \in (0, 1]$ such that (2.7) holds. Then there exists a constant $C_+ > 0$ such that*

$$\sup_{\pi \in \mathcal{A}} E[U^+(A_T^{\pi, t, x, y})] \leq C_+(1 + |(x, y)|^\eta), \quad (t, x, y) \in [0, T] \times \mathcal{O}.$$

(ii) Suppose that there exist constants $K_- \geq 0$ and $\beta > 0$ such that $U^-(x) \leq K_-(1 + x^{-\beta})$. Then there exists a constant $C_- > 0$ such that

$$\sup_{\pi \in \mathcal{A}} E[U^-(A_T^{\pi, t, x, y})] \leq C_-(1 + |x|^{-\beta}), \quad (t, x, y) \in [0, T] \times \mathcal{O}.$$

In particular, in this situation, the value function \mathcal{V} is locally bounded, i.e. there exists a constant $C_{\mathcal{V}} > 0$ such that

$$|\mathcal{V}(t, x, y)| \leq C_{\mathcal{V}}(1 + |x|^{-\beta} + |x|^\eta + |y|^\eta), \quad (t, x, y) \in [0, T] \times \mathcal{O}.$$

Proof. Let us fix $(t, x, y) \in \mathcal{O}$. To prove (i), we apply the estimate (2.7) followed by the a priori estimate (2.4) to obtain the existence of a constant $C_\eta > 0$ with

$$\sup_{\pi \in \mathcal{A}} E[U^+(A_T^{\pi, t, x, y})] \leq K_+ \left(1 + \sup_{\pi \in \mathcal{A}} E[|A_T^{\pi, t, x, y}|^\eta]\right) \leq K_+ \left(1 + C_\eta |(x, y)|^\eta\right).$$

The proof of (ii) works analogously by using (2.3) in place of (2.4): There exists $C_\beta > 0$ such that

$$\sup_{\pi \in \mathcal{A}} E[U^-(A_T^{\pi, t, x, y})] \leq K_- \left(1 + \sup_{\pi \in \mathcal{A}} E[|A_T^{\pi, t, x, y}|^{-\beta}]\right) \leq K_- \left(1 + C_- |x|^{-\beta}\right).$$

The estimate for \mathcal{V} is an immediate consequence of (i) and (ii). \square

Note that for power utility functions U_γ with $\gamma < 0$ the assumption of part (ii) of the proposition is satisfied with $(K_-, \beta) = (-1/\gamma, -\gamma)$ and that for $\gamma < 0$ the positive part and for $0 < \gamma < 1$ the negative part vanishes. The logarithmic utility $U = \log$ is an example of a common utility function for which the result applies and for which neither the positive nor negative part is zero.

In the next step, we derive continuity properties of the value function. This is done under the same assumptions as in the previous proposition together with a suitable growth condition on the derivative of the utility function.

Proposition 2.3 (Continuity). *Suppose that there exist constants $K_- \geq 0$ and $\beta > 0$ as well as*

constants $K' \geq 0$ and $\beta' > 0$ such that

$$(2.8) \quad U^-(x) \leq K_-(1 + x^{-\beta}) \quad \text{and} \quad U'(x) \leq K'x^{-\beta'} \quad \text{for all } x > 0.$$

Then we can find $C' > 0$ such that

$$|\mathcal{V}(t, x, y) - \mathcal{V}(\bar{t}, \bar{x}, \bar{y})| \leq C' \min\{|x|, |\bar{x}|\}^{-\beta'} \left(|(x, y) - (\bar{x}, \bar{y})| + \max\{|(x, y)|, |(\bar{x}, \bar{y})|\} |t - \bar{t}|^{1/2} \right)$$

for all $(t, x, y), (\bar{t}, \bar{x}, \bar{y}) \in [0, T] \times \mathcal{O}$. In particular, \mathcal{V} is locally Lipschitz continuous in (x, y) and locally $1/2$ -Hölder continuous in t .

Proof. Let $(t, x, y), (\bar{t}, \bar{x}, \bar{y}) \in [0, T] \times \mathcal{O}$ and assume, without loss of generality, $\mathcal{V}(t, x, y) \geq \mathcal{V}(\bar{t}, \bar{x}, \bar{y})$. For $\varepsilon > 0$, we let $\pi \in \mathcal{A}$ be an ε -optimal strategy in that

$$\mathcal{V}(t, x, y) \leq \mathbb{E}[U(A_T)] + \varepsilon, \quad \text{where } A_T := A_T^{\pi, t, x, y}.$$

Note that π exists since $\mathcal{V}(t, x, y) > -\infty$. Moreover, Proposition 2.2 guarantees that

$$\mathcal{V}(\bar{t}, \bar{x}, \bar{y}) \geq \mathbb{E}[U(\bar{A}_T)] > -\infty, \quad \text{where } \bar{A}_T := A_T^{\pi, \bar{t}, \bar{x}, \bar{y}}.$$

With this, using the concavity of U , the growth assumption on U' , and finally Hölder's inequality, we obtain the estimate

$$(2.9) \quad \begin{aligned} |\mathcal{V}(t, x, y) - \mathcal{V}(\bar{t}, \bar{x}, \bar{y})| &= \mathcal{V}(t, x, y) - \mathcal{V}(\bar{t}, \bar{x}, \bar{y}) \leq \mathbb{E}[U(A_T) - U(\bar{A}_T)] + \varepsilon \\ &\leq \mathbb{E}[U'(\bar{A}_T)|A_T - \bar{A}_T|] + \varepsilon \\ &\leq K' \mathbb{E}[|\bar{A}_T|^{-\beta'} |A_T - \bar{A}_T|] + \varepsilon \\ &\leq K' \mathbb{E}[|\bar{A}_T|^{-2\beta'}]^{1/2} \mathbb{E}[|A_T - \bar{A}_T|^2]^{1/2} + \varepsilon. \end{aligned}$$

Applying the a priori estimate (2.3) now shows that

$$(2.10) \quad \mathbb{E}[|\bar{A}_T|^{-2\beta'}]^{1/2} \leq |C_{2\beta'}|^{1/2} |\bar{x}|^{-\beta'} \leq |C_{2\beta'}|^{1/2} \min\{|x|, |\bar{x}|\}^{-\beta'}$$

and the a priori estimate (2.5) yields

$$(2.11) \quad \begin{aligned} \mathbb{E}[|A_T - \bar{A}_T|^2]^{1/2} &\leq C \left(|(x, y) - (\bar{x}, \bar{y})|^2 + |(x, y)|^2 |t - \bar{t}| \right)^{1/2} \\ &\leq C \left(|(x, y) - (\bar{x}, \bar{y})| + \max\{|(x, y)|, |(\bar{x}, \bar{y})|\} |t - \bar{t}|^{1/2} \right), \end{aligned}$$

where we have used the subadditivity of the square root function for the last step. Note that neither of the constants $K', C_{2\beta'}, C$ depends on the strategy π and hence $C' := CK'|C_{2\beta'}|^{1/2}$ does not depend on π either. Now plugging (2.10) and (2.11) into (2.9) leads to

$$|\mathcal{V}(t, x, y) - \mathcal{V}(\bar{t}, \bar{x}, \bar{y})| \leq C' \min\{|x|, |\bar{x}|\}^{-\beta'} \left(|(x, y) - (\bar{x}, \bar{y})| + \max\{|(x, y)|, |(\bar{x}, \bar{y})|\} |t - \bar{t}|^{1/2} \right) + \varepsilon.$$

The result follows as ε is chosen arbitrarily and C' does not depend on ε . \square

Note that the power utility function satisfies the assumptions set forth in the previous proposition with $K' = 1$ and $\beta' = \gamma - 1$. For later reference, we gather these results in a corollary.

Corollary 2.4 (Power utility). *Suppose that U is a power utility function, i.e. $U = U_\gamma$ for $\gamma < 1$ with $\gamma \neq 0$. Then there exists a constant $C_\mathcal{V} > 0$ such that*

$$(2.12) \quad |\mathcal{V}(t, x, y)| \leq C_\mathcal{V} \left(1 + |x|^{-\gamma^-} + |x|^{\gamma^+} + |y|^{\gamma^+} \right), \quad (t, x, y) \in [0, T] \times \mathcal{O},$$

where $\gamma^- := \max\{-\gamma, 0\}$ and $\gamma^+ := \max\{\gamma, 0\}$. Moreover, \mathcal{V} is continuous and satisfies

$$|\mathcal{V}(t, x, y) - \mathcal{V}(\bar{t}, \bar{x}, \bar{y})| \leq C' \min\{|x|, |\bar{x}|\}^{1-\gamma} \left(|(x, y) - (\bar{x}, \bar{y})| + \max\{|(x, y)|, |(\bar{x}, \bar{y})|\} |t - \bar{t}|^{1/2} \right)$$

for all $(t, x, y), (\bar{t}, \bar{x}, \bar{y}) \in [0, T] \times \mathcal{O}$ for a constant $C' > 0$.

2.2 Further properties of the value function

Before turning to the HJB equation associated with the optimization problem, we first study some further properties of the value function. In particular, we shall see that \mathcal{V} is concave and monotone in the spatial variables. Moreover, in the special case of power utility, we show that \mathcal{V} is homogeneous of degree γ .

Proposition 2.5 (Monotonicity and concavity). *\mathcal{V} is increasing and jointly concave in the spatial variables. In particular, for every fixed $t \in [0, T]$, the mapping $(x, y) \mapsto \mathcal{V}(t, x, y)$ is locally Lipschitz continuous on the interior of its effective domain $\{(x, y) \in \mathcal{O} : \mathcal{V}(t, x, y) > -\infty\}$.*

We note that, in Proposition 2.3, we have established a stronger continuity result. However, for this we have to assume (2.8), whereas Proposition 2.5 is valid without additional assumptions.

Proof of Prop. 2.5. The proof works along standard lines (cf., e.g., Section 3.6.1 in [31]). Note that continuity follows immediately from concavity, as every concave function is locally Lipschitz continuous in the interior of its effective domain.

Step 1: Monotonicity. Let us fix $t \in [0, T]$, $(x_1, y_1), (x_2, y_2) \in \mathcal{O}$ with $x_1 \leq x_2$ and $y_1 \leq y_2$, and a strategy $\pi \in \mathcal{A}$. Since the endowment process is a geometric Brownian motion we see that

$$c_T^{t, y_1} = y_1 c_T^{t, 1} \leq y_2 c_T^{t, 1} = c_T^{t, y_2}$$

and from the explicit representation of the wealth process we find that

$$A_T^{\pi, t, x_1, y_1} = \left(x_1 + \int_t^T \frac{c_s^{t, y_1}}{Z_s^\pi} ds \right) Z_T^\pi \leq \left(x_2 + \int_t^T \frac{c_s^{t, y_2}}{Z_s^\pi} ds \right) Z_T^\pi = A_T^{\pi, t, x_2, y_2}.$$

Since π was chosen arbitrarily and the utility function U is monotone, this shows that

$$\mathcal{V}(t, x_1, y_1) = \sup_{\pi \in \mathcal{A}} \mathbb{E}[U(A_T^{\pi, t, x_1, y_1})] \leq \sup_{\pi \in \mathcal{A}} \mathbb{E}[U(A_T^{\pi, t, x_2, y_2})] = \mathcal{V}(t, x_2, y_2),$$

i.e. \mathcal{V} is monotone in both its second and third argument.

Step 2: Concavity. Let us again fix $t \in [0, T]$ and $(x_1, y_1), (x_2, y_2) \in \mathcal{O}$. Let moreover $\pi^1, \pi^2 \in \mathcal{A}$, choose $\vartheta \in [0, 1]$, and define

$$x_\vartheta := \vartheta x_1 + (1 - \vartheta)x_2 \quad \text{and} \quad y_\vartheta := \vartheta y_1 + (1 - \vartheta)y_2.$$

Moreover, with $A^i := A^{\pi^i, t, x_i, y_i}$, $i = 1, 2$, we define

$$A^\vartheta := \vartheta A^1 + (1 - \vartheta)A^2 \quad \text{and} \quad \pi^\vartheta := \frac{\vartheta A^1 \pi^1 + (1 - \vartheta)A^2 \pi^2}{A^\vartheta}.$$

As π^ϑ is a convex combination of π^1 and π^2 , it follows that π^ϑ is A -valued and hence π^ϑ is admissible. Let us examine that A^ϑ is the wealth process corresponding to π^ϑ . For this, observe that

$$c^{t, y_\vartheta} = y_\vartheta c^{t, 1} = \vartheta y_1 c^{t, 1} + (1 - \vartheta)y_2 c^{t, 1} = \vartheta c^{t, y_1} + (1 - \vartheta)c^{t, y_2},$$

implying that

$$\begin{aligned} dA_s^\vartheta &= \vartheta dA_s^1 + (1 - \vartheta) dA_s^2 \\ &= r[\vartheta A_s^1 + (1 - \vartheta)A_s^2] ds + \sigma\theta[\vartheta\pi_s^1 A_s^1 + (1 - \vartheta)\pi_s^2 A_s^2] ds + [\vartheta c_s^{t, y_1} + (1 - \vartheta)c_s^{t, y_2}] ds \\ &\quad + \sigma[\vartheta\pi_s^1 A_s^1 + (1 - \vartheta)\pi_s^2 A_s^2] dW_s^1 \\ &= [(r + \sigma\theta\pi_s^\vartheta)A_s^\vartheta + c_s^{t, y_\vartheta}] ds + \sigma\pi_s^\vartheta A_s^\vartheta dW_s^1 \end{aligned}$$

and hence uniqueness of solutions of linear SDEs implies that $A^{\pi^\vartheta, t, x_\vartheta, y_\vartheta} = A^\vartheta = \vartheta A^1 + (1 - \vartheta)A^2$, i.e. A^ϑ is the wealth process corresponding to π^ϑ started in $(t, x_\vartheta, y_\vartheta)$. Using the concavity of the utility function, we get

$$\mathcal{V}(t, x_\vartheta, y_\vartheta) \geq \mathbb{E}[U(A_T^\vartheta)] \geq \vartheta \mathbb{E}[U(A_T^1)] + (1 - \vartheta) \mathbb{E}[U(A_T^2)],$$

but since π^1, π^2 are chosen arbitrary, taking the supremum on the right hand side, we arrive at

$$\mathcal{V}(t, x_\vartheta, y_\vartheta) \geq \vartheta \mathcal{V}(t, x_1, y_1) + (1 - \vartheta)\mathcal{V}(t, x_2, y_2). \quad \square$$

We conclude this section by showing that, in the power utility case $U = U_\gamma$, the value function is homogeneous of degree γ in the spatial variables.

Lemma 2.6 (Homogeneity). *Suppose that $U = U_\gamma$ with $\gamma < 1$, $\gamma \neq 0$. Then the value function \mathcal{V} is homogeneous of degree γ in the spatial variables, i.e.*

$$\mathcal{V}(t, kx, ky) = k^\gamma \mathcal{V}(t, x, y) \quad \text{for all } k > 0 \text{ and } (t, x, y) \in [0, T] \times \mathcal{O}.$$

As a consequence, \mathcal{V} can be represented in separable form as

$$\mathcal{V}(t, x, y) = y^\gamma \mathcal{U}(t, x/y), \quad (t, x, y) \in [0, T] \times \mathcal{O},$$

where

$$(2.13) \quad \mathcal{U} : [0, T] \times (0, \infty) \rightarrow \mathbb{R}, \quad (t, z) \mapsto \mathcal{U}(t, z) := \mathcal{V}(t, z, 1).$$

Proof. The separable representation of \mathcal{V} follows immediately from homogeneity with $k = y$ since

$$\mathcal{V}(t, x, y) = \mathcal{V}(t, y(x/y), y) = y^\gamma \mathcal{V}(t, x/y, 1), \quad (t, x, y) \in [0, T] \times \mathcal{O}.$$

To see that \mathcal{V} is homogeneous, fix $(t, x, y) \in [0, T] \times \mathcal{O}$, $k > 0$, and $\pi \in \mathcal{A}$. Since $c^{t, ky} = kc^{t, y}$ we find that

$$A_T^{\pi, t, kx, ky} = \left(kx + \int_t^T \frac{c_s^{t, ky}}{Z_s^\pi} ds \right) Z_T^\pi = k \left(x + \int_t^T \frac{c_s^{t, y}}{Z_s^\pi} ds \right) Z_T^\pi = k A_T^{\pi, t, x, y}.$$

But then \mathcal{V} must be homogeneous of degree γ since

$$\mathcal{V}(t, kx, ky) = \sup_{\pi \in \mathcal{A}} \mathbb{E} [U_\gamma(A_T^{\pi, t, kx, ky})] = \sup_{\pi \in \mathcal{A}} \mathbb{E} \left[\frac{1}{\gamma} (k A_T^{\pi, t, x, y})^\gamma \right] = k^\gamma \mathcal{V}(t, x, y). \quad \square$$

3 The Hamilton-Jacobi-Bellman equation

Henceforth, we restrict our attention to power utility, i.e. $U = U_\gamma$. Applying well-known principles of stochastic control, see e.g. Chapter 3 in [31], we can write down the HJB equation for the value function of our control problem:

$$(3.1) \quad -\mathcal{V}_t(t, x, y) - \sup_{\pi \in \mathcal{A}} \{ \mathbb{L}^\pi \mathcal{V}(t, x, y) \} = 0, \quad (t, x, y) \in [0, T] \times \mathcal{O},$$

$$\mathcal{V}(t, x, y) = U_\gamma(x), \quad (x, y) \in \mathcal{O},$$

where the linear differential operator \mathbb{L}^π is defined through

$$\mathbb{L}^\pi \mathcal{V}(t, x, y) := [(r + \sigma\theta\pi)x + y] \mathcal{V}_x + \mu_C y \mathcal{V}_y + \frac{1}{2} \sigma^2 \pi^2 x^2 \mathcal{V}_{xx} + \frac{1}{2} \sigma_C^2 y^2 \mathcal{V}_{yy} + \rho \sigma \sigma_C \pi x y \mathcal{V}_{xy}.$$

Here, and in the following, we denote partial derivatives by subscripts and often omit the argument of functions in the equations. The aim of this section is to characterize the value function as the unique viscosity solution of the HJB equation. Moreover, we show that \mathcal{V} is even a classical solution provided that μ_C and σ_C are continuously differentiable.

3.1 A family of supersolutions

Before linking the value function to the HJB equation, let us first study the existence of classical supersolutions of the HJB equation. These play a crucial role for our subsequent analysis and serve, in particular, as abstract boundary/growth conditions for the HJB equation.

We begin by defining two constants

$$K := r + \frac{\theta^2}{2(1-\gamma)}$$

as well as

$$\lambda := \max\left\{-r - \inf_{\pi \in A} \left\{ \sigma\theta\pi - \frac{1}{2}(1+\kappa)\sigma^2\pi^2 \right\}, - \inf_{t \in [0, T]} \left\{ \mu_C(t) - \frac{1}{2}(1+\kappa)\sigma_C(t)^2 \right\}\right\}.$$

With this, we introduce a parametric family of functions

$$\psi_{\gamma, \varepsilon, \iota} : [0, T] \times \mathcal{O} \rightarrow \mathbb{R}, \quad (t, x, y) \mapsto \psi_{\gamma, \varepsilon, \iota}(t, x, y)$$

for $\gamma < 1$ with $\gamma \neq 0$, $\varepsilon \in [0, 1]$, and $\iota \in \{0, 1\}$. We assume that $\psi_{\gamma, \varepsilon, \iota}$ takes the form

$$\psi_{\gamma, \varepsilon, \iota}(t, x, y) := U_\gamma\left(x + y\varphi_\iota(t) + \varepsilon e^{-r(T-t)}\right) e^{\gamma K(T-t)} + \iota(x^{-(1+|\gamma|)} + y^{-(1+|\gamma|)}) e^{(1+|\gamma|)\lambda(T-t)}$$

for all $(t, x, y) \in [0, T] \times \mathcal{O}$, where the function $\varphi_\iota : [0, T] \rightarrow \mathbb{R}$ is given as the unique solution of the ordinary differential equation

$$\dot{\varphi}_\iota(t) = (r - \mu_C(t) + \rho\theta\sigma_C(t))\varphi_\iota(t) - 1, \quad t \in [0, T], \quad \varphi_\iota(T) = \iota.$$

Observe that $\varphi_\iota(t) > 0$ unless $\iota = 0$ and $t = T$. In any case, φ_ι is nonnegative and hence, in particular, $\psi_{\gamma, \varepsilon, \iota}$ well-defined. We proceed to show that $\psi_{\gamma, \varepsilon, \iota}$ is a classical supersolution of the HJB equation.

Proposition 3.1 (Classical supersolution). *For any choice of $\gamma < 1$ with $\gamma \neq 0$, $\varepsilon \in [0, 1]$, and $\iota \in \{0, 1\}$, the function $\psi := \psi_{\gamma, \varepsilon, \iota}$ is a supersolution of the HJB equation, i.e.*

$$-\psi_t(t, x, y) - \sup_{\pi \in A} \{L^\pi \psi(t, x, y)\} \geq 0, \quad (t, x, y) \in [0, T] \times \mathcal{O}.$$

Moreover, it is a strict supersolution if $\iota = 1$.

Proof. We decompose ψ as $\psi = \psi^1 + \psi^2$, where

$$\begin{aligned} \psi^1(t, x, y) &:= U_\gamma\left(x + y\varphi_\iota(t) + \varepsilon e^{-r(T-t)}\right) e^{\gamma K(T-t)}, & (t, x, y) \in [0, T] \times \mathcal{O}, \\ \psi^2(t, x, y) &:= \iota(x^{-(1+|\gamma|)} + y^{-(1+|\gamma|)}) e^{(1+|\gamma|)\lambda(T-t)}, & (t, x, y) \in [0, T] \times \mathcal{O}. \end{aligned}$$

By linearity of the operator L^π , it follows that

$$-\psi_t - \sup_{\pi} \{L^\pi \psi\} \geq \left(-\psi_t^1 - \sup_{\pi} \{L^\pi \psi^1\}\right) + \left(-\psi_t^2 - \sup_{\pi} \{L^\pi \psi^2\}\right).$$

Hence, in order to conclude, it suffices to show that ψ^1 and ψ^2 are supersolutions, and ψ^2 is a strict supersolution if $\iota = 1$.

Step 1: ψ^2 is a (strict) supersolution. If $\iota = 0$, then $\psi^2 \equiv 0$ and it is thus a solution of the HJB

equation. Let us hence assume that $\iota = 1$ and prove that, in this case, ψ^2 is a strict supersolution. Setting $\kappa := (1 + |\gamma|)$, the partial derivatives of ψ^2 are given by

$$\begin{aligned}\psi_t^2 &= -\kappa\lambda(x^{-\kappa} + y^{-\kappa})e^{\kappa\lambda(T-t)}, & \psi_x^2 &= -\kappa x^{-\kappa-1}e^{\kappa\lambda(T-t)}, & \psi_y^2 &= -\kappa y^{-\kappa-1}e^{\kappa\lambda(T-t)}, \\ \psi_{xx}^2 &= \kappa(1 + \kappa)x^{-\kappa-2}e^{\kappa\lambda(T-t)}, & \psi_{yy}^2 &= \kappa(1 + \kappa)y^{-\kappa-2}e^{\kappa\lambda(T-t)}, & \psi_{xy}^2 &= 0.\end{aligned}$$

Plugging these derivatives into the HJB equation and rearranging terms yields

$$\begin{aligned}-\psi_t^2 - \sup_{\pi \in A} \{\mathbf{L}^\pi \psi^2\} &= \kappa \inf_{\pi \in A} \left\{ \lambda + r + \sigma\theta\pi - \frac{1}{2}(1 + \kappa)\sigma^2\pi^2 \right\} x^{-\kappa} e^{\kappa\lambda(T-t)} + \kappa y x^{-\kappa-1} e^{\kappa\lambda(T-t)} \\ &\quad + \kappa \left(\lambda + \mu_C(t) - \frac{1}{2}(1 + \kappa)\sigma_C(t)^2 \right) y^{-\kappa} e^{\kappa\lambda(T-t)} \\ &\geq \kappa \left(\lambda + r + \inf_{\pi \in A} \left\{ \sigma\theta\pi - \frac{1}{2}(1 + \kappa)\sigma^2\pi^2 \right\} \right) x^{-\kappa} e^{\kappa\lambda(T-t)} + \kappa y x^{-\kappa-1} e^{\kappa\lambda(T-t)} \\ &\quad + \kappa \left(\lambda + \inf_{t \in [0, T]} \left\{ \mu_C(t) - \frac{1}{2}(1 + \kappa)\sigma_C(t)^2 \right\} \right) y^{-\kappa} e^{\kappa\lambda(T-t)}.\end{aligned}$$

By the choice of λ , the terms inside the brackets are nonnegative and hence

$$-\psi_t^2 - \sup_{\pi \in A} \{\mathbf{L}^\pi \psi^2\} \geq \kappa y x^{-\kappa-1} e^{\kappa\lambda(T-t)} > 0,$$

i.e. ψ^2 is a strict supersolution of the HJB equation.

Step 2: ψ^1 is a supersolution. Writing $\varphi = \varphi_\iota$ and using the ordinary differential equation for φ , the partial derivatives of ψ^1 are given by

$$\begin{aligned}\psi_t^1 &= -K \left(x + y\varphi(t) + \varepsilon e^{-r(T-t)} \right)^\gamma e^{\gamma K(T-t)} \\ &\quad + \left((r - \mu_C(t) + \rho\theta\sigma_C(t))y\varphi(t) - y + r\varepsilon e^{-r(T-t)} \right) \left(x + y\varphi(t) + \varepsilon e^{-r(T-t)} \right)^{\gamma-1} e^{\gamma K(T-t)}, \\ \psi_x^1 &= \left(x + y\varphi(t) + \varepsilon e^{-r(T-t)} \right)^{\gamma-1} e^{\gamma K(T-t)}, \\ \psi_y^1 &= \varphi(t) \left(x + y\varphi(t) + \varepsilon e^{-r(T-t)} \right)^{\gamma-1} e^{\gamma K(T-t)}, \\ \psi_{xx}^1 &= -(1 - \gamma) \left(x + y\varphi(t) + \varepsilon e^{-r(T-t)} \right)^{\gamma-2} e^{\gamma K(T-t)}, \\ \psi_{yy}^1 &= -(1 - \gamma)\varphi(t)^2 \left(x + y\varphi(t) + \varepsilon e^{-r(T-t)} \right)^{\gamma-2} e^{\gamma K(T-t)}, \\ \psi_{xy}^1 &= -(1 - \gamma)\varphi(t) \left(x + y\varphi(t) + \varepsilon e^{-r(T-t)} \right)^{\gamma-2} e^{\gamma K(T-t)}.\end{aligned}$$

Plugging these derivatives into the HJB equation and rearranging terms yields

$$\begin{aligned}-\psi_t^1(t, x, y) - \sup_{\pi \in A} \{\mathbf{L}^\pi \psi^1(t, x, y)\} \\ = -e^{\gamma K(T-t)} \sup_{\pi \in A} \left\{ -K \left(x + y\varphi(t) + \varepsilon e^{-r(T-t)} \right)^\gamma \right. \\ \left. + \left[(r + \sigma\theta\pi)x + (r + \rho\theta\sigma_C(t))y\varphi(t) + r\varepsilon e^{-r(T-t)} \right] \left(x + y\varphi(t) + \varepsilon e^{-r(T-t)} \right)^{\gamma-1} \right\}\end{aligned}$$

$$-\frac{1}{2}(1-\gamma)\left[\sigma^2\pi^2x^2+2\rho\sigma\sigma_C(t)\pi xy\varphi(t)+\sigma_C(t)^2y^2\varphi(t)^2\right]\left(x+y\varphi(t)+\varepsilon e^{-r(T-t)}\right)^{\gamma-2}\}.$$

Instead of maximizing over A we can maximize over all of \mathbb{R} , making the last expression smaller in doing so. The maximizer in the supremum is given by

$$\hat{\pi} := \frac{1}{(1-\gamma)\sigma x} \left[\theta \left(x + y\varphi(t) + \varepsilon e^{-r(T-t)} \right) - (1-\gamma)\rho\sigma_C(t)y\varphi(t) \right].$$

Plugging this into the above equation and rearranging terms again then yields

$$\begin{aligned} & -\psi_t^1(t, x, y) - \sup_{\pi \in \mathcal{A}} \{ \mathbf{L}^\pi \psi^1(t, x, y) \} \\ & \geq -e^{\gamma K(T-t)} \left[-K \left(x + y\varphi(t) + \varepsilon e^{-r(T-t)} \right)^\gamma \right. \\ & \quad + \left[(r + \sigma\theta\hat{\pi})x + (r + \rho\theta\sigma_C(t))y\varphi(t) + r\varepsilon e^{-r(T-t)} \right] \left(x + y\varphi(t) + \varepsilon e^{-r(T-t)} \right)^{\gamma-1} \\ & \quad \left. - \frac{1}{2}(1-\gamma) \left[\sigma^2\hat{\pi}^2x^2 + 2\rho\sigma\sigma_C(t)\hat{\pi}xy\varphi(t) + \sigma_C(t)^2y^2\varphi(t)^2 \right] \left(x + y\varphi(t) + \varepsilon e^{-r(T-t)} \right)^{\gamma-2} \right] \\ & = \left(K - r - \frac{\theta^2}{2(1-\gamma)} \right) \left(x + y\varphi(t) + \varepsilon e^{-r(T-t)} \right)^\gamma e^{\gamma K(T-t)} \\ & \quad + \frac{1}{2}(1-\gamma)(1-\rho)^2\sigma_C(t)^2y^2\varphi(t)^2 \left(x + y\varphi(t) + \varepsilon e^{-r(T-t)} \right)^{\gamma-2} e^{\gamma K(T-t)} \geq 0, \end{aligned}$$

where the nonnegativity follows from the choice of K . \square

With the existence of classical supersolutions, we can derive tight bounds on the value function.

Proposition 3.2 (Tight bounds on \mathcal{V}). *The value function \mathcal{V} satisfies*

$$U_\gamma(x)e^{\gamma K(T-t)} \leq \mathcal{V}(t, x, y) \leq U_\gamma(x + y\varphi_0(t))e^{\gamma K(T-t)}, \quad (t, x, y) \in [0, T] \times \mathcal{O}.$$

Proof. Step 1: We first prove the lower bound. For this, consider the constant strategy $\pi \equiv \pi_M \in \mathcal{A}$ and recall that $A_T^{\pi_M, t, x, y} \geq A_T^{\pi_M, t, x, 0} = xA_T^{\pi_M, t, 1, 0}$. But then

$$\mathcal{V}(t, x, y) \geq \mathbb{E}[U_\gamma(A_T^{\pi_M, t, x, y})] \geq \mathbb{E}[U_\gamma(xA_T^{\pi_M, t, 1, 0})] = U_\gamma(x) \mathbb{E}[(A_T^{\pi_M, t, 1, 0})^\gamma].$$

The last expectation can be computed explicitly and is given by

$$\begin{aligned} \mathbb{E}[(A_T^{\pi_M, t, 1, 0})^\gamma] &= \mathbb{E}\left[e^{\gamma(r + \sigma\theta\pi_M - \frac{1}{2}\sigma^2\pi_M^2)(T-t) + \gamma\sigma\pi_M(W_T^1 - W_t^1)} \right] \\ &= e^{\gamma(r + \sigma\theta\pi_M - \frac{1}{2}(1-\gamma)\sigma^2\pi_M^2)(T-t)} = e^{\gamma K(T-t)}. \end{aligned}$$

Step 2: To prove the upper bound, let $(t, x, y) \in [0, T] \times \mathcal{O}$ as well as $\pi \in \mathcal{A}$ and write $A := A^{\pi, t, x, y}$ and $c := c^{t, y}$. For $\varepsilon \in (0, 1]$ fixed, set $\psi^\varepsilon := \psi_{\gamma, \varepsilon, 0}$ and denote by ρ_n , $n \in \mathbb{N}$, a localizing sequence of the local martingale

$$M_s := \int_t^s \sigma\pi_u A_u \psi_x^\varepsilon(u, A_u, c_u) dW_u^1 + \int_t^s \sigma_C(u) c_u \psi_y^\varepsilon(u, A_u, c_u) dW_u^C, \quad s \in [t, T].$$

Applying Itô's lemma to $\psi^\varepsilon(\rho_n, A_{\rho_n}, c_{\rho_n})$ and using the fact that ψ^ε is a supersolution of the HJB equation, it follows that

$$\begin{aligned} \psi^\varepsilon(t, x, y) &= \mathbb{E} \left[\psi^\varepsilon(\rho_n, A_{\rho_n}, c_{\rho_n}) + \int_t^{\rho_n} [-\psi_t^\varepsilon(s, A_s, c_s) - L^{\pi_s} \psi^\varepsilon(s, A_s, c_s)] ds - M_{\rho_n} \right] \\ &\geq \mathbb{E} [\psi^\varepsilon(\rho_n, A_{\rho_n}, c_{\rho_n})]. \end{aligned}$$

As $\varepsilon > 0$, we see that ψ^ε is lower bounded. Hence we may apply Fatou's lemma to arrive at

$$\psi^\varepsilon(t, x, y) \geq \liminf_{n \rightarrow \infty} \mathbb{E} [\psi^\varepsilon(\rho_n, A_{\rho_n}, c_{\rho_n})] \geq \mathbb{E} [\psi^\varepsilon(T, A_T, c_T)] = \mathbb{E} [U_\gamma(A_T + \varepsilon)] \geq \mathbb{E} [U_\gamma(A_T)].$$

Since π was chosen arbitrarily, this implies that $\psi^\varepsilon(t, x, y) \geq \mathcal{V}(t, x, y)$ and hence

$$U_\gamma(x + y\varphi_0(t))e^{\gamma K(T-t)} = \lim_{\varepsilon \downarrow 0} \psi^\varepsilon(t, x, y) \geq \mathcal{V}(t, x, y). \quad \square$$

Remark 3.3. Another way to prove estimates on \mathcal{V} is to use financial arguments as follows:

- The lower bound corresponds to the value function of the problem without random endowments and can therefore be improved when random endowments are present;
- An alternative upper bound could be found by comparing the problem to the one of an artificial market model where the endowment can also be traded.

3.2 Viscosity characterization of the value function

The next step is to show that the value function is the unique viscosity solution of the HJB equation. We refer to [7] for the definition of (and all important results on) viscosity solutions of second order PDEs. While the fact that v is a viscosity solution of the HJB equation is standard, the uniqueness result poses significant challenges in the case of $\gamma < 0$ as the value function tends to $-\infty$ near the boundary of the state space. To handle this issue, we require a sufficiently strong comparison principle for the HJB equation.

Before turning to the comparison principle, let us record that the Hamiltonian

$$H : [0, T] \times \mathcal{O} \times \mathbb{R} \times \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}, \quad (t, x, y, p, M) \mapsto H(t, x, y, p, M)$$

given by

$$H(t, x, y, p, M) := \sup_{\pi \in A} \left\{ [(r + \sigma\theta\pi)x + y]p_1 + \mu_C(t)yp_2 + \frac{1}{2} \text{tr} [\Sigma_\pi(t, x, y)\Sigma_\pi(t, x, y)^\top M] \right\}$$

for $(t, x, y) \in [0, T] \times \mathcal{O}$, $p \in \mathbb{R}^2$ and $M \in \mathbb{R}^{2 \times 2}$ with

$$\Sigma_\pi(t, x, y) := \begin{pmatrix} \sigma\pi x & 0 \\ \rho\sigma_C(t)y & \sqrt{1 - \rho^2}\sigma_C(t)y \end{pmatrix},$$

is both finite everywhere and continuous as A is compact. We also note that if φ is a sufficiently

smooth function, it holds that

$$H(t, x, y, D\varphi(t, x, y), D^2\varphi(t, x, y)) = \sup_{\pi \in A} \{L^\pi \varphi(t, x, y)\},$$

where $D\varphi$ and $D^2\varphi$ denote, respectively, the gradient and the Hessian of φ with respect to the spatial variables (x, y) . One important consequence of the continuity of H is that, by Lemma V.6.1 in [17], we are allowed to apply the parabolic version of Ishii's lemma; see Theorem V.6.1 in [17].

Theorem 3.4 (Comparison principle). *Let $u, v : [0, T] \times \mathcal{O} \rightarrow \mathbb{R}$ such that u is an upper semi-continuous viscosity subsolution and v is a lower semi-continuous viscosity supersolution of the HJB equation with*

$$(3.2) \quad U_\gamma(x)e^{\gamma K(T-t)} \leq u(t, x, y), v(t, x, y) \leq U_\gamma(x + y\varphi_0(t))e^{\gamma K(T-t)}, \quad (t, x, y) \in [0, T] \times \mathcal{O}.$$

Suppose furthermore that $u(T, \cdot) \leq v(T, \cdot)$ on \mathcal{O} . Then

$$u(t, x, y) \leq v(t, x, y) \quad \text{for all } (t, x, y) \in [0, T] \times \mathcal{O}.$$

Proof. Step 1: Setup of the proof and definitions. We argue by contradiction and suppose that there exists $(t^*, x^*, y^*) \in [0, T] \times \mathcal{O}$ such that

$$(3.3) \quad u(t^*, x^*, y^*) - v(t^*, x^*, y^*) > 0.$$

For $\delta > 0$ and $n \in \mathbb{N}_0$, we introduce a function $\phi_n : [0, T] \times \mathcal{O} \times \mathcal{O} \rightarrow \mathbb{R}$ given by

$$\phi_n(t, x, y, \bar{x}, \bar{y}) := u(t, x, y) - v(t, \bar{x}, \bar{y}) - \delta\psi(t, x, y) - \delta\psi(t, \bar{x}, \bar{y}) - \frac{n}{2} \left[|x - \bar{x}|^2 + |y - \bar{y}|^2 \right].$$

Here, $\psi := \psi_{\bar{\gamma}, 0, 1}$ for some $\bar{\gamma} \in (0, 1)$ with $\bar{\gamma} > \gamma$. We recall that $\psi : [0, T] \times \mathcal{O} \rightarrow \mathbb{R}$ is a strict classical supersolution of the HJB equation and takes the form

$$\psi(t, x, y) = U_{\bar{\gamma}}(x + y\varphi_1(t))e^{\bar{\gamma}K(T-t)} + (x^{-(1+|\gamma|)} + y^{-(1+|\gamma|)})e^{(1+|\gamma|)\lambda(T-t)}, \quad (t, x, y) \in [0, T] \times \mathcal{O},$$

where φ_1 is strictly positive on $[0, T]$. Finally, we define a function $\phi : [0, T] \times \mathcal{O} \rightarrow \mathbb{R}$ by

$$\phi(t, x, y) := u(t, x, y) - v(t, x, y) - 2\delta\psi(t, x, y), \quad (t, x, y) \in [0, T] \times \mathcal{O},$$

and we shall subsequently assume that $\delta > 0$ is sufficiently small to guarantee that

$$\phi(t^*, x^*, y^*) = u(t^*, x^*, y^*) - v(t^*, x^*, y^*) - 2\delta\psi(t^*, x^*, y^*) > 0,$$

which is possible by (3.3).

Step 2: Maximizers of ϕ_n . For each $n \in \mathbb{N}_0$, we define

$$M_n := \sup_{(t, x, y, \bar{x}, \bar{y}) \in [0, T] \times \mathcal{O} \times \mathcal{O}} \phi_n(t, x, y, \bar{x}, \bar{y}) \quad \text{and} \quad M := \sup_{(t, x, y) \in [0, T] \times \mathcal{O}} \phi(t, x, y).$$

Let us first observe that $M_n \geq M_{n+1} \geq M > 0$ for all $n \in \mathbb{N}_0$ as, clearly, ϕ_n is decreasing in n (implying $M_n \geq M_{n+1}$) and

$$M_n = \sup_{(t,x,y,\bar{x},\bar{y}) \in [0,T] \times \mathcal{O} \times \mathcal{O}} \phi_n(t, x, y, \bar{x}, \bar{y}) \geq \sup_{(t,x,y) \in [0,T] \times \mathcal{O}} \phi(t, x, y) = M \geq \phi(t^*, x^*, y^*) > 0.$$

Moreover, using (3.2), we note that for each $n \in \mathbb{N}_0$ and $(t, x, y, \bar{x}, \bar{y}) \in [0, T] \times \mathcal{O} \times \mathcal{O}$ we have

$$\begin{aligned} \phi_n(t, x, y, \bar{x}, \bar{y}) &\leq u(t, x, y) - v(t, \bar{x}, \bar{y}) - \delta\psi(t, x, y) - \delta\psi(t, \bar{x}, \bar{y}) \\ &\leq \left[U_\gamma(x + y\varphi_0(t)) - \delta U_{\bar{\gamma}}(x + y\varphi_1(t)) - U_\gamma(\bar{x}) - \delta U_{\bar{\gamma}}(\bar{x} + \bar{y}\varphi_1(t)) \right] e^{\bar{\gamma}K(T-t)} \\ &\quad - \delta \left[x^{-(1+|\gamma|)} + y^{-(1+|\gamma|)} + \bar{x}^{-(1+|\gamma|)} + \bar{y}^{-(1+|\gamma|)} \right] e^{(1+|\gamma|)\lambda(T-t)}. \end{aligned}$$

Since $\bar{\gamma} > \max\{\gamma, 0\}$ and $\varphi_1 > 0$, it follows that

$$\sup_{n \in \mathbb{N}_0} \sup_{t \in [0, T]} \phi_n(t, x, y, \bar{x}, \bar{y}) \rightarrow -\infty \quad \text{as } x \rightarrow \infty \text{ or } y \rightarrow \infty \text{ or } \bar{x} \rightarrow \infty \text{ or } \bar{y} \rightarrow \infty.$$

Similarly, we see that

$$\sup_{n \in \mathbb{N}_0} \sup_{t \in [0, T]} \phi_n(t, x, y, \bar{x}, \bar{y}) \rightarrow -\infty \quad \text{as } x \rightarrow 0 \text{ or } y \rightarrow 0 \text{ or } \bar{x} \rightarrow 0 \text{ or } \bar{y} \rightarrow 0.$$

In particular, as ϕ_0 is upper semicontinuous, we find that

$$K := \{(t, x, y, \bar{x}, \bar{y}) \in [0, T] \times \mathcal{O} \times \mathcal{O} : \phi_0(t, x, y, \bar{x}, \bar{y}) \geq 0\} \text{ is compact,}$$

and, as $M_n > 0$ and $\phi_n \leq \phi_0$, any maximizing sequence for any ϕ_n , $n \in \mathbb{N}_0$, is contained in K . Moreover, upper semicontinuity of ϕ_n implies that we find $(t_n, x_n, y_n, \bar{x}_n, \bar{y}_n) \in K$ such that

$$M_n = \sup_{(t,x,y,\bar{x},\bar{y}) \in [0,T] \times \mathcal{O} \times \mathcal{O}} \phi_n(t, x, y, \bar{x}, \bar{y}) = \phi_n(t_n, x_n, y_n, \bar{x}_n, \bar{y}_n).$$

Step 3: Convergence of maximizers. As K is compact and $(t_n, x_n, y_n, \bar{x}_n, \bar{y}_n) \in K$ for all $n \in \mathbb{N}_0$, after passing to a subsequence if necessary, it follows that

$$\lim_{n \rightarrow \infty} (t_n, x_n, y_n, \bar{x}_n, \bar{y}_n) = (t_\infty, x_\infty, y_\infty, \bar{x}_\infty, \bar{y}_\infty)$$

for some $(t_\infty, x_\infty, y_\infty, \bar{x}_\infty, \bar{y}_\infty) \in K$. Moreover, we note that $\psi, M_n \geq 0$ implies

$$\begin{aligned} (3.4) \quad &\frac{n}{2} \left(|x_n - \bar{x}_n|^2 + |y_n - \bar{y}_n|^2 \right) \\ &= u(t_n, x_n, y_n) - v(t_n, \bar{x}_n, \bar{y}_n) - \delta\psi(t_n, x_n, y_n) - \delta\psi(t_n, \bar{x}_n, \bar{y}_n) - M_n \\ &\leq \sup_{(t,x,y,\bar{x},\bar{y}) \in K} [u(t, x, y) - v(t, \bar{x}, \bar{y})] < \infty. \end{aligned}$$

For the finiteness of the last expression, we have used that u and $-v$ are upper semicontinuous and finite-valued on the compact set K . Since (3.4) is finite and does not depend on n , it follows

that $(\bar{x}_\infty, \bar{y}_\infty) = (x_\infty, y_\infty)$. From this, using that $M_n \geq M$ and then upper semicontinuity of both u and $-v$ and continuity of ψ , it follows that

$$\begin{aligned}
 0 &\leq \limsup_{n \rightarrow \infty} \frac{n}{2} (|x_n - \bar{x}_n|^2 + |y_n - \bar{y}_n|^2) \\
 &= \limsup_{n \rightarrow \infty} \left[u(t_n, x_n, y_n) - v(t_n, \bar{x}_n, \bar{y}_n) - \delta\psi(t_n, x_n, y_n) - \delta\psi(t_n, \bar{x}_n, \bar{y}_n) - M_n \right] \\
 &\leq \limsup_{n \rightarrow \infty} \left[u(t_n, x_n, y_n) - v(t_n, \bar{x}_n, \bar{y}_n) - \delta\psi(t_n, x_n, y_n) - \delta\psi(t_n, \bar{x}_n, \bar{y}_n) - M \right] \\
 &\leq u(t_\infty, x_\infty, y_\infty) - v(t_\infty, x_\infty, y_\infty) - \delta\psi(t_\infty, x_\infty, y_\infty) - \delta\psi(t_\infty, x_\infty, y_\infty) - M \\
 &= \phi(t_\infty, x_\infty, y_\infty) - M \leq 0.
 \end{aligned}$$

In particular, all inequalities must actually be equalities and the limsup can be replaced by a proper limit. We have therefore argued that

$$\lim_{n \rightarrow \infty} \frac{n}{2} (|x_n - \bar{x}_n|^2 + |y_n - \bar{y}_n|^2) = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} M_n = M = \phi(t_\infty, x_\infty, y_\infty)$$

as well as

$$\lim_{n \rightarrow \infty} u(t_n, x_n, y_n) = u(t_\infty, x_\infty, y_\infty) \quad \text{and} \quad \lim_{n \rightarrow \infty} v(t, \bar{x}, \bar{y}) = v(t_\infty, x_\infty, y_\infty).$$

Step 4: Application of Ishii's lemma. Let us show that $(t_\infty, x_\infty, y_\infty)$ is not located on the boundary of the state space. As K is compact, we cannot have $x_\infty = 0$ nor $y_\infty = 0$. Moreover, if $t_\infty = T$, we use $\psi \geq 0$ and the assumption that $u(T, \cdot) \leq v(T, \cdot)$ on \mathcal{O} to arrive at the contradiction

$$0 < M = \phi(T, x_\infty, y_\infty) \leq u(T, x_\infty, y_\infty) - v(T, x_\infty, y_\infty) \leq 0.$$

Thus, as $t_\infty < T$, it follows that $t_n < T$ for all $n \in \mathbb{N}$ large enough and hence without loss of generality for all $n \in \mathbb{N}$. We can therefore apply Theorem V.6.1 of [17] (Ishii's lemma) to obtain the existence of $s^n, \bar{s}^n \in \mathbb{R}$ and $M^n, \bar{M}^n \in \mathbb{R}^{2 \times 2}$ symmetric such that $s^n - \bar{s}^n = 0$ and

$$(3.5) \quad \begin{pmatrix} M^n & 0 \\ 0 & -\bar{M}^n \end{pmatrix} \leq 3n \begin{pmatrix} \text{I} & -\text{I} \\ -\text{I} & \text{I} \end{pmatrix} \quad \text{with} \quad \text{I} := \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$

and such that³

$$(q^n, p^n, P^n) \in \overline{\mathcal{F}}^{(1,2),+} u(t_n, x_n, y_n) \quad \text{and} \quad (\bar{q}^n, \bar{p}^n, \bar{P}^n) \in \overline{\mathcal{F}}^{(1,2),-} v(t_n, \bar{x}_n, \bar{y}_n),$$

where

$$(q^n, p^n, P^n) := \left(s^n + \delta\psi_t(t_n, x_n, y_n), n \begin{pmatrix} x_n - \bar{x}_n \\ y_n - \bar{y}_n \end{pmatrix} + \delta\text{D}\psi(t_n, x_n, y_n), M^n + \delta\text{D}^2\psi(t_n, x_n, y_n) \right),$$

³Here, $\overline{\mathcal{F}}^{(1,2),+} w(t, x, y)$ and $\overline{\mathcal{F}}^{(1,2),-} w(t, x, y)$ denote the closures of the second-order parabolic super- and subjets of a function $w : [0, T] \times \mathcal{O} \rightarrow \mathbb{R}$ at (t, x, y) , respectively.

$$(\bar{q}^n, \bar{p}^n, \bar{P}^n) := \left(\bar{s}^n - \delta\psi_t(t_n, \bar{x}_n, \bar{y}_n), n \begin{pmatrix} x_n - \bar{x}_n \\ y_n - \bar{y}_n \end{pmatrix} - \delta D\psi(t_n, \bar{x}_n, \bar{y}_n), \bar{M}^n - \delta D^2\psi(t_n, \bar{x}_n, \bar{y}_n) \right).$$

Step 5: The contradiction. As u and v are viscosity sub- and supersolutions, it follows that

$$-q^n - H(t_n, x_n, y_n, p^n, P^n) \leq 0 \quad \text{and} \quad -\bar{q}^n - H(t_n, \bar{x}_n, \bar{y}_n, \bar{p}^n, \bar{P}^n) \geq 0.$$

Using the elementary inequality

$$\sup\{a + b\} - \sup\{c - d\} \leq \sup\{a - c\} + \sup\{b\} + \sup\{d\},$$

it follows that

$$(3.6) \quad 0 \leq q^n - \bar{q}^n + H(t_n, x_n, y_n, p^n, P^n) - H(t_n, \bar{x}_n, \bar{y}_n, \bar{p}^n, \bar{P}^n) \\ \leq \sup_{\pi \in A} \left\{ (r + \sigma\theta\pi)n|x_n - \bar{x}_n|^2 + n(x_n - \bar{x}_n)(y_n - \bar{y}_n) + \mu_C(t_n)n|y_n - \bar{y}_n|^2 \right. \\ \left. + \frac{1}{2} \text{tr} \left[\Sigma_\pi(t_n, x_n, y_n) \Sigma_\pi(t_n, x_n, y_n)^\top M^n - \Sigma_\pi(t_n, \bar{x}_n, \bar{y}_n) \Sigma_\pi(t_n, \bar{x}_n, \bar{y}_n)^\top \bar{M}^n \right] \right\} \\ + \delta \left(\psi_t(t_n, \bar{x}_n, \bar{y}_n) + \sup_{\pi \in A} \{ \mathbf{L}^\pi \psi(t_n, \bar{x}_n, \bar{y}_n) \} \right) \\ + \delta \left(\psi_t(t_n, x_n, y_n) + \sup_{\pi \in A} \{ \mathbf{L}^\pi \psi(t_n, x_n, y_n) \} \right).$$

Using (3.5), a standard estimate shows that

$$\frac{1}{2} \text{tr} \left[\Sigma_\pi(t_n, x_n, y_n) \Sigma_\pi(t_n, x_n, y_n)^\top M^n - \Sigma_\pi(t_n, \bar{x}_n, \bar{y}_n) \Sigma_\pi(t_n, \bar{x}_n, \bar{y}_n)^\top \bar{M}^n \right] \\ \leq \frac{3n}{2} \left[\pi^2 \sigma^2 |x_n - \bar{x}_n|^2 + \sigma_C(t_n)^2 |y_n - \bar{y}_n|^2 \right].$$

But then this and

$$n(x_n - \bar{x}_n)(y_n - \bar{y}_n) \leq n \max\{|x_n - \bar{x}_n|^2, |y_n - \bar{y}_n|^2\} \leq n[|x_n - \bar{x}_n|^2 + |y_n - \bar{y}_n|^2]$$

allows us to continue to estimate (3.6) as follows:

$$0 \leq \sup_{\pi \in A} \left\{ (|r| + \sigma|\theta||\pi|)n|x_n - \bar{x}_n|^2 + n|x_n - \bar{x}_n|^2 + n|y_n - \bar{y}_n|^2 \right. \\ \left. + |\mu_C(t_n)|n|y_n - \bar{y}_n|^2 + \frac{3}{2}\sigma^2\pi^2n|x_n - \bar{x}_n|^2 + \frac{3}{2}\sigma_C(t_n)^2n|y_n - \bar{y}_n|^2 \right\} \\ + \delta \left(\psi_t(t_n, \bar{x}_n, \bar{y}_n) + \sup_{\pi \in A} \{ \mathbf{L}^\pi \psi(t_n, \bar{x}_n, \bar{y}_n) \} + \psi_t(t_n, x_n, y_n) + \sup_{\pi \in A} \{ \mathbf{L}^\pi \psi(t_n, x_n, y_n) \} \right) \\ \leq nC_x|x_n - \bar{x}_n|^2 + nC_y|y_n - \bar{y}_n|^2 \\ + \delta \left(\psi_t(t_n, \bar{x}_n, \bar{y}_n) + \sup_{\pi \in A} \{ \mathbf{L}^\pi \psi(t_n, \bar{x}_n, \bar{y}_n) \} + \psi_t(t_n, x_n, y_n) + \sup_{\pi \in A} \{ \mathbf{L}^\pi \psi(t_n, x_n, y_n) \} \right),$$

where

$$C_x := 1 + |r| + \sigma|\theta| \sup_{\pi \in A} |\pi| + \frac{3}{2}\sigma^2 \sup_{\pi \in A} |\pi|^2 \quad \text{and} \quad C_y := 1 + \sup_{t \in [0, T]} |\mu_C(t)| + \frac{3}{2} \sup_{t \in [0, T]} |\sigma_C(t)|^2.$$

Sending $n \rightarrow \infty$ and using that ψ is a strict supersolution of the HJB equation therefore yields

$$0 \leq 2\delta \left(\psi_t(t_\infty, x_\infty, y_\infty) + \sup_{\pi \in A} \{ \mathbf{L}^\pi \psi(t_\infty, x_\infty, y_\infty) \} \right) < 0,$$

which is the desired contradiction and hence concludes this proof. \square

With this comparison principle at hand, the stochastic Perron's method [1] immediately implies that the value function \mathcal{V} is the unique continuous viscosity solution of the HJB equation.

Corollary 3.5 (Viscosity characterization). *The value function \mathcal{V} is the unique viscosity solution of the HJB equation (3.1) in the class of continuous functions satisfying the terminal condition*

$$\mathcal{V}(T, x, y) = U_\gamma(x), \quad (x, y) \in \mathcal{O},$$

and the growth condition

$$U_\gamma(x)e^{\gamma K(T-t)} \leq \mathcal{V}(t, x, y) \leq U_\gamma(x + y\varphi_0(t))e^{K(T-t)}, \quad (t, x, y) \in [0, T] \times \mathcal{O}.$$

3.3 Regularity of the value function

We now give sufficient conditions which guarantee that the value function is even a classical solution of the HJB equation. Related but different results can be found in [16] and [15]. The main obstacle to establishing regularity of the value function is the lack of uniform ellipticity of the HJB equation. However, the homotheticity property allows us to consider the transformation

$$(3.7) \quad \mathcal{V}(t, x, y) = y^\gamma e^{-\beta t} w(t, \log x/y), \quad \text{or, equivalently,} \quad w(t, \zeta) := e^{\beta t} \mathcal{V}(t, e^\zeta, 1),$$

where $\zeta := \log x/y$ and $\beta \in \mathbb{R}$. It turns out that w solves a reduced-form HJB equation of the form

$$-w_t - \sup_{\pi \in A} \left\{ a(\pi, t) w_{\zeta\zeta} + b(\pi, t, \zeta, w, w_\zeta) \right\} = 0, \quad (t, \zeta) \in (0, T) \times \mathbb{R},$$

which is uniformly elliptic and admits a classical solution provided that μ_C and σ_C are continuously differentiable. Moreover, observe that by passing to w we have reduced the dimension of the state space by one, which is advantageous for numerical computations.

Theorem 3.6 (Regularity). *Assume that μ_C and σ_C are continuously differentiable. Then $\mathcal{V} \in C^{1,2}((0, T) \times \mathcal{O})$. In particular, \mathcal{V} is a classical solution of the HJB equation.*

Proof. Step 1: The transformed HJB equation. Let us consider the equation

$$(3.8) \quad -w_t - \sup_{\pi \in A} \left\{ a(\pi, t) w_{\zeta\zeta} + b(\pi, t, \zeta, w, w_\zeta) \right\} = 0, \quad (t, \zeta) \in (0, T) \times \mathbb{R},$$

where

$$a : A \times [0, T] \rightarrow \mathbb{R} \quad \text{and} \quad b : A \times [0, T] \times \mathbb{R} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$$

are given by

$$\begin{aligned} a(\pi, t) &:= \frac{1}{2} \left[(\pi\sigma - \rho\sigma_C(t))^2 + (1 - \rho^2)\sigma_C(t)^2 \right], \\ b(\pi, t, \zeta, v, p) &:= \left(r - \mu_C(t) + \frac{1}{2}(1 - 2\gamma)\sigma_C(t)^2 + \pi\sigma\theta - \frac{1}{2}\pi^2\sigma^2 + \gamma\rho\sigma\sigma_C(t)\pi + e^{-\zeta} \right) p \\ &\quad + \left(\gamma\mu_C(t) - \gamma\frac{1}{2}(1 - \gamma)\sigma_C(t)^2 + \beta \right) v, \end{aligned}$$

for $\pi \in A$, $t \in [0, T]$, $\zeta, v, p \in \mathbb{R}$, and where the parameter $\beta \in \mathbb{R}$ is chosen such that

$$(3.9) \quad \beta < \inf_{t \in [0, T]} \left[-\gamma\mu_C(t) + \frac{1}{2}\gamma(1 - \gamma)\sigma_C(t)^2 \right].$$

Formally, Equation (3.8) arises if we consider the transformation

$$w : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}, \quad (t, \zeta) \mapsto w(t, \zeta) := e^{\beta t} \mathcal{V}(t, e^\zeta, 1).$$

Now fix $N \in \mathbb{N}$. Then we claim that Equation (3.8) admits a solution $w^N : (0, T] \times [-N, N] \rightarrow \mathbb{R}$ with $w^N \in C^{1,2}((0, T) \times (-N, N))$ satisfying the boundary and terminal conditions

$$(3.10) \quad w^N(t, \zeta) = e^{\beta t} \mathcal{V}(t, e^\zeta, 1), \quad (t, \zeta) \in \left(\{T\} \times [-N, N] \right) \cup \left((0, T) \times \{-N, N\} \right).$$

Once this is established, we can define a continuous function

$$v^N : (0, T] \times \overline{\mathcal{O}(N)} \rightarrow \mathbb{R}, \quad (t, x, y) \mapsto v^N(t, x, y) := y^\gamma e^{-\beta t} w^N(t, \log x/y),$$

where $\overline{\mathcal{O}(N)}$ denotes the closure of the set

$$\mathcal{O}(N) := \{(x, y) \in \mathcal{O} : \log x/y \in (-N, N)\}.$$

Observe that, by the terminal/boundary condition (3.10) and the homogeneity of \mathcal{V} , we have

$$v^N(t, x, y) = y^\gamma e^{-\beta t} w^N(t, \log x/y) = y^\gamma \mathcal{V}(t, x/y, 1) = \mathcal{V}(t, x, y)$$

whenever $t = T$ or $(x, y) \in \overline{\mathcal{O}(N)} \setminus \mathcal{O}(N)$. Moreover, since w^N solves (3.8) on $(0, T) \times (-N, N)$, a straightforward calculation shows that v^N solves the original HJB equation, i.e.

$$-v_t^N(t, x, y) - \sup_{\pi \in A} \{ \mathbf{L}^\pi v^N(t, x, y) \} = 0, \quad (t, x, y) \in (0, T) \times (-N, N).$$

But then v^N also satisfies this equation in the sense of viscosity solutions and hence $v^N = \mathcal{V}$ on $(0, T] \times \overline{\mathcal{O}(N)}$ by the uniqueness result Theorem V.8.1 in [17]. In particular, we find that $\mathcal{V} \in C^{1,2,2}((0, T) \times \mathcal{O}(N))$ for each $N \in \mathbb{N}$ and we conclude by sending $N \rightarrow \infty$.

Step 2: We are left with showing that Equation (3.8) admits a classical solution w^N on $(0, T) \times (-N, N)$ satisfying the boundary/terminal condition (3.10). For this, it is sufficient to verify the conditions in Theorem A.8 in [16]; see also Theorem 3 in Section 6.4 of [24] for the original result.

- (i) For every $\pi \in A$, it is clear that $a(\pi, \cdot)$ and $b(\pi, \cdot)$ are continuously differentiable and for each pair $(\pi, t) \in A \times [0, T]$ the function $b(\pi, t, \cdot)$ is twice continuously differentiable. Indeed, the non-zero derivatives are given by

$$\begin{aligned} a_t &= \sigma_C(t)\dot{\sigma}_C(t) - \rho\sigma\dot{\sigma}_C(t)\pi, \\ b_t &= (-\dot{\mu}_C(t) + (1 - 2\gamma)\sigma_C(t)\dot{\sigma}_C(t) + \gamma\rho\sigma\dot{\sigma}_C(t)\pi)p + (\gamma\dot{\mu}_C(t) - \gamma(1 - \gamma)\sigma_C(t)\dot{\sigma}_C(t) + \beta)v, \\ b_v &= \gamma\mu_C(t) - \frac{1}{2}\gamma(1 - \gamma)\sigma_C(t)^2 + \beta, \\ b_p &= r - \mu_C(t) + \frac{1}{2}(1 - 2\gamma)\sigma_C(t)^2 + \sigma\theta\pi - \frac{1}{2}\sigma^2\pi^2 + \gamma\rho\sigma\sigma_C(t)\pi + e^{-\zeta}, \\ b_\zeta &= -b_{\zeta\zeta} = -e^{-\zeta}p, \\ b_{\zeta p} &= b_{p\zeta} = e^{-\zeta}. \end{aligned}$$

From this, we also see that a_t, b_t as well as all second-order derivatives of b with respect to ζ and p are bounded on

$$S_M := \{(\pi, t, \zeta, v, p) \in A \times (0, T) \times (-N, N) \times \mathbb{R} \times \mathbb{R} : |v| + |p| \leq M\} \quad \text{for all } M \in \mathbb{N}.$$

- (ii) a is uniformly elliptic on $A \times [0, T]$, i.e.

$$0 < \frac{1}{2}(1 - \rho) \inf_{t \in [0, T]} |\sigma_C(t)|^2 \leq a(\pi, t) \leq \sup_{(t, \pi) \in [0, T] \times A} |a(\pi, t)| < \infty, \quad (\pi, t) \in A \times [0, T].$$

- (iii) For all $(\pi, t, \zeta, v, p) \in A \times (0, T) \times (-N, N) \times \mathbb{R} \times \mathbb{R}$, it holds that

$$(1 + |p|)|a_p(\pi, t)| + |a_v(\pi, t)| + \frac{|a_\zeta(\pi, t)|}{1 + |p|} = 0.$$

Similarly, setting

$$\begin{aligned} C_1 &:= \sup_{(\pi, t, \zeta) \in A \times [0, T] \times [-N, N]} \left| r - \mu_C(t) + \frac{1}{2}(1 - 2\gamma)\sigma_C(t)^2 + \sigma\theta\pi \right. \\ &\quad \left. - \frac{1}{2}\sigma^2\pi^2 + \gamma\rho\sigma\sigma_C(t)\pi + e^{-\zeta} \right|, \\ C_2 &:= \sup_{t \in [0, T]} \left| \gamma\mu_C(t) - \frac{1}{2}\gamma(1 - \gamma)\sigma_C(t)^2 + \beta \right|, \end{aligned}$$

it follows that

$$(1 + |p|)|b_p(\pi, t, \zeta, v, p)| + |b(\pi, t, \zeta, v, p)| + |b_v(\pi, t, \zeta, v, p)| + \frac{|b_\zeta(\pi, t, \zeta, v, p)|}{(1 + |p|)}$$

$$\leq (1 + |p|)C_1 + |p|C_1 + |v|C_2 + C_2 + \frac{|p|}{1 + |p|}e^N \leq h(v)(1 + |p|^2),$$

where $h : \mathbb{R} \rightarrow [0, \infty)$ is the continuous function given by

$$h(v) := |v|C_2 + 3C_1 + C_2 + e^N, \quad v \in \mathbb{R}.$$

(iv) Fix $Q > 0$. Then, since β is chosen to satisfy (3.9), there exists a constant $\delta > 0$ such that

$$\begin{aligned} b(\pi, t, \zeta, -Q, 0) &= - \left(\gamma\mu_C(t) - \frac{1}{2}\gamma(1 - \gamma)\sigma_C(t)^2 + \beta \right) Q \geq \delta > 0, \\ b(\pi, t, \zeta, Q, 0) &= \left(\gamma\mu_C(t) - \frac{1}{2}\gamma(1 - \gamma)\sigma_C(t)^2 + \beta \right) Q \leq -\delta < 0, \end{aligned}$$

for all $(\pi, t, \zeta) \in A \times (0, T) \times (-N, N)$.

Under conditions (i) to (iv), Theorem 3 in Section 6.4 of [24] is applicable. This yields the existence of w^N and the proof is complete. \square

4 The optimal strategy and its asymptotic behavior

In this section, we construct an optimal strategy and study its properties as the ratio of wealth to endowment becomes large.

4.1 Existence and characterization of the optimal strategy

A candidate optimal strategy is readily found by computing the maximizer in the HJB equation. Indeed, a straightforward calculation shows that the unique maximizer of the Hamiltonian H is given by

$$(4.1) \quad \hat{\pi}^*(t, x, y) := \left(\left[-\frac{\theta}{\sigma} \frac{\mathcal{V}_x(t, x, y)}{x\mathcal{V}_{xx}(t, x, y)} - \frac{\rho\sigma_C(t)}{\sigma} \frac{y\mathcal{V}_{xy}(t, x, y)}{x\mathcal{V}_{xx}(t, x, y)} \right] \vee \underline{\pi} \right) \wedge \bar{\pi}, \quad (t, x, y) \in [0, T] \times \mathcal{O}.$$

For any initial configuration (t, x, y) , a candidate optimal strategy $\pi^* \in \mathcal{A}$ is defined on (t, T) by⁴

$$\pi_s^* := \hat{\pi}^*(s, A_s^*, c_s), \quad s \in (t, T)$$

where $c := c^{t, y}$ and A^* is the solution of

$$(4.2) \quad dA_s^* = \left[(r + \sigma\theta\hat{\pi}(s, A_s^*, c_s))A_s^* + c_s \right] ds + \sigma\hat{\pi}(s, A_s^*, c_s)A_s^* dW_s^1, \quad s \in [t, T], \quad A_t^* = x.$$

⁴The strategy π^* should of course be extended to an A -valued process on $[0, T]$. Note, however, that the function $\hat{\pi}^*$ may not be well defined for $t \in \{0, T\}$. An argument as in Remark 3.1 (iv) in [35] shows that $\pi^* \mathbb{1}_{[0, t]}$ can without loss of generality be taken independent of $\pi^* \mathbb{1}_{[t, T]}$, e.g. equal to a constant value.

We note, however, that it is unclear if the drift and diffusion coefficients

$$(x, y) \mapsto (r + \sigma\theta\hat{\pi}(t, x, y))x + y \quad \text{and} \quad (x, y) \mapsto \sigma\hat{\pi}(t, x, y)x$$

satisfy the necessary regularity to ensure the existence of a strong solution of (4.2). We therefore follow a different route: We first argue that for each initial configuration (t, x, y) , an optimal strategy exists and then show that it can be represented in feedback form via the function $\hat{\pi}^*$.

Proposition 4.1 (Existence of optimizers). *Let $(t, x, y) \in [0, T) \times \mathcal{O}$. Then there exists an optimal trading strategy $\pi^* = \pi_{t,x,y}^* \in \mathcal{A}$, i.e.*

$$\mathbb{E}[U_\gamma(A_T^{\pi^*})] = \sup_{\pi \in \mathcal{A}} \mathbb{E}[U_\gamma(A_T^\pi)] = \mathcal{V}(t, x, y).$$

Proof. We employ a classical Komlos-type argument and show that suitable forward-convex combinations of a maximizing sequence for \mathcal{V} converge to an optimal strategy. Denote by $\{\hat{\pi}^n\}_{n \in \mathbb{N}} \subset \mathcal{A}$ such a maximizing sequence for $\mathcal{V}(t, x, y)$, i.e. a sequence of admissible trading strategies with

$$\mathcal{V}(t, x, y) = \lim_{n \rightarrow \infty} \mathbb{E}[U_\gamma(\hat{A}_T^n)], \quad \text{where } \hat{A}^n := A^{\hat{\pi}^n, t, x, y}.$$

For each $n \in \mathbb{N}$, let us denote by $\hat{S}^n := \hat{\pi}^n \hat{A}^n$ the wealth invested in the stock under the strategy $\hat{\pi}^n$. We may extend \hat{S}^n to a process defined on $[0, T]$ by setting $\hat{S}_s^n := \hat{\pi}_s^n x$ for all $s \in [0, t]$. Now observe that, by uniform boundedness of $\{\hat{\pi}^n\}_{n \in \mathbb{N}}$ and the a priori estimate (2.4) in Lemma 2.1, it holds that

$$\sup_{n \in \mathbb{N}} \mathbb{E} \left[\int_0^T |\hat{S}_s^n|^2 ds \right] < \infty,$$

i.e. the sequence $\{\hat{S}^n\}_{n \in \mathbb{N}}$ is bounded in the Hilbert space of progressively measurable and square-integrable processes. We can therefore apply Theorem 15.1.2 in [11] to find $S^* = \{S_s^*\}_{s \in [0, T]}$ and a sequence $\{S^n\}_{n \in \mathbb{N}}$ with $S^n \in \text{conv}\{\hat{S}^n, \hat{S}^{n+1}, \dots\}$ such that

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\int_0^T |S_s^* - S_s^n|^2 ds \right] = 0.$$

Next, for each $n \in \mathbb{N}$, let us introduce processes $A^n = \{A_s^n\}_{s \in [0, T]}$ and $\pi^n = \{\pi_s^n\}_{s \in [0, T]}$ with $\pi^n := S^n/A^n$ and A^n given as the unique solution of

$$dA_s^n = [rA_s^n + \sigma\theta S_s^n + c_s^{t,y}] ds + \sigma S_s^n dW_s^1, \quad s \in [t, T],$$

with $A^n = x$ on $[0, t]$. Observe that A^n is the wealth process corresponding to the trading strategy π^n . Moreover, since $S^n \in \text{conv}\{\hat{S}^n, \hat{S}^{n+1}, \dots\}$, there exist $N_n \in \mathbb{N}$, indices $k_1^n, \dots, k_{N_n}^n \in \{n, n+1, \dots\}$ and convex weights $\lambda_1^n, \dots, \lambda_{N_n}^n$ such that $S^n = \sum_{i=1}^{N_n} \lambda_i^n \hat{S}^{k_i^n}$. With this, we see that

$$d \sum_{i=1}^{N_n} \lambda_i^n \hat{A}_s^{k_i^n} = \sum_{i=1}^{N_n} \lambda_i^n \left[[r\hat{A}_s^{k_i^n} + \sigma\theta \hat{S}_s^{k_i^n} + c_s^{t,y}] ds + \sigma \hat{S}_s^{k_i^n} dW_s^1 \right]$$

$$= \left[r \sum_{i=1}^{N_n} \lambda_i^n \hat{A}_s^{k_i^n} + \sigma \theta S_s^n + c_s^{t,y} \right] ds + \sigma S_s^n dW_s^1, \quad s \in [t, T],$$

from which we conclude $A^n = \sum_{i=1}^{N_n} \lambda_i^n \hat{A}_s^{k_i^n}$ by uniqueness of solutions of linear SDEs. But then

$$\pi^n = \frac{S^n}{A^n} = \sum_{k=1}^{N_n} \frac{\lambda_i^n \hat{A}_s^{k_i^n}}{A^n} \hat{\pi}^{k_i^n},$$

i.e. π^n is a (dynamic) convex combination of $\pi^{k_1^n}, \dots, \pi^{k_{N_n}^n}$ and therefore A -valued and admissible.

Next, consider the wealth process $A^* = \{A_s^*\}_{s \in [0, T]}$ given by

$$dA_s^* = [rA_s^* + \sigma \theta S_s^* + c_s^{t,y}] ds + \sigma S_s^* dW_s^1, \quad s \in [t, T],$$

with $A^* = x$ on $[0, t]$ and observe that, using the dynamics of A^n and A^* , Jensen's inequality, and the Itô isometry, there exists a constant $C > 0$ such that

$$\begin{aligned} \mathbb{E}[|A_s^* - A_s^n|^2] &\leq C \mathbb{E} \left[\int_t^s |A_u^* - A_u^n|^2 du \right] + C \mathbb{E} \left[\int_t^s |S_u^* - S_u^n|^2 du \right] \\ &\leq C \int_t^s \mathbb{E}[|A_u^* - A_u^n|^2] du + C \mathbb{E} \left[\int_0^T |S_u^* - S_u^n|^2 du \right], \quad s \in [t, T]. \end{aligned}$$

Gronwall's inequality therefore yields

$$\lim_{n \rightarrow \infty} \mathbb{E}[|A_s^* - A_s^n|^2] \leq \lim_{n \rightarrow \infty} C \mathbb{E} \left[\int_0^T |S_u^* - S_u^n|^2 du \right] e^{C(T-t)} = 0, \quad s \in [t, T].$$

Setting $\pi^* := S^*/A^*$, using the convergence of (S^n, A^n) to (S^*, A^*) , it follows that π^* is A -valued $dt \otimes \mathbb{P}$ almost everywhere (hence without loss of generality everywhere after possibly redefining π^* on a nullset) and hence admissible. Clearly, A^* is the wealth process corresponding to π^* . Moreover, since $U_\gamma(A_T^n)^2 = |A_T^n|^{2\gamma}/|\gamma|^2$, it follows from Lemma 2.1 that $\{U_\gamma(A_T^n)\}_{n \in \mathbb{N}}$ is bounded in the space of square-integrable random variables and thus uniformly integrable. With this and using the concavity of U_γ , we hence conclude that

$$\mathbb{E}[U_\gamma(A_T^*)] = \lim_{n \rightarrow \infty} \mathbb{E}[U_\gamma(A_T^n)] \geq \lim_{n \rightarrow \infty} \sum_{i=1}^{N_n} \lambda_i^n \mathbb{E}[U_\gamma(\hat{A}_T^{k_i^n})] = \mathcal{V}(t, x, y). \quad \square$$

Next, let us proceed to show that any optimal strategy must necessarily be given in feedback form via the function $\hat{\pi}^*$ defined in (4.1).

Theorem 4.2 (Optimal strategy in feedback form). *Fix $(t, x, y) \in [0, T] \times \mathcal{O}$, let $\pi \in \mathcal{A}$ be an arbitrary strategy, and suppose that $\mathcal{V} \in C^{1,2}((0, T) \times \mathcal{O})$. Then π is optimal if and only if*

$$(4.3) \quad \pi_s(\omega) = \hat{\pi}^*(s, A_s(\omega), c_s(\omega)) \quad \text{for } \mathbb{P} \otimes ds\text{-almost every } (\omega, s) \in \Omega \times ([t, T] \cap (0, T)),$$

where $A := A^{\pi, t, x, y}$ and $c := c^{t, y}$.

Proof. Step 1: Suppose that $\pi \in \mathcal{A}$ is an arbitrary strategy satisfying (4.3). Using that \mathcal{V} solves the HJB equation and $\hat{\pi}^*$ is a pointwise maximizer of the supremum in the HJB equation, an application of Itô's formula shows that

$$\begin{aligned} \mathcal{V}(s, A_s, c_s) &= \mathcal{V}(t, x, y) + \int_t^s \mathcal{V}_t(u, A_u, c_u) + \mathbf{L}^{\pi_u} \mathcal{V}(u, A_u, c_u) du \\ &\quad + \int_t^s \sigma \pi_u A_u \mathcal{V}_x(u, A_u, c_u) dW_u^1 + \int_t^s \sigma_C(u) c_u \mathcal{V}_y(u, A_u, c_u) dW_u^C \\ &= \mathcal{V}(t, x, y) + \int_t^s \sigma \pi_u A_u \mathcal{V}_x(u, A_u, c_u) dW_u^1 + \int_t^s \sigma_C(u) c_u \mathcal{V}_y(u, A_u, c_u) dW_u^C \end{aligned}$$

for all $s \in [t, T]$, i.e. $v(\cdot, A, c)$ is a local martingale. But since by (2.12) and Lemma 2.1

$$(4.4) \quad \mathbf{E} \left[\sup_{s \in [t, T]} |\mathcal{V}(s, A_s, c_s)| \right] \leq C \mathbf{E} \left[1 + \sup_{s \in [t, T]} |A_s|^{-\gamma^-} + \sup_{s \in [t, T]} |A_s|^{\gamma^+} + \sup_{s \in [t, T]} |c_s|^{\gamma^+} \right] < \infty,$$

it follows that $\mathcal{V}(\cdot, A, c)$ is an honest martingale and hence

$$\mathcal{V}(t, x, y) = \mathbf{E}[\mathcal{V}(T, A_T, c_T)] = \mathbf{E}[U_\gamma(A_T)],$$

i.e. π is optimal.

Step 2: Suppose that $\pi \in \mathcal{A}$ is optimal, i.e.

$$(4.5) \quad \mathcal{V}(t, x, y) = \mathbf{E}[U_\gamma(A_T)] = \mathbf{E}[\mathcal{V}(T, A_T, c_T)].$$

An application of Itô's formula and the fact that \mathcal{V} satisfies the HJB equation implies

$$\begin{aligned} \mathcal{V}(s, A_s, c_s) &= \mathcal{V}(t, x, y) + \int_t^s \mathcal{V}_t(u, A_u, c_u) + \mathbf{L}^{\pi_u} \mathcal{V}(u, A_u, c_u) du \\ &\quad + \int_t^s \sigma \pi_u A_u \mathcal{V}_x(u, A_u, c_u) dW_u^1 + \int_t^s \sigma_C(u) c_u \mathcal{V}_y(u, A_u, c_u) dW_u^C \\ (4.6) \quad &\leq \mathcal{V}(t, x, y) + \int_t^s \sigma \pi_u A_u \mathcal{V}_x(u, A_u, c_u) dW_u^1 + \int_t^s \sigma_C(u) c_u \mathcal{V}_y(u, A_u, c_u) dW_u^C \end{aligned}$$

for all $s \in [t, T]$. But as (4.4) is also valid in this situation, it follows that $\mathcal{V}(\cdot, A, c)$ is a supermartingale and thus by (4.5) an honest martingale. Thus we must have equality in (4.6), i.e.

$$\mathcal{V}_t(s, A_s, c_s) + \mathbf{L}^{\pi_s} \mathcal{V}(s, A_s, c_s) = 0 = \mathcal{V}_t(s, A_s, c_s) + \sup_{\pi \in \mathcal{A}} \{ \mathbf{L}^\pi \mathcal{V}(s, A_s, c_s) \}$$

$\mathbb{P} \otimes ds$ -almost everywhere on $\Omega \times ([t, T] \cap (0, T))$. Now the supremum on the right hand side has a unique maximizer given by the function $\hat{\pi}^*$ and thus

$$\pi_s(\omega) = \hat{\pi}^*(s, A_s(\omega), c_s(\omega)) \quad \text{for } \mathbb{P} \otimes ds\text{-almost every } (\omega, s) \in \Omega \times ([t, T] \cap (0, T))$$

and the proof is complete. \square

4.2 Asymptotic behavior of the optimal strategy

In this section, we examine the asymptotic behavior of the value function and the optimal policy as the initial capital converges to infinity. We will see that for this model a sort of “turnpike property” holds (see e.g. [36]) in the sense that for $x \gg y$ (more precisely, as $x/y \rightarrow \infty$ becomes large) the optimal policy approaches the Merton fraction π_M of investing a constant proportion of wealth in the risky asset.

To make this statement precise, we subsequently write

$$f(z) \underset{z \rightarrow \infty}{\sim} g(z) \quad \text{if and only if} \quad \lim_{z \rightarrow \infty} \frac{f(z)}{g(z)} = 1$$

whenever $f, g : \mathbb{R} \rightarrow \mathbb{R}$. Our aim in this section is hence to show that $\hat{\pi}^*(t, x, y) \underset{x/y \rightarrow \infty}{\sim} \pi_M$ for all $t \in (0, T)$, where $\hat{\pi}^*$ is the feedback function defined in (4.1). To prove this, we first recall the reduced value function \mathcal{U} defined in Lemma 2.6 and observe that

$$\begin{aligned} \mathcal{V}_x(t, x, y) &= y^{\gamma-1} \mathcal{U}_z(t, z), & \mathcal{V}_{xx}(t, x, y) &= y^{\gamma-2} \mathcal{U}_{zz}(t, z), \\ \mathcal{V}_{xy}(t, x, y) &= -y^{\gamma-2} [(1-\gamma) \mathcal{U}_z(t, z) + z \mathcal{U}_{zz}(t, z)]. \end{aligned}$$

From this, it follows that $\hat{\pi}^*$ can be rewritten as

$$\hat{\pi}^*(t, z) = \left(\left[-\frac{\theta}{\sigma} \frac{\mathcal{U}_z(t, z)}{z \mathcal{U}_{zz}(t, z)} + \frac{(1-\gamma)\rho\sigma_C(t)}{\sigma} \frac{\mathcal{U}_z(t, z)}{z \mathcal{U}_{zz}(t, z)} + \frac{\rho\sigma_C(t)}{\sigma} \right] \vee \underline{\pi} \right) \wedge \bar{\pi}, \quad (t, z) \in [0, T] \times (0, \infty).$$

Theorem 4.3 (Asymptotics for the optimal strategy). *For any $t \in (0, T)$, we have*

$$\begin{aligned} \mathcal{U}(t, z) &\underset{z \rightarrow \infty}{\sim} \frac{1}{\gamma} z^\gamma e^{\gamma K(T-t)}, & \mathcal{U}_z(t, z) &\underset{z \rightarrow \infty}{\sim} z^{\gamma-1} e^{\gamma K(T-t)}, \\ \mathcal{U}_{zz}(t, z) &\underset{z \rightarrow \infty}{\sim} -(1-\gamma) z^{\gamma-2} e^{\gamma K(T-t)}, \end{aligned}$$

from which it follows that

$$\hat{\pi}^*(t, z) \underset{z \rightarrow \infty}{\sim} \pi_M = \frac{\theta}{(1-\gamma)\sigma}.$$

Proof. The statements on the asymptotic behavior of \mathcal{U} and its derivatives follow from the bounds on \mathcal{V} derived in Proposition 3.2. Indeed, the bounds on \mathcal{V} and homogeneity of U_γ imply that

$$\frac{1}{\gamma} z^\gamma e^{\gamma K(T-t)} \leq \mathcal{U}(t, z) \leq \frac{1}{\gamma} (z + \varphi_0(t))^\gamma e^{\gamma K(T-t)}, \quad (t, z) \in [0, T] \times (0, \infty).$$

From this and using monotonicity and concavity of all functions involved, we see right away that

$$\begin{aligned} \mathcal{U}(t, z) &\underset{z \rightarrow \infty}{\sim} \frac{1}{\gamma} z^\gamma e^{\gamma K(T-t)}, & \mathcal{U}_z(t, z) &\underset{z \rightarrow \infty}{\sim} z^{\gamma-1} e^{\gamma K(T-t)}, \\ \mathcal{U}_{zz}(t, z) &\underset{z \rightarrow \infty}{\sim} -(1-\gamma) z^{\gamma-2} e^{\gamma K(T-t)}, \end{aligned}$$

But then

$$\lim_{z \rightarrow \infty} \frac{\mathcal{U}_z(t, z)}{z\mathcal{U}_{zz}(t, z)} = \lim_{z \rightarrow \infty} \frac{z^{\gamma-1}e^{\gamma K(T-t)}}{-(1-\gamma)z^{\gamma-1}e^{\gamma K(T-t)}} = -\frac{1}{1-\gamma},$$

and it follows that

$$\lim_{z \rightarrow \infty} \left[-\frac{\theta}{\sigma} \frac{\mathcal{U}_z(t, z)}{z\mathcal{U}_{zz}(t, z)} + \frac{(1-\gamma)\rho\sigma_C(t)}{\sigma} \frac{\mathcal{U}_z(t, z)}{z\mathcal{U}_{zz}(t, z)} + \frac{\rho\sigma_C(t)}{\sigma} \right] = \frac{\theta}{(1-\gamma)\sigma} = \pi_M.$$

Since $\pi_M \in A$, we see that $\lim_{z \rightarrow \infty} \hat{\pi}^*(t, z) = \pi_M$ and the proof is complete. □

5 Conclusion

We consider an optimal asset allocation problem in an incomplete market, where exogenous stochastic endowments flow continuously into the portfolio according to a time-inhomogeneous geometric Brownian motion. We analyze the viscosity solution of the HJB PDE, reduce its dimension, and prove that the optimal strategy can be recovered from the optimal policy of a reduced problem. We are also able to describe the asymptotic behavior of the value function, and the strategy when the initial wealth goes to infinity.

References

- [1] E. BAYRAKTAR AND M. SÎRBU, *Stochastic Perron's method for Hamilton-Jacobi-Bellman equations*, SIAM Journal on Control and Optimization, 51 (2013), pp. 4274–4294.
- [2] C. BELAK, O. MENKENS, AND J. SASS, *On the uniqueness of unbounded viscosity solutions arising in an optimal terminal wealth problem with transaction costs*, SIAM Journal on Control and Optimization, 53 (2015), pp. 2878–2897.
- [3] B. BICK, H. KRAFT, AND C. MUNK, *Solving constrained consumption-investment problems by simulation of artificial market strategies*, Management Science, 59 (2013), pp. 483–503.
- [4] D. BROEDERS AND A. CHEN, *Pension regulation and the market value of pension liabilities: A contingent claims analysis using parisian options*, Journal of Banking & Finance, 34 (2010), pp. 1201–1214.
- [5] J. Y. CAMPBELL AND L. M. VICEIRA, *Strategic asset allocation: portfolio choice for long-term investors*, Oxford University Press, 2002.
- [6] J. C. COX AND C.-F. HUANG, *Optimal consumption and portfolio choices when asset prices follow a diffusion process*, Journal of Economic Theory, 49 (1989), pp. 33–83.
- [7] M. G. CRANDALL, H. ISHII, AND P.-L. LIONS, *User's guide to viscosity solutions of second order partial differential equations*, Bulletin of the American Mathematical Society, 27 (1992), pp. 1–67.

- [8] D. CUOCO, *Optimal consumption and equilibrium prices with portfolio constraints and stochastic income*, Journal of Economic Theory, 72 (1997), pp. 33–73.
- [9] J. CVITANIĆ, W. SCHACHERMAYER, AND H. WANG, *Utility maximization in incomplete markets with random endowment*, Finance and Stochastics, 5 (2001), pp. 259–272.
- [10] M. H. DAVIS, *Optimal hedging with basis risk*, in From stochastic calculus to mathematical finance, Y. Kabanov, R. Liptser, and J. Stoyanov, eds., Springer, 2006, pp. 169–187.
- [11] F. DELBAEN AND W. SCHACHERMAYER, *The Mathematics of Arbitrage*, Springer Finance, Springer-Verlag, Berlin, 2006.
- [12] D. DUFFIE, W. FLEMING, H. M. SONER, AND T. ZARIPHOUPOULOU, *Hedging in incomplete markets with HARA utility*, Journal of Economic Dynamics and Control, 21 (1997), pp. 753–782.
- [13] P. H. DYBVIG AND H. LIU, *Lifetime consumption and investment: retirement and constrained borrowing*, Journal of Economic Theory, 145 (2010), pp. 885–907.
- [14] N. EL KAROUI AND M. JEANBLANC-PICQUÉ, *Optimization of consumption with labor income*, Finance and Stochastics, 2 (1998), pp. 409–440.
- [15] S. FEDERICO, P. GASSIAT, AND F. GOZZI, *Utility maximization with current utility on the wealth: regularity of solutions to the hjb equation*, Finance and Stochastics, 19 (2015), pp. 415–448.
- [16] ———, *Impact of time illiquidity in a mixed market without full observation*, Mathematical Finance, 27 (2017), pp. 401–437.
- [17] W. H. FLEMING AND H. M. SONER, *Controlled Markov Processes and Viscosity Solutions*, Stochastic Modelling and Applied Probability, Springer, 2006.
- [18] H. HE AND N. D. PEARSON, *Consumption and portfolio policies with incomplete markets and short-sale constraints: the finite-dimensional case*, Mathematical Finance, 1 (1991), pp. 1–10.
- [19] ———, *Consumption and portfolio policies with incomplete markets and short-sale constraints: The infinite dimensional case*, Journal of Economic Theory, 54 (1991), pp. 259–304.
- [20] U. HORST, Y. HU, P. IMKELLER, A. RÉVEILLAC, AND J. ZHANG, *Forward-backward systems for expected utility maximization*, Stochastic Processes and their Applications, (2014).
- [21] Y. HU, P. IMKELLER, AND M. MÜLLER, *Utility maximization in incomplete markets*, The Annals of Applied Probability, 15 (2005), pp. 1691–1712.
- [22] J. HUGONNIER AND D. KRAMKOV, *Optimal investment with random endowments in incomplete markets*, The Annals of Applied Probability, 14 (2004), pp. 845–864.
- [23] H.-K. KOO, *Consumption and portfolio selection with labor income: a continuous time approach*, Mathematical Finance, 8 (1998), pp. 49–65.

- [24] N. V. KRYLOV, *Nonlinear Elliptic and Parabolic Equations of the Second Order*, Mathematics and its applications (Soviet series), Reidel, Dordrecht, 1987.
- [25] ———, *Controlled Diffusion Processes*, vol. 14 of Stochastic Modelling and Applied Probability, Springer-Verlag, Berlin, 2009.
- [26] R. MERTON, *Lifetime portfolio selection under uncertainty: the continuous-time case*, Review of Economic Statistics, 51 (1969), pp. 247–257.
- [27] ———, *Optimum consumption and portfolio rules in a continuous time model*, Journal of Economic Theory, 3 (1971), pp. 373–413.
- [28] O. MOSTOVYI, *Optimal investment with intermediate consumption and random endowment*, Mathematical Finance, 27 (2017), pp. 96–114.
- [29] O. MOSTOVYI AND M. SÎRBU, *Optimal investment and consumption with labor income in incomplete markets*, The Annals of Applied Probability, 30 (2020), pp. 747–787.
- [30] E. PARDOUX AND A. RĂȘCANU, *Stochastic differential equations, backward SDEs, partial differential equations*, vol. 69 of Stochastic Modelling and Applied Probability, Springer, Cham, 2014.
- [31] H. PHAM, *Continuous-time stochastic control and optimization with financial applications*, vol. 61, Springer, 2009.
- [32] S. R. PLISKA, *A stochastic calculus model of continuous trading: optimal portfolios*, Mathematics of Operational Research, 11 (1986), pp. 371–382.
- [33] M. SONER AND M. VUKELJA, *Utility maximization in an illiquid market in continuous time*, Mathematical Methods of Operations Research, 84 (2016), pp. 285–321.
- [34] S. SUNDARESAN AND F. ZAPATERO, *Valuation, optimal asset allocation and retirement incentives of pension plans*, Review of Financial Studies, 10 (1997), pp. 631–660.
- [35] N. TOUZI, *Deterministic and stochastic control, application to finance*, lecture notes, Université Paris 6, 2012. available at: <http://www.cmap.polytechnique.fr/~touzi>.
- [36] A. J. ZASLAVSKI, *Turnpike Properties in the Calculus of Variations and Optimal Control*, Nonconvex Optimization and Its Applications, Springer, 2006.
- [37] G. ZITKOVIC, *Stability of the utility maximization problem with random endowment in incomplete markets*, Mathematical Finance, 21 (2011), pp. 313–333.