

# On Pimsner Popa bases

Keshab Chandra Bakshi  
 The Institute of Mathematical Sciences  
 Taramani, Chennai 600113,  
 INDIA  
 email: keshabcb@imsc.res.in

## Abstract

In this paper we examine bases for finite index inclusion of  $II_1$  factors and connected inclusion of finite dimensional  $C^*$ -algebras. These bases behave nicely with respect to basic construction towers. As applications we have studied automorphisms of the hyperfinite  $II_1$  factor  $R$  which are ‘compatible with respect to the Jones’ tower of finite dimensional  $C^*$ -algebras’. As a further application, in both Cases we obtain a characterization, in terms of bases, of basic constructions. Finally we use these bases to describe the phenomenon of multistep basic constructions (in both the Cases).

*Mathematics Subject Classification 2010:* 46L37

*Keywords:* Subfactor; basic construction; connected inclusion; Pimsner-Popa bases.

## 1 Introduction

We write  $(N \subseteq M, tr)$  to denote a unital inclusion of finite von Neumann algebras, with ‘ $tr$ ’ a faithful normal tracial state, and write  $N \subset M \stackrel{e_1}{\subset} M_1$  for Jones’ resulting basic construction. The trace  $tr$  is called a *Markov trace of modulus  $\tau$*  if it extends to a positive trace  $Tr : M_1 \rightarrow \mathbb{C}$  such that  $Tr(xe_n) = \tau tr(x)$  for  $x \in M$ .

We confine ourselves to two Cases: (1) when the inclusion is one of  $II_1$  factors with finite index, i.e.,  $[M : N] < \infty$ ; and (2) when we have a connected inclusion of finite dimensional  $C^*$ -algebras. Then it is known that in both Cases ((1) and (2)) there exists a unique Markov trace on  $M$ , and we can iterate the basic construction to obtain a tower,

$$M_1 \subseteq M_2 \subseteq \dots \subseteq M_n \subseteq M_{n+1} \dots$$

where  $M_{n+1} = \langle M_n, e_{n+1} \rangle$  is the result of applying the basic construction for the pair  $M_{n-1} \subseteq M_n$  and  $e_{n+1}$  is the projection implementing the  $tr_{M_n}$  preserving conditional expectation of  $M_n$  onto

$M_{n-1}$ . We then obtain a  $II_1$  factor  $M_\infty$  in both the Cases, which is hyperfinite in Case (2).

Pimsner and Popa have shown (in [11]) that for an inclusion  $N \subset M$  of  $II_1$  factors,  $M$  is a finitely generated projective module over  $N$  if and only if  $[M : N]$  is finite by constructing a family  $\{m_j : 1 \leq j \leq n + 1\}$  of elements in  $M$ , with  $n$  equal to the integer part of  $[M : N]$ , which they called “orthonormal basis” for the pair  $N \subseteq M$ . In a similar manner, we find a slightly less restrictive notion of basis in [7].

In this paper (in section 2) we see that this notion of basis in [7] can also be carried out in our Case (2) of connected inclusions of finite dimensional  $C^*$ -algebras. Further in section 2 we characterize bases, in both Cases (1) and (2), by three equivalent conditions. One advantage of this characterization is a transparent proof of Corollary 2.7. This result has been mentioned for the Case of  $II_1$  factors in [7] (Lemma 4.3.4 (i)), but the proof there seems incomplete. Our characterization of bases now clarifies this point, and also shows that bases behave in a nice way with respect to the Jones’ tower.

As an application we show (in 3.1) how the use of bases leads to a natural proof of existence, in Case (2), (see [13](Theorem 2.1)) of a unique extension of an automorphism on  $M$  which leaves  $N$  globally invariant, to an automorphism on the hyperfinite  $II_1$  factor  $M_\infty$  which is compatible with the tower in the sense of fixing the Jones projections. It has been also proved that the initial automorphism will be automatically trace-preserving.

In [12](Proposition 1.2) Pimsner and Popa have characterized basic construction for  $II_1$  factor inclusion in two equivalent ways. See also [4](section 5). In this paper we have characterized basic construction in terms of basis we introduced(Lemma 3.4). We have succeeded to obtain a simple characterization of  $M_1$  for finite dimensional  $C^*$ -algebra Case also. In [12](Theorem 2.6)Pimsner and Popa have used their characterization of basic construction to describe the  $k$ -th step of the basic construction. In the section 3.2 we have also given another proof of this construction using our characterization of basic construction and have also done the same for connected inclusion of finite dimensional  $C^*$ -algebras.

## 2 Bases

As stated in the Introduction, we assume  $N \subseteq M$  is a unital inclusion of finite von Neumann algebras of one of the following two types.

*Case(1):*  $N$  and  $M$  are  $II_1$  factors with finite index  $[M : N]$  and hence there exists unique Markov trace  $tr$  on  $M$  of modulus  $\tau$  where  $\tau = [M : N]^{-1}$ .

*Case(2):* Let  $N \subseteq M$  be a connected inclusion of finite dimensional  $C^*$ -algebras and hence there exists unique Markov trace  $tr$  on  $M$  of modulus  $\tau$  where  $\tau = \|G\|^{-2}$  where  $G$  is the inclusion matrix for  $N \subseteq M$ .

For both the Cases the following easy but very useful Lemma holds whose proof can be found in [11], (Lemma 1.2) and for Case(2) see [7] (Remark 4.3.2(a)).

LEMMA 2.1. *If  $x_1 \in M_1$ , then there exists unique element  $x_0 \in M$  such that  $x_1 e_1 = x_0 e_1$ , this element is given by  $x_0 = \tau^{-1} E_M(x_1 e_1)$ .*

In the following theorem we give three equivalent descriptions of basis, not necessarily orthonormal in the sense of Pimsner-Popa.

THEOREM 2.2. *Let  $N$  and  $M$  be as in Case(1) or in Case(2). Then for a finite set  $\{\lambda_i : i \in I = 1, 2, \dots, n\} \subseteq M$ , the following are equivalent:*

(1) *Let  $E_N$  be the  $tr$ -preserving conditional expectation of  $M$  onto  $N$  and define a matrix  $Q$  whose  $(i, j)$  entry is given by  $q_{ij} = E_N(\lambda_i \lambda_j^*)$ . Then  $Q$  is a projection in  $M_n(N)$  such that  $tr_{M_n(N)}(Q) = \tau^{-1}/n$ .*

(2)  *$\sum_{i=1}^n \lambda_i^* e_1 \lambda_i = 1$ , where  $e_1$  is the Jones projection.*

(3) *For any  $x \in M$ ,  $x = \sum_{i=1}^n E_N(x \lambda_i^*) \lambda_i$ .*

*Proof.* (1)  $\implies$  (2) : This proof is mainly inspired by [11]. Assume (1) holds. Since  $tr$  on  $M$  is Markov, it extends to a unique trace on  $M_1$ , namely  $tr_{M_1}$ . Put  $v_i = e_1 \lambda_i$  and

$$v = \begin{bmatrix} v_1 & 0 & \dots & 0 \\ v_2 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ v_n & 0 & \dots & 0 \end{bmatrix}.$$

Then,  $v_i v_j^* = e_1 \lambda_i \lambda_j^* e_1 = E_N(\lambda_i \lambda_j^*) e_1 = q_{ij} e_1$ . Thus,

$$v v^* = \begin{bmatrix} q_{11} e_1 & q_{12} e_1 & \dots & q_{1n} e_1 \\ q_{21} e_1 & q_{22} e_1 & \dots & q_{2n} e_1 \\ \vdots & \vdots & \ddots & \vdots \\ q_{n1} e_1 & q_{n2} e_1 & \dots & q_{nn} e_1 \end{bmatrix} = Q E$$

where

$$E = \begin{bmatrix} e_1 & 0 & \dots & 0 \\ 0 & e_1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & e_1 \end{bmatrix}.$$

Thus by property of Jones projection [5] (Proposition 3.1.4),  $vv^* = QE = EQ$  and hence  $v$  is a partial isometry. Thus  $v^*v$  is a projection; i.e.,  $\sum_i v_i^*v_i$  is a projection  $f$  (say) in  $\langle M, e_1 \rangle = M_1$ . But  $f = \sum_i \lambda_i^* e_1 \lambda_i$  satisfies the following equations :

$$\begin{aligned} \text{tr}_{M_1} f &= n \text{tr}_{M_n(M_1)}(vv^*) \\ &= n \text{tr}_{M_n(M_1)}(QE) \\ &= n (1/n) \sum_i \text{tr}_{M_1}(q_{ii}e_1) \\ &= \sum_i \tau \text{tr}(q_{ii}) \quad (\text{Markov property}) \\ &= \tau n \text{tr}_{M_n(N)}(Q) \\ &= 1 \quad (\text{by (1)}). \end{aligned}$$

Thus  $(1 - f) \geq 0$  and  $\text{tr}_{M_1}(f) = 1$ . Then faithfulness of  $\text{tr}$  implies  $f = 1$ . So  $\sum_i \lambda_i^* e_1 \lambda_i = 1$ . Thus (1) implies (2).

(2)  $\implies$  (3): We assume that (2) holds. Let  $x^* \in M$ , then

$$\begin{aligned} x^* e_1 &= \left( \sum_i \lambda_i^* e_1 \lambda_i \right) x^* e_1 \\ &= \sum_i \lambda_i^* E_N(\lambda_i x^*) e_1 \\ &= \left( \sum_i \lambda_i^* E_N(\lambda_i x^*) \right) e_1. \end{aligned}$$

Again applying Lemma 2.1 and then taking adjoint we get (3).

(3)  $\implies$  (2) : We assume (3). Let  $x$  and  $y$  be two arbitrary elements of  $M$ . Then,

$$\begin{aligned} \left( \sum_i \lambda_i^* e_1 \lambda_i \right) (x e_1 y) &= \sum_i \lambda_i^* e_1 \lambda_i x e_1 y \\ &= \sum_i \lambda_i^* E_N(\lambda_i x) e_1 y \\ &= (x e_1 y) \quad (\text{by (3)}). \end{aligned}$$

Similarly,

$$\begin{aligned}
(xe_1y)\left(\sum_i \lambda_i^* e_1 \lambda_i\right) &= \sum_i x e_1 y \lambda_i^* e_1 \lambda_i \\
&= \sum_i x e_1 E_N(y \lambda_i^*) \lambda_i \\
&= (x e_1 y) \quad (\text{by (3)}).
\end{aligned}$$

Then we know the space  $M e_1 M$ , which is linear span of  $\{x e_1 y : x, y \in M\}$ , is a strongly dense \*-subalgebra of  $M_1$ , see for instance [2](Proposition 3.6.1(vii)). Then since multiplication is separately strongly continuous it follows that  $\sum_i \lambda_i^* e_1 \lambda_i = 1$ .

(2)  $\implies$  (1) : Suppose (2) is true. Then,

$$\begin{aligned}
e_1\left(\sum_k q_{ik} q_{kj}\right) &= e_1\left(\sum_k E_N(\lambda_i \lambda_k^*) E_N(\lambda_k \lambda_j^*)\right) \\
&= e_1\left(\sum_k E_N(\lambda_i \lambda_k^* E_N(\lambda_k \lambda_j^*))\right) \\
&= \sum_k e_1 \lambda_i \lambda_k^* E_N(\lambda_k \lambda_j^*) e_1 \\
&= \sum_k e_1 \lambda_i \lambda_k^* e_1 \lambda_k \lambda_j^* e_1 \\
&= e_1 \lambda_i \left(\sum_k \lambda_k^* e_1 \lambda_k\right) \lambda_j^* e_1 \\
&= e_1 \lambda_i \lambda_j^* e_1 \quad (\text{by (2)}) \\
&= e_1 E_N(\lambda_i \lambda_j^*) \\
&= e_1 q_{ij}.
\end{aligned}$$

Thus applying Lemma 2.1 we get  $Q^2 = Q$ . Clearly  $Q^* = Q$ . Hence  $Q$  is a projection in  $M_n(N)$ . Now

$$\begin{aligned}
tr_{M_n(N)}(Q) &= (1/n) \sum_i tr(q_{ii}) \\
&= (1/n) \sum_i tr(E_N(\lambda_i \lambda_i^*)) \\
&= (1/n) \sum_i tr(\lambda_i \lambda_i^*) \\
&= (\tau^{-1}/n) \sum_i tr(e_1 \lambda_i \lambda_i^*) \quad (\text{Markov Property}) \\
&= (\tau^{-1}/n) \sum_i tr(\lambda_i^* e_1 \lambda_i) \\
&= (\tau^{-1}/n).
\end{aligned}$$

Hence (2) implies (1). □

REMARK 2.3. Taking adjoints in (3) it follows that the above three are also equivalent to  $x = \sum_{i=1}^n \lambda_i^* E_N(\lambda_i x)$ , for all  $x \in M$ .

DEFINITION 2.4. A finite set  $\{\lambda_i : i \in I\} \subset M$  satisfying any one of the equivalent conditions (i)-(iii) of Theorem 2.2 will simply be called a **basis** for  $M/N$ .

**Existence of bases:** For Case(1) an explicit construction has been given in [11](Proposition 1.3) while for Case (2) see [7] (Lemma 5.7.3), and [6] (Proposition 2.5). For Case(2) see also [1](section 9.4).

REMARK 2.5. Comparing [11] (Proposition 1.3(c)(2)) and Theorem 2.2 we remark that any Pimsner-Popa basis for  $II_1$  factor inclusions is automatically a basis according to our notion. Also, motivated by [11] and [9], Watatani has introduced (in the memoir [15]) what he calls ‘quasi-basis for conditional expectation  $E$ ’ in a purely algebraic setting. Assuming the existence of quasi-basis he developed index for a conditional expectation of index-finite type, called it Index  $E$ , which he shows to be independent of the choice of quasi-basis. He then investigated Jones’ index theory in  $C^*$ -algebra setting. Observe that, Theorem 2.2 (1) now says that Index  $E$  is same as Jones’ index for Case (1) and equals to  $\|G\|^2$  for Case (2).

REMARK 2.6. The row vector  $[E_N(x\lambda_1^*), \dots, E_N(x\lambda_n^*)] \in M_{1 \times n}(N)Q$  and conversely if  $[x_1, \dots, x_n] \in M_{1 \times n}(N)Q$  satisfies  $x = \sum_{i=1}^n x_i \lambda_i$  then  $x_j = E_N(x\lambda_j^*)$  for all  $j \in I$ .

Exactly the same proof as in [7] (Proposition 4.3.3(b)(ii)) works.

COROLLARY 2.7. Let  $N \subseteq M \subseteq P$  be a tower of  $II_1$  factors with  $[P : N] < \infty$  (or a tower of finite dimensional  $C^*$ -algebras where the two inclusions are connected with inclusion matrices  $G$  and  $H$  respectively). In either Case, let  $\{\lambda_i : 1 \leq i \leq m\}$  be a basis for  $M/N$  and  $\{\mu_j : 1 \leq j \leq n\}$  be a basis for  $P/M$ , then  $\{\lambda_i \mu_j : 1 \leq i \leq m, 1 \leq j \leq n\}$  is a basis for  $P/N$ .

*Proof.* Let  $x \in P$  and as  $\{\mu_j\}$  is a basis for  $P/M$ , we get,  $x = \sum_{j=1}^n E_M(x\mu_j^*)\mu_j$ . Now note  $E_M(x\mu_j^*) \in M$  and  $\{\lambda_i\}$  is a basis for  $M/N$ . Now condition(3) of the Theorem 2.2 yields,

$$E_M(x\mu_j^*) = \sum_{i=1}^m E_N\{E_M(x\mu_j^*)\lambda_i^*\}\lambda_i.$$

Thus we get,

$$\begin{aligned} x &= \sum_{j=1}^n \left[ \sum_{i=1}^m E_N \{ E_M(x\mu_j^*)\lambda_i^* \} \lambda_i \right] \mu_j \\ &= \sum_{i=1}^m \sum_{j=1}^n E_N \{ E_M(x\mu_j^*\lambda_i^*) \} \lambda_i \mu_j. \end{aligned}$$

Thus again applying (3) of the Theorem 2.2 we get that  $\{\lambda_i \mu_j : 1 \leq i \leq m, 1 \leq j \leq n\}$  is a basis for  $P/N$ .  $\square$

**COROLLARY 2.8.** *If  $\{\lambda_i : i \in I = \{1, 2, \dots, n\}\}$  is a basis for  $M/N$ , then  $\{\tau^{-1/2}e_1\lambda_i\}$  is a basis for  $M_1/M$ . Where  $N \subseteq M$  is an inclusion as in Case (1) or Case (2).*

*Proof.* In the Case (1)  $M_1$  is a  $II_1$  factor such that  $[M_1 : M] = [M : N] < \infty$ . In Case (2) inclusion matrix for  $M \subseteq M_1$  is  $G^t$  and hence is a connected inclusion. Now in both the Cases let  $e_2$  be the Jones projection for the inclusion  $M \subseteq M_1$ . Now,

$$\begin{aligned} \sum_{i=1}^n \{ \tau^{-1/2}e_1\lambda_i \}^* e_2 \{ \tau^{-1/2}e_1\lambda_i \} &= \tau^{-1} \sum_{i=1}^n \lambda_i^* e_1 e_2 e_1 \lambda_i \\ &= \tau^{-1} \tau \sum_{i=1}^n \lambda_i^* e_1 \lambda_i \\ &= 1 \quad (\text{since } \{\lambda_i\} \text{ is a basis}). \end{aligned}$$

Now (2) of the Theorem 2.2 yields the result.  $\square$

**REMARK 2.9.** *From Corollary 2.8 every element of  $M_1$  is expressible in the form  $\sum_{i=1}^n x_i e_1 y_i$  for some  $x_i, y_i \in M$  (in fact,  $x = \tau^{-1} \sum_{i=1}^n E_M(x\lambda_i^* e_1) e_1 \lambda_i$ ); however this does not allow us to define a \*-homomorphism on  $M_1$  by merely specifying the image of an element of the form  $x e_1 y$ , as we will need to verify that such a ‘definition’ is unambiguous; but we may define the above canonical decomposition to unambiguously define maps on  $M_1$  once we know where to map elements of  $N$ , the basis vectors  $\lambda_i$  and  $e_1$ . This problem of ambiguity was part of the reason for us the study this notion of bases. The reader need only compare the crisp clarity of the proofs of unambiguity in the definition of  $\alpha_1$  in Theorem 3.2 and of  $\phi$  in Lemma 3.4 with the corresponding proofs of Theorem 2.1 in [13] (actually only to be found in the arXiv version) and of Proposition 1.2 in [12], to appreciate this remark.*

COROLLARY 2.10. *Let  $N \subseteq M$  as in Case (1) or (2) and  $\{\lambda_i : i \in I\}$  be a basis for  $M/N$ . Define  $\widehat{i(k)} = (i_1, i_2, \dots, i_k) \in I^k, k \geq 1$  and  $\lambda_{\widehat{i(k)}} = \tau^{-k(k-1)/4} \lambda_{i_1} e_1 \lambda_{i_2} e_2 e_1 \lambda_{i_3} \dots \lambda_{i_{k-1}} e_{k-1} \dots e_1 \lambda_{i_k}$ . Then  $\{\lambda_{\widehat{i(k)}} : \widehat{i(k)} \in I^k\}$  is a basis for  $M_{k-1}/N$ .*

*Proof.* Clearly the statement is true for  $k = 1$  with the understanding that  $M_0 = M$ . Suppose the statement is true for  $k$ . Now applying Corollary 2.8 recursively we get  $M_k/M_{k-1}$  has basis

$\{\tau^{-k/2} e_k e_{k-1} \dots e_1 \lambda_{i_{k+1}} : i_{k+1} \in I\}$ . Then applying Corollary 2.7 we see that  $M_k/N$  has basis,

$\{\tau^{-k(k-1)/4} \tau^{-k/2} \lambda_{i_1} e_1 \lambda_{i_2} e_2 e_1 \lambda_{i_3} \dots \lambda_{i_{k-1}} e_{k-1} \dots e_1 \lambda_{i_k} e_k e_{k-1} \dots e_1 \lambda_{i_{k+1}}\}$   
 $= \{\lambda_{\widehat{i(k+1)}} : \widehat{i(k+1)} \in I^{k+1}\}$ ; and the proof of the inductive step is complete.  $\square$

### 3 Applications

#### 3.1 Compatible automorphisms of the Hyperfinite $II_1$ factor

Consider an inclusion as in Case (2). Then, we have a unique Markov trace  $tr$  on  $M$ . Next consider the Jones tower  $N \subseteq M \subseteq M_1 \subseteq M_2 \subseteq \dots$  and let  $R$  be the hyperfinite  $II_1$  factor arising from this tower [7]. Suppose further we have an automorphism  $\alpha_0$  on  $M$  such that  $\alpha_0(N) = N$ . In this present section we shall show using our concept of basis how we can construct a unique extension of  $\alpha_0$  to an automorphism  $\alpha$  of the hyperfinite  $II_1$  factor  $R$  which is compatible with respect to the tower in the sense of fixing all the Jones projections and leaving  $M_i$  invariant.

This can be thought of as the finite dimensional  $C^*$ -algebraic version of [10] (Lemma 5.1). In that paper Loi studied automorphisms for a pair of factors using standard form of von Neumann algebras, whereas our treatment is based on basis for the corresponding inclusion. In the similar direction in [8] the author has dealt with automorphisms commuting with a faithful normal conditional expectation for a pair of  $\sigma$ -finite von Neumann algebras and related this with an action of a locally compact abelian group. See also [14], where the author was more concerned with commuting squares.

LEMMA 3.1. *Let  $N, M, tr, \alpha_0$  be as above. Then  $\alpha_0$  is automatically trace preserving, that is  $tr \circ \alpha_0 = tr$ .*

*Proof.* Let the minimal central projections in  $N$  be  $\{p_1, p_2, \dots, p_m\}$  and those in  $M$  be  $\{q_1, q_2, \dots, q_n\}$ . Then the inclusion matrix  $G$  is an



$m \times n$  matrix. Observe  $\alpha_0$  permutes  $p_i$ 's and  $q_j$ 's. Say  $p_i \mapsto p_{\tau(i)}$  and  $q_j \mapsto q_{\sigma(j)}$  for  $\tau \in \Sigma_m$  and  $\sigma \in \Sigma_n$ . As  $\alpha_0$  is an automorphism,  $G(i, j) = G(\tau(i), \sigma(j))$ . Equivalently,  $G = TGS$  for permutation matrices  $T$  and  $S$  of sizes  $m$  and  $n$  respectively. Let  $\vec{t}$  be the trace vector corresponding to  $tr$  for  $M$ . Then it is the unique positive Perron-Frobenius eigenvector of  $G^tG$ , hence also of  $S^{-1}G^tGS$ . But that implies  $S\vec{t}$  is a positive eigenvector of  $G^tG$  with the same eigenvalue as of  $\vec{t}$  and by uniqueness of Perron-Frobenius theory (see chapter XIII [3]) we get  $S\vec{t} = \vec{t}$ . Hence  $tr \circ \alpha_0 = tr$ .  $\square$

The above proof is due to Vijay Kodiyalam. I sincerely thank him for this.

**THEOREM 3.2.** *Let  $\alpha_0$  be an automorphism of  $M$  such that  $\alpha_0(N) = N$ . Then there is a unique (trace preserving) automorphism  $\alpha_1$  of  $M_1$  such that  $\alpha_1(e_1) = e_1$ ,  $\alpha_1(M) = M$  and the restriction of  $\alpha_1$  to  $M$  is  $\alpha_0$ .*

*Proof.* We know there is a basis for  $M/N$ . Fix such a basis  $\{\lambda_i : i \in I\}$ . Then we show  $\{\alpha_0(\lambda_i) : i \in I\}$  is also a basis for  $M/N$ . Let  $Q_1$  be the matrix with  $(i, j)$  entry given by  $q_1(i, j) = E_N\{\alpha_0(\lambda_i \lambda_j^*)\}$ . Now,

$$\begin{aligned} tr_{M_n(N)}(Q_1) &= (1/n) \sum_{i \in I} tr(q_1(i, i)) \\ &= (1/n) \sum_{i \in I} tr[E_N\{\alpha_0(\lambda_i \lambda_i^*)\}] \\ &= (1/n) \sum_{i \in I} tr(\lambda_i \lambda_i^*) && \text{(by Lemma (3.1))} \\ &= tr_{M_n(N)}Q. \end{aligned}$$

Thus it follows from the Theorem 2.2, that  $\{\alpha_0(\lambda_i)\}$  is a basis for  $M/N$ . Observe since  $\alpha_0$  leaves  $N$  invariant it follows that  $E_N(\alpha_0(x)) = \alpha_0(E_N(x))$ . Let  $x \in M_1$ , Corollary 2.8 then implies

$$x = \sum_{i \in I} \tau^{-1} E_M(x \lambda_i^* e_1) e_1 \lambda_i.$$

Then define,

$$\alpha_1(x) = \tau^{-1} \sum_{i \in I} \alpha_0(E_M(x \lambda_i^* e_1)) e_1 \alpha_0(\lambda_i).$$

There is clearly no ambiguity in the definition of  $\alpha_1$ . Next we show that  $\alpha_1$  is a homomorphism. Consider  $y \in M_1$ . Now using the properties of Jones' projection and the fact that  $\alpha_0$  is a homomorphism we get the following series of equations:

$$\begin{aligned}
& \alpha_1(x)\alpha_1(y) \\
&= \tau^{-2} \sum_{i,j} \alpha_0[E_M(x\lambda_i^*e_1)]e_1\alpha_0(\lambda_i)\alpha_0[E_M(y\lambda_j^*e_1)]e_1\alpha_0(\lambda_j) \\
&= \tau^{-2} \sum_{i,j} \alpha_0[E_M(x\lambda_i^*e_1)]E_N(\alpha_0[\lambda_i E_M(y\lambda_j^*e_1)])e_1\alpha_0(\lambda_j) \\
&= \tau^{-2} \sum_{i,j} \alpha_0[E_M(x\lambda_i^*e_1)E_N(\lambda_i E_M(y\lambda_j^*e_1))]e_1\alpha_0(\lambda_j) \\
&\quad (\text{since } \alpha_0 \text{ and } E_N \text{ commute}) \\
&= \tau^{-2} \sum_{i,j} \alpha_0[E_M\{x\lambda_i^*e_1 E_N(\lambda_i E_M(y\lambda_j^*e_1))\}]e_1\alpha_0(\lambda_j) \\
&= \tau^{-2} \sum_j \alpha_0[E_M\{xE_M(y\lambda_j^*e_1)e_1\}]e_1\alpha_0(\lambda_j) \tag{1} \\
&\quad (\text{since } \sum_i \lambda_i^*e_1\lambda_i = 1).
\end{aligned}$$

Similarly,

$$\begin{aligned}
& \alpha_1(xy) \\
&= \tau^{-1} \sum_i \alpha_0[E_M(xy\lambda_i^*e_1)]e_1\alpha_0(\lambda_i) \\
&= \tau^{-1} \sum_i \alpha_0[E_M\{x(\sum_j \tau^{-1}E_M(y\lambda_j^*e_1)e_1\lambda_j)\lambda_i^*e_1\}]e_1\alpha_0(\lambda_i) \\
&= \tau^{-2} \sum_{i,j} \alpha_0[E_M\{xE_M(y\lambda_j^*e_1)E_N(\lambda_j\lambda_i^*)e_1\}]e_1\alpha_0(\lambda_i) \\
&= \tau^{-2} \sum_{i,j} \alpha_0[E_M\{xE_M(y\lambda_j^*e_1)E_N(\lambda_j\lambda_i^*)\}]e_1\alpha_0(\lambda_i) \\
&= \tau^{-2} \sum_i \alpha_0[E_M\{xE_M(y\lambda_i^*e_1)e_1\}]e_1\alpha_0(\lambda_i) \tag{2} \\
&\quad (\text{since } \sum_j \lambda_j^*e_1\lambda_j = 1).
\end{aligned}$$

Now comparing equations (1) and (2) we conclude that  $\alpha_1$  is indeed a homomorphism. Next we show  $\alpha_1$  fixes  $e_1$ . Observe,

$$e_1 = \tau^{-1} \sum_i E_M(e_1\lambda_i^*e_1)e_1\lambda_i.$$

Now using our definition of  $\alpha_1$  and property of Jones' projection it is easy to see that,

$$\begin{aligned}
\alpha_1(e_1) &= \tau^{-1} \sum_i \alpha_0[E_M\{E_N(\lambda_i^*)e_1\}]e_1\alpha_0(\lambda_i) \\
&= \sum_i \alpha_0(E_N(\lambda_i^*))e_1\alpha_0(\lambda_i) \\
&= \sum_i E_N(\alpha_0(\lambda_i^*))e_1\alpha_0(\lambda_i) \quad (\text{as } E_N \text{ and } \alpha_0 \text{ commute}) \\
&= \sum_i e_1\alpha_0(\lambda_i)^*e_1\alpha_0(\lambda_i) \\
&= e_1.
\end{aligned}$$

In the last equation we have used the fact that  $\{\alpha_0(\lambda_i)\}$  is a basis for  $M/N$ .

Next we will show that  $\alpha_1$  agrees with  $\alpha_0$  when it is restricted to  $M$ . Now, since  $\alpha_0$  is a automorphism for  $x \in M$  we find that,

$$\begin{aligned}
\alpha_1(x) &= \tau^{-1} \sum_i \alpha_0\{E_M(x\lambda_i^*)e_1\}e_1\alpha_0(\lambda_i) \\
&= \sum_i \alpha_0(x\lambda_i^*)E_M(e_1)e_1\alpha_0(\lambda_i) \\
&= \sum_i \alpha_0(x)\alpha_0(\lambda_i)^*e_1\alpha_0(\lambda_i) \quad (\text{since } E_M(e_1) = \tau) \\
&= \alpha_0(x) \quad (\text{as } \{\alpha_0(\lambda_i)\} \text{ is a basis for } M/N).
\end{aligned}$$

Now we want to show that  $\alpha_1$  is onto.

Let  $y \in M_1$ . Then,  $y = \sum_i y_i e_1 \alpha_0(\lambda_i)$ , since  $\alpha_0(\lambda_i)$  is a basis for  $M/N$ . As,  $\alpha_0$  is an automorphism there is a unique  $x_i \in M$  such that  $\alpha_0(x_i) = y_i$ . Put  $x = \sum_i x_i e_1 \lambda_i$ . Then  $x$  belongs to  $M_1$ . Now as we have already proved that  $\alpha_1$  is a homomorphism which preserves  $e_1$  and agree with  $\alpha_0$  when restricted to  $M$  it follows trivially that  $\alpha_1(x) = y$ . Thus  $\alpha_1$  is onto.

Lastly we show  $\alpha_1$  is one-one. Observe,  $\alpha_1$  is \*-preserving, since if  $x = \sum_i x_i e_1 \lambda_i$  we find, exactly as above, that

$$\alpha_1(x^*) = \sum_i \alpha_1(\lambda_i)^* e_1 \alpha_1(x_i)^* = \left\{ \sum_i \alpha_1(x_i) e_1 \alpha_1(\lambda_i) \right\}^* = \alpha_1(x)^*.$$

Now,

$$\begin{aligned}
tr(\alpha_1(x)) &= tr\left(\sum_i \alpha_0(x_i)e_1\alpha_0(\lambda_i)\right) \\
&= \sum_i tr\{e_1\alpha_0(\lambda_i)\alpha_0(x_i)\} \\
&= \tau \sum_i tr\{\alpha_0(\lambda_i x_i)\} \quad (\text{Markov property}) \\
&= \tau \sum_i tr(\lambda_i x_i) \quad (\text{by Lemma 3.1}) \\
&= \sum_i tr(x_i e_1 \lambda_i) \quad (\text{Markov property}) \\
&= tr(x).
\end{aligned}$$

so  $\alpha_1$  is  $tr$ -preserving and hence one-one. The uniqueness assertion is obvious since  $M$  and  $e_1$  generate  $M_1$ . Thus  $\alpha_1$  satisfies all the properties mentioned in the Theorem.  $\square$

**COROLLARY 3.3.** *Let  $\alpha_0$  be as in the previous theorem. Then there is a unique (trace preserving) automorphism  $\alpha$  of the hyperfinite  $II_1$  factor  $R$  such that  $\alpha(e_i) = e_i, \alpha(M_i) = M_i$  for all  $i \geq 1$  and  $\alpha|M = \alpha_0$ .*

*Proof.* Apply Theorem 3.2 recursively for the tower of basic construction to get a unique (trace preserving) automorphism  $\alpha_i$  on  $M_i$  which leaves  $M_j$  invariant and fixes all  $e_j$  such that  $1 \leq j \leq i$  and  $\alpha_i|M_j = \alpha_j$ . Thus we can define an automorphism (compatible with respect to the tower)  $\alpha_\infty$  on  $\cup_i M_i$  by,  $\alpha_\infty(x) = \alpha_j(x)$  for  $x \in M_j$ . Now as  $\alpha_\infty$  is bounded it extends to trace preserving automorphism  $\alpha$  (say) on  $R$ . Also since  $M_1$  and  $e_i$  s generate  $R$  uniqueness is straightforward.  $\square$

### 3.2 Iterating basic construction

The following gives a characterization of basic construction using bases, in both Case(1) and Case(2). This would be needed for our proof of the assertion regarding  $k$ -th step basic constructions.

**LEMMA 3.4.** *Let  $N \subseteq M$  be as in Case(1) or Case(2). Assume  $\{\lambda_i : i \in \{1, 2, \dots, n\}\}$  is a basis for  $M/N$  (which exists in both the Cases). Let  $P$  be a  $II_1$  factor in Case(1) or a finite dimensional  $C^*$ -algebra in Case (2) such that  $P$  contains  $M$  and also contains*

a projection  $f$  such that  $\sum_{i=1}^n \lambda_i^* f \lambda_i = 1$  and satisfies further the following two properties :

- 1)  $f x f = E_N(x) f$  for all  $x \in M$  and
- 2)  $\{\tau^{-1/2} f \lambda_i\}$  is a basis for  $P/M$ .

In addition for Case(2)  $P$  satisfies the following property also:

- 3)  $n \longmapsto n f$  is an injective map from  $N$  into  $P$ .

Then there exists an isomorphism from  $M_1 = \langle M, e_1 \rangle$  onto  $P$  which maps  $e_1$  to  $f$ .

In this situation we say that  $P$  is an instance of basic construction applied to the inclusion  $N \subseteq M$  with a choice of projection implementing the conditional expectation being given by  $f$ .

*Proof. Case1 :* Let  $x \in M_1$ . Now from Corollary 2.8 it follows that

$$x = \sum_i \tau^{-1/2} E_M(x \tau^{-1/2} \lambda_i^* e_1) e_1 \lambda_i.$$

Put  $a_i = \tau^{-1} E_M(x \lambda_i^* e_1)$ , then define a map  $\phi : M_1 \mapsto P$  by  $\phi(x) = \sum_i a_i f \lambda_i$ , which is clearly well-defined. Note, if  $y = \sum_i b_i e_1 \lambda_i$  such that  $[b_1, b_2, \dots, b_n] \in M_{1 \times n}(M)Q$ , then by Remark 2.6 we conclude

$$\phi(y) = \sum_i b_i f \lambda_i. \quad (3.1)$$

Since, if  $Q_1$  is the matrix whose  $i - j$  th entry is given by  $q_1(i, j) = E_M((\tau^{-1/2} e_1 \lambda_i)(\tau^{-1/2} e_1 \lambda_j)^*)$ , then  $q_1(i, j) = E_N(\lambda_i \lambda_j^*) = q_{ij}$ . Now, let  $x$  be as above and let  $y \in M_1$ . Put  $y = \sum_i b_i e_1 \lambda_i$  where  $b_i = \tau^{-1} E_M(y \lambda_i^* e_1)$ . Then the following equations follow from properties of Jones' projection,

$$\begin{aligned} \phi(xy) &= \phi\left\{\sum_{i,j} a_i E_N(\lambda_i b_j) e_1 \lambda_j\right\} \\ &= \phi\left\{\sum_{i,j} \tau^{-1} E_M(x \lambda_i^* e_1) E_N(\lambda_i b_j) e_1 \lambda_j\right\} \\ &= \phi\left\{\sum_{i,j} \tau^{-1} E_M(x \lambda_i^* e_1 E_N(\lambda_i b_j)) e_1 \lambda_j\right\} \\ &= \phi\left\{\sum_j \tau^{-1} E_M(x b_j e_1) e_1 \lambda_j\right\} \quad (\text{since } \sum_i \lambda_i^* e_1 \lambda_i = 1) \\ &= \phi\left\{\sum_j \tau^{-1} E_M[x \tau^{-1} E_M(y \lambda_j^* e_1) e_1] e_1 \lambda_j\right\}. \end{aligned}$$

Now it can be easily checked that,

$$[\tau^{-1} E_M\{x \tau^{-1} E_M(y \lambda_1^* e_1) e_1\}, \tau^{-1} E_M\{x \tau^{-1} E_M(y \lambda_2^* e_1) e_1\}, \dots, \tau^{-1} E_M\{x \tau^{-1} E_M(y \lambda_n^* e_1) e_1\}] \in M_{1 \times n}(M)Q.$$

Thus, it follows from equation(3.1) that,

$$\begin{aligned}
\phi(xy) &= \sum_j \tau^{-1} E_M[x\tau^{-1} E_M(y\lambda_j^* e_1) e_1] f \lambda_j \\
&= \sum_{i,j} \tau^{-1} E_M(a_i e_1 \lambda_i b_j e_1) f \lambda_j \\
&= \sum_{i,j} \tau^{-1} a_i E_M(E_N(\lambda_i b_j) e_1) f \lambda_j \\
&= \sum_{i,j} a_i E_N(\lambda_i b_j) f \lambda_j \\
&= \sum_{i,j} a_i f \lambda_i b_j f \lambda_j \quad (\text{by assumption (1)}) \\
&= \phi(x)\phi(y).
\end{aligned}$$

Also, we have,

$$\begin{aligned}
\phi(e_1) &= \tau^{-1} \sum_i E_M(e_1 \lambda_i^* e_1) f \lambda_i \\
&= \sum_i E_N(\lambda_i^*) f \lambda_i \quad (\text{since } E_M(e_1) = \tau) \\
&= \sum_i f \lambda_i^* f \lambda_i \quad (\text{by assumption (1)}) \\
&= f \quad (\text{since, } \sum_i \lambda_i^* f \lambda_i = 1).
\end{aligned}$$

Thus  $\phi$  is a nonzero homomorphism. Now assume,  $x \in M$ , then,

$$\begin{aligned}
\phi(x) &= \sum_i \tau^{-1} E_M(x \lambda_i^* e_1) f \lambda_i \\
&= \sum_i x \lambda_i^* f \lambda_i \quad (\text{since } x \lambda_i^* \in M) \\
&= x.
\end{aligned}$$

$\phi$  is also  $*$ -preserving, as, if  $x = \sum_i a_i e_1 \lambda_i$  is any element of  $M_1$ , then the following identities hold:

$$\begin{aligned}
\phi(\{\sum_i a_i e_1 \lambda_i\}^*) &= \sum_i \lambda_i^* f a_i^* \quad (\text{since } \phi(e_1) = f \text{ and } \phi|_M = id) \\
&= \{\sum_i a_i f \lambda_i\}^* \\
&= \{\phi(\sum_i a_i e_1 \lambda_i)\}^*.
\end{aligned}$$

Thus  $\phi(x^*) = \phi(x)^*$ .

Since we are now in a factor  $\phi$  is automatically injective.

Finally we show  $\phi$  is onto. For this purpose assume  $z \in P$ , assumption(2) then implies  $z = \sum_i c_i f \lambda_i$  for some  $c_i \in M$ . Put,  $y = \sum_i c_i e_1 \lambda_i$  which belongs to  $M_1$  and since  $\phi$  is a homomorphism sending  $e_1$  to  $f$  and whose restriction to  $M$  is identity, we clearly get  $\phi(y) = z$ , proving onto. Thus  $\phi$  is an isomorphism satisfying all the conditions stated in the Lemma.

*Case2* : Note assumption(2) implies  $P = MfM$ . Also this together with assumption(1) imply that  $PfP = MfM$ . Thus  $P = PfP$  which forces  $Z_P(f) = 1$ . Now just applying Corollary 5.3.2 in [7] we get the result.

This completes the Lemma. □

Now we give another proof of  $k$ -th step basic construction for an inclusion of  $II_1$  factors using basis and also we show it can be done for Case(2).

**THEOREM 3.5.** *Let  $N \subseteq M$  be a pair of von Neumann algebras as in Case(1) or (2) and  $N \subseteq M \subseteq M_1 \subseteq \dots$  be the tower of  $II_1$  factors (or finite dimensional  $C^*$ -algebras) in Case(1) (or in Case(2) respectively) which can be obtained by iterating basic construction. Let  $e_i \in M_i$  be the Jones' projections. Then for  $m \geq 0, k \geq -1$ ,  $M_k \subseteq M_{k+m} \subseteq M_{k+2m}$  is an instance of basic construction with a choice of projection implementing the conditional expectation of  $M_{k+m}$  onto  $M_k$  is given by*

$$\begin{aligned} e^{[k,k+m]} &= \tau^{-m(m-1)/2} (e_{k+m+1} e_{k+m} \dots e_{k+2}) (e_{k+m+2} e_{k+m+1} \dots e_{k+3}) \\ &\quad \dots (e_{k+2m} e_{k+2m-1} \dots e_{k+m+1}). \end{aligned}$$

*Proof.* Without loss of generality we shall prove that  $M_{-1} \subseteq M_n \subseteq M_{2n+1}$  is an instance of basic construction with  $e_{[-1,n]}$  is the required projection. Assume  $\{\lambda_i : i \in 1, 2, \dots, n\}$  is a basis for  $M/N$  (which exists in both the Cases). Now from Corollary 2.10 we know that  $\{\lambda_{\widehat{i(n+1)}}\}$  is a basis for  $M_n/N$ .

Now applying Corollary 2.7 and Corollary 2.8 repeatedly we get  $M_{2n+1}/M_n$  has basis,

$$\tau^{-1/2\{(n+1)+(n+2)+\dots+(2n+1)\}} (e_{n+1} \dots e_1) \lambda_{i_1} (e_{n+2} \dots e_1) \times$$

$$\lambda_{i_2} \dots (e_{2n} \dots e_1) \lambda_{i_n} (e_{2n+1} \dots e_1) \lambda_{i_{n+1}}.$$

Observe that,

$$\begin{aligned} & (e_{n+1} \dots e_1) \lambda_{i_1} (e_{n+2} \dots e_1) \lambda_{i_2} \dots (e_{2n+1} \dots e_1) \lambda_{i_{n+1}} \\ &= (e_{n+1} \dots e_1) (e_{n+2} \dots e_2) (e_{n+3} \dots e_3) \dots (e_{2n+1} \dots e_{n+1}) \\ & \quad \lambda_{i_1} e_1 \lambda_{i_2} e_2 e_1 \lambda_{i_3} \dots \lambda_{i_n} e_n \dots e_1 \lambda_{i_{n+1}}. \end{aligned}$$

In other words it shows that,  $M_{2n+1}/M_n$  has basis as,  $\{\tau^{-(n+1)/2} e_{[-1, n]} \lambda_{\widehat{i(n+1)}}\}$ .

Note,  $[M_n : N] = [M : N]^{(n+1)} = \tau^{-(n+1)}$ . Thus condition (2) of the Lemma 3.4 holds for factor Case.

To do the same for finite dimensional  $C^*$ -algebra we break this into two Cases.

*Case1* : Suppose  $n$  is odd. Then the inclusion matrix for  $N \subseteq M$  would be  $(GG^t)^k$  where  $n = (2k - 1)$ . But it is easy to see that  $\|(GG^t)^k\| = \|G\|^{2k} = \|G\|^{(n+1)} = \tau^{-(n+1)/2}$ . Thus condition (2) of the Lemma 3.4 holds in this Case.

*Case2* : Here  $n$  is even  $n = 2m$  (say). Then the inclusion matrix for  $N \subseteq M$  would be  $G(G^tG)^m$ . Then we see,  $\|G^tG(G^tG)^m\| \leq \|G^t\| \|G(G^tG)^m\| = \|G\| \|G(G^tG)^m\|$ . Now applying the Case(1) in left hand side, we get that  $\|G\|^{2m+1} \leq \|G(G^tG)^m\|$ . The opposite inequality is obvious. Thus condition(2) of Lemma 3.4 holds in this Case also.

We need to show that , for all  $k \geq 1$ , (for both Cases),

$$\sum_{i_1, i_2, \dots, i_k} \lambda_{\widehat{i(k)}}^* e_{[-1, k-1]} \lambda_{\widehat{i(k)}} = 1. \quad (3.2)$$

We prove it by induction over  $k \geq 1$ . It is easy to see that,

$$\lambda_{\widehat{i(n)}} (\tau^{-n/2} e_n \dots e_1 \lambda_{i_{n+1}}) = \lambda_{\widehat{i(n+1)}}.$$

and hence,

$$(\tau^{-n/2} \lambda_{i_{n+1}}^* e_1 \dots e_n) \lambda_{\widehat{i(n)}}^* = \lambda_{\widehat{i(n+1)}}^*.$$

Suppose, as induction hypothesis, for  $n \geq 1$ ,

$$\sum_{i_1, i_2, \dots, i_n} \lambda_{\widehat{i(n)}}^* e_{[-1, n-1]} \lambda_{\widehat{i(n)}} = 1. \quad (3.3)$$

Since  $\sum_i \lambda_i^* e_1 \lambda_i = 1$ , we see that equation (3.2) holds for  $k = 1$ . Also we know, for  $n \geq 1$ ,

$$e_{[-1, n]} = \tau^{-n} (e_{n+1} e_{n+2} \dots e_{2n+1}) e_{[-1, n-1]} (e_{2n} e_{2n-1} \dots e_{n+1}).$$



Thus,

$$\begin{aligned}
& \sum_{i_1, i_2, \dots, i_{n+1}} \lambda_{\widehat{i(n+1)}}^* e_{[-1, n]} \lambda_{\widehat{i(n+1)}} \\
&= \sum_{i_1, i_2, \dots, i_{n+1}} \tau^{-2n} \lambda_{i_{n+1}}^* (e_1 e_2 \dots e_n) \lambda_{\widehat{i(n)}}^* (e_{n+1} e_{n+2} \dots e_{2n+1}) e_{[-1, n-1]} \\
&\quad (e_{2n} \dots e_{n+1}) \lambda_{\widehat{i(n)}} (e_n e_{n-1} \dots e_1) \lambda_{i_{n+1}} \\
&= \sum_{i_1, i_2, \dots, i_{n+1}} \tau^{-2n} \lambda_{i_{n+1}}^* (e_1 e_2 \dots e_n) (e_{n+1} \dots e_{2n+1}) \lambda_{\widehat{i(n)}}^* e_{[-1, n-1]} \lambda_{\widehat{i(n)}} \\
&\quad (e_{2n} e_{2n-1} \dots e_{n+1}) (e_n e_{n-1} \dots e_1) \lambda_{i_{n+1}} \\
&= \tau^{-2n} \sum_{i_{n+1}} \lambda_{i_{n+1}}^* (e_1 e_2 \dots e_{2n+1}) (e_{2n} e_{2n-1} \dots e_1) \lambda_{i_{n+1}} \\
&\quad \text{[by equation (3.3)]} \\
&= \sum_{i_{n+1}} \lambda_{i_{n+1}}^* e_1 \lambda_{i_{n+1}} \\
&\quad \text{[since } (e_1 e_2 \dots e_{2n+1}) (e_{2n} e_{2n-1} \dots e_1) = \tau^{2n} e_1 \text{]} \\
&= 1.
\end{aligned}$$

Here, the second equation holds as  $\lambda_{\widehat{i(n)}} \in M_{n-1}$  and  $(e_{n+1} e_{n+2} \dots e_{2n+1})$ ,  $(e_{2n} e_{2n-1} \dots e_{n+1})$  both commutes with  $M_{n-1}$ .

Hence the induction is complete.

Now we show property(1) of the Lemma 3.4.

As induction hypothesis, suppose, for  $n \geq 0$ ,

$$e_{[-1, n]} x_n e_{[-1, n]} = E_N(x_n) e_{[-1, n]} \text{ for } x_n \in M_n.$$

It trivially holds for  $n = 0$ . Then, for  $n \geq 0$ , and for  $x_{n+1} \in M_{n+1}$ ,

we get the following array of equations,

$$\begin{aligned}
& e_{[-1,n+1]}x_{n+1}e_{[-1,n+1]} \\
&= \tau^{-2(n+1)}(e_{n+2}\dots e_{2n+3})e_{[-1,n]}(e_{2n+2}\dots e_{n+2}) \\
&\quad x_{n+1}(e_{n+2}\dots e_{2n+3})e_{[-1,n]}(e_{2n+2}\dots e_{n+2}) \\
&= \tau^{-2(n+1)}(e_{n+2}\dots e_{2n+3})e_{[-1,n]}(e_{2n+2}\dots e_{n+3}) \\
&\quad E_{M_n}(x_{n+1})(e_{n+2}\dots e_{2n+3})e_{[-1,n]}(e_{2n+2}\dots e_{n+2}) \\
&= \tau^{-2(n+1)}(e_{n+2}\dots e_{2n+3})e_{[-1,n]}(e_{2n+2}\dots e_{n+3}) \\
&\quad (e_{n+2}\dots e_{2n+3})E_{M_n}(x_{n+1})e_{[-1,n]}(e_{2n+2}\dots e_{n+2}) \\
&= \tau^{-2(n+1)}(e_{n+2}\dots e_{2n+3})e_{[-1,n]}(\tau^n e_{2n+2}e_{2n+3}) \\
&\quad E_{M_n}(x_{n+1})e_{[-1,n]}(e_{2n+2}\dots e_{n+2}) \\
&= \tau^n \tau^{-2(n+1)}(e_{n+2}\dots e_{2n+2})e_{[-1,n]}(e_{2n+3}e_{2n+2}e_{2n+3}) \\
&\quad E_{M_n}(x_{n+1})e_{[-1,n]}(e_{2n+2}\dots e_{n+2}) \\
&= \tau^{-(n+1)}(e_{n+2}\dots e_{2n+3})e_{[-1,n]}E_{M_n}(x_{n+1})e_{[-1,n]}(e_{2n+2}\dots e_{n+2}) \\
&= \tau^{-(n+1)}(e_{n+2}\dots e_{2n+3})E_N(x_{n+1})e_{[-1,n]}(e_{2n+2}\dots e_{n+2}) \\
&\quad \text{[Induction hypothesis]} \\
&= E_N(x_{n+1})e_{[-1,n+1]}.
\end{aligned}$$

The fourth equation holds because of the almost trivial fact that

$$(e_{2n+2}\dots e_{n+3})(e_{n+2}\dots e_{2n+3}) = \tau^n e_{2n+2}e_{2n+3}. \quad (3.4)$$

It should be mentioned that throughout we have used the fact that, for  $n \geq 0$ ,

$$e_{[-1,n+1]} = \tau^{-(n+1)}(e_{n+2}\dots e_{2n+3})e_{[-1,n]}(e_{2n+2}\dots e_{n+2}).$$

This completes the induction.

Now using Lemma 3.4 we get the desired result for  $II_1$  factor Case. For finite dimensional  $C^*$ - algebra the only remaining thing is to prove that the map  $x \mapsto xe_{[-1,n]}$  for  $x \in N$  is injective. From Lemma 2.1 it follows that  $xe_1 = 0$  implies  $x = 0$  for  $x \in N$ , proving the above fact for  $n = 0$ . Suppose the statement is true for  $(n - 1)$ , that is for  $x \in N, xe_{[-1,n-1]} = 0$  implies  $x = 0$ . Let for  $x \in N, xe_{[-1,n]} = 0$ . Thus,  $(\|xe_{[-1,n]}\|_2)^2 = tr(xe_{[-1,n]}x^*) = 0$ . Note,

$$\begin{aligned}
0 &= tr(e_{[-1,n]}x^*x) \\
&= tr((e_{n+1}e_{n+2}\dots e_{2n+1})e_{[-1,n-1]}(e_{2n}\dots e_{n+1})x^*x) \\
&= tr(e_{[-1,n-1]}(e_{2n}\dots e_{n+1})(e_{n+1}\dots e_{2n+1})x^*x) \quad (\text{since } x^*x \in N) \\
&= tr(e_{[-1,n-1]}(\tau^{n-1}e_{2n}e_{2n+1})x^*x). \quad (\text{by equation(3.4)})
\end{aligned}$$

But as we know  $tr$  is Markov, we conclude from the last equation  $tr(e_{[-1,n-1]}x^*x) = 0$ , that is  $tr(xe_{[-1,n-1]}x^*) = 0$ . In other words,  $x e_{[-1,n-1]} = 0$  and now from induction hypothesis we conclude  $x = 0$ . Hence the induction is complete.

This completes the proof for both the Cases. □

## 4 Acknowledgement

I wish to thank V. S. Sunder for many helpful discussions.

## References

- [1] Evans D. E., and Kawahigashi Y., Quantum symmetries on operator algebras, *Oxford Mathematical Monographs* The Clarendon Press, Oxford University Press, New York, 1998, Oxford Science Publications.
- [2] Frederick M. Goodman, Pierre de La Harpe, and Vaughan F.R. Jones(1989), *Coxeter graphs and towers of algebras*, volume 14, Springer-Verlag New York.
- [3] Gantmacher F.R., The theory of matrices, vol. 2, Chelsea, New York, 1959, *Mathematical Reviews (MathSciNet): MR99f*, 15001, 1979.
- [4] Jolissaint P., Index for pairs of finite von Neumann algebras, *Pacific Journal of Mathematics*, **146(1)**(1990) 43-70.
- [5] Jones V.F.R., Index for subfactors, *Inventiones Mathematicae*, **72(1)**(1983) 1-25.
- [6] Jones V.F.R. and Penneys D., The embedding theorem for finite depth subfactor planar algebras, *Quantum Topol.* **2** Issue (3)(2011) 301-337,
- [7] Jones V.F.R. and Sunder, V.S., *Introduction to subfactors*, volume 234. Cambridge University Press, 1997.
- [8] Kawahigashi Y., Automorphisms commuting with a conditional expectation onto a subfactor with finite index, *J. Operator Theory*, **28**(1992) no. 1, 127-145.
- [9] Kosaki H., Extension of Jones' theory on index to arbitrary factors *Journal of Functional Analysis*, **66**(1986) 123-140.

- [10] Loi P. H., On automorphisms of subfactors, *Journal of Functional Analysis*, **141**(2) (1996) 275-293.
- [11] Pimsner M. and Popa S., Entropy and index for subfactors, In *Annales scientifiques de l'Ecole normale supérieure*, **19** (1986) 57-106.
- [12] Pimsner M. and Popa S., Iterating the basic construction, *Transactions of the American Mathematical Society*, **310**(1) (1988) 127–133 .
- [13] Richard D. Burstein, Group-type subfactors and Hadamard matrices, *Trans. Amer. Math. Soc.* **367**(2015) 6783-6807, see also *arXiv preprint arXiv:0811.1265*(2008)
- [14] Svendsen A.L., Automorphisms of subfactors from commuting squares, *Trans. Amer. Math. Soc.*, **356**(2004) no.6 2515-2543.
- [15] Watatani, Y., Index for  $C^*$ -subalgebras, *Mem. Amer. Math. Soc.*, **424**(1990) vi+117 pp.