

On Alternating and Symmetric Groups Which Are Quasi OD-Characterizable

Ali Reza Moghaddamfar

Faculty of Mathematics, K. N. Toosi University of Technology,
P. O. Box 16315-1618, Tehran, Iran.

E-mails: moghadam@ipm.ir and moghadam@kntu.ac.ir

November 7, 2021

Abstract

Let $\Gamma(G)$ be the prime graph associated with a finite group G and $D(G)$ be the degree pattern of G . A finite group G is said to be k -fold OD-characterizable if there exist exactly k non-isomorphic groups H such that $|H| = |G|$ and $D(H) = D(G)$. The purpose of this article is twofold. First, it shows that the symmetric group S_{27} is 38-fold OD-characterizable. Second, it shows that there exist many infinite families of alternating and symmetric groups, $\{A_n\}$ and $\{S_n\}$, which are k -fold OD-characterizable with $k > 3$.

Keywords: OD-characterization, alternating group, symmetric group, prime graph, spectrum, degree pattern, split extension, subdirect product.

1 Introduction

Throughout the article, all the groups under consideration are *finite* and simple groups are *nonabelian*. For a natural number n , we denote by $\pi(n)$ the set of all prime divisors of n and put $\pi(G) = \pi(|G|)$. The *spectrum* $\omega(G)$ of a group G is the set of orders of elements in G . The set $\omega(G)$ determines the *prime graph* $\Gamma(G)$ whose vertex set is $\pi(G)$, and two vertices p and q are *adjacent* if and only if $pq \in \omega(G)$. For two vertices p and q we will write $(p \sim q)_G$ to indicate that p is adjacent to q in $\Gamma(G)$. Denote by $s(G)$ the number of connected components of $\Gamma(G)$ and by $\pi_i(G)$ ($i = 1, 2, \dots, s(G)$), the set of vertices of its i th connected component. If $2 \in \pi(G)$ then we assume that $2 \in \pi_1(G)$. We recall that the set of vertices of connected components of all finite simple groups are obtained in [5] and [14].

As usual, the *degree* $\deg(p)$ of a vertex $p \in \pi(G)$ in $\Gamma(G)$ is the number of edges incident on p . We denote the set of all vertices of the prime graph $\Gamma(G)$

2000 *Mathematics Subject Classification:* 20D05, 20D06, 20D08.

which are joined to all other vertices by $\Lambda(G)$. If the prime divisors of $|G|$ are $p_1 < p_2 < \dots < p_k$, then we define $D(G) := (\deg(p_1), \deg(p_2), \dots, \deg(p_k))$, which is called the *degree pattern* of G .

Given a group H , denote by $h_{\text{OD}}(H)$ the number of isomorphism classes of groups G such that $|G| = |H|$ and $D(G) = D(H)$. Clearly, there are only finitely many isomorphism types of groups of order n , because there are just finitely many ways that an $n \times n$ multiplication table can be filled in. Hence $1 \leq h_{\text{OD}}(H) < \infty$ for any group H . In terms of function h_{OD} , groups H are classified as follows:

Definition 1.1 *Any group H satisfying $h_{\text{OD}}(H) = k$ is said to be k -fold OD-characterizable. Usually, a 1-fold OD-characterizable group is simply called an OD-characterizable group, and it is called quasi OD-characterizable if it is k -fold OD-characterizable for some $k > 1$.*

This article is a continuation of my investigations of the OD-characterizability of alternating and symmetric groups initiated in [4]. We keep the notation created and the conventions made therein.

In a series of articles (see [4], [8], [9], [11] and [17]), it has been shown that many of the alternating and symmetric groups are n -fold OD-characterizable for $n \in \{1, 2, 3, 8\}$. These results are summarized in the following proposition.

Proposition 1.2 *The following statements hold:*

- (a) *The alternating groups A_p, A_{p+1}, A_{p+2} and the symmetric groups S_p and S_{p+1} , where p is a prime number, are OD-characterizable.*
- (b) *The alternating group A_{10} is 2-fold OD-characterizable, while the symmetric group S_{10} is 8-fold OD-characterizable.*
- (c) *The alternating groups A_{p+3} , where $p \neq 7$ is a prime number less than 100, are OD-characterizable.*
- (d) *The symmetric groups S_{p+3} , where $p \neq 7$ is a prime number less than 100, are OD-characterizable or 3-fold OD-characterizable.*

In addition, it was shown in [4, Corollary 1.5] that all alternating groups A_m for which $m \leq 100$, except A_{10} , are OD-characterizable.

Proposition 1.3 *All alternating groups A_m , where m is a natural number less than 100, except A_{10} , are OD-characterizable.*

These observations convinced us to propose the following conjecture in [4]:

Conjecture 1.4 *All alternating groups A_m , with $m \neq 10$, are OD-characterizable.*

On the other hand, in recent years we have not found any simple group S with $h_{\text{OD}}(S) \geq 3$. Therefore, we asked in [4] the following question:

Problem 1.5 *Is there a simple group S with $h_{\text{OD}}(S) \geq 3$?*

Our recent investigations show that Conjecture 1.4 does not hold in general. Recently, the authors showed in [6] that the alternating group A_{125} satisfying $h_{\text{OD}}(A_{125}) \geq 3$ (see also [7]). Here, we will show that there exist infinite families of alternating groups A_m which are k -fold OD-characterizable with $k \geq 3$. We notice that Problem 1.5 is also answered positively through these examples.

Theorem 1.6 *There are infinitely many alternating groups A_m which satisfy $h_{\text{OD}}(A_m) > 1$. In particular, there is no upper bound for $h_{\text{OD}}(A_m)$.*

It is also worth mentioning that a similar description as Proposition 1.3 is exhibited about OD-characterizability of symmetric groups S_m , where m is a natural number less than 100 (see [4, Theorem 1.7]). Nevertheless, in checking the list of such groups, we found out that it contains a mistake (in fact, Proposition 4.1 in [4] asserts erroneously that the symmetric group S_{27} is 3-fold OD-characterizable). Therefore, another result of the present article can be stated as follows:

Theorem 1.7 *The symmetric group S_{27} is 38-fold OD-characterizable.*

Now, we give a revised list of symmetric groups in question.

Corollary 1.8 *All symmetric groups S_m , where m is a natural number less than 100, except $m = 10, 27$, are OD-characterizable or 3-fold OD-characterizable. Moreover, the symmetric group S_{10} is 8-fold OD-characterizable, while the symmetric group S_{27} is 38-fold OD-characterizable.*

The following conjecture involving symmetric groups is posed in [4]:

Conjecture 1.9 *All symmetric groups S_m , with $m \neq 10$, are OD-characterizable or 3-fold OD-characterizable.*

It turns out that a negative answer to this conjecture is provided by symmetric group S_{27} (see also [6, 7]). In addition, we will get many other examples of symmetric groups which are k -fold OD-characterizable with $k > 3$.

Theorem 1.10 *There are infinitely many symmetric groups S_m which satisfy $h_{\text{OD}}(S_m) > 3$. In particular, there is no upper bound for $h_{\text{OD}}(S_m)$.*

We conclude the introduction with some further notation and definitions. Given a natural number m , we denote by l_m the largest prime less than or equal to m and we let $\Delta(m) = m - l_m$. It is clear that $l_m = m$ (or equivalently $\Delta(m) = 0$) if and only if m is a prime number. Note that if $m > 2$, then l_m is always an odd prime, and so $\Delta(m)$ is even iff m is odd. In addition, from the definition, it is easy to see that

$$l_m = l_{m-1} = l_{m-2} = \dots = l_{m-\Delta(m)+1}.$$

We shall use the notation $\nu(m)$ (resp. $\nu_a(n)$) to denote the number of types of groups (resp. abelian groups) of order m . Clearly, $\nu_a(m) \leq \nu(m)$. We also denote the set of partitions of m by $\text{Par}(m)$. It is known that for any prime p , $\nu_a(p^m) = |\text{Par}(m)|$. Finally, we use A_m and S_m to denote the alternating and symmetric group on m letters, respectively. In the case when $p \geq 5$ is a prime, we denote by \mathcal{S}_p the set of all simple groups with prime divisors at most p .

2 Auxiliary results

In this section we give several definitions and auxiliary results to be used later. The first of them is the following definition of *subdirect products*.

Definition 2.1 *Let $n \geq 2$. A subdirect product of the groups G_1, G_2, \dots, G_n is a subgroup $G \leq G_1 \times G_2 \times \dots \times G_n$ of the direct product such that the canonical projections $G \rightarrow G_i$ are surjective for all i .*

One way of obtaining a subdirect product of *two* groups is via the fibre product construction. This is illustrated here for two groups. Given some groups G_1 and G_2 with normal subgroups N_1 and N_2 such that G_1/N_1 and G_2/N_2 are isomorphic, we want to construct a group G having a normal subgroup N isomorphic to $N_1 \times N_2$ such that G/N_2 is isomorphic to G_1 and G/N_1 is isomorphic to G_2 . Notice that we will usually identify $N_1 \times 1$ with N_1 and $1 \times N_2$ with N_2 . To carry out the construction, let π_1 and π_2 be homomorphisms from G_1 and G_2 onto some group Q . Now let

$$G := \{(g_1, g_2) \in G_1 \times G_2 \mid \pi_1(g_1) = \pi_2(g_2)\}.$$

It is easy to check that G constitutes a subgroup of $G_1 \times G_2$, and the projection maps onto the coordinates map G onto G_1 and G_2 , respectively. We call G the *fibre product* associated with π_1 and π_2 (Remak [12] called it the *meromorphic product* of G_1 and G_2 with normal subgroups N_1 and N_2). Also, $N_1 \times N_2$ is a normal subgroup of G and the map π on G defined by $\pi((g_1, g_2)) = \pi_1(g_1)$ maps G onto Q with kernel $N_1 \times N_2$, so $G/(N_1 \times N_2)$ is isomorphic to Q . In fact, we have

$$G_1/N_1 \cong G/(N_1 \times N_2) \cong G_2/N_2.$$

It is a basic observation that every subdirect product of G_1 and G_2 is a fibre product (or a meromorphic product) of these groups.

Information on the adjacency of vertices in the prime graphs associated with alternating and symmetric groups can be found in [13, 16]. Consider the function $S : \mathbb{N} \rightarrow \mathbb{N}$, defined as follows: $S(1) = 1$ and for $m > 1$ with prime factorization $m = \prod_i p_i^{\alpha_i}$, $S(m) = \sum_i p_i^{\alpha_i}$. Then one has [16, Lemma 4]:

Lemma 2.2 *Let m and n be natural numbers. Then there hold:*

- (1) A_n has an element of order m if and only if $S(m) \leq n$ for odd m and $S(m) \leq n - 2$ for even m .
- (2) S_n has an element of order m if and only if $S(m) \leq n$.

The next two corollaries are the adjacency criteria for two vertices of $\Gamma(A_n)$ and $\Gamma(S_n)$, respectively.

Corollary 2.3 *Let $p, q \in \pi(A_n) \setminus \{2\}$. Then $(p \sim q)_{A_n}$ if and only if $p + q \leq n$, while $(p \sim 2)_{A_n}$ if and only if $p + 2 \leq n - 2$.*

Corollary 2.4 *Let $p, q \in \pi(S_n)$. Then $(p \sim q)_{S_n}$ if and only if $p + q \leq n$.*

The Goldbach conjecture says that every even natural number n greater than 4 can be written as the sum of two odd primes. In what follows, we will need a stronger conjecture:

Strong Goldbach Conjecture. *Every even natural number n greater than six can be written as the sum of two distinct odd primes.*

We can now state the connection between the strong Goldbach conjecture and the adjacency of vertices in the prime graph of a symmetric group:

Theorem 2.5 [2, Theorem 3] *The following statements are equivalent:*

- (1) *the strong Goldbach conjecture is true;*
- (2) *for each even $n > 6$, $\Gamma(S_{n-1}) \subsetneq \Gamma(S_n)$.*

Proof. It follows immediately from Lemma 2.2 (2). \square

The coincidence criterion for pairwise nonisomorphic symmetric groups (statement (1) of the Lemma 2.6) is obtained modulo strong Goldbach conjecture.

Lemma 2.6 *Let m and n be natural numbers with $2 \leq m < n$. The prime graphs of symmetric groups S_m and S_n are equal if and only if $m = n - 1$ and one of the following holds:*

- (1) *both n and $n - 2$ are non-prime odd numbers.*
- (2) *$n = 4$ or 6 .*

*Proof.** (\implies) In the case when $n \leq 6$, it is a straightforward verification. In fact, the equality of the prime graphs $\Gamma(S_n)$ and $\Gamma(S_{n-1})$, for $n \in \{4, 6\}$, can be obviously verified using Corollary 2.4. Assume now that $n > 6$. We first claim that $m = n - 1$. If not, then $m < n - 1$ and hence one of the numbers n and $n - 1$ is even. Since $n \geq 7$, it follows from strong Goldbach conjecture that there exist two distinct odd primes p and q with $m \leq n - 2 < p + q \leq n$. Hence p is adjacent to q in $\Gamma(S_n)$, while p is nonadjacent to q in $\Gamma(S_m)$, so these graphs are not equal. This contradiction shows $m = n - 1$ as claimed.

By Theorem 2.5, we may assume that n is an odd number. Assume that $\Gamma(S_n) \neq \Gamma(S_{n-1})$. The sets of vertices are distinct if and only if n is a prime. If n is a composite number, then the sets of vertices are equal, and so the sets of edges should be distinct. Hence there exist primes $p, q \in \pi(S_n)$ with $n - 1 < p + q \leq n$. Then obviously $p + q = n$, but n is odd. The only possible case is $\{p, q\} = \{2, n - 2\}$, and so $n - 2 \in \pi(S_n)$ and $n - 2$ is a prime.

(\impliedby) The conclusion follows immediately from Corollary 2.4. \square

We need the following lemma to find some infinite families of alternating and symmetric groups which are not OD-characterizable.

Lemma 2.7 *Let p be an odd prime number. There are infinitely many natural numbers n such that $\Delta(p^n) > 4$.*

*The idea of proof was borrowed from [18]

Proof. Take n to be an even natural number. Clearly, $p^n - 4$ is always composite in this case, and so we need only consider $p^n - 2$. Let $a = p^2$ and $b = 2$. We will now show that given any positive integers a, b , we can find an infinite number of values for N such that $a^N - b$ is composite. If $a^N - b$ is always composite, we are done. Otherwise, there exists a positive integer k such that $a^k - b = q > a$ is prime. Then for all positive integers m , we have

$$a^{k+(q-1)m} \equiv b \pmod{q},$$

and certainly $a^{k+(q-1)m} - b > q$ for all $m > 0$, so $a^{k+(q-1)m} - b$ must be composite for all positive integers m . \square

A possible generalization of Lemma 2.7 is the following statement, which seems intuitive to be true. Let p be an odd prime. Then, there are infinitely many positive integers n such that

$$p^n - 2, \quad p^n - 4, \quad p^n - 6, \quad \dots, \quad p^n - (p - 1),$$

are composite. Alternatively, this problem can be formulated as follows.

Problem 2.8 *Let p be an odd prime. Do there exist infinitely many positive integers n such that $\Delta(p^n) > p$.*

3 The symmetric group S_{27}

The aim of this section is to find the number of non-isomorphic groups with the same order and degree pattern as the symmetric group S_{27} . Indeed, we will show that there are 38 such groups.

Proof of Theorem 1.7. Let G be a group satisfying the following conditions:

- (1) $|G| = |S_{27}| = 2^{23} \cdot 3^{13} \cdot 5^6 \cdot 7^3 \cdot 11^2 \cdot 13^2 \cdot 17 \cdot 19 \cdot 23$ and
- (2) $D(G) = D(S_{27}) = (8, 8, 7, 7, 5, 5, 4, 4, 2)$.

Under these conditions, we conclude immediately that $\Gamma(G) = \Gamma(S_{27})$ (see also [4, Lemma 2.15]). Letting R be the solvable radical of G , we break the proof into a number of separate lemmas.

Lemma 3.1 *R is a $\{2, 3, 5, 7, 11\}$ -group. In particular, G is nonsolvable.*

Proof. First, we show that R is a $23'$ -group. Assume the contrary. Let $23 \in \pi(R)$ and let x be an element of R of order 23. Put $C = C_G(x)$ and $N = N_G(\langle x \rangle)$. By the structure of $\Gamma(G)$, it follows that C is a $\{2, 3, 23\}$ -group. Since N/C is embedded in $\text{Aut}(\langle x \rangle) \cong \mathbb{Z}_{22}$, N is a $\{2, 3, 11, 23\}$ -group. Using Frattini argument we get $G = RN$, and so $19 \in \pi(R)$. Thus R contains a Hall $\{19, 23\}$ -subgroup L of order $19 \cdot 23$. Clearly, L is cyclic, and so $(19 \sim 23)_G$, which is a contradiction.

Next, we show that R is a q' -group for $q \in \{13, 17, 19\}$. Let $q \in \pi(R)$, $R_q \in \text{Syl}_q(R)$ and $N = N_G(R_q)$. Again, by Frattini argument $G = RN$ and

since R is a $23'$ -group we deduce that 23 divides the order of N . Let L be a subgroup of N of order 23. Since L normalizes R_q , LR_q is a subgroup of order $23 \cdot |R_q|$, which is abelian. But then $(q \sim 23)_G$, which is a contradiction.

Finally, R is a $\{2, 3, 5, 7, 11\}$ -group, and since $R \neq G$, it follows that G is nonsolvable. This completes the proof of lemma. \square

In what follows, we put $\overline{G} = G/R$ and $S = \text{Soc}(\overline{G})$. Clearly, S is a direct product

$$S = P_1 \times P_2 \times \cdots \times P_m,$$

where the P_i are nonabelian simple groups, and we have

$$S \leq \overline{G} \leq \text{Aut}(S).$$

We show now that S is a simple group, or equivalently $m = 1$.

Lemma 3.2 $m = 1$. *In particular, \overline{G} is an almost simple group.*

Proof. By way of contradiction, let $m \geq 2$. In this case 23 does not divide $|S|$, for otherwise $\deg(23) \geq 3$, which is a contradiction. Hence, for every i we have $P_i \in \mathcal{S}_{19}$. Therefore $23 \in \pi(\overline{G}) \subseteq \pi(\text{Aut}(S))$, and so 23 divides the order of $\text{Out}(S)$. But

$$\text{Out}(S) = \text{Out}(Q_1) \times \text{Out}(Q_2) \times \cdots \times \text{Out}(Q_r),$$

where Q_i is a direct product of n_i isomorphic copies of a simple group P_i such that

$$S \cong Q_1 \times Q_2 \times \cdots \times Q_r.$$

Therefore, for some j , 23 divides the order of the outer automorphism group of Q_j of n_j isomorphic simple groups P_j . Since $P_j \in \mathcal{S}_{19}$, it follows that $|\text{Out}(P_j)|$ is not divisible by 23 (see [3, 15]). Moreover, since $\text{Out}(Q_j) = \text{Out}(P_j) \wr S_{n_j}$, it follows that

$$|\text{Out}(Q_j)| = |\text{Out}(P_j)|^{n_j} \cdot (n_j)!.$$

This forces $n_j \geq 23$, and so 2^{46} must divide the order of G , which is a contradiction. Therefore $m = 1$ and $S = P_1$. \square

Lemma 3.3 *There are exactly 38 possibilities for the group G .*

Proof. By Lemma 3.2, we have $S \leq \overline{G} \leq \text{Aut}(S)$, where S is a nonabelian simple group in \mathcal{S}_{23} , and so $\{13, 17, 19, 23\} \cap \pi(\text{Out}(S)) = \emptyset$, (see [3, 15]). Now, it follows from Lemma 3.1 and condition (1) that

$$|S| = 2^a \cdot 3^b \cdot 5^c \cdot 7^d \cdot 11^e \cdot 13^2 \cdot 17 \cdot 19 \cdot 23,$$

where $2 \leq a \leq 23$, $0 \leq b \leq 13$, $0 \leq c \leq 6$, $0 \leq d \leq 3$ and $0 \leq e \leq 2$. Comparing this with the nonabelian simple groups in \mathcal{S}_{23} , we obtain $S \cong A_{26}$ or A_{27} . We refer to [15] for the list of nonabelian simple groups in \mathcal{S}_{23} . In the sequel, we deal with two cases separately.

- (1) $S \cong A_{27}$. In this case, we have $A_{27} \leq G/R \leq S_{27}$. Thus $G/R \cong A_{27}$ or $G/R \cong S_{27}$. If $G/R \cong A_{27}$, then $|R| = 2$. Clearly, $R \leq Z(G)$ and G is a central extension of $R \cong \mathbb{Z}_2$. If G splits over R , then $G \cong \mathbb{Z}_2 \times A_{27}$, otherwise we have $G \cong \mathbb{Z}_2 \cdot A_{27}$ (non-split extension). Next, we assume $G/R \cong S_{27}$. In this case $R = 1$ and so $G \cong S_{27}$. Finally, in the case when $S \cong A_{27}$ there are three possibilities for G .
- (2) $S \cong A_{26}$. In this case, we have $A_{26} \leq G/R \leq S_{26}$, and so $G/R \cong A_{26}$ or $G/R \cong S_{26}$. If $G/R \cong A_{26}$, then $|R| = 54$. We claim that the only possibilities for G are $G = R \times A_{26}$, where R is an arbitrary group of order 54, and $G = Q \times (\mathbb{Z}_2 \cdot A_{26})$ where Q is an arbitrary group of order 27. In the first case, since there are exactly 15 groups of order 54, there are 15 possibilities for G . In the second case, since there are just 5 groups of order 27, there are 5 possibilities for G . To prove our claim, we first observe that the automorphism group of R has order smaller than $|A_{26}|$. Now, let $C = C_G(R)$. Then G/C is isomorphic to a subgroup of $\text{Aut}(R)$, and so $|G/C| < |A_{26}|$. It follows that C is not contained in R , so $CR > R$ and CR is a normal subgroup of G . Since G/R is simple, it follows that $CR = G$. Let $D = C \cap R$. Since $D \leq R$, we see that C centralizes D , and since $D \leq C$, we see that R centralizes D , and thus $D \leq Z(G)$. Now C/D is isomorphic to A_{26} and $D \leq Z(C)$. Since the Schur multiplier of A_{26} has order 2, the only possibilities are that $C' \cap D$ has order 1 or 2. If $|C' \cap D| = 1$, then $C' = A_{26}$ and $C = D \times A_{26}$, and in this case, A_{26} is a direct factor of G . In the remaining case, $C' = 2 \cdot A_{26}$, and in this case, since $|D|$ is 2 times some power of 3, it follows that $2 \cdot A_{26}$ is a direct factor of C and hence of G .

If $G/R \cong S_{26}$, then $|R| = 27$. Actually, we want to find (up to isomorphism) all groups G having a normal subgroup R of order 27 such that G/R is isomorphic to S_{26} . Let $M < G$ have index 2, where M/R is A_{26} . Note that M is unique. Let C be the centralizer of R in M . Then RC is normal in M , and since M/R is simple, there are only two possibilities: $RC = R$ or $RC = M$. We want to show that $RC = M$. Suppose $RC = R$. Now M/C is embedded in $\text{Aut}(R)$ and this group faithfully permutes the 26 non-identity elements of R . This action must be transitive or else M/C is embedded in a direct product of the form $S_a \times S_b$ where $a + b = 26$ and both a and b are positive. But $|M/C| \geq |A_{26}|$ and

$$|S_a \times S_b| = (a!)(b!) < (26!)/2.$$

This proves that the action is transitive, and it follows that R is elementary abelian. Now

$$|\text{Aut}(R)| = |\text{GL}(3, 3)| < |A_{26}| \leq |M/C|.$$

This is a contradiction, so $RC = M$.

Let $Z = Z(R) = R \cap C$. Then C/Z is isomorphic to M/R , which is A_{26} . The Schur multiplier of A_{26} has order 2, and it follows that $C' \cap Z$

is trivial, and thus C' is a normal subgroup of M isomorphic to A_{26} . It follows that M is the direct product of R and C' . Let us write $A = C'$, so A is isomorphic to A_{26} . Note that A is characteristic in M , so A is normal in G . Now $G/R \cong S_{26}$ and G/A has order 54, so G is a subdirect product of $G_1 = S_{26}$ and $G_2 = G/A$. (Note that, every group of order 54 has a normal subgroup of index 2.) Actually, since each of these groups has a unique homomorphism onto the group of order 2, it follows that G can be constructed as follows: let π_1 and π_2 be the homomorphisms from G_1 and G_2 onto the group \mathbb{Z}_2 . Now, we consider the fibre product associated with π_1 and π_2 , that is

$$G = \{(a_1, a_2) \mid \pi_1(a_1) = \pi_2(a_2)\}.$$

Clearly, $A_{26} \times P$, where P is a group of order 27, is a normal subgroup of G and the map π on G defined by

$$\pi((a_1, a_2)) = \pi_1(a_1),$$

maps G onto \mathbb{Z}_2 with kernel $A_{26} \times P$, so $G/(A_{26} \times P)$ is isomorphic to \mathbb{Z}_2 . This gives 15 groups, including all direct products of S_{26} with groups of order 27.

This completes the proof of lemma and Theorem 1.7. \square

4 Non OD-characterizable alternating groups

We start this section with a result of M. A. Zvezdina [18, Theorem] which is concerning simple groups whose prime graphs coincide with the prime graph of an alternating simple group. More precisely, she proved:

Lemma 4.1 [18, Theorem] *Let G be an alternating group A_n , $n \geq 5$, and let S be a finite simple group. Then the prime graphs of G and S coincide if and only if one of the following holds:*

- (a) $G = A_5$, $S = A_6$;
- (b) $G = A_6$, $S = A_5$;
- (c) $G = A_7$, $S \in \{L_2(49), U_4(3)\}$;
- (d) $G = A_9$, $S \in \{J_2, S_6(2), O_8^+(2)\}$;
- (e) $G = A_n$, $S = A_{n-1}$, n is odd, and the numbers n and $n-4$ are composite.

Although the groups in the statement (e) of this lemma have the same prime graph and so the same degree pattern, but they have different orders. In fact, we have

$$|A_n| = |A_{n-1}| \times n.$$

Now, if we can choose the number n such that $\pi(n)$ is contained in the set of vertices of the prime graph $\Gamma(A_n)$ which are joined to all other vertices, then the groups A_n and $A_{n-1} \times H$, where H is an arbitrary group of order n , have the same order and degree pattern. This will enable us to give a positive answer to problem 1.5.

Let G be a finite group satisfying $|G| = |A_m| = m!/2$ and $D(G) = D(A_m)$. By [4, Lemma 2.15], the prime graph $\Gamma(G)$ coincides with $\Gamma(A_m)$. Simply, $\Gamma(G)$ is a graph with vertex set $\pi(G) = \{2, 3, 5, \dots, l_m\}$ in which two distinct vertices r and s are joined by an edge iff $r + s \leq m$ if r and s are odd primes and $r + 2 \leq m - 2$ if $s = 2$.

If $\Delta(m) \leq 2$ and $p = l_m$, then we will deal with the alternating groups A_p , A_{p+1} and A_{p+2} , which are OD-characterizable (see [10, Theorem 1.5]). Therefore, we may consider the alternating groups A_m for which $\Delta(m) \geq 3$.

Referring to the already mentioned fact (Proposition 1.3) that all alternating groups A_m , $10 \neq m \leq 100$, are OD-characterizable, we restrict our attention to the alternating groups A_m where $m > 100$. In what follows, we prove the following result, which shows that there is an infinite family $\{A_m\}$ of alternating groups such that $h_{\text{OD}}(A_m) \geq 3$.

Proposition 4.2 *Let m be an odd number satisfying $\Delta(m) > 4$ and $\pi(m) \subseteq \pi(\Delta(m)!)$. Then $h_{\text{OD}}(A_m) \geq 2$. In particular, if $\Delta(3^n) > 4$ (resp. $\Delta(5^n) > 4$), then $h_{\text{OD}}(A_{3^n}) \geq 16$ (resp. $h_{\text{OD}}(A_{5^n}) \geq 4$).*

Proof. First of all, it follows by Lemma 2.17 in [4] that $\pi(\Delta(m)!) \subseteq \Lambda(A_m)$, this would mean that every vertex in $\pi(\Delta(m)!)$ and so in $\pi(m)$ is adjacent to all other vertices in $\pi(A_m)$. Furthermore, since m is an odd number with $\Delta(m) > 4$, the prime graphs $\Gamma(A_m)$ and $\Gamma(A_{m-1})$ coincide by Lemma 4.1. Now, if H is an arbitrary group of order m , then the groups A_m and $A_{m-1} \times H$ have the same order and degree pattern, and hence

$$h_{\text{OD}}(A_m) \geq 1 + \nu(m) \geq 2.$$

In the case when $\Delta(3^n) > 4$, it is routine to check that $n \geq 7$, and so

$$h_{\text{OD}}(A_{3^n}) \geq 1 + \nu(3^n) \geq 1 + \nu_a(3^n) = 1 + |\text{Par}(n)| \geq 1 + |\text{Par}(7)| = 16.$$

Similarly, if $\Delta(5^n) > 4$, then $n \geq 3$, and we obtain

$$h_{\text{OD}}(A_{5^n}) \geq 1 + \nu(5^n) \geq 1 + \nu_a(5^n) = 1 + |\text{Par}(n)| \geq 1 + |\text{Par}(3)| = 4.$$

The proof is complete. \square

Proof of Theorem 1.6. The result follows immediately from Lemma 2.7 and Proposition 4.2. Note that, the proof of Proposition 4.2 shows that there is no upper bound to $h_{\text{OD}}(A_m)$. \square

By Proposition 4.2, we can find many examples of alternating groups A_m with $h_{\text{OD}}(A_m) \geq 3$. We point out here some of them.

(4.a) *Some alternating groups A_m , $m \leq 1000$, with $h_{\text{OD}}(A_m) \geq 3$.*

In this case, we obtain the following simple groups amongst A_m with $h_{\text{OD}}(A_m) \geq 3$ (see Table 1):

$$A_{125}, A_{147}, A_{189}, A_{539}, A_{625}, A_{875},$$

and for each of these groups, we have (see Table 1):

- (1) $h_{\text{OD}}(A_{125}) \geq 1 + \nu(125) = 6$, (see also [6]),
- (2) $h_{\text{OD}}(A_{147}) \geq 1 + \nu(147) = 7$,
- (3) $h_{\text{OD}}(A_{189}) \geq 1 + \nu(189) = 14$,
- (4) $h_{\text{OD}}(A_{539}) \geq 1 + \nu(539) = 3$,
- (5) $h_{\text{OD}}(A_{625}) \geq 1 + \nu(625) = 16$,
- (6) $h_{\text{OD}}(A_{875}) \geq 1 + \nu(875) = 6$.

Table 1.

m	$m - 4$	l_m	$\Delta(m)$	$\pi(\Delta(m)!)$	$\nu(m)$
$125 = 5^3$	$121 = 11^2$	113	12	2, 3, 5, 7, 11	5
$147 = 3 \cdot 7^2$	$143 = 11 \cdot 13$	139	8	2, 3, 5, 7	6
$189 = 3^3 \cdot 7$	$185 = 5 \cdot 37$	181	8	2, 3, 5, 7	13
$539 = 7^2 \cdot 11$	$535 = 5 \cdot 107$	523	16	2, 3, 5, 7, 11, 13	2
$625 = 5^4$	$621 = 3^3 \cdot 23$	619	6	2, 3, 5	15
$875 = 5^3 \cdot 7$	$871 = 13 \cdot 67$	863	12	2, 3, 5, 7, 11	5

We shall use Lemma 2.7 to show that there are infinite families of alternating groups A_n , with $h_{\text{OD}}(A_n) \geq 3$.

(4.b) *An infinite family of alternating groups A_{3^n} , with $h_{\text{OD}}(A_{3^n}) \geq 136$.*

The existence of infinite number of values n for which $\Delta(3^n) > 4$ is immediate from Lemma 2.7. Indeed, if we take n to be a natural number such that $n \equiv 14 \pmod{144}$, then it follows directly that $\{7, 17\} \subseteq \pi(3^n - 2)$ and similarly $\{5, 19\} \subseteq \pi(3^n - 4)$, so the numbers $3^n - 2$ and $3^n - 4$ are composite. This shows that $\Delta(3^n) \geq 8$, for all $n \equiv 14 \pmod{144}$ (note that, this provides an alternate proof of Lemma 2.7 for $p = 3$). Reasoning as in the proof of preceding Proposition 4.2, we have

$$h_{\text{OD}}(A_{3^n}) \geq 1 + \nu(3^n) \geq 1 + \nu_a(3^n) = 1 + |\text{Par}(n)| \geq 1 + |\text{Par}(14)| = 136,$$

where $\text{Par}(n)$ denotes the set of all partitions of n . In particular, we have $h_{\text{OD}}(A_{3^{14}}) \geq 136$.

(4.c) *An infinite family of alternating groups A_{5^n} , with $h_{\text{OD}}(A_{5^n}) \geq 4$.*

By Lemma 2.7 again, there exist infinitely many values of n for which $\Delta(5^n) > 4$. Now for every such n , we have $h_{\text{OD}}(A_{5^n}) \geq 4$ by Proposition 4.2 (see also [7]).

5 On the symmetric groups S_m with $h_{\text{OD}}(S_m) \geq 4$

In this section we are looking for finite non-isomorphic groups having the same order and degree pattern as a symmetric group. Suppose that G is a finite group satisfying $|G| = |S_m| = m!$ and $D(G) = D(S_m)$, for some natural number m . First of all, we conclude from [4, Lemma 2.15] that the prime graph $\Gamma(G)$ coincides with $\Gamma(S_m)$. Actually, $\Gamma(G)$ is a graph with vertex set $\pi(G) = \{2, 3, 5, \dots, l_m\}$ in which two distinct vertices r and s are joined by an edge iff $r + s \leq m$. In the case when $\Delta(m) \leq 1$, we deal with the symmetric groups S_p and S_{p+1} , which are OD-characterizable by [10, Theorem 1.5]. We now consider the symmetric groups S_m for which $\Delta(m) \geq 2$, that is

$$S_{p+2}, S_{p+3}, S_{p+4}, \dots, S_{p+\Delta(m)},$$

where $p = l_m$. On the other hand, in view of [4, Theorem 1.7] and Theorem 1.7, it follows that all symmetric groups S_m , where $10, 27 \neq m \leq 100$, are 3-fold OD-characterizable, and for this reason we now restrict our attention to the symmetric groups S_m where $m > 100$. Here, we first prove the following general result:

Proposition 5.1 *Let m be an odd number satisfying $\Delta(m) > 4$ and $\pi(m) \subseteq \pi(\Delta(m)!)$. Then $h_{\text{OD}}(S_m) \geq 4$.*

Proof. It follows by Lemma 2.17 in [4] that $\pi(\Delta(m)!) \subseteq \Lambda(S_m)$, this would mean that every vertex in $\pi(\Delta(m)!)$ and so in $\pi(m)$ is adjacent to all other vertices in $\pi(S_m)$. Moreover, since m is an odd number with $\Delta(m) > 4$, the prime graphs $\Gamma(S_m)$, $\Gamma(S_{m-1})$, $\Gamma(A_m)$ and $\Gamma(A_{m-1})$ coincide by Corollary 2.4 and Lemma 4.1 (see also [4, Table 7]). Now, if H and K are two arbitrary groups of order m and $2m$, respectively, then the groups S_m , $\mathbb{Z}_2 \times A_m$, $\mathbb{Z}_2 \cdot A_m$, $S_{m-1} \times H$, $(\mathbb{Z}_2 \times A_{m-1}) \times H$, $(\mathbb{Z}_2 \cdot A_{m-1}) \times H$ and $A_{m-1} \times K$, have the same order and degree pattern, and hence $h_{\text{OD}}(S_m) \geq 4$. The proof is complete. \square

Proof of Theorem 1.10. The result follows immediately from Lemma 2.7 and Proposition 5.1. Note that, the proof of Proposition 5.1 shows that there is no upper bound to $h_{\text{OD}}(S_m)$. \square

Considering Proposition 5.1, we can now find many examples of symmetric groups S_m satisfying $h_{\text{OD}}(S_m) \geq 4$. We point out here some of them.

(5.a) *Some symmetric groups S_m , $m \leq 1000$, with $h_{\text{OD}}(S_m) \geq 4$.*

As before in (4.a), we can obtain the following symmetric groups amongst S_m , which are k -fold OD-characterizable with $k \geq 4$ (see Table 1):

$$S_{125}, S_{147}, S_{189}, S_{539}, S_{625}, S_{875}.$$

The case S_{125} had already been studied in [6].

(5.b) *There is an infinite family of symmetric groups S_{p^n} , with $p \in \{3, 5\}$, such that $h_{\text{OD}}(S_{p^n}) \geq 4$.*

Actually, reasoning as before in (4.b) and (4.c), there are an infinite number of values n for which $\Delta(p^n) > 4$, and the result is now immediate from Proposition 5.1 (see also [7]).

We conclude this paper with some comments for future works:

- (1) Specify the exact value of $h_{\text{OD}}(A_n)$ (resp. $h_{\text{OD}}(S_n)$) for alternating (resp. symmetric) groups of degree n for $n \in \{125, 147, 189, 539, 625, 875\}$.
- (2) It seems that it should be possible to prove that the value of $h_{\text{OD}}(A_n)$ (resp. $h_{\text{OD}}(S_n)$) is bounded by some function of n .

Acknowledgments

The author would like to express his sincere thanks to Prof. I. M. Isaacs for the very helpful comments and suggestions to improve the contents and presentation of the manuscript. The author also would like to thank Professor Andrei V. Zavarnitsine and Professor Robert Styer for their valuable comments and discussions. Finally, he also wishes to thank Dr. A. Mahmoudifar for pointing out references [6, 7] and for carefully reading the manuscript.

References

- [1] M. Akbari and A. R. Moghaddamfar, Simple groups which are 2-fold OD-characterizable, *Bull. Malays. Math. Sci. Soc.*, 35 (1)(2012), 65–77.
- [2] J. Bamberg, G. Cairns and K. D. Kilminster, The crystallographic restriction, permutations, and Goldbach’s conjecture, *Amer. Math. Monthly*, 110(3) (2003), 202–209.
- [3] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker and R. A. Wilson, *Atlas of Finite Groups*. Oxford, Clarendon Press, 1985.
- [4] R. Kogani-Moghaddam and A. R. Moghaddamfar, Groups with the same order and degree pattern, *Sci. China Math.*, 55(4)(2012), 701–720.
- [5] A. S. Kondratév, Prime graph components of finite simple groups. (Russian) *Math. Sb.*, 180(6)(1989), 787–797.
- [6] A. Mahmoudifar and B. Khosravi, On characterization by order and prime graph for alternating groups, *Siberian Math. J.*, 56(1)(2015), 125–131.
- [7] A. Mahmoudifar and B. Khosravi, The answers to a problem and two conjectures about OD-characterization of finite groups, arXiv: 1409.7903v1, 28 September 2015.
- [8] A. R. Moghaddamfar and S. Rahbariyan, More on the OD-characterizability of a finite group, *Algebra Colloq.*, 18(4)(2011), 663–674.

- [9] A. R. Moghaddamfar and A. R. Zokayi, OD-Characterization of alternating and symmetric groups of degrees 16 and 22, *Front. Math. China*, 4(4)(2009), 669–680.
- [10] A. R. Moghaddamfar and A. R. Zokayi, Recognizing finite groups through order and degree pattern, *Algebra Colloq.*, 15 (3)(2008), 449–456.
- [11] A. R. Moghaddamfar, A. R. Zokayi and M. R. Darafsheh, A characterization of finite simple groups by the degrees of vertices of their prime graphs, *Algebra Colloq.*, 12(3)(2005), 431–442.
- [12] R. Remak, Über die Darstellung der endlichen Gruppen als Untergruppen direkter Produkte, *J. Reine Angew. Math.*, Bd 163, Heft 1 (1930), 1–44.
- [13] A. V. Vasilév and E. P. Vdovin, An adjacency criterion in the prime graph of a finite simple group, *Algebra and Logic*, 44(6)(2005), 381–406.
- [14] J. S. Williams, Prime graph components of finite groups, *J. Algebra*, 69(2)(1981), 487–513.
- [15] A. V. Zavarnitsine, Finite simple groups with narrow prime spectrum, *Sib. Elektron. Mat. Izv.*, 6 (2009), 1–12.
- [16] A. V. Zavarnitsin and V. D. Mazurov, Element orders in coverings of symmetric and alternating groups, *Algebra and Logic*, 38(3)(1999), 159–170.
- [17] L. C. Zhang, W. J. Shi, L. L. Wang and C. G. Shao, OD-Characterization of A_{16} , *Journal of Suzhou University* (Natural Science Edition), 24 (2008), 7–10.
- [18] M. A. Zvezdina, On nonabelian simple groups with the same prime graph as an alternating group, *Sib. Math. J.*, 54(1) (2013), 47–55.