

# A Wick-rotatable metric is purely electric

Christer Helleland and Sigbjørn Hervik

Faculty of Science and Technology,  
University of Stavanger,  
N-4036 Stavanger, Norway

christer.helleland@uis.no  
sigbjorn.hervik@uis.no

December 19, 2021

## Abstract

We show that a metric of arbitrary dimension and signature which allows for a standard Wick-rotation to a Riemannian metric necessarily has a purely electric Riemann and Weyl tensor.

## 1 Introduction

In quantum theories a Wick-rotation is a mathematical trick to relate Minkowski space to Euclidean space by a complex analytic extension to imaginary time. This enables us to relate a quantum mechanical problem to a statistical mechanical one relating time to the inverse temperature. This trick is highly successful and is used in a wide area of physics, from statistical and quantum mechanics to Euclidean gravity and exact solutions.

In spite of its success, there is a question about its range of applicability. A question we can ask is: *Given a spacetime, does there exist a Wick-rotation to transform the metric to a Euclidean one?*

Here we will give a partial answer to this question and will give a necessary condition for a Wick-rotation (as defined below) to exist. However, before we prove our main theorem, we need to be a bit more precise with what we mean by a Wick-rotation. Consider a pseudo-Riemannian metric (of arbitrary dimension and signature). We need to allow for more general coordinate transformations than the real diffeomorphisms preserving the metric signature – namely to complex analytic continuations of the real metric [1, 2].

Consider a point  $p$  and a neighbourhood,  $U$ , of  $p$ . Assume this neighbourhood is an analytic neighbourhood and that  $x^\mu$  are coordinates on  $U$  so that  $x^\mu \in \mathbb{R}^n$ . We will adapt the coordinates to the point  $p$  so that  $p$  is at the origin of this coordinate system. Consider now the complexification of  $x^\mu \mapsto x^\mu + iy^\mu = z^\mu \in \mathbb{C}^n$ . This complexification enables us to consider the complex analytic neighbourhood  $U^{\mathbb{C}}$  of  $p$ .

Furthermore, let  $g_{\mu\nu}^{\mathbb{C}}$  be a complex bilinear form (a holomorphic metric)

induced by the analytic extension of the metric:

$$g_{\mu\nu}(x^\rho)dx^\mu dx^\nu \mapsto g_{\mu\nu}^{\mathbb{C}}(z^\rho)dz^\mu dz^\nu.$$

Next, consider a real analytic submanifold containing  $p$ :  $\bar{U} \subset U^{\mathbb{C}}$  with coordinates  $\bar{x}^\mu \in \mathbb{R}^n$ . The imbedding  $\iota : \bar{U} \mapsto U^{\mathbb{C}}$  enables us to pull back the complexified metric  $\mathbf{g}^{\mathbb{C}}$  onto  $\bar{U}$ :

$$\bar{\mathbf{g}} \equiv \iota^* \mathbf{g}^{\mathbb{C}}. \quad (1)$$

In terms of the coordinates  $\bar{x}^\mu$ :  $\bar{\mathbf{g}} = \bar{g}_{\mu\nu}(\bar{x}^\rho)d\bar{x}^\mu d\bar{x}^\nu$ . This bilinear form may or may not be real. However, *if the bilinear form  $\bar{g}_{\mu\nu}(\bar{x}^\rho)d\bar{x}^\mu d\bar{x}^\nu$  is real (and non-degenerate) then we will call it an analytic extension of  $g_{\mu\nu}(x^\rho)dx^\mu dx^\nu$  with respect to  $p$ , or simply a Wick-rotation of the real metric  $g_{\mu\nu}(x^\rho)dx^\mu dx^\nu$* . This clearly generalises the concept of Wick-rotations from the standard Minkowskian setting to a more general setting [3].

In the following, let us call the Wick-rotation, in the sense above, for  $\bar{\phi}$ ; i.e.,  $\bar{\phi} : U \rightarrow \bar{U}$ . We note that this transformation is complex, and we can assume, since  $U$  is real analytic, that  $\bar{\phi}$  is analytic.

The Wick-rotation in the sense above, leaves the point  $p$  stationary. It therefore induces a linear transformation,  $M$ , between the tangent spaces  $T_p U$  and  $T_p \bar{U}$ . The transformation  $M$  is complex and therefore may change the metric signature; consequently, even if the metric  $\bar{g}_{\mu\nu}$  is real, it does not necessarily need to have the same signature of  $g_{\mu\nu}$ .

Consider now the curvature tensors,  $R$  and  $\nabla^{(k)}R$  for  $g_{\mu\nu}$ , and  $\bar{R}$  and  $\bar{\nabla}^{(k)}\bar{R}$  for  $\bar{g}_{\mu\nu}$ . Since both metrics are real, their curvature tensors also have to be real. The analytic continuation, in the sense above, induces a linear transformation of the tangent spaces; consequently, this would relate the Riemann tensors  $R$  and  $\bar{R}$  through a complex linear transformation. It is useful to introduce an orthonormal frame  $\mathbf{e}_\mu$ . The orthonormal frames  $\mathbf{e}_\mu$  and  $\bar{\mathbf{e}}_\mu$  are related through their complexified frame  $\mathbf{e}_\mu^{\mathbb{C}}$ . We can define a complex orthonormal frame requiring the inner product<sup>1</sup>  $\langle \mathbf{e}_\mu^{\mathbb{C}}, \mathbf{e}_\nu^{\mathbb{C}} \rangle = \delta_{\mu\nu}$ . This inner product is invariant under the complex orthogonal transformations,  $O(n, \mathbb{C})$ . The real frames  $\mathbf{e}_\mu$  and  $\bar{\mathbf{e}}_\mu$  are obtained by restricting the complex frame. As an example, consider the standard holomorphic inner product space  $(\mathbb{C}^n, \mathbf{g}_0^{\mathbb{C}})$  and  $(\mathbf{e}_1^{\mathbb{C}}, \dots, \mathbf{e}_n^{\mathbb{C}})$  the standard basis. Then a real subspace is  $V = \text{span}_{\mathbb{R}}(i\mathbf{e}_1^{\mathbb{C}}, \dots, i\mathbf{e}_p^{\mathbb{C}}, \mathbf{e}_{p+1}^{\mathbb{C}}, \dots, \mathbf{e}_n^{\mathbb{C}})$ , and the corresponding metric (obtained from  $\mathbf{g}_0^{\mathbb{C}}$  by restriction) is real. All such real subspaces  $V$  (of different signatures) are obtained from such identifications and hence different real subspaces  $V$  are related via the action of the complex orthogonal group  $O(n, \mathbb{C})$  (for more details, see e.g. [4, 5]).

Hence, we consider the real vector spaces  $T_p U$  and  $T_p \bar{U}$  as embedded in the complexified vector space  $(T_p U)^{\mathbb{C}} \cong (T_p \bar{U})^{\mathbb{C}}$ . The real frames are thus related through a restriction of a complex frame having an  $O(n, \mathbb{C})$  structure group. If moreover the tangent spaces  $T_p U$  and  $T_p \bar{U}$  are embedded:

$$T_p U, T_p \bar{U} \hookrightarrow (T_p U)^{\mathbb{C}} \cong (T_p \bar{U})^{\mathbb{C}},$$

such that they form a *compatible triple*<sup>2</sup>, then we shall say that the real submanifolds:  $U$  and  $\bar{U}$ , are related through a *standard Wick-rotation*. A standard

<sup>1</sup>This is not really a proper inner product since it is not positive definite, but rather a  $\mathbb{C}$ -bilinear non-degenerate form defining a *holomorphic inner product*.

<sup>2</sup>Let  $W$  and  $\bar{W}$  be real slices of a holomorphic inner product space:  $(E, g)$ . Assume they

Wick-rotation allows us to choose commuting Cartan involutions of the real metrics.

Note the special case where  $\bar{U}$  is Riemannian, then the condition of being Wick-rotated by a standard Wick-rotation, is just the condition that the conjugation maps of  $T_p U$ , and  $T_p \bar{U}$  must commute when embedded into  $(T_p U)^\mathbb{C}$ , i.e.  $T_p U$  and  $T_p \bar{U}$  are compatible real forms.

We refer to the manuscript [5], for more details about standard Wick-rotations and the connection with real GIT, and the special case of  $\bar{U}$  being Riemannian.

By using  $\bar{\phi}$  we can relate the metrics  $\mathbf{g} = \bar{\phi}^* \bar{\mathbf{g}}$ . Since the map is analytic (albeit complex), the curvature tensors are also related via  $\bar{\phi}$ . If  $R$  and  $\bar{R}$  are the Riemann curvature tensors for  $U$  and  $\bar{U}$  respectively, then these are related, using an orthonormal frame, via an  $O(n, \mathbb{C})$  transformation. Consider the components of the Riemann tensor as a vector in some  $\mathbb{R}^N \subset \mathbb{C}^N$ . If there exists a Wick-rotation of the metric at  $p$ , then the (real) Riemann curvature tensors of  $U$  and  $\bar{U}$  must be real restrictions of vectors that lie in the same  $O(n, \mathbb{C})$  orbit in  $\mathbb{C}^N$ .

**Note:** This definition of a Wick-rotation does not include the more general analytic continuations defined by Lozanovski [6]. In particular, we consider *one* particular metric (thus not a family of them) and we require that the point  $p$  is fixed and is therefore more of a complex rotation.

In the following we will utilise the study of real orbits of semi-simple groups, see e.g. [7, 8]. In particular, the considerations made in [9] will be useful. For a more general introduction to the structure of Lie algebras including the Cartan involution, see, for example [10, 11].

## 2 The electric/magnetic parts of a tensor

Following [9], we can introduce the electric and magnetic parts of a tensor by considering the eigenvalue decomposition of the tensor under the Cartan involution  $\theta$  of the real Lie algebras  $\mathfrak{o}(p, q)$ . This involution can be extended to all tensors, and to vectors  $\mathbf{v} \in T_p M$  in particular. Considering an orthonormal frame, so that:

$$\mathbf{g}(\mathbf{e}_\mu, \mathbf{e}_\mu) = \begin{cases} -1, & 1 \leq \mu \leq p \\ +1 & p+1 \leq \mu \leq p+q = n, \end{cases}$$

the  $\theta : T_p M \rightarrow T_p M$ , can be defined as the linear operator:

$$\theta(\mathbf{e}_\mu) = \begin{cases} -\mathbf{e}_\mu, & 1 \leq \mu \leq p \\ +\mathbf{e}_\mu & p+1 \leq \mu \leq p+q = n. \end{cases}$$

---

are both real forms of  $W^\mathbb{C} \subset (E, g)$ . Let  $V$  be another real slice of  $E$ , and a real form of  $W^\mathbb{C}$ , with Euclidean signature. Suppose  $W, \widetilde{W}$  and  $V$  are pairwise compatible (i.e. their conjugation maps commute pairwise), then a triple:  $(W, \widetilde{W}, V)$ , will be called a *compatible triple*. **Examples:**  $(\mathbb{R} \oplus i\mathbb{R}, i\mathbb{R} \oplus \mathbb{R}, \mathbb{R}^2)$  with  $E := \mathbb{C}^2$ , and  $(\mathfrak{o}(p, q), \mathfrak{o}(\bar{p}, \bar{q}), \mathfrak{o}(n))$  with  $E := \mathfrak{o}(n, \mathbb{C})$  and  $g := \kappa(-, -)$  (the Killing form).

Clearly, this implies that the bilinear map:

$$\langle X, Y \rangle_\theta := \mathbf{g}(\theta(X), Y), \quad X, Y \in T_p M$$

defines a positive definite inner-product on  $T_p M$ . This Cartan involution can be extended tensorially to arbitrary tensor products.

Given a Cartan involution  $\theta$ , then since  $\theta^2 = \text{Id}$ , its eigenvalues are  $\pm 1$  and any tensor  $T$  has an eigenvalue decomposition:

$$T = T_+ + T_-, \quad \text{where } \theta(T_\pm) = \pm T_\pm.$$

A space is called *purely electric* (PE) if there exists a Cartan involution so that the Weyl tensor decomposes as  $C = C_+$  [9]. Furthermore, a space is called *purely magnetic* (PM) if the Weyl tensor decomposes as  $C = C_-$ . If this property occurs also for the Riemann tensor, we call the space Riemann purely electric (RPE) or magnetic (RPM), respectively. Clearly, RPE implies PE.

### 3 The Riemann curvature operator

The Riemann curvature tensor can (pointwise) be seen as a bivector operator:

$$\text{Riem} : \wedge^2 \Omega_p(M) \rightarrow \wedge^2 \Omega_p(M).$$

In a pseudo-Riemannian space of signature  $(p, q)$  the metric  $\mathbf{g}$  will provide an isomorphism between the space of bivectors,  $\wedge^2 \Omega_p(M)$ , and the Lie algebra  $\mathfrak{g} = \mathfrak{o}(p, q)$ . This can be seen as follows. The Lie algebra  $\mathfrak{o}(p, q)$  is defined through the action of  $O(p, q)$  on the tangent space  $T_p M$ : For any  $G \in O(p, q)$ ,  $G : T_p M \rightarrow T_p M$  so that  $\mathbf{g}(G \cdot v, G \cdot u) = \mathbf{g}(v, u)$  for all  $v, u \in T_p M$ . Using the exponential map  $\exp : \mathfrak{o}(p, q) \rightarrow O(p, q)$ , we get the requirement that  $\mathbf{g}(X(v), u) + \mathbf{g}(v, X(u)) = 0$  for any  $X \in \mathfrak{o}(p, q)$ . Consequently,  $X$  is antisymmetric with respect to the metric  $\mathbf{g}$ . In terms of the basis vectors, we can write  $X = (X^\mu_\nu)$  and the antisymmetry condition implies that by raising an index we get  $X^{\mu\nu} = -X^{\nu\mu}$  and can therefore be considered as a bivector. Since the dimensions match, the metric thus provides with an isomorphism between the Lie algebra  $\mathfrak{o}(p, q)$  and the space of bivectors  $\wedge^2 \Omega_p(M)$  at a point<sup>3</sup>.

Consequently, the Riemann curvature operator can also be viewed as an endomorphism of  $V := \mathfrak{g}$  treated as a vector space. Consider therefore any  $R \in \text{End}(V)$ :

$$R : V \rightarrow V.$$

This endomorphism can be split in a symmetric and anti-symmetric part,  $R = S + A$ , with respect to the metric induced by  $\mathbf{g}$  (which we also will call  $\mathbf{g}$  and is proportional to the Killing form  $\kappa$  on  $V$ )<sup>4</sup>:

$$\mathbf{g}(S(x), y) = \mathbf{g}(S(y), x), \quad \mathbf{g}(A(x), y) = -\mathbf{g}(A(y), x) \quad \forall x, y \in \mathfrak{g}.$$

<sup>3</sup>Indeed, this is a mere consequence of the fact that there is an  $O(p, q)$ -module isomorphism between  $T_p M$  and  $T_p^* M$ .

<sup>4</sup>That the metric induced by  $\mathbf{g}$  is proportional to the Killing form can be seen either by explicit computation, or from considering  $\kappa$  as a even-ranked tensor over  $V^* \otimes V^*$  which is invariant under the action of  $O(p, q)$ . By, e.g., section 5.3.2 in [12], this tensor is necessarily proportional to the metric tensor on  $V$  induced by  $\mathbf{g}$ .

This metric is invariant under the Lie group action of  $G = O(p, q)$ :

$$\mathbf{g}(h \cdot x, h \cdot y) = \mathbf{g}(x, y),$$

where  $h \cdot x$  is the natural Lie group action on the Lie algebra given by the adjoint:  $h \cdot x := \text{Ad}_h(x) = h^{-1}xh$ .

Consider now a Cartan involution  $\theta : \mathfrak{g} \rightarrow \mathfrak{g}$ . Then we define the inner-product on  $V = \mathfrak{g}$  as follows:

$$\langle x, y \rangle_\theta = \mathbf{g}(\theta(x), y),$$

which is just proportional to  $\kappa_\theta(-, -) := -\kappa(-, \theta(-))$ . We can now, similarly, split any  $R \in \text{End}(V)$  in a symmetric and anti-symmetric part,  $R = R_+ + R_-$ , with respect to the inner-product  $\langle -, - \rangle_\theta$ :

$$\langle R_+(x), y \rangle_\theta = \langle R_+(y), x \rangle_\theta, \quad \langle R_-(x), y \rangle_\theta = -\langle R_-(y), x \rangle_\theta, \quad \forall x, y \in \mathfrak{g}.$$

We shall denote  $V = \mathfrak{t} \oplus \mathfrak{p}$ , for the Cartan decomposition w.r.t  $\theta$ , where  $\mathfrak{t}$  is the compact part and  $\mathfrak{p}$  is the non-compact part.

Suppose now that the real submanifolds  $U$  and  $\bar{U}$  are two Wick-rotatable spaces (of the same dimension) by a standard Wick-rotation at a fixed intersection point  $p$ , but with one of the real slices being Riemannian. So we can set  $V := \mathfrak{o}(p, q)$  as before, and introduce (similarly as with  $V$  above),  $\tilde{V} := \mathfrak{o}(n)$ , a compact real form of  $V^\mathbb{C} := \mathfrak{o}(n, \mathbb{C})$ . These real forms  $V$  and  $\tilde{V}$ , will naturally be compatible when embedded into  $V^\mathbb{C}$ , w.r.t to a standard Wick-rotation, i.e it lets us fix a Cartan involution  $\theta$ , such that  $\mathfrak{t} = V \cap \tilde{V}$ , and  $\mathfrak{p} = V \cap i\tilde{V}$ . Again we refer to the paper [5] for details.

The space of endomorphisms,  $\text{End}(V)$ , is also a vector space with the group action given by conjugation:

$$(g \cdot X)(v) := gX(g^{-1}vg)g^{-1}, \quad X \in \text{End}(V), \quad v \in V, \quad g \in G.$$

Call this action  $\rho$ . We can thus define  $\mathcal{V} := \text{End}(V)$ , and extend the Cartan involution,  $\theta$ , as well as  $\mathbf{g}$  tensorially to  $\mathcal{V}$ . We define analogously an inner product on  $\mathcal{V}$ :

$$\langle\langle X, Y \rangle\rangle_\theta = \mathbf{g}(\theta(X), Y), \quad X, Y \in \mathcal{V}.$$

The inner product can assume to have the following properties (see [7]) w.r.t the action  $\rho$ :

1. The inner product is  $K$ -invariant, where  $K \cong O(p) \times O(q)$  is the maximally compact subgroup of  $G$  with Lie algebra  $\mathfrak{t}$ .
2.  $d\rho(\mathfrak{t}) : \mathcal{V} \rightarrow \mathcal{V}$  consists of skew-symmetric maps w.r.t  $\langle\langle X, Y \rangle\rangle_\theta$ .
3.  $d\rho(\mathfrak{p}) : \mathcal{V} \rightarrow \mathcal{V}$  consists of symmetric maps w.r.t  $\langle\langle X, Y \rangle\rangle_\theta$ .

With such an inner product, enables us to apply the results in [7], i.e we can make use of minimal vectors for determining the closure of real orbits.

Defining  $\tilde{\mathcal{V}} := \text{End}(\tilde{V})$  similarly, we have  $\mathcal{V}, \tilde{\mathcal{V}} \subset \mathcal{V}^\mathbb{C}$  where  $\mathcal{V}^\mathbb{C} := \text{End}(V^\mathbb{C})$ . Now since  $V$  and  $\tilde{V}$  are real forms of  $V^\mathbb{C}$  then  $\mathcal{V}$  and  $\tilde{\mathcal{V}}$  are real forms of  $\mathcal{V}^\mathbb{C}$ .

This is seen in the following way. A map  $R \in \mathcal{V}$  can be extended to the complex linear map  $R^{\mathbb{C}} \in \mathcal{V}^{\mathbb{C}}$  by defining:

$$R^{\mathbb{C}}(x + iy) := R(x) + iR(y), \quad x, y \in V.$$

So we view a map  $R$  as the complex linear map  $R^{\mathbb{C}}$ . Thus regard  $\tilde{\mathcal{V}}$  like this as well. We shall just write  $R$  instead of  $R^{\mathbb{C}}$ .

We thus assume we have two endomorphisms (the Riemann curvature operators):  $R : V \rightarrow V$  (arbitrary pseudo-Riemannian), and  $\tilde{R} : \tilde{V} \rightarrow \tilde{V}$  (Riemannian). Now since we have the two real slices:  $U$  and  $\bar{U}$ , which are Wick-rotated at the point  $p$ , then necessarily  $R \in \mathcal{V}$  and  $\tilde{R} \in \tilde{\mathcal{V}}$  must be conjugated by an element  $g \in G^{\mathbb{C}} := O(n, \mathbb{C})$ .

Set now  $G := O(p, q)$  (with Lie algebra  $V := \mathfrak{o}(p, q)$ ) and  $\tilde{G} := O(n)$  (with Lie algebra  $\tilde{V} := \mathfrak{o}(n)$ ) for the real forms embedded into  $G^{\mathbb{C}}$  (with Lie algebra  $V^{\mathbb{C}} := \mathfrak{o}(n, \mathbb{C})$ ) w.r.t a standard Wick-rotation<sup>5</sup>. Now we have a commutative diagram of conjugation actions:

$$\begin{array}{ccc} G^{\mathbb{C}} & \xrightarrow{\rho^{\mathbb{C}}} & GL(\mathcal{V}^{\mathbb{C}}) \\ i \uparrow & & i \uparrow \\ G & \xrightarrow{\rho} & GL(\mathcal{V}) \end{array} \quad (2)$$

Where  $\rho^{\mathbb{C}}$  is also the action given by conjugation, where  $G^{\mathbb{C}}$  is viewed as a real Lie group, and  $\mathcal{V}^{\mathbb{C}}$  is also viewed as a real vector space. We similarly have such a diagram for the the group  $\tilde{G}$ , where the conjugation action:  $\tilde{\rho}$ , on  $\tilde{\mathcal{V}}$  also extends to  $\rho^{\mathbb{C}}$ .

Now our real Riemann curvature operators from  $U$  and  $\bar{U}$ :  $R$  and  $\tilde{R}$ , will now lie in the same complex orbit, i.e  $G^{\mathbb{C}} \cdot R = G^{\mathbb{C}} \cdot \tilde{R}$ .

So therefore in what follows, we will consider the real orbits,  $G \cdot R$ ,  $\tilde{G} \cdot \tilde{R}$  and its complexified orbit  $G^{\mathbb{C}} \cdot R$  defined by the conjugation action of the group on an endomorphism:  $R \in \mathcal{V}$  and  $\tilde{R} \in \tilde{\mathcal{V}}$ , as follows [7, 8, 9]

$$\begin{aligned} G \cdot R &:= \{h \cdot R \mid h \in O(p, q)\} \subset \mathcal{V} \\ \tilde{G} \cdot \tilde{R} &:= \{h \cdot \tilde{R} \mid h \in O(n)\} \subset \tilde{\mathcal{V}} \\ G^{\mathbb{C}} \cdot R &:= \{h \cdot R \mid h \in O(n, \mathbb{C})\} \subset \mathcal{V}^{\mathbb{C}}. \end{aligned}$$

**Theorem 3.1.** *Suppose  $R = S + A \in \mathcal{V}$  where  $S, A$  are the symmetric/antisymmetric parts w.r.t  $\mathbf{g}$  respectively. Assume that there exists a (real)  $\tilde{R} \in G^{\mathbb{C}} \cdot R$  so that  $\tilde{R} \in \tilde{\mathcal{V}}$  i.e we assume:  $G^{\mathbb{C}} \cdot R = G^{\mathbb{C}} \cdot \tilde{R}$ . Then there exists a Cartan involution  $\theta'$  of  $V$  such that  $R_+ = S$  and  $R_- = A$ , where  $R_+, R_-$  are the symmetric/antisymmetric parts w.r.t  $\langle -, - \rangle_{\theta'}$  respectively.*

*Proof.* Consider the orbits  $G \cdot R$  and  $\tilde{G} \cdot \tilde{R}$ . Since the group  $\tilde{G}$  is compact, the orbit  $\tilde{G} \cdot \tilde{R}$  is necessarily closed in  $\tilde{\mathcal{V}}$ ; consequently,  $G \cdot R$ , is closed as well

<sup>5</sup> $G$  and  $\tilde{G}$  are the structure groups of the real metrics restricting from the holomorphic metric, and thus consist of isometries:  $T_p U \rightarrow T_p U$  and  $T_p \bar{U} \rightarrow T_p \bar{U}$  of the real metrics respectively. These groups are naturally embedded into  $O(n, \mathbb{C})$  as real forms, by complexification:  $f \mapsto f^{\mathbb{C}}$ .

and possesses a minimal vector<sup>6</sup> [7]. Denote by  $\mathcal{M}(G^{\mathbb{C}}, \mathcal{V}^{\mathbb{C}})$  the set of minimal vectors in  $\mathcal{V}^{\mathbb{C}}$ . Assume that  $X \in G \cdot \mathbb{R} \subset \mathcal{V}$  is minimal, then  $X$  is also a minimal vector in the complex orbit:  $G^{\mathbb{C}} \cdot \mathbb{R}$ . However since  $G$  and  $\tilde{G}$  are compatible real forms (i.e  $V$  and  $\tilde{V}$  are compatible<sup>7</sup>), and  $\tilde{G}$  is a compact real form of  $G^{\mathbb{C}}$ , then necessarily:

$$G^{\mathbb{C}} \cdot \mathbb{R} \cap \mathcal{M}(G^{\mathbb{C}}, \mathcal{V}^{\mathbb{C}}) = \tilde{G} \cdot \tilde{\mathbb{R}} \subset \tilde{\mathcal{V}},$$

so we deduce that  $X \in G \cdot \mathbb{R} \cap \tilde{G} \cdot \tilde{\mathbb{R}} \subset \mathcal{V} \cap \tilde{\mathcal{V}}$ .

Now we can choose  $g \in G$  such that  $g \cdot \mathbb{R} = X$ , hence we can conjugate our fixed Cartan involution  $\theta$  using  $g$ , and therefore work with  $\mathbb{R}$  instead of  $X$ . Thus we may assume w.l.o.g that  $X := \mathbb{R}$ . Now  $\mathbb{R}$  leaves invariant both  $V$  and  $\tilde{V}$ , in particular implying that:

$$\mathbb{R}(V \cap \tilde{V}) \subset V \cap \tilde{V} \quad \text{and} \quad \mathbb{R}(V \cap i\tilde{V}) \subset V \cap i\tilde{V}.$$

However again by the compatibility of  $V$  and  $\tilde{V}$  in  $V^{\mathbb{C}}$ , we know that  $V \cap \tilde{V} = \mathfrak{t}$  and  $V \cap i\tilde{V} = \mathfrak{p}$  are the compact/non-compact parts respectively w.r.t our fixed Cartan involution  $\theta$ . So  $\mathbb{R}$  and  $\theta$  commute:  $[\mathbb{R}, \theta] = 0$ , which immediately implies that  $\mathbb{R}_+ = \mathbb{S}$  and  $\mathbb{R}_- = \mathbb{A}$  w.r.t  $\theta$  as required. The theorem is proved.  $\square$

In the case of the Riemann tensor, this is symmetric as a bivector operator with respect to the metric, so we have  $\mathbb{R} = \mathbb{S}$ , consequently, we get the immediate corollary:

**Corollary 3.2.** *A metric (of arbitrary dimension and signature) allowing for a standard Wick-rotation at a point  $p$  to a Riemannian metric, has a purely electric Riemann tensor, and is consequently purely electric, at  $p$ .*

We note that this result applies for a general classes of Wick-rotatable metrics. For example, by complexification of the Lie algebras, it is possible to include Wick-rotations between all of the spaces: de Sitter (dS), anti-de Sitter (AdS), the Riemannian sphere ( $S^n$ ), and hyperbolic space, ( $H^n$ ). These are all group quotients  $G/H$  of different groups  $G$  and  $H$ . This seems at first sight paradoxical since these have different signs of the curvature. Thus if  $\mathbb{R} = g^{-1} \cdot \tilde{\mathbb{R}}$  as claimed in the proof, they would necessarily have the same Ricci scalar<sup>8</sup>. To understand this we first note that when we Wick-rotate to a Riemannian space we may risk to get either a positive definite metric,  $\mathbf{g}(v, v) \geq 0$ , or a negative definite metric,  $\mathbf{g}(v, v) \leq 0$ . The overall sign is conventional and we say that switching the sign using the "anti-isometry",  $\mathbf{g} \mapsto -\mathbf{g}$  is a matter of convention. Note that this switch of the metric gives the same metric for the metric induced by  $\mathbf{g}$  on the Lie algebra.

Consider the simple example of the complex holomorphic metric

$$\mathbf{g}_{\mathbb{C}} = \frac{1}{(1 + z_1^2 + \dots + z_n^2)^2} [dz_1^2 + \dots + dz_n^2] \quad (3)$$

<sup>6</sup>A vector  $X \in V$  is minimal if the norm function  $\| - \| := \sqrt{\langle -, - \rangle_{\theta}}$  along an orbit attains a minimum at  $X$ ; i.e.,  $\|X\| \leq \|h \cdot X\|, \forall h \in G$ .

<sup>7</sup>The conjugation maps of  $V$  and  $\tilde{V}$  in  $V^{\mathbb{C}}$  commute:  $\sigma : V^{\mathbb{C}} \rightarrow V^{\mathbb{C}}$  and  $\tilde{\sigma} : V^{\mathbb{C}} \rightarrow V^{\mathbb{C}}$ , with  $[\sigma, \tilde{\sigma}] = 0$ .

<sup>8</sup>The Riemann endomorphism has components related to the Riemann tensor in  $T_p M \otimes T_p^* M \otimes (T_p M \otimes T_p^* M)^*$ , i.e.,  $R^{\alpha}_{\beta\gamma}{}^{\delta}$ . Thus the Ricci scalar is obtained by taking the double trace showing the Ricci scalar is the same after Wick-rotating.

Locally, the two real slices  $(z_1, \dots, z_n) = (x_1, \dots, x_n)$ , and  $(z_1, \dots, z_n) = (iy_1, \dots, iy_n)$ , give a neighbourhood of  $S^n$  and  $H^n$  respectively. However, note that for hyperbolic space, the induced metric has the "wrong" sign (it is negative definite). Therefore, considering for example the Ricci tensor (by lowering indices appropriately), we get  $R_{\mu\nu} = \lambda g_{\mu\nu}$ ,  $\lambda > 0$ , for both real slices, and the sign of the curvature is encapsured in whether the metric is positive or negative definite.

## 4 Discussion

Using techniques from real invariant theory we have considered a class of metrics allowing for a complex Wick-rotation to a Riemannian space. We have showed that these necessarily are resstricted, in particular, they are purely electric. The result is independent of dimension and signature and shows that if such a Wick rotation is allowable, then we necessarily restrict ourselves to classes of spaces where the "magnetic" degrees of freedom have to vanish (at the point  $p$ ).

There are many examples of purely electric spaces (see [9, 6] and references therein). In particular, a purely electric Lorentzian spacetime is of type G,  $I_i$ , D or O [9]. Thus spacetimes not of these types provide with examples of spaces where such a Wick rotation is not allowed. Non-Wick-rotatable metrics include the classes of Kundt metrics [13] in Lorentzian geometry, and the Walker metrics [14] of more general signature. Also the metrics considered in [15] are in general non-Wick-rotatable metrics. Note that the plane-wave metrics are non-Wick-rotatable metrics.

These results have profound consequences for quantization frameworks where such Wick-rotation is used, since they give a clear restriction of the class of metrics that allows for such a Wick rotation. Clearly, also in the context of quantum gravity, the (real) gravitational degrees of freedom will be restricted by assuming the existence of such a Wick-rotation.

It is worth mentioning that there are quantization procedures which work in the Lorentzian signature all the way through, in particular, there is the algebraic approach to QFT on curved spacetime [16, 17]. For details on renormalization in Lorentzian signature (without Wick rotation), see e.g., [18].

## Acknowledgements

This work was in part supported through the Research Council of Norway, Toppforsk grant no. 250367: *Pseudo-Riemannian Geometry and Polynomial Curvature Invariants: Classification, Characterisation and Applications*.

## References

- [1] C. B.G. McIntosh, M.S. Hickman *Gen.Rel.Grav.*,**17**,111-132 (1985); G.S. Hall, M.S. Hickman, C.B.G. McIntosh, *Gen.Rel.Grav.*,**17**,475-491 (1985); M.S. Hickman, C.B.G. McIntosh, *Gen.Rel.Grav.*,**18**,107-136 (1986); M.S. Hickman, C.B.G. McIntosh, *Gen.Rel.Grav.*,**18**,1275-1290 (1986); C.B.G. McIntosh, M.S. Hickman, A.W.-C. Lun, *Gen.Rel.Grav.*, **20**, 647-657 (1988)

- 
- [2] H. Stephani, D. Kramer, M. MacCallum, C. Hoenselaers, E. Herlt, *Exact solutions of Einsteins field equations*, Cambridge University Press (2003).
  - [3] S. Hervik and A. Coley, 2010, *Class. Quant. Grav.* **27**, 095014 [arXiv:1002.0505]; A. Coley and S. Hervik, 2009, *Class. Quant. Grav.* **27**, 015002 [arXiv:0909.1160].
  - [4] V. Pessers and J. van der Veken, arxiv:1503.07354 [math.DG] .
  - [5] C. Helleland and S. Hervik, *Wick-rotations and real GIT*, arxiv:1703.04576 [math.DG]
  - [6] C. Lozanovski, *Gen. Rel. Grav.* **46**, 1716 (2014).
  - [7] R.W. Richardson and P.J. Slodowy, 1990, *J. London Math. Soc.* (2) **42**: 409-429.
  - [8] P. Eberlein, M. Jablonski, *Contemp. Math.* **491**: 283 (2009).
  - [9] S. Hervik, M. Ortaggio and L. Wylleman, *Class. Quant. Grav.* **30**, 165014 (2013) [arXiv:1203.3563 [gr-qc]].
  - [10] S. Helgason, *Differential Geometry, Lie Groups and Symmetric Spaces*, AMS, 1978.
  - [11] A.W. Knap, *Lie groups beyond an introduction*, Birkhauser, 2005.
  - [12] R. Goodman and N.R. Wallach, *Symmetry, Representations and Invariants*, Springer, 2009.
  - [13] A. Coley, S. Hervik, G. Papadopoulos and N. Pelavas, 2009, *Class. Quant. Grav.* **26**, 105016 [arXiv:0901.0394];
  - [14] A.G. Walker, 1949, *Quart. J. Math. (Oxford)*, **20**, 135-45; A.G. Walker, 1950, *Quart. J. Math. (Oxford)* (2), **1**, 69-79
  - [15] S. Hervik, A. Haarr and K. Yamamoto, arXiv:1410.4347 [math-ph].
  - [16] R. Brunetti, K. Fredenhagen, and R. Verch, *Comm. Math. Phys.* **237** (2003), no. 1-2, 3168.
  - [17] S. Hollands and R. M. Wald, *Comm. Math. Phys.* **223** (2001), no. 2, 289326.
  - [18] R. Brunetti, K. Fredenhagen, *Comm. Math. Phys.* **208** (2000), no. 3, 623661.