

**GEOMETRIC CONVERGENCE OF
THE KÄHLER-RICCI FLOW ON
COMPLEX SURFACES OF GENERAL TYPE**

BIN GUO, JIAN SONG, AND BEN WEINKOVE

ABSTRACT. We show that on smooth minimal surfaces of general type, the Kähler-Ricci flow starting at any initial Kähler metric converges in the Gromov-Hausdorff sense to a Kähler-Einstein orbifold surface. In particular, the diameter of the evolving metrics is uniformly bounded for all time and the Kähler-Ricci flow contracts all the holomorphic spheres with (-2) self-intersection number to isolated orbifold points. Our estimates do not require a priori the existence of an orbifold Kähler-Einstein metric on the canonical model.

1. INTRODUCTION

The Ricci flow, introduced by Hamilton [H], has become a powerful tool to study the topology and geometric structures of Riemannian manifolds. In general, the Ricci flow develops finite time singularities. Hamilton's program of *Ricci flow with surgeries* was carried out by Perelman [P1, P2, P3] to prove Thurston's geometrization conjecture. The minimal model theory in birational geometry can be viewed as the complex analogue of Thurston's geometrization conjecture. It was proposed in [SoT3] that the Ricci flow will carry out an *analytic minimal model program* on projective varieties, analogous to the 3-dimensional Ricci flow with surgeries.

In particular, it was conjectured in [SoT3] that the Ricci flow will perform a canonical surgery, corresponding to a birational transformation such as a divisorial contraction or flip, whenever it encounters a non-volume-collapsing singularity at finite time (see also [T]). This extended earlier conjectures of [FIK], which were established in [SW1, CT]. The theory of the weak Kähler-Ricci flow was developed in [SoT3], where it was shown that the flow can be continued through these birational transformations at the level of potentials. These surgeries are conjectured [SoT3] to be continuous in the Gromov-Hausdorff topology, and this was confirmed for Kähler surfaces [SW2, SW3, SW4]. Indeed it was shown that if the total volume is uniformly bounded below away from zero at a finite time singularity, then the Kähler-Ricci flow converges smoothly outside finitely many distinct holomorphic spheres with self-intersection number (-1) , and these (-1) curves are contracted to distinct points in the Gromov-Hausdorff sense. Furthermore, making use of the work of [SoT3], it was shown that the flow can be extended on the blown-down surfaces, continuously in the Gromov-Hausdorff topology, and

Research supported in part by NSF grants DMS-1406124 and DMS-1406164.

uniquely at the level of potentials. These geometric blow-down procedures terminate in finite time. Higher dimensional metric surgeries via the Kähler-Ricci flow are constructed for certain families of projective manifolds in [SY, S] using a construction of birational cobordism in [W].

Geometric convergence of the Kähler-Ricci flow at infinite time is still largely open. It is conjectured in [SoT3] that if the Kähler-Ricci flow exists for all time, the normalized solution will either converge to a Kähler-Einstein metric space with possible singularities (see [EGZ]) or to a twisted Kähler-Einstein metric space of lower dimension, in the Gromov-Hausdorff topology. In particular, such spaces are expected to be homeomorphic to the canonical models of the original manifolds. This is known to hold if the manifold X admits a Kähler-Einstein metric [C] or if there is a holomorphic submersion $\pi : X \rightarrow B$ with $c_1(B) < 0$ and smooth Calabi-Yau fibers (established in [TWY], building on earlier work of [SoT1, SW3, GTZ, FZ, G]). For more general Kähler manifolds for which the flow exists for all time, weak analytic convergence is proved in [Ts, TZo] when X is of general type and in [SoT1, SoT2] when the flow collapses at infinity. In both cases, the scalar curvature of the normalized solutions of the Kähler-Ricci flow is always uniformly bounded [Z2, SoT4]. However, the problem of global geometric convergence is still open. One of the major obstacles is to obtain a uniform diameter bound for the normalized Kähler-Ricci flow.

In this paper, we will consider minimal surfaces of general type. By definition, a minimal surface of general type is a smooth complex surface X whose canonical bundle K_X is big and nef. Equivalently, there exists a smooth closed $(1, 1)$ -form χ in the first Chern class $c_1(X)$ such that $\chi \geq 0$ and $\int_X \chi^2 > 0$. In particular, X is projective and so it is Kähler. Since there are no (-1) curves on such surfaces, the Kähler-Ricci flow has long time existence and it is proved in [TZo] that the normalized solution will converge weakly to the pullback of the unique orbifold Kähler-Einstein metric on the canonical model of X . More precisely, we will consider the pluricanonical system of X . For sufficiently large m , the holomorphic sections of K_X^m induce a holomorphic map $\Phi : X \rightarrow \mathbb{P}^N$ with image X_{can} , the canonical model of X . X_{can} is an algebraic surface with, at worst, finitely many orbifold A - D - E -singularities (cf. [BHPV]). Moreover, X_{can} admits a unique orbifold Kähler-Einstein metric [Kob] because $K_{X_{\text{can}}}$ is ample, which we will denote by ω_{KE} . The map Φ contracts (-2) curves on X to orbifold points on X_{can} .

We consider a solution $\omega(t)$ to the normalized Kähler-Ricci flow on X starting at any Kähler metric ω_0 . Namely, $\omega = \omega(t)$ solves

$$(1.1) \quad \frac{\partial}{\partial t} \omega = -\text{Ric}(\omega) - \omega, \quad \omega|_{t=0} = \omega_0.$$

A unique solution to (1.1) exists for all time [C, Ts]. It is known by the work of Tsuji [Ts] and Tian-Zhang [TZo] that as $t \rightarrow \infty$, the solution $\omega(t)$ converges in C^∞ on compact subsets of $X \setminus D$ to $\Phi^* \omega_{\text{KE}}$, where D is the union of the (-2) curves on X .

Our result concerns the global Gromov-Hausdorff behavior of $\omega(t)$ as $t \rightarrow \infty$. When X only contains disjoint irreducible (-2) -curves, it was shown in [SW4] that the evolving metrics of the normalized Kähler-Ricci flow converge to the unique Kähler-Einstein orbifold metric on X_{can} in the Gromov-Hausdorff topology. However, when the (-2) curve is not irreducible, the local metric models for the corresponding contracted A - D - E singularities are usually complicated and not explicit, and so the arguments in [SW2, SW4] cannot be immediately applied.

The following is our main result. We establish the global geometric convergence for the normalized Kähler-Ricci flow on all minimal surfaces of general type starting with any initial Kähler metric.

Theorem 1.1. *Let X be a minimal surface of general type and $\omega(t)$ be the solution of the normalized Kähler-Ricci flow (1.1). As $t \rightarrow \infty$, the Kähler manifolds $(X, \omega(t))$ converge in the Gromov-Hausdorff sense to $(X_{\text{can}}, \omega_{\text{KE}})$. In particular, the diameter of $(X, \omega(t))$ is uniformly bounded for all t and the convergence is smooth on the smooth part of X_{can} .*

The local smooth convergence was already proved in [Ts, TZo] away from the (-2) curves on X . Theorem 1.1 implies that all the (-2) curves on X are contracted to orbifold points by the Kähler-Ricci flow as $t \rightarrow \infty$. More precisely, if E is a connected (-2) curve on X , then for any $\varepsilon > 0$, there exists an open neighborhood $U_{E, \varepsilon}$ of E in X such that the diameter of $U_{E, \varepsilon}$ with respect to the evolving metrics $\omega(t)$ is less than ε for all sufficiently large t . Our approach does not rely on the a priori existence of the limiting orbifold Kähler-Einstein metric on X_{can} constructed in [Kob].

A surface of general type is a complex surface whose minimal model is a minimal surface of general type. Therefore, a surface of general type can be obtained by blowing up a minimal surface of general type finitely many times. Combined with the results of [SW2, SW4], we obtain:

Corollary 1.1. *Let X be a general type surface. Then the normalized Kähler-Ricci flow on X starting with any initial Kähler metric g_0 is continuous through finitely many contraction surgeries in the Gromov-Hausdorff topology for $t \in [0, \infty)$ and converges in the Gromov-Hausdorff topology to $(X_{\text{can}}, g_{\text{KE}})$. The convergence is smooth away from the (-2) -curves, where g_{KE} is the unique orbifold Kähler-Einstein metric on X_{can} .*

We refer the readers to [SW4, Theorem 1.1] for details of the canonical surgical contractions by the Kähler-Ricci flow.

As this paper was nearing completion, the authors became aware of a preprint of Gang Tian and Zhenlei Zhang [TZl] where the L^p Cheeger-Colding-Tian theory is used to obtain related results. We remark that the estimates in our proof do not require the a priori existence of the orbifold Kähler-Einstein metric on the canonical model, and we only use the maximum principle and other elementary arguments.

The authors thank the referee for a helpful comment on a previous version of this paper.

2. A PRIORI ESTIMATES

Let X be a minimal surface of general type. Since K_X is big and nef, the pluricanonical linear system $|mK_X|$ is base point free for sufficiently large $m \in \mathbb{Z}^+$ and it induces a birational holomorphic map $\Phi_m = |mK_X| : X \rightarrow X_{\text{can}}$. The sequence Φ_m stabilizes for sufficiently large m and $\Phi = \Phi_m$ contracts all the (-2) -curves on X . In particular, $K_X = \Phi^*K_{X_{\text{can}}}$.

We first reduce the Kähler-Ricci flow to a parabolic complex Monge-Ampère equation, as in [Ts, TZo, SW2]. Define a closed $(1, 1)$ form χ on X by

$$\chi = \frac{1}{m} \Phi^* \omega_{\text{FS}} \in c_1(K_X),$$

where $\Phi : X \rightarrow X_{\text{can}} \subset \mathbb{P}^N$ is the map induced by holomorphic sections of K_X^m , and ω_{FS} is the Fubini-Study metric on \mathbb{P}^N . Choose Ω to be the smooth volume form satisfying

$$\chi = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log \Omega, \quad \int_X \Omega = \int_X \omega_0^2$$

Define a family of reference Kähler metrics $\hat{\omega} = \hat{\omega}(t)$ on X by

$$\hat{\omega} = \chi + e^{-t}(\omega_0 - \chi).$$

Then the Kähler-Ricci flow (1.1) is equivalent to the parabolic complex Monge-Ampère equation for $\varphi = \varphi(t)$ given by

$$(2.1) \quad \frac{\partial \varphi}{\partial t} = \log \frac{(\hat{\omega} + \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \varphi)^2}{\Omega} - \varphi, \quad \varphi|_{t=0} = 0.$$

Namely, if φ solves (2.1) then $\omega(t) := \hat{\omega} + \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \varphi$ solves (1.1). Conversely if we are given a solution to (1.1) then we can obtain a solution $\varphi = \varphi(t)$ of (2.1).

Let $D = \sum_{i=1}^M D_i$ be the (-2) curves contracted by Φ , where D_i is an irreducible component of D and D_i is a smooth rational curve with $D_i \cdot D_i = -2$. All components of D meet transversally with each other (see [BHPV], for example). Since the support of D is the exceptional locus of the pluricanonical map Φ and K_X is big and nef, the Kodaira lemma implies that there exist $a_i \in \mathbb{Q}^+$ for $i = 1, \dots, M$ such that the \mathbf{R} -divisor

$$K_X - \varepsilon D'$$

is ample for all $\varepsilon \in (0, 1)$, where $D' = \sum_{i=1}^M a_i D_i$.

Choose $\sigma_i \in \mathcal{O}(D_i)$, a holomorphic section of the line bundle $[D_i]$ associated to the divisor D_i , vanishing to order 1 along D_i . We fix now $\varepsilon_0 \in (0, 1)$. Then there exist Hermitian metrics h_i on $[D_i]$ such that

$$(2.2) \quad \chi - \varepsilon_0 R_h \geq c_{\varepsilon_0} \omega_0,$$

for some positive constant c_{ε_0} , where we are using the notation

$$R_h := -\frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log h = \sum_{i=1}^M -a_i \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log h_i,$$

for the curvature form of the Hermitian metric $h = \otimes_i (h_i)^{a_i}$ on the \mathbf{Q} -line bundle associated to $[D'] = \sum_{i=1}^M a_i [D_i]$.

In the next lemma we gather together some a priori estimates which are already known by the work of Tsuji, Tian-Zhang and Zhang [Ts, TZo, Z1, Z2].

Lemma 2.1. *There exist uniform constants $C > 0$ and $\lambda > 0$ such that the following hold.*

- (i) $|\varphi| + |\dot{\varphi}| + |R| \leq C$ on $X \times [0, \infty)$.
- (ii) On $X \times [0, \infty)$, we have

$$\omega(t) \leq C |\sigma|_h^{-2\lambda} \omega_0,$$

where $|\sigma|_h^2 = \prod_i |\sigma_i|_{h_i}^2$.

- (iii) For any compact $K \subset X \setminus D$, there exists for every $\ell = 0, 1, 2, \dots$ a constant $C_{K,\ell}$ such that

$$\|\varphi\|_{C^\ell(K)} \leq C_{K,\ell}.$$

Proof. Part (i) is due to Tian-Zhang and Zhang [TZo, Z1, Z2], and makes use of the pluripotential theory estimates of Kołodziej [Kol] (for a recent survey, see [PSoS]). Part (ii) was shown by Tsuji [Ts]. Part (iii) can be proved using a Calabi-type third order estimate and then higher order estimates (see [PSeS] and [ShW], for example). \square

Next, we discuss the local model for an irreducible (-2) -curve in X , following the discussion in [SW4]. Fix an irreducible (-2) -curve D_j in X . Then there exists a holomorphic map $\Psi : X \rightarrow \mathbb{P}^K$ contracting D_j whose image $Y := \Psi(X) \subset \mathbb{P}^K$ is subvariety with single orbifold point $y = \Psi(D_j) \in Y$. Using the notation above, h_j is a Hermitian metric on the line bundle $[D_j]$ and σ_j is a holomorphic section of this line bundle, vanishing to order 1 along D_j .

Write $B \subset \mathbb{C}^2$ for the unit ball centered at the origin. Write $\tilde{B} = B/\mathbb{Z}_2$, where \mathbb{Z}_2 acts on B by

$$(z^1, z^2) \mapsto (-z^1, -z^2).$$

We can then identify a neighborhood of the orbifold point $y \in Y$ with \tilde{B} . Write ω_{Eucl} for the Euclidean metric

$$\omega_{\text{Eucl}} = \sqrt{-1} dz^1 \wedge d\bar{z}^1 + \sqrt{-1} dz^2 \wedge d\bar{z}^2$$

on B , which descends to give a smooth orbifold metric on \tilde{B} . Define r by

$$r = \sqrt{|z^1|^2 + |z^2|^2},$$

on B (and also on \tilde{B}). The map Ψ is a biholomorphism from $\Psi^{-1}(\tilde{B} \setminus \{0\})$ to $\tilde{B} \setminus \{0\}$. The curve D_j is given by $\{\sigma_j = 0\}$ and we have (cf. [SW4, Section 2])

$$(2.3) \quad |\sigma_j|_{h_j}^2 = r^4,$$

where we are identifying r with its pull-back via Ψ . In what follows we will, without comment, identify via Ψ the sets $\Psi^{-1}(\tilde{B} \setminus \{0\})$ and $\tilde{B} \setminus \{0\}$, and similarly for the various functions and $(1, 1)$ forms on these sets. Recalling that ω_0 is a fixed Kähler metric on X , we note that on $\tilde{B} \setminus \{0\}$ we have

$$(2.4) \quad C^{-1}r^2\omega_{\text{Eucl}} \leq \omega_0 \leq \frac{C}{r^2}\omega_{\text{Eucl}}.$$

Moreover, if we define the vector field V on $B \setminus \{0\}$ by

$$(2.5) \quad V = z^1 \frac{\partial}{\partial z^1} + z^2 \frac{\partial}{\partial z^2}$$

then we have

$$(2.6) \quad \frac{r^4}{C} \leq |V|_{\omega_0}^2 \leq Cr^4,$$

for a uniform $C > 0$.

A final remark about the local model is that if p is any point in D_j then we can find holomorphic coordinates u, v in a neighborhood W of p , and centered at p , such that $D_j \cap W = \{u = 0\}$ and the map Ψ from W to \tilde{B} is given by $(z^1, z^2) : W \rightarrow \tilde{B}$ where

$$(2.7) \quad u = (z^1)^2, \quad v = \frac{z^2}{z^1}.$$

Namely, $z^1 = u^{1/2}$ and $z^2 = u^{1/2}v$, which gives a well-defined map into \tilde{B} .

Our next lemma shows that we have estimates for the evolving metric $\omega(t)$ on $\tilde{B} \setminus \{0\}$. However, unlike in [SW4], our (-2) -curve D_j may intersect other (-2) curves. To begin with, we only obtain estimates away from these intersection points. To make this more precise, fix any open tubular neighborhood U_η^j of radius $\eta > 0$, with respect to ω_0 , in X of $\bigcup_{i \neq j} D_i$. This open set U_η^j may intersect $\tilde{B} \setminus \{0\}$.

We have the following estimates, which are analogous to those of Lemma 4.1 and Lemma 10.1 in [SW4].

Lemma 2.2. *Fix $j \in \{1, \dots, M\}$ and $\eta > 0$. With the notation above, there exist constants C and $\delta > 0$ such that on $(\tilde{B} \setminus \{0\}) \cap (X - U_\eta^j)$, depending only on the fixed initial data and $\eta > 0$ such that*

- (i) $\omega \leq \frac{C}{r^2}\omega_{\text{Eucl}}$.
- (ii) $\omega \leq \frac{C}{r^{2(1-\delta)}}(\omega_0 + \omega_{\text{Eucl}})$.
- (iii) $|V|_\omega^2 \leq Cr^{4/3}$, where V is the vector field defined by (2.5).

Proof. For simplicity of notation, we take $j = 1$. The proof is fairly straightforward, given the arguments of [SW2, SW4]. The idea is to modify the quantities in [SW4] by adding a term which tends to negative infinity along the union of D_2, \dots, D_M . After applying the maximum principle, we will obtain a bound away from U_η^1 .

Define on $\tilde{B} \setminus \{0\}$, for $\varepsilon > 0$ and $A \gg 1$ a large constant to be determined,

$$H_\varepsilon = \log \operatorname{tr}_{\omega_0} \omega + A \log \left(|\sigma_1|_{h_1}^{(1+\varepsilon)} \left(\prod_{i=2}^M |\sigma_i|_{h_i}^{4\lambda} \right) \operatorname{tr}_{\omega_{\text{Eucl}}} \omega \right) - A^2 \varphi,$$

where λ is the constant in item (ii) of Lemma 2.1. We wish to use the maximum principle to bound H_ε from above on $\tilde{B} \setminus \{0\}$. Using Lemma 2.1(ii) and the fact that the curve D_1 is completely contained inside \tilde{B} , we see that the quantity H_ε is uniformly bounded from above on the boundary of \tilde{B} . In addition, we see from (2.4) that H_ε tends to $-\infty$ as r tends to 0, and H_ε tends to $-\infty$ on any curves D_2, \dots, D_M which intersect \tilde{B} . We also have an upper bound for H_ε at time $t = 0$. We may suppose then that H_ε attains a maximum at an interior point $(x_0, t_0) \in (\tilde{B} \setminus \{0\}) \times (0, \infty)$, and x_0 does not lie on any of the curves D_2, \dots, D_M . It suffices to bound H_ε at this point (x_0, t_0) .

First recall the following well-known pointwise differential inequality for a solution $\omega = \omega(t)$ of the Kähler-Ricci flow [Y, A, C] (or see [SW3, Prop. 3.2.5] for a recent exposition)

$$(2.8) \quad \left(\frac{\partial}{\partial t} - \Delta \right) \log \operatorname{tr}_\beta \omega \leq C_\beta \operatorname{tr}_\omega \beta - 1,$$

for a fixed metric β , where $-C_\beta$ is the infimum of the bisectional curvature of β . In particular, if the curvature of β vanishes then we can take $C_\beta = 0$.

At (x_0, t_0) , we have, applying (2.8) twice, first with $\beta = \omega_0$ and then with $\beta = \omega_{\text{Eucl}}$,

$$\begin{aligned} 0 &\leq \left(\frac{\partial}{\partial t} - \Delta \right) H_\varepsilon \\ &\leq C_0 \operatorname{tr}_\omega \omega_0 + A \left(\frac{1+\varepsilon}{2} \right) \operatorname{tr}_\omega R_{h_1} + 2A\lambda \operatorname{tr}_\omega \left(\sum_{i=2}^M R_{h_i} \right) \\ &\quad - A^2 \dot{\varphi} + 2A^2 - A^2 \operatorname{tr}_\omega \hat{\omega} \\ &= C_0 \operatorname{tr}_\omega \omega_0 - A^2 \dot{\varphi} + 2A^2 \\ &\quad - A^2 \operatorname{tr}_\omega \left(\hat{\omega} - A^{-1} \left(\frac{1+\varepsilon}{2} \right) R_{h_1} - 2A^{-1} \lambda \sum_{i=2}^M R_{h_i} \right), \end{aligned}$$

where $-C_0$ is the lower bound for the bisectional curvature of ω_0 . Using (2.2) and the definition of $\hat{\omega}$, we may choose A uniformly large enough such

that, for a uniform $c_0 > 0$,

$$\hat{\omega} - A^{-1} \left(\frac{1 + \varepsilon}{2} \right) R_{h_1} - 2A^{-1} \lambda \sum_i R_{h_i} \geq c_0 \omega_0$$

and $A^2 c_0 \geq C_0 + 1$. Hence, using Lemma 2.1, part (i), we have

$$0 \leq -\text{tr}_\omega \omega_0 - A^2 \dot{\varphi} + CA^2,$$

for a uniform constant C . Now recall from Lemma 2.1 that φ and $\dot{\varphi}$ (and hence also $\log(\omega^2/\omega_0^2)$) are uniformly bounded. It follows that we obtain a uniform upper bound of $\text{tr}_\omega \omega_0$ at (x_0, t_0) , and from the inequality $\text{tr}_{\omega_0} \omega \leq (\text{tr}_\omega \omega_0) \frac{\omega^2}{\omega_0^2}$, we also get an upper bound for $\text{tr}_{\omega_0} \omega$. Then H_ε is bounded from above at (x_0, t_0) , and hence has a uniform upper bound on $(\tilde{B} \setminus \{0\}) \times [0, \infty)$. Moreover, our upper bound is independent of ε .

Letting $\varepsilon \rightarrow 0$, we obtain on $\tilde{B} \setminus \{0\}$,

$$(2.9) \quad \left(\text{tr}_{\omega_0} \omega \right)^{1/A} |\sigma_1|_{h_1} \left(\prod_{i=2}^M |\sigma_i|_{h_i}^{4\lambda} \right) \text{tr}_{\omega_{\text{Eucl}}} \omega \leq C.$$

First, using (2.4) we have $|\sigma_1|_{h_1} \text{tr}_{\omega_{\text{Eucl}}} \omega \leq C \text{tr}_{\omega_0} \omega$ and hence

$$\text{tr}_{\omega_{\text{Eucl}}} \omega \leq \frac{C}{r^2} \left(\prod_{i=2}^M |\sigma_i|_{h_i}^{4\lambda} \right)^{-A/(A+1)}.$$

where we have used $|\sigma_1|_{h_1} = r^2$ from (2.3). We then obtain the estimate (i)

since the quantity $\left(\prod_{i=2}^M |\sigma_i|_{h_i}^2 \right)^{-1}$ is uniformly bounded on $X \setminus U_\eta^1$, depending only on the fixed data and η .

For (ii), observe that from (2.9) we also have

$$\left(\text{tr}_{\omega_0 + \omega_{\text{Eucl}}} \omega \right)^{1+1/A} \leq C |\sigma_1|_{h_1}^{-1} \left(\prod_{i=2}^M |\sigma_i|_{h_i}^{4\lambda} \right)^{-1},$$

and so using (2.3) again,

$$\text{tr}_{\omega_0 + \omega_{\text{Eucl}}} \omega \leq Cr^{-2(1-\delta)} \left(\prod_{i=2}^M |\sigma_i|_{h_i}^{4\lambda} \right)^{-\frac{A}{A+1}},$$

for some $\delta \in (0, 1)$ depending only on A . This completes the proof of (ii).

For (iii), we will apply the maximum principle to G_ε , for $\varepsilon > 0$, defined by

$$G_\varepsilon = \log \left(|V|_\omega^{1+\varepsilon} \left(\prod_{i=2}^M |\sigma_i|_{h_i}^{4\lambda} \right) \text{tr}_{\omega_{\text{Eucl}}} \omega \right) + A(|z^1|^2 + |z^2|^2),$$

on $\tilde{B} \setminus \{0\}$, for $A > 0$ to be determined. Note that from the bounds (2.3), (2.4) and (2.6), we see that for each fixed t , the quantity $G_\varepsilon(x)$ tends to $-\infty$ as x approaches zero. Suppose that G_ε achieves a maximum at a point

$(x_0, t_0) \in (\tilde{B} \setminus \{0\}) \times (0, \infty)$ such that x_0 does not lie on any of the curves D_2, \dots, D_M .

Following the computation of [SW2, Lemma 2.6], we compute at (x_0, t_0) ,

$$0 \leq \left(\frac{\partial}{\partial t} - \Delta\right)G_\varepsilon \leq \text{tr}_\omega \left(\sum_{i=2}^M 2\lambda R_{h_i} - A\omega_{\text{Eucl}} \right) < 0,$$

if A is chosen sufficiently large. This is a contradiction, implying that the maximum of G_ε occurs at $t = 0$ or on the boundary of \tilde{B} , where, by Lemma 2.1, we have the upper bound for G_ε . Thus G_ε is bounded from above on $\tilde{B} \setminus \{0\}$, and moreover the bound is independent of ε . Letting $\varepsilon \rightarrow 0$, we obtain

$$|V|_\omega \text{tr}_{\omega_{\text{Eucl}}} \omega \leq C \left(\prod_{i=2}^M |\sigma_i|_{h_i}^{4\lambda} \right)^{-1}.$$

But on $(X \setminus U_\eta^1 \cap (\tilde{B} \setminus \{0\}))$, we have from (i),

$$|V|_\omega \text{tr}_{\omega_{\text{Eucl}}} \omega \leq C.$$

On the other hand,

$$|V|_\omega^2 \leq (\text{tr}_{\omega_{\text{Eucl}}} \omega) |V|_{\omega_{\text{Eucl}}}^2 = r^2 \text{tr}_{\omega_{\text{Eucl}}} \omega,$$

and multiplying both sides by $|V|_\omega$ gives $|V|_\omega^2 \leq Cr^{4/3}$ on $(X \setminus U_\eta^1 \cap (\tilde{B} \setminus \{0\}))$. This completes the proof of the lemma. \square

As an immediate consequence of these estimates, we can obtain the following bound on the evolving metric $\omega = \omega(t)$ in terms of local holomorphic coordinate systems on X .

Corollary 2.1. *Fix $j \in \{1, \dots, M\}$ and $\eta > 0$. We can cover $D_j \cap (X \setminus U_\eta^j)$ by finitely many complex coordinate charts W with complex coordinates u, v so that*

- (i) *On each W , the curve D_j is given by $D_j = \{u = 0\}$.*
- (ii) *There exists a uniform C such that on each W ,*

$$(2.10) \quad \omega(t) \leq C \left(\frac{\sqrt{-1} du \wedge d\bar{u}}{|u|^{4/3}} + \sqrt{-1} dv \wedge d\bar{v} \right).$$

Proof. By compactness, it suffices to prove the result for some neighborhood of a fixed point p in $D_j \cap (X \setminus U_\eta^j)$. By the discussion above the statement of Lemma 2.2, we can find coordinates u, v in a neighborhood W of p such

that $W \cap D_j = \{u = 0\}$ and u, v satisfy (2.7). Compute on $W \setminus D_j$

$$\begin{aligned} \sqrt{-1}(dz^1 \wedge d\bar{z}^1 + dz^2 \wedge d\bar{z}^2) &= \frac{(1 + |v|^2)\sqrt{-1}du \wedge d\bar{u}}{4|u|} + \sqrt{-1}|u|dv \wedge d\bar{v} \\ &\quad + \operatorname{Re} \left(\sqrt{-1} \left(\frac{\bar{u}}{u} \right)^{1/2} v du \wedge d\bar{v} \right) \\ &\leq C \left(\frac{\sqrt{-1}du \wedge d\bar{u}}{|u|} + \sqrt{-1}|u|dv \wedge d\bar{v} \right), \end{aligned}$$

if we assume that v is sufficiently small. Moreover, we have

$$(2.11) \quad r^2 = |z^1|^2 + |z^2|^2 = |u|(1 + |v|^2).$$

Then the estimate (i) of Lemma 2.2 implies that on W we have

$$\omega \leq C \left(\frac{\sqrt{-1}du \wedge d\bar{u}}{|u|^2} + \sqrt{-1}dv \wedge d\bar{v} \right),$$

for a uniform constant C .

We can improve the bound in the $\partial/\partial u$ direction. Indeed, note that

$$u \frac{\partial}{\partial u} = \frac{z^1}{2} \frac{\partial}{\partial z^1} + \frac{z^2}{2} \frac{\partial}{\partial z^2} = \frac{V}{2},$$

and hence from part (iii) of Lemma 2.2 and (2.11),

$$g \left(\frac{\partial}{\partial u}, \frac{\partial}{\partial \bar{u}} \right) = \frac{1}{4|u|^2} |V|_\omega^2 \leq \frac{Cr^{4/3}}{|u|^2} \leq \frac{C'}{|u|^{4/3}},$$

and this gives the estimate (ii) on W , as required. \square

Note that although the estimate (2.10) blows up as we approach $\{u = 0\}$, it is strong enough to show that the diameter of each set U_η^j is uniformly bounded. However at the moment, we have no estimates in neighborhoods of the intersection points $p_{ij} = D_i \cap D_j$ for $i \neq j$. The following lemma shows that our estimates do indeed extend to such neighborhoods.

Lemma 2.3. *Fix an intersection point $p_{ij} = D_i \cap D_j$ for $i \neq j$. There exists a complex coordinate chart W centered at p_{ij} with complex coordinates u, v , such that $D_i \cap W = \{u = 0\}$ and $D_j \cap W = \{v = 0\}$. On $W \setminus (D_i \cup D_j)$, we have*

$$(2.12) \quad \omega(t) \leq C \left(\frac{\sqrt{-1}du \wedge d\bar{u}}{|u|^{4/3}} + \frac{\sqrt{-1}dv \wedge d\bar{v}}{|v|^{4/3}} \right).$$

for a uniform constant C .

Proof. Since $D_i \cdot D_j = 1$ if $i \neq j$ and D_i and D_j meet transversally, we can always find local complex coordinate chart W centered at p_{ij} with complex coordinates u, v , such that $D_i \cap W = \{u = 0\}$ and $D_j \cap W = \{v = 0\}$.

Write

$$\omega_{\text{con}} = \frac{\sqrt{-1}du \wedge d\bar{u}}{|u|^{4/3}} + \frac{\sqrt{-1}dv \wedge d\bar{v}}{|v|^{4/3}}.$$

Note that ω_{con} is a metric with conical singularities with simple normal crossings. The cone angle along $(D_i \cup D_j) \setminus \{p_{ij}\}$ is $2\pi/3$ (cf. [D, DGSW]). A key point is that on $W \setminus (D_i \cup D_j)$, the curvature of ω_{con} vanishes, as can be readily checked by a direct computation, or alternatively, one can apply change of variables by letting $u' = u^{1/3}$, $v' = v^{1/3}$ locally on a well-defined branch.

We will apply the maximum principle to the quantity $G = \log \text{tr}_{\omega_{\text{con}}} \omega$ on $W \setminus (D_i \cup D_j)$. The estimates of Corollary 2.1 imply that the quantity G is uniformly bounded from above on the boundary of the set W . Applying (2.8) with $\beta = \omega_{\text{con}}$ and $C_0 = 0$ we see that on $W \setminus (D_i \cup D_j)$ we have

$$\left(\frac{\partial}{\partial t} - \Delta \right) G \leq -1,$$

which implies that the maximum of G occurs either at $t = 0$ or on the boundary of W . In both cases, we have a uniform upper bound¹ for G . Hence $G \leq C$ on $W \setminus (D_i \cup D_j)$, completing the proof of the lemma. \square

From the estimates now obtained, we can bound distances with respect to $g(t)$ in a neighborhood of the curve D . Denote by $B_{g_0, \delta}(p)$ the g_0 geodesic ball in X centered at $p \in X$ with radius δ , and by \tilde{U}_δ the δ -tubular neighborhood of D with respect to g_0 ,

$$(2.13) \quad \tilde{U}_\delta := \{x \in X \mid d_{g_0}(x, D) < \delta\}.$$

Then we have:

Proposition 2.1. *There exist uniform positive constants C, δ_0 so that the following hold.*

(i) *For every intersection point $p_{ij} = D_i \cap D_j$, and any $\delta \in (0, \delta_0)$, we have*

$$\text{diam}_{g(t)} B_{g_0, \delta}(p_{ij}) \leq C\delta^{1/3}.$$

(ii) *For every $\delta \in (0, \delta_0)$,*

$$d_{g(t)}(D, q) \leq C\delta^{1/3}, \quad \text{for all } q \in \tilde{U}_\delta.$$

Proof. For (i), fix an intersection point p_{ij} , and let W be as in the statement of Lemma 2.3, with p_{ij} corresponding to the point $u = v = 0$ in the complex coordinates u, v . We assume that $\delta_0 > 0$ is sufficiently small so that W contains the ball $B_{g_0, \delta_0}(p_{ij})$. Given a point $(u_0, v_0) \in W$ with both u_0 and v_0 nonzero, we compute using the estimate (2.12) that the length of the path $\gamma : [0, 1] \rightarrow W$ given by $\gamma(\lambda) = (\lambda u_0, \lambda v_0)$, with respect to $g(t)$, satisfies

$$\text{Length}_{g(t)}(\gamma) \leq C(|u_0|^{1/3} + |v_0|^{1/3}) \leq C'\delta^{1/3},$$

for uniform constants C, C' . Since for each t , the metric $\omega(t)$ is smooth, we obtain this bound even when one of u_0, v_0 is zero. This gives (i).

¹To clarify this part of the argument (we thank Valentino Tosatti for pointing out our imprecision here): fix $\varepsilon > 0$ and consider $G_\varepsilon = \log \text{tr}_{\omega_{\text{con}}} \omega + \varepsilon \log \frac{\omega^2}{\omega_{\text{con}}^2}$ which satisfies $(\frac{\partial}{\partial t} - \Delta)G_\varepsilon \leq -1$. Then let $\varepsilon \rightarrow 0$.

For (ii), let $q \in \tilde{U}_\delta$. First assume that q is contained in a coordinate chart W as in the discussion above, and q has coordinates (u_0, v_0) . Since q is in \tilde{U}_δ , at least one of $|u_0|$ or $|v_0|$ is of order $O(\delta)$. Without loss of generality we suppose that $|u_0| = O(\delta)$ and $v_0 \neq 0$. Then consider the path $\gamma : [0, 1] \rightarrow W$ given by $\gamma(\lambda) = (\lambda u_0, v_0)$. The length of this path with respect to $g(t)$ satisfies

$$\text{Length}_{g(t)}(\gamma) \leq C|u_0|^{1/3} \leq C'\delta^{1/3},$$

since when restricted to the $\partial/\partial u$ and $\partial/\partial \bar{u}$ directions, the metric $\omega(t)$ is bounded from above by $C\sqrt{-1}du \wedge d\bar{u}/|u|^{4/3}$.

If $q \in \tilde{U}_\delta$ is not contained in such a neighborhood W of an intersection point, we instead apply the estimate of Corollary 2.1, and we obtain the required bound by an identical argument. This completes the proof of (ii). \square

3. GROMOV-HAUSDORFF CONVERGENCE

Given the estimates above, it is now straightforward to derive the Gromov-Hausdorff convergence of $(X, g(t))$ to $(X_{\text{can}}, g_{\text{KE}})$. First, we have

$$(3.1) \quad \int_D \omega(t) = e^{-t} \int_D \omega_0 \longrightarrow 0, \quad \text{as } t \rightarrow \infty,$$

where $D = \sum_i D_i$ is the (-2) curve as introduced above. Indeed, this follows immediately from the Kähler-Ricci flow equation since

$$\frac{\partial}{\partial t} \int_D \omega(t) = - \int_D \text{Ric}(\omega(t)) - \int_D \omega(t) = - \int_D \omega(t),$$

where we have used the fact that $K_X \cdot D = 0$.

First we prove:

Lemma 3.1. *If \hat{D} is a connected component of D then*

$$\text{diam}_{g(t)} \hat{D} \rightarrow 0, \quad \text{as } t \rightarrow \infty.$$

Proof. For simplicity, assume that $\hat{D} = D_1 \cup D_2$, and that D_1 and D_2 intersect at a point p (the general case is similar). Fix $\varepsilon > 0$. Then by Proposition 2.1 there exists $\delta > 0$ such that the g_0 geodesic ball $B_{g_0, \delta}(p)$ of radius δ centered at p has

$$(3.2) \quad \text{diam}_{g(t)} B_{g_0, \delta}(p) < \varepsilon/3.$$

Define $D_1^{(\varepsilon)} = D_1 \setminus B_{g_0, \delta}(p)$. The estimate of Lemma 2.2, part (ii), together with [SW4, Lemma 5.1] gives the existence of a constant C_ε , depending on ε , such that

$$\omega(t)|_{D_1^{(\varepsilon)}} \leq C_\varepsilon \omega_0|_{D_1^{(\varepsilon)}}.$$

We claim that if a, b are any two points in $D_1^{(\varepsilon)}$, then

$$d_{g(t)}(a, b) \leq C'_\varepsilon e^{-t/3},$$

for another constant C'_ε depending on ε . The claim follows almost verbatim from the argument in [SW2] or [SSW, Lemma 2.4] and the estimate (3.1). Indeed, the only differences are that we have to keep track of the dependence on ε , and we replace the factor $(T - t)$ in [SSW, Lemma 2.4] with e^{-t} .

The proof is now essentially finished. Choose t sufficiently large so that $C'_\varepsilon e^{-t/3} < \varepsilon/3$, so that

$$(3.3) \quad \text{diam}_{g(t)} D_1^{(\varepsilon)} \leq \varepsilon/3.$$

Similarly, if we set $D_2^{(\varepsilon)} = D_2 \setminus B_{g_0, \delta}(p)$ then we obtain

$$(3.4) \quad \text{diam}_{g(t)} D_2^{(\varepsilon)} \leq \varepsilon/3.$$

The result then follows by combining (3.2), (3.3) and (3.4). \square

Combining this lemma with the estimates established in Proposition 2.1, we immediately obtain:

Lemma 3.2. *Let \hat{D} be a connected component of D . For $\delta > 0$, write*

$$\hat{U}_\delta := \{x \in X \mid d_{g_0}(x, \hat{D}) < \delta\},$$

for its δ -tubular neighborhood with respect to g_0 . Then for any $\varepsilon > 0$ there exists $\delta > 0$ and $T > 0$ such that

$$(3.5) \quad \text{diam}_{g(t)} \hat{U}_\delta \leq \varepsilon, \quad \text{for } t \geq T.$$

As a consequence, the diameter of $(X, g(t))$ is uniformly bounded.

Proof. Combine part (ii) of Proposition 2.1 with Lemma 3.1 to obtain (3.5). For the last assertion, fix $\delta > 0$ and use Lemma 2.1 to see that the diameter of $X \setminus V_\delta$ is uniformly bounded, where V_δ is the union of the δ -tubular neighborhoods \hat{U}_δ of all the connected components of D . \square

Finally, we complete the proof of our main result.

Proof of Theorem 1.1. By the results in [Ts] and [TZo], the solution $\varphi(t)$ of the Monge-Ampère flow (2.1) converges smoothly on $X \setminus D$ to a bounded χ -psh function φ_{KE} satisfying

$$\left(\chi + \frac{\sqrt{-1}}{2\pi} \partial\bar{\partial}\varphi_{\text{KE}}\right)^2 = e^{\varphi_{\text{KE}}}\Omega$$

as $t \rightarrow \infty$ and $\varphi_{\text{KE}} \in C^\infty(X \setminus D)$. Since χ vanishes on D and $\varphi_{\text{KE}} \in L^\infty(X) \cap \text{PSH}(X, \chi)$, φ_{KE} is constant along each connected component of D . Immediately $\chi + \frac{\sqrt{-1}}{2\pi} \partial\bar{\partial}\varphi_{\text{KE}} = \Phi^* \omega_{\text{KE}}$ for some Kähler current $\omega_{\text{KE}} \in -c_1(X_{\text{can}})$ on X_{can} with bounded local potentials. We then can apply the smoothing properties of the weak Kähler-Ricci flow [SoT2, Theorem 4.5] on Kähler orbifolds to show that ω'_{KE} must be a smooth orbifold Kähler-Einstein metric on X_{can} . In particular, $\omega_{\text{KE}} = \omega'_{\text{KE}}$ by the uniqueness of orbifold Kähler-Einstein metrics on Kähler orbifolds with ample canonical bundle.

From the local smooth convergence of φ to φ_{KE} on $X \setminus D$, we have

$$\omega(t) \rightarrow \omega_{\text{KE}}, \quad \text{as } t \rightarrow \infty,$$

in $C^\infty(K)$ for any compact subsets $K \subset X \setminus D$. Combining this fact with Lemma 3.2 and the argument of [SW2, Section 3], we immediately obtain the Gromov-Hausdorff convergence of $(X, \omega(t))$ to $(X_{\text{can}}, \omega_{\text{KE}})$. \square

REFERENCES

- [A] Aubin, T. *Equations du type Monge-Ampère sur les variétés kählériennes compactes*, Bull. Sci. Math. (2) 102 (1978), no. 1, 63–95
- [BHPV] Barth, W. P., Hulek, K., Peters, C. A. M. and Van de Ven, A. *Compact complex surfaces*. Second edition. Ergebnisse der Mathematik und ihrer Grenzgebiete. Springer-Verlag, Berlin
- [C] Cao, H.-D. *Deformation of Kähler metrics to Kähler-Einstein metrics on compact Kähler manifolds*, Invent. Math. 81 (1985), no. 2, 359–372
- [CT] Collins, T. and Tosatti, V. *Kähler currents and null loci*, preprint, arXiv:1304.5216
- [DGSW] Datar, V., Guo, B., Song, J. and Wang, X. *Connecting toric conical Kähler-Einstein manifolds*, arXiv:1308.6781
- [D] Donaldson, S.K. *Kähler metrics with cone singularities along a divisor*, Essays in mathematics and its applications, 49–79, Springer, Heidelberg, 2012.
- [EGZ] Eyssidieux, P., Guedj, V. and Zeriahi, A. *Singular Kähler-Einstein metrics*. J. Amer. Math. Soc. 22 (2009), no. 3, 607–639
- [FIK] Feldman, M., Ilmanen, T. and Knopf, D. *Rotationally symmetric shrinking and expanding gradient Kähler-Ricci solitons*, J. Differential Geometry 65 (2003), no. 2, 169–209
- [FZ] Fong, F. T.-H. and Zhang, Z. *The collapsing rate of the Kähler-Ricci flow with regular infinite time singularity*, J. Reine Angew. Math. 703 (2015), 95–113
- [G] Gill, M. *Collapsing of products along the Kähler-Ricci flow*, Trans. Amer. Math. Soc. 366 (2014), no. 7, 3907–3924
- [GTZ] Gross, M., Tosatti, V., Zhang, Y. *Collapsing of abelian fibred Calabi-Yau manifolds*, Duke Math. J. 162 (2013), no. 3, 517–551
- [H] Hamilton, R. S. *Three-manifolds with positive Ricci curvature*, J. Differential Geom. 17 (1982), no. 2, 255–306
- [Kob] Kobayashi, R., *Einstein-Kähler V-metrics on open Satake V-surfaces with isolated quotient singularities*, Math. Ann. 272 (1985), no. 3, 385–398
- [Kol] Kolodziej, S. *The complex Monge-Ampère equation*, Acta Math. 180 (1998), no. 1, 69–117
- [P1] Perelman, G. *The entropy formula for the Ricci flow and its geometric applications*, preprint, arXiv: math.DG/0211159
- [P2] Perelman, G. *Ricci Flow with Surgery on Three-Manifolds*, preprint, arXiv: math.DG/0303109
- [P3] Perelman, G. *Finite Extinction Time for the Solutions to the Ricci Flow on Certain Three-Manifolds*, preprint, arXiv: math.DG/0307245
- [PSeS] Phong, D. H., Sesum, N. and Sturm, J. *Multiplier ideal sheaves and the Kähler-Ricci flow*, Comm. Anal. Geom. 15 (2007), no. 3, 613–632
- [PSoS] Phong, D.H., Song, J. and Sturm, J. *Complex Monge-Ampère equations*, Surveys in differential geometry Vol. XVII, 327–410, Surv. Differ. Geom., 17, Int. Press, Boston, MA, 2012.
- [ShW] Sherman, M. and Weinkove, B. *Interior derivative estimates for the Kähler-Ricci flow*, Pacific J. Math. 257 (2012), no. 2, 491–501
- [S] Song, J. *Ricci flow and birational surgery*, arXiv:1304.2607

- [SSW] Song, J., Székelyhidi, G. and Weinkove, B. *The Kähler-Ricci flow on projective bundles*, Int. Math. Res. Not. 2013 (2013), no. 2, 243–257
- [SoT1] Song, J., and Tian, G. *The Kähler-Ricci flow on surfaces of positive Kodaira dimension*, Invent. Math. 170 (2007), no. 3, 609–653
- [SoT2] Song, J., and Tian, G. *Canonical measures and Kähler-Ricci flow*, J. Amer. Math. Soc. 25 (2012), no. 2, 303–353
- [SoT3] Song, J., and Tian, G. *The Kähler-Ricci flow through singularities*, preprint, arXiv: 0909.4898
- [SoT4] Song, J., and Tian, G. *Bounding scalar curvature for global solutions of the Kähler-Ricci flow*, arXiv:1111.5681, to appear in Amer. J. Math.
- [SW1] Song, J. and Weinkove, B. *The Kähler-Ricci flow on Hirzebruch surfaces*, J. Reine Angew. Math. 659 (2011), 141–168
- [SW2] Song, J. and Weinkove, B. *Contracting exceptional divisors by the Kähler-Ricci flow*, Duke Math. J. 162 (2013), no. 2, 367–415
- [SW3] Song, J. and Weinkove, B. *Introduction to the Kähler-Ricci flow*, Chapter 3 of “Introduction to the Kähler-Ricci flow”, eds S. Boucksom, P. Eyssidieux, V. Guedj, Lecture Notes Math. 2086, Springer 2013
- [SW4] Song, J. and Weinkove, B. *Contracting exceptional divisors by the Kähler-Ricci flow*, Proc. Lond. Math. Soc. (3) 108 (2014), no. 6, 1529–1561
- [SY] Song, J. and Yuan, Y. *Metric flips with Calabi ansatz*, Geom. Func. Anal. 22 (2012), no. 1, 240–265
- [T] Tian, G. *New results and problems on Kähler-Ricci flow*, Géométrie différentielle, physique mathématique, mathématiques et société. II. Astérisque 322 (2008), 71–92
- [TZI] Tian, G. and Zhang, Z. *Convergence of the Kähler-Ricci flow on lower dimensional smooth minimal models of general type*, preprint
- [TZo] Tian, G. and Zhang, Z. *On the Kähler-Ricci flow on projective manifolds of general type*, Chinese Ann. Math. Ser. B 27 (2006), no. 2, 179–192
- [TWY] Tosatti, V., Weinkove, B. and Yang, X. *The Kähler-Ricci flow, Ricci-flat metrics and collapsing limits*, preprint, arXiv: 1408.0161
- [Ts] Tsuji, H. *Existence and degeneration of Kähler-Einstein metrics on minimal algebraic varieties of general type*, Math. Ann. 281 (1988), no. 1, 123–133
- [W] Włodarczyk, J. *Birational cobordisms and factorization of birational maps*, J. Algebraic Geom. 9 (2000), no. 3, 425–449
- [Y] Yau, S.-T. *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation, I*, Comm. Pure Appl. Math. 31 (1978), no. 3, 339–411
- [Z1] Zhang, Z. *On degenerate Monge-Ampère equations over closed Kähler manifolds*, Int. Math. Res. Not. 2006, Art. ID 63640, (2006) 18 pp
- [Z2] Zhang, Z. *Scalar curvature bound for Kähler-Ricci flows over minimal manifolds of general type*, Int. Math. Res. Not. IMRN 2009 (2009), no. 20, 3901–3912

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATAWAY, NJ 08854

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATAWAY, NJ 08854

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, EVANSTON, IL 60208