

# Convergence of Kähler-Ricci flow on lower dimensional algebraic manifolds of general type

Gang Tian\*

Beijing University and Princeton University

Zhenlei Zhang†

Capital Normal University

## Abstract

In this paper, we prove that the  $L^4$ -norm of Ricci curvature is uniformly bounded along a Kähler-Ricci flow on any minimal algebraic manifold. As an application, we show that on any minimal algebraic manifold  $M$  of general type and with dimension  $n \leq 3$ , any solution of the normalized Kähler-Ricci flow converges to the unique singular Kähler-Einstein metric on the canonical model of  $M$  in the Cheeger-Gromov topology.

## 1 Introduction

The purpose of this note is to prove the following

**Theorem 1.1.** *Let  $M$  be a smooth minimal model of general type with dimension  $n \leq 3$  and  $\omega(t)$  be a solution to the normalized Kähler-Ricci flow*

$$\frac{\partial}{\partial t}\omega = -\text{Ric} - \omega. \quad (1.1)$$

*Then  $(M, \omega(t))$  converges in the Cheeger-Gromov sense to the unique singular Kähler-Einstein metric on the canonical model of  $M$ .*

Here, by a smooth minimal model, we mean an algebraic manifold  $M$  with nef canonical bundle  $K_M$ . The theorem should remain true in higher dimensional case; cf. Conjecture 4.1 in [27]. Assuming the uniform bound of Ricci curvature, the conjecture is confirmed by Guo [18]. On the other hand, it has been known since Tsuji [32] and Tian-Zhang [31] the convergence of the Kähler-Ricci flow in the current sense and the smooth convergence on the ample locus of the canonical class.

Applying the  $L^2$  bound of Riemannian curvature (cf. Section 3) and Kähler-Einstein condition, we can say more about the limit singular space  $M_\infty$ . When  $n = 2$ , by a

---

\*Supported partially by NSF and NSFC grants. Email: tian@math.princeton.edu

†Supported partially by NSFC 11431009 and Chinese Scholarship Council. Email: zhleigo@aliyun.com

classical argument of removing isolated singularities following Anderson [1], Bando-Kasue-Nakajima [4] and Tian [29], we can show that the limit is a smooth Kähler-Einstein orbifold with finite orbifold points. When  $n = 3$ , by the argument of Cheeger-Colding-Tian [11] or Cheeger [7], we have that the 2-dimensional Hausdorff measure of the singular set is finite. Moreover, it follows from the parabolic version of the partial  $C^0$ -estimate (see [30]) that the limit  $M_\infty$  is a normal variety and there is a natural holomorphic map from  $M$  onto  $M_\infty$ . This actually implies that  $M_\infty$  is the canonical model of  $M$ .

The proof of our theorem relies on a uniform  $L^4$  bound of Ricci curvature under the Kähler-Ricci flow on a smooth minimal model together with the diameter boundedness of the limit singular Kähler-Einstein space (in the case of general type) proved by Song [26]. From these we derive a uniform local noncollapsing estimate of Kähler-Ricci flow on a minimal model of general type and the Gromov-Hausdorff convergence follows.

In Section 2, we present a short discussion on manifolds with integral bounded Ricci curvature, with emphasis on a uniform local noncollapsing estimate which is essential in extending the regularity theory of Cheeger-Colding and Cheeger-Colding-Tian. Then, in Section 3, we give a proof of our theorem by establishing a uniform  $L^4$  Ricci curvature estimate under the Kähler-Ricci flow.

After we completed the first draft of this paper, we learned that Guo-Song-Weinkove obtained a different proof for the 2-dimensional case of Theorem 1.1 (see [19]).

## 2 Manifolds with $L^p$ bounded Ricci curvature

We recall the relative volume comparison formula relative to the  $L^p$  Ricci curvature,  $p$  bigger than half dimension, which is due to Petersen-Wei [21]. It implies a uniform local noncollapsing property for the Kähler-Ricci flow on lower dimensional minimal models of general type. We will use this to prove the convergence of Kähler-Ricci flow on such manifolds.

Let  $(M, g)$  be a complete Riemannian manifold of (real) dimension  $m$ . For any  $\kappa \in \mathbb{R}$  denote by  $B_r^\kappa$  a metric ball of radius  $r$  in the space form of dimension  $m$  with sectional curvature  $\kappa$  and by  $\text{vol}(B_r^\kappa)$  its volume. Then we have

**Theorem 2.1** ([21]). *For any  $p > \frac{m}{2}$  and  $\Lambda < 0$ , there exists  $C = C(m, p, \Lambda)$  such that the followings hold. First of all, for any  $x \in M$  and  $r > 0$ ,*

$$\frac{d}{dr} \left( \frac{\text{vol}(B_r(x))}{\text{vol}(B_r^\Lambda)} \right)^{\frac{1}{2p}} \leq \frac{Cr^{2p}}{\text{vol}(B_r^\Lambda)} \left( \int_{B_r(x)} |(\text{Ric} - (m-1)\Lambda g)_-|^p dv \right)^{\frac{1}{2p}}, \quad (2.1)$$

where we define

$$(\text{Ric} - (m-1)\Lambda g)_- = \max_{|v|=1} (0, -\text{Ric}(v, v) + (m-1)\Lambda)$$

pointwisely. In particular, for any  $r_2 > r_1 > 0$ ,

$$\left( \frac{\text{vol}(B_{r_2}(x))}{\text{vol}(B_{r_2}^\Lambda)} \right)^{\frac{1}{2p}} - \left( \frac{\text{vol}(B_{r_1}(x))}{\text{vol}(B_{r_1}^\Lambda)} \right)^{\frac{1}{2p}} \leq C \left( \int_{B_{r_2}(x)} |(\text{Ric} - (m-1)\Lambda g)_-|^p dv \right)^{\frac{1}{2p}}. \quad (2.2)$$

Then, by letting  $r_1 \rightarrow 0$  it gives, for any  $r > 0$ ,

$$\frac{\text{vol}(B_r(x))}{\text{vol}(B_r^\Lambda)} \leq 1 + C \int_{B_r(x)} |(\text{Ric} - (m-1)\Lambda g)_-|^p dv. \quad (2.3)$$

The following corollary gives a kind of uniform local noncollapsing property on manifolds with integral Ricci curvature bound; see [22] and [16] for similar volume doubling estimates.

**Corollary 2.2.** *For any  $\Lambda < 0$  and  $p > \frac{m}{2}$ , there exists  $\varepsilon = \varepsilon(m, p, \Lambda) > 0$  such that the following holds. Fix a base point  $x_0 \in M$ . For any  $x \in M$  with  $\text{dist}(x_0, x) = d$ , if*

$$\frac{1}{\text{vol}(B_1(x_0))} \int_{B_{2d+1}(x_0)} |(\text{Ric} - (m-1)\Lambda g)_-|^p dv \leq \frac{\varepsilon}{\text{vol}(B_{d+1}^\Lambda)}, \quad (2.4)$$

then,

$$\frac{\text{vol}(B_r(x))}{r^m} \geq \frac{\text{vol}(B_1(x_0))}{2 \text{vol}(B_{d+1}^\Lambda)}, \quad \forall r \leq 1. \quad (2.5)$$

*Proof.* By (2.2), for any  $r \leq 1$ ,

$$\begin{aligned} \left( \frac{\text{vol}(B_r(x))}{\text{vol}(B_r^\Lambda)} \right)^{\frac{1}{2p}} &\geq \left( \frac{\text{vol}(B_{d+1}(x))}{\text{vol}(B_{d+1}^\Lambda)} \right)^{\frac{1}{2p}} - C \left( \int_{B_{d+1}(x)} |(\text{Ric} - (m-1)\Lambda g)_-|^p dv \right)^{\frac{1}{2p}} \\ &\geq \left( \frac{\text{vol}(B_1(x_0))}{\text{vol}(B_{d+1}^\Lambda)} \right)^{\frac{1}{2p}} - C \left( \int_{B_{2d+1}(x_0)} |(\text{Ric} - (m-1)\Lambda g)_-|^p dv \right)^{\frac{1}{2p}}. \end{aligned}$$

where  $C = C(m, p, \Lambda)$ . If (2.4) holds for some  $\varepsilon = \varepsilon(m, p, \Lambda)$  sufficiently small, then

$$\left( \frac{\text{vol}(B_r(x))}{\text{vol}(B_r^\Lambda)} \right)^{\frac{1}{2p}} \geq \frac{1}{2^{2p}} \left( \frac{\text{vol}(B_1(x_0))}{\text{vol}(B_{d+1}^\Lambda)} \right)^{\frac{1}{2p}},$$

which is exactly the estimate (2.5).  $\square$

The lemma suggests a condition for Gromov-Hausdorff convergence. Suppose we have a sequence of complete Riemannian manifolds  $(M_i, g_i)$  of dimension  $m$  such that

$$\int_M |(\text{Ric}_{g_i} - (m-1)\Lambda g_i)_-|^p dv_{g_i} \rightarrow 0 \quad (2.6)$$

for some  $p > \frac{m}{2}$ ,  $\Lambda > 0$ , and

$$\text{vol}_{g_i}(B_1(x_i)) \geq v \quad (2.7)$$

uniformly for some  $v > 0$  and  $x_i \in M_i$ , then along a subsequence, the manifolds  $(M_i, g_i)$  are uniformly locally noncollapsing on  $B_r(x_i)$  for any fixed  $r > 0$ . Thus, we can apply Gromov precompactness theorem to get a noncollapsing limit of  $(M_i, g_i, x_i)$  in the Gromov-Hausdorff topology. Furthermore, as showed in [30], one can extend the regularity theory

of Colding [12], Cheeger-Colding [8, 9], Cheeger-Colding-Tian [11] and Colding-Naber [13] in this setting. If in addition we replace (2.6) by

$$\int_M |\text{Ric}_{g_i} - (m-1)\Lambda g_i|^p dv_{g_i} \rightarrow 0, \quad (2.8)$$

then Anderson's harmonic radius estimate [2] can also be applied. In summation, we can follow the arguments in [22] and [30] to prove

**Theorem 2.3.** *Let  $(M_i, g_i)$  be a sequence of Riemannian manifolds satisfying (2.7) and (2.8) for some  $p > \frac{m}{2}$  and  $\Lambda, v > 0$ . Then, passing to a subsequence if necessary,  $(M_i, g_i, x_i)$  converges in the Cheeger-Gromov sense to a limit length space  $(M_\infty, d_\infty, x_\infty)$  such that*

- (1) *for any  $r > 0$  and  $y_i \in M_i$  with  $y_i \rightarrow y_\infty \in M_\infty$  we have*

$$\text{vol}(B_r(y_i)) \rightarrow \mathcal{H}^m(B_r(y_\infty)), \quad (2.9)$$

*where  $\mathcal{H}^m$  denotes the  $m$ -dimensional Hausdorff measure;*

- (2)  *$M_\infty = \mathcal{R} \cup \mathcal{S}$  such that  $\mathcal{S}$  is a closed set of codimension  $\geq 2$  and  $\mathcal{R}$  is convex in  $M_\infty$ ;  $\mathcal{R}$  consists of points whose tangent cones are  $\mathbb{R}^m$ ;*
- (3) *the convergence on  $\mathcal{R}$  takes place in the  $C^\alpha$  topology for any  $0 < \alpha < 2 - \frac{m}{p}$ ;*
- (4) *the tangent of any  $y \in M_\infty$  is a metric cone which splits off lines isometrically; the tangent cone has the same properties presented in (2) and (3);*
- (5) *the singular set of  $\mathcal{S}$  has codimension  $\geq 4$  if each  $M_i$  is Kählerian.*

### 3 Kähler-Ricci flow on minimal models

#### 3.1 $L^4$ bound of Ricci curvature under Kähler-Ricci flow on minimal models

Let  $M$  be a smooth minimal model. If the Kodaira dimension equals 0, the manifold is Calabi-Yau and any Kähler-Ricci flow on  $M$  converges smoothly to a Ricci flat metric [6]. We assume from now on that the Kodaira dimension of  $M$  is positive. Then we consider the normalized Kähler-Ricci flow

$$\frac{\partial}{\partial t} \omega(t) = -\text{Ric}_{\omega(t)} - \omega(t), \quad \omega(0) = \omega_0. \quad (3.1)$$

It is well-known that the solution exists for all time  $t \in [0, \infty)$  [31].

We shall prove the following theorem.

**Theorem 3.1.** *Suppose  $M$  has positive Kodaira dimension and  $K_M$  is semi-ample. Then there is a constant  $C$  depending on  $\omega_0$  such that*

$$\int_t^{t+1} \int_M |\text{Ric}_{\omega(s)}|^4 \omega(s)^n ds \leq C, \quad \forall t \in [0, \infty). \quad (3.2)$$

Moreover, for any  $0 < p < 4$  we have

$$\int_t^{t+1} \int_M |\text{Ric}_{\omega(s)} + \omega(s)|^p \omega(s)^n ds \rightarrow 0, \text{ as } t \rightarrow \infty. \quad (3.3)$$

We start with some general set-up following [28]. Since  $K_M$  is semi-ample, a basis of  $H^0(M; K_M^\ell)$  for some large  $\ell$  gives rise to a holomorphic map

$$\pi : M \rightarrow \mathbb{C}P^N, \quad N = \dim H^0(M; K_M^\ell) - 1. \quad (3.4)$$

Let  $\omega_{FS}$  be the Fubini-Study metric on  $\mathbb{C}P^N$  and put

$$\chi = \frac{1}{\ell} \pi^* \omega_{FS} \in [K_M]. \quad (3.5)$$

Choose a smooth volume form  $\Omega$  such that  $\text{Ric}(\Omega) = -\chi$ . Put

$$\hat{\omega}_t = e^{-t} \omega_0 + (1 - e^{-t}) \chi.$$

It represents a Kähler metric in the class  $[\omega(t)]$  and write

$$\omega(t) = \hat{\omega}_t + \sqrt{-1} \partial \bar{\partial} \varphi(t) \quad (3.6)$$

for a family of smooth functions  $\varphi(t)$ . Then the Kähler-Ricci flow (3.1) is equivalent to

$$\frac{\partial \varphi}{\partial t} = \log \frac{e^{(n-\kappa)t} (\hat{\omega}_t + \sqrt{-1} \partial \bar{\partial} \varphi)^n}{\Omega} - \varphi. \quad (3.7)$$

**Lemma 3.2** ([28]). *There exists  $C_i = C_i(\omega_0, \chi)$ ,  $i = 1, 2$ , such that*

$$\|\varphi(t)\|_{C^0} + \|\varphi'(t)\|_{C^0} \leq C_1, \quad \forall t \geq 0 \quad (3.8)$$

and

$$\chi \leq C_2 \omega(t), \quad \forall t \geq 0. \quad (3.9)$$

Let  $u = \varphi + \varphi'$  for any time  $t$ .

**Lemma 3.3** ([28]). *There exists  $C_3 = C_3(\omega_0, \chi)$  such that*

$$\|\nabla u(t)\|_{C^0} + \|\Delta u(t)\|_{C^0} \leq C_3, \quad \forall t \geq 0. \quad (3.10)$$

When the manifold is of general type, these estimates are proved in [35]. We also observe that, from (3.7),

$$\text{Ric}_{\omega(t)} + \sqrt{-1} \partial \bar{\partial} u(t) = -\chi. \quad (3.11)$$

So, by the uniform bound of  $\chi$  in terms of  $\omega(t)$ , to prove the  $L^4$  bound of Ricci curvature, it suffices to show

$$\int_t^{t+1} \int_M |\partial \bar{\partial} u(s)|^4 \omega(s)^n ds \leq C_4, \quad \forall t \geq 0 \quad (3.12)$$

for some  $C_4$  independent of  $t$ . We follow the line in [30] to prove this estimate.

As in [30] we let  $\nabla$  and  $\bar{\nabla}$  denote the (1,0) and (0,1) part of the Levi-Civita connection of  $\omega(t)$ . Then, by the calculations in [30], we have the following lemmas

**Lemma 3.4.** *There exists  $C_5 = C_5(\omega_0, \chi)$  such that*

$$\int_M (|\nabla\nabla u|^2 + |\nabla\bar{\nabla}u|^2 + |\text{Ric}|^2 + |Rm|^2)\omega^n \leq C_5, \forall t \geq 0. \quad (3.13)$$

**Lemma 3.5** ([30]). *There exists a universal constant  $C_6 = C_6(n)$  such that*

$$\int_M |\nabla\bar{\nabla}u|^4 \leq C_6 \int_M |\nabla u|^2 (|\bar{\nabla}\nabla\nabla u|^2 + |\nabla\nabla\bar{\nabla}u|^2); \quad (3.14)$$

$$\int_M (|\bar{\nabla}\nabla\nabla u|^2 + |\nabla\nabla\bar{\nabla}u|^2) \leq C_6 \int_M (|\nabla\Delta u|^2 + |\nabla u|^2 \cdot |Rm|^2). \quad (3.15)$$

The estimates in the last lemma are general facts which remain hold for any smooth function.

The following theorem gives the required estimate (3.2).

**Theorem 3.6.** *There exists  $C_7 = C_7(\omega_0, \chi)$  such that*

$$\int_t^{t+1} \int_M (|\nabla\bar{\nabla}u|^4 + |\nabla\nabla\bar{\nabla}u|^2 + |\bar{\nabla}\nabla\nabla u|^2) \leq C_7, \forall t \geq 0. \quad (3.16)$$

*Proof.* By the previous two lemmas we are sufficient to prove a uniform  $L^2$  bound of  $\nabla\Delta u$ . Recall the evolution of  $\Delta u$ , cf. (3.22) in [28],

$$\frac{\partial}{\partial t}\Delta u = \Delta\Delta u + \Delta u + \langle \text{Ric}, \partial\bar{\partial}u \rangle_\omega + \Delta(\text{tr}_\omega \chi).$$

Thus,

$$\frac{\partial}{\partial t}(\Delta u)^2 = \Delta(\Delta u)^2 - 2|\nabla\Delta u|^2 + 2(\Delta u)^2 + 2\Delta u \langle \text{Ric}, \partial\bar{\partial}u \rangle + 2\Delta u \Delta(\text{tr}_\omega \chi). \quad (3.17)$$

Integrating over the manifolds gives

$$\begin{aligned} \int_M |\nabla\Delta u|^2 &\leq \int_M \left( (\Delta u)^2 + |\Delta u| |\text{Ric}| |\nabla\bar{\nabla}u| - \nabla_i \Delta u \nabla_{\bar{i}}(\text{tr}_\omega \chi) - \frac{1}{2} \frac{\partial}{\partial t} (\Delta u)^2 \right) \\ &\leq \int_M \left( \frac{1}{2} |\nabla\Delta u|^2 + (\Delta u)^2 + (\Delta u)^2 |\text{Ric}|^2 + |\nabla\bar{\nabla}u|^2 + 2|\nabla(\text{tr}_\omega \chi)|^2 \right) \\ &\quad - \frac{1}{2} \int_M (\Delta u)^2 (s+n) - \frac{1}{2} \frac{d}{dt} \int_M (\Delta u)^2. \end{aligned}$$

Applying the uniform bound of  $\Delta u$  and above lemma, we get

$$\int_M |\nabla\Delta u|^2 \leq C \int_M (1 + |\nabla(\text{tr}_\omega \chi)|^2) - \frac{d}{dt} \int_M (\Delta u)^2.$$

Integrating over the time interval we have

$$\int_t^{t+1} \int_M |\nabla\Delta u|^2 \leq C \int_t^{t+1} \int_M (1 + |\nabla(\text{tr}_\omega \chi)|^2) + C, \forall t \geq 0. \quad (3.18)$$

The term  $|\nabla(\text{tr}_\omega \chi)|^2$  can be estimated through the Schwarz lemma. Recall the following formula

$$\Delta(\text{tr}_\omega \chi) \geq -|\text{Ric}| \text{tr}_\omega \chi - C(\text{tr}_\omega \chi)^2 + \frac{|\nabla(\text{tr}_\omega \chi)|^2}{\text{tr}_\omega \chi} \quad (3.19)$$

where  $C$  is a universal constant given by the upper bound of the bisectional curvature of  $\omega_{FS}$  on  $\mathbb{C}P^n$ . Because  $0 \leq \text{tr}_\omega \chi \leq C$  under the flow, we have

$$|\nabla(\text{tr}_\omega \chi)|^2 \leq C \left( \Delta(\text{tr}_\omega \chi) + C|\text{Ric}| + C \right).$$

Thus,

$$\int_M |\nabla(\text{tr}_\omega \chi)|^2 \leq C \int_M (1 + |\text{Ric}|) \leq C$$

uniformly. Substituting into (3.18) we get the desired estimate.  $\square$

To prove (3.3) we use the  $L^2$  estimate to traceless Ricci curvature following the calculation by Y. Zhang [34].

**Lemma 3.7.** *Under the Kähler-Ricci flow,*

$$\int_t^{t+1} \int_M |\text{Ric}_{\omega(s)} + \omega(s)|^2 \omega(s)^n ds \rightarrow 0, \text{ as } t \rightarrow \infty. \quad (3.20)$$

*Proof.* Recall the evolution of scalar curvature  $R = \text{tr}_\omega \text{Ric}$ , cf. [34],

$$\frac{\partial}{\partial t} R = \Delta R + |\text{Ric}|^2 + R = \Delta R + |\text{Ric} + \omega|^2 - (R + n).$$

The maximum principle shows that  $\check{R} = \inf R$  satisfies  $\frac{d}{dt} \check{R} \geq -(\check{R} + n)$ , which implies immediately

$$\check{R}(t) + n \geq e^{-t} \min(\check{R}(0) + n, 0) \geq -Ce^{-t} \quad (3.21)$$

for some  $C = C(\omega_0) > 0$ . Then,

$$\begin{aligned} \int |\text{Ric} + \omega|^2 \omega^n &= \int \left( \frac{\partial}{\partial t} R + R + n \right) \omega^n \\ &= \frac{d}{dt} \int R \omega^n + \int (R + n)(R + 1) \omega^n \\ &= \frac{d}{dt} \int R \omega^n + \int (R + n + Ce^{-t})(R + 1) \omega^n - Ce^{-t} \int (R + 1) \omega^n \\ &\leq \frac{d}{dt} \int R \omega^n + C \int (R + n) \omega^n + Ce^{-t} \end{aligned}$$

where we used the uniform bound of scalar curvature and volume of  $\omega(t)$ . The integration of  $R + n$  can be computed as

$$\int (R + n) \omega^n = n \int (\text{Ric} + \omega) \wedge \omega^{n-1} = n \int (-\chi + \hat{\omega}) \wedge \hat{\omega}^{n-1} = ne^{-t} \int (\omega_0 - \chi) \wedge \hat{\omega}^{n-1}.$$

Thus,  $\int (R + n)\omega^n \leq Ce^{-t}$ . Then we have

$$\int_0^\infty \int |\text{Ric} + \omega|^2 \omega(t)^n dt \leq \lim_{t \rightarrow \infty} \int R(t)\omega(t)^n - \int R(0)\omega_0^n + C \leq C.$$

The lemma is proved by this estimate.  $\square$

The estimate (3.3) when  $2 \leq p < 4$  then is a direct consequence of the Hölder inequality

$$\int_t^{t+1} \int_M |\text{Ric} + \omega|^p \leq \left( \int_t^{t+1} \int_M |\text{Ric} + \omega|^4 \right)^{\frac{p-2}{2}} \left( \int_t^{t+1} \int_M |\text{Ric} + \omega|^2 \right)^{\frac{4-p}{2}}.$$

When  $0 < p < 2$  the estimate (3.3) is obvious.

**Remark 3.8.** *M. Simon presented in [25] an  $L^4$  Ricci curvature estimate under Ricci flow on a four-manifold. Combined with the Kähler condition, his arguments can be adapted to give an alternative proof of our estimate. Another related integral bound of curvature can be found in [3].*

### 3.2 Cheeger-Gromov convergence

When the manifold  $M$  is of general type, the Kähler-Ricci flow (3.1) should converge in the Gromov-Hausdorff topology to a singular Kähler-Einstein metric on the canonical model; cf. Conjecture 4.1 in [27]. In this subsection we confirm this convergence without any curvature assumption in the case of the dimension less than or equal to 3.

Recall the holomorphic map  $\pi : M \rightarrow \mathbb{C}P^N$  by a basis of  $H^0(M; K_M^\ell)$  for some large  $\ell$ . Its image  $M_{\text{can}} = \pi(M)$  is called the canonical model of  $M$ . Let  $E \subset M$  be the exceptional locus of  $\pi$ . Then we have

**Theorem 3.9** ([31, 26]). *Let  $M$  be a smooth minimal model of general type and  $\omega(t)$  be any solution to the Kähler-Ricci flow (3.1). Then,*

- (1)  $\omega(t)$  converges in the current sense to a Kähler-Einstein metric  $\omega_\infty$ , the convergence takes place smoothly outside the exceptional locus  $E$ ;
- (2) the metric completion of  $(M \setminus E, \omega_\infty)$  is homeomorphic to  $M_{\text{can}}$ , so it is compact.

**Remark 3.10.** *It is known that the exceptional locus  $E$  coincides with the non-ample or non-Kähler locus of the canonical class; cf. [31] and [14].*

Suppose  $\dim_{\mathbb{C}} M \leq 3$ . Let  $t_i \rightarrow \infty$  be a sequence of times such that

$$\int_M |\text{Ric}_{\omega(t_i)} + \omega(t_i)|^{\frac{7}{2}} \omega(t_i)^n \rightarrow 0, \text{ as } i \rightarrow \infty. \quad (3.22)$$

Choose a regular point  $x_0 \in M \setminus E$ . The volume of the unit ball  $\text{vol}(B_{1, \omega(t_i)}(x_0))$  has a uniform lower bound. By the  $L^p$  extension of Cheeger-Colding-Tian theory, Theorem 2.3, we may assume that  $(M, \omega(t_i), x_0)$  converges in the Cheeger-Gromov sense to a limit metric space  $(M_\infty, d_\infty, x_\infty)$ . Since the metric  $\omega(t)$  converges smoothly on  $M \setminus E$ , we may view  $(M \setminus E, \omega_\infty)$  as a subset of  $(M_\infty, d_\infty)$  through a locally isometric embedding.

Let  $M_\infty = \mathcal{R} \cup \mathcal{S}$  the regular/singular decomposition of  $M_\infty$ .



**Lemma 3.11.** *Suppose  $x \in \mathcal{R}$ ,  $0 < \alpha < 2 - \frac{4n}{7}$ . There exists  $r = r(x) > 0$  such that any  $x_i \in M$  converging to  $x$  has a holomorphic coordinate  $(z^1, \dots, z^n)$  on  $B_{r, \omega(t_i)}(x_i)$  which satisfies*

$$\frac{1}{2} \leq (g_{k\bar{\ell}}) \leq 2, \|g_{k\bar{\ell}}\|_{C^\alpha} \leq 2$$

where  $g_{k\bar{\ell}} = \omega(t_i)(\frac{\partial}{\partial z^k}, \frac{\partial}{\partial \bar{z}^\ell})$ .

*Proof.* The existence of holomorphic coordinates is well-known. It can be constructed by a slight modification of the local harmonic coordinates. We include a short proof for the reader's convenience. First of all, by the  $C^\alpha$  convergence on  $\mathcal{R}$ , there is a sequence of harmonic coordinate  $v_i = (v_i^1, \dots, v_i^{2n})$  on  $B_{r, \omega(t_i)}(x_i)$  for some  $r > 0$  independent of  $i$  such that

$$\frac{3}{4} \leq (h_{pq}) \leq \frac{4}{3} \text{ and } r^{-\alpha} \|h_{pq}\|_{C^\alpha} \leq \frac{4}{3},$$

where  $h_{pq} = g_i(\frac{\partial}{\partial v_i^p}, \frac{\partial}{\partial v_i^q})$ ,  $g_i$  is the Kähler metric of  $\omega(t_i)$ , and [22]

$$\int_{B_{r, \omega(t_i)}(x_i)} |\langle \frac{\partial}{\partial v^p}, \frac{\partial}{\partial v^q} \rangle - \delta_{pq}| \omega(t_i)^n \rightarrow 0, \text{ as } i \rightarrow \infty.$$

Moreover, since each  $\omega(t_i)$  is Kähler, we may assume that

$$\int_{B_{r, \omega(t_i)}(x_i)} |\langle J \frac{\partial}{\partial v^p}, \frac{\partial}{\partial v^{n+q}} \rangle - \delta_{pq}| \omega(t_i)^n \rightarrow 0, \text{ as } i \rightarrow \infty,$$

for any  $1 \leq p, q \leq n$ , where  $J$  is the complex structure of  $M$ ; see [11, Section 9] for a discussion. Then we choose a pseudoconvex domain  $B_{\frac{1}{2}r, \omega(t_i)}(x_i) \subset \Omega_i \subset B_{r, \omega(t_i)}(x_i)$  and solve the  $\bar{\partial}$  equation in  $\Omega_i$ :

$$\bar{\partial} f_i^p = \bar{\partial}(v_i^p + \sqrt{-1}v^{n+p}), p = 1, \dots, n.$$

The domain can be chosen as the Euclidean ball in the local coordinate. The equation has solution satisfying the  $L^2$  estimate [24, Theorem 4.3.4]

$$\int_{B_{\frac{1}{2}r, \omega(t_i)}(x_i)} |f_i^p|^2 \omega(t_i)^n \leq Cr^2 \int_{B_{r, \omega(t_i)}(x_i)} |\bar{\partial}(v_i^p + \sqrt{-1}v^{n+p})|^2 \omega(t_i)^n$$

for a universal constant  $C$ . This implies  $\int_{B_{\frac{1}{2}r, \omega(t_i)}(x_i)} |f_i^p|^2 \omega(t_i)^n \rightarrow 0$  for all  $1 \leq p \leq n$ .

Then applying the elliptic regularity to  $\Delta_{\omega(t_i)} f_i^p = 0$  we get

$$\sup_{B_{\frac{1}{4}r, \omega(t_i)}(x_i)} (|\partial f_i^p| + |\bar{\partial} f_i^p|) \rightarrow 0, \forall 1 \leq p \leq n.$$

In particular, the function  $w_i^p = v^p + \sqrt{-1}v^{n+p} - f_i^p$ ,  $1 \leq p \leq n$ , gives rise to the desired holomorphic coordinate on  $B_{\frac{1}{4}r, \omega(t_i)}(x_i)$  whenever  $i$  is large enough.  $\square$

**Lemma 3.12.**  *$M_\infty \setminus (M \setminus E)$  is a closed subset of  $M_\infty$  with Hausdoeff codimension  $\geq 2$ .*

*Proof.* Notice that  $M \setminus E \subset \mathcal{R}$ ,  $M_\infty \setminus (M \setminus E) = \mathcal{S} \cup (\mathcal{R} \setminus (M \setminus E))$  where  $\text{Codim } \mathcal{S} \geq 4$  by Theorem 2.3 (5). Therefore, it suffices to show that  $\text{Codim } (\mathcal{R} \setminus (M \setminus E)) \geq 2$ .

For any  $x \in \mathcal{R} \setminus (M \setminus E)$  there exists a sequence of points  $x_i \in E$  which converges to  $x$ . By above lemma, there exists local holomorphic coordinate in  $B_{r, \omega(t_i)}(x_i)$  for some  $r = r(x) > 0$  independent of  $i$  with required  $C^\alpha$  estimate. The intersection  $E \cap B_{r, \omega(t_i)}(x_i)$  is a local subvariety with finite volume, so passing to a subsequence,  $E \cap B_{r, \omega(t_i)}(x_i)$  converges to a limit analytical set  $E_\infty \subset B_{r, \omega_\infty}(x)$ . Thus,  $\mathcal{R} \setminus (M \setminus E)$  is an analytical set,  $\text{Codim } (\mathcal{R} \setminus (M \setminus E)) \geq 2$  as desired.  $\square$

**Lemma 3.13.**  $(M_\infty, d_\infty)$  is isometric to the metric completion of  $(M \setminus E, \omega_\infty)$ .

*Proof.* The lemma follows from the fact that  $\text{Codim } (\mathcal{R} \setminus (M \setminus E)) \geq 2$  and the local isoperimetric constant estimate; see [10] or [23] for details. For the estimate of local isoperimetric constant, one can apply the same argument as Croke [15] by using the volume comparison of geodesic balls by Petersen-Wei (cf. [30, Corollary 2.4] or (2.3)).  $\square$

By the compactness of the limit space by Song [26], see Theorem 3.9 (2) above, the diameters of the sequence  $(M, \omega(t_i))$  are uniformly bounded.

**Lemma 3.14.** *The Kähler-Ricci flow  $\omega(t)$  is uniformly noncollapsing in the sense that: there exists  $\kappa = \kappa(n, \omega_0) > 0$  such that*

$$\text{vol}_{\omega(t)}(B_{r, \omega(t)}(x)) \geq \kappa r^{2n}, \forall x \in M, r \leq 1. \quad (3.23)$$

*Proof.* The lemma follows from Perelman's noncollapsing estimate to Ricci flow [20] together with the scalar curvature estimate by Z. Zhang [35]. Suppose that

$$r_i^{-2n} \text{vol}_{\omega(t_i)}(B_{r_i, \omega(t_i)}(x_i)) \rightarrow 0 \quad (3.24)$$

for a sequence of times  $t_i \rightarrow \infty$  and  $x_i \in M$ ,  $r_i \leq 1$ . Choose  $t'_i \in [t_i - 2, t_i - 1]$  such that (3.22) hold at  $t'_i$ . Then by above lemma we have  $(M, \omega(t'_i))$  converges in the Gromov-Hausdorff topology to the unique limit  $(M_\infty, d_\infty)$ . In particular we have

$$R(\omega(t'_i)) \leq C, \text{diam}(M, \omega(t'_i)) \leq C, C_S(M, \omega(t'_i)) \leq C$$

for some  $C$  independent of  $i$ , where  $R$  is the scalar curvature,  $C_S$  is the Sobolev constant.

Let  $\tilde{\omega}_i(\tilde{t}) = (1 + \tilde{t})\omega(t'_i + \log(1 + \tilde{t}))$  be the solution to the Ricci flow

$$\frac{\partial}{\partial \tilde{t}} \tilde{\omega}_i = -\text{Ric}(\tilde{\omega}_i), \tilde{\omega}_i(0) = \omega(t'_i). \quad (3.25)$$

Under this rescalings, the metric  $\omega(t_i) = (1 + \tilde{t}_i)^{-1} \tilde{\omega}_i(\tilde{t}_i)$  for some  $\tilde{t}_i \in [e - 1, e^2 - 1]$  and  $B_{r_i, \omega(t_i)}(x_i) = B_{\tilde{r}_i, \tilde{\omega}_i(\tilde{t}_i)}(x_i)$  for some  $\tilde{r}_i \leq e$ . Recall the  $\mu$  functional of Perelman [20]

$$\mu(g, \tau) = \inf \int_M [\tau(R + |\nabla f|^2) + f - 2n] (4\pi\tau)^{-n} e^{-f} dv_g$$

for any Riemannian metric  $g$  and  $\tau > 0$ , where the infimum is taken over all  $f \in C^\infty(M; \mathbb{R})$  with restriction  $\int_M (4\pi\tau)^{-n} e^{-f} dv_g = 1$ . The condition (3.24) implies that  $\mu(\tilde{\omega}_i(\tilde{t}_i), \tilde{r}_i^2) \rightarrow$

$-\infty$  as  $i \rightarrow \infty$ ; see [20, Section 4.1]. Then Perelman's monotonicity formula shows  $\mu(\tilde{\omega}_i(0), \tilde{t}_i + \tilde{r}_i^2) \rightarrow -\infty$  as  $i \rightarrow \infty$ , where  $\tilde{t}_i + \tilde{r}_i^2 \in [e - 1, 2e^2 - 1]$ . But this can never happen because of the lower estimate of  $\mu$  (cf. the estimate in [36]):

$$\mu(\tilde{\omega}_i(0), \tau) \geq \tau \inf R(\tilde{\omega}_i(0)) - \frac{n}{2} \ln \tau - n \ln C_S(\tilde{\omega}_i(0)) - C(n), \forall \tau \geq \frac{n}{8}. \quad (3.26)$$

So (3.24) can not hold, the lemma is proved.  $\square$

The global Cheeger-Gromov convergence is a direct corollary of the following proposition.

**Proposition 3.15.** *For any sequence  $t_j \rightarrow \infty$ ,  $(M, \omega_{t_j})$  converges along a subsequence in the Cheeger-Gromov sense to the limit  $(M_\infty, d_\infty)$ .*

*Proof.* By the regularity of manifolds with  $L^p$  bounded Ricci curvature, Theorem 2.3, and the uniform  $L^p$  estimate of Ricci curvature under the Kähler-Ricci flow, (3.2) and (3.3), we can find another sequence of times  $t'_j$  such that

$$t_j - \varepsilon_j \leq t'_j \leq t_j$$

where  $\varepsilon_j \rightarrow 0$  as  $j \rightarrow \infty$ , and

$$(M, \omega_{t'_j}) \xrightarrow{d_{GH}} (M_\infty, d_\infty)$$

along a subsequence. On the other hand, by Gromov precompactness theorem [17] together with the local noncollapsing estimate (3.23), after passing to a subsequence, the manifolds  $(M, \omega(t_j))$  also converge in the Gromov-Hausdorff topology to a compact limit

$$(M, \omega(t_j)) \xrightarrow{d_{GH}} (M'_\infty, d'_\infty).$$

It is remained to show that  $(M'_\infty, d'_\infty)$  is isometric to  $(M_\infty, d_\infty)$ . Actually we have

**Claim 3.16.** *There is a sequence of positive numbers  $\delta_j \rightarrow 0$  as  $j \rightarrow \infty$  such that the identity map defines a  $\delta_j$ -Gromov-Hausdorff approximation from  $(M, \omega(t'_j))$  to  $(M, \omega(t_j))$ .*

*Proof of the Claim.* Recall that by the smooth convergence of  $\omega(t)$  on  $M \setminus E$  and uniform volume noncollapsing (3.23) we have for any  $\delta > 0$  one compact subset  $K \subset M \setminus E$  and  $\epsilon > 0$ ,  $j_0 \gg 1$  such that

$$\epsilon \leq \inf_t \text{dist}_{\omega(t)}(K, E) \leq \sup_t \sup_{x \in E} \text{dist}_{\omega(t)}(x, K) \leq \delta, \quad (3.27)$$

and,

$$\text{vol}_{\omega(t)}(M \setminus K) \leq \delta, \forall t \geq t_{j_0}. \quad (3.28)$$

The later can be seen by simply the derivative estimate to volume

$$\frac{d}{dt} \int_{M \setminus K} \omega(t)^n = - \int_{M \setminus K} (\inf R + n) \omega(t)^n \leq C(\omega_0) e^{-t} \int_{M \setminus K} \omega(t)^n$$

where we used the estimate for the scalar curvature (3.21). For example, one may choose  $j_0$  and  $K$  such that  $C(\omega_0)e^{-j_0} \leq \ln 2$  and  $\text{vol}_{\omega(t_{j_0})}(M \setminus K) \leq \frac{\delta}{2}$ , then the volume estimate holds for any  $t \geq t_{j_0}$ . Moreover, by the convexity of the regular set  $M \setminus E$  in  $M_\infty$  we may assume in addition that any minimal geodesic in  $(M, \omega(t))$  with endpoints in  $K$  does not intersect the  $\epsilon$  neighborhood of  $E$ . Also observe that there exists  $C = C(\epsilon)$  independent of  $j$  such that

$$|\text{Ric}(\omega(t))|(x) \leq C, \forall \text{dist}_{\omega(t)}(x, E) \geq \frac{\epsilon}{2}. \quad (3.29)$$

Then, by the derivative estimate to distance function, cf. Lemma 8.3 in [20],

$$\frac{d}{dt} \text{dist}_{\omega(t)}(x, y) \geq -2(2n-1)(C\epsilon + \epsilon^{-1}), \forall x, y \in K.$$

Thus, we obtain the distance lower variation estimate

$$\text{dist}_{\omega(t_j)}(x, y) \geq \text{dist}_{\omega(t'_j)}(x, y) - 2(2n-1)(C\epsilon + \epsilon^{-1})\epsilon_j \geq \text{dist}_{\omega(t'_j)}(x, y) - \sqrt{\epsilon_j}$$

whenever  $j$  is large enough. On the other hand, let  $\gamma$  be any minimal geodesic in  $(M, \omega(t'_j))$  connecting  $x, y \in K$ . By assumption  $\text{dist}_{\omega(t)}(\gamma, E) \geq \epsilon$  for any  $t$ , so

$$\frac{d}{dt} \int_\gamma |\dot{\gamma}| ds \leq \int_\gamma |\text{Ric} + \omega| |\dot{\gamma}| ds \leq (C+n) \int_\gamma |\dot{\gamma}| ds$$

which gives the upper estimate to the distance distortion

$$\text{dist}_{\omega(t_j)}(x, y) \leq e^{(C+n)\epsilon_j} \text{dist}_{\omega(t'_j)}(x, y) \leq \text{dist}_{\omega(t'_j)}(x, y) + \sqrt{\epsilon_j}$$

whenever  $j$  is large enough. Finally, (3.27) shows that  $K$  is  $\delta$ -dense in any  $(M, \omega(t))$ . Combining with the upper and lower distance variation estimate we get that the identity map defines an  $\delta$ -Gromov-Hausdorff approximation between  $(M, \omega(t_j))$  and  $(M, \omega(t'_j))$  whenever  $j$  is large enough such that  $\sqrt{\epsilon_j} \leq \delta$ . The claim is proved.  $\square$

Proposition 3.15 follows directly from the Claim.  $\square$

We end the paper with a brief discussion about the algebraic structure of  $M_\infty$ . By Song's work [26], the limit space  $M_\infty$  coincides with the canonical model  $M_{can}$ , so it is a normal projective variety. On the other hand, using the Kähler-Ricci flow, we can produce a natural isomorphism from  $M_\infty$  to  $M_{can}$  through holomorphic sections of  $K_M^\ell$  for some  $\ell \gg 1$  such that  $K_M^\ell$  is base point free, and consequently, give an alternative proof of the above result by J. Song. This can be done by choosing a family of orthonormal basis of  $H^0(M, K_M^\ell)$  with respect to the Hermitian metric  $h(t) = \omega(t)^{-n\ell}$ , say  $\{s_{i,t}\}_{i=0}^{N_\ell}$  where  $N_\ell = \dim H^0(M; K_M^\ell) - 1$ , which satisfies the ODE system

$$\frac{\partial}{\partial t} s_{i,t} = \sum_j b_{ij}(t) s_{j,t}. \quad (3.30)$$

In order to preserve the orthonormal property we choose

$$b_{ij} = \bar{b}_{ji} = -\frac{\ell-1}{2} \int_M (R+n) \langle s_{i,t}, s_{j,t} \rangle_{h(t)} \omega(t)^n. \quad (3.31)$$

**Lemma 3.17.** *There exists  $C = C(\omega_0, \ell)$  such that*

$$|b_{ij}| \leq Ce^{-t}.$$

*Proof.* First of all we notice that the section  $s_{i,t}$  admits a uniform  $L^\infty$  bound. This can be seen by the uniform equivalence of the Hermitian metric  $h(t) = e^{-\ell u(t)}\Omega^{-\ell}$ , the volume form  $\omega(t)^n = e^{u(t)}\Omega$  and the  $L^\infty$  estimate of holomorphic sections at time  $t = 0$ . Here we used the uniform  $C^0$  estimate of  $u$  (3.8). Then we can estimate the integration as in the proof of Lemma 3.7, by  $R + n \geq -Ce^{-t}$  and  $\int (R + n)\omega^n \leq Ce^{-t}$ ,

$$|b_{ij}| \leq C \int |R + n|\omega^n \leq C \int (R + n + Ce^{-t})\omega^n + Ce^{-t} \leq Ce^{-t}$$

where  $C = C(\omega_0, \ell)$ . □

Suppose  $s_{i,t} = \sum_j a_{ij}(t)s_{j,0}$  and denote by  $A(t) = (a_{ij}(t))$  a Hermitian matrix, then  $A(t) = e^{\int_0^t B(s)ds}$  where  $B(t) = (b_{ij}(t))$ . Thus the sections  $\{s_{i,t}\}_{i=0}^{N_\ell}$  converge to a set of holomorphic sections  $\{s_{i,\infty}\}_{i=0}^{N_\ell}$  which forms another basis of  $H^0(M; K_M^\ell)$ . It induces a morphism

$$\Phi : M \rightarrow M_{can} \subset \mathbb{C}P^{N_\ell}.$$

On the other hand, we also have a uniform gradient estimate to each  $s = s_{i,0}$ , cf. [26],

$$|\nabla^{h(t)} s|_{h(t) \otimes \omega(t)} = |\nabla^{\Omega^{-\ell}} s + \partial u \otimes s|_{h(t) \otimes \omega(t)} \leq C|\nabla^{\Omega^{-\ell}} s|_{\Omega^{-\ell} \otimes \chi} + C|\partial u|_{\omega(t)}|s|_{\Omega^{-\ell}} \leq C$$

where we used the uniform  $C^1$  estimate of  $u$  and  $\chi \leq C\omega(t)$ , this leads to a convergence of  $\{s_{i,t}\}_{i=0}^{N_\ell}$  under the Cheeger-Gromov convergence, so  $\{s_{i,\infty}\}_{i=0}^{N_\ell}$  can also be seen as a set of holomorphic sections of  $K_{M_\infty}^\ell$ . It is obvious that  $\{s_{i,\infty}\}_{i=0}^{N_\ell}$  is base point free on  $M_\infty$ , so it defines a map

$$\Phi_\infty : M_\infty \rightarrow \mathbb{C}P^{N_\ell}.$$

Finally, through construction of local peak sections, up to rising a power of  $\ell$ , the map  $\Phi_\infty$  separates points of  $M_\infty$ , so it defines a homeomorphism.

## References

- [1] M. Anderson, *Ricci curvature bounds and Einstein metrics on compact manifolds*, J. Amer. Math. Soc. 2 (1989), 455-490.
- [2] M. Anderson, *Convergence and rigidity of manifolds under Ricci curvature bounds*, Invent. Math., 102 (1990), 429-445.
- [3] R. H. Bamler and Q. S. Zhang, *Heat kernel and curvature bounds in Ricci flows with bounded scalar curvature*, arXiv:1501.02191v2
- [4] S. Bando, A. Kasue and H. Nakajima, *On a construction of coordinates at infinity on manifolds with fast curvature decay and maximal volume growth*, Invent. Math. 97 (1989), no. 2, 313-349.

- [5] A. L. Besse, *Einstein manifolds*, Springer-Verlag Berlin Herdelberg GmbH
- [6] H.D. Cao, *Deformation of Kähler metrics to Kähler-Einstein metrics on compact Kähler manifolds*, *Invent. Math.*, 81 (1985), 359-372.
- [7] J. Cheeger, *Integral bounds on curvature, elliptic estimates and rectifiability of singular sets*, *Geom. Funct. Anal.*, 13 (2003), 20-72.
- [8] J. Cheeger and T. H. Colding, *Lower bounds on the Ricci curvature and the almost rigidity of warped products*, *Ann. Math.*, 144 (1996), 189-237.
- [9] J. Cheeger and T. H. Colding, *On the structure of spaces with Ricci curvature bounded below I*, *J. Diff. Geom.*, 46 (1997), 406-480.
- [10] J. Cheeger and T. H. Colding, *On the structure of spaces with Ricci curvature bounded below II*, *J. Diff. Geom.*, 54 (2000), 13-35.
- [11] J. Cheeger, T. H. Colding and G. Tian, *On the singularities of spaces with bounded Ricci curvature*, *Geom. Funct. Anal.*, 12 (2002), 873-914.
- [12] T. H. Colding, *Ricci curvature and volume convergece*, *Ann. of Math.*, 145 (1997), 477-501.
- [13] T. H. Colding and A. Naber, *Sharp Hölder continuity of tangent cones for spaces with a lower Ricci curvature bound and applications*, *Ann. of Math.*, 176 (2012), 1173-1229.
- [14] T. C. Collins and V. Tosatti, *Kähler currents and null loci*, arXiv:1304.5216v5
- [15] C. B. Croke, *Some isoperimetric inequalities and eigenvalue estimates*, *Ann. Sci. Éc. Norm. Sup. Paris*, 13 (1980), 419-435.
- [16] X.Z. Dai and G.F. Wei, *Comparison geometry for Ricci curvature*, preprint.
- [17] M. Gromov, *Metric structures for Riemannian and non-Riemannian spaces*. With appendices by M. Katz, P. Pansu and S. Semmes. Translated from the French by Sean Michael Bates. Birkhäuser Boston, Inc., Boston, MA, 2007.
- [18] B. Guo, *On the Kähler Ricci flow on projective manifolds of general type*, arXiv:1501.04239
- [19] B. Guo, J. song and B. Weinkovw, *Geometric convergence of the Kähler-Ricci flow on complex surfaces of general type*, arXiv:1505.00705
- [20] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications*, arXiv:math.DG/0211159
- [21] P. Petersen and G.F. Wei, *Relative volume comparison with integral curvature bounds*, *Geom. Funct. Anal.*, 7 (1997), 1031-1045.
- [22] P. Petersen and G.F. Wei, *Analysis and geometry on manifolds with integral Ricci curvature bounds, II*, *Trans. AMS.*, 353 (2001), 457-478.

- [23] X.C. Rong and Y.G. Zhang, *Continuity of extremal transitions and flops for Calabi-Yau manifolds*, J. Diff. Geom., 89 (2011), 233-269.
- [24] S.C. Chen and M.C. Shaw, *Partial differential equations in several complex variables*, Amer. Math. Soc., Providence, RI; Intern. Press, Boston, MA, 2001.
- [25] M. Simon, *Some integral curvature estimates for the Ricci flow in four dimensions*, arXiv:1514.02623v1
- [26] J. Song, *Riemannian geometry of Kähler-Einstein currents*, arXiv:1404.0445
- [27] J. Song, *Riemannian geometry of Kähler-Einstein currents II: an analytic proof of Kawamata's base point free theorem*, arXiv:1409.8374
- [28] J. Song and G. Tian, *Bounding scalar curvature for global solutions of the Kähler-Ricci flow*, arXiv:1111.5681v1
- [29] G. Tian, *On Calabi's conjecture for complex surfaces with positive first Chern class*. Invent. Math., 101, (1990), 101-172.
- [30] G. Tian and Z.L. Zhang, *Regularity of Kähler-Ricci flows on Fano manifolds*, arXiv:1310.5897
- [31] G. Tian and Z. Zhang, *On the Kähler-Ricci flow on projective manifolds of general type*, Chinese Ann. Math. Ser. B, 27 (2006), 179-192.
- [32] H. Tsuji, *Existence and degeneration of Kähler-Einstein metrics on minimal algebraic varieties of general type*, Math. Ann., 281 (1988), 123-133.
- [33] Q.S. Zhang, *A uniform Sobolev inequality under Ricci flow*, Inter. math. Res. Notices, 2007.
- [34] Y.G. Zhang, *Miyaoka-Yau inequality for minimal projective manifolds of general type*, Proc. Amer. Math. Soc. 137 (2009), no. 8, 2749C2754.
- [35] Z. Zhang, *Scalar curvature bound for Kähler-Ricci flows over minimal manifolds of general type*, Int. Math. Res. Not., 2009, 3901-3912.
- [36] Z.L. Zhang, *Compact blow-up limits of finite time singularities of Ricci flow are shrinking Ricci solitons*, C. R. Acad. Sci. Paris, Ser. I 345 (2007), 503-506.