

# The Killing tensors on an $n$ -dimensional manifold with $SL(n, \mathbf{R})$ -structure

S. E. Stepanov, I. I. Tsyganok <sup>1</sup>

*Finance University, Moscow, Russia*

## Abstract

In this paper we solve the problem of finding integrals of equations determining the Killing tensors on an  $n$ -dimensional differentiable manifold  $M$  endowed with an equiaffine  $SL(n, \mathbf{R})$ -structure and discuss possible applications of obtained results in Riemannian geometry.

**Keywords:** differentiable manifold,  $SL(n, \mathbf{R})$ -structure, Killing tensors.

**AMS:** 53A15; 53A45

## §1. Introduction

**1.1.** The "structural point of view" of affine differential geometry was introduced by K. Nomizu in 1982 in a lecture at Münster University with the title "What is Affine Differential Geometry?" (see [No 1]). In the opinion of K. Nomizu, the geometry of a manifold  $M$  endowed with an *equiaffine structure* is called affine differential geometry.

In recent years, there has been a new wave of papers devoted to affine differential geometry. Today the number of publications (including monographs) on affine differential geometry reached a considerable level. The main part of these publications is devoted to geometry of hypersurfaces (see [Sch] and [SSV] for the history and references).

**1.2.** In the present paper we solve the problem of finding integrals of equations determining the Killing tensors (see [KSCH] for the definitions, properties and applications) on an  $n$ -dimensional differentiable manifold  $M$  endowed with an equiaffine structure. The paper is a direct continuation of [St 2]. The same notations are used here.

The first of two present theorems proved in our paper is an affine analog of the statement published in the paper [St 1], which appeared in the process of solving problems in General relativity.

## §2. Definitions and results

---

<sup>1</sup>emails: s.e.stepanov@mail.ru; i.i.tsyganok@mail.ru

**2.1.** In order to clarify the approach to problem of finding integrals of equations determining the Killing tensors on an  $n$ -dimensional differentiable manifold  $M$  we shall start with a brief introduction to the subject which emphasizes the notion of an equiaffine  $SL(n, \mathbb{R})$ -structure.

Let  $M$  be a connected differentiable manifold of dimension  $n$  ( $n > 2$ ), and let  $L(M)$  be the corresponding bundle of linear frames with structural group  $GL(n, \mathbb{R})$ . We define  $SL(n, \mathbb{R})$ -structure on  $M$  as a principal  $SL(n, \mathbb{R})$ -subbundle of  $L(M)$ . It is well known that an  $SL(n, \mathbb{R})$ -structure is simply a volume element on  $M$ , i.e. an  $n$ -form  $\eta$  that does not vanishing anywhere (see [Ko], Chapter I, §2).

We recall the famous problem of the existence of a uniquely determined linear connection  $\nabla$  reducible to  $G$  for each given  $G$ -structure on  $M$  (see [C], p. 213). For example, if  $M$  is a manifold with a pseudo-Riemannian metric  $g$  of an arbitrary index  $k$ , then the bundle  $L(M)$  admits a unique linear connection  $\nabla$  without torsion that is reducible to  $O(m, k)$ -structure. Such a connection is called the *Levi-Civita connection*. It is characterized by the following condition  $\nabla g = 0$ .

A linear connection  $\nabla$  having zero torsion and reducible to  $SL(n, \mathbb{R})$  is said to be *equiaffine* and can be characterized by the following equivalent conditions (see [Sch], p 99; [SSV], pp. 57-58):

(1)  $\nabla \eta = 0$ ;

(2) the Ricci tensor  $Ric$  of  $\nabla$  is symmetric; that means  $Ric(X, Y) = Ric(Y, X)$  for any vector fields  $X, Y \in C^\infty TM$ .

An *equiaffine  $SL(n, \mathbb{R})$ -structure* or an *equiaffine structure* on an  $n$ -dimensional differentiable manifold  $M$  is a pair  $(\eta, \nabla)$ , where  $\nabla$  is a linear connection with zero torsion and  $\eta$  is a volume element which is parallel relative to  $\nabla$  (see [No 2], p. 43).

The curvature tensor  $R$  of an equiaffine connection  $\nabla$  admits a point-wise  $SL(n, \mathbb{R})$ -invariant decomposition of the form  $R = (n - 1)^{-1}[id_M \otimes Ric - Ric \otimes id_M] + W$  where  $W$  is the *Weyl projective curvature tensor* (see [SSV], p. 73-74; [E], §40). Then two classes of equiaffine structures can be distinguished in accordance with this decomposition: the *Ricci-flat* equiaffine  $SL(n, \mathbb{R})$ -structures for which  $Ric = 0$ , and the *equiprojective  $SL(n, \mathbb{R})$ -structures* for which

$$R = (n - 1)^{-1}[id_M \otimes Ric - Ric \otimes id_M]. \tag{2.1}$$

**Remark 1.** Recall that a linear connection  $\nabla$  with zero torsion is called *Ricci-flat* if the Ricci tensor  $Ric = 0$  (see [M]). On the another hand, a connection  $\nabla$  is called *equiprojective* if

the Weyl projective curvature tensor  $W = 0$  (see [Sch], §18). In the literature equiprojective connections sometimes are called *projectively flat* (see, for example, [SSV], p. 73).

An autodiffeomorphism of the manifold  $M$  is an automorphism of  $SL(n, \mathbb{R})$ -structure if and only if it preserves the volume element  $\eta$ . Let  $X$  be a vector field on  $M$ . The function  $divX$  defined by the formula  $(divX)\eta = L_X\eta$  where  $L_X$  is the Lie differentiation in the direction of the vector field  $X$  is called the divergence of  $X$  with respect to the  $n$ -form  $\eta$  (see [KN], Appendix no. 6). Obviously,  $X$  is an infinitesimal automorphism of an  $SL(n, \mathbb{R})$ -structure if and only if  $divX = 0$ . Such a vector field  $X$  is said to be *solenoidal*.

For an arbitrary vector field  $X$  on  $M$  with a linear connection  $\nabla$  we can introduce the tensor field  $A_X = L_X - \nabla_X$  regarded as a field of linear endomorphisms of the tangent bundle  $TM$ . If  $M$  is an  $n$ -dimensional with an equiaffine  $SL(n, \mathbb{R})$ -structure then the formula  $traceA_X = -divX$  can be verified directly (see [KN], Appendix no. 6).

We have the  $SL(n, \mathbb{R})$ -invariant decomposition  $A_X = -n^{-1}(divX)id_M + \dot{A}_X$  at every point  $x \in M$ .

Two classes of vector fields on  $M$  endowed with an equiaffine  $SL(n, \mathbb{R})$ -structure can be distinguished in accordance with this decomposition: the *solenoidal vector fields* and the *concircular vector fields* for which, by definition (see [Sc], p. 322; [M]), we have  $A_X = -n^{-1}(divX)id_M$ .

The integrability conditions of the structure equation  $A_X = -n^{-1}(divX)id_M$  of the concircular vector field  $X$  is the Ricci's identity  $Y(divX)Z - Z(divX)Y = nR(Y, Z)X$  for any vector fields  $Y, Z \in C^\infty TM$  (see [E], §11). This identity are equivalent to the condition  $W(Y, Z)X = 0$  for any vector fields  $Y, Z \in C^\infty TM$ . It follows that an equiaffine  $SL(n, \mathbb{R})$ -structure on an  $n$ -dimensional manifold  $M$  is equiprojective if and only if there exist  $n$  linearly independent concircular vector fields  $X_1, X_2, \dots, X_p$  on  $M$  (see also [ST]). This statement is an affine analog of the well known fact for the Riemannian manifold  $M$  of constant sectional curvature (see [F]).

**Remark 2.** A pseudo-Riemannian manifold  $(M, g)$  with a projectively flat Levi-Civita connection  $\nabla$  is a manifold of constant section curvature (see [Sch], §18). Therefore a manifold  $M$  endowed with an equiprojective  $SL(n, \mathbb{R})$ -structure is an affine analog of a pseudo-Riemannian manifold of constant sectional curvature.

**2.2.** We consider an  $n$ -dimensional manifold  $M$  with an equiaffine  $SL(n, \mathbb{R})$ -structure and denote by  $\Lambda^p M$  ( $1 \leq p \leq n-1$ ) the  $p^{th}$  exterior power  $\Lambda^p(T^*M)$  of the cotangent bundle  $T^*M$  of  $M$ . Hence  $C^\infty \Lambda^p M$ , the space of all  $C^\infty$ -sections of  $\Lambda^p M$ , is the space of skew-symmetric covariant tensor fields of degree  $p$  ( $1 \leq p \leq n-1$ ).

Let  $\gamma : J \subset \mathbf{R} \rightarrow M$  be an arbitrary geodesic on  $M$  with affine parameter  $t \in J$ . In this case, we have  $\nabla_{\frac{d\gamma}{dt}} \frac{d\gamma}{dt} = 0$  for the tangent vector  $\frac{d\gamma}{dt}$  of  $\gamma$ .

**Definition 1** (see [St 2]). *A skew-symmetric tensor field  $\omega \in C^\infty \Lambda^p M$  ( $1 \leq p \leq n - 1$ ) on an  $n$ -dimensional manifold  $M$  with an equiaffine  $SL(n, \mathbf{R})$ -structure is called Killing-Yano tensor of degree  $p$  if the tensor  $i_{\frac{d\gamma}{dt}} \omega := \text{trace} \left( \frac{d\gamma}{dt} \otimes \omega \right)$  is parallel along an arbitrary geodesic  $\gamma$  on  $M$ .*

From this definition we conclude that  $\left( \nabla_{\frac{d\gamma}{dt}} \omega \right) \left( \frac{d\gamma}{dt}, X_2, \dots, X_p \right) = 0$  for any vector fields  $X_2, \dots, X_p \in C^\infty TM$ . Since the geodesic  $\gamma$  may be chosen arbitrary, the above relation is possible if and only if  $\nabla \omega \in C^\infty \Lambda^{p+1} M$ , which is equivalent to  $d\omega = (n+1)\nabla \omega$  for the exterior differential operator  $d : C^\infty \Lambda^p M \rightarrow C^\infty \Lambda^{p+1} M$ .

Obviously, the set of Killing-Yano tensors of degree  $p$  ( $1 \leq p \leq n - 1$ ) constitutes an  $\mathbf{R}$ -module of tensor fields on  $M$ , denoted by  $\mathbf{K}^p(M, \mathbf{R})$ .

Let  $X_1, \dots, X_p$  be  $p$  linearly independent concircular vector fields on  $M$  ( $1 \leq p \leq n - 1$ ). Then direct inspection shows that the tensor field  $\omega$  of degree  $n - p$  dual to the tensor field  $\text{alt}\{X_1 \otimes \dots \otimes X_p\}$  relative to the  $n$ -form  $\eta$  is a Killing-Yano tensor (see also [St 2]). Therefore on any  $n$ -manifold  $M$  with equiprojective  $SL(n, \mathbf{R})$ -structure, there exist at least  $n!/[p!(n-p)]^{-1}$  linearly independent Killing-Yano tensors (see [St 2]). Moreover the following theorem is true.

**Theorem 1.** *On an  $n$ -dimensional manifold  $M$  endowed with an equiprojective  $SL(n, \mathbf{R})$ -structure  $(\eta, \nabla)$ , there exist a local coordinate system  $x^1, \dots, x^n$  in which an arbitrary Killing-Yano tensor  $\omega$  of degree  $p$  ( $1 \leq p \leq n - 1$ ) has the components*

$$\omega_{i_1 \dots i_p} = e^{(p+1)\psi} (A_{i_0 i_1 \dots i_p} x^{i_0} + B_{i_1 \dots i_p}) \quad (2.1)$$

where  $A_{i_0 i_1 \dots i_p}$  and  $B_{i_1 \dots i_p}$  are arbitrary constants skew-symmetric w.r.t. all their indices and  $\psi = (n+1)^{-1} \ln(\eta)$ .

From the theorem we conclude that the maximum of linearly independent the Killing-Yano tensors is by calculating the number  $K_n^p$  of independent  $A_{i_0 i_1 \dots i_p}$  and  $B_{i_1 \dots i_p}$  which exist after accounting for the symmetries on the indices. It follows that  $K_n^p = \frac{(n+1)!}{(p+1)!(n-p)!}$  is the maximum number linearly independent the Killing-Yano tensors.

**Corollary 1.** *Let  $M$  be an  $n$ -dimensional manifold endowed with an equiprojective  $SL(n, \mathbf{R})$ -structure then*

$$\dim K^p(M, \mathbf{R}) = \frac{(n+1)!}{(p+1)!(n-p)!}.$$

On our fixed manifold  $M$  with an equiaffine  $SL(n, \mathbf{R})$ -structure, we denote by  $S^p M$  the bundle of symmetric covariant tensor fields of degree  $p$  on  $M$ . Hence  $C^\infty S^p M$ , the space of all  $C^\infty$ -sections of  $S^p M$ , is the space symmetric covariant tensor fields of degree  $p$ .

**Definition 2** (see [St 2]). *A symmetric tensor field  $\varphi \in C^\infty S^p M$  on an  $n$ -dimensional manifold  $M$  with an equiaffine  $SL(n, \mathbf{R})$ -structure is called Killing tensor of degree  $p$  if  $\varphi \left( \frac{d\gamma}{dt}, \dots, \frac{d\gamma}{dt} \right) = \text{const}$  along an arbitrary geodesic  $\gamma$  on  $M$ .*

Let  $\varphi \left( \frac{d\gamma}{dt}, \dots, \frac{d\gamma}{dt} \right) = \text{const}$  along an arbitrary geodesic  $\gamma$  on  $M$  and hence  $\varphi$  is a Killing tensor. Then the above relation is possible if and only if  $\delta^* \varphi := \sum_{\text{cicl}} \{ \nabla \varphi \} = 0$  where for the local components  $\nabla_{i_0} \varphi_{i_1 \dots i_p}$  of  $\nabla \varphi$  we define the sum  $\sum_{\text{cicl}} \{ \nabla_{i_0} \varphi_{i_1 \dots i_p} \}$  as the sum of the terms obtained by a cyclic permutation of the indices  $i_0, i_1, \dots, i_p$ .

Obviously, the set of Killing tensors constitutes an  $\mathbf{R}$ -module of tensor fields on  $M$ , denoted by  $\mathbf{T}^p(M, \mathbf{R})$ .

Let  $M$  be an  $n$ -dimensional manifold endowed with an equiprojective  $SL(n, \mathbf{R})$ -structure  $(\eta, \nabla)$ , and  $\omega_1, \dots, \omega_p$  be  $p$  linearly independent Killing-Yano tensors of degree 1 on  $M$ . Then direct inspection shows that the tensor field  $\varphi := \text{sym}\{\omega_1 \otimes \dots \otimes \omega_p\}$  is a Killing tensor of degree  $p$ . Therefore on any  $n$ -manifold  $M$  with equiprojective  $SL(n, \mathbf{R})$ -structure, there exist at least  $(n + p - 1)! [p!(n - 1)!]^{-1}$  linearly independent Killing tensors (see also [SS]). Moreover the following theorem is true.

**Theorem 2.** *On an  $n$ -dimensional manifold  $M$  endowed with an equiprojective  $SL(n, \mathbf{R})$ -structure  $(\eta, \nabla)$ , there exist a local coordinate system  $x^1, \dots, x^n$  in which the components  $\varphi_{i_1 \dots i_p}$  of an arbitrary Killing tensor  $\varphi$  of degree  $p$  can be expressed in the form of an  $p^{\text{th}}$  degree polynomial in the  $x^i$ 's*

$$\varphi_{i_1 \dots i_p} = e^{2p\psi} \sum_{q=0}^p A_{i_1 \dots i_p j_1 \dots j_q} x^{j_1} \dots x^{j_q} \quad (2.2)$$

where the coefficients  $A_{i_1 \dots i_p j_1 \dots j_q}$  are constant and symmetric in the set of indices  $i_1, \dots, i_p$  and the set of indices  $j_1, \dots, j_q$ . In addition to these properties the coefficients  $A_{i_1 \dots i_p j_1 \dots j_q}$  have the following symmetries

$$\sum_{\text{cicl}} \{ A_{i_1 \dots i_p j_1 \dots j_{p-s}} \}_{j_{p-s+1}} = 0 \quad (2.3)$$

for  $s = 1, \dots, p - 1$  and

$$\sum_{\text{cicl}} \{ A_{i_1 \dots i_p j_1} \} = 0. \quad (2.4)$$

From the theorem we conclude that the maximum number of linearly independent the Killing

tensors is obtained by calculating the number  $T_n^p$  of independent  $A_{i_1 \dots i_p j_1 \dots j_q}$  ( $q = 0, 1, \dots, n$ ) which exist after accounting for the symmetries on the indices the dependence relations (2.3) and (2.4). By [KL] it follows that

$$T_n^p = \frac{p(p+1)^2(p+2)^2 \dots (m+p-1)^2(m+p)}{(p+1)!p!}$$

is the maximum number linearly independent the Killing-Yano tensors. Then we have the following proposition.

**Corollary 2.** *Let  $M$  be an  $n$ -dimensional manifold endowed with an equiprojective  $SL(n, \mathbf{R})$ -structure then*

$$\dim T^p(M, \mathbf{R}) = \frac{p(p+1)^2(p+2)^2 \dots (m+p-1)^2(m+p)}{p!(p+1)!}.$$

### §3. Proofs of theorems

**3.1.** We let  $f : \bar{M} \rightarrow M$  denote the mapping of an  $\bar{n}$ -dimensional manifold  $\bar{M}$  endowed with an equiaffine  $SL(\bar{n}, \mathbf{R})$ -structure onto another an  $n$ -dimensional manifold  $M$  endowed with an equiaffine  $SL(n, \mathbf{R})$ -structure, and let  $f_*$  be the differential of this mapping. For any covariant tensor field  $\omega$  on  $M$ , we can then define the covariant tensor field  $f^*\omega$  on  $\bar{M}$ , where  $f^*$  is the transformation transposed to the transformation  $f_*$ .

If  $\dim \bar{M} = \dim M = n$  and  $f : \bar{M} \rightarrow M$  is a projective diffeomorphism, i.e., a mapping that transforms an arbitrary geodesic in  $\bar{M}$  into a geodesic in  $M$ , then we have the following lemma.

**Lemma 1.** *Let  $f : \bar{M} \rightarrow M$  be a projective diffeomorphism of  $n$ -dimensional manifolds endowed with the equiaffine  $SL(n, \mathbf{R})$ -structures  $(\bar{\eta}, \bar{\nabla})$  and  $(\eta, \nabla)$  respectively. Then for an arbitrary Killing-Yano tensor  $\omega$  of degree  $p$  ( $1 \leq p \leq n-1$ ) on the manifold  $M$  the tensor field  $\bar{\omega} = e^{-(p+1)\psi}(f^*\omega)$  with  $\psi = (n+1)^{-1} \ln(\eta/\bar{\eta})$  will be the Killing-Yano tensor of degree  $p$  on the manifold  $\bar{M}$ .*

**Proof.** It is known that the diffeomorphism  $f : \bar{M} \rightarrow M$  can be realized following the principle of equality of the local coordinates  $\bar{x}^1 = x^1, \dots, \bar{x}^n = x^n$  at the corresponding points  $\bar{x}$  and  $x = f(\bar{x})$  of these manifolds. In this case, we have the equalities (see [Sch], §18; [M] and [YI])

$$\Gamma_{ij}^k = \bar{\Gamma}_{ij}^k + \psi_i \delta_j^k + \psi_j \delta_i^k \quad (3.1)$$

for the objects  $\Gamma_{ij}^k$  and  $\bar{\Gamma}_{ij}^k$  of the a equiaffine connections  $\nabla$  and  $\bar{\nabla}$  in the coordinate system  $x^1, \dots, x^n$  that is common w. r. t. the mapping  $f : \bar{M} \rightarrow M$ , and for the gradient  $\psi_j = (n+1)^{-1} \partial_j \ln[\eta/\bar{\eta}]$ .

Equalities (3.1) imply that the mapping  $f^{-1}$ , which is inverse to the projective diffeomorphism  $f : \bar{M} \rightarrow M$ , is a projective mapping.

We set  $\omega_{i_1 \dots i_p}$  be the local components of a Killing-Yano tensor  $\omega$  of degree  $p$  ( $1 \leq p \leq n-1$ ) arbitrary defined on the manifold  $M$ ; by definition, these components satisfy the equations

$$\nabla_{i_0} \omega_{i_1 \dots i_p} + \nabla_{i_1} \omega_{i_0 \dots i_p} = 0. \quad (3.2)$$

From equalities (3.2) we find directly that the components

$$\bar{\omega}_{i_1 \dots i_p} = e^{-(p+1)\psi} \omega_{i_1 \dots i_p} \quad (3.3)$$

of the tensor field  $\bar{\omega} = e^{-(p+1)\psi} (f^* \omega)$  satisfy the equations

$$\bar{\nabla}_{i_0} \bar{\omega}_{i_1 \dots i_p} + \bar{\nabla}_{i_1} \bar{\omega}_{i_0 \dots i_p} = 0. \quad (3.4)$$

Hence, the tensor field  $\bar{\omega}$  is a Killing-Yano tensor of degree  $p$  ( $1 \leq p \leq n-1$ ) on the manifold  $\bar{M}$ .

**3.2.** Let  $\mathbf{A}^n$  be an  $n$ -dimensional affine space with a volume element given by the determinant:  $\det(e_1, \dots, e_n) = 1$ , where  $\{e_1, \dots, e_n\}$  is the standard basis of the underlying vector space for  $\mathbf{A}^n$ . We denote by  $\nabla$  the standard linear connection in  $\mathbf{A}^n$  relative to which the volume element "det" is parallel (see [No 2]; [SSV], p. 10).

Let  $f : \bar{M} \rightarrow \mathbf{A}^n$  be a projective diffeomorphism from a manifold  $\bar{M}$  endowed with equiaffine  $SL(n, \mathbf{R})$ -structure onto an affine space  $\mathbf{A}^n$  endowed with standard equiaffine  $SL(n, \mathbf{R})$ -structure. It is well known that manifolds endowed with equiprojective  $SL(n, \mathbf{R})$ -structures and only these manifolds are projectively diffeomorphic to an affine space  $\mathbf{A}^n$  (see [Sch], §18; [M]) therefore in our case the  $SL(n, \mathbf{R})$ -structure of the manifold  $\bar{M}$  must be an equiprojective structure.

If  $\mathbf{A}^n$  is an affine space with the Cartesian system of coordinates  $\bar{x}_1, \dots, \bar{x}^n$  then the components  $\bar{\omega}_{i_1 \dots i_p}$  of the Killing-Yano tensor  $\bar{\omega}$  of degree  $p$  ( $(1 \leq p \leq n-1)$ ) in equation (3.4) must now satisfy

$$\partial_j \bar{\omega}_{i_1 \dots i_p} + \partial_i \bar{\omega}_{j i_1 \dots i_p} = 0 \quad (3.5)$$

where  $\partial_j = \frac{\partial}{\partial \bar{x}^j}$ . From (3.5) we conclude the following equations

$$\partial_k \partial_j \bar{\omega}_{i_1 \dots i_p} + \partial_k \partial_i \bar{\omega}_{j i_1 \dots i_p} = 0; \quad (3.6)$$

$$\partial_j \partial_i \bar{\omega}_{ki_1 \dots i_p} + \partial_j \partial_k \bar{\omega}_{ii_1 \dots i_p} = 0; \quad (3.7)$$

$$\partial_i \partial_k \bar{\omega}_{ji_1 \dots i_p} + \partial_i \partial_j \bar{\omega}_{ki_1 \dots i_p} = 0. \quad (3.8)$$

From (3.6), (3.7), (3.8) we find

$$\partial_k \partial_j \bar{\omega}_{i_1 i_2 \dots i_p} = 0, \quad (3.9)$$

by using identities  $\frac{\partial^2 h}{\partial \bar{x}^k \partial \bar{x}^j} = \frac{\partial^2 h}{\partial \bar{x}^j \partial \bar{x}^k}$  which are carried out for an arbitrary smooth function  $h : \mathbf{A}^n \rightarrow \mathbf{R}$ . The integrals of equations (3.9) take the form

$$\bar{\omega}_{i_1 \dots i_p} = A_{i_0 i_1 \dots i_p} \bar{x}^{i_0} + B_{i_1 \dots i_p} \quad (3.10)$$

for any skew-symmetric constants  $A_{i_0 i_1 \dots i_p}$  and  $B_{i_1 \dots i_p}$  (see also [SS] and [St 3]). Taking the components (3.10) of the Killing-Yano tensor  $\bar{\omega}$  in  $\mathbf{A}^n$  and using Lemma 1, we can formulate Theorem 1.

**3.3.** Let  $\bar{M}$  be a manifold of dimension  $n$  endowed with the equiaffine  $SL(n, \mathbf{R})$ -structure  $(\bar{\eta}, \bar{\nabla})$  and  $M$  be a manifold of some dimension endowed with the equiaffine  $SL(n, \mathbf{R})$ -structure  $(\eta, \nabla)$ . Let there is given a projective diffeomorphism  $f : \bar{M} \rightarrow M$ , then we have the following lemma.

**Lemma 2.** *Let  $f : \bar{M} \rightarrow M$  be a projective diffeomorphism of  $n$ -dimensional manifolds endowed with the equiaffine  $SL(n, \mathbf{R})$ -structures  $(\bar{\eta}, \bar{\nabla})$  and  $(\eta, \nabla)$  respectively. Then for an arbitrary Killing tensor  $\varphi$  of degree  $p$  on the manifold  $M$  the tensor field  $\bar{\varphi} = e^{-2p\psi}(f^*\varphi)$  with  $\psi = (n+1)^{-1} \ln(\eta/\bar{\eta})$  will be the Killing tensor of degree  $p$  on the manifold  $\bar{M}$ .*

**Proof.** We set  $\varphi_{i_1 \dots i_p}$  to be components of the Killing tensor  $\varphi$  arbitrary defined on the manifold  $M$ ; by definition, these components satisfy the following equations  $\sum_{cicl} \{\nabla_{i_0} \varphi_{i_1 \dots i_p}\} = 0$ . Then we find directly that the components  $\bar{\varphi}_{i_1 \dots i_p} = e^{-2p\psi} \varphi_{i_1 \dots i_p}$  of the tensor  $\bar{\varphi} = e^{-2p\psi} \varphi$  satisfy the equations

$$\sum_{cicl} \{\bar{\nabla}_{i_0} \bar{\varphi}_{i_1 \dots i_p}\} = e^{-2p\psi} \sum_{cicl} \{\nabla_{i_0} \varphi_{i_1 \dots i_p}\} = 0. \quad (3.11)$$

From (3.11) we conclude that the tensor field  $\bar{\varphi}$  is a Killing tensor of degree  $p$  on the manifold  $\bar{M}$ .

**3.4.** It follows from Nijenhuis (see [Ni]) that in an  $n$ -dimensional affine space  $\mathbf{A}^n$  the components  $\bar{\varphi}_{i_1 \dots i_p}$  of the Killing tensor  $\bar{\varphi}$  of degree  $p$  can be expressed in the form of an  $p^{th}$  degree polynomial in the  $\bar{x}^i$ 's

$$\varphi_{i_1 \dots i_p} = e^{-2p\psi} \sum_{q=0}^p A_{i_1 \dots i_p j_1 \dots j_q} \bar{x}^{j_1} \dots \bar{x}^{j_q}. \quad (3.12)$$

The coefficients  $A_{i_1 \dots i_p j_1 \dots j_q}$  are constant and symmetric in the set of indices  $i_1, \dots, i_p$  and the set of indices  $j_1, \dots, j_q$ . In addition to these properties the coefficients  $A_{i_1 \dots i_p j_1 \dots j_q}$  have the following symmetries  $\sum_{cicl} \{A_{i_1 \dots i_p j_1 \dots j_{p-s}}\}_{j_{p-s+1}} = 0$  for  $s = 1, \dots, p-1$  and  $\sum_{cicl} \{A_{i_1 \dots i_p j_1} = 0\}$ . Taking the components (3.12) of the Killing tensor  $\bar{\varphi}$  in  $\mathbf{A}^n$  and using Lemma 2, we can formulate Theorem 2.

#### §4. Applications to Riemannian geometry

**4.1.** Let  $(M, g)$  be a pseudo-Riemannian manifold of dimensional  $n$ . Then from the present theorems 1 and 2 we conclude that an arbitrary Killing vector  $\omega$  has the following local covariant components  $\omega_i = e^{2\psi}(A_{ik}x^k + B_i)$  where  $\psi = [2(n+1)]^{-1} \ln|\det g|$ ,  $A$ 's and  $B$ 's are constants and  $A_{ik} + A_{ki} = 0$  (see also [St 1]). It follows that the group of infinitesimal isometric transformations has  $\frac{1}{2}n(n+1)$  parameters (see also [E], §71).

**4.2.** Following [T] and [Ka], a skew-symmetric covariant tensor field  $\vartheta$  of degree  $p$  ( $1 \leq p \leq n-1$ ) is called a conformal Killing tensor if  $\vartheta \in \ker D$  for

$$D = \nabla - \frac{1}{p+1}d - \frac{1}{n-p+1}g \wedge d^*$$

where  $d^*$  is the codifferential operator  $d^* : C^\infty \Lambda^{p+1} M \rightarrow C^\infty \Lambda^p M$  and

$$(g \wedge d^* \vartheta)_{i_0 i_1 \dots i_p} = \sum_{a=1}^p (-1)^{a+1} g_{i_0 i_a} (d^* \vartheta)_{i_1 \dots \hat{i}_a \dots i_p}.$$

Obviously, the set of conformal Killing tensors constitutes an vector space of tensor fields on  $(M, g)$ , denoted by  $\mathbf{C}^p(M, \mathbf{R})$  (see [St 5]). If a conformal Killing tensor  $\vartheta$  belongs to  $\ker d^*$ , then it is a Killing-Yano tensor. On the other hand, if a conformal Killing tensor  $\vartheta$  belongs to  $\ker d$ , it is called a closed conformal Killing tensor or a planar tensor (see [St 4] and [St 5]). We denote the vector space of these tensors by  $\mathbf{P}^p(M, \mathbf{R})$ .

By [Ka] on an arbitrary  $n$ -dimensional pseudo-Riemannian manifold  $(M, g)$  of constant nonzero sectional curvature  $C$  ( $C \neq 0$ ) the vector space  $\mathbf{C}^p(M, \mathbf{R})$  of conformal Killing tensors is decomposed uniquely in the form

$$\mathbf{C}^p(M, \mathbf{R}) = \mathbf{K}^p(M, \mathbf{R}) \oplus \mathbf{P}^p(M, \mathbf{R}). \quad (4.1)$$

From (4.1) we conclude that any conformal Killing tensor  $\vartheta$  of degree  $p$  is decomposed uniquely in the form  $\vartheta = \omega + \theta$  where  $\omega$  is a Killing-Yano tensor of degree  $p$  and  $\theta$  is a closed conformal Killing tensor of degree  $p$ .

Following theorem 1, on an  $n$ -dimensional pseudo-Riemannian manifold  $(M, g)$  of constant nonzero sectional curvature  $C$  ( $C \neq 0$ ) there is a local coordinate system  $x^1, \dots, x^n$  in which an arbitrary Killing-Yano tensor  $\omega$  of degree  $p$  ( $2 \leq p \leq n - 1$ ) has the components

$$\omega_{i_1 \dots i_p} = e^{(p+1)\psi} (A_{i_0 i_1 \dots i_p} x^{i_0} + B_{i_1 \dots i_p}) \quad (4.2)$$

where  $\psi = [2(n+1)]^{-1} \ln |\det g|$ ,  $\psi_k = \frac{\partial \psi}{\partial x^k}$  and  $A_{i_0 i_1 \dots i_p}$ ,  $B_{i_1 \dots i_p}$  are arbitrary skew-symmetric constants. On the other hand, by [St 3] on a pseudo-Riemannian manifold  $(M, g)$  of constant nonzero curvature  $C$  ( $C \neq 0$ ) the components  $\theta_{i_1 \dots i_p}$  of a closed conformal Killing tensor  $\theta$  of degree  $p$  ( $1 \leq p \leq n - 1$ ) can be found from the equations

$$\theta_{i_1 i_2 \dots i_p} = -\frac{1}{pC} \nabla_{i_1} \omega_{i_2 \dots i_p} \quad (4.3)$$

where  $\nabla_{i_1} \omega_{i_2 \dots i_p} = \partial_{i_1} \omega_{i_2 \dots i_p} - \omega_{k \dots i_p} \Gamma_{i_2 i_1}^k - \dots - \omega_{i_2 \dots k} \Gamma_{i_p i_1}^k$  is the expression for the covariant derivative  $\nabla \omega$  of the Killing-Yano tensor of degree  $p - 1$ . Moreover, by virtue of (3.1) on a pseudo-Riemannian manifold  $(M, g)$  of constant curvature  $C$  ( $C \neq 0$ ) the Christoffel symbols  $\Gamma_{ij}^k$  have the following form  $\Gamma_{ij}^k = \psi_i \delta_j^k + \psi_j \delta_i^k$  (see also [St 1]). Therefore, we can deduce from (4.2) and (4.3) that

$$\theta_{i_1 \dots i_p} = -\frac{1}{C} e^{p\psi} (\psi_{[i_1} A_{|k| i_2 \dots i_p]} x^k + \psi_{[i_1} B_{i_2 \dots i_p]} + \frac{1}{p} A_{i_1 i_2 \dots i_p}).$$

Consequently we have

**Theorem 3.** *On an  $n$ -dimensional pseudo-Riemannian manifold  $(M, g)$  of constant nonzero sectional curvature  $C$  ( $C \neq 0$ ) there is a local coordinate system  $x^1, \dots, x^n$  in which an arbitrary conformal Killing tensor  $\vartheta$  of degree  $p$  ( $2 \leq p \leq n - 1$ ) has the components*

$$\vartheta_{i_1 \dots i_p} = e^{(p+1)\psi} (A_{k i_1 \dots i_p} x^k + B_{i_1 \dots i_p}) - \frac{1}{C} e^{p\psi} (\psi_{[i_1} C_{|k| i_2 \dots i_p]} x^k + \psi_{[i_1} D_{i_2 \dots i_p]} + \frac{1}{p} C_{i_1 i_2 \dots i_p})$$

where  $\psi = [2(n+1)]^{-1} \ln |\det g|$ ,  $\psi_k = \frac{\partial \psi}{\partial x^k}$  and  $A_{i_0 i_1 \dots i_p}$ ,  $B_{i_1 \dots i_p}$ ,  $C_{i_1 \dots i_p}$  and  $D_{i_1 \dots i_p}$  are arbitrary skew-symmetric constants.

**Remark 3.** For a conformal Killing vector field, see K. Yano and T. Nagano [YN].

## References

- [C] Chern S.S. The geometry of G-structures. *Bull. Amer. Math. Soc.* **72**(1966), 167-219.
- [E] Eisenhart L.P. *Riemannian geometry*, Princeton Univ. Press, Princeton, NJ, 1949.
- [F] Fulton C.M. Parallel vector fields. *Proc. Amer. Math. Soc.* **16**(1965), 136-137.

- [KL] Katzin G.H., Levine J. Note on the number of linearly independent  $m^{\text{th}}$ -order first integrals space of constant curvature. *Tensor*, **19**(1968), no. 1, 42-44.
- [Ka] Kashiwada T. On konformal Killing tensor. *Natural Science Report, Ochanomizu University* **19**(1968), no. 2, 67-74.
- [Ko] Kobayashi Sh. *Transformation groups in differential geometry*. Erbeb. Math. Grenzgeb, Bd. 70, Springer-Verlag, New York-Heidelberg, 1972.
- [KN] Kobayashi and Nomizu K. *Foundations of differential geometry*. Vol. 1, Interscience, New York-London, 1963.
- [KSCH] Kramer D., Stephani H., Mac Callum M.A.H. and Herit E. *Exact solutions of Einstein's field equations*. Cambridge Univ. Press, Cambridge, 1980.
- [M] Mikes J. Geodesic mapping of affine-connected and Riemannian spaces. *J. Math. Sci.*, New York **78**(1996), no. 3, 311-333.
- [Ni] Nijenhuis A. A note on first integrals of geodesics. *Proc. Kon. Ned. Acad. Van. Wetens.*, Ser. A **52**(1967), 141-145.
- [No 1] Nomizu K. *What is affine differential geometry?* Differential Geometry Meeting Univ. Munster 1982. Tagungsbericht, 1982, 42-43.
- [No 2] Nomizu K. On completeness in affine differential geometry. *Geometriae Dedicata*, **20**(1986), no. 1, 43-49.
- [Sc] Schouten J.A. *Ricci-calculus*. Grundlehren Math. Wiss., Bd. **10**, Springer-Verlag, Berlin etc., 1954.
- [Sch] P.A. and A.P. Schirokow. *Affine Differentialgeometrie*. Teubner, Leipzig, 1962.
- [SSV] Simon U., Schwenk-Schellschmidt A., Viesel H. *Introduction to the affine differential geometry of hypersurfaces*. - Science University of Tokyo, Japan, 1991.
- [St 1] Stepanov S.E. The Killing-Yano tensor. *Theoretical and Mathematical Physics* **134**(2003), no. 3, 333-338.
- [St 2] Stepanov S.E. The Bochner technique for an  $m$ -dimensional compact manifold with  $SL(m, \mathbf{R})$ -structure. *St. Petersburg Mathematical Journal* **10**(1999), no. 4, 703-714.
- [St 3] Stepanov S.E. On conformal Killing 2-form of the electromagnetic field. *Journal of Geometry and Physics* **33**(2000), 191-209.
- [St 4] Stepanov S.E. A class of closed forms and special Maxwell's equations. *Tensor, N.S.* **58**(1997), 233-242.
- [St 5] Stepanov S.E. The vector space of conformal Killing forms on a Riemannian manifold.

*Journal of Mathematical Sciences.* **110**(2002), no. 4, 2892-2906.

[SS] Stepanov S.E., Smol'nikova M.V. Fundamental differential operators of orders one on exterior and symmetric forms. *Russ. Math. J.* **46**(2002), no.11, 51-56.

[ST] Stepanov S.E., Tsyganok I.I. *Vector fields in a manifolds with equiaffine connections.* Webs and Quasigroups, Tver Univ. Press, Tver, 1993, 70-77.

[T] Tachibana S.-I. On conformal Killing tensor in a Riemannian space. *Tohoku Math. Journal* **21**(199), 56-64.

[YI] Yano K., Ishihara Sh. Harmonic and relative affine mappings. *J. Differential Geometry* **10**(1975), 501-509.

[YN] Yano K., Nagano T. Some theorems on projective and conformal transformations. *Ind. Math.* **14**(1957), 451-458.