

Reconsideration of De Donder-Weyl theory by covariant analytic mechanics

Satoshi Nakajima*

Graduate School of Pure and Applied Sciences, University of Tsukuba, 1-1-1, Tennodai, Tsukuba, Japan 305-8571

(Dated: March 7, 2022)

We show that the covariant analytic mechanics (CAM) is closely related to the De Donder-Weyl (DW) theory. To treat space and time on an equal footing, the DW theory introduces D conjugate fields (D is the dimension of space-time) for each field and the CAM regards the differential forms as the basic variables. The generalization of the canonical equations is called the DW equations. Although one of the DW equations is not correct for the gauge field and the gravitational field, we show the way to improve it. By rewriting the canonical equations of the CAM, which are manifestly general coordinate covariant and gauge covariant, using the components of the tensors, we show that these are equivalent to the improved DW equations. Additionally, we investigate the Dirac field. We present a modified Hamilton formalism which regards only the Dirac fields as the basic variables and show that it provides the Dirac equations correctly.

PACS numbers:

I. INTRODUCTION

In the traditional analytic mechanics, the Hamilton formalism gives especial weight to time, then, the covariance is not trivial. To solve this problem, the De Donder-Weyl (DW) theory¹⁻³ introduced the conjugate fields $\pi_a^\mu = \partial\mathcal{L}/\partial\partial_\mu\psi^a$ ($\mu = 0, 1, \dots, D-1$) for each field ψ^a . Here, D is the dimension of space-time and \mathcal{L} is the Lagrangian density. The generalization of the canonical equations is called the DW equations. If a includes the tensor index, one of the DW equations is not correct generally.

In the traditional analytic mechanics and the DW theory, the basic variables are the components of the tensors. In contrast, the covariant analytic mechanics (CAM) regards the differential forms as the basic variables. Because the differential form is independent of the coordinate system, the general coordinate covariance is guaranteed manifestly. The Lagrange formalism of the CAM has been used⁴⁻⁷ from a few decades before. Nakamura⁸ generalized this formulation to the Hamilton formalism. In this method, the conjugate form is also a differential form, which treats space and time on an equal footing. Nakamura applied this method to the Proca field and the electromagnetic field with manifest covariance and, in the latter, with the gauge covariance. Kaminaga⁹ formulated strictly mathematically Nakamura's idea and constructed the general theory in arbitrary dimension. Kaminaga studied that the Newtonian mechanics of a harmonic oscillator and the scalar field, the electromagnetic field, the non-abelian gauge fields and 4 dimension gravity (second order formalism) without the Dirac field. On the other hand, Nester¹⁰ also independently investigated the CAM and constructed the general theory in 4 dimension and applied it to the Proca field, the electromagnetic field and the non-abelian gauge field. However, the treatment of the gravitational field (first order formalism) was not complete Hamilton formalism because the quantity corresponding to the Hamiltonian density was given by hand¹¹. We applied the CAM to the gravity with the Dirac field (both first and second order formalism)¹².

In this manuscript, we improve one of the DW equation and show that the improved DW theory is equivalent to the CAM for the first time. We apply the CAM to the Dirac field with the Lagrange multiplier. Moreover, we present a modified Hamilton formalism which regards only the Dirac fields as the basic variables and show that it provides the Dirac equations correctly.

In §2, we discuss the relation between the DW theory and the CAM. We introduce the DW theory in II A and the CAM in II B. In II C, we introduce improved DW equations and show that these are equivalent to the canonical equations of the CAM by rewriting the latter using the components of the tensors. And we discuss the difference between the Poisson bracket of the CAM and that of the DW theory^{2,3}. In §3, we apply the CAM to the Dirac field, which is a constraint system. We treat the Dirac field with the Lagrange multiplier. Moreover, using a modified Hamilton formalism regarding only the Dirac fields as the basic variables, we derive the Dirac equations correctly.

II. DE DONDER-WEYL THEORY AND COVARIANT ANALYTIC MECHANICS

A. De Donder-Weyl theory

Let us consider D dimension space-time. The Lagrangian density is a function of fields $\psi_{\mu_1 \dots \mu_p}^A$ and $\partial_\mu \psi_{\mu_1 \dots \mu_p}^A$. $\mathcal{L} = \mathcal{L}(\psi_{\mu_1 \dots \mu_p}^A, \partial_\mu \psi_{\mu_1 \dots \mu_p}^A)$. Here, A does not include tensor index and p ($= 0, 1, \dots$) can depend on A . The Euler-

Lagrange equation is given by

$$\frac{\partial \mathcal{L}}{\partial \psi_{\mu_1 \dots \mu_p}^A} - \partial_\mu \frac{\partial \mathcal{L}}{\partial \partial_\mu \psi_{\mu_1 \dots \mu_p}^A} = 0, \quad (1)$$

where $\mathcal{L} = \sqrt{-g}\mathcal{L}$ and g is the determinant of the metric $g_{\mu\nu}$. The conjugate fields of $\psi_{\mu_1 \dots \mu_p}^A$ are defined by

$$\pi_A^{\mu, \mu_1 \dots \mu_p} \stackrel{\text{def}}{=} \frac{\partial \mathcal{L}}{\partial \partial_\mu \psi_{\mu_1 \dots \mu_p}^A}, \quad (2)$$

and $\pi_A^{\mu, \mu_1 \dots \mu_p} \stackrel{\text{def}}{=} \pi_A^{\mu, \mu_1 \dots \mu_p} / \sqrt{-g}$. These are also called the generalized momenta or polymomenta. $\pi_A^{0, \mu_1 \dots \mu_p}$ is the traditional conjugate momentum. The DW Hamiltonian density is defined by

$$\mathcal{H}_{\text{DW}}(\psi_{\mu_1 \dots \mu_p}^A, \pi_A^{\mu, \mu_1 \dots \mu_p}) \stackrel{\text{def}}{=} \partial_\mu \psi_{\mu_1 \dots \mu_p}^A \pi_A^{\mu, \mu_1 \dots \mu_p} - \mathcal{L}, \quad (3)$$

and $\mathcal{H}_{\text{DW}} \stackrel{\text{def}}{=} \mathcal{H}_{\text{DW}} / \sqrt{-g}$. The variation of \mathcal{H}_{DW} is given by

$$\delta \mathcal{H}_{\text{DW}} = \partial_\mu \psi_{\mu_1 \dots \mu_p}^A \delta \pi_A^{\mu, \mu_1 \dots \mu_p} - \frac{\partial \mathcal{L}}{\partial \psi_{\mu_1 \dots \mu_p}^A} \delta \psi_{\mu_1 \dots \mu_p}^A. \quad (4)$$

If $\pi_A^{0, \mu_1 \dots \mu_p}, \dots, \pi_A^{D-1, \mu_1 \dots \mu_p}$ are independent with each other (this assumption is not correct generally) and these are independent of $\psi_{\mu_1 \dots \mu_p}^A$, we obtain identities

$$\frac{\partial \mathcal{H}_{\text{DW}}}{\partial \psi_{\mu_1 \dots \mu_p}^A} = -\frac{\partial \mathcal{L}}{\partial \psi_{\mu_1 \dots \mu_p}^A}, \quad \frac{\partial \mathcal{H}_{\text{DW}}}{\partial \pi_A^{\mu, \mu_1 \dots \mu_p}} = \partial_\mu \psi_{\mu_1 \dots \mu_p}^A. \quad (5)$$

Substituting (1) to the first equation, we obtain the DW equations:

$$\partial_\mu \psi_{\mu_1 \dots \mu_p}^A = \frac{\partial \mathcal{H}_{\text{DW}}}{\partial \pi_A^{\mu, \mu_1 \dots \mu_p}}, \quad (6)$$

$$\partial_\mu \pi_A^{\mu, \mu_1 \dots \mu_p} = -\frac{\partial \mathcal{H}_{\text{DW}}}{\partial \psi_{\mu_1 \dots \mu_p}^A}. \quad (7)$$

For instance, the electromagnetic field corresponds to $p = 1$. The Lagrangian density is given by $\mathcal{L}(A_\nu, \partial_\mu A_\nu) = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + A_\nu J^\nu$ with $F_{\mu\nu} = 2\partial_{[\mu}A_{\nu]} = \partial_\mu A_\nu - \partial_\nu A_\mu$. $[\]$ is the anti-symmetrization symbol. A_ν is the vector potential and J^ν is the current density, which is independent of A_μ . All indices are lowered and raised with $g_{\mu\nu}$ or its inverse $g^{\mu\nu}$. The conjugate fields of A_ν are $\pi^{\mu, \nu} = -F^{\mu\nu} = -\pi^{\nu, \mu}$. The DW Hamiltonian density is $\mathcal{H}_{\text{DW}}(A_\nu, \pi^{\mu, \nu}) = -\frac{1}{4}\pi_{\mu, \nu}\pi^{\mu, \nu} - A_\nu J^\nu$. We have $\partial \mathcal{H}_{\text{DW}} / \partial \pi^{\mu, \nu} = -\frac{1}{2}\pi_{\mu, \nu}$ and $\partial \mathcal{H}_{\text{DW}} / \partial A_\nu = -\sqrt{-g}J^\nu$. Then, the DW equations are given by

$$\partial_\mu A_\nu = -\frac{1}{2}\pi_{\mu, \nu}, \quad \partial_\mu \pi^{\mu, \nu} = \sqrt{-g}J^\nu. \quad (8)$$

The latter is equivalent to the Maxwell equation $\partial_\mu(\sqrt{-g}F^{\mu\nu}) = -\sqrt{-g}J^\nu$. However, the former is not correct. This is because $\pi^{\mu, \nu}$ are supposed as independent each other although there is a constraint condition $\pi^{\mu, \nu} = -\pi^{\nu, \mu}$.

B. Covariant analytic mechanics

Suppose a p -form β ($p = 0, 1, \dots, D$) is described by forms $\{\alpha^I\}_{I=1, \dots, k}$. If there exists the form ω_I such that β behaves under variations $\delta\alpha^I$ as

$$\delta\beta = \sum_I \delta\alpha^I \wedge \omega_I, \quad (9)$$

we call ω_I the *derivative* of β by α^I and denote

$$\frac{\partial \beta}{\partial \alpha^I} \stackrel{\text{def}}{=} \omega_I. \quad (10)$$

If α^I is $q_I(\leq p)$ -form, $\partial\beta/\partial\alpha^I$ is $(p - q_I)$ -form. Although the derivative of arbitrary form dose not always exist, as explained in IIC, the derivative of D -forms always exist. The Hodge operator $*$ maps an arbitrary p -form $\omega = \omega_{\mu_1 \dots \mu_p} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_p}$ ($p = 0, 1, \dots, D$) to a $D - p = r$ -form as

$$*\omega = \frac{1}{r!} E_{\nu_1 \dots \nu_r}^{\mu_1 \dots \mu_p} \omega_{\mu_1 \dots \mu_p} dx^{\nu_1} \wedge \dots \wedge dx^{\nu_r}. \quad (11)$$

Here, $\omega = \omega_{\mu_1 \dots \mu_p}$ and $E_{\mu_1 \dots \mu_D}$ are complete anti-symmetric tensors and $E_{01 \dots D-1} = \sqrt{-g}$. In particular, $\Omega = *1$ is the volume form. And $**\omega = -(-1)^{p(D-p)}\omega$ holds.

As the traditional analytic mechanics and the DW theory start from the Lagrangian density, the CAM starts from the *Lagrange D-form* L defined by $L \stackrel{\text{def}}{=} \mathcal{L}\Omega$. L is assumed to be described by forms ψ^A and $d\psi^A$: $L = L(\psi^A, d\psi^A)$. If ψ^A is a p -form ($p = 0, 1, \dots, D - 1$), ψ^A and $d\psi^A$ are expanded using a p -th order complete anti-symmetric tensor $\psi_{\mu_1 \dots \mu_p}^A$ as $\psi^A = \psi_{\mu_1 \dots \mu_p}^A dx^{\mu_1} \wedge \dots \wedge dx^{\mu_p}$ and $d\psi^A = \partial_{[\mu} \psi_{\mu_1 \dots \mu_p]}^A dx^\mu \wedge dx^{\mu_1} \wedge \dots \wedge dx^{\mu_p}$. Namely, the CAM treats a special class of the field theories of which Lagrangian density is described by the complete anti-symmetric tensors $\psi_{[\mu_1 \dots \mu_p]}^A$ and $\partial_{[\mu} \psi_{\mu_1 \dots \mu_p]}^A$. All fundamental field theories belong to this class. For instance, the Lagrange form of the electromagnetic field is given by

$$L(A, dA) = -\frac{1}{2} F \wedge *F + J \wedge A, \quad (12)$$

where $A = A_\mu dx^\mu$, $J = *(J_\mu dx^\mu)$ and $F = dA = \frac{1}{2} F_{\mu\nu} dx^\mu \wedge dx^\nu$.

The Euler-Lagrange equation is given by

$$\frac{\partial L}{\partial \psi^A} - (-1)^p d \frac{\partial L}{\partial d\psi^A} = 0. \quad (13)$$

The *conjugate form* π_A is defined by

$$\pi_A \stackrel{\text{def}}{=} \frac{\partial L}{\partial d\psi^A}. \quad (14)$$

π_A is $D - p - 1 = q$ -form. The *Hamilton D-form* (not $(D - 1)$ -form) is defined by

$$H(\psi^A, \pi_A) \stackrel{\text{def}}{=} d\psi^A \wedge \pi_A - L. \quad (15)$$

The variation of H is given by

$$\delta H = (-1)^{(p+1)q} \delta \pi_A \wedge d\psi^A - \delta \psi^A \wedge \frac{\partial L}{\partial \psi^A}. \quad (16)$$

If ψ^A and π_A are independent, we get identities

$$\frac{\partial H}{\partial \psi^A} = -\frac{\partial L}{\partial \psi^A}, \quad \frac{\partial H}{\partial \pi_A} = (-1)^{(p+1)q} d\psi^A. \quad (17)$$

Here, $(-1)^{(p+1)q} = 1$ if p is an odd number and $(-1)^{(p+1)q} = -(-1)^D$ if p is an even number. By substituting the Euler-Lagrange equation (13) into the first equation of (17), we obtain the *canonical equations*^{9,12}

$$d\psi^A = (-1)^{(p+1)q} \frac{\partial H}{\partial \pi_A}, \quad (18)$$

$$d\pi_A = -(-1)^p \frac{\partial H}{\partial \psi^A}. \quad (19)$$

The inverse transformation of (15) is given by

$$L'(\psi, \nu) \stackrel{\text{def}}{=} \pi_A \wedge \nu^A - H, \quad \nu^A \stackrel{\text{def}}{=} \frac{\partial H}{\partial \pi_A}. \quad (20)$$

From the second equation of (17), we have $\nu^A = (-1)^{(p+1)q} d\psi^A$. Then, $L'(\psi, \nu)$ reduces to the Lagrange form: $L'(\psi, \nu) = L(\psi, d\psi)$.

For the electromagnetic field, the Euler-Lagrange equation is given by $d * F = -(-1)^D J$. The conjugate form, $\pi = \partial L / \partial dA = - * F$, can represent dA as $dA = * \pi$. The Hamilton form is given by

$$H(A, \pi) = \frac{1}{2} \pi \wedge * \pi - J \wedge A. \quad (21)$$

We have $\partial H / \partial \pi = * \pi$ and $\partial H / \partial A = (-1)^D J$. The canonical equations $dA = \partial H / \partial \pi$ and $d\pi = \partial H / \partial A$ are

$$dA = * \pi, \quad d\pi = (-1)^D J. \quad (22)$$

The former is equivalent to the definition of the conjugate form and the latter coincides with the Euler-Lagrange equation.

We introduce two *Poisson brackets* by

$$\{\alpha, \beta\}_0 \stackrel{\text{def}}{=} (-1)^{(p+1)q} \frac{\partial \alpha}{\partial \psi^A} \wedge \frac{\partial \beta}{\partial \pi_A} - (-1)^p \frac{\partial \alpha}{\partial \pi_A} \wedge \frac{\partial \beta}{\partial \psi^A}, \quad (23)$$

$$\{\alpha, \beta\}_1 \stackrel{\text{def}}{=} (-1)^{(p+1)q} \frac{\partial \beta}{\partial \pi_A} \wedge \frac{\partial \alpha}{\partial \psi^A} - (-1)^p \frac{\partial \beta}{\partial \psi^A} \wedge \frac{\partial \alpha}{\partial \pi_A}, \quad (24)$$

for arbitrary forms α and β which have the derivatives by ψ^A and π_A . (23) was introduced in Ref.[12]. The canonical equations (18) and (19) can be written as

$$d\psi^A = \{\psi^A, H\}_i, \quad d\pi_A = \{\pi_A, H\}_i, \quad (25)$$

for $i = 0, 1$. And we have

$$\begin{aligned} \{\psi^A, \psi^B\}_i &= 0, \quad \{\pi_A, \pi_B\}_i = 0, \\ \{\psi^A, \pi_B\}_i &= (-1)^{(p+1)q} \delta_B^A, \quad \{\pi_B, \psi^A\}_i = -(-1)^p \delta_B^A, \end{aligned} \quad (26)$$

for $i = 0, 1$. Moreover, the Leibniz rules

$$\{\alpha, \beta \wedge \gamma\}_0 = \{\alpha, \beta\}_0 \wedge \gamma + (-1)^{b(D+1-a)} \beta \wedge \{\alpha, \gamma\}_0, \quad (27)$$

$$\{\alpha \wedge \beta, \gamma\}_1 = \{\alpha, \gamma\}_1 \wedge \beta + (-1)^{a(D+1-c)} \alpha \wedge \{\beta, \gamma\}_1 \quad (28)$$

hold for arbitrary a, b and c forms α, β and γ which have the derivatives by ψ^A and π_A . Using (28) and (25), the equation of motions of $\alpha \wedge \beta = \psi^A \wedge \psi^B, \psi^A \wedge \pi_B, \pi_A \wedge \pi_B$ are given by

$$d(\alpha \wedge \beta) = \{\alpha \wedge \beta, H\}_1. \quad (29)$$

In the derivation of (27) and (28), we used

$$\frac{\partial(\alpha \wedge \beta)}{\partial \psi^A} = \frac{\partial \alpha}{\partial \psi^A} \wedge \beta + (-1)^{ap} \alpha \wedge \frac{\partial \beta}{\partial \psi^A}, \quad \frac{\partial(\alpha \wedge \beta)}{\partial \pi_A} = \frac{\partial \alpha}{\partial \pi_A} \wedge \beta + (-1)^{aq} \alpha \wedge \frac{\partial \beta}{\partial \pi_A}. \quad (30)$$

If the derivative of α or β by $\psi^A(\pi_A)$ does not exist, the right hand side of the above first (second) equation is meaningless. For instance, if $\alpha \wedge \beta = \delta^{BC} \pi_B \wedge * \pi_C$, although $\partial(\alpha \wedge \beta) / \partial \pi_A = 2 * \pi_A$ holds, the above second equation does not hold since $\partial * \pi_B / \partial \pi_A$ does not exist (except for $D = 1$).

C. Relation between the two theories

In the CAM, the derivative of the fields is included as $\partial_{[\mu} \psi_{\mu_1 \dots \mu_p]}^A$. Then, the conjugate fields defined by (2) satisfy

$$\pi_A^{\mu, \mu_1 \dots \mu_p} = \pi_A^{[\mu, \mu_1 \dots \mu_p]}. \quad (31)$$

Considering this condition at (4), we obtain an improved equation of the second equation of (5) and of the DW equation (6):

$$\partial_{[\mu} \psi_{\mu_1 \dots \mu_p]}^A = \frac{\partial \mathcal{H}_{\text{DW}}}{\partial \pi_A^{\mu, \mu_1 \dots \mu_p}}. \quad (32)$$

For the electromagnetic field, this equation becomes $\partial_{[\mu} A_{\nu]} = -\frac{1}{2}\pi_{\mu,\nu}$. This is correct and corresponds to the first equation of (22). In the following, we show that (18) and (19) are respectively equivalent to (32) and (7).

We put $*\partial L/\partial\psi^A = m_{A\mu_1\cdots\mu_p} dx^{\mu_1} \wedge \cdots \wedge dx^{\mu_p}$. From $\delta\psi^A \wedge \partial L/\partial\psi^A = -*(*\partial L/\partial\psi^A) \wedge \delta\psi^A$, we have

$$\delta\psi^A \wedge \frac{\partial L}{\partial\psi^A} = -p! m_A^{\mu_1\cdots\mu_p} \delta\psi_{\mu_1\cdots\mu_p}^A \Omega. \quad (33)$$

On the other hand, δL is given by

$$\delta L = \left[\frac{\partial \mathcal{L}}{\partial\psi_{\mu_1\cdots\mu_p}^A} \delta\psi_{\mu_1\cdots\mu_p}^A + \pi_A^{\mu_1\mu_2\cdots\mu_p} \delta\partial_{[\mu} \psi_{\mu_1\cdots\mu_p]}^A \right] \frac{\Omega}{\sqrt{-g}}. \quad (34)$$

Then, we get $m_A^{\mu_1\cdots\mu_p} = -\frac{1}{p!} \frac{1}{\sqrt{-g}} \frac{\partial \mathcal{L}}{\partial\psi_{\mu_1\cdots\mu_p}^A}$. It leads

$$\frac{\partial L}{\partial\psi^A} = \frac{(-1)^{p(D-p)}}{p!} \frac{1}{\sqrt{-g}} \frac{\partial \mathcal{L}}{\partial\psi_{\mu_1\cdots\mu_p}^A} * dx_{\mu_1} \wedge \cdots \wedge dx_{\mu_p}, \quad (35)$$

with $dx_\mu = g_{\mu\nu} dx^\nu$. Similarly, we obtain

$$\begin{aligned} \pi_A &= \frac{(-1)^{(p+1)q}}{(p+1)!} \pi_A^{\mu_1\mu_2\cdots\mu_p} * dx_{\mu_1} \wedge dx_{\mu_2} \wedge \cdots \wedge dx_{\mu_p} \\ &= \frac{(-1)^{(p+1)q}}{(p+1)! q!} \pi_A^{\mu_1\mu_2\cdots\mu_p} \frac{1}{\sqrt{-g}} E_{\nu_1\cdots\nu_q, \mu_1\cdots\mu_p} dx^{\nu_1} \wedge \cdots \wedge dx^{\nu_q}. \end{aligned} \quad (36)$$

It leads

$$d\pi_A = \frac{(-1)^{(p+1)q}}{(p+1)! q!} \partial_\lambda \pi_A^{\mu_1\mu_2\cdots\mu_p} \frac{1}{\sqrt{-g}} E_{\nu_1\cdots\nu_q, \mu_1\cdots\mu_p} dx^\lambda \wedge dx^{\nu_1} \wedge \cdots \wedge dx^{\nu_q}. \quad (37)$$

From this equation, we obtain

$$*d\pi_A = -\frac{(-1)^p}{p!} \frac{1}{\sqrt{-g}} \partial_\mu \pi_A^{\mu_1\mu_2\cdots\mu_p} dx_{\mu_1} \wedge \cdots \wedge dx_{\mu_p}, \quad (38)$$

using a formula $E_{\nu_1\cdots\nu_a \mu_1\cdots\mu_b} E^{\nu_1\cdots\nu_a \rho_1\cdots\rho_b} = -a!b! \delta_{[\mu_1}^{\rho_1} \cdots \delta_{\mu_b]}^{\rho_b}$ ($a+b=D$). The above equation is equivalent to

$$d\pi_A = \frac{(-1)^{pq}}{p!} \frac{1}{\sqrt{-g}} \partial_\mu \pi_A^{\mu_1\mu_2\cdots\mu_p} * dx_{\mu_1} \wedge \cdots \wedge dx_{\mu_p}. \quad (39)$$

Because of (35) and (39), the Euler-Lagrange equation (13) is equivalent to (1). By the way, $d\psi^A \wedge \pi_A = \partial_{[\mu} \psi_{\mu_1\cdots\mu_p]}^A \pi_A^{\mu_1\mu_2\cdots\mu_p} \Omega = \partial_\mu \psi_{\mu_1\cdots\mu_p}^A \pi_A^{\mu_1\mu_2\cdots\mu_p} \Omega$ holds. It leads

$$H = \mathcal{H}_{\text{DW}} \Omega = \mathcal{H}_{\text{DW}} \frac{\Omega}{\sqrt{-g}}. \quad (40)$$

Using this equation, we get an equation similar to (35):

$$\frac{\partial H}{\partial\psi^A} = \frac{(-1)^{p(D-p)}}{p!} \frac{1}{\sqrt{-g}} \frac{\partial \mathcal{H}_{\text{DW}}}{\partial\psi_{\mu_1\cdots\mu_p}^A} * dx_{\mu_1} \wedge \cdots \wedge dx_{\mu_p}. \quad (41)$$

Because of the above equation and (39), the canonical equation (19) is equivalent to (7). If we put $\partial H/\partial\pi_A = l_{\mu\mu_1\cdots\mu_p}^A dx^\mu \wedge dx^{\mu_1} \wedge \cdots \wedge dx^{\mu_p}$, we have

$$\delta\pi_A \wedge \frac{\partial H}{\partial\pi_A} = (-1)^{(p+1)q} \delta\pi_A^{\mu_1\mu_2\cdots\mu_p} l_{\mu\mu_1\cdots\mu_p}^A \frac{\Omega}{\sqrt{-g}}. \quad (42)$$

It means $l_{\mu\mu_1\cdots\mu_p}^A = (-1)^{(p+1)q} \partial \mathcal{H}_{\text{DW}} / \partial \pi_A^{\mu_1\mu_2\cdots\mu_p}$. Then, we obtain

$$\frac{\partial H}{\partial\pi_A} = (-1)^{(p+1)q} \frac{\partial \mathcal{H}_{\text{DW}}}{\partial \pi_A^{\mu_1\mu_2\cdots\mu_p}} dx^\mu \wedge dx^{\mu_1} \wedge \cdots \wedge dx^{\mu_p}. \quad (43)$$

Because of this equation, the canonical equation (18) becomes (32).

In the DW theory, the Poisson-Gerstenhaber bracket (PGB), which is similar with (23) and (24), had been introduced using forms and multivector fields on the extended polymomentum phase space² of which coordinate is $(\psi^a, \pi_a^\mu, x^\mu)$. Here, a denotes both A and $\mu_1 \cdots \mu_p$. While our Poisson brackets are defined using the forms on space-times, the PGB is defined using the components of the tensors. Because the constraint (31) is not considered, the PGB does describe incorrect equation (6). Because of use of the the extended polymomentum phase space, the PGB has abundant structure and the scope of application of the PGB is wider than (23) and (24). The PGB satisfies only the graded Leibniz rule but also the graded anticommutativity and the graded Jacobi identity. While a generalization of the PGB to the Dirac bracket had been studied³, the generalization of our Poisson bracket is not known yet.

In the DW thoery, (6) is incorrect since the constraint (31) is not considered. However, decreasing independent variables using (31), we obtain correct equation (32). This fact suggests that one can obtain correct equation of motions from the Hamilton form by the variation using only independent variables determined by constraints among the original variables $\psi_{\mu_1 \cdots \mu_p}^A$ and $\pi_A^{[\mu, \mu_1 \cdots \mu_p]}$. We confirm this expectation for the Dirac field in the next section.

III. DIRAC FIELD

To obtain (17) from (16), we supposed that ψ^A and π_A are independent. However, if $\pi_A^{\mu, \mu_1 \cdots \mu_p}$ is a function of $\psi_{\mu_1 \cdots \mu_p}^A$ or identically zero, π_A is not an independent variable. The Dirac field and the first order formalism of the gravitational field correspond to this case. In this section, we study the Dirac field based on the CAM. First we use the method of Lagrange multiplier, which is the standard method for constraint systems. And next, we present a modified Hamilton formalism regarding only the Dirac fields as the basic variables to confirm the expectation mentioned in II C. In Ref.[13], the Dirac field had been studied based on the DW theory without using the Lagrange multiplier.

In the following of this paper, we set $D = 4$. Let $\{\theta^a\}_{a=0,1,2,3}$ denote an orthonormal frame. θ^a can be expanded as $\theta^a = \theta^a_\mu dx^\mu$ with the vielbein θ^a_μ . We have $g_{\mu\nu} = \eta_{ab} \theta^a_\mu \theta^b_\nu$ with $\eta_{ab} = \text{diag}(-+++)$. All indices are lowered and raised with η_{ab} or its inverse η^{ab} . Let ω^a_b be the connection 1-form. $\omega_{ba} = -\omega_{ab}$ holds. The torsion 2-form $\Theta^a = \frac{1}{2} C^a_{bc} \theta^b \wedge \theta^c$ satisfies $d\theta^a + \omega^a_b \wedge \theta^b = \Theta^a$.

The Lagrange form of the Dirac field $\psi = {}^t(\psi^1, \psi^2, \psi^3, \psi^4)$ is given by

$$L_D^\beta = -\frac{1+\beta}{2} \bar{\psi} \gamma_c e^c \wedge (d\psi + \frac{1}{4} \gamma_{ab} \omega^{ab} \psi) + \frac{1-\beta}{2} e^c \wedge (d\bar{\psi} - \frac{1}{4} \bar{\psi} \gamma_{ab} \omega^{ab}) \gamma_c \psi - m \bar{\psi} \psi \Omega - \frac{\beta}{2} C_a \bar{\psi} \gamma^a \psi \Omega, \quad (44)$$

with $\bar{\psi} \stackrel{\text{def}}{=} i\psi^\dagger \gamma^0 = (\bar{\psi}_1, \bar{\psi}_2, \bar{\psi}_3, \bar{\psi}_4)$, $e^a \stackrel{\text{def}}{=} * \theta^a$, $C_a \stackrel{\text{def}}{=} C^b_{ab}$ and $\gamma_{ab} \stackrel{\text{def}}{=} \gamma_{[a} \gamma_{b]}$. Here, γ^a is the gamma matrix, which satisfies $\gamma^a \gamma^b + \gamma^b \gamma^a = 2\eta^{ab}$. β is an arbitrary real number. m is the mass. L_D^β is given by $L_D^{\beta=0} + \frac{\beta}{2} d(e^a \bar{\psi} \gamma_a \psi)$ because of

$$de_a = -(\omega_a + C_a) \Omega, \quad \gamma_c e^c \wedge \frac{1}{4} \gamma_{ab} \omega^{ab} = \frac{1}{4} e^c \wedge \gamma_{ab} \omega^{ab} \gamma_c + \gamma^a \omega_a \Omega, \quad (45)$$

with $\omega_a \stackrel{\text{def}}{=} \omega^b_{ab}$. Because the Dirac field is a representation of the Lorentz transformation and a spinor field is defined in the tangent Minkowski space, the frame is necessary to write down the Lagrange form even in the flat space-time. The connection ω^a_b is the gauge field for the local Lorentz transformations. C_a is regarded as independent of ψ and $\bar{\psi}$. For simplicity, we treat the Dirac field as a usual number (not the Grassmann number). The Euler-Lagrange equations $\partial L_D^\beta / \partial \psi^A - d(\partial L_D^\beta / \partial d\psi^A) = 0$ and $\partial L_D^\beta / \partial \bar{\psi}_A - d(\partial L_D^\beta / \partial d\bar{\psi}_A) = 0$ ($A = 1, 2, 3, 4$) are respectively given by

$$(d\bar{\psi} - \frac{1}{4} \bar{\psi} \gamma_{ab} \omega^{ab}) \wedge \gamma_c e^c + m \bar{\psi} \Omega - \frac{1}{2} C_a \bar{\psi} \gamma^a \Omega = 0, \quad (46)$$

$$\gamma_c e^c \wedge (d + \frac{1}{4} \gamma_{ab} \omega^{ab}) \psi + m \psi \Omega + \frac{1}{2} C_a \gamma^a \psi \Omega = 0, \quad (47)$$

using (45)¹². (46) is the Hermitian conjugate of (47).

The conjugate forms of ψ^A and $\bar{\psi}_A$ are given by

$$\Pi_A^\beta = \frac{\partial L_D^\beta}{\partial d\psi^A} = \frac{1+\beta}{2} (\bar{\psi} \gamma_c)_A e^c, \quad \bar{\Pi}^{\beta A} = \frac{\partial L_D^\beta}{\partial d\bar{\psi}_A} = -\frac{1-\beta}{2} e^c (\gamma_c \psi)^A. \quad (48)$$

Corresponding conjugate fields $\pi_{(\beta)A}^\mu$ and $\bar{\pi}_{(\beta)}^{A\mu}$ are given by

$$\pi_{(\beta)A}^\mu = -\frac{1+\beta}{2}(\bar{\psi}\gamma_c)_A\theta^{c\mu}, \quad \bar{\pi}_{(\beta)}^{A\mu} = \frac{1-\beta}{2}\theta^{c\mu}(\gamma_c\psi)^A. \quad (49)$$

$\pi_{(\beta)}^0, \dots, \pi_{(\beta)}^3$ ($\bar{\pi}_{(\beta)}^0, \dots, \bar{\pi}_{(\beta)}^3$) are not independent each other and not independent from $\bar{\psi}(\psi)$. Then, we need to use the method of Lagrange multiplier. The Hamilton form is given by

$$H_{D,\text{tot}}^\beta = H_D^\beta + \left[\Pi_A^\beta - \frac{1+\beta}{2}(\bar{\psi}\gamma_c)_A e^c \right] \wedge \lambda^A + \bar{\lambda}_A \wedge \left[\bar{\Pi}^{\beta A} + \frac{1-\beta}{2}e^c(\gamma_c\psi)^A \right], \quad (50)$$

with

$$\begin{aligned} H_D^\beta &= d\psi^A \wedge \Pi_A^\beta + d\bar{\psi}_A \wedge \bar{\Pi}^{\beta A} - L_D^\beta \\ &= \frac{1+\beta}{2}\bar{\psi}\gamma_c e^c \wedge \frac{1}{4}\gamma_{ab}\omega^{ab}\psi + \frac{1-\beta}{2}e^c \wedge \frac{1}{4}\bar{\psi}\gamma_{ab}\omega^{ab}\gamma_c\psi + m\bar{\psi}\psi\Omega + \frac{\beta}{2}C_a\bar{\psi}\gamma^a\psi\Omega. \end{aligned} \quad (51)$$

Here, λ^A and $\bar{\lambda}_A$ are the Lagrange multiplier 1-forms. While the traditional Hamiltonian density includes $\partial_i\psi$, $\partial_i\bar{\psi}$ ($i = 1, 2, 3$), the Hamilton form does not include the exterior derivative of the Dirac fields. At $\beta = 1$, $\bar{\pi}_{(\beta)}^\mu$ becomes zero. In the traditional analytic mechanics at $\beta = 1$, the treatment which regards only ψ and $\pi_{(1)}^0 = -\bar{\psi}\gamma_c\theta^{c0}$ as the arguments of the Hamiltonian density goes well. However, in the CAM and the DW theory, one need to use the Lagrange multiplier even at $\beta = 1$. If we regard $\Pi_A^{\beta=1}$ as independent variable, $d\Pi_A^{\beta=1} = -\partial H_D^{\beta=1}/\partial\psi^A$ provides (46) correctly, however, $d\psi^A = -\partial H_D^{\beta=1}/\partial\Pi_A^{\beta=1}$ leads incorrect equation¹²

$$d\psi + \frac{1}{4}\gamma_{ab}\omega^{ab}\psi + \frac{m}{D}\gamma_a\theta^a\psi + \frac{1}{2}C_a\theta^a\psi = 0. \quad (52)$$

Applying $\gamma_b e^b$ to the above equation from the left and using $e^b \wedge \theta^a = \eta^{ab}\Omega$, we obtain (47). From (50), we obtain

$$\frac{\partial H_{D,\text{tot}}^\beta}{\partial\Pi^\beta} = \lambda, \quad \frac{\partial H_{D,\text{tot}}^\beta}{\partial\bar{\Pi}^\beta} = -\bar{\lambda}, \quad (53)$$

$$\begin{aligned} \frac{\partial H_{D,\text{tot}}^\beta}{\partial\psi} &= \frac{1+\beta}{2}\bar{\psi}\gamma_c e^c \wedge \frac{1}{4}\gamma_{ab}\omega^{ab} + \frac{1-\beta}{2}e^c \wedge \frac{1}{4}\bar{\psi}\gamma_{ab}\omega^{ab}\gamma_c + m\bar{\psi}\Omega \\ &\quad + \frac{\beta}{2}C_a\bar{\psi}\gamma^a\Omega + \bar{\lambda} \wedge \frac{1-\beta}{2}e^c\gamma_c, \end{aligned} \quad (54)$$

$$\begin{aligned} \frac{\partial H_{D,\text{tot}}^\beta}{\partial\bar{\psi}} &= \frac{1+\beta}{2}\gamma_c e^c \wedge \frac{1}{4}\gamma_{ab}\omega^{ab}\psi + \frac{1-\beta}{2}e^c \wedge \frac{1}{4}\gamma_{ab}\omega^{ab}\gamma_c\psi + m\psi\Omega \\ &\quad + \frac{\beta}{2}C_a\gamma^a\psi\Omega - \frac{1+\beta}{2}\gamma_c e^c \wedge \lambda. \end{aligned} \quad (55)$$

Then, the canonical equations $d\psi = -\partial H_{D,\text{tot}}^\beta/\partial\Pi^\beta$ and $d\bar{\psi} = -\partial H_{D,\text{tot}}^\beta/\partial\bar{\Pi}^\beta$ respectively become

$$d\psi = -\lambda, \quad d\bar{\psi} = \bar{\lambda}. \quad (56)$$

The canonical equations $d\Pi^\beta = -\partial H_{D,\text{tot}}^\beta/\partial\psi$ and $d\bar{\Pi}^\beta = -\partial H_{D,\text{tot}}^\beta/\partial\bar{\psi}$ respectively become

$$d\Pi^\beta = \frac{1}{4}\bar{\psi}\gamma_{ab}\omega^{ab} \wedge e^c\gamma_c - \frac{1+\beta}{2}\bar{\psi}\gamma^c\omega_c\Omega - m\bar{\psi}\Omega - \frac{\beta}{2}C_a\bar{\psi}\gamma^a\Omega - \frac{1-\beta}{2}d\bar{\psi} \wedge e^c\gamma_c, \quad (57)$$

$$d\bar{\Pi}^\beta = -\gamma_c e^c \wedge \frac{1}{4}\gamma_{ab}\omega^{ab}\psi + \frac{1-\beta}{2}\gamma^c\omega_c\psi\Omega - m\psi\Omega - \frac{\beta}{2}C_a\gamma^a\psi\Omega - \frac{1+\beta}{2}\gamma_c e^c \wedge d\psi, \quad (58)$$

using (54), (55), the second equation of (45) and (56). From (48) and the first equation of (45), the left hand sides of the above two equations are given by

$$d\Pi^\beta = \frac{1+\beta}{2}d\bar{\psi}\gamma_c \wedge e^c - \frac{1+\beta}{2}\bar{\psi}\gamma^c(\omega_c + C_c)\Omega, \quad (59)$$

$$d\bar{\Pi}^\beta = \frac{1-\beta}{2}(\omega_c + C_c)\gamma^c\psi\Omega + \frac{1-\beta}{2}e^c\gamma_c \wedge d\psi. \quad (60)$$

Substituting these into (57) and (58), we obtain (46) and (47) respectively.

We confirm the expectation mentioned at the last of IIC. We take only ψ^A and $\bar{\psi}_A$ as the independent variables. In contrast to the traditional analytic mechanics, the conjugate forms are the dependent variables. The variation of Π_A^β and $\bar{\Pi}^{\beta A}$ are respectively given by $\delta\bar{\psi}_B \frac{\partial \Pi_A^\beta}{\partial \psi_B}$ and $\frac{\partial \bar{\Pi}^{\beta A}}{\partial \bar{\psi}_B} \delta\psi^B$. Then, the variation of the Hamilton form $H_D^\beta(\psi, \bar{\psi}) = d\psi^A \wedge \Pi_A^\beta + d\bar{\psi}_A \wedge \bar{\Pi}^{\beta A} - L_D^\beta$ is given by

$$\delta H_D^\beta(\psi, \bar{\psi}) = \delta\psi^A \left[d\bar{\psi}_B \wedge \frac{\partial \bar{\Pi}^{\beta B}}{\partial \psi^A} - \frac{\partial L_D^\beta}{\partial \psi^A} \right] + \delta\bar{\psi}_A \left[-\frac{\partial \Pi_B^\beta}{\partial \bar{\psi}_A} \wedge d\psi^B - \frac{\partial L_D^\beta}{\partial \bar{\psi}_A} \right]. \quad (61)$$

From this equation, we obtain

$$d\Pi_A^\beta = -\frac{\partial H_D^\beta}{\partial \psi^A} + d\bar{\psi}_B \wedge \frac{\partial \bar{\Pi}^{\beta B}}{\partial \psi^A}, \quad (62)$$

$$d\bar{\Pi}^{\beta A} = -\frac{\partial H_D^\beta}{\partial \bar{\psi}_A} - \frac{\partial \Pi_B^\beta}{\partial \bar{\psi}_A} \wedge d\psi^B, \quad (63)$$

using the Euler-Lagrange equations. We call this method the modified Hamilton formalism. $\frac{\partial H_D^\beta}{\partial \psi^A}$ and $\frac{\partial H_D^\beta}{\partial \bar{\psi}_A}$ are respectively $\bar{\lambda} \rightarrow 0, \lambda \rightarrow 0$ limits of (54) and (55). The second terms of the right hand sides of (62) and (63) are given by

$$d\bar{\psi}_B \wedge \frac{\partial \bar{\Pi}^{\beta B}}{\partial \psi^A} = -\frac{1-\beta}{2} (d\bar{\psi} \wedge e^c \gamma_c)_A, \quad -\frac{\partial \Pi_B^\beta}{\partial \bar{\psi}_A} \wedge d\psi^B = -\frac{1+\beta}{2} (\gamma_c e^c \wedge d\psi)^A. \quad (64)$$

Then, the right hand sides of (62),(63) are given by

$$-\frac{\partial H_D^\beta}{\partial \psi^A} + d\bar{\psi}_B \wedge \frac{\partial \bar{\Pi}^{\beta B}}{\partial \psi^A} = -\frac{\partial H_{D,\text{tot}}^\beta}{\partial \psi^A} \Big|_{\bar{\lambda}=d\bar{\psi}}, \quad (65)$$

$$-\frac{\partial H_D^\beta}{\partial \bar{\psi}_A} - \frac{\partial \Pi_B^\beta}{\partial \bar{\psi}_A} \wedge d\psi^B = -\frac{\partial H_{D,\text{tot}}^\beta}{\partial \bar{\psi}_A} \Big|_{\lambda=-d\psi}. \quad (66)$$

Therefore, (62) and (63) are respectively equivalent to (57) and (58). For the Dirac field, we confirmed that one can obtain correct equation of motions from the Hamilton form by the variation using only independent variables.

IV. SUMMARY

We showed that the covariant analytic mechanics (CAM) is closely related to the De Donder-Weyl (DW) theory. Because the DW theory does not consider the constraint that the conjugate fields are complete anti-symmetric tensors, one of the DW equations is not correct generally. Decreasing independent variables using this constraint, we obtained an improved DW equation (IIC). By rewriting the canonical equations of the CAM using the components of the tensors, we showed that these are equivalent to the improved DW equations (IIC). So, we showed that the CAM is equivalent to the improved DW theory for the first time. While the Poisson bracket of the DW theory is defined using the components of the tensors and may describe the incorrect DW equation, our Poisson brackets are defined using the differential forms and do not lead to the incorrect DW equation. To generalize our Poisson bracket to the Dirac bracket and to apply it to the Dirac field are future works.

In III, we applied the CAM to the Dirac field, which is a constraint system. We treated the Dirac field with Lagrange multipliers. Getting a hint from the relation between the CAM and the DW theory, we presented the modified Hamilton formalism which regards only the Dirac fields as the basic variables and showed it provides the Dirac equations correctly.

Acknowledgments

We acknowledge helpful discussions with Y. Tokura, S. Tanimura and I. V. Kanatchikov.

* Electronic address: subarusatosi@gmail.com

- ¹ H. Weyl, “Observations on Hilbert’s Independence Theorem and Born’s Quantization of Field Equations”, *Phys. Rev.* **46**, 505 (1934).
- ² I. V. Kanatchikov, “Canonical structure of classical field theory in the polymomentum phase space”, *Rept. Math. Phys.* **41**, 49 (1998).
- ³ I. Kanatchikov, “On a generalization of the Dirac bracket in the De Donder-Weyl Hamiltonian formalism”, arXiv:0807.3127v1.
- ⁴ B. Kuchowicz, “Cosmology with spin and torsion. Part I. Physical and mathematical foundations”, *Acta Cosmologica, Zesz.* **3**, 109 (1975).
- ⁵ A. Trautman, “RECENT ADVANCES IN THE EINSTEIN-CARTAN THEORY OF GRAVITY”, *Annals of the New York Academy of Sciences* **262**, 241 (1975).
- ⁶ R. P. Wallner, “Notes on Recent U_4 Theories of Gravitation”, *General Relativity and Gravitation* **12**, 719 (1980).
- ⁷ F. W. Hehl, J. D. McCrea, E. W. Mielke and Y. Ne’eman, “Metric-affine gauge theory of gravity: field equations, Noether identities, world spinors, and breaking of dilation invariance”, *Phys. Rep.* **258**, 1 (1995).
- ⁸ T. Nakamura, *Bussei Kenkyu* **79**, 2 (2002) [in Japanese].
- ⁹ Y. Kaminaga, “Covariant Analytic Mechanics with Differential Forms and Its Application to Gravity”, *EJTP* **9**, 199 (2012).
- ¹⁰ J. M. Nester, “General pseudotensors and quasilocal quantities”, *Classical and Quantum Gravity* **21**, S261 (2004).
- ¹¹ J. M. Nester, “A covariant Hamiltonian for gravity theories”, *Mod. Phys. Lett. A* **06**, 2655 (1991).
- ¹² S. Nakajima, “Application of covariant analytic mechanics with differential forms to gravity with Dirac field”, *EJTP* **13**, 95 (2016).
- ¹³ G. Sardanashevily, “Generalized Hamiltonian Formalism for Field Theory: Constraint Systems” (World Scientific Pub. Co. Pte. Ltd., 1995).