

Quantum walk on spin network

M. M. Amaral^{a*}, R. Aschheim^{b†}, Klee Irwin^{b‡}

^a*Institute for Gravitation and the Cosmos,*
The Pennsylvania State University, University Park, PA 16802, USA
^b*Quantum Gravity Research*
Topanga, CA.

Abstract

We apply a discrete quantum walk from a quantum particle on a discrete quantum spacetime from loop quantum gravity and show that the related Entanglement Entropy can drive an entropic force. We apply these concepts to propose a model of a walker position topologically encoded on a spin network.

1 Introduction

One of the principal results from Loop Quantum Gravity (LQG) is a discrete spacetime, a network of loops implemented by spin networks [1] which is a digital/computational substrate of reality. So in order to better understand this substrate, it is natural to use tools from quantum information / quantum computation. Gravity, from a general perspective, has been studied with thermodynamic methods and in recent years numerous questions on black hole entropy and entanglement entropy have made these intense fields of research. On the side of quantum information and quantum computation there have been large improvements, including the development of many tools and we will herein consider here the framework of the coined Discrete time Quantum Walk (DQW). We will see that the problem of a quantum particle on a fixed spin network background from LQG can be worked out with the DQW. This gives rise to a new understanding of entanglement entropy and entropic force, permitting the proposal of a model for dynamics. From an ontological viewpoint we will see that dynamics and mass emerge from the spin network topology, and that quantum walk implements it.

In summary we will reinterpret some results from LQG trying to approach a quantum information perspective of a quantum geometric spacetime. One more step in direction of Wheeler's 'It from Bit', or 'It from Qubit'.

2 Methods

2.1 Particle interacting LQG

We start by considering a quantum particle on a quantum gravitational field from LQG [2]. In LQG, spin networks define quantum states of the gravitational field. To consider a quantum particle on this gravitational field we consider the state space built from tensor product of the gravity state \mathcal{H}_{LQG} and the particle state \mathcal{H}_P , $\mathcal{H} = \mathcal{H}_{LQG} \otimes \mathcal{H}_P$. The LQG state space can be spanned by a spin network basis $|s\rangle$ that is a spin network graph $\Gamma = (V(\Gamma), L(\Gamma))$, with $V(\Gamma) \rightarrow v_1, v_2, \dots$, vertices (nodes) colouring and $L(\Gamma) \rightarrow l_1, l_2, \dots, (\{\frac{1}{2}, 1, \frac{3}{2}, \dots\})$ links colouring. For the particle state space the relevant contribution comes from the position on the nodes of graph Γ , spanned by $|x_n\rangle$ where ($n = 1, 2, \dots$). The quantum state of the particle captures information from discrete geometry and cannot be considered independently from it. Therefore the Hilbert space is spanned by $|s, x_n\rangle$ and the Hamiltonian operator can be derived by fixing a spin network, and calculating the matrix element of this operator $\langle\psi|H|\psi\rangle$ with $\psi \in \mathcal{H}_P$. Accordingly, for the interaction between the particle and the fixed gravity state space [2] we have,

$$\langle\psi|H|\psi\rangle = \kappa \sum_l j_l(j_l + 1) (\psi(l_f) - \psi(l_i))^2, \quad (1)$$

*email: mramaciel@gmail.com

†email: raymond@quantumgravityresearch.org

‡email: klee@quantumgravityresearch.org

where κ is a constant that we can initially take as equal to one, l_f are the final points of the link l and l_i the initial points. So if the link l starts at node m and ends at node n we can change the notation, relabelling the color of this link l between m and n as j_{mn} and the wave function on the end points as $\psi(v_m), \psi(v_n)$. Now, for H , we have

$$\langle \psi | H | \psi \rangle = \kappa \sum_l j_{mn} (j_{mn} + 1) (\psi(v_n) - \psi(v_m))^2, \quad (2)$$

The operator H is positive semi-definite, and the ground state of H corresponds to the case where the particle is maximally delocalized. This leads to entropy [2]. In reference [3] it was considered that the classical random walk is associated with (2), a Markov chain. The transition probabilities for this random walk are

$$P_{mn} = \frac{j_{mn}(j_{mn} + 1)}{\sum_k j_{mk}(j_{mk} + 1)}. \quad (3)$$

and [3] shows that this reproduces an entropic force. This random walk is implemented with the Laplacian read off from (2) and differs from the Laplacians coming from the discrete calculus calculate in [4]. We will consider DQW related with those more general Laplacians in other work.

2.2 DQW

As we are studying the influence of a discrete geometry on a quantum field, it is natural to consider the quantum version of this random walk. To have a DQW we need an auxiliary subspace, the coin space, to define the unitary evolution

$$U = S(C \otimes I), \quad (4)$$

where S is a swap operation to change the position to a neighbour node and C is the coin operator related to this auxiliary subspace. On some graphs its clear how to define the coin Hilbert space [6] but for the spin network considered above its not clear exactly what auxiliary space to use. For a more general approach we can make use of Szegedy's DQW [7, 8], which we can work in two ways; 1- consider a bipartite walk; or 2 - a walk with memory. For a bipartite walk, if we consider a graph Γ we can simply make an operation of duplication to obtain a second graph $\bar{\Gamma}$, but for our purpose here, it's better to consider the second option: for the walk with memory, consider the N_v -dimensional Hilbert space $\mathcal{H}_n, \{|n\rangle, n = 1, 2, \dots, N_v\}$ and $\mathcal{H}_m, \{|m\rangle, m = 1, 2, \dots, N_v\}$, where N_v is the number of the nodes $V(\Gamma)$. The state of the walk is given in the product $\mathcal{H}_n^{N_v} \otimes \mathcal{H}_m^{N_v}$ spanned by these bases, by states at the previous $|m\rangle$ and current $|n\rangle$ steps, defined by

$$|\psi_n(t)\rangle = \sum_m^{N_v} \sqrt{P_{mn}} |n\rangle \otimes |m\rangle, \quad (5)$$

where P_{mn} is the transition probabilities that define a classical random walk, a Markov Chain, which is a discrete time stochastic process without a memory, with

$$\sum_n P_{mn} = 1. \quad (6)$$

Note that (6) is implied by definition of (3). For the evolution we can consider a simplified version of Szegedy's DQW [9], by defining a reflection, which we can interpret with the unitary "coin" operator¹,

$$C = 2 \sum_n |\psi_n\rangle \langle \psi_n| - I, \quad (7)$$

and a reflection with inverse action of the P swap (previous and current step) that can be implemented by a generalized swap operation

$$S = \sum_{n,m} |m, n\rangle \langle n, m|, \quad (8)$$

¹In literature we find usually that Szegedy's DQW don't have a "coin" operator.

and we have the unitary evolution

$$U = CS, \quad (9)$$

that defines the DQW.

So using Szegedy's approach is a straightforward way to obtain the discrete quantum walk on the spin network Γ considered above. It is given by equations (5 - 9) with P given by (3). Namely, for equation (5)

$$|\psi_n(t)\rangle = \sum_m^{N_v} \sqrt{\frac{j_{mn}(j_{mn} + 1)}{\sum_k j_{mk}(j_{mk} + 1)}} |n\rangle \otimes |m\rangle. \quad (10)$$

We can interpret that the coin space, the space of decisions encapsulating a non deterministic process and getting memory so that we have a unitary evolution (deterministic), makes the usual Entropy convert into Entanglement Entropy between steps in time. Szegedy's DQW is a general algorithm to quantize a Markov chain defined by transition probabilities $P_{n,m}$. These transition probabilities are obtained directly from the Hamiltonian of the system, so that Szegedy's DQW is used to simulate this system with quantum computation [10].

2.3 Entanglement Entropy and Entropic Force

We turn now to calculate Entanglement Entropy. Consider the Schmidt decomposition. Take a Hilbert space \mathcal{H} and decompose it into two subspaces \mathcal{H}_1 of dimension N_1 and \mathcal{H}_2 of dimension $N_2 \geq N_1$, so

$$\mathcal{H} = \mathcal{H}_1 \otimes \mathcal{H}_2. \quad (11)$$

Let $|\psi\rangle \in \mathcal{H}_1 \otimes \mathcal{H}_2$, and $\{|\psi_i^1\rangle\} \subset \mathcal{H}_1$, $\{|\psi_i^2\rangle\} \subset \mathcal{H}_2$, and positive real numbers $\{\lambda_i\}$, then the Schmidt decomposition read

$$|\psi\rangle = \sum_i^{N_1} \sqrt{\lambda_i} |\psi_i^1\rangle \otimes |\psi_i^2\rangle, \quad (12)$$

where λ_i are the Schmidt coefficients and the number of the terms in the sum is the Schmidt rank, which we label N . With this we can calculate the Entanglement Entropy between the two subspaces

$$S_E(C) = S_E(P) = -\sum_{i \in N} \lambda_i \log \lambda_i. \quad (13)$$

We can now calculate the local Entanglement Entropy between the previous step and the current n step (similar for current and next). Identify the Schmidt coefficients λ_i with P_{mn} given by (3) and from (10) we see that the Schmidt rank N is the valence of the node. Then insert in (12, 13), and the local Entanglement Entropy on current step is

$$S_{E_n} = -\sum_m^N P_{mn} \log P_{mn}. \quad (14)$$

By maximizing Entanglement Entropy

$$S_{E_n} = \log N_{max}, \quad (15)$$

where N_{max} is the largest valence. Which gives the entropic force for gravity worked out in [3, 5] with $\frac{dS}{dx} = |S_{E_n} - S_{E_m}|$ proportional to a small number identified with the particle mass M

$$\frac{dS}{dx} = |S_{E_n} - S_{E_m}| = \alpha M, \quad (16)$$

where, if we take the logarithm of (15) to be base 2, α is a constant of dimension $[bit/mass]$.

Here we have different interpretations and applications. With the DQW we have unitary evolution by encoding a non-deterministic part of the classical Markov Chain. This gives an internal structure for the particle, and Entanglement Entropy. In this digital physics substrate the particle is walking to maximizing Entanglement Entropy related to a memory of its walking path (a measurement of the system on n implies the state at previous m) and this generates an entropic force.

In fact we can consider von Neumann Entropy (14) in the context of quantum information. The probabilities are the Markov Chain connecting the steps: so the particles are constructing "words" while walking and this entropy measures the information needed. From which we can see the Entropic force emerge.

3 Results

3.1 Entropy of Black Hole

The framework above simulates a quantum field interacting with a fixed discrete geometry from quantum gravity which can be related to a gravitational entropic force. Consider now a regime from pure quantum gravity, like the black hole quantum horizon, so that there are no quantum fields, only quantum geometry. We can try to use this DQW to simulate this regime.

Note that DWQ implies a local Entanglement Entropy (14) in the sense that the chain/network is always under construction in this pixelated animated view. This resembles the isolated quantum horizons formulation from LQG [11, 12, 13, 14, 15] that gives an understanding of the origins of black hole entropy in LQG. In this scenario the Horizon and your area is emergent from the stepwise animated reality at the Planck scale, that in our case is simulated by DQW. We can propose that this DQW is the black hole quantum horizon, where the particle mass is the black hole mass in a random quantum walk on a fixed spin network.

In the isolated quantum horizons formulation the entropy is usually calculated by considering the eigenvalues of the area operator $A(j)$ and introducing an area interval $\delta a = [A(j) - \delta, A(j) + \delta]$ of the order of the Planck length, with relation to the classical area a of the horizon. $A(j)$ is given by

$$A(j) = 8\pi\gamma l_p^2 \sum_l \sqrt{j_l(j_l + 1)}, \quad (17)$$

where γ is the Barbero-Immirzi parameter and l_p the Planck length. The entropy, in adimensional form, is

$$S_{BH} = \ln N(A), \quad (18)$$

with $N(A)$ the number of microstates of quantum geometry on the horizon (area interval a) implemented, combinatorially, by considering states with links sequences that implement the two conditions

$$8\pi\gamma l_p^2 \sum_{l=1}^{N_a} \sqrt{j_l(j_l + 1)} \leq a, \quad (19)$$

related to the area, where N_a is the number of admissible j that puncture the horizon area, and

$$\sum_{l=1}^{N_a} m_l = 0 \quad (20)$$

related to the flux with m_l the magnetic quantum number satisfying the condition $-j_l \leq m_l \leq j_l$. To compare with Bekenstein-Hawking entropy only the first condition is important.

The detailed calculation [12, 15] shows that the dominant contribution to entropy comes from states in which there is a very large number of punctures. Thus, it is very attractive to interpret this entropy by quantum informational entropy (14). Let's investigate how the horizon area and related entropy can be emergent from maximal entanglement entropy of a DQW. Condition (19) is associated with this DQW. From (14 and 15), considering edge coloring, the maximal entanglement entropy occurs for states on nodes of large valence N_{max} and sequence with $j_l = \frac{l}{2}$ ($l = 1, 2, \dots, N_{max}$). So for the entropy calculation we can consider that admissible j that puncture the horizon area respect this sequences and the nodes associated have large valence. So we can rewrite condition (19) as

$$\sum_{i=1}^{N_{a_i}} a_i = a_c, \quad (21)$$

where N_{a_i} is the number of admissible nodes, with

$$a_c = \frac{a}{4\pi\gamma l_p^2}. \quad (22)$$

and each a_i is calculated from

$$a_i = \sqrt{l(l+2)}, \quad (23)$$

and considering the dominant contributions given by the over-estimate a_c and counting $N(a_c)$ that will give $N(A)$, each a_i is a integer strictly greater than 1

$$\sqrt{l(l+2)} = \sqrt{(l+1)^2 - 1} \approx l+1, \quad (24)$$

which means that the combinatorial problem we need to solve is to find $N(a_c)$ such that (21) holds, i.e. counting the ordered N_{a_i} -tuples of integers strictly greater than 1 summing to a_c . This was discussed in a similar problem in [11] and it is straightforward² to see that

$$\log N(A) = \frac{\log(\phi)}{\pi\gamma} \frac{a}{4l_p^2}, \quad (25)$$

where $\phi = \frac{1+\sqrt{5}}{2}$ is the golden ratio³.

We conjecture now that this entropy, relating to states that give maximal entanglement entropy, is just the entanglement entropy of the DQW. So the holographic quantum horizon can be simulated by the DQW with maximal internal entanglement entropy. The walker move is from node m to node n both with large valence so that equation (16) works. If we consider a walker mass spanning I with Planck mass m_p , we can choose the proportionality constant so that

$$\Delta S_E = |S_{E_n} - S_{E_m}| = I, \quad (26)$$

so $M = Im_p$ gives the amount of information on horizon, and ΔS_E is given in *bit*. We can make explicit the \log_2 on (25) so that it is given in *bit* too and propose that $\log_2 N(A) = \Delta S_E$, and that this measure of information gives the emergent Bekenstein-Hawking entropy. DQWs capture the black hole information, so to infer your entropy, someone needs to make contact with some frame of DQW, the particle that simulates black holes will be more probable on a node of maximal entanglement entropy, this encapsulates $\log N(A)$. Note that $\log N(A)$ is really not the usual statistical entropy because we have not taken into account all of the microstates, nor the flux condition. From (18) the black hole entropy, changing the logarithm, is

$$S_{BH} = \log_2 N(A) = \frac{\log_2(\phi)}{\pi\gamma} \frac{a}{4l_p^2}. \quad (27)$$

So Bekenstein-Hawking entropy is recovered by setting the Barbero-Immirzi parameter

$$\gamma = \frac{\log_2(\phi)}{\pi}, \quad (28)$$

showing that the states of maximal entanglement entropy are dominant to black hole entropy. So we can think of Bekenstein-Hawking entropy as emergent from the local Entanglement Entropy above. A horizon area a spanning $4I$ Planck areas has I bits like (26).

4 Discussion

4.1 A model of walker position topologically encoded on spin network


The Clebsch-Gordan condition at each node is implied by covering the graph with loops. From (3) and (14) we can compute the local entropy from a node as

$$S_{E_n} = \log \sigma - \frac{1}{\sigma} \sum_m^N j_{mn} (j_{mn} + 1) \log(j_{mn} (j_{mn} + 1)), \quad (29)$$

where $\sigma = \sum_m^N j_{mn} (j_{mn} + 1)$ of neighbour links. For example at a node $\{2, 3, 3\}$, $j = \{1, \frac{3}{2}, \frac{3}{2}\}$ gives $\sigma = \frac{19}{2}$ and $S_{E_n} = 1.06187$. At a node $\{2, 2, 2\}$, $j = \{1, 1, 1\}$ gives $\sigma = 6$, $S_{E_n} = 1.09861$ which is the maximal possible local entropy. Note (29) is the local entropy formula given in [3] as 14, and in accordance with well known LQG formulas for quantized length and quantized area.

²Because the cardinal $N(a_c)$ of the set of ordered tuples of integers strictly greater than 1 summing to a_c is the a_c^{th} Fibonacci number $F(a_c)$.

³The golden ratio is a irrational number (1.61803398875...) solution of the equation $\phi^2 = \phi + 1$.



The local entropy at each node is color coded. From equation (16) a massless particle move on same color and a massive particle moves along constant absolute color differences.

In the following figure :

(blue) $\{l_1, l_2, l_3\} = \{0, 1, 1\}$ or $\{0, 2, 2\}$ (side effect)

(white) ... $\{1, 2, 3\}$


(yellow) ... $\{1, 1, 2\}$

(orange) ... $\{2, 3, 3\}$


(red) ... $\{2, 2, 2\}$.

Dynamics:

Photons could orbit on orange hexagons. No possible particles with constant mass. Possible travel of massive particle, with $mass = |S_{(red)} - S_{(orange)}|$, ($= 0.037$) interacting with photons at some orange node.



The walker position, or the presence of a particle at one node is encoded by a triangle. Its move is a couple of 3-1 and 1-3 Pachner moves on neighbor positions, piloted by the walk probability



5 Conclusions

Results from Loop Quantum Gravity suggest the interesting idea that we can apply the results and tools from quantum information and quantum computation to a quantum spacetime. This is a field of research very promising. In this work we start a project to apply this tools like DQW to spacetime. We considered a DQW of a quantum particle on a quantum gravitational field and studied applications of related Entanglement Entropy. This memory time related Entanglement Entropy can drive an entropic force and give contribution to black hole entropy. With this we proposed a model for a walker position topologically encoded on spin network. This results in anomaly cancellation because the particle is no longer a point but has Planck scale size. We note that more complex models can be build, with more dimension and 8 quantum numbers in E8 model [16] and models with realistic emergent masses can be explored by computation.

A better understanding of Entanglement Entropy on black hole and application on cosmology are under investigation, as well as relating this model with QuasiCrystal E8 model [17]: the spin network can be chosen as the dual of a quasicrystal and the digital physics rules can be implemented by the quantum walk.

References

- [1] C. Rovelli; F. Vidotto, “Covariant Loop Quantum Gravity,” Cambridge University Press 1 edition, 2014.
- [2] C. Rovelli; F. Vidotto, “Single particle in quantum gravity and Braunstein-Ghosh-Severini entropy of a spin network,” *Phys. Rev. D* 2010, 81, arXiv:0905.2983v2 [gr-qc].
- [3] J. M. Garcia-Islas, “Entropic Motion in Loop Quantum Gravity,” ARXIV gr-qc/1411.4383.
- [4] G. Calcagni, D. Oriti and J. Thurigen, “Laplacians on discrete and quantum geometries,” *Class. Quantum Grav.* 30 (2013) 125006, arXiv:1208.0354 [hep-th].
- [5] E. Verlinde, “On the Origin of Gravity and the Laws of Newton,” *JHEP* 1104 (2011) 029, arXiv:1001.0785 [hep-th].
- [6] D. Aharonov, A. Ambainis, J. Kempe, U. Vazirani, “Quantum Walks On Graphs,” *Proceedings of ACM Symposium on Theory of Computation (STOC’01)*, July 2001, p. 50-59, arXiv:quant-ph/0012090.
- [7] M. Szegedy, “Quantum speed-up of Markov chain based algorithms,” *Proc. 45th IEEE Symposium on Foundations of Computer Science*, pp. 32-41, 2004, arXiv:quant-ph/0401053.
- [8] M. Szegedy, “Spectra of quantized walks and a rule”, arXiv:quant-ph/0401053v1.
- [9] B. Kollar, “Disorder and entropy rate in discrete time quantum walks,” PhD thesis, 2014, Pecs, Hungary.
- [10] A. Childs, “On the relationship between continuous and discrete time quantum walk,” *Commun.Math.Phys.*,294,581, 2010, arXiv:0810.0312 [quant-ph].
- [11] C. Rovelli, “Black hole entropy from loop quantum gravity,” *Phys. Rev. Lett.* **77**, 3288 (1996), arXiv:gr-qc/9603063.
- [12] A. Ashtekar, J. C. Baez and K. Krasnov, “Quantum geometry of isolated horizons and black hole entropy,” *Adv. Theor. Math. Phys.* **4**, 1 (2000), arXiv:gr-qc/0005126.
- [13] M. Domagala and J. Lewandowski, “Black hole entropy from quantum geometry,” *Class. Quant. Grav.* **21**, 5233 (2004), arXiv:gr-qc/0407051.
- [14] A. Ghosh and P. Mitra, “An Improved lower bound on black hole entropy in the quantum geometry approach,” *Phys. Lett. B* **616**, 114 (2005), arXiv:gr-qc/0411035.
- [15] J. F. Barbero G. and A. Perez, “Quantum Geometry and Black Holes”, arXiv:1501.02963 [gr-qc].
- [16] R. Aschheim, “SpinFoam with topologically encoded tetrad on trivalent spin networks,” Loops11 talk, slide 10, 24 may 2011, 19h10, Madrid, arXiv:1212.5473.
- [17] F. Fang; K. Irwin, “A Chiral Icosahedral QC and its Mapping to an E8 QC”, Aperiodic2015 poster, arXiv:1511.07786 [math.MG].