

Pressure Type Metrics on Spaces of Metric Graphs

Lien-Yung Kao*

Abstract

In this note, we consider two Riemannian metrics on a moduli space of metric graphs. Each of them could be thought of as an analogue of the Weil-Petersson metric on the moduli space of metric graphs. We discuss and compare geometric features of these two metrics with the “classic” Weil-Petersson metric in Teichmüller theory. This paper is motivated by Pollicott and Sharp’s work [PS14]. Moreover, we fix some errors in [PS14].

Contents

1	Introduction	1
2	Preliminaries	4
2.1	Symbolic dynamics and the thermodynamic formalism	4
2.2	The Weil-Petersson metric on moduli spaces	8
3	Pressure metrics on the space of metric graphs	9
3.1	From undirected graphs to directed graphs	9
3.2	Two pressure type metrics	10
4	Examples	12
4.1	Figure 8 graphs	13
4.2	Belt buckles	14
4.3	Dumbbells	18
4.4	Three-petal roses	21
5	Appendix	24

1 Introduction

This note is a further study of a dynamical-system-theoretically defined metric on deformation spaces– the pressure metric. The study of pressure metrics is ignited by McMullen’s study of the Weil-Petersson metric on Teichmüller space. In [Mcm08], McMullen proved that one can realize the Weil-Petersson metric on Teichmüller space by the pressure metric (on a certain functional space). The pressure metric has been a great tool for defining and studying the Weil-Petersson metrics on a variety of contexts: Teichmüller spaces [Mcm08], Anosov representations [BCLS13] and Blaschke products [Mcm08]. However, at this point, most of our understandings of the geometry of the pressure metric are coming from its relation with the “classic” Weil-Petersson metric. In this work, carrying over ideas from Pollicott–Sharp’s work [PS14], we focus on investigating the pressure metric geometry from a more dynamical approach.

More explicitly, we follow Pollicott–Sharp’s construction of pressure type metrics on the moduli space of metric graphs, and we consider two “natural”¹ pressure type metrics on the moduli space of graphs. We correct a formula in [PS14], and

*Department of Mathematics, University of Notre Dame, Notre Dame, IN 46545 USA. *E-mail*: lkao@nd.edu

¹ They are natural in the sense that the constructions of these two metrics are close to the construction of the “classic” Weil-Petersson metric on Teichmüller spaces introduced by McMullen.

using the revised formulas we examine geometric features of these two “natural” pressure type metrics on a moduli space of “typical”² graphs, and these examples show that geometric behaviors of these two “natural” pressure type metrics are very far from what we know about the “classic” Weil-Petersson metric (i.e. the pressure metric) on Teichmüller spaces.

To state our result more precisely and to put it in context, we first review basic setups in Teichmüller theory (cf. Section 2 for more details). Let S be a compact topological surface with negative Euler characteristics. Teichmüller space $\mathcal{T}(S)$ could be thought of as the set of isotopy classes of Riemannian metrics with constant curvature -1 , and the moduli space $\mathcal{M}(S)$ could be described as the set of isometry classes of Riemannian metrics with constant curvature -1 . Moreover, the moduli space $\mathcal{M}(S)$ is obtained by quotienting Teichmüller space $\mathcal{T}(S)$ by the mapping class group $\text{MCG}(S)$. The Weil-Petersson metric is a naturally defined and well-studied MCG-invariant metric on Teichmüller space (thus on the moduli space) with several striking features:

- the Weil-Petersson metric is negatively curved,
- the sectional curvature are neither bounded away from 0 nor $-\infty$, and
- the Weil-Petersson metric is incomplete.

McMullen’s result in [Mcm08] shows that on Teichmüller space we can define a Riemannian metric, *the pressure metric*, via the thermodynamic formalism, and which is exactly the Weil-Petersson metric. In other words, the pressure metric shares these notable geometric features with the Weil-Petersson metric on Teichmüller space.

Definition. Given a *undirected finite graph* \mathcal{G} with *edge set* \mathcal{E} . The *edge weighting function* $l : \mathcal{E} \rightarrow \mathbb{R}_{>0}$ assigns to each edge a length, which endows a metric structure onto \mathcal{G} . We call the pair (\mathcal{G}, l) a *metric graph*.

From a dynamical point of view, the metric graphs possess very similar structures as Riemann surfaces. Dynamics of paths on metric graphs is analogous to the dynamics of the geodesic flow for Riemann surfaces. It is because the length weighting function l on \mathcal{G} plays the same role as a Riemannian metric on surfaces. Hence, it is natural to begin the study of the pressure metric geometry from deformation spaces of metric graphs. Here, our deformation space corresponding to the graph \mathcal{G} is the space $\mathcal{M}_{\mathcal{G}}$ of all edge weighting functions.

Definition. For a graph \mathcal{G} and an edge weighting function l the *entropy* $h(l)$ is defined by

$$h(l) = \lim_{T \rightarrow \infty} \frac{1}{T} \log \#\{\gamma; l(\gamma) < T\}$$

where $\gamma = (e_0, e_1, \dots, e_n = e_0)$ is a closed cycle of edges in \mathcal{G} (without backtracking) and $l(\gamma) = \sum_{i=0}^{n-1} l(e_i)$.

From the dynamical perspective, the entropy $h(l)$ for metric graphs, as we have seen in surface cases, is an important and informative quality. We recall that the moduli space $\mathcal{M}(S)$ is the collection (up to isometry) of Riemannian metric on S with constant curvature -1 . We notice that the constant negative curvature condition of $\mathcal{M}(S)$ could be interpreted dynamically by using the constant (topological) entropy (of the geodesic flow on S) condition. More precisely, because when S has a constant negative curvature (say $K(S)$), the topological entropy of the geodesic flow for S is equal to $\sqrt{|K(S)|}$. Thus, to derive a close analogy to the moduli space $\mathcal{M}(S)$, it is natural and dynamical meaningful to consider the condition that entropy $h(l)$ equal to 1. Moreover, there is one more reason for us to concentrate on the space $\mathcal{M}_{\mathcal{G}}^1$; however, this reason is more technical and it’s from the nature of pressure type metrics. We will explain this reason in Remark 3.1. In what follows, we will focus on the deformation space $\mathcal{M}_{\mathcal{G}}^1$ of metric graphs with entropy 1. i.e.

$$\mathcal{M}_{\mathcal{G}}^1 = \{l : \mathcal{E} \rightarrow \mathbb{R}_{>0}; h(l) = 1\}.$$





² Graphs whose the fundamental group is a free group of rank 2.

Inspired by McMullen [Mcm08], Pollicott and Sharp constructed a pressure type metric (they call it a Weil-Petersson type metric) for metric graphs [PS14]. We follow their work and construct another pressure type metric for $\mathcal{M}_{\mathcal{G}}^1$. We call these two pressure type metrics the *pressure metric* and the *Weil-Petersson metric* for $\mathcal{M}_{\mathcal{G}}^1$, and denote them by $\|\cdot\|_P$ and $\|\cdot\|_{WP}$, respectively. The Weil-Petersson metric $\|\cdot\|_{WP}$ for $\mathcal{M}_{\mathcal{G}}^1$ is indeed conformal to the pressure metric $\|\cdot\|_P$ and the scaling function could be thought of a volume term of \mathcal{G} . Through normalizing the pressure metric $\|\cdot\|_P$ by the volume term of \mathcal{G} , the resulting metric $\|\cdot\|_{WP}$ is a closer analogy to McMullen's definition of the Weil-Petersson metric on Teichmüller space and the definition of the pressure metric on Hitchin components (cf. Theorem 2.6 and Section 3 for more details). Moreover, in Section 3.2, we propose several formulas and properties of pressure type metrics, which help to illustrate the usefulness of the definition.

Using formulas that we get in Section 3.1, we discuss four typical metric graphs: a figure 8 graph, a belt buckle, a dumbbell, and a three-petal rose. The first three graphs share the same fundamental group F_2 (the free group of rank 2). This makes an interesting connection with the *outer space* [CV86] in rank 2. For brevity, throughout this note, we denote the figure 8 graph, belt buckle, dumbbell, and the three-petal rose by \mathcal{G}_1 , \mathcal{G}_2 , \mathcal{G}_3 , and \mathcal{G}_4 , respectively. We summarize several results in Section 4 in the following.

Proposition (Proposition 4.1, 4.2, 4.5, 4.8, and Observation 4.1, 4.2, 4.3, 4.4). -

1. There exist examples of graphs for which the metric $\|\cdot\|_P$ is not complete.
2. There exist examples of graphs for which the curvature of the metric $\|\cdot\|_P$ is positive.
3. There exist examples of graphs for which the metric $\|\cdot\|_{WP}$ is complete.
4. There exist examples of graphs for which the curvature of the metric $\|\cdot\|_{WP}$ take positive and negative values.

			
Figure 8	Belt buckle	Dumbbell	Three-petal rose
$\ \cdot\ _P$	incomplete	incomplete $0 < C_2 \leq K \leq C_1$	incomplete $0 < K$, unbounded
$\ \cdot\ _{WP}$	complete	$C_2 \leq K \leq C_1 < 0$	K takes + and -

where K denotes the Gaussian curvature.

Remark 1.1. -

1. One need to be careful that our pressure metric $\|\cdot\|_P$ is the Weil-Petersson type metric defined in [PS14].
2. One should compare these results with results in Pollicott-Sharp's paper [PS14]. Although, Pollicott-Sharp and we are working on the same examples, the results are not quite match. It is because Lemma 3.3 in [PS14] (the main formula for calculating the pressure metric used in [PS14]) is true only if we input an extra condition that f is normalized with respect to the transfer operator (i.e. $\mathcal{L}_f 1 = 1$). For more details, one can check to our Remark 3.4.
3. The difference between **Propositions** and **Observations** is that for **Propositions** we give proofs and for **Observations** we give computer computation evidences. The reason that we skip proofs for **Observations** is because computations of the curvature with respect to $\|\cdot\|_{WP}$ and $\|\cdot\|_P$ are not hard but tedious and complex, which make no sense to analyze them in great detail rather than using the help from computers.

4. It is surprising that for all of our examples the sectional curvatures with respect to the pressure metric $\|\cdot\|_P$ are positive.

From the above proposition, we can see that both the pressure metric and the Weil-Petersson metric for metric graphs shares some common features with the Weil-Petersson metric on Teichmüller space. However, through working on these explicit examples, we can conclude that, from a pure dynamical setting, the pressure type metrics for metric graphs don't behave like the Weil-Petersson metric on Teichmüller space. In other words, the topology and the geometry of closed surfaces force the pressure metric to be the Weil-Petersson metric on Teichmüller space, but for graphs, without these special structures, the pressure type metrics cannot reflect these well-known features in the Weil-Petersson geometry.

This note is organized as the following. In the sake of completeness, we will review from basic knowledge of the symbolic dynamics, thermodynamic formalism and Teichmüller theory in Section 2. In Section 3, we relate the deformation space of metric graphs with a subshift of finite type and construct the pressure type metrics for metric graph via this relation. Furthermore, we give several useful formulas of them. In the last section, we discuss the pressure (type) metric geometry through working on explicit examples.

Acknowledgments

The author is extremely grateful to his Ph.D. advisor Prof. François Ledrappier and Prof. Mark Pollicott. They lead the author into the beautiful world of ergodic theory and dynamical systems, especially the thermodynamic formalism. This work would have never been possible without their support, guidance, and sharing of their insightful ideas.

2 Preliminaries

2.1 Symbolic dynamics and the thermodynamic formalism

We begin by recalling definitions and facts in symbolic dynamical systems and the thermodynamic formalism. An excellent reference on these two topics (concerning our approach) is the book [PP90] written by Parry and Pollicott.

A $k \times k$ matrix A is called *irreducible*, if for each pair (i, j) , $1 \leq i, j \leq k$, there exists $n \geq 1$ such that $A^n(i, j) > 0$, where A^n is an n -fold product of A with itself. Suppose A is an irreducible matrix consisting of 0 and 1, we define

$$\Sigma_A^+ = \{\underline{x} = (x_n)_{n \in \mathbb{N}} \in \{1, 2, \dots, k\}^{\mathbb{N}}; A(x_n, x_{n+1}) = 1\}.$$

We consider the shift map $\sigma : \Sigma_A^+ \rightarrow \Sigma_A^+$ by $(\sigma(\underline{x}))_n = x_{n+1}$, and then call (Σ_A^+, σ) a *(one-sided) subshift of finite type*. We notice that Σ_A^+ is a compact zero dimensional space with respect to the Tychonoff product topology, and we can endow a metric d on Σ_A^+ . More precisely, this space is compact with respect to the metric

$$d(\underline{x}, \underline{y}) = \sum_{n=0}^{\infty} \frac{1 - \delta(x_n, y_n)}{2^n},$$

where $\delta(i, j)$ is the standard Kronecker delta.

Remark 2.1. A is irreducible implies that (Σ_A^+, σ) is (topological) transitive (i.e. $\sigma : \Sigma_A^+ \rightarrow \Sigma_A^+$ has a dense orbit).

We denote the set of continuous functions on Σ_A^+ by $C(\Sigma_A^+)$. A function $f : \Sigma_A^+ \rightarrow \mathbb{R}$ is called α -Hölder continuous if there exists $C > 0$ and $\alpha \in (0, 1]$ such that for all $\underline{x}, \underline{y} \in \Sigma_A^+$ we have $|f(\underline{x}) - f(\underline{y})| \leq Cd(\underline{x}, \underline{y})^\alpha$, and we denote by $C^\alpha(\Sigma_A^+)$ the space of α -Hölder continuous functions on Σ_A^+ . We call a function $f : \Sigma_A^+ \rightarrow \mathbb{R}$ is Hölder continuous if it is α -Hölder continuous for some α . Two function $f, g \in C(\Sigma_A^+)$ are called *Livšic cohomologous* ($f \sim g$), if there exists $h \in C(\Sigma_A^+)$ such that

$$f - g = h \circ \sigma - h.$$

2.1.1 Transfer operator, pressure and variance

Most of the following definitions could be generalized to continuous functions $C(\Sigma_A^+)$ on Σ_A^+ . However, in this note we don't need such a generality, so we will only focus on Hölder continuous functions on Σ_A^+ .

Definition 2.1. The *transfer operator* (or *Ruelle operator*) $\mathcal{L}_f : C(\Sigma_A^+) \rightarrow \mathbb{R}$ of a Hölder continuous function f is defined by

$$(\mathcal{L}_f w)(\underline{x}) = \sum_{\underline{y}=\sigma^{-1}(\underline{x})} e^{f(\underline{y})} w(\underline{y}).$$

Theorem 2.1 (Ruelle-Perron-Frobenius Theorem, Theorem 2.2 [PP90]). *Let f be a Hölder continuous function on Σ_A^+ and suppose A is irreducible.*

1. *There is a simple maximal positive eigenvalue β_f of $\mathcal{L}_f : C(\Sigma_A^+) \rightarrow C(\Sigma_A^+)$ with a corresponding strictly positive eigenfunction $v_f \in C^\alpha(\Sigma_A^+)$, and β_f realizes the spectral radius of $\mathcal{L}_f : C(\Sigma_A^+) \rightarrow C(\Sigma_A^+)$.*
2. *There exists a unique probability measure μ_f such that $\mathcal{L}_f^* \mu_f = \beta_f \mu_f$. i.e.*

$$\int \mathcal{L}_f \phi d\mu_f = \beta_f \int \phi d\mu_f \text{ for all } \phi \in C(\Sigma_A^+).$$

By the Ruelle-Perron-Frobenius theorem, we are now ready to define the pressure.

Definition 2.2. For every $f \in C^\alpha(\Sigma_A^+)$, the *pressure of f* , $P(f)$, is the log of the spectral radius of \mathcal{L}_f . i.e.

$$P(f) = \log \beta_f,$$

where β_f is a simple maximal positive eigenvalue of \mathcal{L}_f given in Theorem 2.1 (the Ruelle-Perron-Frobenius theorem).

Remark 2.2. One can prove that we can characterize the pressure via periodic points of (Σ_A^+, σ) . More precisely, for every $f \in C^\alpha(\Sigma_A^+)$,

$$P(f) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \left(\sum_{\sigma^n \underline{x} = \underline{x}} e^{f^n(\underline{x})} \right),$$

where $f^n(\underline{x}) = f(\underline{x}) + f(\sigma \underline{x}) + \dots + f(\sigma^{n-1} \underline{x})$.

Definition 2.3. Let f be a Hölder continuous function, and suppose $P(f) = 0$. We define the *equilibrium state* of f to be the measure

$$m_f = \frac{v_f \cdot \mu_f}{\int v_f d\mu_f},$$

where μ_f is the unique probability measure associated with f given in Theorem 2.1 (the R-P-F theorem).

We notice that the equilibrium state is an ergodic, σ -invariant probability measure with positive entropy, and it is unique up to Livšic cohomology for Hölder continuous functions.

Definition 2.4. The *variance* of $w \in C(\Sigma_A^+)$ with respect to a σ -invariant probability measure μ is defined by

$$\text{Var}(w, \mu) = \lim_{n \rightarrow \infty} \frac{1}{n} \int \left(w^n(\underline{x}) - n \int w d\mu \right)^2 d\mu(\underline{x}),$$

where $w^n(\underline{x}) = \sum_{i=0}^{n-1} w \circ \sigma^i(\underline{x})$.

In this note, we only consider variances with respect to equilibrium states, which simplifies the discussion. In the sequel, we first recall some facts and useful formulas in calculating variances with respect to equilibrium states.

Theorem 2.2 (Lemma 3.4 [JP07]). *Suppose $f \in C^\alpha(\Sigma_A^+)$, $\mathcal{L}_f 1 = 1$ and $\int w dm_f = 0$, then*

$$\text{Var}(w, m_f) = \int_{\Sigma_A^+} w^2 dm_f,$$

where m_f is the equilibrium state with respect to f .

We also recall several useful properties of the pressure.

Theorem 2.3 (Analyticity of the Pressure, Prop. 4.7, 4.12 [PP90]). *If $f, g \in C^\alpha(\Sigma_A^+)$ and if m_f is the equilibrium state of f , then*

1. *The function $t \rightarrow P(f + tg)$ is analytic.*
2. *$\text{Var}(g, m_f) = 0$ if and only if g is Livšic cohomologous to a constant.*

By the analyticity of the pressure, we are able to differentiate the pressure, and more importantly derivatives of the pressure give us some handy formulas.

Theorem 2.4 (Derivatives of the pressure, Prop. 4.10, 4.11 [PP90], Theorem 2.2 [Mcm08]). *Let ψ_t be a smooth path in $C^\alpha(\Sigma_A^+)$, $m_0 = m_{\psi_0}$ be the equilibrium state of ψ_0 , and $\dot{\psi}_0 := \left. \frac{d\psi_t}{dt} \right|_{t=0}$. We then have*

$$\left. \frac{dP(\psi_t)}{dt} \right|_{t=0} = \int_{\Sigma_A^+} \dot{\psi}_0 dm_0 \quad (2.1)$$

and, if the first derivative is zero, i.e. $\int_{\Sigma_A^+} \dot{\psi}_0 dm_0 = 0$, then

$$\left. \frac{d^2 P(\psi_t)}{dt^2} \right|_{t=0} = \text{Var}(\dot{\psi}_0, m_0) + \int_{\Sigma_A^+} \ddot{\psi}_0 dm_0. \quad (2.2)$$

We now state an important property found by Bowen, so-called Bowen's formula, which relates the growth rate of weighted periodic orbits (or topological entropy of a certain system) with the pressure.

Theorem 2.5 (Bowen's formula, Prop. 6.1 [PP90]). *If $f : \Sigma_A^+ \rightarrow \mathbb{R}$ is a positive continuous function, then*

$$P(-s \cdot f) = 0$$

if and only if $s = h_f$ where

$$h_f = \lim_{T \rightarrow \infty} \frac{1}{T} \log \#\{\underline{x} \in \Sigma_A^+; \sigma^n \underline{x} = \underline{x} \text{ and } f^n(\underline{x}) < T \text{ for some } n \in \mathbb{N}\}.$$

Now we change gear and focus on a particular type of Hölder continuous functions: functions only depending on first two coordinates, i.e. $f : \Sigma_A^+ \rightarrow \mathbb{R}$ and $f(\underline{x}) = f(x_0, x_1)$, where $\underline{x} = x_0 x_1 x_2 \dots$. For every such function f , we have explicit formulas of the eigenvalue β_f , the eigenfunction v_f , and the measure μ_f given in Theorem 2.1.

Proposition 2.1 (Remark 1, p.27 [PP90]). *Let $f : \Sigma_A^+ \rightarrow \mathbb{R}$ be a function depending only on first two coordinates and $\mathcal{L}_f v_f = \beta_f v_f$. Suppose A_f is the matrix defined by*

$$A_f(i, j) = A(i, j) e^{f(i, j)}, \quad 1 \leq i, j \leq k,$$

then

1. β_f is a maximal eigenvalue of A_f .
2. v_f is the (left) eigenvector \mathbf{v}_f of A_f w.r.t. β_f , i.e.

$$\sum_i \mathbf{v}_f(i) A(i, j) e^{f(i, j)} = \beta_f \cdot \mathbf{v}_f(j).$$

3. If we define $g(i, j) = \log \mathbf{v}_f(i) - \log \mathbf{v}_f(j) - \log \beta_f + f(i, j)$, then $\mathcal{L}_g \mathbf{1} = 1$ and the matrix P_f corresponding to \mathcal{L}_g is

$$P_f(i, j) = A(i, j) e^{g(i, j)} = \frac{A(i, j) \mathbf{v}_f(i) e^{f(i, j)}}{\beta_f \cdot \mathbf{v}_f(j)}.$$

Moreover, P_f is column stochastic, i.e. $\sum_i A(i, j) e^{g(i, j)} = 1$, and the equilibrium state m_g w.r.t. g is given by

$$m_g[i_0, \dots, i_n] = P_f(i_0, i_1) \cdot \dots \cdot P_f(i_{n-1}, i_n) \mathbf{p}_f(i_n),$$

where $P_f \mathbf{p}_f = \mathbf{p}_f$ and $\sum_i \mathbf{p}_f(i) = 1$ and we use the notation $[i_0, i_1, \dots, i_n] = \{x \in \Sigma_A^+; x_j = i_j, j = 0, 1, \dots, n\}$.

Remark 2.3. We call $[i_0, i_1, \dots, i_n]$ a cylinder set, and in fact cylinder sets form a basis of the topology on Σ_A^+ .

2.1.2 Pressure type metric on subshifts of finite type

Here we keep the same setting as in the previous subsection that (Σ_A^+, σ) is a subshift of finite type and A is irreducible. We consider the space $\mathcal{P}(\Sigma_A^+)$ of Livšic cohomology classes of pressure zero Hölder continuous on Σ_A^+ , i.e.

$$\mathcal{P}(\Sigma_A^+) := \{f; f \in C^\alpha(\Sigma_A^+) \text{ for some } \alpha \text{ and } P(f) = 0\} / \sim,$$

where \sim denotes the Livšic cohomology relation.

The tangent space of $\mathcal{P}(\Sigma_A^+)$ at f is defined by

$$T_f \mathcal{P}(\Sigma_A^+) = \ker \mathbf{D}P(f) = \{\phi; \phi \in C^\alpha(\Sigma_A^+) \text{ for some } \alpha \text{ and } \int \phi dm_f = 0\} / \sim,$$

where $\mathbf{D}P(f)$ is derivative of P at f and the m_f is the equilibrium state of f .

Since the variance vanishes only on functions that are cohomologous to zero, by using the variance we can define two *pressure type metrics* on $T_f \mathcal{P}(\Sigma_A^+)$ as what stated below. The main reason why we call these metrics “pressure type metrics” is that these metrics are defined via the variance which is indeed the second derivative of the pressure (cf. Theorem 2.4).

Definition 2.5 (Pressure type metrics for subshifts of finite type). Let $f \in \mathcal{P}(\Sigma_A^+)$ and $\phi \in T_f \mathcal{P}(\Sigma_A^+)$ then we define two *pressure type metrics*:

$$\|\phi\|_P^2 := \text{Var}(\phi, m_f)$$

and

$$\|\phi\|_{WP}^2 := \frac{\text{Var}(\phi, m_f)}{-\int f dm_f}.$$

The former one is called the *pressure metric* on $\mathcal{P}(\Sigma_A^+)$ and denoted by $\|\cdot\|_P$, and the latter one is called the *Weil-Petersson metric* on $\mathcal{P}(\Sigma_A^+)$ and denoted by $\|\cdot\|_{WP}$.

One can immediately see from the definition that $\|\cdot\|_P$ and $\|\cdot\|_{WP}$ are conformal and they only differ by a normalization. We call $\|\cdot\|_{WP}$ the Weil-Petersson metric, because McMullen points out that after normalizing by the variance, one can get the Weil-Petersson metric of Teichmüller space (cf. the Theorem 2.6).

The following theorem is our main formula for computing the pressure metric and the Weil-Petersson metric.

Proposition 2.2. *If $\{\phi_t\}_{t \in (-1,1)}$ is a smooth one parameter family contained in $\mathcal{P}(\Sigma_A^+)$, then*

$$\left\| \dot{\phi}_0 \right\|_{WP}^2 = \frac{\int \ddot{\phi}_0 dm_{\phi_0}}{\int \phi_0 dm_{\phi_0}} \text{ and } \left\| \dot{\phi}_0 \right\|_P^2 = - \int \ddot{\phi}_0 dm_{\phi_0}$$

where $\dot{\phi}_0 = \frac{d}{dt} \phi_t \Big|_{t=0}$ and $\ddot{\phi}_0 = \frac{d^2}{dt^2} \phi_t \Big|_{t=0}$.

Proof. This follows the direct computation of the (Gâteaux) second derivative of $P(\phi_t)$:

$$\begin{aligned} \frac{d^2}{dt^2} P(\phi_t) \Big|_{t=0} &= (\mathbf{D}^2 P)(\phi_0)(\dot{\phi}_0, \dot{\phi}_0) + (\mathbf{D} P)(\phi_0)(\ddot{\phi}_0) \\ &= \text{Var}(\dot{\phi}_0, m_{\phi_0}) + \int \ddot{\phi}_0 dm_{\phi_0}. \end{aligned}$$

Since $P(\phi_t) = 0$, we have

$$\left\| \dot{\phi}_0 \right\|_{WP}^2 := \frac{\text{Var}(\dot{\phi}_0, m_{\phi_0})}{-\int \phi_0 dm_{\phi_0}} = \frac{\int \ddot{\phi}_0 dm_{\phi_0}}{\int \phi_0 dm_{\phi_0}}.$$

□

2.2 The Weil-Petersson metric on moduli spaces

In this subsection, we recall several facts of the Weil-Petersson metric on moduli spaces. These geometric features of the Weil-Petersson metric are critical clues for us to investigate the pressure metric geometry. The following approach to the Weil-Petersson metric is not traditional. The classic construction of the Weil-Petersson metric is built on the complex structure of Teichmüller space. Whereas, without employing the complex structure, McMullen in [Mcm08] gave a new characterization of the Weil-Petersson metric via the thermodynamic formalism. Our study in this work is inspired by this point of view. We summarize McMullen's theorem in Theorem 2.6. Before we jump into the statement, we recall some definitions.

Given a compact topological surface S with negative Euler characteristic, Teichmüller space $\mathcal{T}(S)$ describes the marked Riemannian metrics on S . i.e. the conformal classes of Riemannian metric on S with constant Gaussian curvature -1 . The moduli space $\mathcal{M}(S)$ describes the unmarked Riemannian metrics on S and is obtained by quotienting $\mathcal{T}(S)$ by the Mapping Class Group of S .

We consider a C^1 family of metric $g_\lambda \in \mathcal{M}(S)$, $0 \leq \lambda \leq 1$. Let $T^1 S$ be the unit tangent bundle of the surface S with respect to the metric g_{λ_0} . Let μ_{λ_0} be the corresponding Liouville measure on $T^1 S$. We denote by $\phi_t^{(\lambda_0)} : T^1 S \rightarrow T^1 S$ the geodesic flow. Since g_λ , for $0 \leq \lambda \leq 1$, is a volume preserving deformation we have

$$\int \dot{g}_{\lambda_0}(v, v) d\mu_{\lambda_0}(v) = 0,$$

where \dot{g}_{λ_0} is defined via the expansion

$$g_\lambda = g_{\lambda_0} + \dot{g}_{\lambda_0}(\lambda - \lambda_0) + O((\lambda - \lambda_0)^2).$$

(cf. Lemma 7. (a) and (c) [Pol94].)

Definition 2.6. The *variance* for $\dot{g}_{\lambda_0}(v, v)$ is given by

$$\text{Var}(\dot{g}_{\lambda_0}, \mu_{\lambda_0}) := \lim_{t \rightarrow \infty} \frac{1}{t} \int \left(\int_0^t \dot{g}_{\lambda_0}(\phi_s^{(\lambda_0)}(v), \phi_s^{(\lambda_0)}(v)) ds \right)^2 d\mu_{\lambda_0}$$

Theorem 2.6 (McMullen, Theorem 1.12 [Mcm08]). *The Weil-Petersson metric is proportional to the variance. More precisely,*

$$\|\dot{g}_{\lambda_0}\|_{Pressure}^2 := \frac{\text{Var}(\dot{g}_{\lambda_0}, \mu_{\lambda_0})}{\int_{T^1S} g_{\lambda_0}(v, v) d\mu_{\lambda_0}} = \frac{4}{3} \cdot \frac{\|\dot{g}_{\lambda_0}\|_{WP}^2}{\text{area}(S, g_{\lambda_0})}.$$

3 Pressure metrics on the space of metric graphs

In what follows, \mathcal{G} denotes a finite, connected, nontrivial (i.e. which contain at least two distinct closed path) and undirected graph with edge set \mathcal{E} . The length of each edge is given by the edge weighting function $l : \mathcal{E} \rightarrow \mathbb{R}_{>0}$.

Definition 3.1. Let $\mathcal{M}_{\mathcal{G}}$ denote the space of all edge weightings $l : \mathcal{E} \rightarrow \mathbb{R}_{>0}$ on \mathcal{G} .

Definition 3.2. The entropy $h(l)$ of the metric graph (\mathcal{G}, l) is defined by

$$h(l) = \lim_{T \rightarrow \infty} \frac{1}{T} \log \#\{\gamma; l(\gamma) < T\},$$

where $\gamma = (e_0, e_1, \dots, e_n = e_0)$ is a closed cycle of edges in \mathcal{G} (without backtracking) and $l(\gamma) = \sum_{i=0}^{n-1} l(e_i)$.

Definition 3.3. $\mathcal{M}_{\mathcal{G}}^1 = \{l : \mathcal{E} \rightarrow \mathbb{R}_{>0}; h(l) = 1\}$ is the space of all edge weightings with entropy $h(l) = 1$.

Remark 3.1. One might consider taking other normalizations on the space $\mathcal{M}_{\mathcal{G}}$, for example normalize $\mathcal{M}_{\mathcal{G}}$ by the volume of the graph, i.e. consider the moduli space $\mathcal{M}_{\mathcal{G}}^v := \{l : \mathcal{E} \rightarrow \mathbb{R}_{>0}; \text{Vol}(\mathcal{G}) = \sum l(e_i) = 1\}$. Notice that in the construction of pressure type metrics for subshifts of finite type we only consider functions of pressure zero. In general, for any positive Hölder functions f , there exists a positive number h_f such that $-h_f f$ is pressure zero (Bowen's formula cf. Theorem 2.5). Therefore, for each edge weighting function l , we can always scale l to be the pressure zero function $-h_l l$. So, for example the pressure type metrics on $\mathcal{M}_{\mathcal{G}}^v$ are the pullback the pressure type metrics on $\mathcal{M}_{\mathcal{G}}^1$ by the map $S : \mathcal{M}_{\mathcal{G}}^v \rightarrow \mathcal{M}_{\mathcal{G}}^1$ where $S(l) = -h_l l$. Hence, it is enough to study pressure type metrics on $\mathcal{M}_{\mathcal{G}}^1$.

3.1 From undirected graphs to directed graphs

In this subsection, we closely follow the construction of a symbolic model associating with metric graphs given in Pollicott and Sharp's work [PS14]. It is well-known that we can associate each directed graph with an adjacency matrix which records the directed edges connecting vertices to vertices.

To put directions on the undirected graph, we do the following. Given a undirected graph \mathcal{G} , for each edge $e \in \mathcal{E}$ we associate e with two directed edges which, abusing notation, we shall denote by e and \bar{e} , i.e. two opposite directions $\cdot \xleftrightarrow{\bar{e}} \cdot$. We denote by \mathcal{E}^0 the set of all directed edges. We say $e' \in \mathcal{E}^o$ follows $e \in \mathcal{E}^0$ if e' begins at the terminal endpoint of e , i.e. $\begin{matrix} \nearrow \\ e' \end{matrix} \cdot \begin{matrix} \searrow \\ e \end{matrix}$. We then define a $|\mathcal{E}^o| \times |\mathcal{E}^o|$ matrix A , with rows and columns indexed by \mathcal{E}^0 , by

$$A(e, e') = \begin{cases} 1 & \text{if } e' \text{ follows } e \text{ and } e' \neq \bar{e} \\ 0 & \text{otherwise.} \end{cases}$$

Then the shift space

$$\Sigma_A = \{\underline{e} = (e_n)_{n \in \mathbb{Z}} \in (\mathcal{E}^o)^{\mathbb{Z}}; A(e_n, e_{n+1}) = 1 \forall n \in \mathbb{Z}\}$$

can be naturally identified with the space of all two-sided infinite path (with a distinguished zeroth edge) in the graph \mathcal{G} . We call A the *adjacency matrix* of the undirected graph \mathcal{G} .

Remark 3.2. It is not hard to see A is irreducible. Recall that the graph \mathcal{G} is connected, so after we associate two (opposite) directions to each edge of \mathcal{G} , we know that for each pair of vertices of \mathcal{G} there exists a directed path connecting them.

Besides, Bowen showed that each two-sided subshift of finite type (Σ_A, σ) can be characterized by an one-sided subshift of finite type (Σ_A^+, σ) . More precisely, each Hölder function on Σ_A (two-sided sequences) is cohomologous to a Hölder function only depending on the future coordinates (one-sided sequences). Thus, it is enough to study the one-sided subshift of finite type (Σ_A^+, σ) , which is what we will do in the following.

The edge weighting function $l : \mathcal{E} \rightarrow \mathbb{R}_{>0}$, our main player, is clearly well-defined under the directed graph setting. Moreover, each edge weighting could be identified as a locally constant function on Σ_A^+ by, abusing the notation, $l(\underline{e}) := l(e_0)$ where $\underline{e} = e_0 e_1 \dots \in \Sigma_A^+$. Therefore, $\mathcal{M}_{\mathcal{G}}$ can be naturally regarded as a set of positive locally constant functions on Σ_A^+ . By definition, we can rewrite the entropy of the metric graph (\mathcal{G}, l) by

$$\begin{aligned} h(l) &:= \lim_{T \rightarrow \infty} \frac{1}{T} \log \# \{ \gamma; l(\gamma) < T \} \\ &= \lim_{T \rightarrow \infty} \frac{1}{T} \log \# \left\{ \underline{e} \in \Sigma_A^+; \sigma^n(\underline{e}) = \underline{e}, \text{ and } \sum_{i=0}^{n-1} l(e_i) < T \text{ for some } n \in \mathbb{N} \right\} \\ &= h_l. \end{aligned}$$

Hence, we have $\mathcal{M}_{\mathcal{G}}^1 = \{ l \in C(\Sigma_A^+); l > 0, l(\underline{e}) = l(e_0) \text{ and } h_l = 1 \}$.

Lemma 3.1 (Lemma 3.1 [PS14]). .

1. The pressure function is analytic on the space of locally continuous functions.
2. The entropy $h(l)$ of (\mathcal{G}, l) is characterized by $P(-h(l) \cdot l) = 0$.
3. The entropy function $\mathcal{M}_{\mathcal{G}} \ni l \mapsto h(l) \in \mathbb{R}_{>0}$ varies analytically for $l > 0$.

Proof. The first two assertions are coming from Theorem 2.3 and Theorem 2.5, respectively. The last one is a consequence of the implicit function theorem (for Banach spaces). \square

3.2 Two pressure type metrics

Following the discussion in Section 2.1.2, we can similarly define the pressure type metrics on $\mathcal{M}_{\mathcal{G}}^1$. Specifically, because $l \in \mathcal{M}_{\mathcal{G}}^1$, we know $h(l) = 1$ and $P(-h(l)l) = P(-l) = 0$. Hence for each $l \in \mathcal{M}_{\mathcal{G}}^1$ there exists an equilibrium state m_{-l} with respect to the function $-l$.

Definition 3.4. We define the tangent space to $\mathcal{M}_{\mathcal{G}}^1$ at l by

$$T_l \mathcal{M}_{\mathcal{G}}^1 = \left\{ \phi : \Sigma_A^+ \rightarrow \mathbb{R}; \phi(\underline{e}) = \phi(e_0) \text{ for all } e \in \mathcal{E}^0, \text{ and } \int \phi dm_{-l} = 0 \right\}.$$

Definition 3.5. For $\phi \in T_l \mathcal{M}_{\mathcal{G}}^1$ then we define two pressure type metrics:

$$\|\phi\|_P^2 := \text{Var}(\phi, m_{-l})$$

and

$$\|\phi\|_{WP}^2 := \frac{\text{Var}(\phi, m_{-l})}{\int l dm_{-l}}.$$

The former one is called the *pressure metric* on $\mathcal{M}_{\mathcal{G}}^1$ and denoted by $\|\cdot\|_P$, and the latter one is called the *Weil-Petersson metric* on $\mathcal{M}_{\mathcal{G}}^1$ and denoted by $\|\cdot\|_{WP}$. We can then define the length of every continuously differentiable

curve $\gamma : [0, 1] \rightarrow \mathcal{M}_G^1$ by

$$L_P(\gamma) = \int_0^1 \|\dot{\gamma}\|_P dt \text{ and } L_{WP}(\gamma) = \int_0^1 \|\dot{\gamma}\|_{WP} dt$$

and thus define two path space metrics on \mathcal{M}_G^1 by $d_{WP}(l_1, l_2) = \inf_{\gamma} \{L_{WP}(\gamma)\}$ and $d_P(l_1, l_2) = \inf_{\gamma} \{L_P(\gamma)\}$, where the infimum is taken over all continuously differentiable curves with $\gamma(0) = l_1$ and $\gamma(1) = l_2$.

Remark 3.3. -

1. One should notice that our Weil-Petersson metric for graphs is different from the Weil-Petersson metric for graphs mentioned in Pollicott and Sharp's paper [PS14]. In fact, their Weil-Petersson type metric is our pressure metric. Readers should be careful about this difference.
2. Another reason that we consider the new pressure type, $\|\cdot\|_{WP}$, is because it also coincides with the definition of the pressure metric on the Hitchin component introduced by Bridgeman, Canary, Labourie and Sambarino in [BCLS13].

Because l is a locally constant function, most of the above formulas of pressure type metrics, variances, and equilibrium states could be simplified quite a bit. The remaining of this subsection dedicates to expressing those quantities in simpler forms.

By Proposition 2.1, we consider the matrix A_{-l} associating to $-l$ defined by $A_{-l}(i, j) = A(i, j)e^{-l(i)}$. Notice that because $h(l) = 1$, A_{-l} has a simple maximum eigenvalue 1. Let \mathbf{v}_{-l} be the left eigenvector of A_{-l} w.r.t. 1, i.e. $\mathbf{v}_{-l} \cdot A_{-l} = \mathbf{v}_{-l}$. Then we construct the column stochastic matrix P_{-l} associating with $-l$:

$$P_{-l}(i, j) = \frac{A(i, j)\mathbf{v}_{-l}(i)e^{f(i)}}{\mathbf{v}_{-l}(j)}, \quad (3.1)$$

and let \mathbf{p}_{-l} be the unit right eigenvector of P_{-l} w.r.t. 1, i.e. $P_{-l}\mathbf{p}_{-l} = \mathbf{p}_{-l}$ and $\sum_i \mathbf{p}_{-l}(i) = 1$. Thus we know that the equilibrium state m_{-l} is given by

$$m_{-l}[i_0, i_1, \dots, i_n] = P_{-l}(i_0, i_1) \cdot \dots \cdot P_{-l}(i_{n-1}, i_n) \mathbf{p}_{-l}(i_n).$$

When there is no ambiguity, for short, we will drop the subscript l from \mathbf{v}_{-l}, P_{-l} and \mathbf{p}_{-l} and denote them by \mathbf{v} , P and \mathbf{p} , respectively.

Proposition 3.1. *We have*

1. For each $l \in \mathcal{M}_G^1$, the tangent space to \mathcal{M}_G^1 at l is

$$T_l \mathcal{M}_G^1 = \left\{ \phi \in C(\Sigma_A^+); \phi(\underline{e}) = \phi(e_0) \text{ and } \sum_{e \in \mathcal{E}^o} \phi(e) \mathbf{p}_e = 0 \right\},$$

where $\mathbf{p}_e = \mathbf{p}(e)$.

2. Let l_t be a smooth path in \mathcal{M}_G^1 , and g_t be the (up to cohomology) normalized function cohomologous to $-l_t$, i.e. $\mathcal{L}_{g_t} \mathbf{1} = 1$ and $-l_t = g_t + h_t \circ \sigma - h_t$, and let $P(i, j) = P_{-l}(i, j) = A(i, j)e^{g_0(i, j)}$, and \mathbf{p} be the right eigenvector of P w.r.t. 1 (i.e. $P\mathbf{p} = \mathbf{p}$) then

$$\text{Var}(-\dot{l}_0, m_{-l_0}) = \int (\ddot{l}_0) dm_{-l_0} = \sum_i \ddot{l}_0(i) \mathbf{p}_i \quad (3.2)$$

$$= \int (\dot{g}_0)^2 dm_{g_0} = \sum_{i, j} (\dot{g}_0(i, j))^2 P_{ij} \mathbf{p}_j. \quad (3.3)$$

Proof. (1): It is obvious.

(2): By Proposition 2.1, we know that $m_{-l_0}[e_i] = \mathbf{p}_{e_i}$ for each i and $m_{-l_0}[e_i, e_j] = P(e_i, e_j)\mathbf{p}(e_j)$ for all $e_i, e_j \in \mathcal{E}^0$.

For equation (3.2), by Theorem 2.4, we know

$$\text{Var}(-\dot{l}_0, m_{-l_0}) = \int (\ddot{l}_0) dm_{-l_0} = \sum_i \int_{[i]} (\ddot{l}_0) dm_{-l_0} = \sum_i \ddot{l}_0(i) \mathbf{p}_i.$$

For equation (3.3), since $P(-h(l)l) = 0$ and by Theorem 2.4, we have $\int \dot{l}_0 dm_{-l_0} = 0$. Moreover, because for each Hölder continuous function the equilibrium state is unique in each Livšic cohomology class we know that $m_{-l_0} = m_{g_0}$. Since $-\dot{l}_0 \sim \dot{g}_0$ and $-\ddot{l}_0 \sim \ddot{g}_0$, we have $\int \dot{l}_0 dm_{-l_0} = \int \dot{g}_0 dm_{g_0} = 0$ and $\int -\ddot{l}_0 dm_{-l_0} = \int \ddot{g}_0 dm_{g_0}$. Because $\mathcal{L}_{g_0} 1 = 1$ and $\int \dot{g}_0 dm_{g_0} = 0$, by Theorem 2.2 we have

$$\text{Var}(-\dot{l}_0, m_{-l_0}) = \text{Var}(\dot{g}_0, m_{g_0}) = \int (\dot{g}_0)^2 dm_{g_0} = \sum_{i,j} \int_{[i,j]} (\dot{g}_0)^2 dm_{g_0} = \sum_{i,j} (\dot{g}_0(i,j))^2 P_{ij} \mathbf{p}_j.$$

□

Remark 3.4. Equation (3.3) is very close to the formula given in Lemma 3.3 [PS14]. However, Lemma 3.3 in [PS14] is NOT true for all functions in the tangent space $T_l \mathcal{M}_{\mathcal{G}}^1$. It is only true for normalized functions, i.e. $f \in T_l \mathcal{M}_{\mathcal{G}}^1$ and $\mathcal{L}_f 1 = 1$.

Notice that it is convenient to interpret $\mathcal{M}_{\mathcal{G}}^1$ as a hypersurface in $\mathcal{M}_{\mathcal{G}}$. Because each edge weighting function l is depending on the weighting (i.e. length) of each edge, it can be regarded as a k -variable function $l = l(e_1, e_2, \dots, e_k)$ where e_1, e_2, \dots, e_k are edges of \mathcal{G} . Thus, in this perspective $\mathcal{M}_{\mathcal{G}}$ is $\mathbb{R}_{>0}^k$. Moreover, the condition $h(l) = 1$ gives us an equation of e_1, e_2, \dots , and e_k , so $\mathcal{M}_{\mathcal{G}}^1$ is a co-dimension one submanifold in $\mathcal{M}_{\mathcal{G}}$.

4 Examples

For examples in this section, we follow the recipe below to calculate the pressure type metrics. Let \mathcal{G} be a undirected finite graph and l be an edge weighting function then


- First, associate two opposite directions to each edge as we defined in Section 3.1, then write down the adjacency matrix A and the wighted adjacency matrix $A_{-h(l)l}$ associated with l .
- Second, solve the equation $h(l) = 1$. Explicitly, since $h(l) = 1$ means that 1 is the an eigenvalue of the matrix A_{-l} , the characteristic polynomial of A_{-l} is a function of edges e_1, \dots, e_k of \mathcal{G} , i.e. $\det(A_{-l} - \text{Id}) = 0$. By solving the characteristic polynomial, we can write $l = (l(e_1), l(e_2), \dots, l(e_k))$ where $l(e_k) = w(l(e_1), \dots, l(e_{k-1}))$ is an analytic function depending $l(e_1), \dots, l(e_{k-1})$.
- Third, compute the right eigenvector \mathbf{v}_{-l} of A_{-l} w.r.t the eigenvalue 1.
- Fourth, write down the normalized weighted matrix P_{-l} in (3.1), and compute the equilibrium state \mathbf{p} of $-l$, i.e. the unit left eigenvector \mathbf{p} of P_{-l} w.r.t. 1.
- Fifth, consider $l = (l(e_1), l(e_2), \dots, w(l(e_1), l(e_2), \dots, l(e_k)))$ as a parametrization of $\mathcal{M}_{\mathcal{G}}^1$, and then compute the tangent vectors $\frac{\partial}{\partial l(e_i)}$ for $1 \leq i \leq k$.
- Last, compute the pressure metric and the Weil-Petersson metric of $\frac{\partial}{\partial l(e_i)}$ for $1 \leq i \leq k$.

We notice that graphs of a belt buckle \mathcal{G}_2 , a dumbbell \mathcal{G}_3 and a three-petal rose \mathcal{G}_3 are graphs with 3 edges. Therefore the moduli spaces $\mathcal{M}_{\mathcal{G}_2}^1$, $\mathcal{M}_{\mathcal{G}_3}^1$ and $\mathcal{M}_{\mathcal{G}_3}^1$ are two-dimensional manifolds sitting in \mathbb{R}^3 . Moreover, in the Riemannian geometry setting, it is natural to study the important geometric invariants—curvatures. In dimension two, it is the Gaussian curvature.

Before we start working on examples, we recall two useful asymptotic notations that $A(x) \sim B(x)$ means $\lim \frac{A(x)}{B(x)} = 1$ and $A(x) \asymp B(x)$ means there exist positive constants c and c' such that $c < \frac{A(x)}{B(x)} < c'$.

4.1 Figure 8 graphs

The first example is a figure 8 graph which we denote by \mathcal{G}_1 . The picture below is a figure 8. In the sake of brevity, we denote $l(e_1)$ and $l(e_2)$ by x and y , respectively.



Following the recipe, we have

- $A = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \end{pmatrix}$ and $A_{-l} = \begin{pmatrix} e^{-x} & e^{-x} & 0 & e^{-x} \\ e^{-y} & e^{-y} & e^{-y} & 0 \\ 0 & e^{-x} & e^{-x} & e^{-x} \\ e^{-y} & 0 & e^{-y} & e^{-y} \end{pmatrix}$.
- $h(l) = 1 \implies \det(\text{Id} - A_{-l}) = 0 \implies e^{-y} = \frac{1-e^{-x}}{1+3e^{-x}}$.
- $\mathbf{v} = \left(\frac{2}{1+3e^{-x}}, 1, \frac{2}{1+3e^{-x}}, 1 \right)$.
- $P = \begin{pmatrix} e^{-x} & \frac{2e^{-x}}{1+3e^{-x}} & 0 & \frac{2e^{-x}}{1+3e^{-x}} \\ \frac{1-e^{-x}}{2} & \frac{1-e^{-x}}{1+3e^{-x}} & \frac{1-e^{-x}}{2} & 0 \\ 0 & \frac{2e^{-x}}{1+3e^{-x}} & e^{-x} & \frac{2e^{-x}}{1+3e^{-x}} \\ \frac{1-e^{-x}}{2} & 0 & \frac{1-e^{-x}}{2} & \frac{1-e^{-x}}{1+3e^{-x}} \end{pmatrix}$ and $\mathbf{p} = \begin{pmatrix} \mathbf{p}_1(x) \\ \mathbf{p}_2(x) \\ \mathbf{p}_3(x) \\ \mathbf{p}_4(x) \end{pmatrix} = \begin{pmatrix} \frac{2e^x}{6e^x + e^{2x} - 3} \\ \frac{2e^x + e^{2x} - 3}{2(6e^x + e^{2x} - 3)} \\ \mathbf{p}_1(x) \\ \mathbf{p}_2(x) \end{pmatrix}$.
- Consider the path $\mathcal{M}_{\mathcal{G}_1}^1 \ni l = (l_1, l_2) = (x, -\log \frac{1-e^{-x}}{1+3e^{-x}}) =: c(x)$.
- $\dot{c}(x) = \left(1, -\frac{4e^x}{(e^x-1)(e^x+3)} \right)$ and $\ddot{c}(x) = \left(0, \frac{4e^x(e^{2x}+3)}{(e^x-1)^2(e^x+3)^2} \right)$.

Proposition 4.1. *The moduli space of figure 8 graphs is **incomplete** under the pressure metric $\|\cdot\|_P$. i.e., $(\mathcal{M}_{\mathcal{G}_1}^1, \|\cdot\|_P)$ is incomplete.*

Proof. By Proposition 3.1, we have

$$\|\dot{c}(x)\|_P^2 = \text{Var}(\dot{c}(x), m_{c(x)}) = 2 \cdot \frac{4e^x(e^{2x}+3)}{(e^x-1)^2(e^x+3)^2} \cdot \mathbf{p}_2(x) = \frac{4e^x(e^{2x}+3)}{-24e^x + 6e^{2x} + 8e^{3x} + e^{4x} + 9}.$$

When x is close to zero, we have the expansion

$$\|\dot{c}(x)\|_P^2 = \frac{1}{x} - \frac{5}{4} + \frac{85x}{48} + o(x^2).$$

This estimate shows that when $x \rightarrow 0$

$$\sqrt{\int \|\dot{c}(x)\|_P^2 dm_{c(x)}} \asymp \frac{1}{\sqrt{x}}.$$

Finally, since $\int_0^1 \frac{1}{\sqrt{x}} dx$ is convergent we see that the metric is incomplete. i.e. the curve arrives at $l_1 = x = 0$ in finite time with respect to this metric. \square

Remark 4.1. There is a hidden natural condition $l_2 > 0$ that we have to take into account. This condition was missing in Section 6 [PS14]. For \mathcal{G}_1 the condition is $1 - e^{-x} > 0$ and $\frac{1-e^{-x}}{1+3e^{-x}} < 1$, which is equivalent to $x > 0$.

Proposition 4.2. *The moduli space of figure 8 graphs is **complete** under the Weil-Petersson metric $\|\cdot\|_{WP}$. i.e., $(\mathcal{M}_{\mathcal{G}_1}^1, \|\cdot\|_{WP})$ is complete.*

Proof. Continue with the computation in the previous proposition, we know that

$$\|\dot{c}(x)\|_{WP}^2 = \frac{\text{Var}(\dot{c}(x), m_{c(x)})}{2x\mathbf{p}_1(x) + 2c(x)\mathbf{p}_2(x)} = -\frac{4e^x(e^{2x} + 3)}{(e^x - 1)(e^x + 3) \left((2e^x + e^{2x} - 3) \log\left(\frac{e^x - 1}{e^x + 3}\right) - 4e^x x \right)}.$$

When x is close to zero, we know that

$$\|\dot{c}(x)\|_{WP}^2 \asymp \frac{-1}{x^2 \log x},$$


and when x tends to infinity we have

$$\|\dot{c}(x)\|_{WP}^2 \asymp \frac{1}{x}.$$

This shows that the curve arrives the boundary of $\mathcal{M}_{\mathcal{G}_1}^1$, i.e. $x = 0$ and $x = \infty$, in infinite time. Because $\mathcal{M}_{\mathcal{G}_1}^1 = \{(x, c(x)); x \in (0, \infty)\}$ is a one dimensional smooth manifold, we can conclude it is complete. \square

4.2 Belt buckles

The second example is a graph with two vertices, connected to each other by three edges which we denote by \mathcal{G}_2 and call it a belt buckle. The picture below is a picture of a belt buckle. For brevity, we denote $l(e_1)$, $l(e_2)$ and $l(e_3)$ by x , y and z , respectively.



Following the recipe, we have

$$\bullet A = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 \end{pmatrix} A_{-l} = \begin{pmatrix} 0 & 0 & 0 & 0 & e^{-x} & e^{-x} \\ 0 & 0 & 0 & e^{-y} & 0 & e^{-y} \\ 0 & 0 & 0 & e^{-z} & e^{-z} & 0 \\ 0 & e^{-x} & e^{-x} & 0 & 0 & 0 \\ e^{-y} & 0 & e^{-y} & 0 & 0 & 0 \\ e^{-z} & e^{-z} & 0 & 0 & 0 & 0 \end{pmatrix}.$$

$$\bullet h(l) = 1 \implies \det(A_{-l} - \text{Id}) = 0 \implies e^{-z} = \frac{1 - e^{-x-y}}{2e^{-x-y} + e^{-x} + e^{-y}}.$$

- $\mathbf{v} = \left(\frac{e^{-y}+1}{2e^{-x-y}+e^{-x}+e^{-y}}, \frac{e^{-x}+1}{2e^{-x-y}+e^{-x}+e^{-y}}, 1, \frac{e^{-y}+1}{2e^{-x-y}+e^{-x}+e^{-y}}, \frac{e^{-x}+1}{2e^{-x-y}+e^{-x}+e^{-y}}, 1 \right)$.
- $P = \begin{pmatrix} 0 & 0 & 0 & 0 & \frac{e^{-x}(e^{-y}+1)}{e^{-x}+1} & \frac{e^{-x}(e^{-y}+1)}{2e^{-x-y}+e^{-x}+e^{-y}} \\ 0 & 0 & 0 & \frac{(e^{-x}+1)e^{-y}}{e^{-y}+1} & 0 & \frac{(e^{-x}+1)b}{2e^{-x-y}+e^{-x}+e^{-y}} \\ 0 & 0 & 0 & \frac{1-e^{-x-y}}{e^{-y}+1} & \frac{1-e^{-x-y}}{e^{-x}+1} & 0 \\ 0 & \frac{e^{-x}(e^{-y}+1)}{e^{-x}+1} & \frac{e^{-x}(e^{-y}+1)}{2e^{-x-y}+e^{-x}+e^{-y}} & 0 & 0 & 0 \\ \frac{(e^{-x}+1)e^{-y}}{e^{-y}+1} & 0 & \frac{(e^{-x}+1)b}{2e^{-x-y}+e^{-x}+e^{-y}} & 0 & 0 & 0 \\ \frac{1-e^{-x-y}}{e^{-y}+1} & \frac{1-e^{-x-y}}{e^{-x}+1} & 0 & 0 & 0 & 0 \end{pmatrix}$.
- $\mathbf{p} = \begin{pmatrix} \mathbf{p}_1(x, y) \\ \mathbf{p}_2(x, y) \\ \mathbf{p}_3(x, y) \\ \mathbf{p}_4(x, y) \\ \mathbf{p}_5(x, y) \\ \mathbf{p}_6(x, y) \end{pmatrix} = \begin{pmatrix} \frac{e^x(e^y+1)^2}{4(3e^{x+y}+e^{2x+y}+e^{x+2y}-1)} \\ \frac{(e^x+1)^2e^y}{4(3e^{x+y}+e^{2x+y}+e^{x+2y}-1)} \\ \frac{(e^x+e^y+2)(e^{x+y}-1)}{4(3e^{x+y}+e^{2x+y}+e^{x+2y}-1)} \\ \mathbf{p}_1(x, y) \\ \mathbf{p}_2(x, y) \\ \mathbf{p}_3(x, y) \end{pmatrix}$.

- Now we consider the surface $\mathcal{M}_{\mathcal{G}_2}^1 \ni l = (l_1, l_2, l_3) = (x, y, -\log\left(\frac{e^{x+y}-1}{e^x+e^y+2}\right)) = (x, y, S(x, y)) = l(x, y)$.
- $\frac{\partial}{\partial x} l = (1, 0, -\frac{e^x(e^y+1)^2}{(e^x+e^y+2)(e^{x+y}-1)})$, $\frac{\partial}{\partial y} l = (0, 1, -\frac{(e^x+1)^2e^y}{(e^x+e^y+2)(e^{x+y}-1)})$,
 $\frac{\partial^2}{\partial x^2} l = (0, 0, \frac{e^x(e^y+1)^2(e^{2x+y}+e^y+2)}{(e^x+e^y+2)^2(e^{x+y}-1)^2})$, $\frac{\partial^2}{\partial x \partial y} l = (0, 0, -\frac{(e^x+1)(e^y+1)e^{x+y}(e^{x+y}-e^x-e^y-3)}{(e^x+e^y+2)^2(e^{x+y}-1)^2})$,
and $\frac{\partial^2}{\partial y^2} l = (0, 0, \frac{(e^x+1)^2e^y(e^{x+2y}+e^x+2)}{(e^x+e^y+2)^2(e^{x+y}-1)^2})$.

Remark 4.2. For this graph, the hidden natural condition $l_3 > 0$ is equivalent to

$$e^{x+y} < 3 + e^x + e^y.$$

In this example, we are interested in the curvature of $\mathcal{M}_{\mathcal{G}_2}^1$ with respect to different metrics $\|\cdot\|_P$ and $\|\cdot\|_{WP}$. Since $\mathcal{M}_{\mathcal{G}_2}^1$ is a two-dimensional manifold in \mathbb{R}^3 , in order to calculate the Gaussian curvature, we first need to derive the corresponding first fundamental forms

$$ds^2 = E(x, y)dx^2 + F(x, y)dxdy + G(x, y)dy^2$$

where $x = l_1$ and $y = l_2$.

Lemma 4.1 (Brioschi formula). *If a metric has local coordinates*

$$ds^2 = E(x, y)dx^2 + F(x, y)dxdy + G(x, y)dy^2,$$

then the curvature is given by

$$K(x, y) = \frac{\det \begin{vmatrix} -\frac{E_{yy}}{2} + F_{xy} - \frac{G_{xx}}{2} & \frac{E_x}{2} & F_x - \frac{E_y}{2} \\ F_y - \frac{G_x}{2} & E & F \\ \frac{G_y}{2} & F & G \end{vmatrix} - \det \begin{vmatrix} 0 & \frac{E_y}{2} & \frac{G_x}{2} \\ \frac{E_y}{2} & E & F \\ \frac{G_x}{2} & F & G \end{vmatrix}}{(EG - F^2)^2}.$$

By direction computations, we have the following propositions.

Proposition 4.3. *The first fundamental form of $(\mathcal{M}_{\mathcal{G}_2}^1, \|\cdot\|_P)$ is*

$$\begin{aligned}
E_P(x, y) &= \text{Var}\left(\frac{\partial l}{\partial x}, m_{-l}\right) = 2 \cdot \frac{\partial^2 S}{\partial x^2} \cdot \mathbf{p}_3(x, y) \\
&= \frac{e^x (e^y + 1)^2 (e^{2x+y} + e^y + 2)}{2(e^x + e^y + 2)(e^{x+y} - 1)(3e^{x+y} + e^{2x+y} + e^{x+2y} - 1)}, \\
F_P(x, y) &= \frac{1}{2} \left(\text{Var}\left(\frac{\partial l}{\partial x} + \frac{\partial l}{\partial y}, m_{-l}\right) - \text{Var}\left(\frac{\partial l}{\partial x}, m_{-l}\right) - \text{Var}\left(\frac{\partial l}{\partial y}, m_{-l}\right) \right) \\
&= 2 \cdot \frac{\partial^2 S}{\partial x \partial y} \cdot \mathbf{p}_3(x, y) \\
&= \frac{(e^x + 1)(e^y + 1)e^{x+y}(-e^{x+y} + e^x + e^y + 3)}{2(e^x + e^y + 2)(e^{x+y} - 1)(3e^{x+y} + e^{2x+y} + e^{x+2y} - 1)}, \\
G_P(x, y) &= \text{Var}\left(\frac{\partial l}{\partial y}, m_{-l}\right) = 2 \cdot \frac{\partial S}{\partial y} \cdot \mathbf{p}_3(x, y) \\
&= \frac{(e^x + 1)^2 e^y (e^{x+2y} + e^x + 2)}{2(e^x + e^y + 2)(e^{x+y} - 1)(3e^{x+y} + e^{2x+y} + e^{x+2y} - 1)}.
\end{aligned}$$

where $S(x, y) = -\log\left(\frac{e^{x+y}-1}{e^x+e^y+2}\right)$ and $l(x, y) = (x, y, S(x, y)) \in \mathcal{M}_{\mathcal{G}_2}^1$.

Proof. Only $F_P(x, y)$ needs some elaboration. By the parallelogram formula and Proposition 3.1, we have

$$\begin{aligned}
F_P(x, y) &= \frac{1}{2} \left(\left\| \frac{\partial l}{\partial x} + \frac{\partial l}{\partial y} \right\|_P^2 - \left\| \frac{\partial l}{\partial x} \right\|_P^2 - \left\| \frac{\partial l}{\partial y} \right\|_P^2 \right) \\
&= \left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} \right)^2 S(x, y) \cdot \mathbf{p}_3(x, y) - \left(\frac{\partial}{\partial x} \right)^2 S(x, y) \cdot \mathbf{p}_3(x, y) - \left(\frac{\partial}{\partial y} \right)^2 S(x, y) \cdot \mathbf{p}_3(x, y) \\
&= 2 \frac{\partial^2}{\partial x \partial y} S(x, y) \cdot \mathbf{p}_3(x, y).
\end{aligned}$$

□

We notice that the same computation holds for Weil-Petersson metric $(\|\cdot\|_{WP})$. Therefore, we have the following result.

Proposition 4.4. *The first fundamental form of $(\mathcal{M}_{\mathcal{G}_2}^1, \|\cdot\|_{WP})$ is*

$$\begin{aligned}
E_{WP}(x, y) &= \frac{\text{Var}\left(\frac{\partial l}{\partial x}, m_{-l}\right)}{V(l)} \\
&= \frac{e^x (e^y + 1)^2 (e^{2x+y} + e^y + 2)}{(e^x + e^y + 2)(e^{x+y} - 1) f(x, y)}, \\
F_{WP}(x, y) &= \frac{\frac{1}{2} \left(\text{Var}\left(\frac{\partial l}{\partial x} + \frac{\partial l}{\partial y}, m_{-l}\right) - \text{Var}\left(\frac{\partial l}{\partial x}, m_{-l}\right) - \text{Var}\left(\frac{\partial l}{\partial y}, m_{-l}\right) \right)}{V(l)} \\
&= -\frac{(e^x + 1)(e^y + 1)e^{x+y}(e^{x+y} - e^x - e^y - 3)}{(e^x + e^y + 2)(e^{x+y} - 1) f(x, y)}, \\
G_{WP}(x, y) &= \frac{\text{Var}\left(\frac{\partial l}{\partial y}, m_{-l}\right)}{V(l)} \\
&= \frac{(e^x + 1)^2 e^y (e^{x+2y} + e^x + 2)}{(e^x + e^y + 2)(e^{x+y} - 1) f(x, y)},
\end{aligned}$$

where $V(l) = \int l dm_{-l} = 2(l_1 \cdot \mathbf{p}_1 + l_2 \cdot \mathbf{p}_2 + l_3 \cdot \mathbf{p}_3)$ and

$$f(x, y) = \left(x e^{x+2y} + y e^{2x+y} + 2e^{x+y}(x+y) + (-2e^{x+y} - e^{2x+y} - e^{x+2y} + e^x + e^y + 2) \log \left(\frac{e^{x+y} - 1}{e^x + e^y + 2} \right) + e^x x + e^y y \right)$$

Proposition 4.5. *The moduli space of belt buckles is bounded positively curved under the pressure metric $\|\cdot\|_P$. i.e., $(\mathcal{M}_{\mathcal{G}_2}^1, \|\cdot\|_P)$ is positively curved and the Gaussian curvature is bounded. Moreover, $(\mathcal{M}_{\mathcal{G}_2}^1, \|\cdot\|_P)$ is incomplete.*

Proof. By the Brioschi formula, we can write down the curvature explicitly as the following

$$K_P(x, y) = \frac{1}{4(e^x + 1)^2 (e^y + 1)^2 (3e^{x+y} + e^{2x+y} + e^{x+2y} - 1)} \cdot (5 + 6e^x + 3e^{2x} + 6e^y + 3e^{2y} + 3e^{x+y} + 45e^{2(x+y)} + 19e^{3(x+y)} + 11e^{2x+y} + 9e^{3x+y} + 3e^{4x+y} + 11e^{x+2y} + 33e^{3x+2y} + 8e^{4x+2y} + 9e^{x+3y} + 33e^{2x+3y} + 3e^{4x+3y} + 3e^{x+4y} + 8e^{2x+4y} + 3e^{3x+4y}).$$

We observe that the numerator and the denominator of $K(x, y)$ have the same highest exponents e^{3x+4y} and e^{4x+3y} . Consider the polar coordinate $e^x = r \cos \theta$ and $e^y = r \sin \theta$ where $1 < r < \infty$ and $\theta \in (0, \frac{\pi}{2})$

$$K_P(x, y) = \frac{6r^7(\cos^3 \theta \sin^4 \theta + \cos^4 \theta \sin^3 \theta) + HOT(r^6)}{8r^7(\cos^3 \theta \sin^4 \theta + \cos^4 \theta \sin^3 \theta) + HOT(r^6)}.$$

Therefore, we know $K_P(x, y)$ is close to $\frac{3}{4}$ when r is large, and it is easy see that $K_p(x, y) > 0$ for $x, y \geq 0$. Because $B := \{(x, y); x \geq 0, y \geq 0, e^{2x} + e^{2y} \leq r^2\}$ is a compact set, we know $c_1 \geq K_p(x, y) \geq c_0 > 0$.

To prove the second assertion, we consider the path $c(x) = l(x, x) = (x, x, -\log(\frac{1-e^{-2x}}{2e^{-2x}+2e^{-x}}))$. We repeat the argument that we used in Proposition 4.1. Because when x goes to 0, $c(x)$ goes to the boundary of $\mathcal{M}_{\mathcal{G}_2}^1$, to prove the incompleteness we only need to show that $c(x)$ goes to boundary in finite time. Since


$$\begin{aligned} \|\dot{c}(x)\|_P^2 &= P_3(x, x) \cdot \frac{d^2}{dx^2} \left(-\log \left(\frac{1 - e^{-2x}}{2e^{-2x} + 2e^{-x}} \right) \right) + P_6(x, x) \cdot \frac{d^2}{dx^2} \left(-\log \left(\frac{1 - e^{-2x}}{2e^{-2x} + 2e^{-x}} \right) \right) \\ &= \frac{e^x}{-3e^x + 2e^{2x} + 1}, \end{aligned}$$

it is clear that $\int_0^1 \|\dot{c}(x)\|_P dx$ is convergent. Hence the geodesic distance from $c(1)$ to the boundary point $c(0)$ is finite. \square

Observation 4.1. *The sectional curvature of $(\mathcal{M}_{\mathcal{G}_2}^1, \|\cdot\|_{WP})$ is negative and bounded.*

Evidence. By the Brioschi formula, we can explicitly write down the curvature. However, it is too long and unnecessary to state it in the proof. We put the explicit expression of $K_{WP}(x, y)$ in the appendix for whom is interested in that.


The following is the graph of $K_{WP}(x, y)$ provided $0 < x, y < 5$ and the hidden condition $e^{x+y} < 3 + e^x + e^y$. With the explicit formula of K_{WP} , we can easily plot the picture and estimate the maximum and the minimal of K_{WP} through softwares, and here we use Mathematica. By Mathematica, we know the maximum of $K_{WP}(x, y)$ for $0 < x, y$ and $e^{x+y} < 3 + e^x + e^y$ happens around $(x_0, y_0) \approx (1.09861, 1.09861)$ where $K_{WP}(x_0, y_0) \approx -0.485025$, and the lower bound of K_{WP} is around -0.564958 .



□

4.3 Dumbbells

In this subsection, we discuss the graph of a dumbbell. We denote this graph by \mathcal{G}_3 and the picture below is what it looks like. Likewise, for short, we denote $l(e_1)$, $l(e_2)$ and $l(e_3)$ by x , y and z , respectively.



Following the same recipe, we have

$$\bullet A = \begin{pmatrix} 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 & 0 & 0 \end{pmatrix} \quad A_{-l} = \begin{pmatrix} e^{-x} & 0 & e^{-x} & 0 & 0 & 0 \\ 0 & e^{-y} & 0 & 0 & 0 & e^{-y} \\ 0 & e^{-z} & 0 & 0 & e^{-z} & 0 \\ 0 & 0 & e^{-x} & e^{-x} & 0 & 0 \\ 0 & 0 & 0 & 0 & e^{-y} & e^{-y} \\ e^{-z} & 0 & 0 & e^{-z} & 0 & 0 \end{pmatrix}.$$

$$\bullet h(l) = 1 \implies \det(A_{-l} - \text{Id}) = 0 \implies e^{-z} = \sqrt{\frac{e^{-x-y} - e^{-x} - e^{-y} + 1}{4e^{-x-y}}}.$$

$$\bullet \mathbf{v} = \left(\frac{\sqrt{\frac{(e^{-x}-1)(e^{-y}-1)}{e^{-x-y}}}}{2-2e^{-x}}, \frac{1}{2e^{-y}}, -\frac{e^{-x}\sqrt{\frac{(e^{-x}-1)(e^{-y}-1)}{e^{-x-y}}}}{e^{-x}-1}, \frac{\sqrt{\frac{(e^{-x}-1)(e^{-y}-1)}{e^{-x-y}}}}{2-2e^{-x}}, \frac{1}{2e^{-y}}, 1 \right).$$

$$\bullet P = \begin{pmatrix} e^{-x} & 0 & \frac{1}{2} & 0 & 0 & 0 \\ 0 & e^{-y} & 0 & 0 & 0 & \frac{1}{2} \\ 0 & 1 - e^{-y} & 0 & 0 & 1 - e^{-y} & 0 \\ 0 & 0 & \frac{1}{2} & a & 0 & 0 \\ 0 & 0 & 0 & 0 & e^{-y} & \frac{1}{2} \\ 1 - e^{-x} & 0 & 0 & 1 - e^{-x} & 0 & 0 \end{pmatrix}.$$

$$\bullet \mathbf{p} = \begin{pmatrix} \mathbf{p}_1(x, y) \\ \mathbf{p}_2(x, y) \\ \mathbf{p}_3(x, y) \\ \mathbf{p}_4(x, y) \\ \mathbf{p}_5(x, y) \\ \mathbf{p}_6(x, y) \end{pmatrix} = \begin{pmatrix} \frac{e^x(-1+e^y)}{4-6e^x-6e^y+8e^{x+y}} \\ \frac{e^y(-1+e^x)}{4-6e^x-6e^y+8e^{x+y}} \\ \frac{(-1+e^x)(-1+e^y)}{2-3e^x-3e^y+4e^{x+y}} \\ \mathbf{p}_1(x, y) \\ \mathbf{p}_2(x, y) \\ \mathbf{p}_3(x, y) \end{pmatrix}.$$

• Now we consider the surface $\mathcal{M}_{\mathcal{G}_3}^1 \ni l = (l_1, l_2, l_3) = (x, y, \log(2) - \frac{1}{2} \log((e^x - 1)(e^y - 1))) = l(x, y)$.

• $\frac{\partial}{\partial x} l = (1, 0, \frac{e^x}{2-2e^x})$, $\frac{\partial}{\partial y} l = (0, 1, \frac{e^y}{2-2e^y})$, $\frac{\partial^2}{\partial x^2} l = (0, 0, \frac{e^x}{2(e^x-1)^2})$, $\frac{\partial^2}{\partial x \partial y} l = (0, 0, 0)$, and $\frac{\partial^2}{\partial y^2} l = (0, 0, \frac{e^y}{2(e^y-1)^2})$.

Remark 4.3. There is a hidden condition $l_3 > 0$. In this case, the condition is equivalent to

$$4 > (e^x - 1)(e^y - 1).$$

Set $x = l_1$ and $y = l_2$. Since $\mathcal{M}_{\mathcal{G}_3}^1$ is also a two-dimensional manifold in \mathbb{R}^3 , so we repeat the same argument as for $\mathcal{M}_{\mathcal{G}_2}^1$. Following propositions are coming from direct computations.

Proposition 4.6. *The first fundamental form of $(\mathcal{M}_{\mathcal{G}_3}^1, \|\cdot\|_P)$ is*

$$\begin{aligned} E_P(x, y) &= \text{Var}\left(\frac{\partial l}{\partial x}, m_{-l}\right) = 2 \cdot \frac{\partial l}{\partial x} \cdot \mathbf{p}_3(x, y) \\ &= \frac{e^x(e^y - 1)}{(e^x - 1)(4e^{x+y} - 3e^x - 3e^y + 2)}, \\ F_P(x, y) &= \frac{1}{2} \left(\text{Var}\left(\frac{\partial l}{\partial x} + \frac{\partial l}{\partial y}, m_{-l}\right) - \text{Var}\left(\frac{\partial l}{\partial x}, m_{-l}\right) - \text{Var}\left(\frac{\partial l}{\partial y}, m_{-l}\right) \right) \\ &= 0, \\ G_P(x, y) &= \text{Var}\left(\frac{\partial l}{\partial y}, m_{-l}\right) = 2 \cdot \frac{\partial l}{\partial y} \cdot \mathbf{p}_3(x, y) \\ &= \frac{(e^x - 1)e^y}{(e^y - 1)(4e^{x+y} - 3e^x - 3e^y + 2)}. \end{aligned}$$

Proposition 4.7. *The first fundamental form of $(\mathcal{M}_{\mathcal{G}_3}^1, \|\cdot\|_{WP})$ is*

$$\begin{aligned} E_{WP}(x, y) &= \frac{\text{Var}\left(\frac{\partial l}{\partial x}, m_{-l}\right)}{V(l)} \\ &= \frac{e^x(e^y - 1)}{(e^x - 1)f(x, y)}, \\ F_{WP}(x, y) &= \frac{\frac{1}{2} \left(\text{Var}\left(\frac{\partial l}{\partial x} + \frac{\partial l}{\partial y}, m_{-l}\right) - \text{Var}\left(\frac{\partial l}{\partial x}, m_{-l}\right) - \text{Var}\left(\frac{\partial l}{\partial y}, m_{-l}\right) \right)}{V(l)} \\ &= 0, \\ G_{WP}(x, y) &= \frac{\text{Var}\left(\frac{\partial l}{\partial y}, m_{-l}\right)}{V(l)} \\ &= \frac{(e^x - 1)e^y}{(e^y - 1)f(x, y)}, \end{aligned}$$

where $V(l) = \int l dm_{-l} = 2(l_1 \cdot \mathbf{p}_1 + l_2 \cdot \mathbf{p}_2 + l_3 \cdot \mathbf{p}_3)$ and

$$f(x, y) = \left(xe^{x+y} + ye^{x+y} - \log((e^x - 1)(e^y - 1)) + 2(-e^{x+y} + e^x + e^y) \log\left(\frac{1}{2}\sqrt{(e^x - 1)(e^y - 1)}\right) - e^x x - e^y y + \log(4) \right)$$

Proposition 4.8. *The moduli space of dumbbells is positively curved under the pressure metric $\|\cdot\|_P$. i.e., $(\mathcal{M}_{\mathcal{G}_3}^1, \|\cdot\|_P)$ is positively curved. More precisely, the Gaussian curvature is strictly bigger than zero, but has no upper bound. Moreover, $(\mathcal{M}_{\mathcal{G}_3}^1, \|\cdot\|_P)$ is incomplete.*

Proof. Applying the Brioschi formula, we can write down the curvature explicitly:

$$K_P(x, y) = \frac{2e^{x+y} - 1}{4e^{x+y} - 3e^x - 3e^y + 2}$$

where $x = l_1$ and $y = l_2$. It is clear that $K_P(x, y) > 0$ for all $x, y > 0$, and $\lim_{(x,y) \rightarrow (0,0)} K_P(x, y) = \infty$. Moreover, set $a = e^x$ and $b = e^y$,

$$K_P(a, b) = \frac{2ab - 1}{4ab - 3a - 3b + 2}$$

provided $a > 1$, $b > 1$ and $(a - 1)(b - 1) < 4$. Since $\partial_a K_P(a, b) = 0$ and $\partial_b K_P(a, b) = 0$ has no real solution, we know the extreme values of $K_P(a, b)$ could only happen on the boundaries $a = 1$, $b = 1$ or $(a - 1)(b - 1) < 4$. Easy computation shows that $K_P(a, b)$ is strictly bigger than 0 on these boundaries. Therefore, we can conclude that $K_P(x, y) > c_0 > 0$ for some c_0 . And Mathematica computation indicates that c_0 is about 0.85.


To prove the last assertion, we consider the path $c(x) = l(x, x) = (x, x, \log(2) - \frac{1}{2} \log((e^x - 1)(e^y - 1)))$. We repeat the argument that we used in Proposition 4.1. Because when x goes to 0, $c(x)$ goes to the boundary of $\mathcal{M}_{\mathcal{G}_2}^1$, to prove the incompleteness we only need to show that $c(x)$ goes to boundary in finite time. Since

$$\begin{aligned} \|\dot{c}(x)\|_P^2 &= P_3(x, x) \cdot \frac{d^2}{dx^2} (\log(2) - \log(e^x - 1)) + P_6(x, x) \cdot \frac{d^2}{dx^2} (\log(2) - \log(e^x - 1)) \\ &= \frac{e^x}{-3e^x + 2e^{2x} + 1}, \end{aligned}$$

it is clear that $\int_0^1 \|\dot{c}(x)\|_P dx$ is convergent. Hence the geodesic distance from $c(1)$ to the boundary point $c(0)$ is finite. \square

Observation 4.2. *The sectional curvature of $(\mathcal{M}_{\mathcal{G}_3}^1, \|\cdot\|_{WP})$ takes positive and negative values.*


Evidence. It is the same as in the belt buckle case, using the Brioschi formula, we can have the explicit expression of $K_{WP}(x, y)$ which, in the sake of brevity, is stated the appendix. The following figure is the graph of $K_{WP}(x, y)$ produced by Mathematica for $0.05 < x, y < 3$ and. Moreover, Mathematica computation shows that K_{WP} takes positive (e.g. when x is close to 0.05 and y is close to 3) and negative values (e.g. when x and y are close to 0.05).



\square

4.4 Three-petal roses

The fourth example is a three-petal rose \mathcal{G}_4 , which could be thought of as a generalization of the figure 8. The picture below is a picture of a three-petal rose. For brevity, we denote $l(e_1)$, $l(e_2)$ and $l(e_3)$ by x , y and z , respectively.



Follow the recipe, we have:

$$\bullet A = \begin{pmatrix} 1 & 1 & 1 & 0 & 1 & 1 \\ 1 & 1 & 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 1 & 1 & 1 \end{pmatrix}, A_{-l} = \begin{pmatrix} e^{-x} & e^{-x} & e^{-x} & 0 & e^{-x} & e^{-x} \\ e^{-y} & e^{-y} & e^{-y} & e^{-y} & 0 & e^{-y} \\ e^{-z} & e^{-z} & e^{-z} & e^{-z} & e^{-z} & 0 \\ 0 & e^{-x} & e^{-x} & e^{-x} & e^{-x} & e^{-x} \\ e^{-y} & 0 & e^{-y} & e^{-y} & e^{-y} & e^{-y} \\ e^{-z} & e^{-z} & 0 & e^{-z} & e^{-z} & e^{-z} \end{pmatrix}.$$

$$\bullet h(l) = 1 \implies \det(A_{-l} - Id) = 0 \implies e^{-z} = -\frac{-e^{x+y} + e^x + e^y + 3}{e^{x+y} + 3e^x + 3e^y + 5}.$$

$$\bullet \mathbf{v} = \left(\frac{2(e^{-y}+1)}{e^{-x}(5e^{-y}+3)+3e^{-y}+1}, \frac{2(e^{-x}+1)}{e^{-x}(5e^{-y}+3)+3e^{-y}+1}, 1, \frac{2(e^{-y}+1)}{e^{-x}(5e^{-y}+3)+3e^{-y}+1}, \frac{2(e^{-x}+1)}{e^{-x}(5e^{-y}+3)+3e^{-y}+1}, 1 \right).$$

$$\bullet P = \begin{pmatrix} \frac{2(1+e^{-y})}{e^{-x}(3+5e^{-y})+3e^{-y}+1} & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{2(1+e^{-x})}{e^{-x}(3+5e^{-y})+3e^{-y}+1} & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{2(1+e^{-y})}{e^{-x}(3+5e^{-y})+3e^{-y}+1} & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{2(1+e^{-x})}{e^{-x}(3+5e^{-y})+3e^{-y}+1} & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

$$\bullet \mathbf{p} = \begin{pmatrix} \mathbf{p}_1(x, y) \\ \mathbf{p}_2(x, y) \\ \mathbf{p}_3(x, y) \\ \mathbf{p}_4(x, y) \\ \mathbf{p}_5(x, y) \\ \mathbf{p}_6(x, y) \end{pmatrix} = \begin{pmatrix} \frac{2e^x(e^y+1)^2}{12e^{x+y}+e^{2(x+y)}+6e^{2x+y}+6e^{x+2y}-10e^x-3e^{2x}-10e^y-3e^{2y}-15} \\ \frac{2(e^x+1)^2e^y}{12e^{x+y}+e^{2(x+y)}+6e^{2x+y}+6e^{x+2y}-10e^x-3e^{2x}-10e^y-3e^{2y}-15} \\ \frac{(e^{x+y}-e^x-e^y-3)(e^{x+y}+3e^x+3e^y+5)}{2(12e^{x+y}+e^{2(x+y)}+6e^{2x+y}+6e^{x+2y}-10e^x-3e^{2x}-10e^y-3e^{2y}-15)} \\ \mathbf{p}_1(x, y) \\ \mathbf{p}_2(x, y) \\ \mathbf{p}_3(x, y) \end{pmatrix}.$$

$$\bullet \text{ Now we consider the surface } \mathcal{M}_{\mathcal{G}_4}^1 \ni l = (l_1, l_2, l_3) = (x, y, -\log(-\frac{-e^{x+y} + e^x + e^y + 3}{e^{x+y} + 3e^x + 3e^y + 5})) = l(x, y).$$

$$\bullet \frac{\partial}{\partial x} l = (1, 0, \frac{4e^x(e^y+1)^2}{(-e^{x+y}+e^x+e^y+3)(e^{x+y}+3e^x+3e^y+5)}), \frac{\partial l}{\partial y} = (0, 1, \frac{4(e^x+1)^2e^y}{(-e^{x+y}+e^x+e^y+3)(e^{x+y}+3e^x+3e^y+5)}),$$

$$\frac{\partial^2}{\partial x^2} l = (0, 0, \frac{4e^x(e^y+1)^2(e^y+3)(e^{2x+y}-e^{2x}+3e^y+5)}{(-e^{x+y}+e^x+e^y+3)^2(e^{x+y}+3e^x+3e^y+5)^2}), \frac{\partial^2}{\partial x \partial y} l = (0, 0, \frac{32(e^x+1)(e^y+1)e^{x+y}(e^x+e^y+2)}{(-e^{x+y}+e^x+e^y+3)^2(e^{x+y}+3e^x+3e^y+5)^2}),$$

$$\text{ and } \frac{\partial^2}{\partial y^2} l = (0, 0, \frac{4(e^x+1)^2(e^x+3)e^y(e^{x+2y}+3e^x-e^{2y}+5)}{(-e^{x+y}+e^x+e^y+3)^2(e^{x+y}+3e^x+3e^y+5)^2}).$$

Remark 4.4. There is a hidden condition $l_3 > 0$. In this case, the condition is equivalent to

$$-e^{x+y} + e^x + e^y + 3 < 0.$$

Since $\mathcal{M}_{\mathcal{G}_4}^1$ is still a two-dimensional manifold in \mathbb{R}^3 , so we repeat the same argument as for $\mathcal{M}_{\mathcal{G}_2}^1$. Set $x = l_1$, $y = l_2$ we have the following results of the first fundamental form with respect to $\|\cdot\|_P$ and $\|\cdot\|_{WP}$.

Proposition 4.9. *The first fundamental form of $(\mathcal{M}_{\mathcal{G}_4}^1, \|\cdot\|_P)$ is*

$$\begin{aligned} E_P(x, y) &= \text{Var}\left(\frac{\partial l}{\partial x}, m_{-l}\right) = 2 \cdot \frac{\partial l}{\partial x} \cdot \mathbf{p}_3(x, y) \\ &= -\frac{4e^x (e^y + 1)^2 (e^y + 3) (e^{2x+y} - e^{2x} + 3e^y + 5)}{(-e^{x+y} + e^x + e^y + 3) (e^{x+y} + 3e^x + 3e^y + 5) w(x, y)}, \\ F_P(x, y) &= \frac{1}{2} \left(\text{Var}\left(\frac{\partial l}{\partial x} + \frac{\partial l}{\partial y}, m_{-l}\right) - \text{Var}\left(\frac{\partial l}{\partial x}, m_{-l}\right) - \text{Var}\left(\frac{\partial l}{\partial y}, m_{-l}\right) \right) \\ &= -\frac{32 (e^x + 1) (e^y + 1) e^{x+y} (e^x + e^y + 2)}{(-e^{x+y} + e^x + e^y + 3) (e^{x+y} + 3e^x + 3e^y + 5) w(x, y)}, \\ G_P(x, y) &= \text{Var}\left(\frac{\partial l}{\partial y}, m_{-l}\right) = 2 \cdot \frac{\partial l}{\partial y} \cdot \mathbf{p}_3(x, y) \\ &= -\frac{4 (e^x + 1)^2 (e^x + 3) e^y (e^{x+2y} + 3e^x - e^{2y} + 5)}{(-e^{x+y} + e^x + e^y + 3) (e^{x+y} + 3e^x + 3e^y + 5) w(x, y)}. \end{aligned}$$

where $w(x, y) = (12e^{x+y} + e^{2(x+y)} + 6e^{2x+y} + 6e^{x+2y} - 10e^x - 3e^{2x} - 10e^y - 3e^{2y} - 15)$

Proposition 4.10. *The first fundamental form of $(\mathcal{M}_{\mathcal{G}_4}^1, \|\cdot\|_{WP})$ is*


$$\begin{aligned} E_{WP}(x, y) &= \frac{\text{Var}\left(\frac{\partial l}{\partial x}, m_{-l}\right)}{V(l)} \\ &= \frac{4e^x (e^y + 1)^2 (e^y + 3) (e^{2x+y} - e^{2x} + 3e^y + 5)}{f_1(x, y) f_2(x, y) f_3(x, y)}, \\ F_{WP}(x, y) &= \frac{\frac{1}{2} \left(\text{Var}\left(\frac{\partial l}{\partial x} + \frac{\partial l}{\partial y}, m_{-l}\right) - \text{Var}\left(\frac{\partial l}{\partial x}, m_{-l}\right) - \text{Var}\left(\frac{\partial l}{\partial y}, m_{-l}\right) \right)}{V(l)} \\ &= \frac{32 (e^x + 1) (e^y + 1) e^{x+y} (e^x + e^y + 2)}{f_1(x, y) f_2(x, y) f_3(x, y)}, \\ G_{WP}(x, y) &= \frac{\text{Var}\left(\frac{\partial l}{\partial y}, m_{-l}\right)}{V(l)} \\ &= \frac{4 (e^x + 1)^2 (e^x + 3) e^y (e^{x+2y} + 3e^x - e^{2y} + 5)}{f_1(x, y) f_2(x, y) f_3(x, y)}, \end{aligned}$$

where $V(l) = \int l dm_{-l} = 2(l_1 \cdot \mathbf{p}_1 + l_2 \cdot \mathbf{p}_2 + l_3 \cdot \mathbf{p}_3)$ and

$$\begin{aligned} f_1(x, y) &= -e^{x+y} + e^x + e^y + 3, \\ f_2(x, y) &= e^{x+y} + 3e^x + 3e^y + 5 \\ f_3(x, y) &= \left[-4 (xe^{x+2y} + ye^{2x+y} + 2e^{x+y}(x+y) + e^x x + e^y y) + \right. \\ &\quad \left. (-4e^{x+y} + e^{2(x+y)} + 2e^{2x+y} + 2e^{x+2y} - 14e^x - 3e^{2x} - 14e^y - 3e^{2y} - 15) \cdot \right. \\ &\quad \left. \log \left(-\frac{-e^{x+y} + e^x + e^y + 3}{e^{x+y} + 3e^x + 3e^y + 5} \right) \right] \end{aligned}$$

Observation 4.3. *The moduli space of three-petal roses is positively curved under the pressure metric $\|\cdot\|_P$. i.e., $(\mathcal{M}_{\mathcal{G}_4}^1, \|\cdot\|_P)$ is positively curved and the Gaussian curvature is bounded. Moreover, $(\mathcal{M}_{\mathcal{G}_4}^1, \|\cdot\|_P)$ is incomplete.*

Evidence. Applying the Brioschi formula, we can write down the curvature explicitly; however, in this note we only give the figure of the curvature and avoid stating lengthy results.




Moreover, numerical results indicate that $0.2 < K_P(x, y) < 1$.

The second assertion is because each three-petal rose contains a figure 8, and we know figure 8 is incomplete with respect to the pressure metric.

□

Observation 4.4. *The sectional curvature of $(\mathcal{M}_{\mathcal{G}_4}^1, \|\cdot\|_{WP})$ takes positive and negative values.*

Evidence. The explicit formula of K_{WP} is complex and make no sense to state in here. The following figure of K_{WP} for $0.5 < x, y < 20$ is produced by Mathematica , which indicates that K_{WP} is positive when $(x, y) = (5, 15)$ and negative when $(x, y) = (19, 19)$.



□

5 Appendix

Proposition 5.1. *The curvature of $(\mathcal{M}_{\mathcal{G}_2}^1, \|\cdot\|_{WP})$ could be written explicitly as the following. For $0 < x, y$ and $e^{x+y} < 3 + e^x + e^y$, we have*

$$\begin{aligned}
K_{WP}(x, y) = & \frac{1}{4(e^x + 1)^2 (e^y + 1)^2 f(x, y)} \cdot (e^x x(x - y - 4) + 2e^{2x} x(x - y - 4) + 2x(y - 2)e^{3x+4y} + 2(x - 2)ye^{4x+3y} - \\
& (x + 4)e^{3y}y - e^{3x}x(y + 4) + e^y y(-x + y - 4) + 2e^{2y}y(-x + y - 4) + ye^{4x+y}(2x + y - 4) + \\
& 2ye^{4x+2y}(2x + y - 4) + xe^{x+4y}(x + 2y - 4) + 2xe^{2x+4y}(x + 2y - 4) + 4e^{x+y}(x^2 - 4x + (y - 4)y) + \\
& 2e^{x+3y}(2x^2 + x(3y - 8) - 8y) + 2e^{3x+y}(x(3y - 8) + 2(y - 4)y) + 4e^{3(x+y)}(x(3y - 4) - 4y) + \\
& e^{2x+y}(8x^2 + x(5y - 32) + 6(y - 4)y) + e^{x+2y}(6x^2 + x(5y - 24) + 8(y - 4)y) + \\
& 4e^{2(x+y)}(3x^2 + x(5y - 12) + 3(y - 4)y) + e^{2x+3y}(8x^2 + x(17y - 32) - 24y) + e^{3x+2y}(x(17y - 24) + 8(y - 4)y) - \\
& (5(x + 4)e^y + 4(x + 4)e^{2y} + (x + 4)e^{3y} + 4e^{2x+4y}(x + y - 2) + 5e^x(y + 4) + 4e^{2x}(y + 4) + e^{3x}(y + 4) + \\
& 4e^{4x+2y}(x + y - 2) + 2(x + y + 4) + 2e^{x+y}(x + y + 20) + e^{4x+y}(2x + y - 4) + e^{4x+3y}(2x + y - 4) + 4e^{3x+y}(2x + y - 2) + \\
& e^{x+4y}(x + 2y - 4) + e^{3x+4y}(x + 2y - 4) + 4e^{x+3y}(x + 2y - 2) + 2e^{3(x+y)}(5x + 5y - 12) + e^{2x+y}(11x + 5y + 16) + \\
& e^{x+2y}(5x + 11y + 16) + e^{2x+3y}(19x + 17y - 32) + e^{3x+2y}(17x + 19y - 32) + e^{2(x+y)}(26x + 26y - 24)) \\
& (e^{x+y} - 1)^2 (6e^{x+y} + e^{2x+y} + e^{x+2y} + 7e^x + 2e^{2x} + 7e^y + 2e^{2y} + 6) \log^2 \left(\frac{e^{x+y} - 1}{e^x + e^y + 2} \right) + \log \left(\frac{e^{x+y} - 1}{e^x + e^y + 2} \right)
\end{aligned}$$

where

$$f(x, y) = \left(xe^{x+2y} + ye^{2x+y} + 2e^{x+y}(x + y) + (-2e^{x+y} - e^{2x+y} - e^{x+2y} + e^x + e^y + 2) \log \left(\frac{e^{x+y} - 1}{e^x + e^y + 2} \right) + e^x x + e^y y \right).$$

Proposition 5.2. *The curvature of $(\mathcal{M}_{\mathcal{G}_3}^1, \|\cdot\|_{WP})$ could be written explicitly as the following. For $0 < x, y$ and $4 > (e^x - 1)(e^y - 1)$, we have*

$$\begin{aligned}
K_{WP}(x, y) = & -\{4(e^x - 1)(e^y - 1) \cdot \\
& \left(xe^{x+y} + ye^{x+y} - \log((e^x - 1)(e^y - 1)) + 2(-e^{x+y} + e^x + e^y) \log \left(\frac{1}{2} \sqrt{(e^x - 1)(e^y - 1)} \right) - e^x x - e^y y + \log(4) \right) \}^{-1} \cdot \\
& \left\{ -4e^x x + 4e^{2x} x + 8xe^{x+y} + 4xe^{2(x+y)} - 8xe^{2x+y} - 4xe^{x+2y} + e^x x^2 - e^{2x} x^2 - 2x^2 e^{x+y} - x^2 e^{2(x+y)} + 2x^2 e^{2x+y} + \right. \\
& x^2 e^{x+2y} - 4e^y y + 4e^{2y} y + 8ye^{x+y} + 4ye^{2(x+y)} - 4ye^{2x+y} - 8ye^{x+2y} - e^x xy + e^{2x} xy - xe^y y + xe^{2y} y - \\
& - 2y^2 e^{x+y} - y^2 e^{2(x+y)} + y^2 e^{2x+y} + 2y^2 e^{x+2y} + 2xye^{2(x+y)} + xye^{2x+y} + xye^{x+2y} + e^y y^2 - e^{2y} y^2 - \\
& x \log(4) + xe^y \log(4) - y \log(4) + e^x y \log(4) - (x(e^y - 1) + (e^x - 1)y) \log((e^x - 1)(e^y - 1)) + \\
& 2[-2(x - 4)e^{2x+y} - 16e^{x+y} + (3x + 8)e^y + e^{2x}(x - y - 4) - 2(y - 4)e^{x+2y} + e^{2y}(-x + y - 4) \\
& \left. \left. (+e^{2(x+y)}(x + y - 4) - 2(x + y + 2) + e^x(3y + 8)) \right] \log \left(\frac{1}{2} \sqrt{(e^x - 1)(e^y - 1)} \right) \right\}
\end{aligned}$$

References

- [BCLS13] Martin Bridgeman, Richard Canary, François Labourie, and Andrés Sambarino, *The pressure metric for Anosov representations*, arXiv.org (2013).
- [CV86] Marc Culler and Karen Vogtmann, *Moduli of graphs and automorphisms of free groups*, *Inventiones Mathematicae* **84** (1986), no. 1, 91–119.
- [JP07] Thomas Jordan and Mark Pollicott, *Multifractal analysis and the variance of Gibbs measures*, *Journal of the London Mathematical Society. Second Series* **76** (2007), no. 1, 57–72.
- [Mcm08] Curtis McMullen, *Thermodynamics, dimension and the Weil-Petersson metric*, *Inventiones Mathematicae* **173** (2008), no. 2, 365–425.
- [Pol94] Mark Pollicott, *Derivatives of topological entropy for Anosov and geodesic flows*, *Journal of Differential Geometry* **39** (1994), no. 3, 457–489.
- [PP90] William Parry and Mark Pollicott, *Zeta functions and the periodic orbit structure of hyperbolic dynamics*, *Astérisque* (1990), no. 187-188, 268.
- [PS14] Mark Pollicott and Richard Sharp, *A Weil-Petersson type metric on spaces of metric graphs*, *Geometriae Dedicata* **172** (2014), no. 1, 229–244.