

Complex reflection groups, logarithmic connections and bi-flat F-manifolds

Alessandro Arsie* and Paolo Lorenzoni**

*Department of Mathematics and Statistics

The University of Toledo, 2801 W. Bancroft St., 43606 Toledo, OH, USA

**Dipartimento di Matematica e Applicazioni

Università di Milano-Bicocca, Via Roberto Cozzi 53, I-20125 Milano, Italy

*alessandro.arsie@utoledo.edu, **paolo.lorenzoni@unimib.it

Abstract

We show that bi-flat F -manifolds can be interpreted as natural geometrical structures encoding the almost duality for Frobenius manifolds without metric. Using this framework, we extend Dubrovin's duality between orbit spaces of Coxeter groups and Veselov's \vee -systems, to the orbit spaces of exceptional well-generated complex reflection groups of rank 2 and 3. On the Veselov's \vee -systems side, we provide a generalization of the notion of \vee -systems that gives rise to a dual connection which coincides with a Dunkl-Kohno-type connection associated with such groups. In particular, this allows us to treat on the same ground several different examples including Coxeter and Shephard groups. Remarkably, as a byproduct of our results, we prove that in some examples basic flat invariants are not uniquely defined. As far as we know, such a phenomenon has never been pointed out before.

Contents

1	Introduction	4
2	Frobenius manifolds and their almost dual	9
2.1	Frobenius manifolds and space of orbits of Coxeter groups	10
2.1.1	The case of A_3	11
2.1.2	The case of $H_3 (G_{23})$	12

2.2	Frobenius manifolds and space of orbits of Shephard groups	13
2.2.1	The case of G_{26}	13
2.2.2	Shephard groups of rank 2 and 3	14
3	V-systems	15
3.1	An example: A_3	17
4	Flat and bi-flat F-manifolds	17
4.1	Almost duality and bi-flat F -manifolds	18
5	Bi-flat F-structure on the space of orbits of complex reflection groups	21
5.1	The case of G_4	24
5.2	The case of G_5	25
5.3	The case of G_6	25
5.4	The case of G_8	26
5.5	The case of G_9	26
5.6	The case of G_{10}	27
5.7	The case of G_{14}	27
5.8	The case of G_{16}	28
5.9	The case of G_{17}	28
5.10	The case of G_{18}	29
5.11	The case of G_{20}	30
5.12	The case of G_{21}	30
5.13	The case of G_{24}	31
5.14	The case of G_{25}	32
5.15	The case of G_{26}	32
5.16	The case of G_{27}	33
5.17	The case of $G(m, 1, 2)$	34
5.18	The case of $G(m, 1, 3)$	34
6	A universal formula for the dual product	35
6.1	Remarks	37
6.2	An analogue of Veselov's potential	37
6.3	Generalized Saito flat coordinates: the standard case	38
7	Bi-flat F-manifold structure on the space of orbits of groups revisited	38
7.1	The case of G_{26}	39
7.2	The case of G_5	41
7.3	The case of G_6	41
7.4	The case of G_9	41

7.5	The case of G_{10}	42
7.6	The case of G_{14}	42
7.7	The case of G_{17}	43
7.8	The case of G_{18}	43
7.9	The case of G_{21}	43
7.10	The case of $G(3, 1, 2)$	44
7.11	The case of $G(3, 1, 3)$	44
7.12	The bi-flat F -structures of associated Coxeter groups	46
8	Bi-flat F-manifolds and generalized WDVV equations	48
9	Appendix 1. Reflecting mirrors	50
9.1	The G_4 case	50
9.2	The G_5 case	51
9.3	The G_6 case	51
9.4	The G_8 case	51
9.5	The G_9 case	52
9.6	The G_{10} case	52
9.7	The G_{14} case	53
9.8	The G_{16} case	54
9.9	The G_{17} case	55
9.10	The G_{18} case	56
9.11	The G_{20} case	57
9.12	The G_{21} case	58
9.13	The G_{23} case	59
9.14	The G_{24} case	60
9.15	The G_{25} case	61
9.16	The G_{26} case	61
9.17	The G_{27} case	62
9.18	The $G(3, 1, 2)$ case	64
9.19	The $G(3, 1, 3)$ case	64
9.20	The case of $I_2(6)$	65
9.21	The case of $I_2(8)$	65
9.22	The case of $I_2(10)$	65
9.23	The case of B_2	66
9.24	The case of B_3	66
10	Appendix 2. Saito coordinates in the cases of the groups G_{29}, G_{32}, G_{33}	66
10.1	The case of G_{32}	67
10.2	The case of G_{33}	68

1 Introduction

A complex (pseudo)-reflection is a unitary transformation of \mathbb{C}^n of finite period that leaves invariant a hyperplane. It is characterized by the property that all the eigenvalues of the associated matrix representation are equal to 1, except for one. The remaining eigenvalue is a k -th primitive root of unity, where k is the period of the transformation.

A finite subgroup of the group of unitary transformations is a finite complex reflection group if it is generated by complex reflections. Irreducible finite complex reflection groups were classified by Shephard and Todd in [44], and consist in an infinite family depending on 3 positive integers and 34 exceptional cases. They proved that the ring of invariant polynomials of a complex reflection group is generated by n algebraically independent invariant polynomials, where n is the dimension of the complex vector space on which the group acts. Well-generated irreducible complex reflection groups are irreducible complex reflection groups of rank n , whose minimal generating set consists of n reflections. In the sequel we will restrict our attention only to well-generated complex reflection groups, even if it is not explicitly stated.

In this paper, we consider two affine flat connections naturally associated with an irreducible well-generated complex reflection group G :

1. The first one is defined by the Saito coordinates, which correspond to a special choice of basic invariant polynomials. They were originally introduced for Coxeter groups in [40, 41] and play a crucial role in the construction of the Frobenius manifolds associated with Coxeter groups [10] (see also [42] for the case of singularity theory). The notion of Saito flat coordinates (also called flat basic invariants) was extended to Shephard groups in [36] (see also [37]). Recently, the existence of a system of flat basic invariants for well-generated complex reflection groups has been proved in [20].
2. The second one is a flat torsionless logarithmic connection associated with the arrangement of hyperplanes H fixed by some elements of G , given by

$$\tilde{\nabla} = \nabla - \frac{1}{N} \sum_{H \in \mathcal{H}} \frac{d\alpha_H}{\alpha_H} \otimes \kappa_H \pi_H, \quad (1.1)$$

where ∇ is the standard flat connection on \mathbb{C}^n , α_H is a linear form defining the mirror H , \mathcal{H} is the collection of the mirrors H , π_H denotes the unitary

projection onto the unitary complement of the hyperplane H , computed with respect to a suitable Hermitian metric, the weights κ_H are suitable complex numbers and N is a normalizing factor chosen in such a way that

$$\sum_{H \in \mathcal{H}} \kappa_H \pi_H = \text{Id}_{\mathbb{C}^n}. \quad (1.2)$$

The above connection is a particular case of *Dunkl-Kohno-type* connections considered in [7] (see also similar structures considered in [5] and [12]). If the collection $\{\kappa_H\}_{H \in \mathcal{H}}$ of complex weights is G -invariant, then the connection defined in (1.1) is flat (see [26]). In a more general setting, different flatness conditions for this kind of connections were first studied by Kohno (see [23, 24]). In the case of Coxeter groups, these Dunkl-Kohno-type connections coincide with the flat connection appearing in the theory of \vee -system [45, 46]. In this case Kohno's flatness conditions are equivalent to Veselov's definition of \vee -system [3, 15]. Important examples of these logarithmic connections appear also in the Physics literature and are called KZ connections [22].

The aim of this paper is to show how the two flat structures above are related.

The main results of the paper are the following.

1. We show that the two affine flat connections described above endow the space of orbits of well-generated irreducible complex reflection groups with the structure of a bi-flat F -manifold. The notion of bi-flat F manifold has been introduced in [2] as a natural generalization of Frobenius manifolds in the framework of integrable dispersionless PDEs. It was later showed that tridimensional semisimple bi-flat F are parameterized by solutions of generic Painlevé VI equation [28] (see also [2]) and, in the non-semisimple (regular) case, by solutions of generic Painlevé IV and Painlevé V equations, depending on the Jordan canonical form of the linear endomorphism given by multiplication by the Euler vector field [4]. We believe that these results combined with the results of the present paper unveil the relevance of the notion of bi-flat F -manifold as a meaningful generalization of the notion of Frobenius manifold.

Due to computational difficulties we focus on the rank 2 and rank 3 cases. However we believe that our results hold true also for higher rank groups.

2. We prove that for the Shephard groups $G_5, G_6, G_9, G_{10}, G_{14}, G_{17}, G_{18}, G_{21}, G_{26}$ the bi-flat F -structure is not uniquely defined but depends on a parameter. For each of these groups, there is a single value of the parameter that corresponds to the associated Frobenius manifold structure. In other words, the

Frobenius structure of each of these orbit spaces is naturally embedded in a one-parameter family of bi-flat F -manifold structures, which can be viewed as a kind of deformation of the Frobenius structure itself.

3. It is a well-known result that the Frobenius structure of each Shephard group coincides with the Frobenius structure of the underlying Coxeter group ([9]) as a consequence of Hertling's theorem [16]. In complete analogy, we prove that on the orbit spaces of all Shephard groups mentioned above, the bi-flat F -manifold structure coincide with the bi-flat F -structure of the orbit space of the underlying Coxeter group. More precisely:
 - the one parameter families of bi-flat F -structures on the orbit space of the groups G_5, G_{10}, G_{18} coincide with the one parameter family of bi-flat F -structures on the orbit space of B_2 .
 - the one parameter families of bi-flat F -structures on the orbit space of the groups G_6, G_9, G_{17} coincide with the one parameter family of bi-flat F -structures on the orbit space of $I_2(6)$.
 - the one parameter family of bi-flat F -structures on the orbit space of G_{14} coincides with the one parameter family of bi-flat F -structures on the orbit space of $I_2(8)$.
 - the one parameter family of bi-flat F -structures on the orbit space of G_{21} coincides with the one parameter family of bi-flat F -structures on the orbit space of $I_2(10)$.
 - the one parameter family of bi-flat F -structures on the orbit space of G_{26} coincides with the one parameter family of bi-flat F -structures on the orbit space of B_3 .

In all these cases, for specific values of the parameter we recover the standard Frobenius manifold structure on the space of orbits of Coxeter groups. This fact leads us to introduce the notion of *generalized Saito flat coordinates*. Remarkably, there is a standard choice of the bi-flat structure on the orbit space of well-generated complex reflection groups such that the structure constants of the dual product and the Christoffel symbols of the dual connection have the following form:

$$c_{jk}^{*i}(p) = -\Gamma_{jk}^{*i}(p) = \frac{1}{d_l - 1} \frac{\partial^2 u^l}{\partial p^j \partial p^k} (J^{-1})_l^i, \quad (1.3)$$

where $\{p^1, \dots, p^n\}$ are standard coordinates on \mathbb{C}^n , $\{u^1, \dots, u^n\}$ are the set of basic invariant polynomials defining the generalized Saito flat coordinates, $J_j^i = \frac{\partial u^i}{\partial p^j}$ and d_i are the degrees of the invariant polynomials.

4. Inspired by Veselov's \vee -system idea, we prove that in all rank 2 and 3 exceptional well-generated complex reflection groups, the dual connection can be expressed as a connection of Dunkl-Kohno-type built from the reflecting hyperplanes. As a byproduct, our construction provides also a generalization of \vee -systems to the case of complex roots in \mathbb{C}^n and to the case of *Hermitian metrics*. Indeed, while in the case of \vee -systems one uses a non-degenerate symmetric bilinear form (either in \mathbb{R}^n or in \mathbb{C}^n , see [15]), in our case, to identify the relevant connection as a connection of Dunkl-Kohno-type, one is forced to use a non-degenerate sesquilinear form (in all cases we considered it is just the standard Hermitian form on \mathbb{C}^n , except in the case of G_{27}). Remarkably, in all the examples the dual product has the form

$$X * Y_p = \frac{1}{N} \left(\sum_{s=1}^n \frac{\kappa_s}{\|\alpha_s\|^2} \frac{\alpha_s(X) \alpha_s(Y) \check{\alpha}_s}{\alpha_s(p)} \right) \quad (1.4)$$

where κ_s is the order of the corresponding (pseudo)-reflection, X and Y are arbitrary vector fields, α_s are the covectors defining the hyperplanes and $\check{\alpha}_s$ are vectors obtained from α_s using a suitable Hermitian metric. Combining the above formula with the formula (1.3) we obtain the following system of PDEs for the standard generalized Saito flat coordinates:

$$\frac{\partial^2 u^i}{\partial p^j \partial p^k} = (d_i - 1) c_{jk}^{*s} \frac{\partial u^i}{\partial p^s}, \quad (1.5)$$

where the dual product is given by the formula (1.4). In other words *each Saito flat coordinate u^i is also a flat coordinate of the flat connection $\nabla^{(0)} + (d_i - 1)*$* . Using conditions (1.5) one can easily find standard flat basic invariants starting from general basic invariants. A similar formula holds true also in the non-standard case. As far as we know, many of the explicit formulas obtained in this paper for flat basic invariants have never appeared in literature before.

As it is often the case in the mathematical literature concerning well-generated complex reflection groups, our results have been obtained via a case by case analysis for each group. This is consistent with the fact that very often there does not seem to exist an overarching strategy to tackle properties of well-generated complex reflection groups (see for instance the comments in [38]).

In our case this is also due to the fact that while we present general statements, there are always some "misbehaved" groups. For instance, when we prove that the dual connection can be expressed as a connection of Dunkl-Kohno type using a sort of generalization of Veselov's \vee -systems, the Hermitian metric appearing

in the construction is always the standard one, except for G_{27} (see Table 3). Similarly, the (generalized) flat Saito coordinates are in some case uniquely determined, while in other cases they appear in families depending on a parameter. These facts suggest that it might be unlikely to find general a priori proofs of most of the results appearing in this work.

The paper is organized as follows. In Section 2, we introduce the notion of Frobenius structures and their almost dual and we recall the Dubrovin-Saito construction on the space of orbits of Coxeter groups, and Dubrovin's construction on the space of orbits of Shephard groups. In Section 3, we introduce Veselov's \vee -systems focusing our attention on the case of Coxeter groups. In Section 4, we briefly recall the definition of flat and bi-flat F -manifolds and we show how the bi-flat structure arises naturally when one tries to extend Dubrovin's almost duality to Frobenius manifolds "without metric". In Section 5 we show how to endow with a bi-flat F -structure the space of orbits of well-generated complex reflection groups. In Section 6, we show that for all rank 2 and 3 exceptional well-generated irreducible complex reflection groups, the standard dual connection coincides with the Dunkl-Kohno connection obtained choosing as weights k_H the orders of the corresponding (pseudo)-reflections.

In general, in the case of Shephard groups the standard bi-flat F -structure does not coincide with the structure of Frobenius manifolds introduced by Dubrovin. In Section 7 we explain how to recover the Dubrovin's structure considering a family of deformed Saito flat coordinates (generalized Saito Coordinates). It turns out that, also in this case, the dual connection is of Dunkl-Kohno type but with a different choice of the weights. In the final Section 8, we show that semisimple bi-flat F -manifolds are parametrized in flat coordinates by solutions of generalized WDVV equations. These equations are the oriented associativity equations [29] with an additional homogeneity condition. They have been recently introduced in [19] in the study of Saito structures without metrics [39]. These structures are completely characterized by a flat meromorphic connection on the bundle π^*TM on $\mathbb{P} \times M$. The result of Section 8 implies that this connection plays the same role in the theory of semisimple bi-flat F -manifolds as the role played by Dubrovin's extended connection in the theory of Frobenius manifolds. It also implies that one can construct a bi-flat structure on the orbit space of a well-generated complex reflection group, starting from any solution of generalized WDVV equations. The paper ends with three Appendices. In the first one we list all the reflecting hyperplanes for all exceptional well-generated complex reflection groups up to G_{27} , in the second one, using our procedure, we provide (conjectural) Saito flat coordinates for G_{29} , G_{32} and G_{33} . In the last one we briefly discuss what happens in the non-well-generated cases.

Acknowledgements

We would like to thank Vic Reiner for having suggested reference [25] and Alexander Veselov for reference [14] and for his interest in this work. This research was partially supported by GNFM through the 2015 Visitors Program.

2 Frobenius manifolds and their almost dual

Originally introduced by Dubrovin as a geometric framework to study the WDVV equations of topological field theory, Frobenius manifolds play nowadays an important role in several areas of mathematics including singularity theory, quantum cohomology and integrable systems. Let us recall their definition.

Definition 2.1 *A Frobenius manifold (M, \circ, η, e, E) is a manifold equipped with an associative commutative product \circ on sections of its tangent bundle, two distinguished vector fields e (unit) and E (Euler vector field) and a flat pseudo-metric η satisfying the following requirements:*

- *η is invariant with respect to the product: $\eta_{il}c_{jk}^l = \eta_{jl}c_{ik}^l$, where c_{jk}^i are the structure constants for \circ .*
- *the Levi-Civita connection ∇ associated to η is compatible with the product:*

$$\nabla_k c_{jl}^i = \nabla_j c_{kl}^i.$$

- *e is the unit of the product and it is flat: $\nabla e = 0$.*
- *Furthermore, the following conditions must hold:*

$$\nabla \nabla E = 0, \quad [e, E] = e, \quad \text{Lie}_E c_{jk}^i = c_{jk}^i, \quad \text{Lie}_E \eta = D\eta,$$

where D is a constant.

The product is called *semisimple* if there exist a distinguished coordinates system such that $c_{jk}^i = \delta_j^i \delta_k^i$.

The existence of the Euler vector field allows one to define a second commutative associative product on sections of the tangent bundle, called *the dual product* and defined as

$$X * Y := E^{-1} \circ X \circ Y, \tag{2.1}$$

and a second contravariant flat metric g , called the intersection form, and defined as $g^{ij} = \eta^{il} c_{lk}^j E^k$.

Furthermore, the very definition of Frobenius manifold is constructed in such a way that in flat coordinates $\{u_1, \dots, u_n\}$ for η , there exists a smooth function $F(u_1, \dots, u_n)$ such that $c_{jk}^i = \eta^{il} \frac{\partial^3 F}{\partial u_l \partial u_j \partial u_k}$. This function is called *Frobenius potential*.

In the next Section we will study in details the case of Frobenius manifolds associated with Coxeter groups.

2.1 Frobenius manifolds and space of orbits of Coxeter groups

One of the main examples of Frobenius manifold is provided by the space of orbits of Coxeter groups [10]. The realization of Frobenius manifold structure on these spaces is based on the notion of *flat pencil of metrics* and relies on the existence of a distinguished set of basic invariants of the group, called *Saito flat coordinates* [41].

A flat pencil of contravariant metrics is a one-parameter family of contravariant metrics $g_\lambda = g - \lambda\eta$ satisfying the following two properties [11]:

- g_λ is flat for any λ ,
- the pencil of Christoffel symbols coincides with the Christoffel symbols of the pencil.

In the case of Frobenius manifolds, it is well known that the inverse η^{ij} of the invariant (pseudo)-metric η_{ij} and the intersection form g^{ij} satisfy the above conditions (see [11] for details). They satisfy also two additional properties called *quasi-homogeneity* and *exactness*:

- Quasi-homogeneity implies the existence of a constant c such that $\text{Lie}_E \eta = c\eta$ (here and below Lie_X denotes the Lie derivative with respect to a vector field X).
- Exactness means that there exists a vector field (the unit vector field e in the Frobenius case) such that

$$\text{Lie}_e g = \eta, \quad \text{Lie}_e \eta = 0. \quad (2.2)$$

Since the unit vector field is flat (i.e. $\nabla e = 0$), one can choose flat coordinates $\{u^1, \dots, u^n\}$ such that $e = \frac{\partial}{\partial u^1}$. In such coordinates one obtains η just shifting g along the variable u^1 .

Conversely, given an exact quasi-homogeneous flat pencil of metrics, one can reconstruct a Frobenius manifold. In particular the Frobenius potential $F(u_1, \dots, u_n)$ is obtained in flat coordinates (u^1, \dots, u^n) for η (the Saito flat coordinates) by means of the following formula [10]

$$\eta^{il} \eta^{jm} \frac{\partial^2 F}{\partial u^l \partial u^m} = \frac{g^{ij}}{\deg(g^{ij})} \quad (2.3)$$

where $\deg(g^{ij}) = \frac{E(g^{ij})}{g^{ij}}$.

Given a finite Coxeter group, the crucial observation that allowed Dubrovin to define an exact flat pencil of metrics is that, due to the properties of the degrees of basic invariant polynomials, the Euclidean metric g , written with respect to a set of basic invariants, depends linearly on the invariant polynomial of highest degree. This implies that the metrics g and $\eta := \text{Lie}_e g$ (where e is the generator of the shift along the highest invariant polynomial) form an exact flat pencil of metrics. Moreover, using the homogeneity properties of basic invariant polynomials, one can easily prove that this pencil of metrics is quasi-homogeneous. Starting from this pencil one can easily reconstruct the full Forbenius structure using the formula (2.3). In order to illustrate Dubrovin's construction we consider two simple examples.

2.1.1 The case of A_3

A_3 is a finite Coxeter group that can be realized as a real reflection group. Any real reflection group gives rise to a well-generated complex reflection group. A_3 can be viewed as a well-generated complex reflection group of rank three. Since it also coincides with the symmetric group on 4 letters $\text{Sym}(4)$ (with a suitable representation) its invariants are constructed starting from \mathbb{C}^4 , considering polynomials in p_0, p_1, p_2, p_3 that are invariants under $\text{Sym}(4)$ and then restricting them to the hyperplane given by the equation $p_0 + p_1 + p_2 + p_3 = 0$. A general basis for the ring of invariants, restricted already to the hyperplane $p_0 + p_1 + p_2 + p_3 = 0$ is provided by the following polynomials of degree 2, 3, 4, where an arbitrary constant c has been introduced:

$$\begin{aligned} u_1 &= -p_1^2 - p_2^2 - p_3^2 - p_1p_2 - p_1p_3 - p_2p_3, \\ u_2 &= p_1^2(p_2 + p_3) + p_2^2(p_1 + p_3) + p_3^2(p_1 + p_2) + 2p_1p_2p_3, \\ u_3 &= -p_1^2p_2p_3 - p_1p_2^2p_3 - p_1p_2p_3^2 - cu_1^2. \end{aligned}$$

The Euclidean covariant metric $dp_0^2 + dp_1^2 + dp_2^2 + dp_3^2$ restricted to the hyperplane $p_0 + p_1 + p_2 + p_3 = 0$ is expressed via the following matrix (in the coordinates p_1, p_2, p_3):

$$g = \begin{bmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{bmatrix},$$

and inverting it, one gets the contravariant metric

$$g^{-1} = \begin{bmatrix} \frac{3}{4} & -\frac{1}{4} & -\frac{1}{4} \\ -\frac{1}{4} & \frac{3}{4} & -\frac{1}{4} \\ -\frac{1}{4} & -\frac{1}{4} & \frac{3}{4} \end{bmatrix}.$$

Using the tensorial properties of the metric, we can express the metric g in the coordinates u_1, u_2, u_3 as follows:

$$g = \begin{bmatrix} -2u_1 & -3u_2 & -4u_3 \\ -3u_2 & -4u_3 + (1-4c)u_1^2 & (6c + \frac{1}{2})u_1u_2 \\ -4u_3 & (6c + \frac{1}{2})u_1u_2 & \frac{3}{4}u_2^2 + (16c-2)u_1u_3 + (8c^2-2c)u_1^3 \end{bmatrix}.$$

In particular in the Saito flat coordinates, which correspond to the value $c = \frac{1}{8}$, we obtain

$$g = \begin{pmatrix} -2u_1 & -3u_2 & -4u_3 \\ -3u_2 & -4u_3 + \frac{1}{2}u_1^2 & \frac{5}{4}u_1u_2 \\ -4u_3 & \frac{5}{4}u_1u_2 & \frac{3}{4}u_2^2 - \frac{1}{8}u_1^3 \end{pmatrix}.$$

Up to an inessential constant factor the metric η is given by

$$\eta = \text{Lie}_e g = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix},$$

and, solving the equations (2.3) one obtains the following Frobenius potential:

$$F = \frac{1}{8}u_3^2u_1 + \frac{1}{8}u_3u_2^2 - \frac{1}{64}u_1^2u_2^2 + \frac{1}{3840}u_1^5. \quad (2.4)$$

2.1.2 The case of H_3 (G_{23})

A second example is the Coxeter group H_3 . It corresponds to the group G_{23} of the Shephard-Todd list. The basic invariants are given by ([33, 18])

$$\begin{aligned} U_1 &= p_1^2 + p_2^2 + p_3^2 \\ U_2 &= 2(p_1^6 + p_2^6 + p_3^6) - 15(p_1^4p_2^2 + p_1^4p_3^2 + p_2^4p_1^2 + p_2^4p_3^2 + p_3^4p_1^2 + p_3^4p_2^2) + 180p_1^2p_2^2p_3^2 \\ &\quad + 21\sqrt{5}\Delta \\ U_3 &= 5(p_1^{10} + p_2^{10} + p_3^{10}) - 45(p_1^8p_2^2 + p_1^8p_3^2 + p_2^8p_1^2 + p_2^8p_3^2 + p_3^8p_1^2 + p_3^8p_2^2) \\ &\quad + 42(p_1^6p_2^4 + p_1^6p_3^4 + p_2^6p_1^4 + p_2^6p_3^4 + p_3^6p_1^4 + p_3^6p_2^4) \\ &\quad + 1008(p_1^6p_2^2p_3^2 + p_2^6p_1^2p_3^2 + p_3^6p_1^2p_2^2) - 1260(p_1^4p_2^4p_3^2 + p_1^4p_3^4p_2^2 + p_2^4p_3^4p_1^2) + \\ &\quad - 33\sqrt{5}\Delta(3(p_1^4 + p_2^4 + p_3^4) - 11(p_1^2p_2^2 + p_1^2p_3^2 + p_2^2p_3^2)). \end{aligned}$$

where $\Delta = (p_1^2 - p_2^2)(p_1^2 - p_3^2)(p_2^2 - p_3^2)$. It turns out that Saito flat coordinates in this case are provided by the following polynomials:

$$u_1 = U_1, \quad u_2 = U_2 + c_1 U_1^3, \quad u_3 = U_3 + c_2 U_1^2 U_2 + c_3 U_1^5,$$

where $c_1 = -2$, $c_2 = \frac{33}{7}$ and $c_3 = -\frac{2091}{175}$.

The Euclidean contravariant metric written in the Saito flat coordinates is given by

$$g = \begin{pmatrix} 4u_1 & 12u_2 & 20u_3 \\ 12u_2 & -\frac{3528}{187}u_3 - 168u_2u_1^2 + \frac{3528}{25}u_1^5 & \frac{748}{3}u_2^2u_1 - \frac{5236}{5}u_2u_1^4 \\ 20u_3 & \frac{748}{3}u_2^2u_1 - \frac{5236}{5}u_2u_1^4 & -\frac{349690}{3087}u_2^3 + \frac{139876}{49}u_2^2u_1^3 + \frac{139876}{25}u_1^9 \end{pmatrix},$$

and the metric $\eta = \text{Lie}_e g$ is given by

$$\eta = \begin{pmatrix} 0 & 0 & 20 \\ 0 & -\frac{3528}{187} & 0 \\ 20 & 0 & 0 \end{pmatrix}.$$

Solving the equations (2.3) one obtains the Frobenius potential for this case, namely

$$F = \frac{1}{400}u_1u_3^2 - \frac{187}{70560}u_3u_2^2 + \frac{34969}{1764000}u_2^2u_1^5 - \frac{34969}{4445280}u_3^3u_1^2. \quad (2.5)$$

2.2 Frobenius manifolds and space of orbits of Shephard groups

A Shephard group is the symmetry group of a regular complex polytope [43]. The space of orbits of a Shephard group is endowed with the structure of a Frobenius manifold. The definition of this structure was given in [9]. The crucial observation in this case is that, due to the results of [35], the inverse of the Hessian of the basic invariant polynomial of lowest degree defines a flat (pseudo)-metric which depends linearly on the highest degree polynomial (in a coordinate system given by basic invariant polynomials). Using this fact, one can use the same arguments that work in the case of Coxeter groups. In particular, in the case of Coxeter groups, the lowest degree basic invariant polynomial is the sum of the squares of the coordinates and its Hessian coincides with the standard Euclidean metric. In this way one recovers the standard construction presented in the examples above. We present as an example the case of G_{26} , which is a Shephard group.

2.2.1 The case of G_{26}

Starting from the invariant polynomial of lowest degree

$$u_1 = p_1^6 - 10p_1^3p_2^3 - 10p_1^3p_3^3 + p_2^6 - 10p_2^3p_3^3 + p_3^6,$$

one can define the covariant (pseudo)-metric given by the Hessian:

$$\frac{\partial^2 u_1}{\partial p^i \partial p^j} = \begin{pmatrix} 30p_1^4 - 60p_1p_2^3 - 60p_1p_3^3 & -90p_1^2p_2^2 & -90p_1^2p_3^2 \\ -90p_1^2p_2^2 & -60p_1^3p_2 + 30p_2^4 - 60p_2p_3^3 & -90p_2^2p_3^2 \\ -90p_1^2p_3^2 & -90p_2^2p_3^2 & -60p_1^3p_3 - 60p_2^3p_3 + 30p_3^4 \end{pmatrix}.$$

The inverse of this metric in the Saito coordinates (u_1, u_2, u_3) with u_2 and u_3 defined as

$$\begin{aligned} u_2 &= (p_1^3 + p_2^3 + p_3^3)(216p_1^3p_2^3p_3^3 + (p_1^3 + p_2^3 + p_3^3)^3) - \frac{1}{4}u_1^2, \\ u_3 &= (p_1^3 + p_2^3 + p_3^3)^6 - 540p_1^3p_2^3p_3^3(p_1^3 + p_2^3 + p_3^3)^3 - 5832p_1^6p_2^6p_3^6 - \frac{3}{4}u_1u_2 + \frac{3}{16}u_1^3. \end{aligned}$$

reads

$$g = \begin{pmatrix} \frac{6}{5}u_1 & \frac{9}{5}u_2 & \frac{12}{5}u_3 \\ \frac{9}{5}u_2 & -\frac{1}{160}u_3 + \frac{3}{1280}u_1^2 & -\frac{9}{4}u_1u_2 \\ \frac{12}{5}u_3 & -\frac{9}{4}u_1u_2 & \frac{27}{40}u_1^3 + \frac{2592}{5}u_2^2 \end{pmatrix}.$$

Notice that g does indeed depend linearly on u_3 , the highest degree invariant. Then defining $\eta := \text{Lie}_e g$ (where e in this case is the vector field $\frac{\partial}{\partial u_3}$), it is easy to check that the pencil

$$g - \lambda\eta = \begin{pmatrix} \frac{6}{5}u_1 & \frac{9}{5}u_2 & \frac{12}{5}u_3 \\ \frac{9}{5}u_2 & -\frac{1}{160}u_3 + \frac{3}{1280}u_1^2 & -\frac{9}{4}u_1u_2 \\ \frac{12}{5}u_3 & -\frac{9}{4}u_1u_2 & \frac{27}{40}u_1^3 + \frac{2592}{5}u_2^2 \end{pmatrix} - \lambda \begin{pmatrix} 0 & 0 & \frac{12}{5} \\ 0 & -\frac{1}{160} & 0 \\ \frac{12}{5} & 0 & 0 \end{pmatrix},$$

is an exact flat pencil of metrics. From the formula (2.3) one easily obtains the Frobenius potential

$$F = \frac{1}{96768}u_1^7 + \frac{5}{3456}u_2^2u_1^3 + \frac{1}{290304}(840u_2^3 + 2240u_3^2)u_1 + \frac{5}{432}u_3u_2^2.$$

2.2.2 Shephard groups of rank 2 and 3

Applying Dubrovin's procedure one can easily reconstruct the Frobenius potential for each Shephard group. Below we present the list of the Frobenius potential for *all* Shephard groups of rank 2 and 3, not just for the exceptional ones.

Table 1: Frobenius potential for all Shephard groups of rank 2 and 3.

Type	Potential
G_4	$F = \frac{1}{24}u_1u_2^2 - \frac{i}{4608}\sqrt{3}u_1^4$
G_5	$F = -\frac{1}{4}u_1^5 + \frac{5}{288}u_1u_2^2$
G_6	$F = \frac{1}{96}u_2^2u_1 - \frac{1}{1935360}u_1^7$

G_8	$F = \frac{7}{288}u_2^2u_1 + \frac{7}{1536}u_1^4$
G_9	$F = \frac{7}{1152}u_1u_2^2 + \frac{1}{7680}u_1^7$
G_{10}	$F = \frac{11}{1152}u_1u_2^2 + \frac{11}{34560}u_1^5$
G_{14}	$F = \frac{5}{1152}u_1u_2^2 + \frac{45}{56}u_1^9$
G_{16}	$F = -\frac{19}{2880000}\sqrt{5}u_1^4 + \frac{19}{1800}u_2^2u_1$
G_{17}	$F = \frac{19}{7200}u_2^2u_1 + \frac{19}{6048000000}u_1^7$
G_{18}	$F = \frac{29}{7200}u_1u_2^2 + \frac{29}{12}u_1^5$
G_{20}	$F = \frac{11}{1800}u_2^2u_1 + \frac{11}{5184000}\sqrt{5}u_1^6$
G_{21}	$F = \frac{11}{7200}u_1u_2^2 + \frac{1}{933120000}u_1^{11}$
G_{23}	$F = \frac{1}{400}u_1u_3^2 - \frac{187}{70560}u_3u_2^2 + \frac{34969}{1764000}u_2^2u_1^5 - \frac{34969}{4445280}u_3^3u_1^2$
G_{25}	$F = \frac{5}{288}u_1u_3^2 - \frac{20}{3}u_3u_2^2 + \frac{5}{2}u_1^2u_2^2 + \frac{1}{3072}u_1^5$
G_{26}	$F = \frac{1}{96768}u_1^7 + \frac{5}{3456}u_2^2u_1^3 + \frac{1}{290304}(840u_2^3 + 2240u_3^2)u_1 + \frac{5}{432}u_3u_2^2$
$G(m, 1, 2)$	$F = \frac{1}{24}u_1u_2^2 + \frac{1}{2880}u_1^5$
$G(m, 1, 3)$	$F = \frac{1}{54}u_1u_3^2 + \frac{1}{72}u_2^2u_3 + \frac{1}{2592}u_1u_2^2(u_1^2 + 3u_2) + \frac{1}{816480}u_1^7$

Recall that any Shephard group G admits a presentation of the form [8]

$$s_i^{p_i} = 1$$

$$s_i s_j s_i \cdots = s_j s_i s_j \cdots$$

where s_i are the generating (pseudo)-reflections and p_i are integers ≥ 2 . The Coxeter group obtained replacing p_i with 2 in the above presentation is usually called the Coxeter group associated with G . Due to the Hertling's theorem [16], since the Frobenius manifold structure on the space of orbits of a Shephard group is polynomial, it must be isomorphic to the Frobenius manifold structure on the space of orbits of a Coxeter group. For each Shephard group G , this Coxeter group is exactly the Coxeter group associated with G [9]. This means that the above Frobenius potentials coincide up to rescaling of the variables with the potentials of the associated Coxeter group. For instance, the Frobenius potentials in the last two examples ($G_{(m,1,2)}$ and $G_{(m,1,3)}$) do not depend on m and thus coincide with the Frobenius potentials associated with the Coxeter groups B_2 and B_3 , respectively.

3 \vee -systems

\vee -systems were introduced by A. Veselov in [45] to construct new solutions of WDVV associativity equations, starting from a special set of covectors. The con-

ditions defining a \vee -system are precisely the conditions that guarantee that a particular function constructed from a special set of covectors satisfies the WDVV equations.

Now we recall the notion of \vee -systems (see [45]). Let V be a finite dimensional real vector space (the notion has been generalized also to complex vector spaces in [15]), and denote with \mathcal{V} a finite set of non-collinear covectors $\alpha \in V^*$. We assume that the symmetric bilinear form defined by $g := \sum_{\alpha \in \mathcal{V}} \alpha \otimes \alpha$ is non-degenerate. We will denote by $\check{\alpha}$ the vector uniquely defined by: $(\check{\alpha}, \cdot) = \alpha(\cdot)$ where (\cdot, \cdot) is the bilinear form defined by g .

Definition 3.1 *We say that \mathcal{V} is a \vee -system if for each two-dimensional plane $\Pi \subset V^*$ we have*

$$\sum_{\beta \in \Pi \cap \mathcal{V}} \beta(\check{\alpha})\check{\beta} = \mu\check{\alpha}, \quad (3.1)$$

for each $\alpha \in \Pi \cap \mathcal{V}$ and for some μ , which may depend on Π and α .

Let \mathcal{H} be the family of hyperplanes associated with the covectors contained in \mathcal{V} . By definition, $H \in \mathcal{H}$ if and only if $\ker(\alpha) = H$ for one of the covectors in \mathcal{V} . We denote by α_H a covector associated to H and by $\pi_H : V \rightarrow H^\perp$ is the linear map having as kernel H and range H^\perp , the orthogonal complement of H in V , where the orthogonality is defined via g . Due to the results of [3, 15] the definition of \vee -system is equivalent to the requirement that the one parameter family of connections

$$\nabla - \lambda \sum_{H \in \mathcal{H}} \frac{d\alpha_H}{\alpha_H} \otimes \pi_H, \quad (3.2)$$

(where ∇ is the trivial affine structure defined by the coordinates $\{p^1, \dots, p^n\}$) is flat for any value of the parameter λ .

Given a set of covectors defining a \vee -system the corresponding solution of WDVV associativity equations is given by the formula:

$$F(p) := \frac{1}{2} \sum_{H \in \mathcal{H}} (\alpha_H(p))^2 \log \alpha_H(p). \quad (3.3)$$

One of the main example of \vee -systems is given by Coxeter systems [45]. It was proved in [9] that Veselov's solutions of WDVV equations constructed from a Coxeter group define the almost dual structure of the Frobenius manifolds defined on the space of orbits of these groups. Notice that the dual product has the form

$$* = \sum_{H \in \mathcal{H}} \frac{d\alpha_H}{\alpha_H} \otimes \pi_H. \quad (3.4)$$

Similarly, we will see in Section 6 that for all exceptional well-generated complex reflection groups of rank 2 and 3 the dual connection (and the dual product) of a

bi-flat F -manifold constructed on the relevant orbit spaces can be expressed using a Dunkl-Kohno-type connection. This connection is built using a generalization of the notion of \vee -system.

3.1 An example: A_3

The \vee -system data that give rise to A_3 are the following covectors:

$$\begin{aligned}\alpha_1 &= [1, -1, 0], & \alpha_2 &= [1, 0, -1], & \alpha_3 &= [0, 1, -1] \\ \alpha_4 &= [2, 1, 1], & \alpha_5 &= [1, 2, 1], & \alpha_6 &= [1, 1, 2],\end{aligned}$$

and the associated projections:

$$\begin{aligned}\pi_1 &= \begin{pmatrix} 1 & -1 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & \pi_2 &= \begin{pmatrix} 1 & 0 & -1 \\ 0 & 0 & 0 \\ -1 & 0 & 1 \end{pmatrix}, & \pi_3 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & -1 & 1 \end{pmatrix}, \\ \pi_4 &= \begin{pmatrix} 2 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & \pi_5 &= \begin{pmatrix} 0 & 0 & 0 \\ 1 & 2 & 1 \\ 0 & 0 & 0 \end{pmatrix}, & \pi_6 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 1 & 2 \end{pmatrix}.\end{aligned}$$

4 Flat and bi-flat F -manifolds

F -manifolds with compatible flat structure (for short *flat F -manifolds*) have been introduced by Manin as a generalization of Frobenius manifolds. They are a particular instance of a more general class of manifolds, called F -manifolds, introduced by Hertling and Manin in [17]. Let us recall their definition.

Definition 4.1 *A flat F -manifold (M, \circ, ∇, e) is a manifold M equipped with the following data:*

1. *a commutative associate product $\circ : TM \times TM \rightarrow TM$ with flat unit e .*
2. *a flat torsionless affine connection ∇ compatible with the product:*

$$\nabla_k c_{jl}^i = \nabla_j c_{kl}^i,$$

where c_{jk}^i are the structure constants of \circ .

In flat coordinates for ∇ , as a consequence of the axioms defining a flat F -manifold, we have that the structure constants can be expressed via the second partial derivatives of a vector potential:

$$c_{jk}^i = \partial_j \partial_k A^i.$$

Furthermore, the vector potential A^i satisfies the following equations, where one can choose as unit vector field $e = \partial_1$:

$$\partial_j \partial_l A^i \partial_k \partial_m A^l = \partial_k \partial_l A^i \partial_k \partial_m A^l \quad (4.1)$$

$$\partial_1 \partial_i A^j = \delta_i^j. \quad (4.2)$$

They are called *oriented associativity equations* [29]. Flat F -manifolds share several properties with Frobenius manifolds. Dubrovin's deformed connections, Dubrovin's duality and Dubrovin's principal hierarchy are well defined also for these manifolds (for the last point see [27]). The missing data are the invariant metric η and the Euler vector field E . A natural generalization of Frobenius manifolds with Euler vector field can be obtained replacing the flat invariant metric η with a flat connection ∇ satisfying the usual conditions.

Definition 4.2 A Frobenius manifold without metric (M, \circ, ∇, e, E) is a manifold equipped with an associative commutative product \circ on sections of its tangent bundle, two distinguished vector fields e (unit) and E (Euler vector field) and a flat connection ∇ satisfying the following requirements:

- the connection ∇ is compatible with the product:

$$\nabla_k c_{jl}^i = \nabla_j c_{kl}^i.$$

- e is the unit of the product and it is flat: $\nabla e = 0$.
- Furthermore, the following conditions must hold:

$$\nabla \nabla E = 0, \quad [e, E] = e, \quad \text{Lie}_E c_{jk}^i = c_{jk}^i.$$

4.1 Almost duality and bi-flat F -manifolds

In the semisimple case, Dubrovin's almost dual structure can be extended to Frobenius manifolds without metric using the following data:

- a dual product $*$ defined in the usual way as $X * Y = E^{-1} \circ X \circ Y$ for any pair of vector fields X and Y .
- a dual connection $\nabla^{(2)}$ satisfying the following properties
 1. $\nabla^{(2)} E = 0$.
 2. $\nabla^{(2)}$ is compatible with $*$.

3. $\nabla^{(1)}$ and $\nabla^{(2)}$ are almost hydrodynamically equivalent [1]:

$$(d_{\nabla^{(1)}} - d_{\nabla^{(2)}})(X \circ) = 0,$$

for every vector field X , where d_{∇} denotes the exterior covariant derivative associated to connection ∇ .

Remark 4.3 *In the case of Frobenius manifolds the dual connection, in general, does coincide with the Levi-Civita connection of the intersection form. The difference between these two connections is proportional to the dual product.*

Theorem 4.4 *The dual connection defined above is uniquely defined in terms of $\nabla^{(1)}$. Moreover the flatness of the dual connection is equivalent to the linearity of the Euler vector field: $\nabla^{(1)}\nabla^{(1)}E = 0$.*

Proof. From the above definition it follows that, in canonical coordinates for \circ (the product compatible with $\nabla^{(1)}$), we have (see [1, 2]):

$$\begin{aligned} c_{jk}^i &= \delta_j^i \delta_k^i, & c_{jk}^{*i} &= \frac{1}{u^i} \delta_j^i \delta_k^i, \\ e &= \sum_k \partial_k, & E &= \sum_k u^k \partial_k, \\ \Gamma_{ij}^{(1)i} &= \Gamma_{ij}^{(2)i} = \Gamma_{ij}^i, & i &\neq j \end{aligned}$$

Moreover (see [2]):

$$\begin{aligned} \Gamma_{jk}^{(1)i} &:= 0, & \Gamma_{jk}^{(2)i} &:= 0, & \forall i \neq j \neq k \neq i, \\ \Gamma_{jj}^{(1)i} &:= -\Gamma_{ij}^{(1)i}, & \Gamma_{jj}^{(2)i} &:= -\frac{u^i}{u^j} \Gamma_{ij}^{(2)i}, & i \neq j, \\ \Gamma_{ii}^{(1)i} &:= -\sum_{l \neq i} \Gamma_{li}^{(1)i}, & \Gamma_{ii}^{(2)i} &:= -\sum_{l \neq i} \frac{u^l}{u^i} \Gamma_{li}^{(2)i} - \frac{1}{u^i}. \end{aligned} \tag{4.3}$$

This proves the first part of the theorem. Let us prove the second part. The conditions expressing zero curvature are given by:

$$\partial_j \Gamma_{ik}^i + \Gamma_{ij}^i \Gamma_{ik}^i - \Gamma_{ik}^i \Gamma_{kj}^k - \Gamma_{ij}^i \Gamma_{jk}^j = 0, \quad \text{if } i \neq k \neq j \neq i, \tag{4.4}$$

$$e^l \partial_l \Gamma_{ik}^i = 0, \tag{4.5}$$

$$E^l \partial_l \Gamma_{ik}^i = -\Gamma_{ik}^i. \tag{4.6}$$

The conditions (4.4) and (4.5) provide the flatness of the natural connection and thus are equivalent to oriented associativity equations (4.1) and (4.2). The flatness of the dual connection follows from the condition (4.6).

On the other hand, using the vanishing of the curvature for $\nabla^{(1)}$, after a straightforward computation, we obtain

$$(\nabla^{(1)}\nabla^{(1)}E)_{kj}^i = \partial_k\partial_j E^i + \Gamma_{jl}^i\partial_k E^l + \Gamma_{km}^i\partial_j E^m - \Gamma_{kj}^m\partial_m E^i + E^l\partial_l\Gamma_{kj}^i = 0. \quad (4.7)$$

In canonical coordinates, the condition (4.7) reads

$$\delta_k^i\delta_j^i + \Gamma_{jk}^i + E^l\partial_l\Gamma_{kj}^i = 0,$$

and one can easily check that it is equivalent to the condition (4.6). \blacksquare

Bi-flat F -manifolds are Frobenius manifolds without metric endowed with the almost dual structure introduced above. They have been introduced in [2] motivated by the theory of integrable systems of hydrodynamic type. The definition is as follows:

Definition 4.5 *A bi-flat F -manifold $(M, \nabla^{(1)}, \nabla^{(2)}, \circ, *, e, E)$ is a manifold M equipped with a pair of flat connections $\nabla^{(1)}$ and $\nabla^{(2)}$, a pair of products \circ and $*$ on sections of the tangent bundle TM and a pair of vector fields e and E satisfying the following axioms:*

- *E behaves like a Euler vector field: $[e, E] = e$, $\text{Lie}_E c_{jk}^i = c_{jk}^i$, where c_{jk}^i are the structure constants of \circ .*
- *the product \circ is commutative, associative and with unity e . Moreover $\nabla^{(1)}e = 0$.*
- *The product $*$ is commutative, associative and with unity E . It is defined as: $X*Y = E^{-1} \circ X \circ Y$, $\forall X, Y$ vector fields on M . Moreover $\nabla^{(2)}E = 0$.*
- *$\nabla^{(1)}$ is compatible with \circ and $\nabla^{(2)}$ is compatible with $*$ in the following sense:*

$$\nabla_X^{(l)} \circ_{(l)}(Y, Z) = \nabla_Y^{(l)} \circ_{(l)}(X, Z), \quad l = 1, 2,$$

*for all X, Y, Z are vector fields on M , where $\circ_{(1)} = \circ$ and $\circ_{(2)} = *$.*

- *$\nabla^{(1)}$ and $\nabla^{(2)}$ are almost hydrodynamically equivalent.*

The first connection $\nabla^{(1)}$ is often called *the natural connection*. Frobenius manifolds are bi-flat F -manifolds equipped with an invariant metric η .

Remark 4.6 *The manifold M in the above definition might be a real or complex manifold. In the latter case TM is intended as the holomorphic tangent bundle and all the geometric data are supposed to be holomorphic. In the present paper we will deal with complex manifolds.*

5 Bi-flat F -structure on the space of orbits of complex reflection groups

In this Section we present a procedure that allows us to reconstruct (in principle) the bi-flat F -manifold structure and the vector potential starting from a well-generated complex reflection group (recall that any finite real reflection group is automatically well-generated). Let us consider the pair (V, G) , where V is a complex vector space of complex dimension n and G is a well-generated complex reflection group. We identify V with \mathbb{C}^n with standard coordinates $\{p^1, \dots, p^n\}$. We also consider the coordinates given by a set of basic invariant polynomials $\{u^1, \dots, u^n\}$ in the ring of invariant polynomials $\mathbb{C}[V]^G$, which can be viewed as coordinates on the space of orbits \mathbb{C}^n/G . Since the algorithm is based on the interplay between the standard coordinates $\{p^1, \dots, p^n\}$ on \mathbb{C}^n and the coordinates $\{u^1, \dots, u^n\}$, we denote with a superscript tilde the components written in the coordinates $\{u^1, \dots, u^n\}$.

Although $\{p^1, \dots, p^n\}$ and $\{u^1, \dots, u^n\}$ are coordinates on different spaces, since the map $\phi : \mathbb{C}^n \rightarrow \mathbb{C}^n/G$ is a diffeomorphism on any fundamental domain of the action of G on \mathbb{C}^n and since we are working locally, we will treat $\{p^1, \dots, p^n\}$ and $\{u^1, \dots, u^n\}$ just as different coordinates systems. Notice also that the choice of basic invariant polynomials $\{u^1, \dots, u^n\}$ is not unique in general.

Our construction relies on the hypotheses that the dual connection $\nabla^{(2)}$ and the dual product $*$ are related by the formula

$$\nabla^{(2)} = \nabla^{(0)} + \lambda*, \quad (5.1)$$

where $\nabla^{(0)}$ is the standard affine flat connection on \mathbb{C}^n and λ is a suitable constant. Taking into account the condition $\nabla^{(2)}E = 0$ and the form of the Euler vector field E , it is immediate to prove that $\lambda = -1$ (see remark 5.4 below).

The hypotheses above were chosen because this is exactly what happens in many examples of \vee -systems related to Coxeter groups, which constitute our inspiring model. The fact that with these assumptions one indeed can equip the orbit space with a bi-flat F -manifold structure is then verified a posteriori.

In this Section we explain how to define the dual connection and the dual product under the assumption (5.1). We assume that the flat coordinates for the natural connection we are looking for are basic invariants for the complex reflection groups and we call them *generalized Saito flat coordinates*. Combining this assumption with the assumption (5.1) we obtain an explicit formula for the Christoffel symbols of the dual connection and an explicit expression for the structure constants of the dual product, in terms of flat basic invariants. In practice, to select the generalized

Saito flat coordinates among basic invariants, we have to impose that the dual connection defined by this formula is flat and that the natural connection associated to generalized Saito flat coordinates is compatible with the natural product. We then show that, in all the examples, these two conditions fix uniquely the natural connection. We will see later in Section 7 that dropping the hypothesis (5.1) the natural connection and the associated generalized Saito flat coordinates in general are no longer uniquely determined.

Before explaining the procedure we present the following fundamental Lemma which provides an important consequence of the fact that $\nabla^{(1)}$ and $\nabla^{(2)}$ are forced to be almost hydrodynamically equivalent:

Lemma 5.1 *Let $\nabla^{(1)}$ and $\nabla^{(2)}$ be two almost hydrodynamically equivalent flat connections associated to the products \circ and $*$ respectively. Then in flat coordinates for $\nabla^{(1)}$, we have:*

$$\tilde{\Gamma}_{il}^{(2)k} \tilde{c}_{mj}^{*l} = \tilde{\Gamma}_{jl}^{(2)k} \tilde{c}_{mi}^{*l}. \quad (5.2)$$

Proof: By definition of almost hydrodynamical equivalence, we have $(d_{\nabla^{(1)}} - d_{\nabla^{(2)}})(X*) = 0$ for every vector fields X (and equivalently with $X\circ$). By straightforward computation we get

$$\Gamma_{lj}^{(1)k}(X*)_i^l - \Gamma_{li}^{(1)k}(X*)_j^l = \Gamma_{lj}^{(2)k}(X*)_i^l - \Gamma_{li}^{(2)k}(X*)_j^l.$$

In flat coordinates for $\nabla^{(1)}$ the above condition reduces to (5.2). ■

We are now able to find a general formula for the dual product and the dual connection under the assumption (5.1).

Theorem 5.2 *In the generalized Saito flat coordinates $\{u^1, \dots, u^n\}$, the Christoffel symbols of the dual connection and the Christoffel symbols of the trivial connection on \mathbb{C}^n are related by the formula*

$$\tilde{\Gamma}_{jk}^{(2)i} = -\frac{d_i}{d_i - 1} \frac{\partial^2 u^i}{\partial p^m \partial p^n} (J^{-1})_j^m (J^{-1})_k^n, \quad (5.3)$$

where $J_j^i = \frac{\partial u^i}{\partial p^j}$ and $d_i = \text{degree}(u^i)$, the degree of the invariant polynomial u^i .

Proof: Since $\nabla^{(1)}$ and $\nabla^{(2)}$ are almost hydrodynamically equivalent, relation (5.2) holds. Moreover, it is immediate to prove that the Euler vector field $E = \sum_k p^k \frac{\partial}{\partial p^k}$ expressed in the Saito flat coordinates is given by $\tilde{E} = \sum_k d_k u^k \frac{\partial}{\partial u^k}$, where d_k are the degrees of the invariant polynomials.

From the condition $\nabla^{(2)} E = 0$, we obtain:

$$\tilde{\Gamma}_{jl}^{(2)k} \tilde{E}^l = -d_k \delta_j^k. \quad (5.4)$$

If we multiply equation (5.2) by \tilde{E}^i on both sides we get

$$\tilde{\Gamma}_{il}^{(2)k} \tilde{E}^i \tilde{c}_{mj}^{*l} = \tilde{\Gamma}_{jl}^{(2)k} \tilde{c}_{mi}^{*l} \tilde{E}^i.$$

Taking into account that E is the unity of the dual product, the right hand side reduces to $\tilde{\Gamma}_{jm}^{(2)k}$, while, due to (5.4), the left hand side reduces to $-d_k \tilde{c}_{mj}^{*k}$.

In this way we get:

$$\tilde{\Gamma}_{jm}^{(2)k} = -d_k \tilde{c}_{mj}^{*k}. \quad (5.5)$$

The Christoffel symbols of the dual connection are also given by the formula

$$\tilde{\Gamma}_{jk}^{(2)i} = -\frac{\partial^2 u^i}{\partial p^m \partial p^n} (J^{-1})_j^m (J^{-1})_k^n - \tilde{c}_{jk}^{*i},$$

which is simply obtained by (5.1) applying the transformation formula for Christoffel symbols passing from the coordinates $\{p^1, \dots, p^n\}$ to the coordinates $\{u^1, \dots, u^n\}$. Combining these two relations we get

$$\tilde{c}_{jk}^{*i} = -\frac{1}{1 - d_i} \frac{\partial^2 u^i}{\partial p^m \partial p^n} (J^{-1})_j^m (J^{-1})_k^n,$$

and from this using (5.5) and the standing hypothesis (5.1), we find the fundamental formula (5.3). ■

It is important to observe that the fundamental relation (5.3) holds only in the generalized Saito flat coordinates: indeed in deriving it we used relation (5.2), which is true only in a coordinate system given by a distinguished set of basic of invariant polynomials in which the Christoffel symbols of $\nabla^{(1)}$ vanish identically, i.e. the generalized Saito flat coordinates. This means that (5.3) can be used to define the dual connection only in this special set of coordinates. On the other hand, we can use this fact to select the generalized Saito coordinates among other coordinate systems. Indeed, among the basic invariant polynomials we choose those for which the formula (5.3) *defines a flat* connection and for which the structure constants \tilde{c}_{jk}^i of \circ , defined via

$$\tilde{c}_{jk}^i = \tilde{c}_{jl}^{*i} \tilde{c}_{km}^{*l} (e^{-1})^m,$$

are compatible with the natural connection $\nabla^{(1)}$. In this way, in the Saito flat coordinates, we have that the Christoffel symbols of the natural connection, the Christoffel symbols of the dual connection and the structure constants of the dual product can all be reconstructed directly and unambiguously from the Christoffel symbols of the trivial connection expressed in such coordinates:

$$\tilde{\Gamma}_{ij}^{(1)k} = 0, \quad \tilde{\Gamma}_{ij}^{(2)k} = \frac{d_i}{1 - d_i} \frac{\partial^2 u^i}{\partial p^m \partial p^n} (J^{-1})_j^m (J^{-1})_k^n, \quad \tilde{c}_{jk}^{*i} = \frac{1}{d_i - 1} \frac{\partial^2 u^i}{\partial p^m \partial p^n} (J^{-1})_j^m (J^{-1})_k^n.$$

Coming back to the original coordinates on \mathbb{C}^n given by $\{p^1, \dots, p^n\}$ we get the following theorem, whose proof is a straightforward application of tensorial transformations.

Theorem 5.3 *In the coordinates $\{p^1, \dots, p^n\}$ the structure constants of the dual product and the Christoffel symbols of the dual connection are given by the formulas*

$$c_{jk}^{*i} = -\Gamma_{jk}^{(2)i} = \frac{1}{d_l - 1} \frac{\partial^2 u^l}{\partial p^j \partial p^k} (J^{-1})_l^i, \quad (5.6)$$

where $J_j^i = \frac{\partial u^i}{\partial p^j}$.

In practice, to select the Saito flat coordinates it is convenient to impose the flatness of the dual connection and the compatibility between $\nabla^{(1)}$ and \circ in the coordinates $\{p^1, \dots, p^n\}$, where the Christoffel symbols of $\nabla^{(2)}$ are given by the simple formula (5.6). Once the generalized Saito coordinates have been selected, one can easily reconstruct the full bi-flat F -manifold structure. It turns out that in all the cases we have investigated the unit of the product \circ is the vector field $e = \frac{\partial}{\partial u^n}$ where u^n is the highest degree basic invariant polynomial.

Remark 5.4 *With this choice, one has $[e, E] = d_n e$. In order to obtain $[e, E] = e$ one should normalize the Euler vector field and the dual product as $E \rightarrow \frac{E}{d_n}$ and $c_{jk}^{*i} \rightarrow d_n c_{jk}^{*i}$. Using this normalization the r.h.s of (5.1) must be replaced by $-\frac{1}{d_n} X^*$.*

Below we present the results of the procedure just described for *all* well-generated exceptional complex reflection groups of rank 2 and 3 and for the families $G(m, 1, 2)$ and $G(m, 1, 3)$. In particular, for each case we provide the Saito flat coordinates (the distinguished basis of the invariant polynomials) and the vector potential for the product \circ in the Saito flat coordinates. The other data for the bi-flat F -structure can be easily reconstructed from these data using the formulas above.

5.1 The case of G_4

In this case the ring of invariants is generated by (see [25]):

$$U_1 = p_1^4 + 2i\sqrt{3}p_1^2 p_2^2 + p_2^4, \quad U_2 = p_1^5 p_2 - p_1 p_2^5.$$

Here U_1 and U_2 coincide with the generalized Saito flat coordinates.

Up to inessential linear terms the vector potential for \tilde{c}_{jk}^i is given by

$$A^1 = u_1 u_2, \quad A^2 = \frac{1}{2} u_2^2 - \frac{1}{96} i \sqrt{3} u_1^3.$$

Lowering the indices with the metric

$$\eta = \begin{pmatrix} 0 & \frac{1}{12} \\ \frac{1}{12} & 0 \end{pmatrix}$$

we get an exact 1-form whose potential coincides with the Frobenius potential for G_4 .

5.2 The case of G_5

In this case the ring of invariants is generated by (see [25]):

$$U_1 = p_1^5 p_2 - p_1 p_2^5, \quad U_2 = (p_1^4 + 2i\sqrt{3}p_1^2 p_2^2 + p_2^4)^3.$$

We look for generalized Saito flat coordinates of the form $u_1 = U_1$, $u_2 = U_2 + cU_1^2$. Under the assumption of the general algorithm we presented, we have that c is fixed by the compatibility of the product \circ with the natural connection $\nabla^{(1)}$ (in the coordinates p), and it is given by $c_1 = -6i\sqrt{3}$. Up to inessential linear terms the vector potential for \circ in the generalized Saito flat coordinates is given by

$$A^1 = u_1^2, \quad A^2 = \frac{1}{2}u_2^2 - \frac{141}{4}u_1^4.$$

It is easy to check that the above vector potential does not come from a Frobenius potential.

5.3 The case of G_6

G_6 is a well-generated complex reflection group of rank two. Its ring of invariants is generated by (see [25]):

$$U_1 = p_1^4 + 2i\sqrt{3}p_1^2 p_2^2 + p_2^4, \quad U_2 = p_1^{10} p_2^2 - 2p_1^6 p_2^6 + p_1^2 p_2^{10}.$$

We look for generalized Saito flat coordinates of the form $u_1 = U_1$, $u_2 = U_2 + cU_1^3$. The compatibility between the natural connection $\nabla^{(1)}$ and \circ fixes a unique value of c , namely $c = \frac{5}{288}i\sqrt{3}$ and consequently a distinguished set of basic invariants. From these we get immediately the vector potential A^k , $k = 1, 2$ for \circ in the flat coordinates is given by (up to inessential linear terms):

$$A^1 = u_1 u_2 - \frac{1}{576}i\sqrt{3}u_1^4, \quad A^2 = \frac{1}{2}u_2^2 - \frac{1}{6144}u_1^6.$$

It is easy to check that the above vector potential does not come from a Frobenius potential.

5.4 The case of G_8

G_8 is a well-generated complex reflection group of rank two. Basic invariants are given by (see [25]):

$$u_1 = p_1^8 + 14p_1^4p_2^4 + p_2^8, \quad u_2 = p_1^{12} - 33p_1^8p_2^4 - 33p_1^4p_2^8 + p_2^{12}.$$

These are automatically generalized Saito coordinates and there are no arbitrary constants to be fixed. The compatibility between the natural connection $\nabla^{(1)}$ and \circ is automatically satisfied, as well all the other properties characterizing a bi-flat F structure. The vector potential A^k , $k = 1, 2$ for \circ in the flat coordinates is given by (up to inessential linear terms)

$$A^1 = u_1u_2, \quad A^2 = \frac{1}{2}u_2^2 + \frac{3}{8}u_1^3.$$

Lowering the indices with the metric:

$$\eta = \begin{pmatrix} 0 & \frac{1}{12} \\ \frac{1}{12} & 0 \end{pmatrix}$$

we get an exact 1-form, whose potential coincides with the Frobenius potential for G_8 .

5.5 The case of G_9

G_9 is a well-generated complex reflection group of rank two. Basic polynomial invariants are (see [25]):

$$U_1 = p_1^8 + 14p_1^4p_2^4 + p_2^8, \quad U_2 = (p_1^{12} - 33p_1^8p_2^4 - 33p_1^4p_2^8 + p_2^{12})^2.$$

We look for Saito flat coordinates of the form $u_1 = U_1$, $u_2 = U_2 + cU_1^3$. Imposing the compatibility of the natural connection $\nabla^{(1)}$ with \circ we obtain a unique value of c , namely $c = -11/16$ and consequently a distinguished basis of invariant polynomials (generalized Saito coordinates). In this way we obtain the vector potential for \circ (up to inessential linear terms) given by:

$$A^1 = u_1u_2 + \frac{3}{32}u_1^4, \quad A^2 = \frac{1}{2}u_2^2 + \frac{33}{512}u_1^6.$$

It is easy to check that the above vector potential does not come from a Frobenius potential.

5.6 The case of G_{10}

G_{10} is a well-generated complex reflection group of rank two. Basic polynomial invariants are given by (see [25]):

$$U_1 = p_1^{12} - 33p_1^8p_2^4 - 33p_1^4p_2^8 + p_2^{12}, \quad U_2 = (p_1^8 + 14p_1^4p_2^4 + p_2^8)^3.$$

We look for generalized Saito flat coordinates of the form $u_1 = U_1$, $u_2 = U_2 + cU_1^2$. The compatibility between the natural connection $\nabla^{(1)}$ and \circ determines a unique value of c , namely $c = -\frac{7}{12}$ and consequently a distinguished basis of invariant polynomials (Saito coordinates). In this way we obtain the vector potential (up to inessential linear terms)

$$A^1 = u_1u_2 + \frac{1}{18}u_1^3, \quad A^2 = \frac{1}{2}u_2^2 + \frac{35}{432}u_1^4.$$

It is easy to check that the above vector potential does not come from a Frobenius potential.

5.7 The case of G_{14}

G_{14} is a well-generated complex reflection group of rank two. Basic polynomial invariants are provided by (see [25]):

$$U_1 = p_1^5p_2 - p_1p_2^5, \quad U_2 = (p_1^{12} - 33p_1^8p_2^4 - 33p_1^4p_2^8 + p_2^{12})^2.$$

We look for generalized Saito flat coordinates of the form $u_1 = U_1$, $u_2 = U_2 + cU_1^4$. The compatibility between the natural connection $\nabla^{(1)}$ and \circ determines a unique value of c , namely $c = 66$, and consequently a distinguished set of basic invariant polynomials. In this way we obtain the vector potential (up to inessential linear terms)

$$A^1 = -\frac{24}{5}u_1^5 + u_1u_2, \quad A^2 = \frac{1}{2}u_2^2 + 792u_1^8.$$

It is easy to check that the above vector potential does not come from a Frobenius potential.

5.8 The case of G_{16}

G_{16} is a well-generated complex reflection group of rank two. Basic polynomial invariants are given by (see [25]):

$$\begin{aligned}
u_1 &= p_1^{20} - \frac{38}{3}\sqrt{5}p_1^{18}p_2^2 - 19p_1^{16}p_2^4 - 152\sqrt{5}p_1^{14}p_2^6 - 494p_1^{12}p_2^8 + \frac{988}{3}\sqrt{5}p_1^{10}p_2^{10} + p_2^{20} + \\
&\quad - \frac{38}{3}\sqrt{5}p_2^{18}p_1^2 - 19p_2^{16}p_1^4 - 152\sqrt{5}p_2^{14}p_1^6 - 494p_2^{12}p_1^8, \\
u_2 &= p_1^{29}p_2 - \frac{116}{45}\sqrt{5}p_1^{27}p_2^3 + \frac{1769}{25}p_1^{25}p_2^5 + \frac{464}{5}\sqrt{5}p_1^{23}p_2^7 + \frac{2001}{5}p_1^{21}p_2^9 - \frac{2668}{15}\sqrt{5}p_1^{19}p_2^{11} \\
&\quad + \frac{12673}{5}p_1^{17}p_2^{13} - p_2^{29}p_1 + \frac{116}{45}\sqrt{5}p_2^{27}p_1^3 - \frac{1769}{25}p_2^{25}p_1^5 - \frac{464}{5}\sqrt{5}p_2^{23}p_1^7 - \frac{2001}{5}p_2^{21}p_1^9 \\
&\quad + \frac{2668}{15}\sqrt{5}p_2^{19}p_1^{11} - \frac{12673}{5}p_2^{17}p_1^{13}.
\end{aligned}$$

There are no arbitrary constants in this case, and these invariant polynomials are automatically generalized Saito coordinates. All the conditions defining a bi-flat F structure are automatically satisfied. For the vector potential of \circ we obtain (up to inessential linear terms)

$$A^1 = u_1u_2, \quad A^2 = \frac{1}{2}u_2^2 - \frac{1}{800}\sqrt{5}u_1^3.$$

Lowering the indices with the metric:

$$\eta = \begin{pmatrix} 0 & \frac{19}{900} \\ \frac{19}{900} & 0 \end{pmatrix}$$

we get an exact 1-form whose potential coincides with the Frobenius potential for G_{16} .

5.9 The case of G_{17}

G_{17} is a well-generated complex reflection group of rank two. Basic polynomial invariants are given by (see [25]):

$$\begin{aligned}
U_1 &= p_1^{20} - \frac{38}{3}\sqrt{5}p_1^{18}p_2^2 - 19p_1^{16}p_2^4 - 152\sqrt{5}p_1^{14}p_2^6 - 494p_1^{12}p_2^8 + \frac{988}{3}\sqrt{5}p_1^{10}p_2^{10} + p_2^{20} \\
&\quad - \frac{38}{3}\sqrt{5}p_2^{18}p_1^2 - 19p_2^{16}p_1^4 - 152\sqrt{5}p_2^{14}p_1^6 - 494p_2^{12}p_1^8, \\
U_2 &= (p_1^{29}p_2 - \frac{116}{45}\sqrt{5}p_1^{27}p_2^3 + \frac{1769}{25}p_1^{25}p_2^5 + \frac{464}{5}\sqrt{5}p_1^{23}p_2^7 + \frac{2001}{5}p_1^{21}p_2^9 - \frac{2668}{15}\sqrt{5}p_1^{19}p_2^{11} \\
&\quad + \frac{12673}{5}p_1^{17}p_2^{13} - p_2^{29}p_1 + \frac{116}{45}\sqrt{5}p_2^{27}p_1^3 - \frac{1769}{25}p_2^{25}p_1^5 - \frac{464}{5}\sqrt{5}p_2^{23}p_1^7 - \frac{2001}{5}p_2^{21}p_1^9 \\
&\quad + \frac{2668}{15}\sqrt{5}p_2^{19}p_1^{11} - \frac{12673}{5}p_2^{17}p_1^{13})^2.
\end{aligned}$$

We look for generalized Saito flat coordinates of the form $u_1 = U_1$, $u_2 = U_2 + cU_1^3$. Imposing the compatibility between the natural connection $\nabla^{(1)}$ and \circ we obtain a unique value of c , namely $c = \frac{29}{12000}\sqrt{5}$ and consequently a distinguished basis of invariant polynomials (Saito coordinates). All the other conditions defining a bi-flat F structure are satisfied. In this way we obtain the vector potential for \circ (up to inessential linear terms)

$$A^1 = u_1 u_2 - \frac{3}{8000}\sqrt{5}u_1^4, \quad A^2 = \frac{1}{2}u_2^2 + \frac{319}{96000000}u_1^6.$$

It is easy to check that the above vector potential does not come from a Frobenius potential.

5.10 The case of G_{18}

G_{18} is a well-generated complex reflection group of rank two. Basic polynomial invariants are given by (see [25]):

$$\begin{aligned} U_1 &= p_1^{29} p_2 - \frac{116}{45}\sqrt{5} p_1^{27} p_2^3 + \frac{1769}{25} p_1^{25} p_2^5 + \frac{464}{5}\sqrt{5} p_1^{23} p_2^7 + \frac{2001}{5} p_1^{21} p_2^9 - \frac{2668}{15}\sqrt{5} p_1^{19} p_2^{11} \\ &\quad + \frac{12673}{5} p_1^{17} p_2^{13} - p_2^{29} p_1 + \frac{116}{45}\sqrt{5} p_2^{27} p_1^3 - \frac{1769}{25} p_2^{25} p_1^5 - \frac{464}{5}\sqrt{5} p_2^{23} p_1^7 - \frac{2001}{5} p_2^{21} p_1^9 \\ &\quad + \frac{2668}{15}\sqrt{5} p_2^{19} p_1^{11} - \frac{12673}{5} p_2^{17} p_1^{13}, \\ U_2 &= (p_1^{20} - \frac{38}{3}\sqrt{5} p_1^{18} p_2^2 - 19 p_1^{16} p_2^4 - 152\sqrt{5} p_1^{14} p_2^6 - 494 p_1^{12} p_2^8 + \frac{988}{3}\sqrt{5} p_1^{10} p_2^{10} + p_2^{20} + \\ &\quad - \frac{38}{3}\sqrt{5} p_2^{18} p_1^2 - 19 p_2^{16} p_1^4 - 152\sqrt{5} p_2^{14} p_1^6 - 494 p_2^{12} p_1^8)^3. \end{aligned}$$

We look for generalized Saito flat coordinates of the form $u_1 = U_1$, $u_2 = U_2 + cU_1^2$. Imposing the compatibility of the natural connection $\nabla^{(1)}$ with \circ , we obtain a unique value of c , namely $c = 38\sqrt{5}$ and consequently a distinguished basis of invariant polynomials (Saito coordinates). In this way we obtain the vector potential for \circ (up to inessential linear terms)

$$A^1 = -\frac{16}{3}\sqrt{5}u_1^3 + u_1 u_2, \quad A^2 = \frac{1}{2}u_2^2 + \frac{4180}{3}u_1^4.$$

It is easy to check that the above vector potential does not come from a Frobenius potential.

5.11 The case of G_{20}

G_{20} is a well-generated complex reflection group of rank two. Basic polynomial invariants are given by (see [25]):

$$\begin{aligned} u_1 &= p_1^{12} + \frac{22}{5}\sqrt{5}p_1^{10}p_2^2 - 33p_1^8p_2^4 - \frac{44}{5}\sqrt{5}p_1^6p_2^6 - 33p_1^4p_2^8 + \frac{22}{5}\sqrt{5}p_1^2p_2^{10} + p_2^{12}, \\ u_2 &= p_1^{29}p_2 - \frac{116}{45}\sqrt{5}p_1^{27}p_2^3 + \frac{1769}{25}p_1^{25}p_2^5 + \frac{464}{5}\sqrt{5}p_1^{23}p_2^7 + \frac{2001}{5}p_1^{21}p_2^9 - \frac{2668}{15}\sqrt{5}p_1^{19}p_2^{11} \\ &\quad + \frac{12673}{5}p_1^{17}p_2^{13} - p_2^{29}p_1 + \frac{116}{45}\sqrt{5}p_2^{27}p_1^3 - \frac{1769}{25}p_2^{25}p_1^5 - \frac{464}{5}\sqrt{5}p_2^{23}p_1^7 - \frac{2001}{5}p_2^{21}p_1^9 \\ &\quad + \frac{2668}{15}\sqrt{5}p_2^{19}p_1^{11} - \frac{12673}{5}p_2^{17}p_1^{13}. \end{aligned}$$

In this case, the general base does not depend on arbitrary constants, and all the conditions defining a bi-flat F -structure are automatically satisfied. The computation of the vector potential for \circ gives (up to inessential linear terms):

$$A^1 = u_1u_2, \quad A^2 = \frac{1}{2}u_2^2 + \frac{1}{960}\sqrt{5}u_1^5.$$

Lowering the indices with the metric:

$$\eta = \begin{pmatrix} 0 & \frac{11}{900} \\ \frac{11}{900} & 0 \end{pmatrix}$$

we get an exact 1-form whose potential coincides with the Frobenius potential for G_{20} .

5.12 The case of G_{21}

G_{21} is a well-generated complex reflection group of rank two. Basic polynomial invariants are given by (see [25]):

$$\begin{aligned} U_1 &= p_1^{12} + \frac{22}{5}\sqrt{5}p_1^{10}p_2^2 - 33p_1^8p_2^4 - \frac{44}{5}\sqrt{5}p_1^6p_2^6 - 33p_1^4p_2^8 + \frac{22}{5}\sqrt{5}p_1^2p_2^{10} + p_2^{12}, \\ U_2 &= (p_1^{29}p_2 - \frac{116}{45}\sqrt{5}p_1^{27}p_2^3 + \frac{1769}{25}p_1^{25}p_2^5 + \frac{464}{5}\sqrt{5}p_1^{23}p_2^7 + \frac{2001}{5}p_1^{21}p_2^9 - \frac{2668}{15}\sqrt{5}p_1^{19}p_2^{11} \\ &\quad + \frac{12673}{5}p_1^{17}p_2^{13} - p_2^{29}p_1 + \frac{116}{45}\sqrt{5}p_2^{27}p_1^3 - \frac{1769}{25}p_2^{25}p_1^5 - \frac{464}{5}\sqrt{5}p_2^{23}p_1^7 - \frac{2001}{5}p_2^{21}p_1^9 \\ &\quad + \frac{2668}{15}\sqrt{5}p_2^{19}p_1^{11} - \frac{12673}{5}p_2^{17}p_1^{13})^2. \end{aligned}$$

We look for generalized Saito flat coordinates of the form $u_1 = U_1$, $u_2 = U_2 + cU_1^5$. The compatibility of the natural connection $\nabla^{(1)}$ with the product \circ determines a unique value of c , namely $c = -\frac{29}{14400}\sqrt{5}$ and consequently a distinguished basis

of invariant polynomials (Saito coordinates). All the other conditions defining a bi-flat F structure are automatically satisfied.

Computing the vector potential for \circ , we obtain (up to inessential linear terms)

$$A^1 = u_1 u_2 + \frac{1}{8640} \sqrt{5} u_1^6, \quad A^2 = \frac{1}{2} u_2^2 + \frac{551}{149299200} u_1^{10}.$$

It is easy to check that the above vector potential does not come from a Frobenius potential.

5.13 The case of G_{24}

The complex reflection group G_{24} is a rank three well-generated complex reflection group. Basic invariants are given by the following homogeneous polynomials of degree 4, 6, 14 (see for instance Section 6 of [21] or [13]):

$$\begin{aligned} U_1 &= p_1^3 p_2 + p_1 p_3^3 + p_2^3 p_3, \\ U_2 &= p_1^5 p_3 + p_1 p_2^5 + p_2 p_3^5 - 5 p_1^2 p_2^2 p_3^2, \\ U_3 &= p_1^{14} + p_2^{14} + p_3^{14} + 18 p_1^7 p_2^7 + 18 p_1^7 p_3^7 + 18 p_2^7 p_3^7 \\ &\quad - 126 p_1^6 p_2^3 p_3^5 - 126 p_1^5 p_2^6 p_3^3 - 126 p_1^3 p_2^5 p_3^6 + 375 p_1^8 p_2^4 p_3^2 + 375 p_1^4 p_2^2 p_3^8 + 375 p_1^2 p_2^8 p_3^4 \\ &\quad - 250 p_1^9 p_2 p_3^4 - 250 p_1^4 p_2^9 p_3 - 250 p_1 p_2^4 p_3^9 - 34 p_1^{11} p_2^2 p_3 - 34 p_1^2 p_2 p_3^{11} - 34 p_1 p_2^{11} p_3^2. \end{aligned}$$

Notice that U_2 is equal (up to a constant factor) to the determinant of the Hessian of U_1 , while U_3 is constructed using a differential determinant involving U_1 and U_2 (see formula (1.14) in [13]). We look for generalized Saito flat coordinates of the form

$$u_1 = U_1, \quad u_2 = U_2, \quad u_3 = U_3 + c U_1^2 U_2.$$

Imposing the flatness of the dual connection (in the original coordinates of \mathbb{C}^n)

$$\Gamma_{jk}^{(2)i} = \frac{1}{1 - d_l} \frac{\partial^2 u^l}{\partial p^j \partial p^k} (J^{-1})^i_l,$$

we get $c = -34$. In the flat coordinates we obtain that the vector potential for \circ is given by (up to inessential linear terms)

$$\begin{aligned} A^1 &= -\frac{14}{3} u_1^3 u_2 - u_1 u_2 - \frac{14}{3} u_2^3, \\ A^2 &= -\frac{14}{5} u_1^5 + 14 u_1^2 u_2^2 - u_2 u_3, \\ A^3 &= -\frac{1}{2} u_3^2 - 56 u_1^7 - 294 u_1^4 u_2^2 + 196 u_1 u_2^4. \end{aligned}$$

Rescaling the variables $v_1 = -2^{-\frac{1}{3}} u_1$, $v_2 = u_2$, $v_3 = -2^{\frac{1}{3}} \frac{u_3}{56}$, we obtain the vector potential found in [20] (Section 7.2). It is easy to check that the above vector potential does not come from a Frobenius potential.

5.14 The case of G_{25}

Basic invariants are given by the following homogeneous polynomials of degree 6, 9, 12 (see for instance [31]):

$$\begin{aligned} U_1 &= p_1^6 + p_2^6 + p_3^6 - 10p_1^3p_2^3 - 10p_1^3p_3^3 - 10p_2^3p_3^3 \\ U_2 &= (p_1^3 - p_2^3)(p_1^3 - p_3^3)(p_2^3 - p_3^3) \\ U_3 &= (p_1^3 + p_2^3 + p_3^3)((p_1^3 + p_2^3 + p_3^3)^3 + 216p_1^3p_2^3p_3^3). \end{aligned}$$

We look for generalized Saito flat coordinates of the form:

$$u_1 = U_1, \quad u_2 = U_2, \quad u_3 = U_3 + c_1U_1^2.$$

Imposing the flatness of the dual connection in the original coordinates of \mathbb{C}^n

$$\Gamma_{jk}^{(2)i} = \frac{1}{1 - d_l} \frac{\partial^2 u^l}{\partial p^j \partial p^k} (J^{-1})_l^i,$$

we obtain $c = -\frac{5}{8}$. In the Saito flat coordinates, the vector potential for \circ is given by (up to inessential linear terms)

$$A^1 = \frac{1}{8}(3u_1^2 - 8u_3)u_2, \quad A^2 = -\frac{1}{2}u_3^2 - 144u_1u_2^2 - \frac{3}{64}u_1^4, \quad A^3 = -u_1u_3 + 192u_2^2.$$

Lowering the indices with the metric:

$$\eta = \begin{pmatrix} 0 & 0 & -\frac{5}{144} \\ 0 & \frac{40}{3} & 0 \\ -\frac{5}{144} & 0 & 0 \end{pmatrix}$$

we get an exact 1-form whose potential coincides with the Frobenius potential for G_{20} .

5.15 The case of G_{26}

The basic invariants are:

$$\begin{aligned} U_1 &= p_1^6 - 10p_1^3p_2^3 - 10p_1^3p_3^3 + p_2^6 - 10p_2^3p_3^3 + p_3^6 \\ U_2 &= (p_1^3 + p_2^3 + p_3^3)(216p_1^3p_2^3p_3^3 + (p_1^3 + p_2^3 + p_3^3)^3) \\ U_3 &= (p_1^3 + p_2^3 + p_3^3)^6 - 540p_1^3p_2^3p_3^3(p_1^3 + p_2^3 + p_3^3)^3 - 5832p_1^6p_2^6p_3^6 \end{aligned}$$

We look for generalized Saito flat coordinates of the form:

$$u_1 = U_1, \quad u_2 = U_2 + c_1U_1^2, \quad u_3 = U_3 + c_2U_1U_2 + c_3U_1^3.$$

Imposing the flatness of the dual connection in the original coordinates of \mathbb{C}^n

$$\Gamma_{jk}^{(2)i} = \frac{1}{1 - d_l} \frac{\partial^2 u^l}{\partial p^j \partial p^k} (J^{-1})_l^i,$$

we obtain

$$c_1 = -\frac{1}{6}, \quad c_2 = -\frac{1}{2}, \quad c_3 = \frac{1}{9}.$$

In the Saito flat coordinates the vector potential for \circ is given by

$$\begin{aligned} A^1 &= u_1 u_3 + \frac{3}{4} u_2^2 - \frac{1}{4} u_1^2 u_2 + \frac{1}{48} u_1^4, \\ A^2 &= \frac{1}{4} u_1 u_2^2 + \frac{1}{12} u_1^3 u_2 + u_2 u_3 - \frac{1}{144} u_1^5, \\ A^3 &= \frac{1}{2} u_3^2 + \frac{1}{4} u_2^3 + \frac{1}{4} u_2^2 u_1^2 - \frac{1}{48} u_1^4 u_2 + \frac{1}{216} u_1^6 \end{aligned}$$

It is easy to check that this vector potential does not come from a Frobenius potential. In other words we do not recover the Frobenius manifold associated with the Shephard group G_{26} . We will see later how to obtain this structure modifying slightly the above procedure.

5.16 The case of G_{27}

G_{27} is a well-generated complex reflection group. As a basis for its invariants, we use the invariants of Wiman [47].

$$\begin{aligned} U_1 &= 10p_1^3 p_2^3 + 9p_3(p_1^5 + p_2^5) - 45p_1^2 p_2^2 p_3^2 - 135p_1 p_2 p_3^4 + 27p_3^6 \\ U_2 &= \frac{1}{20250} \det(\text{hessian}(U_1)) \\ U_3 &= \frac{1}{24300} \text{Bord}(U_1, U_2), \end{aligned}$$

where Bord is a differential determinantal expression involving U_1 and U_2 . Using the same procedure adopted in the previous cases involving rank three well-generated complex reflection groups, we obtain generalized Saito flat coordinates

$$u^1 = U^1, \quad u^2 = U_2 + c_1 u_1^2, \quad u_3 = U_3 + c_2 U_2^2 u_1 + c_3 U_2 u_1^3 + c_4 u_1^5,$$

with

$$c_1 = -\frac{1}{9}, \quad c_2 = -\frac{17}{18}, \quad c_3 = \frac{29}{27}, \quad c_4 = -\frac{43}{486}.$$

The vector potential for \circ is provided by:

$$\begin{aligned} A^1 &= u_1 u_3 - \frac{5}{6} u_2^3 - \frac{5}{18} u_2^2 u_1^2 + \frac{5}{81} u_2 u_1^4 + \frac{2}{2187} u_1^6, \\ A^2 &= u_2 u_3 + \frac{25}{54} u_2^3 u_1 - \frac{25}{81} u_2^2 u_1^3 - \frac{10}{729} u_2 u_1^5 - \frac{100}{19683} u_1^7, \\ A^3 &= \frac{1}{2} u_3^2 - \frac{35}{72} u_2^5 + \frac{875}{648} u_1^2 u_2^4 + \frac{875}{6561} u_2^2 u_1^6 + \frac{24000}{1417176} u_1^8 u_2 - \frac{350}{177147} u_1^{10}. \end{aligned}$$

Rescaling the variables we obtain the vector potential found in [20] (Section 7.3). It is easy to check that the above vector potential does not come from a Frobenius potential.

5.17 The case of $G(m, 1, 2)$

We can choose as basic invariants

$$U_1 = p_1^m + p_2^m, \quad U_2 = p_1^{2m} + p_2^{2m}.$$

Generalized Saito flat coordinates are $u_1 = U_1$ and $u_2 = U_2 + cU_1^2$ with $c = -\frac{1}{2}\frac{2m-1}{m}$ and the vector potential is

$$\begin{aligned} A^1 &= u_1 u_2 + \frac{1}{6} \frac{m-2}{m} u_1^3, \\ A^2 &= \frac{1}{2} u_2^2 + \frac{1}{12} \frac{m-1}{m^2} u_1^4. \end{aligned}$$

It is easy to check that the above vector potential does not come from a Frobenius potential.

5.18 The case of $G(m, 1, 3)$

We can choose as basic invariants

$$U_1 = p_1^m + p_2^m + p_3^m, \quad U_2 = p_1^{2m} + p_2^{2m} + p_3^{2m}, \quad U_3 = p_1^{3m} + p_2^{3m} + p_3^{3m}.$$

Generalized Saito flat coordinates are

$$u_1 = U_1, \quad u_2 = U_2 + c_1 U_1^2, \quad u_3 = U_3 + c_2 U_1 U_2 + c_3 U_1^3$$

with

$$c_1 = -\frac{1}{3} \frac{2m-1}{m}, \quad c_2 = -\frac{1}{2} \frac{3m-1}{m}, \quad c_3 = \frac{2}{9} c_2^2$$

and the vector potential is

$$\begin{aligned} A^1 &= u_1 u_3 + \left(\frac{1}{2} \frac{3m-1}{m} - \frac{5}{4} \right) u_1^2 u_2 + \left(\frac{1}{12} \frac{(3m-1)^2}{m^2} - \frac{31}{72} \frac{3m-1}{m} + \frac{5}{9} \right) u_1^4 + \frac{3}{8} u_2^2, \\ A^2 &= u_2 u_3 + \frac{1}{180} \frac{m^3 - 2m^2 - m + 2}{m^3} u_1^5 + \frac{1}{180} \frac{10m^3 - 10m^2}{m^3} u_1^3 u_2 + \\ &\quad + \frac{1}{180} \frac{45m^3 - 45m^2}{m^3} u_1 u_2^2 \\ A^3 &= \frac{1}{2160} \frac{4m^3 - 12m^2 + 16m - 8}{m^4} u_1^6 + \frac{1}{2160} \frac{45m^3 - 135m^2 + 90m}{m^4} u_1^4 u_2 + \\ &\quad + \frac{1}{2160} \frac{270m^3 - 270m^2}{m^4} u_1^2 u_2^2 + \frac{1}{16} \frac{u_2^3}{m} + \frac{1}{2} u_3^2. \end{aligned}$$

6 A universal formula for the dual product

It is well-known that in the case of the Frobenius structure associated to any Coxeter group, the dual product is immediately recovered via Veselov's \vee -system structure associated with the roots of the Coxeter group itself. In this Section we show that, similarly to the Coxeter case, in the case of irreducible well-generated complex reflection groups of rank 2 and 3 the dual product and the dual connection are given by a universal formula of Dunkl-Kohno type. This is a generalization of the \vee -system construction, since the formula (6.1) below is obtained considering unitary projections onto the unitary complement of the hyperplanes defined by the covectors α_s , computed with respect to a suitable Hermitian metric (in all cases analyzed it is standard Hermitian metric, except for the case of G_{27}).

In order to obtain this formula, it is necessary to identify the reflecting hyperplanes in all cases under consideration. This is achieved using the fact that Shephard and Todd proved that the determinant of the Jacobian matrix $\det \frac{\partial u^i}{\partial p^j}$ factorizes into the product of linear forms defining the reflecting hyperplanes. The form corresponding to a (pseudo)-reflection of order p appears exactly with multiplicity $p - 1$ in $\det \frac{\partial u^i}{\partial p^j}$. Remarkably, for all rank 2 and rank 3 well-generated complex reflection groups of the Shephard Todd list the "weights" κ_s assigned at each hyperplane coincide with the order of the (pseudo)-reflection having as mirror exactly that hyperplane.

Theorem 6.1 *In the cases $G_4, G_5, G_6, G_8, G_9, G_{10}, G_{14}, G_{16}, G_{17}, G_{18}, G_{20}, G_{21}, G_{23}, G_{24}, G_{25}, G_{26}, G_{27}$ the dual product and the dual connection at a point p are expressed by the following formula (in the coordinates $\{p^1, \dots, p^n\}$):*

$$c_{jk}^{*i}(p) = -\Gamma_{jk}^{*i}(p) = h^{im} \left\{ \frac{1}{N} \sum_{s=1}^M \frac{\kappa_s}{\|\alpha_s\|^2} \frac{(\alpha_s)_j (\alpha_k)_p (\bar{\alpha}_s)_m}{\alpha_s(p)} \right\}, \quad (6.1)$$

where

- α_s are constant covectors in \mathbb{C}^n ;
- h^{im} are the components of the inverse of a suitable Hermitian metric. In all the cases a part from one (G_{27}) h is the standard Hermitian metric;
- N is a normalizing factor chosen in such a way that $c_{jk}^{*i} E^k = \delta_j^i$;
- M is the number of distinct factors of $\det \frac{\partial u^i}{\partial p^j}$ (the number of reflecting hyperplanes);
- κ_s is the order of the (pseudo)-reflection defined by the hyperplane $\ker(\alpha_s)$.

The list of values of M, N, κ_s and the choice of the Hermitian metric are given below. The complete list of 1-forms α obtained factorizing $\det \frac{\partial u^i}{\partial p^j}$ is given in the Appendix 1.

Table 2: Rank two complex reflection groups.

Type	Metric	M	N	κ_s
G_4	$h = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	4	4	2 (s=1..4)
G_5	$h = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	8	8	2 (s=1..8),
G_6	$h = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	10	12	2 (s=1..6), 3 (s=7..10)
G_8	$h = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	6	6	2 (s=1..6).
G_9	$h = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	18	24	2 (s=1..12), 4 (s=13..18)
G_{10}	$h = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	14	24	3 (s=1..8), 4 (s=9..14)
G_{14}	$h = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	20	24	2 (s=1..12), 3 (s=13..20)
G_{16}	$h = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	12	12	2 (s=1..12)
G_{17}	$h = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	42	60	2 (s=1..30), 5 (s=31..42)
G_{18}	$h = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	32	60	3 (s=1..20), 5 (s=21..32)
G_{20}	$h = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	20	20	2 (s=1..20)
G_{21}	$h = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	50	60	2 (s=1..30) 3 (s=31..50)

Table 3: Rank three complex reflection groups.

Type	Metric	n	N	κ_s
G_{24}	$h = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$	21	42	2 ($s=1..21$)
G_{25}	$h = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$	12	24	2 ($s=1..12$),
G_{26}	$h = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$	21	18	2 ($s=1..9$), 3 ($s=10..21$)
G_{27}	$h = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 3 \end{pmatrix}$	45	15	2 ($s=1..45$)

Proof: The proof boils down to check with a direct computation that the following identity holds true in all the cases examined

$$\frac{1}{d_l - 1} \frac{\partial^2 u^l}{\partial p^j \partial p^k} (J^{-1})_l^i = \frac{1}{N} \sum_{H \in \mathcal{H}} \frac{d\alpha_H}{\alpha_H} \otimes \kappa_H \pi_H. \quad (6.2)$$

where \mathcal{H} is the collection of reflecting hyperplanes, the weights κ_H coincide with the order of the corresponding (pesudo)-reflection fixing hyperplane H and π_H is a unitary projection onto a suitable unitary complement of H in \mathbb{C}^n . The long symbolic computations have been performed using the symbolic capabilities of the program Maple. ■

6.1 Remarks

1. In all the cases examined except for G_{27} the Hermitian metric is the standard one.
2. The proof is based on long symbolic computations. We believe that it is a general fact that the dual product and the dual connection associated to well-generated complex groups can be expressed in this way, a fact not limited to the exceptional cases of rank 2 and 3.

6.2 An analogue of Veselov's potential

In the case of \vee -systems, there is a natural potential F that allows one to recover the dual product (see formula (3.3)). In complete analogy, in our case, we have a

similar formula for a *vector potential* F^i .

Proposition 6.2 *The structure constants of the dual product c_{jk}^{*i} in the cases appearing in Theorem 6.1, can be expressed as*

$$c_{jk}^{*i}(p) = \frac{\partial F^i(p)}{\partial p^j \partial p^k},$$

where the dual potential is given by the formula

$$F^i(p) = \frac{1}{N} \sum_{H \in \mathcal{H}} \frac{\kappa_H}{\|\alpha_H\|^2} \alpha_H(p) \ln(\alpha_H(p)) \check{\alpha}_H^i. \quad (6.3)$$

The weights κ_H coincide with the order of the (pesudo)-reflection fixing hyperplane H and $\check{\alpha}_H^i$ is the i -component of the vector obtained from α_H via the Hermitian metric.

Proof: This is a straightforward computation. ■

6.3 Generalized Saito flat coordinates: the standard case

We have seen that, in all the examples, the dual product is given by the formula (6.1). Given the dual product, the fundamental formula (5.6) allows one to uniquely reconstruct the flat basic invariants. Indeed, the freedom in the choice of these invariants disappears if one requires that

$$\frac{\partial^2 u^i}{\partial p^j \partial p^k} = (d_i - 1) c_{jk}^{*s} \frac{\partial u^i}{\partial p^s}. \quad (6.4)$$

This means that u^i must be a flat coordinate of the flat connection $\nabla^{(0)} + (d_i - 1)*$. We will call basic invariants satisfying condition (6.4) the *standard generalized Saito flat coordinates*. Non standard bi-flat structures and related flat coordinates will be constructed in the next Section.

Remark 6.3 *It was observed in [14] that the condition (6.4) is satisfied by standard Saito polynomials.*

7 Bi-flat F -manifold structure on the space of orbits of groups revisited

If we look back at the results of Section 5, we observe that in the case of Shephard groups, the standard bi-flat F -structure does not coincide, in general, with

the Dubrovin's structure. In this Section we show that the bi-flat structure associated with a complex reflection group in general is not unique. In particular, we prove that in all the examples of Shephard groups where the standard structure does coincide with the Frobenius manifold structure, there is an underlying one-parameter family including both of them for special values of the parameter. As a consequence, we have a *one parameter family of generalized Saito flat coordinates*.

7.1 The case of G_{26}

We treat in details the case of G_{26} and we provide the main steps for the remaining groups. All the results of this part have been obtained with the help of the symbolic package Maple.

In this case our starting point is the dual product obtained in the Section 6

$$c_{lp}^{*i}(u) = \frac{2}{18} \sum_{s=1}^9 \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)} + \frac{3}{18} \sum_{s=10}^{21} \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)}.$$

The Christoffel symbols of the natural connection are given by:

$$\tilde{\Gamma}_{jk}^{(1)i} = \frac{\partial^2 u^m}{\partial p^j \partial p^k} (J^{-1})_m^i,$$

where (u_1, u_2, u_3) are easily obtained from the basic invariants (U_1, U_2, U_3) of Section 5:

$$u_1 = U_1, \quad u_2 = U_2 + c_1 U_1^2, \quad u_3 = U_3 + c_2 U_1 U_2 + c_3 U_1^3. \quad (7.1)$$

By construction (u_1, u_2, u_3) are flat coordinates of the flat connection $\nabla^{(1)}$.

The product \circ can be also be defined from the dual product \star , specifically its structure constants are given by:

$$\tilde{c}_{jk}^i = \tilde{c}_{jl}^{*i} \tilde{c}_{km}^{*l} (e^{-1})^m,$$

where e^{-1} is the inverse of the unit of \circ , where the inverse is computed with respect to \star .

The compatibility of \circ with the natural connection $\nabla^{(1)}$ imposes the following constraints on the constants c_i appearing in the expressions of (u_1, u_2, u_3) as a function of the basic invariants (U_1, U_2, U_3) :

$$c_1 = c_1, \quad c_2 = 3c_1, \quad c_3 = \frac{1}{2}c_1(2c_1 - 1).$$

Once these constraints are imposed to (7.1), we effectively obtain a one-parameter family (depending on c_1) of Saito flat coordinates. In particular, it is straightforward to check that the product \circ in the generalized Saito flat coordinates is given

by $\tilde{c}_{jk}^i = \frac{\partial^2 A^i}{\partial u^j \partial u^k}$ where

$$\begin{aligned} A^1 &= \frac{1}{8}(12c_1^2 + 7c_1 + 1)u_1^4 + \frac{1}{8}(-24c_1 - 6)u_2u_1^2 + u_3u_1 + \frac{3}{4}u_2^2, \\ A^2 &= \frac{1}{20}(12c_1^3 + 15c_1^2 + 3c_1)u_1^5 - \frac{1}{2}u_2u_1^3c_1 - \frac{3}{2}c_1u_2^2u_1 + u_3u_2, \\ A^3 &= \frac{1}{2}u_3^2 + \frac{1}{80}(-96c_1^4 - 96c_1^3 - 36c_1^2 - 6c_1)u_1^6 + \frac{9}{2}(c_1 + \frac{1}{2})c_1(c_1 + \frac{1}{4})u_2u_1^4 + \\ &\quad - \frac{9}{2}(c_1 + \frac{1}{2})c_1u_2^2u_1^2 + \frac{3}{4}u_2^3c_1 + \frac{3}{8}u_2^3. \end{aligned}$$

In other words, we have equipped the orbit space of G_{26} with a one parameter family of bi-flat F -structures. Among them there is only one that admits an invariant metric given by (up to multiplicative constant):

$$\eta = \begin{pmatrix} 0 & 0 & \frac{5}{18} \\ 0 & \frac{5}{12} & 0 \\ \frac{5}{18} & 0 & 0 \end{pmatrix}.$$

This corresponds to special choice $c_1 = -\frac{1}{4}$. In this case we have a Frobenius potential

$$F = \frac{5}{36}u_1u_3^2 + \frac{5}{24}u_3u_2^2 + \frac{5}{96}u_1u_2^3 + \frac{5}{192}u_2^2u_1^3 + \frac{1}{5376}u_1^7,$$

that coincides with the Frobenius potential obtained in Section 2.

In the coordinates $\{p^1, \dots, p^n\}$, for generic values of c_1 , the Christoffel symbols of the dual connection can be obtained imposing the almost-hydrodynamical equivalence between $\nabla^{(1)}$ and $\nabla^{(2)}$. This gives rise to the following equation

$$(\Gamma_{il}^{(2)k} - \Gamma_{il}^{(1)k})c_{mj}^l - (\Gamma_{jl}^{(2)k} - \Gamma_{jl}^{(1)k})c_{mi}^l,$$

which we can solve for $\Gamma_{jk}^{(2)i}$. In this case, the Christoffel symbols of $\nabla^{(2)}$ are not related anymore to the structure constants of the dual product \star by the simple formula

$$\Gamma_{jk}^{(2)i} = -c_{jk}^{*i}.$$

However, the Christoffel symbols of $\nabla^{(2)}$ can still be expressed via a Dunkl-Kohn-type connection associated with a different choice of the weights. We have

$$\Gamma_{jk}^{(2)i} = -c_{jk}^{*i} + \left(\frac{2}{3}c_1 + \frac{1}{9}\right)C_{jk}^i,$$

where

$$C_{lp}^i(u) = 4 \sum_{s=1}^9 \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)} - 3 \sum_{s=10}^{21} \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)}.$$

Let us briefly summarize the remaining cases.

7.2 The case of G_5

The Christoffel symbols of the dual connection are

$$\Gamma_{jk}^{(2)i} = -c_{jk}^{*i} + \left(-\frac{1}{4} + \frac{i}{72}\sqrt{3}c \right) C_{jk}^i,$$

where

$$C_{lp}^i(u) = 3 \sum_{s=1}^4 \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)} - 3 \sum_{s=5}^8 \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)}.$$

Up to inessential linear terms the vector potential is given by:

$$\begin{aligned} A^1 &= -(4i)\sqrt{3}u_1^3 - \frac{2}{3}u_1^3c + u_1u_2, \\ A^2 &= -(4i)\sqrt{3}u_1^4c - \frac{1}{3}u_1^4c^2 + \frac{1}{2}u_2^2. \end{aligned}$$

Finally, for $c = -6i\sqrt{3}$ one recovers the Frobenius structure listed in Table 1.

7.3 The case of G_6

The Christoffel symbols of the dual connection are

$$\Gamma_{jk}^{(2)i} = -c_{jk}^{*i} + \left(-8ic\sqrt{3} - \frac{5}{12} \right) C_{jk}^i,$$

where

$$C_{lp}^i(u) = 2 \sum_{s=1}^6 \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)} - 3 \sum_{s=7}^{10} \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)}.$$

Up to inessential linear terms the vector potential is given by:

$$\begin{aligned} A^1 &= u_2u_1 + \frac{1}{144}u_1^4(i\sqrt{3} - 72c), \\ A^2 &= \frac{1}{2}u_2^2 + \frac{1}{120}u_1^6c(i\sqrt{3} - 36c). \end{aligned}$$

Finally, for $c = \frac{1}{72}i\sqrt{3}$ one recovers the Frobenius structure listed in Table 1.

7.4 The case of G_9

The Christoffel symbols of the dual connection are

$$\Gamma_{jk}^{(2)i} = -c_{jk}^{*i} + \left(-\frac{4}{3}c - \frac{11}{12} \right) C_{jk}^i,$$

where

$$C_{lp}^i(u) = \sum_{s=1}^{12} \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)} - 2 \sum_{s=13}^{18} \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)}.$$

Up to inessential linear terms the vector potential is given by:

$$\begin{aligned} A^1 &= \frac{1}{4}(-2c-1)u_1^4 + u_2u_1, \\ A^2 &= \frac{1}{10}(-3c-3)u_1^6 + \frac{1}{2}u_2^2. \end{aligned}$$

Finally, for $c = -\frac{1}{2}$ one recovers the Frobenius structure listed in Table 1.

7.5 The case of G_{10}

The Christoffel symbols of the dual connection are

$$\Gamma_{jk}^{(2)i} = -c_{jk}^{*i} + \left(-\frac{1}{2}c - \frac{7}{24}\right) C_{jk}^i,$$

where

$$C_{lp}^i(u) = 3 \sum_{s=1}^8 \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)} - 4 \sum_{s=9}^{14} \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)}.$$

Up to inessential linear terms the vector potential is given by:

$$\begin{aligned} A^1 &= \frac{1}{3}(-2c-1)u_1^3 + u_2u_1, \\ A^2 &= \frac{1}{6}(-2c^2-2c)u_1^4 + \frac{1}{2}u_2^2. \end{aligned}$$

Finally, for $c = -\frac{1}{2}$ one recovers the Frobenius structure listed in Table 1.

7.6 The case of G_{14}

The Christoffel symbols of the dual connection are

$$\Gamma_{jk}^{(2)i} = -c_{jk}^{*i} + \left(\frac{1}{144}c - \frac{11}{24}\right) C_{jk}^i,$$

where

$$C_{lp}^i(u) = 2 \sum_{s=1}^{12} \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)} - 3 \sum_{s=13}^{20} \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)}.$$

Up to inessential linear terms the vector potential is given by:

$$\begin{aligned} A^1 &= \frac{1}{5}(-2c+108)u_1^5 + u_2u_1, \\ A^2 &= \frac{1}{14}(-4c^2+432c)u_1^8 + \frac{1}{2}u_2^2. \end{aligned}$$

Finally, for $c = 54$ one recovers the Frobenius structure listed in Table 1.

7.7 The case of G_{17}

The Christoffel symbols of the dual connection are

$$\Gamma_{jk}^{(2)i} = -c_{jk}^{*i} + \left(\frac{29}{60} - 40\sqrt{5}c \right) C_{jk}^i,$$

where

$$C_{lp}^i(u) = 2 \sum_{s=1}^{30} \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)} - 5 \sum_{s=31}^{42} \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)}.$$

Up to inessential linear terms the vector potential is given by:

$$\begin{aligned} A^1 &= u_1 u_2 + \frac{1}{1200} (\sqrt{5} - 600c) u_1^4, \\ A^2 &= \frac{1}{2} u_2^2 + \frac{1}{1000} c u_1^6 (\sqrt{5} - 300c). \end{aligned}$$

Finally, for $c = \frac{1}{600} \sqrt{5}$ one recovers the Frobenius structure listed in Table 1.

7.8 The case of G_{18}

The Christoffel symbols of the dual connection are

$$\Gamma_{jk}^{(2)i} = -c_{jk}^{*i} + \left(-\frac{19}{60} + \frac{1}{600} \sqrt{5}c \right) C_{jk}^i,$$

where

$$C_{lp}^i(u) = 3 \sum_{s=1}^{20} \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)} - 5 \sum_{s=21}^{32} \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)}.$$

Up to inessential linear terms the vector potential is given by:

$$\begin{aligned} A^1 &= 20u_1^3 \sqrt{5} - \frac{2}{3} u_1^3 c + u_1 u_2, \\ A^2 &= 20c u_1^4 \sqrt{5} - \frac{1}{3} c^2 u_1^4 + \frac{1}{2} u_2^2. \end{aligned}$$

Finally, for $c = -6i\sqrt{3}$ one recovers the Frobenius structure listed in Table 1.

7.9 The case of G_{21}

The Christoffel symbols of the dual connection are

$$\Gamma_{jk}^{(2)i} = -c_{jk}^{*i} + \left(-\frac{1}{300} (29\sqrt{5} + 14400c) \sqrt{5} \right) C_{jk}^i,$$

where

$$C_{lp}^i(u) = 2 \sum_{s=1}^{30} \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)} - 3 \sum_{s=31}^{50} \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)}.$$

Up to inessential linear terms the vector potential is given by:

$$\begin{aligned} A^1 &= -\frac{1}{1800} u_1^6 (\sqrt{5} + 600c) + u_1 u_2, \\ A^2 &= -\frac{1}{1080} c (\sqrt{5} + 300c) u_1^{10} + \frac{1}{2} u_2^2. \end{aligned}$$

Finally, for $c = -\frac{1}{600} \sqrt{5}$ one recovers the Frobenius structure listed in Table 1.

7.10 The case of $G(3, 1, 2)$

The dual connection is

$$\Gamma_{jk}^{(2)i} = -c_{jk}^{*i} + \left(c + \frac{5}{6}\right) C_{jk}^i,$$

where

$$c_{lp}^{*i}(u) = \frac{1}{6} \left(3 \sum_{s=1}^2 \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)} + 2 \sum_{s=3}^5 \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)} \right)$$

and

$$C_{lp}^i(u) = 3 \sum_{s=1}^2 \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)} - 2 \sum_{s=3}^5 \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)}.$$

Up to inessential linear terms the vector potential is

$$\begin{aligned} A^1 &= \frac{1}{6} (-3 - 4c) u_1^3 + u_1 u_2, \\ A^2 &= \frac{1}{6} (-2c^2 - 3c - 1) u_1^4 + \frac{1}{2} u_2^2. \end{aligned}$$

Finally, for $c = -\frac{3}{4}$ one recovers the Frobenius structure listed in Table 1 with $m = 3$.

7.11 The case of $G(3, 1, 3)$

The flat coordinates are

$$\begin{aligned} u_1 &= p_1^3 + p_2^3 + p_3^3 \\ u_2 &= p_1^6 + p_2^6 + p_3^6 + c_1 (p_1^3 + p_2^3 + p_3^3)^2 \\ u_3 &= p_1^9 + p_2^9 + p_3^9 + c_2 (p_1^3 + p_2^3 + p_3^3) (p_1^6 + p_2^6 + p_3^6) + c_3 (p_1^3 + p_2^3 + p_3^3)^3 \end{aligned}$$

with $c_2 = \frac{3}{2}c_1 - \frac{1}{2}$ and $c_3 := \frac{1}{2}c_1^2 - \frac{1}{3}c_1 + \frac{1}{18}$. The dual connection is

$$\Gamma_{jk}^{(2)i} = -c_{jk}^{*i} + \left(2c_1 + \frac{10}{9}\right) C_{jk}^i,$$

where

$$c_{lp}^{*i}(u) = \frac{1}{9} \left(3 \sum_{s=1}^3 \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)} + 2 \sum_{s=4}^{12} \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)} \right)$$

and

$$C_{lp}^i(u) = 3 \sum_{s=1}^3 \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)} - \sum_{s=4}^{12} \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)}.$$

Up to inessential linear terms the vector potential is

$$\begin{aligned} A^1 &= \frac{1}{24}(18c_1^2 + 19c_1 + 5)u_1^4 + \frac{1}{24}(-36c_1 - 18)u_2u_1^2 + u_1u_3 + \frac{3}{8}u_2^2, \\ A^2 &= \frac{1}{180}(54c_1^3 + 99c_1^2 + 54c_1 + 9)u_1^5 + \frac{1}{180}(-30c_1 - 10)u_2u_1^3 + \\ &\quad \frac{1}{180}(-135c_1 - 45)u_2^2u_1 + u_2u_3, \\ A^3 &= \frac{1}{2}u_3^2 + \frac{1}{2160}(-648c_1^4 - 1296c_1^3 - 972c_1^2 - 324c_1 - 40)u_1^6 + \\ &\quad + \frac{9}{8} \left(c_1 + \frac{1}{3}\right) \left(c_1 + \frac{2}{3}\right) u_2 \left(c_1 + \frac{1}{2}\right) u_1^4 - \frac{9}{8} \left(c_1 + \frac{1}{3}\right) \left(c_1 + \frac{2}{3}\right) u_2^2u_1^2 + \\ &\quad \frac{1}{2160}(405c_1 + 270)u_2^3 \end{aligned}$$

Finally, for $c_1 = -\frac{1}{2}$ one recovers the Frobenius structure listed in Table 1.

Remark 7.1 *In all the cases we dealt with in this Section, the dual connection has the general form*

$$\Gamma_{jk}^{(2)i} = -c_{jk}^{*i} + \lambda C_{jk}^i,$$

where C_{jk}^i are the Christoffel symbol of a Dunkl-Kohno-type connection

$$\nabla + \frac{1}{N} \sum_{H \in \mathcal{H}} \frac{d\alpha_H}{\alpha_H} \otimes \kappa_H \pi_H. \quad (7.2)$$

where the projectors π_H and the weights κ_H satisfy the condition

$$\sum_{H \in \mathcal{H}} \kappa_H \pi_H = 0. \quad (7.3)$$

This follows from the fact that the dual connection has to satisfy the condition $\nabla^{(2)}E = 0$. Moreover the collection $\{\kappa_H\}_{H \in \mathcal{H}}$ of complex weights define a G -invariant function. It

would be interesting to prove that in general the dual structure is always defined by the standard dual product for a suitable choice of the hermtian metric and by a dual connection of the form above. Indeed this would have as an immediate consequence the existence of an upper bound for the number of parameters: taking into account the condition (7.3), the number of parameters should be strictly less than the number of G -orbits in \mathcal{H} .

7.12 The bi-flat F -structures of associated Coxeter groups

The Frobenius manifold structure on the space of orbits of Shephard groups coincides with the Frobenius structure on the space of orbits of the associated Coxeter group. We prove a similar theorem for the bi-flat F -structures of the exceptional Shephard groups $G_5, G_6, G_9, G_{10}, G_{14}, G_{17}, G_{18}, G_{21}, G_{26}$. In other words, the Frobenius structure of each orbit space of these groups sits inside a one-parameter family of bi-flat F -manifold structures, which depends only on the associated Coxeter groups and can be viewed as a natural deformation of the Frobenius structure itself.

Theorem 7.2 *The bi-flat F -structure on the space of orbits of each exceptional Shephard groups of rank 2 and 3 coincides with the bi-flat F -structure on the space of orbits of the associated Coxeter group.*

Proof: The proof is not trivial only for the groups $G_5, G_6, G_9, G_{10}, G_{14}, G_{17}, G_{18}, G_{21}, G_{26}$.

For the groups G_5, G_{10}, G_{18} we have to prove that the vector potential coincides with the vector potential of B_2 . The generalized Saito flat coordinates are $u_1 = p_1^2 + p_2^2$ and $u_2 = p_1^4 + p_2^4 + c_1 u_1^2$ and in these coordinates we obtain:

$$\begin{aligned} A_{B_2}^1 &= -\left(\frac{2}{3}c_1 + \frac{1}{2}\right)u_1^3 + u_1u_2, \\ A_{B_2}^2 &= -\left(\frac{1}{3}c_1^2 + \frac{1}{2}c_1 + 1\right)u_1^4 + \frac{1}{2}u_2^2. \end{aligned}$$

These coincides with the vector potential of G_5, G_{10}, G_{18} up to rescaling of the variables.

For the groups G_6, G_9, G_{17} we have to prove that the vector potential coincides with the vector potential of $I_2(6)$. In the generalized Saito flat coordinates

$$\begin{aligned} u_1 &= p_1^2 + p_2^2, \\ u_2 &= 2p_1^6 - 30p_1^4p_2^2 + 30p_1^2p_2^4 - 2p_2^6 + c_1u_1^3 \end{aligned}$$

we obtain

$$\begin{aligned} A_{I_2(6)}^1 &= -\left(\frac{2}{3}c_1 + \frac{1}{2}\right)u_1^3 + u_1u_2, \\ A_{I_2(6)}^2 &= -\left(\frac{1}{3}c_1^2 + \frac{1}{2}c_1 + 1\right)u_1^4 + \frac{1}{2}u_2^2, \end{aligned}$$

that coincides with the vector potential of G_6, G_9, G_{17} up to rescaling of the variables.

For the group G_{14} we have to prove that the vector potential coincides with the vector potential of $I_2(8)$. In the generalized Saito flat coordinates

$$\begin{aligned} u_1 &= p_1^2 + p_2^2, \\ u_2 &= 2p_1^8 - 56p_1^6p_2^2 + 140p_1^4p_2^4 - 56p_1^2p_2^6 + 2p_2^8 + c_1u_1^4, \end{aligned}$$

we get

$$\begin{aligned} A_{I_2(8)}^1 &= -\frac{2}{5}c_1u_1^5 + u_1u_2, \\ A_{I_2(8)}^2 &= -\frac{2}{7}u_1^8c_1^2 + \frac{8}{7}u_1^8 + \frac{1}{2}u_2^2, \end{aligned}$$

that coincides with the vector potential of G_{14} up to rescaling of the variables.

For the group G_{21} we have to prove that the vector potential coincides with the vector potential of $I_2(10)$. In the generalized Saito flat coordinates

$$\begin{aligned} u_1 &= p_1^2 + p_2^2, \\ u_2 &= 2p_1^{10} - 90p_1^8p_2^2 + 420p_1^6p_2^4 - 420p_1^4p_2^6 + 90p_1^2p_2^8 - 2p_2^{10} + c_1u_1^5, \end{aligned}$$

we obtain

$$\begin{aligned} A_{I_2(10)}^1 &= -\left(\frac{2}{3}c_1 + \frac{1}{2}\right)u_1^3 + u_1u_2, \\ A_{I_2(10)}^2 &= -\left(\frac{1}{3}c_1^2 + \frac{1}{2}c_1 + 1\right)u_1^4 + \frac{1}{2}u_2^2, \end{aligned}$$

that coincides with the vector potential of G_{21} up to rescaling of the variables.

For the group G_{26} we have to prove that the vector potential coincides with the vector potential of B_3 . The generalized Saito flat coordinates are

$$\begin{aligned} u_1 &= p_1^6 + p_2^6 + p_3^6, \\ u_2 &= p_1^4 + p_2^4 + p_3^4 + c_1(p_1^2 + p_2^2 + p_3^2)^2, \\ u_3 &= p_1^6 + p_2^6 + p_3^6 + c_2(p_1^2 + p_2^2 + p_3^2)(p_1^4 + p_2^4 + p_3^4) + c_3(p_1^2 + p_2^2 + p_3^2)^3, \end{aligned}$$

with $c_1 = \frac{1}{3} + \frac{2}{3}c_2$ and $c_3 = \frac{2}{9}c_2^2$. After some computations we get

$$\begin{aligned} A_{B_3}^1 &= \frac{1}{72}(24c_2^2 + 62c_2 + 40)u_1^4 + \frac{1}{72}(-72c_2 - 90)u_2u_1^2 + u_1u_3 + \frac{3}{8}u_2^2, \\ A_{B_3}^2 &= \frac{1}{90}(8c_2^3 + 34c_2^2 + 46c_2 + 20)u_1^5 - \frac{1}{9}(c_2 + 1)u_2u_1^3 + \frac{1}{90}(-45c_2 - 45)u_2^2u_1 + u_2u_3, \\ A_{B_3}^3 &= \frac{1}{2}u_3^2 + \frac{1}{2160}(-128c_2^4 - 640c_2^3 - 1200c_2^2 - 1000c_2 - 312)u_1^6 + \frac{1}{3}\left(c_2 + \frac{5}{4}\right)(c_2 + 1) + \\ &\quad \left(c_2 + \frac{3}{2}\right)u_2u_1^4 - \frac{1}{2}(c_2 + 1)\left(c_2 + \frac{3}{2}\right)u_2^2u_1^2 + \frac{1}{2160}(270c_2 + 405)u_2^3, \end{aligned}$$

that coincides with the vector potential of G_6, G_9, G_{17} up to rescaling of the variables. \blacksquare

Remark 7.3 *The dual connection has the form*

$$\Gamma_{jk}^{(2)i} = -c_{jk}^{*i} + \lambda C_{jk}^i,$$

where

$$c_{lp}^{*i}(u) = \frac{2}{N} \left(\sum_{s=1}^n \frac{1}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)} \right)$$

and

$$C_{lp}^i(u) = \sum_{s=1}^k \frac{\kappa_s}{\|\alpha_s\|^2} \frac{(\alpha_s)_l (\alpha_s)_p (\check{\alpha}_s)^i}{\alpha_s(u)}.$$

In the case of B_2 we have $N = 4, n = 4, \kappa_s = 1$ if $s = 1, 2, \kappa_s = -1$ if $s = 3, 4$ and $\lambda = 2c_1 + \frac{3}{2}$.

In the case of $I_2(6)$ we have $N = 6, n = 6, \kappa_s = 1$ if $s = 1, 4, 6, \kappa_s = -1$ if $s = 2, 3, 5$ and $\lambda = -\frac{1}{3}c_1$.

In the case of $I_2(8)$ we have $N = 8, n = 8, \kappa_s = 1$ if $s = 1, 2, 3, 4, \kappa_s = -1$ if $s = 5, 6, 7, 8$ and $\lambda = -\frac{3}{8}c_1$.

In the case of $I_2(10)$ we have $N = 10, n = 10, \kappa_s = 1$ if $s = 1, 7, 8, 9, 10, \kappa_s = -1$ if $s = 2, 3, 4, 5, 6$ and $\lambda = \frac{2}{5}c_1$.

In the case of B_3 we have $N = 6, n = 9, \kappa_s = 2$ if $s = 1, 2, 3, \kappa_s = -1$ if $s = 4, 5, 6, 7, 8, 9$ and $\lambda = 2c_1 + \frac{3}{2}$.

In all cases the metric is the standard hermitean metric. The list of α is given in the Appendix 2.

8 Bi-flat F -manifolds and generalized WDVV equations

In this Section we prove that under a mild assumption semisimple bi-flat F -manifold are described in flat coordinates for the natural connection by solutions of gener-

alized WDVV equations

$$\partial_j \partial_l A^i \partial_k \partial_m A^l = \partial_k \partial_l A^i \partial_k \partial_m A^l, \quad (8.1)$$

$$\partial_1 \partial_i A^j = \delta_i^j, \quad (8.2)$$

$$E(A^i) = (1 + w_i)A^i, \quad (8.3)$$

where $w_1 = 1$. This system of equations was obtained recently by Kato, Mano and Sekiguchi in the study of Saito structures without metrics [20].

Comparing equations (8.1), (8.2), (8.3) with (4.1) and (4.2), we see that we have only the extra condition (8.3). We want to show that this extra condition is related to the presence of a second compatible flat structure.

We have seen that bi-flat F -manifolds, in the semisimple case, are equivalent to Frobenius manifolds without metric. This means that one can replace the conditions involving the dual connection with the condition

$$\nabla^{(1)} \nabla^{(1)} E = 0.$$

Assuming $e = \frac{\partial}{\partial t^1}$ (this is not restrictive) and E of the form

$$E = \sum_i w^i t^i \frac{\partial}{\partial t^i}, \quad (8.4)$$

where (t^1, \dots, t^n) are flat coordinates (this is not too restrictive if $\nabla^{(1)} E$ is semisimple) and imposing the conditions on the Euler vector field (see Definition 4.5):

$$[e, E] = e, \quad (\text{Lie}_{EC})_{jk}^i = c_{jk}^i,$$

we obtain $w_1 = 1$ and

$$\begin{aligned} (\text{Lie}_{EC})_{jk}^i &= E^l \partial_l c_{jk}^i - (\partial_l E^i) c_{jk}^l + (\partial_j E^l) c_{lk}^i + (\partial_k E^l) c_{jl}^i = \\ &= E^l \partial_l c_{jk}^i - w_i c_{jk}^i + w_j c_{jk}^i + w_k c_{jk}^i = c_{jk}^i. \end{aligned}$$

From this relation we have

$$E(c_{jk}^i) = (1 + w_i - w_j - w_k) c_{jk}^i.$$

Writing c_{jk}^i in terms of derivative of the vector potential, we obtain

$$E^l \partial_l \partial_j \partial_k A^i = (1 + w_i - w_j - w_k) \partial_j \partial_k A^i.$$

It is immediate to check that

$$\partial_j \partial_k (E(A^i)) = w_k c_{jk}^i + w_j c_{jk}^i + E(c_{jk}^i),$$

and substituting in the previous expression we find

$$\partial_j \partial_k E(A^i) = (1 + w_i) \partial_j \partial_k A^i,$$

or

$$\partial_j \partial_k (E(A^i) - (1 + w_i) A^i) = 0,$$

which is exactly equation (8.3) up to affine terms in t^i . But since the vector potential itself is defined up to affine terms in t^i we are done.

All the vector potentials obtained in this paper satisfy by construction the generalized WDVV equations (8.1), (8.2) and (8.3) where in (8.2) one has to replace 1 with n (since the unit vector field is $\frac{\partial}{\partial u_n}$) and in (8.3) one has to replace E with the normalized Euler vector field $\frac{E}{d_n}$ and, as a consequence, $w_i = \frac{d_i}{d_n}$. Vector potentials associated with rank > 2 well-generated exceptional complex reflection groups have been found in [19] (with the exception of Shephard groups). The results of the present Section, combined with the results of [20] imply the existence of a bi-flat structure for any well-generated complex reflection group. In particular, given the flat structure (∇, \circ, e) in flat coordinates (u^1, \dots, u^n) one can immediately reconstruct the dual structure using the standard formula for the dual product and the formula (5.5). In the present paper we have adopted a “dual” approach starting from the dual structure that has a universal beautiful form in the coordinates (p^1, \dots, p^n) . This approach leads naturally to discover the presence of parameters in several examples. As far as we know, the presence of these parameters has never been observed before. The results of the present Section also prove that the relation between generalized WDVV equations and the full family of Painlevé VI equation (in standard form) obtained in [20] is the counterpart in flat coordinates of the relation between three dimensional semisimple bi-flat F -manifolds and the full family of Painlevé VI equation (in sigma form) obtained in canonical coordinates in [28, 4] (see also [2]).

9 Appendix 1. Reflecting mirrors

We give the list, case by case, of the covectors α_s obtained factorizing $\det \frac{\partial u^i}{\partial p^j}$.

9.1 The G_4 case

We have 4 mirrors defining reflections of order 3:

$$\begin{aligned} \alpha_1 &= \left[-1 - i, \sqrt{3} - 1 \right], & \alpha_2 &= \left[1 - i, \sqrt{3} + 1 \right], \\ \alpha_3 &= \left[-1 + i, \sqrt{3} + 1 \right], & \alpha_4 &= \left[1 + i, \sqrt{3} - 1 \right]. \end{aligned}$$

9.2 The G_5 case

We have 8 mirrors defining reflections of order 3:

$$\begin{aligned}\alpha_1 &= \left[1, -\frac{1}{2}\sqrt{3} + \frac{1}{2} + \frac{1}{2}i - \frac{1}{2}i\sqrt{3}\right], & \alpha_2 &= \left[1, \frac{1}{2}\sqrt{3} + \frac{1}{2} - \frac{1}{2}i - \frac{1}{2}i\sqrt{3}\right], \\ \alpha_3 &= \left[1, \frac{1}{2}\sqrt{3} - \frac{1}{2} - \frac{1}{2}i + \frac{1}{2}i\sqrt{3}\right], & \alpha_4 &= \left[1, -\frac{1}{2}\sqrt{3} - \frac{1}{2} + \frac{1}{2}i + \frac{1}{2}i\sqrt{3}\right], \\ \alpha_5 &= \left[1, \frac{1}{2}\sqrt{3} + \frac{1}{2} + \frac{1}{2}i + \frac{1}{2}i\sqrt{3}\right], & \alpha_6 &= \left[1, -\frac{1}{2}\sqrt{3} + \frac{1}{2} - \frac{1}{2}i + \frac{1}{2}i\sqrt{3}\right], \\ \alpha_7 &= \left[1, \frac{1}{2}\sqrt{3} - \frac{1}{2} + \frac{1}{2}i - \frac{1}{2}i\sqrt{3}\right], & \alpha_8 &= \left[1, -\frac{1}{2}\sqrt{3} - \frac{1}{2} - \frac{1}{2}i - \frac{1}{2}i\sqrt{3}\right].\end{aligned}$$

9.3 The G_6 case

We have 6 mirrors defining reflections of order 2:

$$\alpha_1 = [1, -1], \alpha_2 = [1, -i], \alpha_3 = [1, i], \alpha_4 = [1, 1], \alpha_5 = [1, 0], \alpha_6 = [0, 1],$$

and 4 mirrors defining reflections of order 3:

$$\begin{aligned}\alpha_7 &= [-1 - i, \sqrt{3} - 1], & \alpha_8 &= [-1 + i, \sqrt{3} + 1], \\ \alpha_9 &= [1 + i, \sqrt{3} - 1], & \alpha_{10} &= [1 - i, \sqrt{3} + 1].\end{aligned}$$

9.4 The G_8 case

We have 6 mirrors defining reflections of order 4:

$$\alpha_1 = [1, 1], \alpha_2 = [1, -1], \alpha_3 = [0, 1], \alpha_4 = [1, -i], \alpha_5 = [1, i], \alpha_6 = [1, 0].$$

9.5 The G_9 case

We have 12 mirrors defining reflections of order 2:

$$\begin{aligned}
 \alpha_1 &= \left[1, \frac{1}{2}\sqrt{2} - \frac{1}{2}i\sqrt{2}\right], & \alpha_2 &= \left[1, \frac{1}{2}\sqrt{2} + \frac{1}{2}i\sqrt{2}\right], \\
 \alpha_3 &= \left[1, -\frac{1}{2}\sqrt{2} - \frac{1}{2}i\sqrt{2}\right], & \alpha_4 &= \left[1, -\frac{1}{2}\sqrt{2} + \frac{1}{2}i\sqrt{2}\right], \\
 \alpha_5 &= [1, -\sqrt{2} + 1], & \alpha_6 &= [1, \sqrt{2} + 1], \\
 \alpha_7 &= [1, i\sqrt{2} + i], & \alpha_8 &= [1, -i\sqrt{2} - i], \\
 \alpha_9 &= [1, i\sqrt{2} - i], & \alpha_{10} &= [1, -i\sqrt{2} + i], \\
 \alpha_{11} &= [1, \sqrt{2} - 1], & \alpha_{12} &= [1, -\sqrt{2} - 1],
 \end{aligned}$$

and 6 mirrors defining reflections of order 4:

$$\alpha_{13} = [1, 1], \alpha_{14} = [0, 1], \alpha_{15} = [1, -1], \alpha_{16} = [1, i], \alpha_{17} = [1, -i], \alpha_{18} = [1, 0].$$

9.6 The G_{10} case

We have 8 mirrors defining reflections of order 3:

$$\begin{aligned}
 \alpha_1 &= \left[1, -\frac{1}{2}i\sqrt{3} - \frac{1}{2}\sqrt{2 + 2i\sqrt{3}} - \frac{1}{2}\right], \\
 \alpha_2 &= \left[1, \frac{1}{2}i\sqrt{3} + \frac{1}{4}\sqrt{2 + 2i\sqrt{3}} - \frac{1}{2} - \frac{1}{4}i\sqrt{3}\sqrt{2 + 2i\sqrt{3}}\right], \\
 \alpha_3 &= \left[1, \frac{1}{2}i\sqrt{3} - \frac{1}{4}\sqrt{2 + 2i\sqrt{3}} - \frac{1}{2} + \frac{1}{4}i\sqrt{3}\sqrt{2 + 2i\sqrt{3}}\right], \\
 \alpha_4 &= \left[1, -\frac{1}{2}i\sqrt{3} + \frac{1}{2}\sqrt{2 + 2i\sqrt{3}} - \frac{1}{2}\right], \\
 \alpha_5 &= \left[1, \frac{1}{2}i\sqrt{3} + \frac{1}{2}\sqrt{2 + 2i\sqrt{3}} + \frac{1}{2}\right], \\
 \alpha_6 &= \left[1, -\frac{1}{2}i\sqrt{3} - \frac{1}{4}\sqrt{2 + 2i\sqrt{3}} + \frac{1}{2} + \frac{1}{4}i\sqrt{3}\sqrt{2 + 2i\sqrt{3}}\right], \\
 \alpha_7 &= \left[1, -\frac{1}{2}i\sqrt{3} + \frac{1}{4}\sqrt{2 + 2i\sqrt{3}} + \frac{1}{2} - \frac{1}{4}i\sqrt{3}\sqrt{2 + 2i\sqrt{3}}\right], \\
 \alpha_8 &= \left[1, \frac{1}{2}i\sqrt{3} - \frac{1}{2}\sqrt{2 + 2i\sqrt{3}} + \frac{1}{2}\right],
 \end{aligned}$$

and 6 mirrors defining reflections of order 4:

$$\begin{aligned}\alpha_9 &= [1, 1], \alpha_{10} = [1, 0], \alpha_{11} = [1, -1], \alpha_{12} = [0, 1], \\ \alpha_{13} &= \left[1, -\frac{1}{4}\sqrt{2+2i\sqrt{3}} - \frac{1}{4}i\sqrt{3}\sqrt{2+2i\sqrt{3}}\right], \\ \alpha_{14} &= \left[1, \frac{1}{4}\sqrt{2+2i\sqrt{3}} + \frac{1}{4}i\sqrt{3}\sqrt{2+2i\sqrt{3}}\right].\end{aligned}$$

9.7 The G_{14} case

We have 12 mirrors defining reflections of order 2:

$$\begin{aligned}\alpha_1 &= \left[1, -\frac{1}{2}\sqrt{2} + \frac{1}{2}i\sqrt{2}\right], & \alpha_2 &= \left[1, -\frac{1}{2}\sqrt{2} - \frac{1}{2}i\sqrt{2}\right], \\ \alpha_3 &= \left[1, \frac{1}{2}\sqrt{2} - \frac{1}{2}i\sqrt{2}\right], & \alpha_4 &= \left[1, \frac{1}{2}\sqrt{2} + \frac{1}{2}i\sqrt{2}\right], \\ \alpha_5 &= [1, \sqrt{2} - 1], & \alpha_6 &= [1, -\sqrt{2} - 1], \\ \alpha_7 &= [1, -i\sqrt{2} + i], & \alpha_8 &= [1, i\sqrt{2} - i], \\ \alpha_9 &= [1, -i\sqrt{2} - i], & \alpha_{10} &= [1, i\sqrt{2} + i], \\ \alpha_{11} &= [1, -\sqrt{2} + 1], & \alpha_{12} &= [1, \sqrt{2} + 1],\end{aligned}$$

and 8 mirrors defining reflections of order 3:

$$\begin{aligned}\alpha_{13} &= \left[1, -\frac{1}{2} + \frac{1}{2}i + \frac{1}{2}\sqrt{3} - \frac{1}{2}i\sqrt{3}\right], & \alpha_{14} &= \left[1, -\frac{1}{2} + \frac{1}{2}i - \frac{1}{2}\sqrt{3} + \frac{1}{2}i\sqrt{3}\right], \\ \alpha_{15} &= \left[1, -\frac{1}{2}\sqrt{3} - \frac{1}{2}i\sqrt{3} - \frac{1}{2} - \frac{1}{2}i\right], & \alpha_{16} &= \left[1, \frac{1}{2}\sqrt{3} + \frac{1}{2}i\sqrt{3} - \frac{1}{2} - \frac{1}{2}i\right], \\ \alpha_{17} &= \left[1, \frac{1}{2}\sqrt{3} + \frac{1}{2}i\sqrt{3} + \frac{1}{2} + \frac{1}{2}i\right], & \alpha_{18} &= \left[1, -\frac{1}{2}\sqrt{3} - \frac{1}{2}i\sqrt{3} + \frac{1}{2} + \frac{1}{2}i\right], \\ \alpha_{19} &= \left[1, \frac{1}{2} - \frac{1}{2}i + \frac{1}{2}\sqrt{3} - \frac{1}{2}i\sqrt{3}\right], & \alpha_{20} &= \left[1, \frac{1}{2} - \frac{1}{2}i - \frac{1}{2}\sqrt{3} + \frac{1}{2}i\sqrt{3}\right],\end{aligned}$$

9.8 The G_{16} case

We have 12 mirrors defining reflections of order 2:

$$\begin{aligned}
\alpha_1 &= \left[1, -\frac{1}{10}\mu - \frac{1}{10}\sigma \right], & \alpha_2 &= \left[1, -\left(\frac{1}{10}\mu + \frac{1}{10}\sigma\right)^3 - \left(\frac{1}{25}\mu + \frac{1}{25}\sigma\right)\sqrt{5} \right], \\
\alpha_3 &= \left[1, \left(\frac{1}{25}\mu + \frac{1}{25}\sigma\right)\sqrt{5} + \left(\frac{1}{10}\mu + \frac{1}{10}\sigma\right)^3 \right], & \alpha_4 &= \left[1, \frac{1}{10}\mu + \frac{1}{10}\sigma \right], \\
\alpha_5 &= \left[1, -\left(-\frac{1}{2}i - \frac{1}{2}i\sqrt{5} + \frac{1}{2}\nu\right)^3 - (-i - i\sqrt{5} + \nu)\sqrt{5} + 4i + 4i\sqrt{5} - 4\nu \right], \\
\alpha_6 &= \left[1, \left(-\frac{1}{2}i - \frac{1}{2}i\sqrt{5} + \frac{1}{2}\nu\right)^3 + (-i - i\sqrt{5} + \nu)\sqrt{5} - 4i - 4i\sqrt{5} + 4\nu \right], \\
\alpha_7 &= \left[1, \frac{1}{2}i + \frac{1}{2}i\sqrt{5} - \frac{1}{2}\nu \right], & \alpha_8 &= \left[1, -\frac{1}{2}i - \frac{1}{10}i\sqrt{5} + \frac{1}{2}\nu \right], \\
\alpha_9 &= \left[1, \frac{1}{2}\sqrt{5} - \frac{1}{2} - \frac{1}{2}\lambda \right], & \alpha_{10} &= \left[1, \frac{1}{2}\sqrt{5} - \frac{1}{2} + \frac{1}{2}\lambda \right], \\
\alpha_{11} &= \left[1, -\frac{1}{2}\sqrt{5} + \frac{1}{2} - \frac{1}{2}\lambda \right], & \alpha_{12} &= \left[1, -\frac{1}{2}\sqrt{5} + \frac{1}{2} + \frac{1}{2}\lambda \right],
\end{aligned}$$

with

$$\nu = \sqrt{-10 - 2\sqrt{5}}, \quad \lambda = \sqrt{10 - 2\sqrt{5}}, \quad \sigma = \sqrt{5}\nu, \mu = \sqrt{5}\lambda.$$

9.9 The G_{17} case

We have 30 mirrors defining reflections of order 2:

$$\begin{aligned}
\alpha_1 &= \left[1, \frac{1}{2}\sqrt{5} - \frac{1}{2} + \frac{1}{2}i + \frac{1}{2}i\sqrt{5}\right], & \alpha_2 &= \left[1, \frac{1}{2}\sqrt{5} - \frac{1}{2} - \frac{1}{2}i - \frac{1}{2}i\sqrt{5}\right], \\
\alpha_3 &= \left[1, -\frac{1}{2}\sqrt{5} + \frac{1}{2} + \frac{3}{2}i - \frac{1}{2}i\sqrt{5}\right], & \alpha_4 &= \left[1, -\frac{1}{2}\sqrt{5} + \frac{1}{2} - \frac{3}{2}i + \frac{1}{2}i\sqrt{5}\right], \\
\alpha_5 &= \left[1, \frac{1}{2}\sqrt{5} - \frac{1}{2} + \frac{3}{2}i - \frac{1}{2}i\sqrt{5}\right], & \alpha_6 &= \left[1, \frac{1}{2}\sqrt{5} - \frac{1}{2} - \frac{3}{2}i + \frac{1}{2}i\sqrt{5}\right], \\
\alpha_7 &= [1, -i], & \alpha_8 &= [1, i], \\
\alpha_9 &= \left[1, \frac{1}{10}\sqrt{5} - \frac{1}{2} - \frac{1}{2}i + \frac{3}{10}i\sqrt{5}\right], & \alpha_{10} &= \left[1, \frac{1}{10}\sqrt{5} - \frac{1}{2} + \frac{1}{2}i - \frac{3}{10}i\sqrt{5}\right], \\
\alpha_{11} &= \left[1, -\frac{1}{6}\sqrt{5} + \frac{1}{6} + \frac{1}{6}i + \frac{1}{6}i\sqrt{5}\right], & \alpha_{12} &= \left[1, -\frac{1}{6}\sqrt{5} + \frac{1}{6} - \frac{1}{6}i - \frac{1}{6}i\sqrt{5}\right], \\
\alpha_{13} &= \left[1, -\frac{1}{2}\sqrt{5} - \frac{3}{2} - \frac{1}{2}i - \frac{1}{2}i\sqrt{5}\right], & \alpha_{14} &= \left[1, -\frac{1}{2}\sqrt{5} - \frac{3}{2} + \frac{1}{2}i + \frac{1}{2}i\sqrt{5}\right], \\
\alpha_{15} &= \left[1, -\frac{3}{10}\sqrt{5} - \frac{1}{2} - \frac{1}{2}i - \frac{1}{10}i\sqrt{5}\right], & \alpha_{16} &= \left[1, -\frac{3}{10}\sqrt{5} - \frac{1}{2} + \frac{1}{2}i + \frac{1}{10}i\sqrt{5}\right], \\
\alpha_{17} &= \left[1, \frac{3}{10}\sqrt{5} + \frac{1}{2} - \frac{1}{2}i - \frac{1}{10}i\sqrt{5}\right], & \alpha_{18} &= \left[1, \frac{3}{10}\sqrt{5} + \frac{1}{2} + \frac{1}{2}i + \frac{1}{10}i\sqrt{5}\right], \\
\alpha_{19} &= [0, 1], & \alpha_{20} &= [1, -1], \\
\alpha_{21} &= \left[1, \frac{1}{2}\sqrt{5} + \frac{3}{2} + \frac{1}{2}i + \frac{1}{2}i\sqrt{5}\right], & \alpha_{22} &= \left[1, \frac{1}{2}\sqrt{5} + \frac{3}{2} - \frac{1}{2}i - \frac{1}{2}i\sqrt{5}\right], \\
\alpha_{23} &= \left[1, -\frac{1}{10}\sqrt{5} + \frac{1}{2} + \frac{1}{2}i - \frac{3}{10}i\sqrt{5}\right], & \alpha_{24} &= \left[1, -\frac{1}{10}\sqrt{5} + \frac{1}{2} - \frac{1}{2}i + \frac{3}{10}i\sqrt{5}\right], \\
\alpha_{25} &= [1, 1], & \alpha_{26} &= \left[1, -\frac{1}{2}\sqrt{5} + \frac{1}{2} - \frac{1}{2}i - \frac{1}{2}i\sqrt{5}\right], \\
\alpha_{27} &= \left[1, -\frac{1}{2}\sqrt{5} + \frac{1}{2} + \frac{1}{2}i + \frac{1}{2}i\sqrt{5}\right], & \alpha_{28} &= \left[1, \frac{1}{6}\sqrt{5} - \frac{1}{6} - \frac{1}{6}i - \frac{1}{6}i\sqrt{5}\right], \\
\alpha_{29} &= \left[1, \frac{1}{6}\sqrt{5} - \frac{1}{6} + \frac{1}{6}i + \frac{1}{6}i\sqrt{5}\right], & \alpha_{30} &= [1, 0],
\end{aligned}$$

and 12 mirrors defining reflections of order 5:

$$\begin{aligned}
\alpha_{31} &= \left[1, \frac{1}{2}\sqrt{5} - \frac{1}{2} - \frac{1}{2}\sigma \right], & \alpha_{32} &= \left[1, \frac{1}{2}\sqrt{5} - \frac{1}{2} + \frac{1}{2}\sigma \right], \\
\alpha_{33} &= \left[1, -\frac{1}{2}\sqrt{5} + \frac{1}{2} + \frac{1}{2}\sigma \right], & \alpha_{34} &= \left[1, -\frac{1}{2}\sqrt{5} + \frac{1}{2} - \frac{1}{2}\sigma \right], \\
\alpha_{35} &= \left[1, \frac{1}{10}\mu + \frac{1}{10}\nu \right], & \alpha_{36} &= \left[1, -\left(\frac{1}{25}\mu + \frac{1}{25}\nu\right)\sqrt{5} - \left(\frac{1}{10}\mu + \frac{1}{10}\nu\right)^3 \right], \\
\alpha_{37} &= \left[1, -\frac{1}{10}\mu - \frac{1}{10}\nu \right], & \alpha_{38} &= \left[1, \left(\frac{1}{10}\mu + \frac{1}{10}\nu\right)^3 + \left(\frac{1}{25}\mu + \frac{1}{25}\nu\right)\sqrt{5} \right], \\
\alpha_{39} &= \left[1, \frac{1}{2}i + \frac{1}{2}i\sqrt{5} - \frac{1}{2}\lambda \right], & \alpha_{42} &= \left[1, -\frac{1}{2}i - \frac{1}{2}i\sqrt{5} + \frac{1}{2}\lambda \right], \\
\alpha_{40} &= \left[1, \left(-\frac{1}{2}i - \frac{1}{2}i\sqrt{5} + \frac{1}{2}\lambda\right)^3 + (-i - i\sqrt{5} + \lambda)\sqrt{5} - 4i - 4i\sqrt{5} + 4\lambda \right], \\
\alpha_{41} &= \left[1, -\left(-\frac{1}{2}i - \frac{1}{2}i\sqrt{5} + \frac{1}{2}\lambda\right)^3 - (-i - i\sqrt{5} + \lambda)\sqrt{5} + 4i + 4i\sqrt{5} - 4\lambda \right],
\end{aligned}$$

with

$$\sigma = \sqrt{10 - 2\sqrt{5}}, \quad \lambda = \sqrt{-10 - 2\sqrt{5}}, \quad \mu = \sqrt{5}\sigma, \quad \nu = \sqrt{5}\lambda.$$

9.10 The G_{18} case

We have 20 mirrors defining reflections of order 3:

$$\begin{aligned}
\alpha_1 &= \left[1, -\frac{1}{2}\sqrt{3} + \frac{1}{2} - \frac{1}{2}i + \frac{1}{2}i\sqrt{3} \right], & \alpha_2 &= \left[1, \frac{1}{2}\sqrt{3} + \frac{1}{2} - \frac{1}{2}i - \frac{1}{2}i\sqrt{3} \right], \\
\alpha_3 &= \left[1, -\frac{1}{2}\sqrt{3} + \frac{1}{2} + \frac{1}{2}i - \frac{1}{2}i\sqrt{3} \right], & \alpha_4 &= \left[1, \frac{1}{2}\sqrt{3} + \frac{1}{2} + \frac{1}{2}i + \frac{1}{2}i\sqrt{3} \right], \\
\alpha_5 &= \left[1, \frac{1}{2}\sqrt{3} - \frac{1}{2} - \frac{1}{2}i + \frac{1}{2}i\sqrt{3} \right], & \alpha_6 &= \left[1, -\frac{1}{2}\sqrt{3} - \frac{1}{2} - \frac{1}{2}i - \frac{1}{2}i\sqrt{3} \right], \\
\alpha_7 &= \left[1, -\frac{1}{2}\sqrt{3} - \frac{1}{2} + \frac{1}{2}i + \frac{1}{2}i\sqrt{3} \right], & \alpha_8 &= \left[1, \frac{1}{2}\sqrt{3} - \frac{1}{2} + \frac{1}{2}i - \frac{1}{2}i\sqrt{3} \right], \\
\alpha_9 &= \left[1, \frac{1}{2}\sqrt{3} + \frac{1}{2}\sqrt{5}\sqrt{3} - \frac{1}{2}\sqrt{5} - \frac{3}{2} \right], & \alpha_{10} &= \left[1, -\frac{1}{2}\sqrt{3} - \frac{1}{2}\sqrt{5}\sqrt{3} - \frac{1}{2}\sqrt{5} - \frac{3}{2} \right], \\
\alpha_{11} &= \left[1, -\frac{1}{2}i(\sqrt{5}\sqrt{3} - \sqrt{3} + \sqrt{5} - 3) \right], & \alpha_{12} &= \left[1, \frac{1}{2}i(\sqrt{5}\sqrt{3} - \sqrt{3} - \sqrt{5} + 3) \right], \\
\alpha_{13} &= \left[1, \frac{1}{2}i(\sqrt{5}\sqrt{3} - \sqrt{3} + \sqrt{5} - 3) \right], & \alpha_{14} &= \left[1, -\frac{1}{2}i(\sqrt{5}\sqrt{3} - \sqrt{3} - \sqrt{5} + 3) \right], \\
\alpha_{15} &= \left[1, \frac{3}{2} - \frac{1}{2}\sqrt{5}\sqrt{3} - \frac{1}{2}\sqrt{3} + \frac{1}{2}\sqrt{5} \right], & \alpha_{16} &= \left[1, \frac{1}{2}\sqrt{3} + \frac{1}{2}\sqrt{5}\sqrt{3} + \frac{1}{2}\sqrt{5} + \frac{3}{2} \right], \\
\alpha_{17} &= \left[1, -\frac{1}{6}\sqrt{5}\sqrt{3} - \frac{1}{6}\sqrt{3} - \frac{1}{6}i\sqrt{3} + \frac{1}{6}i\sqrt{3}\sqrt{5} \right], & \alpha_{18} &= \left[1, -\frac{1}{6}\sqrt{5}\sqrt{3} - \frac{1}{6}\sqrt{3} + \frac{1}{6}i\sqrt{3} - \frac{1}{6}i\sqrt{3}\sqrt{5} \right], \\
\alpha_{19} &= \left[1, \frac{1}{6}\sqrt{5}\sqrt{3} + \frac{1}{6}\sqrt{3} + \frac{1}{6}i\sqrt{3} - \frac{1}{6}i\sqrt{3}\sqrt{5} \right], & \alpha_{20} &= \left[1, \frac{1}{6}\sqrt{5}\sqrt{3} + \frac{1}{6}\sqrt{3} - \frac{1}{6}i\sqrt{3} + \frac{1}{6}i\sqrt{3}\sqrt{5} \right],
\end{aligned}$$

and 12 hyperplanes defining reflections of order 5:

$$\begin{aligned}
\alpha_{21} &= \left[1, \frac{1}{20}i\sqrt{5}\mu + \frac{1}{4}i\mu + \frac{1}{10}\sqrt{5}\mu\right], & \alpha_{22} &= \left[1, -\frac{1}{20}i\sqrt{5}\mu - \frac{1}{4}i\mu - \frac{1}{10}\sqrt{5}\mu\right], \\
\alpha_{23} &= \left[1, -\frac{1}{20}i\sqrt{5}\mu - \frac{1}{4}i\mu + \frac{1}{10}\sqrt{5}\mu\right], & \alpha_{24} &= \left[1, \frac{1}{20}i\sqrt{5}\mu + \frac{1}{4}i\mu - \frac{1}{10}\sqrt{5}\mu\right], \\
\alpha_{25} &= \left[1, -\frac{1}{2}\sqrt{5} + \frac{1}{2}\mu + \frac{1}{2}\right], & \alpha_{26} &= \left[1, -\frac{1}{2}\sqrt{5} - \frac{1}{2}\mu + \frac{1}{2}\right], \\
\alpha_{27} &= \left[1, \frac{1}{2}\sqrt{5} - \frac{1}{2}\mu - \frac{1}{2}\right], & \alpha_{28} &= \left[1, \frac{1}{2}\sqrt{5} + \frac{1}{2}\mu - \frac{1}{2}\right], \\
\alpha_{29} &= \left[1, -\frac{1}{16}\nu(i\sqrt{3} + 1)(\mu + 2)(1 + \sqrt{5})\right], & \alpha_{30} &= \left[1, \frac{1}{16}\nu(i\sqrt{3} + 1)(\mu - 2)(1 + \sqrt{5})\right], \\
\alpha_{31} &= \left[1, \frac{1}{16}\nu(i\sqrt{3} + 1)(\mu + 2)(1 + \sqrt{5})\right], & \alpha_{32} &= \left[1, -\frac{1}{16}\nu(i\sqrt{5} + 1)(\mu - 2)(1 + \sqrt{5})\right],
\end{aligned}$$

with

$$\mu = \sqrt{10 - 2\sqrt{5}}, \quad \nu = \sqrt{2 + 2i\sqrt{3}}.$$

9.11 The G_{20} case

We have 20 mirrors defining reflections of order 2:

$$\begin{aligned}
\alpha_1 &= \left[1, -i\sqrt{-2\sqrt{5}\sqrt{3} + 4\sqrt{3} - 3\sqrt{5} + 8}\right], & \alpha_2 &= \left[1, i\sqrt{-2\sqrt{5}\sqrt{3} + 4\sqrt{3} - 3\sqrt{5} + 8}\right], \\
\alpha_3 &= \left[1, -i\sqrt{2\sqrt{5}\sqrt{3} - 4\sqrt{3} - 3\sqrt{5} + 8}\right], & \alpha_4 &= \left[1, i\sqrt{2\sqrt{5}\sqrt{3} - 4\sqrt{3} - 3\sqrt{5} + 8}\right], \\
\alpha_5 &= \left[1, \frac{1}{2}i\sqrt{3} + \frac{1}{2}\sqrt{2 - 2i\sqrt{3}} - \frac{1}{2}\right], & \alpha_6 &= \left[1, \frac{1}{2}i\sqrt{3} - \frac{1}{2}\sqrt{2 - 2i\sqrt{3}} - \frac{1}{2}\right], \\
\alpha_7 &= \left[1, -\frac{1}{2}i\sqrt{3} + \frac{1}{4}\sqrt{2 - 2i\sqrt{3}} - \frac{1}{2} + \frac{1}{4}i\sqrt{3}\sqrt{2 - 2i\sqrt{3}}\right], \\
\alpha_8 &= \left[1, -\frac{1}{2}i\sqrt{3} - \frac{1}{4}\sqrt{2 - 2i\sqrt{3}} - \frac{1}{2} - \frac{1}{4}i\sqrt{2 - 2i\sqrt{3}}\sqrt{3}\right], \\
\alpha_9 &= \left[1, -\frac{1}{2}\sqrt{5} - \frac{3}{2} + \frac{1}{2}\sqrt{5}\sqrt{3} + \frac{1}{2}\sqrt{3}\right], & \alpha_{10} &= \left[1, -\frac{1}{2}\sqrt{5} - \frac{3}{2} - \frac{1}{2}\sqrt{5}\sqrt{3} - \frac{1}{2}\sqrt{3}\right], \\
\alpha_{11} &= \left[1, -\frac{1}{3}\sqrt{6i + 3\sqrt{5}}\right], & \alpha_{12} &= \left[1, \frac{1}{3}\sqrt{6i + 3\sqrt{5}}\right], \\
\alpha_{13} &= \left[1, -\frac{1}{3}\sqrt{-6i + 3\sqrt{5}}\right], & \alpha_{14} &= \left[1, \frac{1}{3}\sqrt{-6i + 3\sqrt{5}}\right], \\
\alpha_{15} &= \left[1, \frac{1}{2}i\sqrt{3} - \frac{1}{4}\sqrt{2 - 2i\sqrt{3}} + \frac{1}{2} - \frac{1}{4}i\sqrt{2 - 2i\sqrt{3}}\sqrt{3}\right], \\
\alpha_{16} &= \left[1, -\frac{1}{2}i\sqrt{3} - \frac{1}{2}\sqrt{2 - 2i\sqrt{3}} + \frac{1}{2}\right], & \alpha_{17} &= \left[1, -\frac{1}{2}i\sqrt{3} + \frac{1}{2}\sqrt{2 - 2i\sqrt{3}} + \frac{1}{2}\right], \\
\alpha_{18} &= \left[1, \frac{1}{2}i\sqrt{3} + \frac{1}{4}\sqrt{2 - 2i\sqrt{3}} + \frac{1}{2} + \frac{1}{4}i\sqrt{3}\sqrt{2 - 2i\sqrt{3}}\right], \\
\alpha_{19} &= \left[1, \frac{1}{2}\sqrt{5} + \frac{3}{2} - \frac{1}{2}\sqrt{5}\sqrt{3} - \frac{1}{2}\sqrt{3}\right], & \alpha_{20} &= \left[1, \frac{1}{2}\sqrt{5} + \frac{3}{2} + \frac{1}{2}\sqrt{5}\sqrt{3} + \frac{1}{2}\sqrt{3}\right].
\end{aligned}$$

9.12 The G_{21} case

We have 30 mirrors defining reflections of order 2:

$$\begin{aligned}
\alpha_1 &= \left[1, -\frac{1}{2}\sqrt{5} + \frac{1}{2} + \frac{3}{2}i - \frac{1}{2}i\sqrt{5}\right], & \alpha_2 &= \left[1, -\frac{1}{2}\sqrt{5} + \frac{1}{2} - \frac{3}{2}i + \frac{1}{2}i\sqrt{5}\right], \\
\alpha_3 &= [1, 1], & \alpha_4 &= \left[1, \frac{1}{6}\sqrt{5} - \frac{1}{6} - \frac{1}{6}i - \frac{1}{6}i\sqrt{5}\right], \\
\alpha_5 &= \left[1, \frac{1}{6}\sqrt{5} - \frac{1}{6} + \frac{1}{6}i + \frac{1}{6}i\sqrt{5}\right], & \alpha_6 &= \left[1, -\frac{3}{10}\sqrt{5} - \frac{1}{2} + \frac{1}{2}i + \frac{1}{10}i\sqrt{5}\right], \\
\alpha_7 &= \left[1, -\frac{3}{10}\sqrt{5} - \frac{1}{2} - \frac{1}{2}i - \frac{1}{10}i\sqrt{5}\right], & \alpha_8 &= [1, -1], \\
\alpha_9 &= [0, 1], & \alpha_{10} &= \left[1, -\frac{1}{10}\sqrt{5} + \frac{1}{2} + \frac{1}{2}i - \frac{3}{10}i\sqrt{5}\right], \\
\alpha_{11} &= \left[1, -\frac{1}{10}\sqrt{5} + \frac{1}{2} - \frac{1}{2}i + \frac{3}{10}i\sqrt{5}\right], & \alpha_{12} &= \left[1, -\frac{1}{6}\sqrt{5} + \frac{1}{6} + \frac{1}{6}i + \frac{1}{6}i\sqrt{5}\right], \\
\alpha_{13} &= \left[1, -\frac{1}{6}\sqrt{5} + \frac{1}{6} - \frac{1}{6}i - \frac{1}{6}i\sqrt{5}\right], & \alpha_{14} &= \left[1, \frac{3}{10}\sqrt{5} + \frac{1}{2} - \frac{1}{2}i - \frac{1}{10}i\sqrt{5}\right], \\
\alpha_{15} &= \left[1, \frac{3}{10}\sqrt{5} + \frac{1}{2} + \frac{1}{2}i + \frac{1}{10}i\sqrt{5}\right], & \alpha_{16} &= \left[1, \frac{1}{2}\sqrt{5} - \frac{1}{2} - \frac{1}{2}i - \frac{1}{2}i\sqrt{5}\right], \\
\alpha_{17} &= \left[1, \frac{1}{2}\sqrt{5} - \frac{1}{2} + \frac{1}{2}i + \frac{1}{2}i\sqrt{5}\right], & \alpha_{18} &= \left[1, \frac{1}{2}\sqrt{5} + \frac{3}{2} + \frac{1}{2}i + \frac{1}{2}i\sqrt{5}\right], \\
\alpha_{19} &= \left[1, \frac{1}{2}\sqrt{5} + \frac{3}{2} - \frac{1}{2}i - \frac{1}{2}i\sqrt{5}\right], & \alpha_{20} &= \left[1, -\frac{1}{2}\sqrt{5} - \frac{3}{2} - \frac{1}{2}i - \frac{1}{2}i\sqrt{5}\right], \\
\alpha_{21} &= \left[1, -\frac{1}{2}\sqrt{5} - \frac{3}{2} + \frac{1}{2}i + \frac{1}{2}i\sqrt{5}\right], & \alpha_{22} &= \left[1, -\frac{1}{2}\sqrt{5} + \frac{1}{2} + \frac{1}{2}i + \frac{1}{2}i\sqrt{5}\right], \\
\alpha_{23} &= \left[1, -\frac{1}{2}\sqrt{5} + \frac{1}{2} - \frac{1}{2}i - \frac{1}{2}i\sqrt{5}\right], & \alpha_{24} &= [1, 0], \\
\alpha_{25} &= [1, -i], & \alpha_{26} &= [1, i], \\
\alpha_{27} &= \left[1, \frac{1}{2}\sqrt{5} - \frac{1}{2} + \frac{3}{2}i - \frac{1}{2}i\sqrt{5}\right], & \alpha_{28} &= \left[1, \frac{1}{2}\sqrt{5} - \frac{1}{2} - \frac{3}{2}i + \frac{1}{2}i\sqrt{5}\right], \\
\alpha_{29} &= \left[1, -\frac{1}{2} + \frac{1}{2}i + \frac{1}{10}\sqrt{5} - \frac{3}{10}i\sqrt{5}\right], & \alpha_{30} &= \left[1, -\frac{1}{2} - \frac{1}{2}i + \frac{1}{10}\sqrt{5} + \frac{3}{10}i\sqrt{5}\right],
\end{aligned}$$

and 20 mirrors defining reflections of order 3:

$$\begin{aligned}
\alpha_{31} &= \left[1, \frac{1}{2} + \frac{1}{2}i - \frac{1}{2}\sqrt{6i}\right], & \alpha_{32} &= \left[1, \frac{1}{2} + \frac{1}{2}i + \frac{1}{2}\sqrt{6i}\right], \\
\alpha_{33} &= \left[1, -\frac{1}{2}i\sqrt{6i} + \frac{1}{2} - \frac{1}{2}i\right], & \alpha_{34} &= \left[1, \frac{1}{2}i\sqrt{6i} + \frac{1}{2} - \frac{1}{2}i\right], \\
\alpha_{35} &= \left[1, -\frac{1}{2}\sqrt{5} - \frac{3}{2} - \frac{1}{2}\sqrt{5}\sqrt{3} - \frac{1}{2}\sqrt{3}\right], & \alpha_{36} &= \left[1, -\frac{3}{2} - \frac{1}{2}\sqrt{5} + \frac{1}{2}\sqrt{5}\sqrt{3} + \frac{1}{2}\sqrt{3}\right], \\
\alpha_{37} &= [1, \sigma], & \alpha_{38} &= [1, -\nu\sqrt{5} + \sigma^3], \\
\alpha_{39} &= [1, -\sigma^3 + \nu\sqrt{5}], & \alpha_{40} &= \left[1, -\frac{1}{6}\sqrt{5}\sqrt{3} - \frac{1}{6}\sqrt{3} - \frac{1}{6}i\sqrt{5}\sqrt{3} + \frac{1}{6}i\sqrt{3}\right], \\
\alpha_{41} &= \left[1, -\frac{1}{2}i\sqrt{6i} - \frac{1}{2} + \frac{1}{2}i\right], & \alpha_{42} &= \left[1, \frac{1}{2}i\sqrt{6i} - \frac{1}{2} + \frac{1}{2}i\right], \\
\alpha_{43} &= \left[1, -\frac{1}{2} - \frac{1}{2}i + \frac{1}{2}\sqrt{6i}\right], & \alpha_{44} &= \left[1, -\frac{1}{2} - \frac{1}{2}i - \frac{1}{2}\sqrt{6i}\right], \\
\alpha_{45} &= [1, -\mu^3 + \lambda\sqrt{5} + 24i - 8i\sqrt{5} - 8i\sqrt{5}\sqrt{3} + 8i\sqrt{3}], & \alpha_{46} &= [1, \mu], \\
\alpha_{47} &= [1, \mu^3 - \lambda\sqrt{5} - 24i + 8i\sqrt{5} + 8i\sqrt{5}\sqrt{3} - 8i\sqrt{3}], & \alpha_{48} &= \left[1, \frac{3}{2}i - \frac{1}{2}i\sqrt{5} - \frac{1}{2}i\sqrt{5}\sqrt{3} + \frac{1}{2}i\sqrt{3}\right], \\
\alpha_{49} &= \left[1, \frac{1}{2}\sqrt{5} + \frac{3}{2} - \frac{1}{2}\sqrt{5}\sqrt{3} - \frac{1}{2}\sqrt{3}\right], & \alpha_{50} &= \left[1, \frac{1}{2}\sqrt{5} + \frac{3}{2} + \frac{1}{2}\sqrt{5}\sqrt{3} + \frac{1}{2}\sqrt{3}\right],
\end{aligned}$$

with

$$\mu = -\frac{3}{2}i + \frac{1}{2}i\sqrt{5} + \frac{1}{2}i\sqrt{5}\sqrt{3} - \frac{1}{2}i\sqrt{3}, \quad \nu = \frac{1}{9}\sqrt{5}\sqrt{3} + \frac{1}{9}\sqrt{3} + \frac{1}{9}i\sqrt{5}\sqrt{3} - \frac{1}{9}i\sqrt{3}, \quad \sigma = \frac{3}{2}\nu, \quad \lambda = 6\mu.$$

9.13 The G_{23} case

We have 15 mirrors defining reflections of order 2:

$$\begin{aligned}
\alpha_1 &= [1, 0, 0] & \alpha_2 &= [0, 1, 0] & \alpha_3 &= [0, 0, 1] \\
\alpha_4 &= [-2, \sqrt{5} + 3, -\sqrt{5} - 1], & \alpha_5 &= [-2, \sqrt{5} - 1, -\sqrt{5} + 3], & \alpha_6 &= [2, \sqrt{5} - 1, \sqrt{5} - 3] \\
\alpha_7 &= [-2, \sqrt{5} - 1, \sqrt{5} - 3], & \alpha_8 &= [2, \sqrt{5} + 3, -\sqrt{5} - 1], & \alpha_9 &= [-2, \sqrt{5} + 3, \sqrt{5} + 1] \\
\alpha_{10} &= [2, \sqrt{5} - 1, -\sqrt{5} + 3], & \alpha_{11} &= [2, \sqrt{5} + 3, \sqrt{5} + 1], & \alpha_{12} &= [2, \sqrt{5} - 1, -\sqrt{5} - 1] \\
\alpha_{13} &= [-2, \sqrt{5} - 1, -\sqrt{5} - 1], & \alpha_{14} &= [2, \sqrt{5} - 1, \sqrt{5} + 1], & \alpha_{15} &= [-2, \sqrt{5} - 1, \sqrt{5} + 1]
\end{aligned}$$

9.14 The G_{24} case

We have 21 mirrors defining reflections of order 2:

$$\begin{aligned}
\alpha_1 &= \left[1, -\frac{1}{9}\mu^2 - \frac{1}{9}\mu + \frac{11}{9}, \frac{1}{9}\mu^2 + \frac{1}{9}\mu - \frac{20}{9} \right] \\
\alpha_2 &= \left[1, \frac{1}{3}\mu - \frac{1}{3}, -\frac{1}{3}\mu - \frac{2}{3} \right] \\
\alpha_3 &= \left[1, \frac{1}{9}\mu^2 - \frac{2}{9}\mu - \frac{17}{9}, -\frac{1}{9}\mu^2 + \frac{2}{9}\mu + \frac{8}{9} \right] \\
\alpha_4 &= \left[1, \frac{1}{27}\nu\mu^2 - \frac{2}{27}\nu\mu + \frac{1}{6}\mu - \frac{8}{27}\nu + \frac{5}{6}, \frac{1}{2} + \frac{1}{27}\nu\mu^2 - \frac{17}{27}\nu - \frac{2}{27}\nu\mu \right] \\
\alpha_5 &= \left[1, -\frac{1}{27}\nu\mu^2 - \frac{2}{27}\nu\mu + \frac{1}{6}\mu + \frac{8}{27}\nu + \frac{5}{6}, \frac{1}{2} - \frac{1}{27}\nu\mu^2 + \frac{17}{27}\nu + \frac{2}{27}\nu\mu \right] \\
\alpha_6 &= \left[1, \frac{1}{27}\nu\mu^2 - \frac{1}{18}\mu^2 - \frac{5}{27}\nu\mu - \frac{1}{18}\mu + \frac{4}{27}\nu + \frac{29}{18}, -\frac{4}{9}\nu\mu + \frac{1}{9}\nu\mu^2 - \frac{1}{3}\nu + \frac{1}{2} \right] \\
\alpha_7 &= \left[1, -\frac{1}{27}\nu\mu^2 - \frac{1}{18}\mu^2 + \frac{5}{27}\nu\mu - \frac{1}{18}\mu - \frac{4}{27}\nu + \frac{29}{18}, \frac{4}{9}\nu\mu - \frac{1}{9}\nu\mu^2 + \frac{1}{3}\nu + \frac{1}{2} \right] \\
\alpha_8 &= \left[1, \frac{1}{27}\nu\mu^2 + \frac{1}{18}\mu^2 - \frac{5}{27}\nu\mu - \frac{1}{9}\mu - \frac{5}{27}\nu + \frac{1}{18}, \frac{2}{27}\nu\mu - \frac{1}{27}\nu\mu^2 - \frac{1}{27}\nu + \frac{1}{2} \right] \\
\alpha_9 &= \left[1, -\frac{1}{27}\nu\mu^2 + \frac{1}{18}\mu^2 + \frac{5}{27}\nu\mu - \frac{1}{9}\mu + \frac{5}{27}\nu + \frac{1}{18}, -\frac{2}{27}\nu\mu + \frac{1}{27}\nu\mu^2 + \frac{1}{27}\nu + \frac{1}{2} \right] \\
\alpha_{10} &= \left[1, \frac{2}{27}\nu\mu^2 + \frac{1}{18}\mu^2 - \frac{7}{27}\nu\mu + \frac{1}{18}\mu - \frac{13}{27}\nu - \frac{10}{9}, -\frac{1}{27}\nu\mu^2 + \frac{1}{18}\mu^2 + \frac{5}{27}\mu\nu - \frac{5}{18}\mu - \frac{4}{27}\nu - \frac{5}{18} \right] \\
\alpha_{11} &= \left[1, -\frac{2}{27}\nu\mu^2 + \frac{1}{18}\mu^2 + \frac{7}{27}\nu\mu + \frac{1}{18}\mu + \frac{13}{27}\nu - \frac{10}{9}, \frac{1}{27}\nu\mu^2 + \frac{1}{18}\mu^2 - \frac{5}{27}\mu\nu - \frac{5}{18}\mu + \frac{4}{27}\nu - \frac{5}{18} \right] \\
\alpha_{12} &= \left[1, -\frac{1}{6}\mu - \frac{1}{3} + \frac{1}{3}\nu, \frac{1}{27}\nu\mu^2 - \frac{1}{9}\mu^2 - \frac{2}{27}\nu\mu + \frac{1}{18}\mu - \frac{8}{27}\nu + \frac{37}{18} \right] \\
\alpha_{13} &= \left[1, -\frac{1}{6}\mu - \frac{1}{3} - \frac{1}{3}\nu, -\frac{1}{27}\nu\mu^2 - \frac{1}{9}\mu^2 + \frac{2}{27}\nu\mu + \frac{1}{18}\mu + \frac{8}{27}\nu + \frac{37}{18} \right] \\
\alpha_{14} &= \left[1, -\frac{2}{27}\nu\mu^2 - \frac{1}{18}\mu^2 + \frac{7}{27}\nu\mu + \frac{1}{9}\mu + \frac{4}{27}\nu + \frac{4}{9}, -\frac{1}{27}\nu\mu^2 + \frac{1}{18}\mu^2 + \frac{5}{27}\mu\nu + \frac{2}{9}\mu + \frac{5}{27}\nu - \frac{5}{18} \right] \\
\alpha_{15} &= \left[1, \frac{2}{27}\nu\mu^2 - \frac{1}{18}\mu^2 - \frac{7}{27}\nu\mu + \frac{1}{9}\mu - \frac{4}{27}\nu + \frac{4}{9}, \frac{1}{27}\nu\mu^2 + \frac{1}{18}\mu^2 - \frac{5}{27}\mu\nu + \frac{2}{9}\mu - \frac{5}{27}\nu - \frac{5}{18} \right] \\
\alpha_{16} &= \left[1, \frac{1}{18}\mu^2 + \frac{1}{9}\nu\mu + \frac{1}{18}\mu - \frac{1}{9}\nu - \frac{10}{9}, \frac{1}{18}\mu^2 - \frac{1}{9}\nu\mu - \frac{1}{9}\mu + \frac{4}{9}\nu - \frac{13}{9} \right] \\
\alpha_{17} &= \left[1, \frac{1}{18}\mu^2 - \frac{1}{9}\nu\mu + \frac{1}{18}\mu + \frac{1}{9}\nu - \frac{10}{9}, \frac{1}{18}\mu^2 + \frac{1}{9}\nu\mu - \frac{1}{9}\mu - \frac{4}{9}\nu - \frac{13}{9} \right] \\
\alpha_{18} &= \left[1, \frac{2}{27}\nu\mu^2 - \frac{10}{27}\nu\mu - \frac{1}{6}\mu - \frac{1}{27}\nu - \frac{1}{3}, -\frac{1}{18}\mu^2 - \frac{1}{18}\mu + \frac{1}{9}\mu\nu - \frac{1}{9}\nu + \frac{1}{9} \right] \\
\alpha_{19} &= \left[1, -\frac{2}{27}\nu\mu^2 + \frac{10}{27}\nu\mu - \frac{1}{6}\mu + \frac{1}{27}\nu - \frac{1}{3}, -\frac{1}{18}\mu^2 - \frac{1}{18}\mu - \frac{1}{9}\mu\nu + \frac{1}{9}\nu + \frac{1}{9} \right] \\
\alpha_{20} &= \left[1, -\frac{1}{18}\mu^2 - \frac{1}{9}\nu\mu + \frac{1}{9}\mu + \frac{4}{9}\nu + \frac{4}{9}, -\frac{2}{27}\nu\mu^2 + \frac{10}{27}\nu\mu + \frac{1}{6}\mu + \frac{1}{27}\nu - \frac{2}{2} \right] \\
\alpha_{21} &= \left[1, -\frac{1}{18}\mu^2 + \frac{1}{9}\nu\mu + \frac{1}{9}\mu + \frac{4}{9}\nu - \frac{4}{9}, \frac{2}{27}\nu\mu^2 - \frac{10}{27}\nu\mu + \frac{1}{6}\mu - \frac{1}{27}\nu - \frac{2}{2} \right]
\end{aligned}$$

where μ is a root of $z^3 - 21z - 7 = 0$ and $\nu = \pm\sqrt{\frac{\mu^2}{4} - 2\mu - 14}$.

9.15 The G_{25} case

We have 12 mirrors defining reflections of order 2:

$$\begin{aligned}\alpha_1 &= [1, 0, 0], \alpha_2 = [0, 1, 0], \alpha_3 = [0, 0, 1], \\ \alpha_4 &= [1, 1, 1] \alpha_5 = [-2, 1 + \mu, 1 - \mu], \alpha_6 = [2, -1 + \mu - 1 - \mu], \\ \alpha_7 &= [-2, -2, 1 + \mu], \alpha_8 = [2, 2, -1 + \mu], \alpha_9 = [-2, 1 + \mu, -2], \\ \alpha_{10} &= [2, -1 + \mu, 2], \alpha_{11} = [2, -1 + \mu, -1 + \mu], \alpha_{12} = [-2, 1 + \mu, 1 + \mu]\end{aligned}$$

where $\mu = \pm i\sqrt{3}$.

9.16 The G_{26} case

We have 9 mirrors defining reflections of order 2:

$$\begin{aligned}\alpha_1 &= \left[0, 1, -\frac{1}{2}\mu + \frac{1}{2}\right], \alpha_2 = \left[0, 1, \frac{1}{2}\mu + \frac{1}{2}\right], \alpha_3 = \left[1, 0, \frac{1}{2}\mu + \frac{1}{2}\right], \\ \alpha_4 &= \left[1, 0, -\frac{1}{2}\mu + \frac{1}{2}\right], \alpha_5 = \left[1, -\frac{1}{2}\mu + \frac{1}{2}, 0\right], \alpha_6 = \left[1, \frac{1}{2}\mu + \frac{1}{2}, 0\right], \\ \alpha_7 &= [1, -1, 0], \alpha_8 = [1, 0, -1], \alpha_9 = [0, 1, -1],\end{aligned}$$

and 12 hyperplanes defining reflections of order 3:

$$\begin{aligned}\alpha_{10} &= [1, 0, 0], \alpha_{11} = [0, 1, 0], \alpha_{12} = [0, 0, 1], \\ \alpha_{13} &= [1, 1, 1], \alpha_{14} = \left[1, \frac{1}{2}\mu - \frac{1}{2}, -\frac{1}{2}\mu - \frac{1}{2}\right], \alpha_{15} = \left[1, -\frac{1}{2}\mu - \frac{1}{2}, \frac{1}{2}\mu - \frac{1}{2}\right], \\ \alpha_{16} &= \left[1, \frac{1}{2}\mu - \frac{1}{2}, 1\right], \alpha_{17} = \left[1, -\frac{1}{2}\mu - \frac{1}{2}, 1\right], \alpha_{18} = \left[1, 1, \frac{1}{2}\mu - \frac{1}{2}\right], \\ \alpha_{19} &= \left[1, 1, -\frac{1}{2}\mu - \frac{1}{2}\right], \alpha_{20} = \left[1, \frac{1}{2}\mu - \frac{1}{2}, \frac{1}{2}\mu - \frac{1}{2}\right], \alpha_{21} = \left[1, -\frac{1}{2}\mu - \frac{1}{2}, -\frac{1}{2}\mu - \frac{1}{2}\right]\end{aligned}$$

where $\mu = \pm i\sqrt{3}$.

9.17 The G_{27} case

We have 45 mirrors defining reflections of order 2:

$$\begin{aligned}
\alpha_1 &= [1, -1, 0] \\
\alpha_2 &= [-384, \mu^3 - \mu^2 - 165\mu + 2\sigma - 291, 0] \\
\alpha_3 &= [-384, \mu^3 - \mu^2 - 165\mu - 2\sigma - 291, 0] \\
\alpha_4 &= [-36864, \mu^3\sigma - 96\mu^3 - \mu^2\sigma + 96\mu^2 - 165\mu\sigma + 15840\mu - 291\sigma + 9504, 0] \\
\alpha_5 &= [36864, \mu^3\sigma + 96\mu^3 - \mu^2\sigma - 96\mu^2 - 165\mu\sigma - 15840\mu - 291\sigma - 9504, 0] \\
\alpha_6 &= [36864, 36864, \\
&\quad \mu^3\sigma - 288\mu^3 - 9\mu^2\sigma + 288\mu^2 - 117\mu\sigma + 47520\mu + 405\sigma + 28512] \\
\alpha_7 &= [-36864, -36864, \\
&\quad \mu^3\sigma + 288\mu^3 - 9\mu^2\sigma - 288\mu^2 - 117\mu\sigma - 47520\mu + 405\sigma - 28512] \\
\alpha_8 &= [-36864, \mu^3\sigma + 96\mu^3 - \mu^2\sigma - 96\mu^2 - 165\mu\sigma - 15840\mu - 291\sigma - 9504, \\
&\quad -3\mu^3\sigma - 288\mu^3 + 3\mu^2\sigma + 1440\mu^2 + 447\mu\sigma + 36000\mu + 441\sigma - 2592] \\
\alpha_9 &= [36864, \mu^3\sigma - 96\mu^3 - \mu^2\sigma + 96\mu^2 - 165\mu\sigma + 15840\mu - 291\sigma + 9504, \\
&\quad -3\mu^3\sigma + 288\mu^3 + 3\mu^2\sigma - 1440\mu^2 + 447\mu\sigma - 36000\mu + 441\sigma + 2592] \\
\alpha_{10} &= [36864, \mu^3\sigma - 96\mu^3 - \mu^2\sigma + 96\mu^2 - 165\mu\sigma + 15840\mu - 291\sigma + 9504, \\
&\quad -288\mu^3 + 1440\mu^2 + 48\mu\sigma + 36000\mu - 144\sigma - 57888] \\
\alpha_{11} &= [-36864, \mu^3\sigma + 96\mu^3 - \mu^2\sigma - 96\mu^2 - 165\mu\sigma - 15840\mu - 291\sigma - 9504, \\
&\quad 288\mu^3 - 1440\mu^2 + 48\mu\sigma - 36000\mu - 144\sigma + 57888] \\
\alpha_{12} &= [36864, \mu^3\sigma - 96\mu^3 - \mu^2\sigma + 96\mu^2 - 165\mu\sigma + 15840\mu - 291\sigma + 9504, \\
&\quad 4\mu^3\sigma - 96\mu^3 + 864\mu^2 - 588\mu\sigma + 15840\mu - 1224\sigma + 2592] \\
\alpha_{13} &= [-36864, \mu^3\sigma + 96\mu^3 - \mu^2\sigma - 96\mu^2 - 165\mu\sigma - 15840\mu - 291\sigma - 9504, \\
&\quad 4\mu^3\sigma + 96\mu^3 - 864\mu^2 - 588\mu\sigma - 15840\mu - 1224\sigma - 2592] \\
\alpha_{14} &= [-36864, \mu^3\sigma + 96\mu^3 - \mu^2\sigma - 96\mu^2 - 165\mu\sigma - 15840\mu - 291\sigma - 9504, \\
&\quad -\mu^3\sigma + 192\mu^3 - 3\mu^2\sigma + 576\mu^2 + 93\mu\sigma - 31680\mu + 351\sigma - 136512] \\
\alpha_{15} &= [36864, \mu^3\sigma - 96\mu^3 - \mu^2\sigma + 96\mu^2 - 165\mu\sigma + 15840\mu - 291\sigma + 9504, \\
&\quad -\mu^3\sigma - 192\mu^3 - 3\mu^2\sigma - 576\mu^2 + 93\mu\sigma + 31680\mu + 351\sigma + 136512] \\
\alpha_{16} &= [768, 2\mu^3 - 2\mu^2 - 330\mu + 4\sigma - 582, 8\mu^3 - \mu\sigma - 1080\mu - 9\sigma - 2736] \\
\alpha_{17} &= [768, 2\mu^3 - 2\mu^2 - 330\mu - 4\sigma - 582, 8\mu^3 + \mu\sigma - 1080\mu + 9\sigma - 2736] \\
\alpha_{18} &= [-18432, -18432, \\
&\quad \mu^3\sigma - 144\mu^3 + 3\mu^2\sigma + 144\mu^2 - 141\mu\sigma + 23760\mu - 495\sigma + 41904]
\end{aligned}$$

$$\begin{aligned}
\alpha_{19} &= [18432, 18432, \\
&\quad \mu^3\sigma + 144\mu^3 + 3\mu^2\sigma - 144\mu^2 - 141\mu\sigma - 23760\mu - 495\sigma - 41904] \\
\alpha_{20} &= [-12288, -32\mu^3 + 32\mu^2 + 5280\mu + 64\sigma + 9312, \\
&\quad \mu^3\sigma + 128\mu^3 - \mu^2\sigma - 149\mu\sigma - 17280\mu - 339\sigma - 25344] \\
\alpha_{21} &= [12288, 32\mu^3 - 32\mu^2 - 5280\mu + 64\sigma - 9312, \\
&\quad \mu^3\sigma - 128\mu^3 - \mu^2\sigma - 149\mu\sigma + 17280\mu - 339\sigma + 25344] \\
\alpha_{22} &= [-18432, -48\mu^3 + 48\mu^2 + 7920\mu - 96\sigma + 13968, \\
&\quad \mu^3\sigma - 240\mu^3 - 3\mu^2\sigma + 432\mu^2 - 129\mu\sigma + 33840\mu - 189\sigma + 11664] \\
\alpha_{23} &= [18432, 48\mu^3 - 48\mu^2 - 7920\mu - 96\sigma - 13968, \\
&\quad \mu^3\sigma + 240\mu^3 - 3\mu^2\sigma - 432\mu^2 - 129\mu\sigma - 33840\mu - 189\sigma - 11664] \\
\alpha_{24} &= [-18432, -48\mu^3 + 48\mu^2 + 7920\mu + 96\sigma + 13968, \\
&\quad \mu^3\sigma + 96\mu^3 - 3\mu^2\sigma - 288\mu^2 - 129\mu\sigma - 10080\mu + 99\sigma - 25056] \\
\alpha_{25} &= [18432, 48\mu^3 - 48\mu^2 - 7920\mu + 96\sigma - 13968, \\
&\quad \mu^3\sigma - 96\mu^3 - 3\mu^2\sigma + 288\mu^2 - 129\mu\sigma + 10080\mu + 99\sigma + 25056] \\
\alpha_{26} &= [-9216, \mu^2\sigma + 96\mu^2 + 6\mu\sigma - 576\mu + 21\sigma + 864, \\
&\quad 24\mu^3 + 72\mu^2 + 12\mu\sigma - 3384\mu - 36\sigma - 4968] \\
\alpha_{27} &= [-9216, -\mu^2\sigma + 96\mu^2 - 6\mu\sigma - 576\mu - 21\sigma + 864, \\
&\quad 24\mu^3 + 72\mu^2 - 12\mu\sigma - 3384\mu + 36\sigma - 4968] \\
\alpha_{28} &= [-96, \mu^3 - 3\mu^2 - 153\mu - 21, -\mu^3 + 3\mu^2 + 153\mu + 117] \\
\alpha_{29} &= [-192, \mu^3 + 3\mu^2 - 189\mu - 447, -\mu^3 - 3\mu^2 + 189\mu + 639] \\
\alpha_{30} &= [36864, 2\mu^3\sigma - 96\mu^3 + 2\mu^2\sigma + 480\mu^2 - 306\mu\sigma + 13536\mu - 1074\sigma - 42336, \\
&\quad 3\mu^3\sigma + 96\mu^3 - 3\mu^2\sigma + 288\mu^2 - 447\mu\sigma - 13536\mu - 441\sigma - 75168] \\
\alpha_{31} &= [-36864, 2\mu^3\sigma + 96\mu^3 + 2\mu^2\sigma - 480\mu^2 - 306\mu\sigma - 13536\mu - 1074\sigma + 42336, \\
&\quad 3\mu^3\sigma - 96\mu^3 - 3\mu^2\sigma - 288\mu^2 - 447\mu\sigma + 13536\mu - 441\sigma + 75168] \\
\alpha_{32} &= [36864, 3\mu^3\sigma - 192\mu^3 - 3\mu^2\sigma - 192\mu^2 - 447\mu\sigma + 29376\mu - 633\sigma + 103104, \\
&\quad -\mu^3\sigma + 192\mu^3 - 3\mu^2\sigma - 576\mu^2 + 141\mu\sigma - 29376\mu + 783\sigma + 32832] \\
\alpha_{33} &= [-36864, 3\mu^3\sigma + 192\mu^3 - 3\mu^2\sigma + 192\mu^2 - 447\mu\sigma - 29376\mu - 633\sigma - 103104, \\
&\quad -\mu^3\sigma - 192\mu^3 - 3\mu^2\sigma + 576\mu^2 + 141\mu\sigma + 29376\mu + 783\sigma - 32832] \\
\alpha_{34} &= [36864, 384\mu^2 - 48\mu\sigma + 2304\mu - 48\sigma + 8064, \\
&\quad \mu^3\sigma + 96\mu^3 + 3\mu^2\sigma + 288\mu^2 - 141\mu\sigma - 18144\mu - 207\sigma - 6048] \\
\alpha_{35} &= [-36864, -384\mu^2 - 48\mu\sigma - 2304\mu - 48\sigma - 8064, \\
&\quad \mu^3\sigma - 96\mu^3 + 3\mu^2\sigma - 288\mu^2 - 141\mu\sigma + 18144\mu - 207\sigma + 6048] \\
\alpha_{36} &= [-4, \mu + 1, -\mu + 3]
\end{aligned}$$

$$\begin{aligned}
\alpha_{37} &= [64, \mu^3 - \mu^2 - 149\mu - 211, -\mu^3 + \mu^2 + 149\mu + 147] \\
\alpha_{38} &= [-18432, \mu^3\sigma + 48\mu^3 - 3\mu^2\sigma - 48\mu^2 - 153\mu\sigma - 5616\mu - 21\sigma + 6768, \\
&\quad 96\mu^3 - 288\mu^2 + 24\mu\sigma - 12384\mu + 216\sigma + 9504] \\
\alpha_{39} &= [18432, \mu^3\sigma - 48\mu^3 - 3\mu^2\sigma + 48\mu^2 - 153\mu\sigma + 5616\mu - 21\sigma - 6768, \\
&\quad -96\mu^3 + 288\mu^2 + 24\mu\sigma + 12384\mu + 216\sigma - 9504] \\
\alpha_{40} &= [-36864, \mu^3\sigma - 192\mu^3 + 3\mu^2\sigma + 192\mu^2 - 189\mu\sigma + 27072\mu - 447\sigma + 69696, \\
&\quad -3\mu^3\sigma - 192\mu^3 + 3\mu^2\sigma + 576\mu^2 + 447\mu\sigma + 24768\mu + 1017\sigma + 36288] \\
\alpha_{41} &= [36864, \mu^3\sigma + 192\mu^3 + 3\mu^2\sigma - 192\mu^2 - 189\mu\sigma - 27072\mu - 447\sigma - 69696, \\
&\quad -3\mu^3\sigma + 192\mu^3 + 3\mu^2\sigma - 576\mu^2 + 447\mu\sigma - 24768\mu + 1017\sigma - 36288] \\
\alpha_{42} &= [-36864, \mu^3\sigma - 288\mu^3 - \mu^2\sigma + 672\mu^2 - 117\mu\sigma + 40608\mu + 141\sigma - 9504, \\
&\quad -2\mu^3\sigma + 6\mu^2\sigma + 306\mu\sigma - 4608\mu + 234\sigma - 41472] \\
\alpha_{43} &= [36864, \mu^3\sigma + 288\mu^3 - \mu^2\sigma - 672\mu^2 - 117\mu\sigma - 40608\mu + 141\sigma + 9504, \\
&\quad -2\mu^3\sigma + 6\mu^2\sigma + 306\mu\sigma + 4608\mu + 234\sigma + 41472] \\
\alpha_{44} &= [36864, 2\mu^3\sigma - 96\mu^3 - 2\mu^2\sigma + 480\mu^2 - 282\mu\sigma + 8928\mu - 726\sigma + 26784, \\
&\quad \mu^3\sigma - 288\mu^3 + 3\mu^2\sigma + 288\mu^2 - 189\mu\sigma + 42912\mu - 639\sigma + 97632] \\
\alpha_{45} &= [-36864, 2\mu^3\sigma + 96\mu^3 - 2\mu^2\sigma - 480\mu^2 - 282\mu\sigma - 8928\mu - 726\sigma - 26784, \\
&\quad \mu^3\sigma + 288\mu^3 + 3\mu^2\sigma - 288\mu^2 - 189\mu\sigma - 42912\mu - 639\sigma - 97632]
\end{aligned}$$

where μ is a root of $\mu^4 - 150\mu^2 - 360\mu + 45 = 0$, and $\sigma = \sqrt{-48\mu^3 + 48\mu^2 + 7920\mu - 13680}$.

9.18 The $G(3, 1, 2)$ case

We have 2 mirrors defining reflections of order 3:

$$\alpha_1 = [1, 0], \quad \alpha_2 = [0, 1],$$

and 3 mirrors defining reflections of order 2:

$$\alpha_3 = [1, -1], \quad \alpha_4 = \left[1, \frac{1}{2}i\sqrt{3} + \frac{1}{2}\right], \quad \alpha_5 = \left[1, -\frac{1}{2}i\sqrt{3} + \frac{1}{2}\right].$$

9.19 The $G(3, 1, 3)$ case

We have 3 mirrors defining reflections of order 3:

$$\alpha_1 = [1, 0, 0], \quad \alpha_2 = [0, 1, 0], \quad \alpha_3 = [0, 0, 1].$$

and 9 mirrors defining reflections of order 2:

$$\begin{aligned}\alpha_4 &= [1, -1, 0], & \alpha_5 &= [1, 0, -1], & \alpha_6 &= [0, 1, -1] \\ \alpha_7 &= \left[1, \frac{1}{2}i\sqrt{3} + \frac{1}{2}, 0\right], & \alpha_8 &= \left[1, 0, \frac{1}{2}i\sqrt{3} + \frac{1}{2}\right], & \alpha_9 &= \left[0, 1, \frac{1}{2}i\sqrt{3} + \frac{1}{2}\right] \\ \alpha_{10} &= \left[1, -\frac{1}{2}i\sqrt{3} + \frac{1}{2}, 0\right], & \alpha_{11} &= \left[1, 0, -\frac{1}{2}i\sqrt{3} + \frac{1}{2}\right], & \alpha_{12} &= \left[0, 1, -\frac{1}{2}i\sqrt{3} + \frac{1}{2}\right]\end{aligned}$$

9.20 The case of $I_2(6)$

We have 6 mirrors defining reflections of order 2:

$$\begin{aligned}\alpha_1 &= [1, 0], & \alpha_2 &= [0, 1], & \alpha_3 &= \left[1, \frac{1}{\sqrt{3}}\right] \\ \alpha_4 &= [1, -\sqrt{3}], & \alpha_5 &= \left[1, -\frac{1}{\sqrt{3}}\right], & \alpha_6 &= [1, \sqrt{3}]\end{aligned}$$

9.21 The case of $I_2(8)$

We have 8 mirrors defining reflections of order 2:

$$\begin{aligned}\alpha_1 &= [1, 0], & \alpha_2 &= [0, 1], & \alpha_3 &= [1, -1] & \alpha_4 &= [1, 1] \\ \alpha_5 &= [1, -\sqrt{2} - 1], & \alpha_6 &= [1, \sqrt{2} - 1], & \alpha_7 &= [1, -\sqrt{2} + 1], & \alpha_8 &= [1, \sqrt{2} + 1]\end{aligned}$$

9.22 The case of $I_2(10)$

We have 10 mirrors defining reflections of order 2:

$$\begin{aligned}\alpha_1 &= [1, 0], & \alpha_2 &= [0, 1], \\ \alpha_3 &= \left[1, \frac{1}{2}\sqrt{5 - 2\sqrt{5}}(-4 - 2\sqrt{5})\right], & \alpha_4 &= \left[1, -\sqrt{5 - 2\sqrt{5}}\right], \\ \alpha_5 &= \left[1, \sqrt{5 - 2\sqrt{5}}\right], & \alpha_6 &= -\left[1, \frac{1}{2}\sqrt{5 - 2\sqrt{5}}(-4 - 2\sqrt{5})\right], \\ \alpha_7 &= \left[1, -\frac{1}{5}\sqrt{5 - 2\sqrt{5}}(-5 - 2\sqrt{5})\right], & \alpha_8 &= \left[1, \frac{1}{5}\sqrt{5 - 2\sqrt{5}}\sqrt{5}\right], \\ \alpha_9 &= \left[1, \frac{1}{5}\sqrt{5 - 2\sqrt{5}}(-5 - 2\sqrt{5})\right], & \alpha_{10} &= \left[1, -\frac{1}{5}\sqrt{5 - 2\sqrt{5}}\sqrt{5}\right]\end{aligned}$$

9.23 The case of B_2

We have 4 mirrors defining reflections of order 2:

$$\alpha_1 = [1, 0], \quad \alpha_2 = [0, 1], \quad \alpha_3 = [1, -1] \quad \alpha_4 = [1, 1]$$

9.24 The case of B_3

We have 9 mirrors defining reflections of order 2:

$$\begin{aligned} \alpha_1 &= [1, 0, 0], & \alpha_2 &= [0, 1, 0], & \alpha_3 &= [0, 0, 1] \\ \alpha_4 &= [1, 0, -1], & \alpha_6 &= [0, 1, -1], & \alpha_7 &= [1, -1, 0] \\ \alpha_7 &= [1, 1, 0], & \alpha_8 &= [1, 0, 1], & \alpha_9 &= [0, 1, 1]. \end{aligned}$$

10 Appendix 2. Saito coordinates in the cases of the groups G_{29}, G_{32}, G_{33}

For higher rank complex reflection groups the computations become cumbersome. In the case of G_{29}, G_{32}, G_{33} not all conditions have been checked. However, imposing the flatness conditions at some special points, it is sufficient to fix uniquely the parameters in the choice of the basic invariants. For this reason we conjecture that the basic invariants obtained in this way coincide with the generalized Saito flat coordinates of a bi-flat structure. We expect also that the vector potentials of the natural product of this structure coincide with the vector potentials obtained in [20] with a different approach.

10.1 The case of G_{32}

Basic invariants are (see [32])

$$u_1 = F_{12}$$

$$u_2 = -54p_4^{18} + (17)(54)C_6p_4^{12} + (54)(1870)C_9p_4^9 + \frac{1}{2}(17)(27)(19C_6^2 - 15C_{12})p_4^6 + (54)(170)C_6C_9p_4^3 + C_6^3 - 30C_6C_{12} - 25C_{18} = F_{18}$$

$$u_3 = 1728C_6p_4^{18} - (36)(1728)C_9p_4^{15} + (15)(144)(7C_{12} + C_6^2)p_4^{12} - (10)(1728)C_6C_9p_4^9 + (36)(178C_{18} - 135C_6C_{12} + 5C_6^3)p_4^6 + 432(41C_{12} - C_6^2)C_9p_4^3 + C_6^4 + 6C_6^2C_{12} - 16C_6C_{18} + 9C_{12}^2 = F_{24}$$

$$u_4 = F_{30} = -2(6^4)C_6p_4^{24} + 312(6^4)C_9p_4^{21} + 216(715C_{12} - 127C_6^2)p_4^{18} + 272(6^4)C_6C_9p_4^{15} + 18(1306C_{18} + 6045C_6C_{12} - 295C_6^3)p_4^{12} + 216(73C_6^2 - 5473C_{12})C_9p_4^9 + \frac{3}{2}(16648C_6C_{18} + 2334C_6^2C_{12} - 20709C_{12}^2 - C_6^4)p_4^6 - 36(1370C_{18} - 657C_6C_{12} + 7C_6^3)C_9p_4^3 + C_6^5 - 19C_6^3C_{12} + 29C_6^2C_{18} - 6C_6C_{12}^2 - 5C_{12}C_{18},$$

with

$$\begin{aligned} C_6 &= p_1^6 - 10p_1^3p_2^3 - 10p_1^3p_3^3 + p_2^6 - 10p_2^3p_3^3 + p_3^6 \\ C_9 &= (p_1^3 - p_2^3)(p_2^3 - p_3^3)(-p_1^3 + p_3^3) \\ C_{12} &= (p_1^3 + p_2^3 + p_3^3)((p_1^3 + p_2^3 + p_3^3)^3 + 216p_1^3p_2^3p_3^3) \\ C_{18} &= (p_1^3 + p_2^3 + p_3^3)^6 - 540p_1^3p_2^3p_3^3(p_1^3 + p_2^3 + p_3^3)^3 - 5832p_1^6p_2^6p_3^6 \end{aligned}$$

We conjecture that Saito flat coordinates are

$$u_1 = F_{12}, \quad u_2 = F_{18}, \quad u_3 = F_{24} + c_1F_{12}^2, \quad u_4 = F_{30} + c_2F_{12}F_{18}.$$

with

$$c_1 = -\frac{21}{25}, \quad c_2 = -\frac{11}{25}.$$

Up to an inessential constant factor they coincide with the choice of basic invariants of Orlik and Terao.

10.2 The case of G_{33}

The basic invariants are (see [6] with corrections of [34])

$$\begin{aligned}
U_1 &= p_1^4 - 8a_1p_1 + 48p_2p_3p_4p_5 \\
U_2 &= p_1^6 + (-20p_2^3 - 20p_3^3 - 20p_4^3 - 20p_5^3)p_1^3 + 360p_1^2p_2p_3p_4p_5 - 8p_2^6 + 80p_2^3p_3^3 + 80p_2^3p_4^3 \\
&\quad + 80p_2^3p_5^3 - 8p_3^6 + 80p_3^3p_4^3 + 80p_3^3p_5^3 - 8p_4^6 + 80p_4^3p_5^3 - 8p_5^6 \\
U_3 &= \frac{1}{63700992} \det(H(u_1)) \\
U_4 &= 5a_2p_1^6 + (99a_3 + a_1a_2)p_1^3 + 216a_4 - 36a_1a_3 + 24a_2^2 - 4a_1^2a_2 + \\
&\quad p_2p_3p_4p_5(3p_1^8 + 33a_1p_1^5 + (18a_2 + 30a_1^2)p_1^2) + (p_2p_3p_4p_5)^2(243p_1^4 + 108a_1p_1) \\
U_5 &= 4a_3p_1^9 + (54a_4 + 12a_1a_3 - a_2^2)p_1^6 + (162a_1a_4 - 18a_2a_3 + 12a_1^2a_3 - 2a_1a_2^2)p_1^3 + \\
&\quad 27a_3^2 - 18a_1a_2a_3 + 4a_1^3a_3 + 4a_2^3 - a_1^2a_2^2 + \\
&\quad p_2p_3p_4p_5(6a_2p_1^8 + p_1^5(54a_3 + 12a_1a_2) + p_1^2(243a_4 + 54a_1a_3 - 36a_2^2 + 6a_1^2a_2)) \\
&\quad + (p_2p_3p_4p_5)^2(3p_1^{10} + 18a_1p_1^7 + p_1^4(54a_2 + 27a_1^2) + p_1(162a_3 - 54a_1a_2 + 12a_1^3)).
\end{aligned}$$

with

$$\begin{aligned}
a_1 &= -p_2^3 - p_3^3 - p_4^3 - p_5^3 \\
a_2 &= p_2^3p_3^3 + p_2^3p_4^3 + p_2^3p_5^3 + p_3^3p_4^3 + p_3^3p_5^3 + p_4^3p_5^3 \\
a_3 &= -p_2^3p_3^3p_4^3 - p_2^3p_3^3p_5^3 - p_2^3p_4^3p_5^3 - p_3^3p_4^3p_5^3 \\
a_4 &= p_2^3p_3^3p_4^3p_5^3
\end{aligned}$$

We conjecture that Saito flat coordinates are

$$\begin{aligned}
u_1 &= U_1, \quad u_2 = U_2, \quad u_3 = U_3 + c_1U_1U_2, \quad u_4 = U_4 + c_2U_1^3 + c_3U_2^2, \\
u_5 &= U_5 + c_4U_1^3U_2 + c_5U_1^2U_3 + c_6U_2^3 + c_7U_2U_4.
\end{aligned}$$

with

$$\begin{aligned}
c_1 &= -\frac{1}{768}, \quad c_2 = -\frac{5}{3072}, \quad c_3 = -\frac{5}{2304}, \quad c_4 = \frac{11}{884736}, \quad c_5 = -\frac{1}{128}, \\
c_6 &= \frac{11}{1990656}, \quad c_7 = -\frac{1}{288}.
\end{aligned}$$

11 Appendix 3. The non-well generated cases

The procedure we have introduced works only partially for non-well generated complex reflection groups in the sense that it allows us to reconstruct flat connections, but it fails to provide a compatible product \circ . Let us present what happens if one tries to implement the algorithm in the case of G_7 , a non-well generated

complex reflection group of rank 2, whose ring of invariants is generated by the following polynomials (see [25]):

$$u_1 = (p_1^4 + (2i)\sqrt{3}p_1^2p_2^2 + p_2^4)^3, \quad u_2 = (p_1^5p_2 - p_1p_2^5)^2.$$

Since u_1 and u_2 have the same degree, the connection $\nabla^{(1)}$ is uniquely determined. It turns out that it is almost hydrodynamically equivalent to the dual connection. Unfortunately, defining the unit of the product as a flat vector field and defining the product \circ in the standard way, we obtain that the compatibility is no longer satisfied. Similar results hold true in the case of the groups G_{11} , G_{12} , G_{13} , G_{19} and G_{22} . In the case of G_{15} , there is one parameter in the basic invariants. It turns out that the connection $\nabla^{(1)}$ is almost hydrodynamically equivalent to the dual connection for each value of this parameter and that the compatibility with the product \circ is never satisfied. To conclude let us consider the case of G_{31} . The basic invariants are

$$u_1 = U_1, \quad u_2 = U_2, \quad u_3 = U_3 + c_1U_1U_2, \quad u_4 = U_4 + c_2U_1^3 + c_3U_2^2,$$

where U_2 and U_3 coincide respectively with $U_3 = F_{12}$ and $U_4 = F_{20}$ of G_{29} and

$$\begin{aligned} U_1 &= p_1^8 + 14p_1^4p_2^4 + 14p_1^4p_3^4 + 14p_1^4p_4^4 + 168p_1^2p_2^2p_3^2p_4^2 + p_2^8 + 14p_2^4p_3^4 \\ &\quad + 14p_2^4p_4^4 + p_3^8 + 14p_3^4p_4^4 + p_4^8 \\ U_4 &= \frac{1}{265531392} \det(H(U_1)). \end{aligned}$$

The values of these parameters are fixed using the same procedure applied to G_{29} , G_{32} and G_{33} . They are

$$c_1 = -\frac{3}{5}, \quad c_2 = -\frac{1}{6480}, \quad c_3 = -\frac{1}{4860}.$$

References

- [1] A. Arsie and P. Lorenzoni *F-manifolds with eventual identities, bidifferential calculus and twisted Lenard-Magri chains*, Int. Math. Res. Not. IMRN 2012 doi.10.1093/imrn/rns172.
- [2] A. Arsie and P. Lorenzoni *From Darboux-Egorov system to bi-flat F-manifolds*, Journal of Geometry and Physics, Vol 70, (2013), 98-116.
- [3] A. Arsie and P. Lorenzoni, *Purely non-local Hamiltonian formalism, Kohno connections and \vee -systems*, J. Math. Phys. 55 (2014).

- [4] A. Arsie and P. Lorenzoni *F-manifolds, multi-flat structures and Painlevé transcendents*, arXiv:1501.06435.
- [5] M. Broué, G. Malle, and R. Rouquier, *Complex Reflection groups, braid groups, Hecke Algebras*, Journal für die reine and angewandte Mathematik, **500**, (1998), 127-190.
- [6] H. Burkhardt, *Untersuchungen auf dem Gebiete der hyper-elliptischen Modulfunctionen (Zweiter Teil)*, Math. Annalen, 38 (1891), 161-224.
- [7] W. Couwenberg, G. Heckman and E. Looijenga *Geometric Structures on the Complement of a Projective Arrangement Publications Mathematiques de l'Institut des Hautes tudes Scientifiques Volume 101, Issue 1 , pp 69–161*
- [8] H.S.M. Coxeter *Regular Polytopes*, 3rd ed., Dover, New York 1973.
- [9] B.A. Dubrovin, *On almost duality for Frobenius manifolds*, Geometry, topology and mathematical physics, 75-132, Amer. Math. Soc. Transl. Ser. 2, **212**.
- [10] B.A. Dubrovin, *Differential geometry of the space of orbits of a Coxeter group*, Surv. Diff. Geom. IV (1999), 181–211.
- [11] B. Dubrovin, *Flat pencils of metrics and Frobenius manifolds*, Integrable systems and algebraic geometry (Kobe/Kyoto), 1997), 47-72, World Sci. Publishing, River Edge, NJ (1998).
- [12] C.F. Dunkl, E.M. Opdam, *Dunkl Operators for Complex Reflection Groups*, Proc. London Math. Soc. (2003) 86 (1): 70-108.
- [13] N. Elkies, *The Klein quartic in number theory*, in The Eightfold Way: The Beauty of Klein's Quartic Curve, Mathematical Sciences Research Institute publications, 35, Cambridge University Press.
- [14] M. Feigin and A. Silantyev *Singular polynomials from orbit spaces*, Compositio Mathematica, 2012, v. 148 (6), pp. 1867-1879
- [15] M. V. Feigin and A. P. Veselov, *\vee -systems, holonomy Lie algebras and logarithmic vector fields*, arXiv:1409.2424.
- [16] C. Hertling, *Frobenius manifolds and moduli spaces for singularities*, Cambridge Tracts in Mathematics 151, Cambridge University Press (2002).
- [17] C. Hertling, Y. Manin, *Weak Frobenius manifolds*, Internat. Math. Res. Notices **1999**, no. 6, 277–286.

- [18] K. Iwasaki, A. Kenma and K. Matsumoto, *Polynomial Invariants and Harmonic Functions Related to Exceptional Regular Polytopes*, Experiment. Math. 11 (2002), no. 2, 313319.
- [19] M. Kato, T. Mano and J. Sekiguchi, *Flat Structures without Potentials*, Rev. Roumaine Math. Pures Appl. **60** (2015), no. 4, 481–505
- [20] M. Kato, T. Mano and J. Sekiguchi *Flat structure on the space of isomonodromic deformations*, arXiv:1511.01608v1
- [21] F. Klein, *Ueber die transformationen siebenter Ordnung der elliptischen Funktionen*, Math. Ann. 14 (1879) 428–471.
- [22] V. G. Knizhnik and A. B. Zamolodchikov, *Current algebra and Wess- Zumino models in two dimensions*, Nuclear Phys. B **247** (1984), 83–103.
- [23] T. Kohno *Holonomy Lie algebras, logarithmic connections and the lower central series of fundamental groups*. Singularities (Iowa City, IA, 1986), 171182, Contemp. Math., 90, Amer. Math. Soc., Providence, RI, 1989.
- [24] T. Kohno, *On the holonomy Lie algebra and the nilpotent completion of the fundamental group of the complement of hypersurfaces*, Nagoya Math. J., **92** (1983), 21–37.
- [25] G. I. Lehrer and D. E. Taylor, *Unitary reflection groups*, Australian Mathematical Society Lecture Series 20, (2009), Cambridge University Press.
- [26] E. Looijenga, *Arrangements, KZ systems and Lie algebra homology*, in: Singularity Theory, B. Bruce and D. Mond eds., London Math. Soc. Lecture Note Series 263, CUP (1999), 109130.
- [27] P. Lorenzoni, M. Pedroni and A. Raimondo, *F-manifolds and integrable systems of hydrodynamic type*, Archivum Mathematicum **47** (2011), 163-180.
- [28] P. Lorenzoni, *Darboux-Egorov system, bi-flat F-manifolds and Painlevé VI*, IMRN (2014), Vol. 12, 3279-3302.
- [29] A. Losev, Y.I. Manin, *Extended modular operads*, in: Frobenius manifolds, quantum cohomology and singularities (C. Hertling and M. Marcoli, Eds.), Aspects of Math., E36, 181-211 (2004).
- [30] Y.I. Manin, *F-manifolds with flat structure and Dubrovin’s duality*, Adv. Math. **198** (2005), no. 1, 5–26.

- [31] H. Maschke, *Ueber die quaternäre, endliche, lineare Substitutionsgruppe der Borchardt'schen Moduln*, Math. Ann. **30** (1887) 496-515
- [32] H. Maschke, *Aufstellung des vollen Formensystems einer quaternären Gruppe von 51840 linearen Substitutionen*, Mathematische Annalen **33** (1888) 496-515..
- [33] M. L. Mehta, *Basic sets of invariant polynomials for finite reflection groups*, Comm. Algebra **16** (1988), 1083-1098.
- [34] P. Orlik. *Basic derivations for unitary reflection groups*, Contemporary Mathematics **90** (1989), 211-228.
- [35] P. Orlik and L. Solomon, *The hessian map in the invariant theory of reflection groups* Nagya Math. J. Vol. 109 (1988), 1-21
- [36] P. Orlik and L. Solomon, *Discriminants in the invariant theory of reflection groups*, Nagoya Math. J. **109** (1988) 23-45.
- [37] P. Orlik and H. Terao, *Arrangements of Hyperplanes*, Grundlehren der mathematischen Wissenschaften 300, Berlin, Springer-Verlag, 1992.
- [38] V. Reiner, *Reflection groups counting and q-counting*, Summer School on Algebraic and Enumerative Combinatorics, S. Miguel de Seide 2012 (available at <http://www.math.umn.edu/~reiner/Talks/Portugal2012/>).
- [39] C. Sabbah, *Isomonodromic deformations and Frobenius manifolds, An Introduction*, Universitext, Springer Verlag, (2008).
- [40] K. Saito K., T. Yano and J. Sekiguchi, *On a certain generator system of the ring of invariants of a finite reflection group*, Comm. in Algebra **8**(4) (1980) 373 - 408.
- [41] K. Saito, *On a linear structure of a quotient variety by a finite reflexion group*, RIMS Kyoto preprint 288, 1979.
- [42] K. Saito, *Period mapping associated to a primitive form*, Publ RIMS, Kyoto Univ., **19** (1983), 1231-1264.
- [43] G. C. Shephard, *Regular Complex Polytopes*, Proc. London Math. Soc, (3) **2** (1952), 82-97.
- [44] G.C. Shephard and J.A. Todd, *Finite unitary reflection groups*, Canad. J. Math. **6** (1954) 274-304
- [45] A. P. Veselov, *Deformations of the root systems and new solutions to generalised WDVV equations*, Physics Letters A, **261**, (1999), 297-302.

- [46] A. P. Veselov, *On geometry of a special class of solutions to generalized WDVV equations*, Integrability: the Seiberg-Witten and Whitham equations (Edinburgh, 1998), 125–135, Gordon and Breach.
- [47] A. Wiman, *Ueber eine einfache Gruppe von 360 ebenen Collineationen*, Math. Ann. 47 (1896) 531–556.