

Pseudo-bundles of exterior algebras as diffeological Clifford modules

Ekaterina Pervova

February 7, 2022

Abstract

We consider the diffeological pseudo-bundles of exterior algebras, and the Clifford action of the corresponding Clifford algebras, associated to a given finite-dimensional and locally trivial diffeological vector pseudo-bundle, as well as the behavior of the former three constructions (exterior algebra, Clifford action, Clifford algebra) under the diffeological gluing of pseudo-bundles. Despite these being our main object of interest, we dedicate significant attention to the issues of compatibility of pseudo-metrics, and the gluing-dual commutativity condition, that is, the condition ensuring that the dual of the result of gluing together two pseudo-bundles can equivalently be obtained by gluing together their duals, which is not automatic in the diffeological context. We show that, assuming that the dual of the gluing map, which itself does not have to be a diffeomorphism, on the total space is one, the commutativity condition is satisfied, via a natural map, which in addition turns out to be an isometry for the natural pseudo-metrics on the pseudo-bundles involved.

MSC (2010): 53C15 (primary), 57R35, 57R45 (secondary).

Introduction

This work is intended as a supplement to [8], dealing with some issues regarding pseudo-bundles of exterior algebras associated to finite-dimensional diffeological vector pseudo-bundles, viewed as pseudo-bundles of diffeological Clifford modules, so endowed with the action of the corresponding diffeological pseudo-bundles of Clifford algebras. Let us explain as briefly as possible what all these objects are; the precise definitions are to be found in a dedicated section, or else in the references given therein.

First of all, the *diffeology*. The notion is due to J.M. Souriau [14], [15] and is a categorical extension of the notion of a smooth structure; in essence, or maybe as an example, it is a way to consider a topological space which is in no way a manifold, as if it were one. In and of itself, a diffeology on a set is a collection of maps into this set which are declared to be smooth; this collection must satisfy certain conditions. The set is then a diffeological space, and all the basic constructions follow; there is a notion of smooth maps between two diffeological spaces, that of the underlying topology, the so-called *D-topology* (introduced in [4], see also [2] for a recent treatment), and so on. The notion builds on existing ones, such as those of *differentiable spaces*, *V-manifolds*, and so on (see, for instance, [13], [1], to name a few). A particularly important aspect of diffeology is that all the usual topological constructions, notably subsets and quotients, have an inherited diffeological structure (unlike the case of smooth manifolds, where subsets and quotients are quite rarely smooth manifolds themselves).

The basic object for us, though, is not just a diffeological space, but a *diffeological vector pseudo-bundle* ([4], [17], [5], [10]). The difference with respect to the standard notion is not only in the fact that the smooth structure is replaced by a diffeological one (under some respects this would be a minor difference), but also in that it does not have to be locally trivial, although in many contexts we do add this assumption, see the discussion of *pseudo-metrics*, which replace the usual Riemannian metrics — diffeological pseudo-bundles frequently do not carry the latter.

On the other hand, as explained in the references listed above, the usual operations on vector bundles have their diffeological counterparts; in particular, direct sums, tensor products, and taking duals all apply. From this, obtaining pseudo-bundles of tensor algebras, those of exterior algebras, or defining, in the abstract, pseudo-bundles of Clifford modules is automatic.

The point of view that we take in this paper has to do with studying the behavior of these concepts under the operation of *diffeological gluing*. This procedure is one of the many possible extensions of the concept of an atlas on a smooth manifold, and the resulting spaces are among the more obvious extensions of smooth manifolds and include some well-known singular spaces; for instance, a manifold with a conical singularity can be seen as a result of gluing of a usual smooth manifold to a single-point space.

For the basic operations on diffeological vector pseudo-bundles, the behavior under diffeological gluing was considered in [10] (see also [11] for some details); for tensor algebras and pairs of given Clifford modules, in [8]. What is lacking is a study of gluing of pseudo-bundles of the exterior algebras, in particular, the covariant version. This paper aims to fill this void (another motivation for it is to provide some necessary building blocks for defining the notion of a diffeological Dirac operator and studying its behavior under gluing, see [12]).

The content Section 1 goes over the main definitions used and introduces notation. In Section 2 we consider the compatibility of pseudo-metrics in terms of the assumptions on the gluing map(s); in Section 3 we relate this to the gluing-dual commutativity condition, and in Section 4 we show that the compatibility of dual pseudo-metrics implies that the commutativity condition must be satisfied. In Section 5 we show that the gluing-dual commutativity diffeomorphism is an isometry. All these allow us to consider, in Sections 6-8, Clifford algebras (the covariant case), the exterior algebras, and the corresponding Clifford actions; in particular, in Section 8 we establish, where appropriate, several equivalences showing that, again under the gluing-dual commutativity assumption, everything reduces to two cases, the contravariant case and the covariant one. Section 9 contains a couple of simple (but necessarily lengthy) examples.

Acknowledgments This work benefitted from some assistance, for which I would like to thank Prof. Riccardo Zucchi, even if he always says that he is being overvalued, some of his doctorate students (Martina and Leonardo), although they do not expect this at all, and also Prof. Mario Petrini (he most definitely will be surprised).

1 Main definitions and known facts

We now go, as briefly as possible, over the main definitions that appear in what follows (for terms whose use is not as frequent, we will provide definitions as we go along).

1.1 Diffeology and diffeological vector spaces

Let X be a set. A **diffeology** on X (see [14], [15]) is a set $\mathcal{D} = \{p : U \rightarrow X\}$ of maps into X , each defined on a domain of some \mathbb{R}^n (with varying n), that satisfies the following conditions: 1) it includes all constant maps, *i.e.*, maps of form $U \rightarrow \{x_0\}$, for all open (possibly disconnected) sets $U \subseteq \mathbb{R}^n$ and for all points $x_0 \in X$; 2) for any $\mathcal{D} \ni p : U \rightarrow X$ and for any usual smooth map $g : V \rightarrow U$ (again defined on some domain $V \subseteq \mathbb{R}^m$) we have $p \circ g \in \mathcal{D}$; and 3) if a set map $p : U \rightarrow X$ is such that its domain of definition $U \subseteq \mathbb{R}^n$ has an open cover $U = \cup_{i \in I} U_i$ for which $p|_{U_i} \in \mathcal{D}$ then $p \in \mathcal{D}$.

The maps composing \mathcal{D} are called **plots**. A standard example of diffeology/ diffeological space is a usual smooth manifold M , with diffeology composed of all usual smooth maps into M . On the other hand, any set (usually at a least a topological space) admits plenty of non-standard diffeologies, obtained via the concept of a *generated diffeology*.

Generated diffeologies Given a fixed set X , various diffeologies on it can be compared with respect to the inclusion;¹ for two diffeologies \mathcal{D} and \mathcal{D}' such that $\mathcal{D} \subset \mathcal{D}'$ one says that \mathcal{D} is **finer** than \mathcal{D}' , whereas \mathcal{D}' is said to be **coarser**. Frequently, for a given property P which a diffeology might possess, there is the finest and/or the coarsest diffeology with the property P (see [5], Sect. 1.25); this fact is often used in describing concrete diffeologies, or defining a class of them. A specific example of the former is the **generated diffeology**: for a set X and a set $A = \{p : U \rightarrow X\}$ of maps into X , the diffeology

¹In fact, each diffeology is just a set of maps, which may wholly contain another diffeology, or be contained in one; in fact, there is a complete lattice on them on any X .

generated by A is the smallest diffeology on X that contains A . Notice that A can be *any* set; it might include non-differentiable maps, discontinuous ones, and so on.

Smooth maps A map $f : X \rightarrow Y$ between two diffeological spaces X and Y is considered **smooth** if for every plot p of X the composition $f \circ p$ is a plot of Y . Note that it might easily happen that all $f \circ p$ are plots of Y , but that *vice versa* is not true: Y may have plots that do not have form $f \circ p$, whatever the plot p of X . On the other hand, if for every plot $q : U \rightarrow Y$ of Y and for every point $u \in U$ there is a plot p_u of X such that in a neighborhood of u we have $q = f \circ p_u$ then we say that the diffeology of Y is the **pushforward** of the diffeology of X via f , and conversely, the diffeology of X is the **pullback** of the diffeology of Y by f .

Subset, quotient, product, and disjoint union diffeologies All typical topological constructions admit diffeological counterparts. If X' is any subset of a diffeological space X , it carries the **subset diffeology** that consists of all plots of X whose range is contained in X' ; and if X/\sim is any² quotient of X , with $\pi : X \rightarrow X/\sim$ being the natural projection, then the standard choice of diffeology on X/\sim is the **quotient diffeology**, defined as the pushforward of the diffeology of X by π . As we said above, this means that locally each plot of X/\sim has form $\pi \circ p$, where p is a plot of X .

Let us now have several (a finite number of, although the definition can be stated more broadly) diffeological spaces X_1, \dots, X_n . Their usual direct product also has its standard diffeology, the **product diffeology**, defined as the coarsest diffeology such that the projection on each term is smooth. Locally any plot of this diffeology is just an n -tuple (p_1, \dots, p_n) , where each p_i is a plot of the corresponding X_i . Finally, the disjoint union $\sqcup_{i=1}^n X_i$ of these spaces has the **disjoint union diffeology**, this being the finest diffeology such that the inclusion of each term into the disjoint union is smooth. Locally, any plot of this diffeology is a plot of precisely one of terms X_i .

Functional diffeology The space $C^\infty(X, Y)$ of all smooth (in the diffeological sense) maps between two diffeological spaces X and Y also has its standard diffeology, called the **functional diffeology**. It consists of all possible maps $q : U \rightarrow C^\infty(X, Y)$ such that for every plot $p : U' \rightarrow X$ of X the natural evaluation map $U \times U' \ni (u, u') \mapsto q(u)(p(u')) \in Y$ is smooth (with respect to the diffeology of Y ; the product $U \times U'$ is still a domain, therefore asking for the evaluation map to be smooth is equivalent to asking it to be a plot of Y).

Diffeological vector spaces This is one specific instance of a diffeological space endowed also with an algebraic structure whose operations are smooth for the diffeology involved.³ A (real) **diffeological vector space** is a vector space V endowed with a diffeology such that the addition map $V \times V \rightarrow V$ and the scalar multiplication map $\mathbb{R} \times V \rightarrow V$ are smooth (for the product diffeology on $V \times V$ and $\mathbb{R} \times V$ respectively). For a fixed V there can be many such diffeologies; any one of them is called a **vector space diffeology** (on V). If V is finite-dimensional, and so as just a vector space is isomorphic to some \mathbb{R}^n , then the finest of all vector space diffeologies is the one consisting of all usual smooth maps into it.⁴ This diffeology is called the **standard diffeology**; endowed with it, V is called a **standard space**.

All usual operations on vector spaces (taking subspaces, quotients, direct sums, tensor products, and duals) admit their natural diffeological counterparts ([17], [19]; see also [7]), via the more general diffeological constructions described above. Thus, any vector subspace of a diffeological vector space V is automatically endowed with the subset diffeology; every quotient space carries the quotient diffeology; the direct sum carries the product diffeology (relative to the diffeologies of its terms); and the tensor product has the quotient diffeology of the finest vector space diffeology on the free product of the factors that contains the product diffeology on their direct product. Finally, the diffeological dual V^* of V is defined as the space of all diffeologically smooth linear maps $L^\infty(V, \mathbb{R})$ (where \mathbb{R} is standard) endowed with the functional diffeology. Notice that, unless a finite-dimensional V is a standard space, we have

²We stress that this can indeed be any quotient, with no restrictions on the equivalence relation \sim .

³There are analogous notions of a diffeological group, diffeological algebra, and so on.

⁴In contrast with non-vector space diffeologies, the finest of which is always the discrete one, consisting of constant maps only.

$\dim(V^*) < \dim(V)$ (and in general, the space of smooth linear maps between two diffeological vector spaces is strictly smaller than the space of all linear maps).

Pseudo-metrics and characteristic subspaces A finite-dimensional diffeological vector space V does not admit a smooth scalar product, unless it is standard (see [5]). The best possible substitute for it is any smooth symmetric semi-definite positive bilinear form of rank $\dim(V^*)$; (at least one) such a form exists on any finite-dimensional V and is called a **pseudo-metric**.

A pseudo-metric g on a finite-dimensional diffeological vector space V allows to identify in V the unique vector subspace V_0 which is maximal, with respect to inclusion, for the following two properties: the subset diffeology of V_0 is that of a standard space, and V_0 splits off smoothly in V , which means there is a usual vector space direct sum decomposition $V = V_0 \oplus V_1$ such that the corresponding direct sum diffeology on V relative to the subset diffeologies on V_0 and V_1 coincides with the initial diffeology of V .⁵ This subspace can be described as the subspace generated by all the eigenvectors of g that are relative to the positive eigenvalues; however, it does not actually depend on the specific choice of a pseudo-metric and is instead an invariant of V itself. It is called the **characteristic subspace** of V .

1.2 Diffeological vector pseudo-bundles and pseudo-metrics on them

A *diffeological vector pseudo-bundle* is a diffeological counterpart of a usual smooth vector bundle.⁶ Apart from the diffeological smoothness replacing the usual concept of smooth maps, they lack an atlas of local trivializations, although in many contexts, and in most of what follows, we do add this assumption.

Diffeological vector pseudo-bundles Let V and X be diffeological spaces, and let $\pi : V \rightarrow X$ be a smooth surjective map. The map π , or the total space V , is called a **diffeological vector pseudo-bundle** if for each $x \in X$ the pre-image $\pi^{-1}(x)$ carries a vector space structure such that the following three maps are smooth: the addition map $V \times_X V \rightarrow V$ (where $V \times_X V$ is endowed with the subset diffeology as a subset of $V \times V$), the scalar multiplication map $\mathbb{R} \times V \rightarrow V$, and the zero section $X \rightarrow V$. An example of a diffeological vector pseudo-bundle which is not locally trivial, can be found in [3] (see Example 4.3).

The fibrewise operations and fibrewise diffeologies Since each fibre of a diffeological vector pseudo-bundle is a diffeological vector space, all the usual operations on vector bundles (direct sums, tensor products, dual bundles) can be performed on/with pseudo-bundles (see [17], and also [10] for some details), although not in an entirely similar way (the lack of local trivializations prevents that). Instead, these operations are performed by first carrying out the operation in question to each fibre, defining the total space of the new pseudo-bundle as the union of the resulting diffeological vector spaces (with the obvious fibering over the base), and finally defining the diffeology of this total space as the finest that induces on each fibre its existing diffeology.⁷

As an example, and also because this instance will be particularly important for us, let us consider dual pseudo-bundles. Let $\pi : V \rightarrow X$ be a diffeological vector pseudo-bundle with finite-dimensional fibres. The dual pseudo-bundle of V is

$$V^* = \cup_{x \in X} (\pi^{-1}(x))^*,$$

where $(\pi^{-1}(x))^* = L^\infty(\pi^{-1}(x), \mathbb{R})$ is the diffeological dual of the diffeological vector space $\pi^{-1}(x)$, the pseudo-bundle projection π^* is given by $\pi^*((\pi^{-1}(x))^*) = \{x\}$ for all $x \in X$, and the diffeology on V^* is characterized as follows: a map $q : \mathbb{R}^l \supseteq U' \rightarrow V^*$ is a plot of V^* if and only if for every plot $p : \mathbb{R}^m \supseteq U \rightarrow V$ of V the evaluation map $(u', u) \mapsto q(u')(p(u)) \in \mathbb{R}$ is smooth for the subset diffeology

⁵ *A priori*, this direct-sum-diffeology-from-subset-diffeologies is finer, so not all usual direct sum decompositions of diffeological vector spaces are smooth; see an example in [9].

⁶The choice of the term *diffeological vector pseudo-bundle* is ours; the same object is called just a *diffeological fibre bundle* in [4], a *regular vector bundle* in [17], and a *diffeological vector space over X* in [3]. The choice of the term pseudo-bundle is meant to distinguish these objects from the numerous other versions of bundles that have appeared so far.

⁷This obviously poses the question of the existence of such diffeology; this was considered in [17]; see also [3], Proposition 4.6 for a relevant methodology.

on its domain of definition $\{(u', u) \mid \pi^*(q(u')) = \pi(p(u))\} \subseteq \mathbb{R}^{l+m}$ and the standard diffeology on \mathbb{R} . The collection of all possible maps q satisfying this property does form a diffeology, equipped with which, V^* becomes a diffeological vector pseudo-bundle, and furthermore, the corresponding subset diffeology on each fibre $(\pi^*)^{-1}(x)$ coincides with the usual (functional) diffeology on $(\pi^{-1}(x))^*$.

Finally, another useful observation (and maybe a peculiarity of diffeology) is that any collection of vector subspaces, one per fibre, in a diffeological vector pseudo-bundle is again a diffeological vector pseudo-bundle (called a **diffeological sub-bundle**), for the subset diffeology. Likewise, any collection of quotients, one of each fibre, is a diffeological vector pseudo-bundle for the quotient diffeology. We call these facts a peculiarity since they go well beyond what happens for the usual smooth vector bundles.

Pseudo-metrics on diffeological vector pseudo-bundles Let $\pi : V \rightarrow X$ be a diffeological vector pseudo-bundle with finite-dimensional fibres. A **pseudo-metric** on it is a smooth section of the pseudo-bundle $V^* \otimes V^*$ such that for each $x \in X$ the bilinear form $g(x)$ is a pseudo-metric, in the sense of diffeological vector spaces, on the fibre $\pi^{-1}(x)$. Not all pseudo-bundles admit a pseudo-metric (see [10]; it is not quite clear yet under which conditions a pseudo-bundle admits a pseudo-metric), although if a pseudo-bundle is locally trivial with a finite atlas of local trivializations, the reasoning similar to that in the case of usual smooth vector bundles would allow to conclude its existence on any pseudo-bundle with the above two properties.

1.3 Diffeological gluing

On the level of the underlying topological⁸ spaces, diffeological gluing (introduced in [10]) is just the usual topological gluing. The result is endowed with a canonical diffeology, called the gluing diffeology. It is the finest diffeology for several properties, and is usually finer than other natural diffeologies on the same space.

1.3.1 Gluing of spaces, maps, and pseudo-bundles

The basic ingredient in the definition of the diffeological gluing procedure is the operation of gluing of two diffeological spaces, where we must essentially specify which diffeology is assigned to the space obtained by the usual topological gluing. This basic construction is then extended to gluing of smooth maps between diffeological spaces, a particularly important instance of which is the gluing of diffeological vector pseudo-bundles.

Diffeological spaces Let X_1 and X_2 be two diffeological spaces, and let $f : X_1 \supseteq Y \rightarrow X_2$ be a map defined on a subset Y of X_1 and smooth for the subset diffeology on Y . Let

$$X_1 \cup_f X_2 := (X_1 \sqcup X_2) / \sim,$$

where the equivalence relation \sim is given by, $X_1 \sqcup X_2 \ni x_1 \sim x_2 \in X_1 \sqcup X_2$ if and only if either $x_1 = x_2$ or $x_1 \in Y$ and $x_2 = f(x_1)$. Denote by $\pi : X_1 \sqcup X_2 \rightarrow X_1 \cup_f X_2$ the quotient projection, and define the **gluing diffeology** on $X_1 \cup_f X_2$ to be the pushforward of the disjoint union diffeology on $X_1 \sqcup X_2$ by π .⁹

This construction is particularly adapted to endowing piecewise-linear objects with a diffeology (among others). An easiest example is a wedge of two lines, which can be identified with the union of the two coordinate axes in \mathbb{R}^2 . It is interesting to notice that the gluing diffeology on this union is finer than the subset diffeology relative to its inclusion into \mathbb{R}^2 , as demonstrated by an example due to Watts, see Example 2.67 in [18]. (From this, it is easy to extrapolate the existence of similar examples in other dimensions). Indeed, relative to the gluing diffeology the natural inclusions¹⁰

$$i_1 : X_1 \setminus Y \hookrightarrow X_1 \cup_f X_2 \text{ and } i_2 : X_2 \hookrightarrow X_1 \cup_f X_2$$

⁸This means the topology underlying the diffeological structure, the so-called D-topology, see [4]. In most significant examples, however, the diffeology is put on a space already carrying a topological structure, and in way such that the D-topology coincides with it.

⁹Notice that in the case when the gluing map f is invertible as map from its domain to its range, and only in this case, $X_1 \cup_f X_2$ is a span of the spaces X_1 and X_2 . However, *a priori* the gluing construction is more general.

¹⁰They are defined by composition of the obvious inclusions into $X_1 \sqcup X_2$, with the quotient projection π .

are smooth, and their ranges form a disjoint cover of $X_1 \cup_f X_2$. Notice in particular that X_1 does not in general inject into $X_1 \cup_f X_2$, while X_2 always does; in other words, *a priori* the operation of gluing is not symmetric.¹¹

This asymmetry is demonstrated, for instance, by the following (useful in practice) description of plots of the gluing diffeology. Since the latter is a pushforward diffeology, any plot of it lifts to a plot of the covering space $X_1 \sqcup X_2$. By the properties of the disjoint union diffeology, this means that if $p : U \rightarrow X_1 \cup_f X_2$ is a plot and U is connected then it either lifts to a plot p_1 of X_1 or a plot p_2 of X_2 . By construction of $X_1 \cup_f X_2$, we obtain that in the former case

$$p(u) = \begin{cases} i_1(p_1(u)) & \text{if } p_1(u) \in X_1 \setminus Y, \\ i_2(f(p_1(u))) & \text{if } p_1(u) \in Y, \end{cases}$$

while in the latter case we simply have $p = i_2 \circ p_2$.

Gluing of maps The operation of gluing of diffeological spaces, when performed on the domains and possibly the ranges of some given smooth maps, defines a gluing of these maps, provided the maps themselves satisfy a natural compatibility condition.¹² More precisely, suppose first that X_1, X_2, Z are diffeological spaces and $\varphi_i : X_i \rightarrow Z$ for $i = 1, 2$ are smooth maps. Let $f : X_1 \supseteq Y \rightarrow X_2$ be a smooth map, and suppose that $\varphi_2(f(y)) = \varphi_1(y)$ for all $y \in Y$ (the maps φ_1 and φ_2 are then said to be **f -compatible**). Then the map $\varphi_1 \cup_f \varphi_2 : X_1 \cup_f X_2 \rightarrow Z$ given by

$$(\varphi_1 \cup_f \varphi_2)(x) = \begin{cases} \varphi_1(i_1^{-1}(x)) & \text{if } x \in \text{Range}(i_1), \\ \varphi_2(i_2^{-1}(x)) & \text{if } x \in \text{Range}(i_2) \end{cases}$$

is well-defined and smooth. In fact, assigning the map $\varphi_1 \cup_f \varphi_2 \in C^\infty(X_1 \cup_f X_2, Z)$ to each pair (φ_1, φ_2) , with $\varphi_i \in C^\infty(X_i, Z)$ for $i = 1, 2$, of f -compatible maps yields a map

$$C^\infty(X_1, Z) \times_{\text{comp}} C^\infty(X_2, Z) \rightarrow C^\infty(X_1 \cup_f X_2, Z)$$

that is smooth for the functional diffeology on $C^\infty(X_1 \cup_f X_2, Z)$ and the subset diffeology¹³ on the set of f -compatible pairs $C^\infty(X_1, Z) \times_{\text{comp}} C^\infty(X_2, Z)$ (see [11] for details).

Finally, all of this extends to the case of two maps with distinct ranges, that is, $\varphi_1 : X_1 \rightarrow Z_1$ and $\varphi_2 : X_2 \rightarrow Z_2$, with appropriate gluings of X_1 to X_2 and Z_1 to Z_2 . Specifically, let again $f : X_1 \supseteq Y \rightarrow X_2$ be smooth, but consider also a smooth $g : Z_1 \supseteq \varphi_1(Y) \rightarrow Z_2$; assume that $\varphi_2(f(y)) = g(\varphi_1(y))$ for all $y \in Y$ (the maps φ_1 and φ_2 are said to be **(f, g) -compatible**). Notice that the counterparts of i_1 and i_2 for the space $Z_1 \cup_g Z_2$ are the natural inclusion maps $j_1 : Z_1 \setminus \varphi_1(Y) \hookrightarrow Z_1 \cup_g Z_2$ and $j_2 : Z_2 \rightarrow Z_1 \cup_g Z_2$. We define the map $\varphi_1 \cup_{(f,g)} \varphi_2$ by setting

$$(\varphi_1 \cup_{(f,g)} \varphi_2)(x) = \begin{cases} j_1(\varphi_1(i_1^{-1}(x))) & \text{if } x \in \text{Range}(i_1), \\ j_2(\varphi_2(i_2^{-1}(x))) & \text{if } x \in \text{Range}(i_2). \end{cases}$$

All the analogous statements, in particular, the smoothness of the thus-defined map $C^\infty(X_1, Z_1) \times_{\text{comp}} C^\infty(X_2, Z_2) \rightarrow C^\infty(X_1 \cup_f X_2, Z_1 \cup_g Z_2)$, continue to hold (see [11]).

Pseudo-bundles Let us now turn to gluing of two pseudo-bundles; this is of course a specific instance of gluing of (f, g) -compatible maps. We specifically indicate it in order to fix notation and standard terminology, and also to give conditions under which the result of gluing is again a diffeological vector pseudo-bundle.

¹¹Which is a natural consequence of the construction and also its merit, as it allows to treat, for instance, conical singularities as the results of gluing to a one-point space. For this reason, although in most cases we deal with gluings along invertible maps, hence symmetric ones, we treat them as if they were not, to keep the discussion as general as possible.

¹²According to a personal preference, one could use the language of category theory and describe the gluing operation as the pushforward of the pair of maps $\text{Id} : X_1 \rightarrow X_1$ and $f : X_1 \supseteq Y \rightarrow X_2$ to the category of smooth maps between diffeologies, that enjoys the bifunctoriality property.

¹³Relative to the product diffeology on $C^\infty(X_1, Z) \times C^\infty(X_2, Z)$ which in turn comes from the functional diffeologies on $C^\infty(X_1, Z)$ and $C^\infty(X_2, Z)$.

Let $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$ be two such pseudo-bundles, let $f : X_1 \supseteq Y \rightarrow X_2$ be a smooth map, and let $\tilde{f} : V_1 \supseteq \pi_1^{-1}(Y) \rightarrow V_2$ be any smooth lift of f that is linear on each fibre (where it is defined). The gluing of these pseudo-bundles consists in the already-defined operations of gluing V_1 to V_2 along \tilde{f} , gluing X_1 to X_2 along f , and π_1 to π_2 along (\tilde{f}, f) . It is easy to check (see [10]) that the resulting map $\pi_1 \cup_{(\tilde{f}, f)} \pi_2 : V_1 \cup_{\tilde{f}} V_2 \rightarrow X_1 \cup_f X_2$ is a diffeological vector pseudo-bundle for the gluing diffeologies on $V_1 \cup_{\tilde{f}} V_2$ and $X_1 \cup_f X_2$; in particular, the vector space structure on its fibres is inherited from either V_1 or V_2 (more precisely, it is inherited from V_1 on fibres over the points in $i_1(X_1 \setminus Y)$, and from V_2 on fibres over the points in $i_2(X_2)$).

Standard notation for gluing of pseudo-bundles We now fix some standard notation that applies to pseudo-bundles specifically. We have already described the standard inclusions

$$i_1 : X_1 \setminus Y \hookrightarrow X_1 \cup_f X_2, \quad i_2 : X_2 \hookrightarrow X_1 \cup_f X_2,$$

$$j_1 : V_1 \setminus \pi_1^{-1}(Y) \hookrightarrow V_1 \cup_{\tilde{f}} V_2, \quad j_2 : V_2 \hookrightarrow V_1 \cup_{\tilde{f}} V_2.$$

When dealing with more than one gluing at a time, we will need a more complicated notation, which is as follows. Let $\chi_1 \cup_{(\tilde{h}, h)} \chi_2 : W_1 \cup_{\tilde{h}} W_2 \rightarrow Z_1 \cup_h Z_2$ be any pseudo-bundle obtained by gluing; denote by $Y' \subset Z_1$ the domain of definition of h . Then the counterparts of i_1 and i_2 will be denoted by¹⁴

$$i_1^{Z_1} : Z_1 \setminus Y' \hookrightarrow Z_1 \cup_h Z_2 \quad \text{and} \quad i_2^{Z_2} : Z_2 \hookrightarrow Z_1 \cup_h Z_2 \quad \text{for} \quad Z_1 \cup_h Z_2$$

$$j_1^{W_1} : W_1 \setminus \chi_1^{-1}(Y') \hookrightarrow W_1 \cup_{\tilde{h}} W_2 \quad \text{and} \quad j_2^{W_2} : W_2 \hookrightarrow W_1 \cup_{\tilde{h}} W_2 \quad \text{for} \quad W_1 \cup_{\tilde{h}} W_2.$$

Obviously, $i_1^{W_1}$ and $j_1^{W_1}$ would mean the same thing, and the same goes for $i_2^{W_2}$ and $j_2^{W_2}$; however, we use the different letters so that there always be a clear distinction between the base space and the total space. Finally, since the base space will be the same for all our pseudo-bundles, and so we will usually use the abbreviated notation i_1, i_2 for it.

The switch map As we mentioned many times already, the operation of gluing for diffeological spaces is asymmetric. However, if we assume that gluing map f is a diffeomorphism with its image then obviously, we can use its inverse to perform the gluing in the reverse order, with the two results, $X_1 \cup_f X_2$ and $X_2 \cup_{f^{-1}} X_1$, being canonically diffeomorphic via the so-called **switch map**

$$\varphi_{X_1 \leftrightarrow X_2} : X_1 \cup_f X_2 \rightarrow X_2 \cup_{f^{-1}} X_1.$$

Using the notation just introduced, this map can be described by

$$\begin{cases} \varphi_{X_1 \leftrightarrow X_2}(i_1^{X_1}(x)) = i_2^{X_1}(x) & \text{for } x \in X_1 \setminus Y, \\ \varphi_{X_1 \leftrightarrow X_2}(i_2^{X_2}(f(x))) = i_1^{X_1}(x) & \text{for } x \in Y, \\ \varphi_{X_1 \leftrightarrow X_2}(i_2^{X_2}(x)) = i_1^{X_2}(x) & \text{for } x \in X_2 \setminus f(Y). \end{cases}$$

This is well-defined, not only because the maps $i_1^{X_1}$ and $i_2^{X_2}$ are injective with disjoint ranges covering $X_1 \cup_f X_2$, but also because f is a diffeomorphism with its image.

1.3.2 Gluing and operations

Diffeological gluing of pseudo-bundles is relatively well-behaved with respect to the usual operations on vector bundles. More precisely, it commutes with the direct sum and the tensor product, while the situation is somewhat more complicated for the dual pseudo-bundles, see [10] (the facts needed are recalled below).

¹⁴The rule-of-thumb is that i_1^{smth} and i_2^{smth} are used respectively for the left-hand and the right-hand factor in the base space of the pseudo-bundle under consideration, while j_1^{smth} and j_2^{smth} refer to the left-hand and the right-hand factor of the total space.

Direct sum Gluing of diffeological vector pseudo-bundles commutes with the direct sum in the following sense. Given a gluing along (\tilde{f}, f) of a pseudo-bundle $\pi_1 : V_1 \rightarrow X_1$ to a pseudo-bundle $\pi_2 : V_2 \rightarrow X_2$, as well as a gluing along (\tilde{f}', f) of a pseudo-bundle $\pi'_1 : V'_1 \rightarrow X_1$ to a pseudo-bundle $\pi'_2 : V'_2 \rightarrow X_2$, there are two natural pseudo-bundles that can be formed from these by applying the operations of gluing and direct sum. These are the pseudo-bundles

$$\begin{aligned} &(\pi_1 \cup_{(\tilde{f}, f)} \pi_2) \oplus (\pi'_1 \cup_{(\tilde{f}', f)} \pi'_2) : (V_1 \cup_{\tilde{f}} V_2) \oplus (V'_1 \cup_{\tilde{f}'} V'_2) \rightarrow X_1 \cup_f X_2 \text{ and} \\ &(\pi_1 \oplus \pi'_1) \cup_{(\tilde{f} \oplus \tilde{f}', f)} (\pi_2 \oplus \pi'_2) : (V_1 \oplus V'_1) \cup_{\tilde{f} \oplus \tilde{f}'} (V_2 \oplus V'_2) \rightarrow X_1 \cup_f X_2; \end{aligned}$$

they are diffeomorphic as pseudo-bundles, that is, there exists a fibrewise linear diffeomorphism

$$\Phi_{\cup, \oplus} : (V_1 \cup_{\tilde{f}} V_2) \oplus (V'_1 \cup_{\tilde{f}'} V'_2) \rightarrow (V_1 \oplus V'_1) \cup_{\tilde{f} \oplus \tilde{f}'} (V_2 \oplus V'_2)$$

(see below) that covers the identity map on the base $X_1 \cup_f X_2$.

Tensor product What has just been said about the direct sum, applies equally well to the tensor product. Specifically, the two possible pseudo-bundles are

$$\begin{aligned} &(\pi_1 \cup_{(\tilde{f}, f)} \pi_2) \otimes (\pi'_1 \cup_{(\tilde{f}', f)} \pi'_2) : (V_1 \cup_{\tilde{f}} V_2) \otimes (V'_1 \cup_{\tilde{f}'} V'_2) \rightarrow X_1 \cup_f X_2 \text{ and} \\ &(\pi_1 \otimes \pi'_1) \cup_{(\tilde{f} \otimes \tilde{f}', f)} (\pi_2 \otimes \pi'_2) : (V_1 \otimes V'_1) \cup_{\tilde{f} \otimes \tilde{f}'} (V_2 \otimes V'_2) \rightarrow X_1 \cup_f X_2, \end{aligned}$$

and again, they are pseudo-bundle-diffeomorphic via

$$\Phi_{\cup, \otimes} : (V_1 \cup_{\tilde{f}} V_2) \otimes (V'_1 \cup_{\tilde{f}'} V'_2) \rightarrow (V_1 \otimes V'_1) \cup_{\tilde{f} \otimes \tilde{f}'} (V_2 \otimes V'_2)$$

covering the identity on $X_1 \cup_f X_2$.

The dual pseudo-bundle The case of dual pseudo-bundles is substantially different. For one thing, to even make sense of the commutativity question, we must assume that f is invertible (which in general it does not have to be). Moreover, even with this assumption, in general the operation of gluing does *not* commute with that of taking duals. The reason of this is easy to explain. Let $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$ be two diffeological vector pseudo-bundles, and let (\tilde{f}, f) be a gluing between them; consider the pseudo-bundle $\pi_1 \cup_{(\tilde{f}, f)} \pi_2 : V_1 \cup_{\tilde{f}} V_2 \rightarrow X_1 \cup_f X_2$ and the corresponding dual pseudo-bundle

$$(\pi_1 \cup_{(\tilde{f}, f)} \pi_2)^* : (V_1 \cup_{\tilde{f}} V_2)^* \rightarrow X_1 \cup_f X_2;$$

compare it with the result of the induced gluing (performed along the pair (\tilde{f}^*, f)) of $\pi_2^* : V_2^* \rightarrow X_2$ to $\pi_1^* : V_1^* \rightarrow X_1$, that is, the pseudo-bundle

$$\pi_2^* \cup_{(\tilde{f}^*, f)} \pi_1^* : V_2^* \cup_{\tilde{f}^*} V_1^* \rightarrow X_2 \cup_{f^{-1}} X_1.$$

It then follows from the construction itself that for any $y \in Y$ (the domain of gluing) we have

$$((\pi_1 \cup_{(\tilde{f}, f)} \pi_2)^*)^{-1}(i_2^{X_2}(f(y))) \cong (\pi_2^{-1}(f(y)))^* \text{ and } (\pi_2^* \cup_{(\tilde{f}^*, f)} \pi_1^*)^{-1}(i_2^{X_1}(y)) \cong (\pi_1^{-1}(y))^*;$$

since $i_2^{X_2}(f(y))$ and $i_2^{X_1}(y)$ are related by the switch map, for the two pseudo-bundles to be diffeomorphic in a natural way¹⁵ the two vector spaces $(\pi_2^{-1}(f(y)))^*$ and $(\pi_1^{-1}(y))^*$ must be diffeomorphic, and *a priori* they are not.¹⁶

Thus, we obtain the necessary condition (for the pseudo-bundles $(V_1 \cup_{\tilde{f}} V_2)^*$ and $V_2^* \cup_{\tilde{f}^*} V_1^*$ to be diffeomorphic), which is that $(\pi_2^{-1}(f(y)))^* \cong (\pi_1^{-1}(y))^*$ for all $y \in Y$. We do note right away that this condition may not be sufficient, in the sense that two pseudo-bundles over the same base may have all respective fibres diffeomorphic without being diffeomorphic themselves (this can be illustrated by the standard example of open annulus and open Möbius strip, both of which, equipped with the standard diffeology¹⁷ can be seen as pseudo-bundles over the circle). We will leave it at that for now, returning to the issue of the gluing-dual commutativity later in the paper.

¹⁵For us this means, via a diffeomorphism covering the switch map.

¹⁶They may have different dimensions.

¹⁷That is, one determined by their usual smooth structure.

1.3.3 The commutativity diffeomorphisms

We now say more about the commutativity diffeomorphisms mentioned in the previous section.¹⁸

The diffeomorphism $\Phi_{\cup, \oplus}$ We have already mentioned the existence of this diffeomorphism, which is a pseudo-bundle map

$$\Phi_{\cup, \oplus} : (V_1 \cup_{\tilde{f}} V_2) \oplus (V'_1 \cup_{\tilde{f}',} V'_2) \rightarrow (V_1 \oplus V'_1) \cup_{\tilde{f} \oplus \tilde{f}',} (V_2 \oplus V'_2)$$

that covers the identity map on $X_1 \cup_f X_2$. We now add that this map can be described (in fact, fully defined) by the following identities:

$$\Phi_{\cup, \oplus} \circ (j_1^{V_1} \oplus j_1^{V'_1}) = j_1^{V_1 \oplus V'_1} \quad \text{and} \quad \Phi_{\cup, \oplus} \circ (j_2^{V_2} \oplus j_2^{V'_2}) = j_2^{V_2 \oplus V'_2}.$$

The diffeomorphism $\Phi_{\cup, \otimes}$ Once again, the case of the tensor product is very similar to that of the direct sum. The already-mentioned diffeomorphism

$$\Phi_{\cup, \otimes} : (V_1 \cup_{\tilde{f}} V_2) \otimes (V'_1 \cup_{\tilde{f}',} V'_2) \rightarrow (V_1 \otimes V'_1) \cup_{\tilde{f} \otimes \tilde{f}',} (V_2 \otimes V'_2)$$

is uniquely determined by the identities

$$\Phi_{\cup, \otimes} \circ (j_1^{V_1} \otimes j_1^{V'_1}) = j_1^{V_1 \otimes V'_1} \quad \text{and} \quad \Phi_{\cup, \otimes} \circ (j_2^{V_2} \otimes j_2^{V'_2}) = j_2^{V_2 \otimes V'_2};$$

notice that these, themselves, suffice to ensure that it covers the identity map on $X_1 \cup_f X_2$.

The gluing-dual commutativity conditions, and diffeomorphism $\Phi_{\cup, *}$ We have already described the situation regarding the gluing-dual commutativity, first of all the fact, that it is far from being always present. At this moment we concentrate on what it actually means for the gluing to commute with taking duals (once again leaving aside the question when this does happen). Specifically, we say that the **gluing-dual commutativity condition** holds, if there exists a diffeomorphism

$$\Phi_{\cup, *} : (V_1 \cup_{\tilde{f}} V_2)^* \rightarrow V_2^* \cup_{\tilde{f}^*} V_1^*$$

that covers the switch map, that is,

$$(\pi_2^* \cup_{(\tilde{f}^*, f^{-1})} \pi_1^*) \circ \Phi_{\cup, *} = \varphi_{X_1 \leftrightarrow X_2} \circ (\pi_1 \cup_{(\tilde{f}, f)} \pi_2)^*,$$

and such that the following are true:

$$\left\{ \begin{array}{ll} \Phi_{\cup, *} \circ ((j_1^{V_1})^*)^{-1} = j_2^{V_1^*} & \text{on } (\pi_2^* \cup_{(\tilde{f}^*, f^{-1})} \pi_1^*)^{-1}(i_2^{X_1}(X_1 \setminus Y)), \\ \Phi_{\cup, *} \circ ((j_2^{V_2})^*)^{-1} = j_2^{V_1^*} \circ \tilde{f}^* & \text{on } (\pi_2^* \cup_{(\tilde{f}^*, f^{-1})} \pi_1^*)^{-1}(i_2^{X_1}(Y)), \\ \Phi_{\cup, *} \circ ((j_2^{V_2})^*)^{-1} = j_1^{V_2^*} & \text{on } (\pi_2^* \cup_{(\tilde{f}^*, f^{-1})} \pi_1^*)^{-1}(i_1^{X_2}(X_2 \setminus f(Y))). \end{array} \right.$$

1.3.4 Gluing and pseudo-metrics

The behavior of pseudo-metrics under gluing depends significantly on whether the gluing-dual commutativity condition is satisfied. More precisely, if we glue together two pseudo-bundles carrying a pseudo-metric each, then under a certain natural compatibility condition (see below) for these pseudo-metrics, the new pseudo-bundle carries a pseudo-metric as well; but how the latter is constructed depends on the existence of the commutativity diffeomorphism $\Phi_{\cup, *}$.

¹⁸If one wishes, these could be described as universal factorization properties in the category of diffeological pseudo-bundles

The compatibility notion for pseudo-metrics Let $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$ be two finite-dimensional diffeological vector pseudo-bundles, endowed with pseudo-metrics g_1 and g_2 respectively, and let (\tilde{f}, f) be a pair of smooth maps that defines gluing of X_1 to X_2 ; let $Y \subseteq X_1$ be the domain of definition of f . The pseudo-metrics g_1 and g_2 are said to be **compatible with (the gluing along) the pair (\tilde{f}, f)** if for all $y \in Y$ and for all $v, w \in \pi_1^{-1}(y)$ the following is true:

$$g_1(y)(v, w) = g_2(f(y))(\tilde{f}(v), \tilde{f}(w)).$$

If f is invertible, this means that g_2 and g_1 are $(f^{-1}, \tilde{f}^* \otimes \tilde{f}^*)$ -compatible as smooth maps $X_2 \rightarrow V_2^* \otimes V_2^*$ and $X_1 \rightarrow V_1^* \otimes V_1^*$ respectively.

As can be expected,¹⁹ the existence of compatible pseudo-metrics on two pseudo-bundles imposes substantial restrictions on their fibres over the domain of gluing. Later in the paper we will make precise statements to this effect.

The induced pseudo-metric in the presence of $\Phi_{U,*}$ If we assume that the gluing-dual commutativity condition is satisfied, this implies also that f is invertible (with smooth inverse). In this case we can use the map

$$g_2 \cup_{(f^{-1}, \tilde{f}^* \otimes \tilde{f}^*)} g_1 : X_2 \cup_{f^{-1}} X_1 \rightarrow (V_2^* \otimes V_2^*) \cup_{\tilde{f}^* \otimes \tilde{f}^*} (V_1^* \otimes V_1^*)$$

to construct a pseudo-metric on $V_1 \cup_{\tilde{f}} V_2$ by taking the following composition of it with the switch map and the commutativity diffeomorphisms:

$$\tilde{g} = (\Phi_{U,*}^{-1} \otimes \Phi_{U,*}^{-1}) \circ \Phi_{\otimes,U} \circ (g_2 \cup_{(f^{-1}, \tilde{f}^* \otimes \tilde{f}^*)} g_1) \circ \varphi_{X_1 \leftrightarrow X_2},$$

where $\varphi_{X_1 \leftrightarrow X_2}$ is the switch map, $\Phi_{U,*}$ (of which we need the inverse) is the just-seen gluing-dual commutativity diffeomorphism, while $\Phi_{\otimes,U} : (V_2^* \otimes V_2^*) \cup_{\tilde{f}^* \otimes \tilde{f}^*} (V_1^* \otimes V_1^*) \rightarrow (V_2^* \cup_{\tilde{f}^*} V_1^*) \otimes (V_2^* \cup_{\tilde{f}^*} V_1^*)$ is the appropriate version of the tensor product-gluing commutativity diffeomorphism.

Constructing a pseudo-metric on $V_1 \cup_{\tilde{f}} V_2$ when $\Phi_{U,*}$ does not exist Although we will mostly deal with the cases where the gluing-dual commutativity condition is present (and so the above definition of the pseudo-metric \tilde{g} on $V_1 \cup_{\tilde{f}} V_2$ is sufficient), we briefly mention that even if such condition does not hold, the flexibility of diffeology allows for a direct construction of a pseudo-metric on $V_1 \cup_{\tilde{f}} V_2$. This construction uses the fact that each fibre of $V_1 \cup_{\tilde{f}} V_2$ is naturally identified with one of either V_1 or V_2 ; accordingly, \tilde{g} can be defined to coincide with either g_1 or g_2 on each fibre individually. The surprising fact is that \tilde{g} coming from this construction is still diffeologically smooth across the fibres; see [11] for details.

1.4 The pseudo-bundle of smooth linear maps

The last more-or-less standard construction that we need is that of **pseudo-bundle of smooth linear maps**. Let $\pi_1 : V_1 \rightarrow X$ and $\pi_2 : V_2 \rightarrow X$ be two finite-dimensional diffeological vector pseudo-bundles with the same space X . For every $x \in X$ the space $L^\infty(\pi_1^{-1}(x), \pi_2^{-1}(x))$ of smooth linear maps $\pi_1^{-1}(x) \rightarrow \pi_2^{-1}(x)$ is a (finite-dimensional) diffeological vector space for the functional diffeology. The union

$$\mathcal{L}(V_1, V_2) = \cup_{x \in X} L^\infty(\pi_1^{-1}(x), \pi_2^{-1}(x))$$

of all these spaces has the obvious projection (denoted π^L) to X , and the pre-image of each point under this projection has vector space structure. It becomes a diffeological vector pseudo-bundle when endowed with the **pseudo-bundle functional diffeology**, that is defined as the finest diffeology containing all maps $p : U \rightarrow \mathcal{L}(V_1, V_2)$, with $U \subseteq \mathbb{R}^m$ an arbitrary domain, that possess the following property: for every plot $q : \mathbb{R}^{m'} \supseteq U' \rightarrow V_1$ of V_1 the corresponding evaluation map $(u, u') \mapsto p(u)(q(u')) \in V_2$ defined on $Y' = \{(u, u') \mid \pi^L(p(u)) = \pi_1(q(u'))\} \subset U \times U'$ is smooth for the subset diffeology of Y' . This is the type of object where the Clifford actions live.

¹⁹The notion of a pseudo-metric is designed to be a generalization from a scalar product on a vector space, where the compatibility with a given f is equivalent to f being an isometry (of its domain with its range).

1.5 The pseudo-bundles of Clifford algebras and Clifford modules

For diffeological pseudo-bundles, these have already been described in the abstract setting (see [8]). We briefly summarize the main points that appear therein, noting that the main conclusions do not differ from the usual case, or are as expected anyhow.

1.5.1 The pseudo-bundle $cl(V, g)$

Let $\pi : V \rightarrow X$ be a finite-dimensional diffeological pseudo-bundle endowed with a pseudo-metric g . The construction of the corresponding pseudo-bundle $cl(V, g)$ of Clifford algebras is the immediate one, since all the operations involved have already been described. Specifically, the **pseudo-bundle of Clifford algebras** $\pi^{Cl} : cl(V, g) \rightarrow X$ is given by

$$cl(V, g) := \cup_{x \in X} cl(\pi^{-1}(x), g(x))$$

and is endowed with the quotient diffeology coming from **pseudo-bundle of tensor algebras** $\pi^{T(V)} : T(V) \rightarrow X$. The latter pseudo-bundle has total space given by $T(V) := \cup_{x \in X} T(\pi^{-1}(x))$, where $T(\pi^{-1}(x)) := \bigoplus_r (\pi^{-1}(x))^{\otimes r}$ is the usual tensor algebra of the diffeological vector space $\pi^{-1}(x)$ (in particular, it is endowed with the vector space direct sum diffeology relative to the tensor product diffeology on each factor).

Remark 1.1. *We will not make much use of the algebra structure on this pseudo-bundle, and so will actually consider all the direct sums involved to be finite (limited by the maximum of the dimensions of fibres of V in question — this includes the assumption that such maximum exists), thus considering, instead of the whole $T(V)$ its finite-dimensional sub-bundle $T_{\leq n}(V)$, with fibre at x the space $T_{\leq n}(\pi^{-1}(x))$ consisting of all tensors in $T(\pi^{-1}(x))$ of degree at most n . These fibres are not algebras, but of course each of them is a vector subspace of the corresponding fibre of $T(V)$.*

Recall that the subset diffeology on each fibre of $T(V)$ is that of the tensor algebra²⁰ of the individual fibre $\pi^{-1}(x)$. In each such fibre we choose the subspace W_x that is the kernel of the universal map $T(\pi^{-1}(x)) \rightarrow Cl(\pi^{-1}(x), g(x))$. Then, as is generally the case, $W = \cup_{x \in X} W_x \subset T(V)$ endowed with the subset diffeology relative to this inclusion is a sub-bundle of $T(V)$. The fibre of the corresponding quotient pseudo-bundle at any given point $x \in X$ is $cl(\pi^{-1}(x), g(x))$, and the quotient diffeology on the fibre is that of the Clifford algebra over the vector space $\pi^{-1}(x)$. This is exactly the diffeology that we endow $Cl(V, g)$ with.

1.5.2 The pseudo-bundle $Cl(V_1 \cup_{\tilde{f}} V_2, \tilde{g})$ as the result of a gluing

The main result, that we immediately state and that appears in [8], is the following one.

Theorem 1.2. *Let $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$ be two diffeological vector pseudo-bundles, let (\tilde{f}, f) be two maps defining a gluing between these two pseudo-bundles, both of which are diffeomorphisms, and let g_1 and g_2 be pseudo-metrics on V_1 and V_2 respectively, compatible with this gluing. Let \tilde{g} be the pseudo-metric on $V_1 \cup_{\tilde{f}} V_2$ induced by g_1 and g_2 . Then there exists a map \tilde{F}^{Cl} defining a gluing of the pseudo-bundles $Cl(V_1, g_1)$ and $Cl(V_2, g_2)$, and a diffeomorphism Φ^{Cl} between the pseudo-bundles $Cl(V_1, g_1) \cup_{\tilde{F}^{Cl}} Cl(V_2, g_2)$ and $Cl(V_1 \cup_{\tilde{f}} V_2, \tilde{g})$ covering the identity on $X_1 \cup_f X_2$.*

Let us briefly describe the maps \tilde{F}^{Cl} and Φ^{Cl} . The construction of \tilde{F}^{Cl} is the immediately obvious one. It is defined on each fibre over a point $y \in Y$ as the map $Cl(\pi_1^{-1}(y), g_1|_{\pi_1^{-1}(y)}) \rightarrow Cl(\pi_2^{-1}(f(y)), g_2|_{\pi_2^{-1}(f(y))})$ induced by \tilde{f} via the universal property of Clifford algebras. In practice, this means that on each fibre \tilde{F}^{Cl} is linear and multiplicative (with respect to the tensor product), so if $v_1 \otimes \dots \otimes v_k$ is a representative of an equivalence class in $Cl(\pi_1^{-1}(y), g_1|_{\pi_1^{-1}(y)})$ (viewed as the appropriate quotient of $T(\pi_1^{-1}(y))$) then by this definition

$$\tilde{F}^{Cl}(v_1 \otimes \dots \otimes v_k) = \tilde{F}^{Cl}(v_1) \otimes \dots \otimes \tilde{F}^{Cl}(v_k) = \tilde{f}(v_1) \otimes \dots \otimes \tilde{f}(v_k).$$

²⁰Or of its appropriate vector subspace, see the remark above.

That this is well-defined as a map $\mathcal{C}\ell(\pi_1^{-1}(y), g_1|_{\pi_1^{-1}(y)}) \rightarrow \mathcal{C}\ell(\pi_2^{-1}(f(y)), g_2|_{\pi_2^{-1}(f(y))})$ follows from the compatibility of pseudo-metrics g_1 and g_2 . Indeed, if $v, w \in \pi_1^{-1}(y)$ then

$$\tilde{F}^{\mathcal{C}\ell}(v \otimes w + w \otimes v + 2g_1(y)(v, w)) = \tilde{f}(v) \otimes \tilde{f}(w) + \tilde{f}(w) \otimes \tilde{f}(v) + 2g_1(y)(v, w)$$

by the above formula, and $2g_1(y)(v, w) = 2g_2(f(y))(\tilde{f}(v), \tilde{f}(w))$ by the compatibility. Thus, $\tilde{F}^{\mathcal{C}\ell}$ preserves the defining relation for Clifford algebras, so indeed it is well-defined (on each fibre; hence on the whole pseudo-bundle).

The diffeomorphism $\Phi^{\mathcal{C}\ell}$, which we specify to be $\mathcal{C}\ell(V_1, g_1) \cup_{\tilde{F}^{\mathcal{C}\ell}} \mathcal{C}\ell(V_2, g_2) \rightarrow \mathcal{C}\ell(V_1 \cup_{\tilde{f}} V_2, \tilde{g})$, is then the natural identification; namely, by definitions of the gluing operation and that of the induced pseudo-metric \tilde{g} , over a point of form $x = i_1^{X_1}(x_1)$ both the fibre of $\mathcal{C}\ell(V_1, g_1) \cup_{\tilde{F}^{\mathcal{C}\ell}} \mathcal{C}\ell(V_2, g_2)$ and that of $\mathcal{C}\ell(V_1 \cup_{\tilde{f}} V_2, \tilde{g})$ are naturally identified with $\mathcal{C}\ell(\pi_1^{-1}(x), g_1|_{\pi_1^{-1}(x)})$, while over any point of form $x = i_2^{X_2}(x_2)$ they are identified with $\mathcal{C}\ell(\pi_2^{-1}(x), g_2|_{\pi_2^{-1}(x)})$.

1.5.3 Gluing of pseudo-bundles of Clifford modules

A statement similar to that of the Theorem cited in the previous section can also be obtained for given Clifford modules over the algebras $\mathcal{C}\ell(V_1, g_1)$ and $\mathcal{C}\ell(V_2, g_2)$. This requires some additional assumptions on these modules, and the appropriate notion of the compatibility of the actions.

The two Clifford modules Let $\pi_1 : V_1 \rightarrow X_1$, $\pi_2 : V_2 \rightarrow X_2$, (\tilde{f}, f) , g_1 , and g_2 be as in Theorem 1.2. Recall that this yields the following pseudo-bundles, $\pi_1^{\mathcal{C}\ell} : \mathcal{C}\ell(V_1, g_1) \rightarrow X_1$, $\pi_2^{\mathcal{C}\ell} : \mathcal{C}\ell(V_2, g_2) \rightarrow X_2$, and $\pi_1^{\mathcal{C}\ell} \cup_{(\tilde{F}^{\mathcal{C}\ell}, f)} \pi_2^{\mathcal{C}\ell} : \mathcal{C}\ell(V_1, g_1) \cup_{\tilde{F}^{\mathcal{C}\ell}} \mathcal{C}\ell(V_2, g_2) = \mathcal{C}\ell(V_1 \cup_{\tilde{f}} V_2, \tilde{g}) \rightarrow X_1 \cup_f X_2$.

Now suppose that we are given two pseudo-bundles of Clifford modules, $\chi_1 : E_1 \rightarrow X_1$ and $\chi_2 : E_2 \rightarrow X_2$ (over $\mathcal{C}\ell(V_1, g_1)$ and $\mathcal{C}\ell(V_2, g_2)$ respectively), that is, there is a smooth pseudo-bundle map $c_i : \mathcal{C}\ell(V_i, g_i) \rightarrow \mathcal{L}(E_i, E_i)$ that covers the identity on the bases. Suppose further that there is a smooth fibrewise linear map $\tilde{f}' : \chi_1^{-1}(Y) \rightarrow \chi_2^{-1}(f(Y))$ that covers f . We describe the pseudo-bundle $E_1 \cup_{\tilde{f}'} E_2$ as a Clifford module over $\mathcal{C}\ell(V_1 \cup_{\tilde{f}} V_2, \tilde{g})$, with respect to an action induced by c_1 and c_2 .

Compatibility of c_1 and c_2 Similar to how it occurs for smooth maps, there is always an action of $\mathcal{C}\ell(V_1 \cup_{\tilde{f}} V_2, \tilde{g})$ on $E_1 \cup_{\tilde{f}'} E_2$ induced by c_1 and c_2 , it is smooth on each fibre but in general, it might not be smooth across the fibres. For it to be so, we need a notion of compatibility for Clifford actions, which is as follows.

Definition 1.3. *The actions c_1 and c_2 are **compatible** (with respect to $\tilde{F}^{\mathcal{C}\ell}$ and \tilde{f}') if for all $y \in Y$, for all $v \in (\pi_1^{\mathcal{C}\ell})^{-1}(y)$, and for all $e_1 \in \chi_1^{-1}(y)$ we have*

$$\tilde{f}'(c_1(v)(e_1)) = c_2(\tilde{F}^{\mathcal{C}\ell}(v))(\tilde{f}'(e_1)).$$

We note that the compatibility of c_1 and c_2 as it has been just defined, does not automatically translate into their $(\tilde{F}^{\mathcal{C}\ell}, \tilde{f}')$ compatibility as smooth maps in $\mathcal{C}\ell(V_i, g_i) \rightarrow \mathcal{L}(E_i, E_i)$; see [8] for a discussion on this.

The induced action Assuming now that the two given actions c_1 and c_2 are compatible in the sense just stated, we can define an induced action on $E_1 \cup_{\tilde{f}'} E_2$, that is, a smooth homomorphism

$$c : \mathcal{C}\ell(V_1 \cup_{\tilde{f}} V_2, \tilde{g}) \rightarrow \mathcal{L}(E_1 \cup_{\tilde{f}'} E_2, E_1 \cup_{\tilde{f}'} E_2).$$

Using the already-mentioned identification, via the diffeomorphism $\Phi^{\mathcal{C}\ell}$, of $\mathcal{C}\ell(V_1 \cup_{\tilde{f}} V_2, \tilde{g})$ with $\mathcal{C}\ell(V_1, g_1) \cup_{\tilde{F}^{\mathcal{C}\ell}} \mathcal{C}\ell(V_2, g_2)$, the action c can be described by defining first

$$c'(v)(e) = \begin{cases} j_1^{E_1} \left(c_1((j_1^{\mathcal{C}\ell(V_1, g_1)})^{-1}(v))((j_1^{E_1})^{-1}(e)) \right) & \text{if } v \in \text{Im}(j_1^{\mathcal{C}\ell(V_1, g_1)}) \Rightarrow e \in \text{Im}(j_1^{E_1}), \\ j_2^{E_2} \left(c_2((j_2^{\mathcal{C}\ell(V_2, g_2)})^{-1}(v))((j_2^{E_2})^{-1}(e)) \right) & \text{if } v \in \text{Im}(j_2^{\mathcal{C}\ell(V_2, g_2)}) \Rightarrow e \in \text{Im}(j_2^{E_2}). \end{cases}$$

Since the images of the inductions $j_1^{Cl(V_1, g_1)}$ and $j_2^{Cl(V_2, g_2)}$ are disjoint and cover $Cl(V_1, g_1) \cup_{\tilde{F}Cl} Cl(V_2, g_2)$, and those of $j_1^{E_1}$ and $j_2^{E_2}$ cover $E_1 \cup_{\tilde{F}} E_2$ (and are disjoint as well), this is a well-defined fibrewise action of the former on the latter. From the formal point of view, we must also pre-compose it with the inverse of Φ^{Cl} , to obtain an action

$$c = c' \circ (\Phi^{Cl})^{-1} : Cl(V_1 \cup_{\tilde{F}} V_2, \tilde{g}) \rightarrow \mathcal{L}(E_1 \cup_{\tilde{F}} E_2, E_1 \cup_{\tilde{F}} E_2).$$

Then, the following is true (see [8]).

Theorem 1.4. *The action c is smooth as a map $Cl(V_1 \cup_{\tilde{F}} V_2, \tilde{g}) \rightarrow \mathcal{L}(E_1 \cup_{\tilde{F}} E_2, E_1 \cup_{\tilde{F}} E_2)$.*

2 The induced pseudo-metrics on dual pseudo-bundles

Let $\pi : V \rightarrow X$ be a locally trivial finite-dimensional diffeological vector pseudo-bundle endowed with a pseudo-metric g ; then its dual pseudo-bundle $\pi^* : V^* \rightarrow X$ admits an induced pseudo-metric g^* (see [11]; we also recall the definition below). In this section we consider gluings of two pseudo-bundles endowed with compatible pseudo-metrics, and the corresponding dual constructions; starting from indicating which restrictions are imposed in the initial pseudo-bundles by the existence of compatible pseudo-metrics on them, we proceed to discuss when the induced pseudo-metrics on the dual pseudo-bundles are compatible in their turn, and finally, how it relates to the gluing-dual commutativity condition.

2.1 The induced pseudo-metric on V^*

Let $\pi : V \rightarrow X$ be a locally trivial finite-dimensional diffeological vector pseudo-bundle, and let g be a pseudo-metric on it. Under the assumption of local triviality of V ,²¹ the dual pseudo-bundle V^* carries an induced pseudo-metric g^* , which is obtained by what can be considered a diffeological counterpart of the usual natural pairing.

Specifically, consider the pseudo-bundle map $\Phi : V \rightarrow V^*$ defined by

$$\Phi(v) = g(\pi(v))(v, \cdot) \text{ for all } v \in V.$$

It is not hard to show (see [9] for the case of a single diffeological vector space, and then [11] for the case of pseudo-bundles) that Φ is surjective, smooth, and linear on each fibre.

The **induced pseudo-metric** g^* is given by the following equality:

$$g^*(x)(\Phi(v), \Phi(w)) := g(x)(v, w) \text{ for all } x \in X \text{ and for all } v, w \in V \text{ such that } \pi(v) = \pi(w) = x.$$

This is well-defined, because whenever $\Phi(v) = \Phi(v')$ (which obviously can occur only for v, v' belonging to the same fibre), the vectors v and v' differ by an element of the isotropic subspace of the fibre to which they belong. Furthermore, since each fibre of the dual pseudo-bundle (in the finite-dimensional case) carries the standard diffeology (see [9]), $g^*(x)$ is always a scalar product. Notice that without the requirement of local triviality, we cannot guarantee that g^* is indeed a pseudo-metric, and more precisely, that it is smooth (the map Φ always has a right inverse, which *a priori* may not be smooth).

Lemma 2.1. *Let $\pi : V \rightarrow X$ be a finite-dimensional diffeological vector pseudo-bundle endowed with a pseudo-metric g , let $\pi^* : V^* \rightarrow X$ be the dual pseudo-bundle, and let g^* be the induced pseudo-metric. Then for all $x \in X$ the symmetric bilinear form $g^*(x)$ on $(\pi^*)^{-1}(x)$ is non-degenerate.*

2.2 Existence of compatible pseudo-metrics: the case of a diffeological vector space

Before treating various issues regarding the induced pseudo-metrics, it makes sense to consider in more detail what the compatibility of two pseudo-metrics means. We do so starting with the case of just

²¹Which at the moment does not appear to be particularly limiting, in the sense that we do not know of any non-locally-trivial pseudo-bundles that admit pseudo-metrics in the first place.

diffeological vector spaces (we consider the duals of vector spaces in the section that immediately follows, and pseudo-bundles in the one after that).

Let V and W be finite-dimensional diffeological vector spaces, let g_V be a pseudo-metric on V , and let g_W be a pseudo-metric on W . We assume that we are given a smooth linear map $f : V \rightarrow W$, with respect to which g_V and g_W are compatible, $g_V(v_1, v_2) = g_W(f(v_1), f(v_2))$. We show that, quite similarly to usual vector spaces and scalar products, there are pairs of diffeological ones such that no pair of pseudo-metrics is compatible with respect to any smooth linear f . The similarity that we are referring to has to do with the fact that the compatibility of two pseudo-metrics with respect to f essentially amounts to f being a diffeological analogue of an isometry onto a subspace. As is well-known, between two usual vector spaces such isometry may not exist (it is necessary that the dimension of the domain space must be less or equal to that of the target space), and something similar happens for diffeological vector spaces; and then further conditions are added in terms of their diffeological structures.

2.2.1 The characteristic subspaces of V and W

Assuming that two given pseudo-metrics g_V and g_W on diffeological vector spaces V and W respectively are compatible with a given $f : V \rightarrow W$ has several implications for the diffeological structures of V and W ; describing these requires the following notion.

Given a pseudo-metric g on a finite-dimensional diffeological vector space V , the subspace $V_0 \leq V$ generated by all the eigenvectors of g relative to the non-zero eigenvalues has subset diffeology that is standard; and among all subspaces of V whose diffeology is standard, it has the maximal dimension, which is equal to $\dim(V^*)$. In general, V contains more than one subspace of dimension $\dim(V^*)$ whose diffeology is standard. But the subspace V_0 is the only one that also splits off as a smooth direct summand.²² Thus, V_0 does not actually depend on the choice of a pseudo-metric and is an invariant of the space itself (see [9]). We call this subspace the **characteristic subspace** of V .

Let us now return to the two diffeological vector spaces V and W above. Let V_0 and W_0 be their characteristic subspaces, and let $V_1 \leq V$ and $W_1 \leq W$ be the isotropic subspaces relative to g_V and g_W respectively, such that $V = V_0 \oplus V_1$ and $W = W_0 \oplus W_1$ with each of these decompositions being smooth. We also recall [9] that V_0 not only has the same dimension as V^* , but for any fixed pseudo-metric is diffeomorphic to it, via (the restriction to V_0 of) the map $\Phi_V : v \mapsto g_V(v, \cdot)$; likewise, W_0 is diffeomorphic to W^* via $\Phi_W : w \mapsto g_W(w, \cdot)$.

2.2.2 The necessary conditions

Let us now assume that the given g_V , g_W , and f satisfy the compatibility condition. The corollaries of this assumption can be described in terms of the characteristic subspaces of V and W , and therefore in terms of their diffeological duals.

The kernel of f The first corollary is quite trivial, and starts with a simple linear algebra argument. Let $v \in V$ belong to the kernel of f . Then by the compatibility assumption for g_V and g_W we have

$$g_V(v, v') = g_W(0, f(v')) = 0 \text{ for any } v' \in V.$$

Thus, the kernel of f is contained in the maximal isotropic subspace V_1 , therefore the restriction of f to V_0 is a bijection with its image. This restriction is of course a smooth map, and since V_0 splits off as a smooth direct summand, it is an induction (that is, a diffeomorphism with its image). Finally, f itself is a diffeomorphism of $V_0 \oplus (V_1/\text{Ker}(f))$ with its image in W . In particular, we have the following.

Lemma 2.2. *Let V and W be finite-dimensional diffeological vector spaces, and let $f : V \rightarrow W$ be a smooth linear map. If V and W admit pseudo-metrics compatible with f then the $\text{Ker}(f) \cap V_0 = \{0\}$.*

Notice that in the standard case V and W would be vector spaces, g_V and g_W scalar products on them, and f an isometry of V with its image in W . In particular, f would be injective; Lemma 2.2 is the diffeological counterpart of that.

²²Which means that the direct sum diffeology coincides with V 's or W 's own diffeology, or, alternatively, that the composition of each plot of V (respectively W) with the projection on V_0 (respectively W_0) is a plot of the latter.

The dimensions of V and W , and those of V^* and W^* Continuing the analogy with the standard case, we observe that the standard inequality $\dim(V) \leq \dim(W)$ does not have to hold in the diffeological setting. What instead is true, is the corresponding inequality for the dimensions of their dual spaces, which follows from the lemma below.

Lemma 2.3. *Let V and W be finite-dimensional diffeological vector spaces, let $f : V \rightarrow W$ be a smooth linear map, and suppose that V and W carry pseudo-metrics g_V and g_W respectively, compatible with respect to f . Then the subset diffeology of $f(V_0)$ is the standard one.*

Proof. Let e_1, \dots, e_n be a g_V -orthonormal basis of V_0 ; then by compatibility $f(e_1), \dots, f(e_n)$ is a g_W -orthonormal basis of $f(V_0)$, which can be completed to a basis of eigenvectors of g_W . It suffices to show that the projection of W on the line generated by each $f(e_i)$ is a usual smooth function. Since this projection is given by $w \mapsto g_W(f(e_i), w)$, the claim follows from the smoothness of g_W . \square

Now, the fact that $f(V_0)$ carries the standard diffeology, does not automatically imply that it is contained in W_0 — there are standard subspaces that are not (we will however show later on that this inclusion does hold for $f(V_0)$). However, $f(V_0)$ is still a standard subspace of W , and since W_0 has maximal dimension among such subspaces, we have

$$\dim(V^*) = \dim(V_0) = \dim(f(V_0)) \leq \dim(W_0) = \dim(W^*).$$

Therefore we have the following statement.

Proposition 2.4. *Let V and W be finite-dimensional diffeological vector spaces. If there exist a smooth linear map $f : V \rightarrow W$ and pseudo-metrics g_V and g_W on V and W respectively, compatible with respect to f , then*

$$\dim(V^*) \leq \dim(W^*).$$

In other words, if $\dim(V^*) > \dim(W^*)$, then no two pseudo-metrics on V and W are compatible, whatever the map f (which obviously mimics the standard situation: there is no isometry from the space of a bigger dimension to one of smaller dimension).²³

The subspace $f(V_0)$ in W We now show that the *a priori* case when $f(V_0)$ is not contained in W_0 is actually impossible, that is, if $f : V \rightarrow W$ is such that V and W admit compatible pseudo-metrics then f sends the characteristic subspace of V to the characteristic subspace of W .

Lemma 2.5. *Let V and W be finite-dimensional diffeological vector spaces, let $f : V \rightarrow W$ be a smooth linear map, and suppose that V and W admit compatible pseudo-metrics g_V and g_W respectively. Then $f(V_0)$ splits off smoothly in W .*

Proof. Let e_1, \dots, e_n be a g_V -orthonormal basis of V_0 . Then by assumption $f(e_1), \dots, f(e_n)$ is a g_W -orthonormal basis of $f(V_0)$. This can be completed to an orthogonal basis of W composed of eigenvectors of g_W ; denote by u_1, \dots, u_k the elements added, ordered in such a way that the eigenvectors corresponding to the zero eigenvalue are the last m vectors. Let us show that the usual direct sum decomposition $W = f(V_0) \oplus \text{Span}(u_1, \dots, u_k)$ is a smooth one.

Let $p : U \rightarrow W$ be a plot of W , and let p' be its composition with the projection (associated to the direct sum decomposition just mentioned) of W to $f(V_0)$. It suffices to show that p' is a plot of $f(V_0)$. Notice that by the choice of the basis $f(e_1), \dots, f(e_n), u_1, \dots, u_k$ of W (more precisely, by the g_W -orthogonality of said basis) we have

$$p'(u) = g_W(f(e_1), p(u))f(e_1) + \dots + g_W(f(e_n), p(u))f(e_n),$$

where each coefficient $g_W(f(e_i), p(u))$ is an ordinary smooth function $U \rightarrow \mathbb{R}$ by the smoothness of the pseudo-metric g_W . This means precisely that p' is a plot of $f(V_0)$, whence the claim. \square

From the lemma just proven, we can now easily draw the following conclusion.

²³The choices of f however could be plenty; it suffices to take V the standard \mathbb{R}^n and W any other diffeological vector space of dimension strictly smaller than n . Any linear map from V to W is then going to be smooth (see Section 3.9 in [5]).

Corollary 2.6. *Under the assumptions of Lemma 2.5, there is the inclusion $f(V_0) \leq W_0$.*

Proof. The subspace W_0 is the only subspace of dimension equal to that of W^* that has standard diffeology and splits off smoothly. Since $f(V_0) \oplus W'_0$ has all the same properties, we obtain that $f(V_0) \oplus W'_0 = W_0$. \square

The summary of necessary conditions We collect the conclusions of this section in the following statement.

Theorem 2.7. *Let V and W be two finite-dimensional diffeological vector spaces, and let $f : V \rightarrow W$ be a smooth linear map. If there exist pseudo-metrics g_V and g_W on V and W respectively that are compatible with respect to f then the following are true:*

1. $\dim(V^*) \leq \dim(W^*)$;
2. $\text{Ker}(f) \cap V_0 = \{0\}$, where V_0 is the characteristic subspace of V ;
3. The subset diffeology on $f(V_0)$ relative to its inclusion into W is standard;
4. $f(V_0)$ splits off smoothly in W .

2.2.3 Sufficient conditions

Suppose now that V and W are such that the just-mentioned necessary condition is satisfied, and let $f : V \rightarrow W$ be a smooth linear map such that $\text{Ker}(f) \cap V_0 = \{0\}$ (where V_0 is the characteristic subspace of V). By definition of a pseudo-metric, if $V = V_0 \oplus V_1$ is a smooth decomposition of V^{24} then g_V is defined by its restriction to V_0 (which is a scalar product) and is extended by zero elsewhere. The same is true of g_W and the corresponding smooth decomposition $W_0 \oplus W_1$. In this way we obtain the following.

Proposition 2.8. *Let V and W be finite-dimensional diffeological vector spaces such that $\dim(V^*) \leq \dim(W^*)$, and let $f : V \rightarrow W$ be a smooth linear map such that $\text{Ker}(f) \cap V_0 = \{0\}$ and $f(V_0) \leq W_0$. Then V and W admit pseudo-metrics compatible with respect to f .*

Proof. Let us fix smooth decompositions $V = V_0 \oplus V_1$ and $W = W_0 \oplus W_1$. To construct a pseudo-metric g_V , choose a basis v_1, \dots, v_k of V_0 and a basis v_{k+1}, \dots, v_n of V_1 ; then set

$$g_V(v_i, v_j) = \delta_{i,j} \text{ for } i, j = 1, \dots, k \text{ and } g_V(v_i, v_{k+j}) = 0 \text{ for } i = 1, \dots, n, j = 1, \dots, n - k,$$

and extend by bilinearity and symmetricity. To define g_W , then, consider $f(v_1), \dots, f(v_k) \in W_0$; notice that they are linearly independent by the assumption on $\text{Ker}(f)$. Add first $u_1, \dots, u_l \in W_0$ to obtain the basis $f(v_1), \dots, f(v_k), u_1, \dots, u_l$ of W_0 . Finally, choose a basis w_1, \dots, w_m of W_1 to obtain the basis $f(v_1), \dots, f(v_k), u_1, \dots, u_l, w_1, \dots, w_m$ of the whole W . It then suffices to define g_W to be

$$g_W(f(v_i), f(v_j)) = \delta_{i,j}, \quad g_W(u_i, u_j) = \delta_{i,j}, \quad g_W(f(v_i), u_j) = 0, \quad g_W(f(v_i), w_p) = 0, \quad g_W(u_i, w_p) = 0$$

and extend by bilinearity and symmetry. The bilinear maps g_V and g_W thus obtained are smooth, because each of the characteristic subspaces V_0 and W_0 splits off as a smooth direct summand, and by construction g_V and g_W are zero maps outside of V_0 and W_0 respectively. Finally, that they are pseudo-metrics and are compatible with each other is immediate from their definitions, whence the conclusion. \square

We are ready to establish the final criterion of the existence of compatible pseudo-metrics on a pair of diffeological vector spaces, that we state in the following form.

Theorem 2.9. *Let V and W be two finite-dimensional diffeological vector spaces, and let $f : V \rightarrow W$ be a smooth linear map. Then V and W admit compatible pseudo-metrics if and only if $\text{Ker}(f) \cap V_0 = \{0\}$ and $f(V_0) \leq W_0$.*

²⁴As we have said already, V_0 is uniquely defined by V ; this is not necessarily true of V_1 , which is defined uniquely only when the pseudo-metric has been fixed already. However, there is always at least one choice of V_1 ; what we mean at the moment is that such a choice is fixed arbitrarily.

Proof. The fact that these two conditions are necessary follows from Lemma 2.2 and Lemma 2.5, so let us show that they are sufficient. Let $V = V_0 \oplus V_1$ be a smooth decomposition, let g_V be any pseudo-metric on V , and let $W = f(V_0) \oplus W'$ be a smooth decomposition that exists by assumption. Notice that, since W' is just another instance of a finite-dimensional diffeological space, it has its own smooth decomposition of form $W'_0 \oplus W_1$, where W'_0 is standard and W_1 has trivial diffeological dual; so the whole of W smoothly decomposes as $W = (f(V_0) \oplus W'_0) \oplus W_1$. Notice that this implies that $W^* = (f(V_0) \oplus W'_0)^*$ (by the smoothness of the decomposition), in particular, they have the same dimension.

Let us define a pseudo-metric g_W , by setting it to coincide with g_V (in the obvious sense) on $f(V_0)$, choosing any scalar product for its restriction on W'_0 , while requiring W'_0 to be orthogonal to $f(V_0)$, and finally setting W_1 to be an isotropic subspace. That this is indeed a pseudo-metric follows from the considerations above, so it remains to show that g_W is indeed compatible with g_V . This essentially follows from the construction, more precisely, from the fact that $f(V_0)$ is orthogonal to any its direct complement. Indeed, if $v' = v'_0 + v'_1$ and $v'' = v''_0 + v''_1$ are any two elements of V , then

$$g_V(v', v'') = g_V(v'_0, v''_0) = g_W(f(v'_0), f(v''_0)) = g_W(f(v'_0) + f(v'_1), f(v''_0) + f(v''_1)) = g_W(f(v'), f(v'')),$$

where the third equality is by the orthogonality just mentioned. This means that g_V and g_W are compatible with f , and the proof is finished. \square

2.3 Compatibility of the dual pseudo-metrics: diffeological vector spaces

We now consider the induced pseudo-metrics on the duals of diffeological vector spaces; the main question that we aim to answer is, under what conditions the pair of pseudo-metrics dual to (induced by) two compatible ones is in turn compatible.

The induced pseudo-metric g^* on V^* : definition Recall ([9]) that, given a finite-dimensional diffeological vector space V endowed with a pseudo-metric g , the diffeological dual of V carries the induced pseudo-metric g^* (actually, a scalar product, since the diffeological dual of any finite-dimensional diffeological vector space is standard) defined by

$$g^*(v_1^*, v_2^*) := g(v_1, v_2),$$

where $v_i \in V$ is any element such that $v_i^*(\cdot) = g(v_i, \cdot)$ for $i = 1, 2$. That this is well-defined, *i.e.*, the result does not depend on the choice of v_i (as long as $g(v_i, \cdot)$ remains the same), and that v_i^* always admits such a form, was shown in [9].

The compatibility for the induced pseudo-metrics Let g_V and g_W be pseudo-metrics on V and W respectively, compatible with respect to f . Let $w_1^*, w_2^* \in W^*$; then there exist $w_1, w_2 \in W$, defined up to the cosets of the isotropic subspace of g_W , such that $w_i^*(\cdot) = g_W(w_i, \cdot)$ for $i = 1, 2$, by definition of the dual pseudo-metric

$$g_W^*(w_1^*, w_2^*) = g_W(w_1, w_2),$$

and finally, $f^*(w_i^*)(\cdot) = w_i^*(f(\cdot)) = g_W(w_i, f(\cdot))$.

The compatibility condition that we need to check is the following one:

$$g_W^*(w_1^*, w_2^*) = g_V^*(f^*(w_1^*), f^*(w_2^*)).$$

Now, in order to calculate the right-hand term in this expression, we must choose v_1 and v_2 , again defined up to their cosets with respect to the isotropic subspace of g_V , such that $g_V(v_i, v') = f^*(w_i^*)(v') = g_W(w_i, f(v'))$, for all elements of $v' \in V$ and for $w_1^*, w_2^* \in W^*$. The term on the right then becomes $g_V^*(f^*(w_1^*), f^*(w_2^*)) = g_V(v_1, v_2)$.

The dual pseudo-metrics and compatibility Let us now consider the pseudo-metrics on V^* and W^* dual to a pair of compatible pseudo-metrics on V and W . We observe right away that in general, the induced pseudo-metrics are not compatible. This follows from Lemma 2.2, as well as from the standard theory, all diffeological constructions being in fact extensions of the standard ones.

Example 2.10. Let V be the standard \mathbb{R}^n , with the canonical basis denoted by e_1, \dots, e_n , and let W be the standard \mathbb{R}^{n+k} , with the canonical basis denoted by $u_1, \dots, u_n, u_{n+1}, \dots, u_{n+k}$. Let $f : V \rightarrow W$ be the embedding of V via the identification of V with the subspace generated by u_1, \dots, u_n , given by $e_i \mapsto u_i$ for $i = 1, \dots, n$. Let g_V be any scalar product on \mathbb{R}^n ; this trivially induces a scalar product on $f(V) = \text{Span}(u_1, \dots, u_n) \leq W_0$, and let g_W be any extension of it to a scalar product on the whole W .

Let us consider the dual map on the dual the standard complement of the subspace $\text{Span}(u_1, \dots, u_n)$, that is, on the dual of $\text{Span}(u_{n+1}, \dots, u_{n+k})$. This dual is the usual dual, so it is $\text{Span}(u^{n+1}, \dots, u^{n+k})$. Let v be any element of V ; since $f(v) \in \text{Span}(u_1, \dots, u_n)$, we have

$$f^*(u^{n+i})(v) = u^{n+i}(f(v)) = 0,$$

so in the end we obtain that $\text{Ker}(f^*) = \text{Span}(u^{n+1}, \dots, u^{n+k})$.

Finally, let us consider the compatibility condition. We observe that

$$g_W^*(u^{n+i}, u^{n+i}) = g_W(u_{n+i}, u_{n+i}) > 0,$$

since g_W is a scalar product, while, of course,

$$g_V^*(f^*(u^{n+i}), f^*(u^{n+i})) = 0.$$

Quite evidently, the compatibility condition cannot be satisfied (unless $k = 0$).

Sufficient conditions for compatibility of the induced pseudo-metrics It can be inferred from the above example that the induced pseudo-metrics on the duals of standard spaces are compatible only if the spaces have the same dimension (which is not surprising, since in this case the notion of the induced pseudo-metric itself coincides with the standard one). This can be generalized to the following statement.

Theorem 2.11. Let V and W be two finite-dimensional diffeological vector spaces, and let $f : V \rightarrow W$ be a smooth linear map such that $\text{Ker}(f) \cap V_0 = \{0\}$ and $f(V_0) \leq W_0$. Let g_V and g_W be compatible pseudo-metrics on V and W respectively. Then the induced pseudo-metrics g_W^* and g_V^* are compatible with f^* if and only if $f^* : W^* \rightarrow V^*$ is a diffeomorphism.

Proof. The *only if* part of the statement, illustrated by the example above, follows from standard reasoning. Indeed, g_W^* and g_V^* are usual scalar products on standard spaces W^* and V^* respectively, and their compatibility means that f^* is a usual isometry, whose existence implies that W^* and V^* have the same dimension, and being standard spaces, this means that they are diffeomorphic as diffeological vector spaces.

Let us prove the *if* part, namely, that g_W^* and g_V^* are compatible under the assumptions of the proposition. Let e_1, \dots, e_n be a g_V -orthogonal basis of V_0 . Since g_V and g_W are compatible, $f(e_1), \dots, f(e_n)$ is a g_W -orthogonal basis of $f(V_0)$. Now, $(e_1)^*, \dots, (e_n)^*$ (recall here that v^* for $v \in V$ stands for the map $v^*(\cdot) = g_V(v, \cdot)$) form a basis of V^* (which is also orthogonal with respect to the induced pseudo-metric g_V^*), while $(f(e_1))^*, \dots, (f(e_n))^*$ are linearly independent elements of W^* . Since V^* and W^* have the same dimension, $(f(e_1))^*, \dots, (f(e_n))^*$ actually forms a basis of W^* ; and so, g_W^* is entirely determined by its values on pairs $(f(e_i))^*, (f(e_j))^*$, and moreover, we have

$$g_W^*((f(e_i))^*, (f(e_j))^*) = g_W(f(e_i), f(e_j)) = g_V(e_i, e_j) = g_V^*(e_i^*, e_j^*).$$

Finally, $f^*((f(e_i))^*)(v) = (f(e_i))^*(f(v)) = g_W(f(e_i), f(v)) = g_V(e_i, v) = e_i^*(v)$ for all $v \in V$. Therefore

$$g_V^*(e_i^*, e_j^*) = g_V^*(f^*((f(e_i))^*), f^*((f(e_j))^*)) = g_W^*((f(e_i))^*, (f(e_j))^*),$$

at which point the compatibility, with respect to f^* , of the pseudo-metrics g_W^* and g_V^* follows from $(f(e_1))^*, \dots, (f(e_n))^*$ being a basis of W^* . \square

2.4 Compatibility of the dual pseudo-metrics: diffeological vector pseudo-bundles

Let us now consider the following question: if two given pseudo-metrics g_1 and g_2 are compatible with respect to the gluing along a given pair of maps (f, f) , when is it true that g_2^* and g_1^* are compatible with the gluing defined by (f^{-1}, f^*) ? (Obviously, we assume here that f is invertible).

The compatibility condition for g_2^* and g_1^* Let $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$ be locally trivial finite-dimensional diffeological vector pseudo-bundles, let (f, f) be a gluing of the former to the latter such that f is a diffeomorphism with its image, and let g_1 and g_2 be pseudo-metrics on V_1 and V_2 respectively, compatible with respect to the given gluing. The latter induces a well-defined gluing, along the maps \tilde{f}^* and f^{-1} , of the dual pseudo-bundle $\pi_2^* : V_2^* \rightarrow X_2$ to the pseudo-bundle $\pi_1^* : V_1^* \rightarrow X_1$, the result of which is the pseudo-bundle $\pi_2^* \cup_{(\tilde{f}^*, f^{-1})} \pi_1^* : V_2^* \cup_{\tilde{f}^*} V_1^* \rightarrow X_2 \cup_{f^{-1}} X_1$, while g_2 and g_1 induce pseudo-metrics $g_2^* : X_2 \rightarrow (V_2^*)^* \otimes (V_2^*)^*$ and $g_1^* : X_1 \rightarrow (V_1^*)^* \otimes (V_1^*)^*$ on the dual pseudo-bundles. They satisfy the usual compatibility condition if

$$g_1^*(f^{-1}(y'))(\tilde{f}^*(v^*), \tilde{f}^*(w^*)) = g_2^*(y')(v^*, w^*)$$

for all $y' \in Y' = f(Y)$ and for all $v^*, w^* \in (\pi_2^{-1}(y'))^*$.

The necessary condition The compatibility between g_2^* and g_1^* implies in particular that for all $y \in Y$ the pseudo-metrics $g_2^*(f(y))$ and $g_1^*(y)$ are compatible with the smooth linear map $\tilde{f}^*|_{\pi_1^{-1}(y)}$ between diffeological vector spaces $(\pi_2^{-1}(f(y)))^*$ and $(\pi_1^{-1}(y))^*$. Thus, Theorem 2.11 implies that \tilde{f}^* is a diffeomorphism on each fibre.

Proposition 2.12. *Let $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$ be diffeological vector pseudo-bundles, locally trivial and with finite-dimensional fibres, and let (\tilde{f}, f) be an invertible gluing between them. Suppose that g_1 and g_2 are two pseudo-metrics on these pseudo-bundles compatible with the gluing along (\tilde{f}, f) . If the induced pseudo-metrics g_2^* and g_1^* are compatible with the gluing along (\tilde{f}^*, f^{-1}) then the restriction of \tilde{f}^* on each fibre in its domain of definition is a diffeomorphism.*

Criterion of compatibility The statement that follows shows that, for the two induced pseudo-metrics to be compatible with the induced gluing, the map dual to the gluing map \tilde{f} must satisfy a rather stringent condition (although an expected one).

Theorem 2.13. *Let $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$ be two diffeological vector pseudo-bundles, locally trivial and with finite-dimensional fibres, let (\tilde{f}, f) be a gluing between them, and let g_1 and g_2 be compatible pseudo-metrics on V_1 and V_2 respectively. Then the induced pseudo-metrics g_2^* and g_1^* on the corresponding dual pseudo-bundles are compatible if and only if \tilde{f}^* is a pseudo-bundle diffeomorphism of its domain with its image.*

Notice that diffeological vector spaces may have diffeomorphic duals without being diffeomorphic themselves, and the same is true for diffeological vector pseudo-bundles.

Proof. By assumption, g_1 and g_2 are compatible with the gluing given by the pair (\tilde{f}, f) , that is

$$g_2(f(y))(\tilde{f}(v), \tilde{f}(w)) = g_1(y)(v, w) \text{ for all } y \in Y \text{ and for all } v, w \in \pi_1^{-1}(y).$$

Suppose first that \tilde{f}^* is a diffeomorphism with its image. Then, first of all, by the definition of g_2^* we have

$$g_2^*(y')(v^*, w^*) = g_2(y')(v, w)$$

for all $y' \in f(Y)$, for all $v^*, w^* \in (\pi_2^{-1}(y'))^*$, and for $v, w \in (\pi_2^{-1}(y'))_0$. Notice that v and w are uniquely defined by the latter condition; and they are such that $v^*(\cdot) = g_2(y')(v, \cdot)$ and $w^*(\cdot) = g_2(y')(w, \cdot)$. Notice also (we will need this immediately below) that this means

$$\tilde{f}^*(v^*)(\cdot) = v^*(\tilde{f}(\cdot)) = g_2(y')(v, \tilde{f}(\cdot)) = g_2(y')(\tilde{f}(v_1), \tilde{f}(\cdot)) = g_1(f^{-1}(y'))(v_1, \cdot),$$

where $v_1 \in (\pi_1^{-1}(f^{-1}(y')))_0$ is such that $v = \tilde{f}(v_1)$; such an element exists and is uniquely defined because \tilde{f}^* being a diffeomorphism is equivalent to \tilde{f} being a diffeomorphism between each pair of subspaces $(\pi_1^{-1}(f^{-1}(y')))_0$ and $(\pi_2^{-1}(y'))_0$. Similarly, we have

$$\tilde{f}^*(w^*)(\cdot) = g_2(y')(\tilde{f}(w_1), \tilde{f}(\cdot)) = g_1(f^{-1}(y'))(w_1, \cdot)$$

for $w_1 \in (\pi_1^{-1}(f^{-1}(y')))_{0}$ such that $w = \tilde{f}(w_1)$. It remains now to consider the left-hand part of the compatibility condition. We have:

$$g_1^*(f^{-1}(y'))(\tilde{f}^*(v^*), \tilde{f}^*(w)) = g_1(f^{-1}(y'))(v_1, w_1) = g_2(y')(\tilde{f}(v_1), \tilde{f}(w_1)) = g_2(y')(v, w) = g_2^*(y')(v^*, w^*),$$

as wanted.

Let us prove the *vice versa* of the statement, that is, let us assume that g_2^* and g_1^* are compatible, and let us show that \tilde{f}^* is a diffeomorphism. We notice first of all that it follows from the considerations made for individual vector spaces that \tilde{f}^* is bijective and, as is the case for any dual map, it is smooth. Finally, the smoothness of its inverse follows from the fact that V_1^* and V_2^* are locally trivial and have standard fibres. \square

3 Compatibility of pseudo-metrics and the gluing-dual commutativity conditions

In this section we consider which correlations there might be between the notion of compatible pseudo-metrics on two given pseudo-bundles (with a specified gluing), and the gluing-dual commutativity conditions. We start by taking our two usual pseudo-bundles, $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$, and a gluing along (\tilde{f}, f) between them. Assuming that these pseudo-bundles admit pseudo-metrics g_1 and g_2 compatible with the gluing, we consider the following questions: 1) what are the implications of the existence of compatible pseudo-metrics for the pseudo-bundles themselves? 2) if the gluing-dual commutativity condition holds for V_1, V_2 , and (\tilde{f}, f) , does it necessarily hold for V_2^*, V_1^* , and (f^*, f^{-1}) ? 3) if g_2^* and g_1^* exist, under what conditions are they compatible with (\tilde{f}^*, f^{-1}) , in particular, is their compatibility equivalent to the gluing-dual commutativity? 4) does taking the dual pseudo-metric commute (in the notation to be introduced, this will be the equality $\tilde{g}^* = \tilde{g}^*$) with the gluing of pseudo-metrics (as defined in [11])? We consider these questions in order, after quickly introducing a preliminary notion.

3.1 The characteristic sub-bundle of a finite-dimensional vector pseudo-bundle

Let $\pi : V \rightarrow X$ be a finite-dimensional diffeological vector pseudo-bundle, and let $(\pi^{-1}(x))_0$ be the characteristic subspace of the fibre $\pi^{-1}(x)$. Denote by V_0 the sub-bundle of V defined as

$$V_0 := \cup_{x \in X} (\pi^{-1}(x))_0.$$

We say that V_0 is the **characteristic sub-bundle** of the pseudo-bundle V . It is evident from the construction that the characteristic sub-bundle of a locally trivial pseudo-bundle is itself locally trivial. Furthermore, every pseudo-metric on V is uniquely defined by its restriction to V_0 . Finally, the *vice versa* of the latter statement is also true, if we assume V to be locally trivial.

Lemma 3.1. *Let $\pi : V \rightarrow X$ be a locally trivial diffeological vector pseudo-bundle, let $\pi_0 : V_0 \rightarrow X$ be its characteristic sub-bundle, and let g_0 be a pseudo-metric on V_0 . Then there exists one, and only one, pseudo-metric g on V whose restriction on V_0 coincides with g_0 .*

Proof. It suffices to define $g(x)(v', v'') = \begin{cases} g_0(x)(v', v'') & \text{if } v', v'' \in \pi_0^{-1}(x) \\ 0 & \text{otherwise;} \end{cases}$ the conclusion then follows from the definitions of a pseudo-metric and that of the characteristic sub-bundle. \square

Proposition 3.2. *Let $\pi : V \rightarrow X$ be a locally trivial finite-dimensional diffeological vector pseudo-bundle that admits a pseudo-metric g . Then its characteristic sub-bundle $\pi_0 : V_0 \rightarrow X$ is diffeomorphic to its dual pseudo-bundle $\pi^* : V^* \rightarrow X$ via the natural pairing map associated to g .*

Proof. Let $\psi_g : V_0 \rightarrow V^*$ be the natural pairing associated to g , that is,

$$\psi_g(v)(\cdot) = g(\pi(v))(v, \cdot).$$

That this is a bijection follows from its fibrewise nature and it being a bijection on each individual fibre (see [9]); furthermore (see the same source), as a map on the characteristic subspace it is a diffeomorphism

with the dual fibre. By the assumption of local triviality this implies that ψ_g , as well as its inverse, are smooth across the fibres as well, so they are smooth as a whole, whence the conclusion.²⁵ \square

3.2 Implications of compatibility of g_1 and g_2 for V_1 and V_2

This extends the criterion for diffeological vector spaces (Theorem 2.9). The pseudo-bundle version is an immediate consequence and is as follows.

Proposition 3.3. *Let $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$ be two diffeological vector pseudo-bundles, locally trivial and with finite-dimensional fibres, let (\tilde{f}, f) be a gluing between them, and let g_1 and g_2 be compatible pseudo-metrics. Then \tilde{f} determines, over the domain of gluing, a smooth embedding of the characteristic sub-bundle of V_1 into the characteristic sub-bundle of V_2 .*

Proof. This follows directly from the already-mentioned Theorem 2.9, applied to the restriction of \tilde{f} on each fibre in its domain of definition; the theorem affirms that such restriction is an embedding of each characteristic subspace, so the fibre of the characteristic sub-bundle, of V_1 into that of V_2 . We should only add that the restriction of \tilde{f} onto the intersection of its domain of definition with the characteristic sub-bundle of V_1 is smooth across the fibres, because \tilde{f} is so. \square

3.3 The gluing-dual commutativity condition and gluing along a diffeomorphism

We now recall a statement (which essentially appears in [10], Lemma 5.17) that (together with some results from the previous sections) will allow us to deduce the gluing-dual commutativity in a number of cases. The statement basically is that if the gluing of two pseudo-bundles is performed along a diffeomorphism, then the gluing-dual commutativity condition always holds. We also add the explicit construction of the commutativity diffeomorphism (which was not specified in the above source).

Theorem 3.4. *Let $\chi_1 : W_1 \rightarrow X_1$ and $\chi_2 : W_2 \rightarrow X_2$ be two diffeological vector pseudo-bundles, let $h : X_1 \supseteq Y \rightarrow X_2$ be a smooth invertible map with smooth inverse, and let \tilde{h} be its smooth fibrewise linear lift that is a diffeomorphism of its domain with its image. Then the map*

$$\Psi_{\cup,*} : (W_1 \cup_{\tilde{h}} W_2)^* \rightarrow W_2^* \cup_{\tilde{h}^*} W_1^*$$

defined by

$$\Psi_{\cup,*} = \begin{cases} j_2^{W_1^*} \circ (j_1^{W_1})^* & \text{on } ((\chi_1 \cup_{(\tilde{h},h)} \chi_2)^*)^{-1} (i_1^{X_1} (X_1 \setminus Y)) \\ j_2^{W_1^*} \circ \tilde{h}^* \circ (j_2^{W_2})^* & \text{on } ((\chi_1 \cup_{(\tilde{h},h)} \chi_2)^*)^{-1} (i_2^{X_2} (h(Y))) \\ j_1^{W_2^*} \circ (j_2^{W_2})^* & \text{on } ((\chi_1 \cup_{(\tilde{h},h)} \chi_2)^*)^{-1} (i_2^{X_2} (X_2 \setminus h(Y))) \end{cases}$$

is a pseudo-bundle diffeomorphism covering the switch map $\varphi_{X_1 \leftrightarrow X_2}$.

Proof. It is easy to see that $\Psi_{\cup,*}$ is a bijection, so let us show that it is smooth (the proof that its inverse is smooth is then analogous). Let us first consider the general shape of an arbitrary plot q^* of $(W_1 \cup_{\tilde{h}} W_2)^*$, and that of an arbitrary plot s^* of $W_2^* \cup_{\tilde{h}^*} W_1^*$. Let $q^* : U \rightarrow (W_1 \cup_{\tilde{h}} W_2)^*$ be any plot; we can however assume that U is connected, so $(\chi_1 \cup_{(\tilde{h},h)} \chi_2) \circ q$, which is a plot of $X_1 \cup_h X_2$, lifts to a plot of X_1 or to a plot of X_2 . This means that q^* acts only on fibres of $W_1 \cup_{\tilde{h}} W_2$ that pullback to W_1 or to W_2 , respectively. In the former case we have that there exists a plot q_1^* of W_1^* such that

$$q^* = \begin{cases} ((j_1^{W_1})^*)^{-1} \circ q_1^* & \text{over } i_1^{X_1} (X_1 \setminus Y) \\ ((j_2^{W_2})^*)^{-1} \circ (\tilde{h}^*)^{-1} \circ q_1^* & \text{over } i_2^{X_2} (h(Y)); \end{cases}$$

in the latter case there exists a plot q_2^* of W_2^* such that

$$q^* = ((j_2^{W_2})^*)^{-1} \circ q_2^*.$$

²⁵It is easy prove that ψ_g is smooth even if we do not assume V to be locally trivial.

For analogous reasons, if $s^* : U' \rightarrow W_2^* \cup_{\tilde{h}^*} W_1^*$ defined on a connected domain U' , then either there exists a plot s_2^* of W_2^* such that

$$s^* = \begin{cases} j_1^{W_2^*} \circ s_2^* & \text{over } i_1^{X_2}(X_2 \setminus h(Y)) \\ j_2^{W_1^*} \circ \tilde{h}^* \circ s_2^* & \text{over } i_2^{X_1}(h(Y)), \end{cases}$$

or else there exists a plot s_1^* of W_1^* such that

$$s^* = j_2^{W_2^*} \circ s_1^*.$$

Let us consider $\Psi_{\cup,*} \circ q^*$ for q^* of the first and the second type. Assume that q^* is of the first type. Then by direct calculation we obtain

$$\Psi_{\cup,*} \circ q^* = \begin{cases} j_2^{W_1^*} \circ q_1^* & \text{over } i_2^{X_1}(X_1 \setminus Y), \\ j_2^{W_1^*} \circ \tilde{h}^* \circ (j_2^{W_2^*})^* \circ ((j_2^{W_2^*})^*)^{-1} \circ (\tilde{h}^*)^{-1} \circ q_1^* & \text{over } i_2^{X_1}(Y), \end{cases}$$

that is, $\Psi_{\cup,*} \circ q^* = j_2^{W_1^*} \circ q_1^*$ over the whole of $i_2^{X_1}(X_1)$, which corresponds to a plot s^* of the second type, for $s_1^* := q_1^*$.

Similarly, if q^* has its second possible form, we obtain

$$\Psi_{\cup,*} \circ q^* = \begin{cases} j_2^{W_1^*} \circ \tilde{h}^* \circ q_2^* & \text{over } i_2^{X_1}(Y) \\ j_1^{W_2^*} \circ q_2^* & \text{over } i_1^{X_2}(X_2 \setminus h(Y)), \end{cases}$$

that is, the first possible form of a plot s^* , with $s_2^* := q_2^*$.

Finally, the smoothness of $(\Psi_{\cup,*})^{-1}$, whose formula

$$(\Psi_{\cup,*})^{-1} = \begin{cases} ((j_1^{W_1^*})^*)^{-1} \circ (j_2^{W_1^*})^{-1} & \text{on } (\chi_2^* \cup_{(\tilde{h},h)} \chi_1^*)^{-1}(i_2^{X_1}(X_1 \setminus Y)) \\ ((j_2^{W_2^*})^*)^{-1} \circ (\tilde{h}^*)^{-1} \circ (j_2^{W_1^*})^{-1} & \text{on } (\chi_2^* \cup_{(\tilde{h},h)} \chi_1^*)^{-1}(i_2^{X_1}(Y)) \\ ((j_2^{W_2^*})^*)^{-1} \circ (j_1^{W_2^*})^{-1} & \text{on } (\chi_2^* \cup_{(\tilde{h},h)} \chi_1^*)^{-1}(i_1^{X_2}(X_2 \setminus h(Y))) \end{cases}$$

is given by the inverses of the three maps that $\Psi_{\cup,*}$ itself, is established in a completely analogous fashion. \square

3.4 Gluing-dual commutativity conditions for (\tilde{f}, f) and (\tilde{f}^*, f^{-1})

We now consider the gluing-dual commutativity condition for V_2^* , V_1^* , and (\tilde{f}^*, f^{-1}) , under the assumption that such condition holds for V_1 , V_2 , and (\tilde{f}, f) . For the duals, this condition takes form of the existence of a diffeomorphism

$$\Phi_{\cup,*}^{(*)} : (V_2^* \cup_{\tilde{f}^*} V_1^*)^* \rightarrow V_1^{**} \cup_{\tilde{f}^{**}} V_2^{**}$$

covering the inverse of the switch map $\varphi_{X_1 \leftrightarrow X_2}$ and satisfying the following:

$$\begin{cases} \Phi_{\cup,*}^{(*)} \circ ((j_2^{V_2^*})^*)^{-1} = j_1^{V_1^{**}} & \text{on } (\pi_1^{**} \cup_{(\tilde{f}^{**},f)} \pi_2^{**})^{-1}(i_1^{X_1}(X_1 \setminus Y)), \\ \Phi_{\cup,*}^{(*)} \circ ((j_1^{V_1^*})^*)^{-1} = j_2^{V_2^{**}} \circ \tilde{f}^* & \text{on } (\pi_1^{**} \cup_{(\tilde{f}^{**},f)} \pi_2^{**})^{-1}(i_2^{X_2}(f(Y))), \\ \Phi_{\cup,*}^{(*)} \circ ((j_2^{V_1^*})^*)^{-1} = j_2^{V_2^{**}} & \text{on } (\pi_1^{**} \cup_{(\tilde{f}^{**},f)} \pi_2^{**})^{-1}(i_2^{X_2}(X_2 \setminus f(Y))). \end{cases}$$

Notice that this formula can serve as a definition of a certain map $\Phi_{\cup,*}^{(*)}$ between the domain and the range indicated; what we really need to do is to show that it is a diffeomorphism.

Theorem 3.5. *Let $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$ be two locally trivial finite-dimensional diffeological vector pseudo-bundles, let (\tilde{f}, f) be a pair of smooth maps that defines a gluing of the former pseudo-bundle to the latter, and let $\Phi_{\cup,*} : (V_1 \cup_{\tilde{f}} V_2)^* \rightarrow V_2^* \cup_{\tilde{f}^*} V_1^*$ be the canonical gluing-dual commutativity diffeomorphism. Let g_1 and g_2 be pseudo-metrics on V_1 and V_2 respectively, compatible with respect to the gluing. Then there exists a diffeomorphism*

$$\Phi_{\cup,*}^{(*)} : (V_2^* \cup_{\tilde{f}^*} V_1^*)^* \rightarrow V_1^{**} \cup_{\tilde{f}^{**}} V_2^{**}$$

covering the map $(\varphi_{X_1 \leftrightarrow X_2})^{-1} : X_2 \cup_{f^{-1}} X_1 \rightarrow X_1 \cup_f X_2$.

The claim of the theorem could be restated by saying that the dual pseudo-bundles V_2^* and V_1^* satisfy the gluing-dual commutativity condition for the gluing pair (\tilde{f}^*, f^{-1}) .

Proof. Recall that the gluing-dual commutativity condition for the initial pseudo-bundles, that is, for V_1 and V_2 , implies that \tilde{f}^* is a diffeomorphism of its domain with its image. It is then a direct consequence of Theorem 3.4 that the following map

$$\Phi_{\cup,*}^{(*)} = \begin{cases} j_2^{V_2^{**}} \circ (j_1^{V_2^*})^* & \text{on } \left((\pi_2^* \cup_{(\tilde{f}^*, f^{-1})} \pi_1^*)^* \right)^{-1} (i_1^{X_2}(X_2 \setminus f(Y))), \\ j_2^{V_2^{**}} \circ \tilde{f}^* \circ (j_2^{V_1^*})^* & \text{on } \left((\pi_2^* \cup_{(\tilde{f}^*, f^{-1})} \pi_1^*)^* \right)^{-1} (i_2^{X_1}(Y)), \\ j_1^{V_1^{**}} \circ (j_2^{V_1^*})^* & \text{on } \left((\pi_2^* \cup_{(\tilde{f}^*, f^{-1})} \pi_1^*)^* \right)^{-1} (i_2^{X_1}(X_1 \setminus Y)) \end{cases}$$

is the desired gluing-dual commutativity diffeomorphism. \square

4 Compatibility of g_2^* and g_1^* implies the gluing-dual commutativity condition for V_1 and V_2

So far we have spoken of the gluing-dual commutativity condition as a prerequisite to obtaining a canonical construction of the induced pseudo-metric on the pseudo-bundle obtained by gluing. In principle, it is not a necessary condition (a pseudo-metric on $V_1 \cup_{\tilde{f}} V_2$ can be constructed directly out of compatible pseudo-metrics on V_1 and V_2 , using the flexibility of diffeology in piecing together smooth maps); however, if we want at the same time to consider the dual pseudo-bundles V_2^* and V_1^* , and to ensure that the induced pseudo-metrics on them are again compatible, the reasoning involved starts to come rather close to the gluing-dual commutativity. Indeed, in this section we show that there is essentially an equivalence between the compatibility of g_2^* with g_1^* , and the existence of a gluing-dual commutativity diffeomorphism $\Phi_{\cup,*}$.

4.1 From $\Phi_{\cup,*}$ to the compatibility of g_2^* , g_1^* , and (\tilde{f}^*, f^{-1})

It is not difficult to show that if we assume the gluing-dual commutativity, and more precisely, the existence of the specific diffeomorphism $\Phi_{\cup,*} : (V_1 \cup_{\tilde{f}} V_2)^* \rightarrow V_2^* \cup_{\tilde{f}^*} V_1^*$ given by

$$\Phi_{\cup,*} = \begin{cases} j_2^{V_1^*} \circ (j_1^{V_1})^* & \text{on } ((\pi_1 \cup_{(\tilde{f}, f)} \pi_2)^*)^{-1} (i_1^{X_1}(X_1 \setminus Y)) \\ j_2^{V_1^*} \circ \tilde{f}^* \circ (j_2^{V_2})^* & \text{on } ((\pi_1 \cup_{(\tilde{f}, f)} \pi_2)^*)^{-1} (i_2^{X_2}(f(Y))) \\ j_1^{V_2^*} \circ (j_2^{V_2})^* & \text{on } ((\pi_1 \cup_{(\tilde{f}, f)} \pi_2)^*)^{-1} (i_2^{X_2}(X_2 \setminus f(Y))) \end{cases}$$

then the dual pseudo-metrics, if they exist, are compatible.

Theorem 4.1. *Let $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$ be two finite-dimensional locally trivial diffeological vector pseudo-bundles, let (\tilde{f}, f) be a pair of smooth maps defining a gluing of V_1 to V_2 , and let g_1 and g_2 be pseudo-metrics on V_1 and V_2 respectively, compatible with this gluing; assume that V_1 , V_2 , and (\tilde{f}, f) satisfy the gluing-dual commutativity condition. Then g_2^* and g_1^* are compatible with the gluing of V_2^* to V_1^* along the pair (\tilde{f}^*, f^{-1}) .*

Proof. By Theorem 2.13 it suffices to show that \tilde{f}^* is a diffeomorphism of its domain with its image, and this is a trivial consequence of the form in which we stated the gluing-dual commutativity condition, namely, as the smoothness of the specific diffeomorphism $\Phi_{\cup,*}$. Indeed, denoting for brevity $Z_2^* := ((\pi_1 \cup_{(\tilde{f}, f)} \pi_2)^*)^{-1} (i_2^{X_2}(f(Y)))$, we immediately obtain

$$\tilde{f}^* = (j_2^{V_1^*})^{-1} \circ \Phi_{\cup,*}|_{Z_2^*} \circ ((j_2^{V_2})^*)^{-1};$$

thus, \tilde{f}^* is a composition of three diffeomorphisms, so a diffeomorphism itself. \square

4.2 The *vice versa*: the smoothness of $\Phi_{U,*}$ out of compatibility of g_2^* and g_1^*

The reverse implication, that is, obtaining a smooth $\Phi_{U,*}$ assuming the compatibility of g_2^* and g_1^* , is now easily obtained by applying Theorem 3.4, and the criteria for compatibility of the dual pseudo-metrics. Do note that the application of Theorem 3.4 is not straightforward; indeed, it speaks of gluing along a diffeomorphism, and the assumptions formulated in terms of various pseudo-metrics do not provide for \tilde{f} being one. Therefore we need some preliminary considerations.

4.2.1 Characteristic sub-bundles and the respective dual pseudo-bundles

In order to obtain our desired conclusion, namely, that the compatibility of pseudo-metrics dual to a given pair of compatible pseudo-metrics provides for the gluing-dual commutativity, we need several preliminary statements. We collect them in this section; they are more or less of independent interest.

The pseudo-metric \tilde{g} on the pseudo-bundle $W_1 \cup_{\tilde{h}} W_2$ Assuming that W_1 and W_2 admit compatible pseudo-metrics g_1 and g_2 allows (without any further assumptions on \tilde{h}) for a direct construction of a pseudo-metric \tilde{g} on $W_1 \cup_{\tilde{h}} W_2$, which fibrewise coincides with either g_1 or g_2 , as appropriate. This pseudo-metric is defined by the following formula:

$$\tilde{g}(x) = \begin{cases} g_1((i_1^{X_1})^{-1}(x)) \circ ((j_1^{W_1})^{-1} \otimes (j_1^{W_1})^{-1}) & \text{for } x \in i_1^{X_1}(X_1 \setminus Y) \\ g_2((i_2^{X_2})^{-1}(x)) \circ ((j_2^{W_2})^{-1} \otimes (j_2^{W_2})^{-1}) & \text{for } x \in i_2^{X_2}(X_2). \end{cases}$$

The switch map for characteristic sub-bundles As we have already commented, a pseudo-metric on a pseudo-bundle is essentially determined by its behavior on the characteristic sub-bundle; and furthermore, assuming the local triviality and the existence of a pseudo-metric, the characteristic sub-bundle is diffeomorphic to the dual pseudo-bundle. Thus, we can expect significant correlations between these three notions; and in this paragraph we specify some of them, as needed to reach the final aim of this section.

Lemma 4.2. *Let $\chi_1 : W_1 \rightarrow X_1$ and $\chi_2 : W_2 \rightarrow X_2$ be two locally trivial diffeological vector pseudo-bundles, let W_1^0 and W_2^0 be their characteristic sub-bundles, let $h : X_1 \supseteq Y \rightarrow X_2$ be a smooth invertible map with smooth inverse, and let \tilde{h} be its smooth fibrewise linear lift such that its restriction \tilde{h}_0 on $\text{Domain}(\tilde{h}) \cap W_1^0$ is a diffeomorphism. Then there is a canonical diffeomorphism*

$$\varphi_{W_1^0 \leftrightarrow W_2^0} : W_1^0 \cup_{\tilde{h}_0} W_2^0 \rightarrow W_2^0 \cup_{\tilde{h}_0^{-1}} W_1^0$$

covering the switch map $\varphi_{X_1 \leftrightarrow X_2}$.

Proof. The diffeomorphism in question is in fact the same concept as the switch map (which we implied is a diffeomorphism, but did not prove that). Indeed, we denote the map obtained by analogy with $\varphi_{X_1 \leftrightarrow X_2}$, by $\varphi_{W_1^0 \leftrightarrow W_2^0}$ and define it to be

$$\varphi_{W_1^0 \leftrightarrow W_2^0} = \begin{cases} j_2^{W_1^0} \circ (j_1^{W_1^0})^{-1} & \text{on } (\chi_1^0 \cup_{(\tilde{h}_0, h)} \chi_2^0)^{-1}(i_1^{X_1}(X_1 \setminus Y)) \\ j_2^{W_1^0} \circ (\tilde{h}_0)^{-1} \circ (j_2^{W_2^0})^{-1} & \text{on } (\chi_1^0 \cup_{(\tilde{h}_0, h)} \chi_2^0)^{-1}(i_2^{X_2}(h(Y))) \\ j_1^{W_2^0} \circ (j_2^{W_2^0})^{-1} & \text{on } (\chi_1^0 \cup_{(\tilde{h}_0, h)} \chi_2^0)^{-1}(i_2^{X_2}(X_2 \setminus f(Y))), \end{cases}$$

where by χ_i^0 we denote the restriction of χ_i to W_i^0 . Notice that it is its own inverse; let us show that it is smooth.

Let $p : U \rightarrow W_1^0 \cup_{\tilde{h}_0} W_2^0$; it suffices to consider the case when U is connected. Under such assumption, p lifts to either a plot p_1 of W_1^0 or to a plot p_2 of W_2^0 , therefore p itself either has form

$$p = \begin{cases} j_1^{W_1^0} \circ p_1 & \text{on } p_1^{-1}(W_1^0 \setminus \text{Domain}(\tilde{h}_0)) \\ j_2^{W_2^0} \circ \tilde{h}_0 \circ p_1 & \text{on } p_1^{-1}(\text{Domain}(\tilde{h}_0)) \end{cases}$$

in the former case, or it has form $p = j_2^{W_2^0} \circ p_2$ in the latter case. Accordingly, by direct calculation we obtain that $\varphi_{W_1^0 \leftrightarrow W_2^0} \circ p = j_2^{W_2^0} \circ p_1$ for p that lifts to p_1 and

$$\varphi_{W_1^0 \leftrightarrow W_2^0} \circ p = \begin{cases} j_2^{W_1^0} \circ (\tilde{h}_0)^{-1} \circ p_2 & \text{on } p_2^{-1}(\text{Range}(\tilde{h}_0)) \\ j_1^{W_2^0} \circ p_2 & \text{on } p_2^{-1}(W_2^0 \setminus \text{Range}(\tilde{h}_0)) \end{cases}$$

for p that lifts for p_2 . Clearly, in both cases the result is a plot of $W_2^0 \cup_{(\tilde{h}_0)^{-1}} W_1^0$, whence the conclusion. \square

The lemma just proven is a preliminary statement which will be needed to establish a link between the following two statements; all three put together will allow us to relate the gluing-dual commutativity to compatibility of (dual) pseudo-metrics.

The triple $W_1^0 \cup_{\tilde{h}_0} W_2^0 \cong (W_1 \cup_{\tilde{h}} W_2)^0 \cong (W_1 \cup_{\tilde{h}} W_2)^*$ The facts that we prove here ensure a kind of total commutativity between $()^0$ (characteristic) and $()^*$ (dual); this phrase is of course very informal, what we really mean shall be clear from the two statements that follow.

Proposition 4.3. *Let $\chi_1 : W_1 \rightarrow X_1$ and $\chi_2 : W_2 \rightarrow X_2$ be two locally trivial diffeological vector pseudo-bundles, let W_1^0 and W_2^0 be their characteristic sub-bundles, let $h : X_1 \supseteq Y \rightarrow X_2$ be a smooth invertible map with smooth inverse, and let \tilde{h} be its smooth fibrewise linear lift. Let g_1 and g_2 be pseudo-metrics on W_1 and W_2 respectively, compatible with the gluing along (\tilde{h}, h) , and assume that the dual pseudo-metrics g_2^* and g_1^* are compatible with (\tilde{h}^*, h^{-1}) . Let \tilde{h}_0 be the restriction of \tilde{h} on $\text{Domain}(\tilde{h}) \cap W_1^0$. Then:*

1. \tilde{h}_0 is a diffeomorphism with values in W_2^0 ;
2. There is a pseudo-bundle diffeomorphism

$$\Phi^0 : W_1^0 \cup_{\tilde{h}_0} W_2^0 \rightarrow (W_1 \cup_{\tilde{h}} W_2)^0$$

covering the identity map $X_1 \cup_f X_2 \rightarrow X_1 \cup_f X_2$;

3. The natural pairing map $\Psi_g^0 : (W_1 \cup_{\tilde{h}} W_2)^0 \rightarrow (W_1 \cup_{\tilde{h}} W_2)^*$ associated to the pseudo-metric \tilde{g} on $W_1 \cup_{\tilde{h}} W_2$ is a diffeomorphism of the characteristic sub-bundle $(W_1 \cup_{\tilde{h}} W_2)^0$ with the dual pseudo-bundle $(W_1 \cup_{\tilde{h}} W_2)^*$.

Proof. 1. The compatibility of g_1 and g_2 means that their restrictions $g_1(y)$ and $g_2(h(y))$ on each fibre in the domain of definition and, respectively, the range of \tilde{h} are compatible pseudo-metrics on the diffeological vector spaces $\chi_1^{-1}(y)$ and $\chi_2^{-1}(h(y))$. By Theorem 2.9 this means that the restriction \tilde{h}_0 of \tilde{h} to $\text{Domain}(\tilde{h}) \cap W_1^0$ is a smooth injection, whose range is contained in $\text{Range}(\tilde{h}) \cap W_2^0$. Furthermore, by compatibility of g_2^* and g_1^* and Theorem 2.11, the dual spaces $(\chi_2^{-1}(h(y)))^*$ and $(\chi_1^{-1}(y))^*$ have the same dimension, which is equal to the dimension of the corresponding characteristic subspaces; therefore \tilde{h}_0 is also surjective. Finally, that its inverse is smooth, follows from local triviality and the fact that its restriction onto each fibre is a smooth linear map between finite-dimensional vector spaces.

2. This is a direct consequence of the definition of a characteristic sub-bundle. The diffeomorphism $\Phi^0 : W_1^0 \cup_{\tilde{h}_0} W_2^0 \rightarrow (W_1 \cup_{\tilde{h}} W_2)^0$ is defined by

$$\Phi^0 = \begin{cases} j_1^{W_1} \circ (j_1^{W_1^0})^{-1} & \text{over } i_1^{X_1}(X_1 \setminus Y) \\ j_2^{W_2} \circ (j_2^{W_2^0})^{-1} & \text{over } i_2^{X_2}(X_2); \end{cases}$$

it is essentially the natural inclusion map.

3. The diffeomorphism in question is the pairing map

$$\Psi_g^0 : (W_1 \cup_{\tilde{h}} W_2)^0 \rightarrow (W_1 \cup_{\tilde{h}} W_2)^*$$

restricted to the characteristic sub-bundle and defined in the usual way, *i.e.*, by

$$\text{if } w \in (W_1 \cup_{\tilde{h}} W_2)^0 \Rightarrow \Psi_g^0(w)(\cdot) = \tilde{g}((\chi_1 \cup_{(\tilde{h}, h)} \chi_2)(w))(w, \cdot).$$

That it is smooth, follows from smoothness of \tilde{g} ; that it is bijective, is easily obtained by examining its restriction on each fibre, where, since the fibres of characteristic sub-bundle have standard diffeology, it becomes the usual isomorphism-by-duality on some standard \mathbb{R}^n . Finally, the smoothness of its inverse follows from the assumption of local triviality. \square

Under the same assumptions as those of Proposition 4.3, we also have the following:

Proposition 4.4. *There is a canonical diffeomorphism*

$$\Psi_{g_2}^0 \cup_{(\tilde{h}_0^{-1}, \tilde{h}^*)} \Psi_{g_1}^0 : W_2^0 \cup_{\tilde{h}_0^{-1}} W_1^0 \rightarrow W_2^* \cup_{\tilde{h}^*} W_1^*,$$

whose restrictions onto the factors of gluing coincide with the natural pairing maps associated to a pair of compatible pseudo-metrics g_2 and g_1 .

Proof. It follows from the assumptions, specifically, the assumption of compatibility of g_2^* and g_1^* , that \tilde{h}^* is a diffeomorphism; by Proposition 4.3 the \tilde{h}_0^{-1} is also a diffeomorphism. Thus, the desired diffeomorphism of the two pseudo-bundles in the statement is the result $\Psi_{g_2}^0 \cup_{(\tilde{h}_0^{-1}, \tilde{h}^*)} \Psi_{g_1}^0$ of a gluing (see Section 1.3.1 for definitions) of the natural pairing maps $\Psi_{g_2}^0 : W_2^0 \rightarrow W_2^*$ and $\Psi_{g_1}^0 : W_1^0 \rightarrow W_1^*$. Since the gluing of two diffeomorphisms along a pair of diffeomorphisms yields again a diffeomorphism, we only need to check that $\Psi_{g_2}^0$ and $\Psi_{g_1}^0$ are $(\tilde{h}_0^{-1}, \tilde{h}^0)$ -compatible, that is, that for any $w_2 \in \text{Domain}(\tilde{h}_0^{-1})$ we have

$$\tilde{h}^*(\Psi_{g_2}^0(w_2)) = \Psi_{g_1}^0(\tilde{h}_0^{-1}(w_2)).$$

Let us verify why this is true.

The left-hand side of the expression is by definition

$$\tilde{h}^*(\Psi_{g_2}^0(w_2))(\cdot) = \tilde{h}^*(g_2(\chi_2(w_2))(w_2, \cdot)) = g_2(\chi_2(w_2))(w_2, \tilde{h}(\cdot)) = g_1(h^{-1}(\chi_2(w_2)))(\tilde{h}_0^{-1}(w_2), \cdot),$$

where the last equality follows from the compatibility of pseudo-metrics g_1 and g_2 . The right-hand side of the expression is

$$\Psi_{g_1}^0(\tilde{h}_0^{-1}(w_2))(\cdot) = g_1(\chi_1(\tilde{h}_0^{-1}(w_2)))(\tilde{h}_0^{-1}(w_2), \cdot),$$

and it remains to observe that $h^{-1}(\chi_2(w_2)) = \chi_1(\tilde{h}_0^{-1}(w_2))$, simply because \tilde{h}_0 is a lift of h , and the statement is proven. \square

The map $\Psi_{g_2}^0 \cup_{(\tilde{h}_0^{-1}, \tilde{h}^*)} \Psi_{g_1}^0$ We conclude this section by giving the precise formula for the map $\Psi_{g_2}^0 \cup_{(\tilde{h}_0^{-1}, \tilde{h}^*)} \Psi_{g_1}^0$, which we will need in the next section. As follows from the general construction of gluing of two smooth maps, we have

$$\left(\Psi_{g_2}^0 \cup_{(\tilde{h}_0^{-1}, \tilde{h}^*)} \Psi_{g_1}^0 \right) (w^0) = \begin{cases} j_1^{W_2^*} \circ \Psi_{g_2}^0 \circ (j_1^{W_2^0})^{-1} & \text{on } (\chi_2^0 \cup_{(\tilde{h}_0^{-1}, h^{-1})} \chi_1^0)^{-1}(i_1^{X_2}(X_2 \setminus h(Y))) \\ j_2^{W_1^*} \circ \Psi_{g_1}^0 \circ (j_2^{W_1^0})^{-1} & \text{on } (\chi_2^0 \cup_{(\tilde{h}_0^{-1}, h^{-1})} \chi_1^0)^{-1}(i_2^{X_1}(X_1)). \end{cases}$$

4.2.2 Proving the gluing-dual commutativity

We now give our final statement, which is a sufficient condition (and, together with Theorem 4.1, a criterion) for the gluing-dual commutativity condition to be satisfied.

Theorem 4.5. *Let $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$ be diffeological vector pseudo-bundles, locally trivial and with finite-dimensional fibres, and let (\tilde{f}, f) be a gluing of V_1 to V_2 , with an invertible f . Suppose that V_1 and V_2 admit pseudo-metrics compatible with this gluing, and let g_1 and g_2 be a fixed choice of such pseudo-metrics. Assume, finally, that the induced pseudo-metrics g_2^* and g_1^* on the dual pseudo-bundles V_2^* and V_1^* are compatible with the gluing along (\tilde{f}^*, f^{-1}) . Then V_1, V_2 , and (\tilde{f}, f) satisfy the gluing-dual commutativity condition.*

Proof. Let us first show that $(V_1 \cup_{\tilde{f}} V_2)^*$ and $V_2^* \cup_{\tilde{f}^*} V_1^*$ are diffeomorphic, and then explain why there is a canonical diffeomorphism between them. Applying Proposition 4.3(3) and then (2), we obtain

$$(V_1 \cup_{\tilde{f}} V_2)^* \cong (V_1 \cup_{\tilde{f}} V_2)^0 \cong V_1^0 \cup_{\tilde{f}_0} V_2^0;$$

next, by Lemma 4.2 and Proposition 4.4 we obtain

$$V_1^0 \cup_{\tilde{f}_0} V_2^0 \cong V_2^0 \cup_{\tilde{f}_0^{-1}} V_1^0 \cong V_2^* \cup_{\tilde{f}^*} V_1^*,$$

as wanted. It remains to see that the diffeomorphisms involved produce in the end the canonical commutativity diffeomorphism $\Phi_{\cup, *}$.

Let us now specify the final diffeomorphism that we obtain from the above sequence. Let \tilde{g} be the pseudo-metric on $V_1 \cup_{\tilde{f}} V_2$ induced by g_1 and g_2 . Then the first diffeomorphism of the sequence is $(\Psi_{\tilde{g}}^0)^{-1}$; the second is $(\Phi^0)^{-1}$, the inverse of the diffeomorphism described in the proof of Proposition 4.3, the third is the switch-like map $\varphi_{V_1^0 \leftrightarrow V_2^0}$, and the fourth is $\Psi_{g_2}^0 \cup_{(\tilde{f}_0^{-1}, \tilde{f}^*)} \Psi_{g_1}^0$. We need to see that the composition

$$\Phi = \left(\Psi_{g_2}^0 \cup_{(\tilde{f}_0^{-1}, \tilde{f}^*)} \Psi_{g_1}^0 \right) \circ \left(\varphi_{V_1^0 \leftrightarrow V_2^0} \right) \circ (\Phi^0)^{-1} \circ (\Psi_{\tilde{g}}^0)^{-1}$$

coincides with the appropriate canonically defined map $\Phi_{\cup, *}$.

Indeed, after some obvious cancelations the pointwise description of the diffeomorphism Φ is as follows:

$$\Phi(v) = \begin{cases} \left(j_2^{V_1^*} \circ \Psi_{g_1}^0 \circ (j_1^{V_1})^{-1} \circ (\Psi_{\tilde{g}}^0)^{-1} \right) (v) & \text{if } (\pi_1 \cup_{(\tilde{f}, f)} \pi_2)^*(v) \in i_1^{X_1}(X_1 \setminus Y) \\ \left(j_2^{V_1^*} \circ \Psi_{g_1}^0 \circ \tilde{f}_0^{-1} \circ (j_2^{V_2})^{-1} \circ (\Psi_{\tilde{g}}^0)^{-1} \right) (v) & \text{if } (\pi_1 \cup_{(\tilde{f}, f)} \pi_2)^*(v) \in i_2^{X_2}(f(Y)) \\ \left(j_1^{V_2^*} \circ \Psi_{g_2}^0 \circ (j_2^{V_2})^{-1} \circ (\Psi_{\tilde{g}}^0)^{-1} \right) (v) & \text{if } (\pi_1 \cup_{(\tilde{f}, f)} \pi_2)^*(v) \in i_2^{X_2}(X_2 \setminus f(Y)). \end{cases}$$

Let us consider the three cases indicated.

Let first v be such that $(\pi_1 \cup_{(\tilde{f}, f)} \pi_2)^*(v) \in i_1^{X_1}(X_1 \setminus Y)$; then we have

$$\Phi(v) = \left(j_2^{V_1^*} \circ \Psi_{g_1}^0 \circ (j_1^{V_1})^{-1} \right) \left((\Psi_{\tilde{g}}^0)^{-1}(v) \right), \text{ where } (\Psi_{\tilde{g}}^0)^{-1}(v) = v^0 \in j_1^{V_1}(\pi_1^{-1}(X_1 \setminus Y)) \cap (V_1 \cup_{\tilde{f}} V_2)^0$$

$$\text{such that } v(\cdot) = g_1(x)((j_1^{V_1})^{-1}(v^0), (j_1^{V_1})^{-1}(\cdot)) \text{ for } x = \pi_1((j_1^{V_1})^{-1}(v^0)),$$

where (\cdot) stands for the argument of $v \in (V_1 \cup_{\tilde{f}} V_2)^*$, that is being taken in $j_1^{V_1}(V_1 \setminus \pi_1^{-1}(Y))$. Hence we obtain

$$\Phi(v)(\cdot) = j_2^{V_1^*} \left(g_1(x)((j_1^{V_1})^{-1}(v^0), \cdot) \right).$$

This time (\cdot) stands for an element of V_1^0 ; it is related to the argument of $v(\cdot)$ by the map $j_1^{V_1^0}$, so we have in the end

$$\Phi(v)(\cdot) = j_2^{V_1^*} \left(v(j_1^{V_1^0}(\cdot)) \right) \Rightarrow \Phi(v) = \left(j_2^{V_1^*} \circ (j_1^{V_1})^* \right) (v) \Rightarrow \Phi = j_2^{V_1^*} \circ (j_1^{V_1})^*,$$

i.e., the canonical form of the gluing-dual commutativity diffeomorphism.

The other two cases are rather similar. Let $v \in (V_1 \cup_{\tilde{f}} V_2)^*$ be such that $(\pi_1 \cup_{(\tilde{f}, f)} \pi_2)^*(v) \in i_2^{X_2}(f(Y))$; we then have

$$\Phi(v) = \left(j_2^{V_1^*} \circ \Psi_{g_1}^0 \circ \tilde{f}_0^{-1} \circ (j_2^{V_2})^{-1} \right) \left((\Psi_{\tilde{g}}^0)^{-1}(v) \right), \text{ where } (\Psi_{\tilde{g}}^0)^{-1}(v) = v^0 \in j_2^{V_2}(\pi_2^{-1}(f(Y))) \cap (V_1 \cup_{\tilde{f}} V_2)^0$$

$$\text{such that } v(\cdot) = g_2(x)((j_2^{V_2})^{-1}(v^0), (j_2^{V_2})^{-1}(\cdot)) \text{ for } x = \pi_2((j_2^{V_2})^{-1}(v^0)).$$

From this, we obtain

$$\Phi(v) = \left(j_2^{V_1^*} \circ \Psi_{g_1}^0 \circ \tilde{f}_0^{-1} \circ (j_2^{V_2})^{-1} \right) (v^0)(\cdot) = j_2^{V_1^*} \left(g_2(f^{-1}(x))((\tilde{f}_0^{-1} \circ (j_2^{V_2})^{-1})(v^0), \cdot) \right).$$

Once again, we should relate this to the expression for $v(\cdot)$, this time keeping in mind the compatibility of the pseudo-metrics g_1 and g_2 . It suffices to recall that the argument (\cdot) in this case is being taken in

V_1^0 and is related to that of $v(\cdot)$ by the map $j_2^{V_2} \circ \tilde{f}_0$. Therefore we can rewrite the expression for $\Phi(v)(\cdot)$ as follows:

$$\Phi(v)(\cdot) = j_2^{V_1^*} \left(g_1(f^{-1}(x))((\tilde{f}_0^{-1} \circ (j_2^{V_2})^{-1})(v^0), \cdot) \right) = (j_2^{V_1^*} \circ \tilde{f}_0^*) \left(g_2(x)((j_2^{V_2})^{-1}(v^0), \tilde{f}_0(\cdot)) \right),$$

which then allows us to conclude that

$$\Phi(v)(\cdot) = \left(j_2^{V_1^*} \circ \tilde{f}_0^* \circ (j_2^{V_2})^* \right) (v)(\cdot) \Rightarrow \Phi = j_2^{V_1^*} \circ \tilde{f}_0^* \circ (j_2^{V_2})^*.$$

It remains to consider the third part of the definition of Φ . Let $v \in (V_1 \cup_{\tilde{f}} V_2)^*$ be such that $(\pi_1 \cup_{(\tilde{f}, f)} \pi_2)^*(v) \in i_2^{X_2}(X_2 \setminus f(Y))$; we then have

$$\Phi(v) = \left(j_1^{V_2^*} \circ \Psi_{g_2}^0 \circ (j_2^{V_2})^{-1} \right) \left((\Psi_g^0)^{-1}(v) \right), \text{ where } (\Psi_g^0)^{-1}(v) = v^0 \in j_2^{V_2}(\pi_2^{-1}(X_2 \setminus f(Y))) \cap (V_1 \cup_{\tilde{f}} V_2)^0$$

such that $v(\cdot) = g_2(x)((j_2^{V_2})^{-1}(v^0), (j_2^{V_2})^{-1}(\cdot))$ for $x = \pi_2((j_2^{V_2})^{-1}(v^0))$.

We therefore have

$$\Phi(v)(\cdot) = \left(j_1^{V_2^*} \circ \Psi_{g_2}^0 \circ (j_2^{V_2})^{-1} \right) (v^0)(\cdot) = j_1^{V_2^*} \left(g_2(x)((j_2^{V_2})^{-1}(v^0), \cdot) \right).$$

By the same considerations regarding the argument (\cdot) , which in this case is related to that of $v(\cdot)$ by the map $j_2^{V_2}$, we obtain

$$\Phi(v)(\cdot) = j_1^{V_2^*} \left(v(j_2^{V_2}(\cdot)) \right) \Rightarrow \Phi = j_1^{V_2^*} \circ (j_2^{V_2})^*,$$

therefore Φ has the canonical form also in the third case, whence the final claim. \square

4.3 Final remarks on the gluing-commutativity condition

To conclude the discussion on the gluing-dual commutativity, we stress that the crucial point²⁶ throughout was the dual map \tilde{f}^* being a diffeomorphism (of its domain with its image). As follows from our proofs, it is this condition that is equivalent to both

- the *specific* map $\Phi_{\cup, *}$ being a diffeomorphism; and
- the dual pseudo-metrics g_2^* and g_1^* being compatible.

It remains to observe that this also justifies our choice to state right away the gluing-dual commutativity condition in terms of $\Phi_{\cup, *}$ being smooth, rather than just asking for the existence of *some* diffeomorphism between $(V_1 \cup_{\tilde{f}} V_2)^*$ and $V_2^* \cup_{\tilde{f}^*} V_1^*$: the two are equivalent.

5 The pseudo-metrics \tilde{g}^* and \tilde{g}^* on pseudo-bundles $(V_1 \cup_{\tilde{f}} V_2)^*$ and $V_2^* \cup_{\tilde{f}^*} V_1^*$

Assuming that V_1 , V_2 , and (\tilde{f}, f) satisfy the gluing-dual commutativity condition implies, in particular, that there are two ways to construct a pseudo-metric on the pseudo-bundle $(V_1 \cup_{\tilde{f}} V_2)^* \cong V_2^* \cup_{\tilde{f}^*} V_1^*$, which correspond, respectively, to the left-hand side and the right-hand side of this expression. Specifically, the (specific expression for the) pseudo-bundle on the left carries the pseudo-metric \tilde{g} that is induced by, or dual to, the pseudo-metric \tilde{g} . The pseudo-bundle on the right is obtained from a given gluing of two pseudo-bundles carrying compatible pseudo-metrics; it therefore carries a pseudo-metric \tilde{g}^* that corresponds to this gluing (we mentioned this in Section 1; the details can be found in [11], and we recall what we need immediately below). We show that this is actually the same pseudo-metric.

²⁶Under the assumption that the pseudo-bundles involved are finite-dimensional and locally trivial.

5.1 The pseudo-metric \tilde{g}^* on $(V_1 \cup_{\tilde{f}} V_2)^*$

This is the pseudo-metric dual to²⁷ the pseudo-metric \tilde{g} on the pseudo-bundle $V_1 \cup_{\tilde{f}} V_2$; the latter is defined as the composition

$$\tilde{g} = (\Phi_{\cup, *}^{-1} \otimes \Phi_{\cup, *}^{-1}) \circ \Phi_{\otimes, \cup} \circ (g_2 \cup_{(f^{-1}, \tilde{f}^* \otimes \tilde{f}^*)} g_1) \circ \varphi_{X_1 \leftrightarrow X_2},$$

where $\varphi_{X_1 \leftrightarrow X_2}$ is the switch map, and

$$\Phi_{\cup, *} : (V_1 \cup_{\tilde{f}} V_2)^* \rightarrow V_2^* \cup_{\tilde{f}^*} V_1^* \quad \text{and} \quad \Phi_{\otimes, \cup} : (V_2^* \otimes V_2^*) \cup_{\tilde{f}^* \otimes \tilde{f}^*} (V_1^* \otimes V_1^*) \rightarrow (V_2^* \cup_{\tilde{f}^*} V_1^*) \otimes (V_2^* \cup_{\tilde{f}^*} V_1^*)$$

are the appropriate versions of the commutativity diffeomorphisms (see Section 4.2.1 for the explicit formula).

The pseudo-metric \tilde{g}^* is then defined as the pseudo-metric dual to \tilde{g} , which by the usual definition means that, if $\Psi_{\tilde{g}} : V_1 \cup_{\tilde{f}} V_2 \rightarrow (V_1 \cup_{\tilde{f}} V_2)^*$ is the (already-seen) pairing map relative to \tilde{g} , that is, $\Psi_{\tilde{g}}(v) = \tilde{g}((\pi_1 \cup_{(\tilde{f}, f)} \pi_2)(v))(v, \cdot)$, then for any $x \in X_1 \cup_f X_2$ and any $v^*, w^* \in ((\pi_1 \cup_{(\tilde{f}, f)} \pi_2)^*)^{-1}(x)$ we have by definition

$$\tilde{g}^*(x)(v^*, w^*) := \tilde{g}(x)(v, w),$$

where $v, w \in (\pi_1 \cup_{(\tilde{f}, f)} \pi_2)^{-1}(x)$ are any two elements such that $\Psi_{\tilde{g}}(v) = v^*$ and $\Psi_{\tilde{g}}(w) = w^*$. We can avail ourselves of the already-mentioned restriction $\Psi_{\tilde{g}}^0$ of $\Psi_{\tilde{g}}$ to the characteristic sub-bundle $(V_1 \cup_{\tilde{f}} V_2)^0$ and thus define

$$\tilde{g}^*(x)(v^*, w^*) := \tilde{g}(x) \left((\Psi_{\tilde{g}}^0)^{-1}(v^*), (\Psi_{\tilde{g}}^0)^{-1}(w^*) \right).$$

Finally, an even more explicit formula for \tilde{g}^* is

$$\tilde{g}^*(x)(v^*, w^*) = \begin{cases} g_1((i_1^{X_1})^{-1}(x))(v_1, w_1), & \text{if } x \in \text{Range}(i_1^{X_1}), \\ g_2((i_2^{X_2})^{-1}(x))(v_2, w_2), & \text{if } x \in \text{Range}(i_2^{X_2}), \end{cases}$$

where we have by definition

$$\begin{cases} v_1 = \left((j_1^{V_1})^{-1} \circ (\Psi_{\tilde{g}}^0)^{-1} \right) (v^*), & w_1 = \left((j_1^{V_1})^{-1} \circ (\Psi_{\tilde{g}}^0)^{-1} \right) (w^*) \\ v_2 = \left((j_2^{V_2})^{-1} \circ (\Psi_{\tilde{g}}^0)^{-1} \right) (v^*), & w_2 = \left((j_2^{V_2})^{-1} \circ (\Psi_{\tilde{g}}^0)^{-1} \right) (w^*). \end{cases}$$

5.2 The pseudo-metric \tilde{g}^* on $V_2^* \cup_{\tilde{f}^*} V_1^*$

The pseudo-metric \tilde{g}^* is defined on the pseudo-bundle $V_2^* \cup_{\tilde{f}^*} V_1^*$ fibrewise, by imposing it to coincide with g_2^* or g_1^* on the appropriate subsets. Specifically, for $i = 1, 2$ let, as before, $\Psi_{g_i} : V_i \rightarrow V_i^*$ be the pairing map associated to g_1 and g_2 respectively; then for any given $x \in X_2 \cup_{f^{-1}} X_1$ and any two $v^*, w^* \in (\pi_2^* \cup_{(\tilde{f}^*, f^{-1})} \pi_1^*)^{-1}(x)$ we define

$$\tilde{g}^*(x)(v^*, w^*) = \begin{cases} g_2^*((i_1^{X_2})^{-1}(x))((j_1^{V_2^*})^{-1}(v^*), (j_1^{V_2^*})^{-1}(w^*)) = g_2((i_1^{X_2})^{-1}(x))(v_2, w_2) & \text{if } x \in \text{Range}(i_1^{X_2}), \\ g_1^*((i_2^{X_1})^{-1}(x))((j_2^{V_1^*})^{-1}(v^*), (j_2^{V_1^*})^{-1}(w^*)) = g_1((i_2^{X_1})^{-1}(x))(v_1, w_1) & \text{if } x \in \text{Range}(i_2^{X_1}), \end{cases}$$

where again we make reference to the characteristic sub-bundles and the corresponding invertible restrictions $\Psi_{g_2}^0$ of Ψ_{g_2} and $\Psi_{g_1}^0$ of Ψ_{g_1} , since by construction

$$\begin{cases} v_2 = \left((\Psi_{g_2}^0)^{-1} \circ (j_1^{V_2^*})^{-1} \right) (v^*), & w_2 = \left((\Psi_{g_2}^0)^{-1} \circ (j_1^{V_2^*})^{-1} \right) (w^*), \\ v_1 = \left((\Psi_{g_1}^0)^{-1} \circ (j_2^{V_1^*})^{-1} \right) (v^*), & w_1 = \left((\Psi_{g_1}^0)^{-1} \circ (j_2^{V_1^*})^{-1} \right) (w^*). \end{cases}$$

²⁷It would be more precise to say, induced by duality.

5.3 Comparing \tilde{g}^* and \tilde{g}^*

In the case we are considering, we have assumed²⁸ that V_1 , V_2 , and (\tilde{f}, f) satisfy the gluing-dual commutativity condition. By the very definition of the latter, this means that the pseudo-bundles $(V_1 \cup_{\tilde{f}} V_2)^*$ and $V_2^* \cup_{\tilde{f}^*} V_1^*$ are diffeomorphic, and in a canonical way. Since both of these pseudo-bundles also carry the canonical pseudo-metrics, described in the two sections above, it is natural to ask next whether their canonical identification, via the gluing-dual commutativity diffeomorphism $\Phi_{U,*}$, is an isometry relative to these pseudo-metrics. In this section we prove that it is.

More precisely, since \tilde{g}^* and \tilde{g}^* are maps

$$\tilde{g}^* : X_1 \cup_f X_2 \rightarrow (V_1 \cup_{\tilde{f}} V_2)^{**} \otimes (V_1 \cup_{\tilde{f}} V_2)^{**} \quad \text{and} \quad \tilde{g}^* : X_2 \cup_{f^{-1}} X_1 \rightarrow (V_2^* \cup_{\tilde{f}^*} V_1^*)^* \otimes (V_2^* \cup_{\tilde{f}^*} V_1^*)^*,$$

we show that by adding the diffeomorphism

$$(\Phi_{U,*}^{-1})^* \otimes (\Phi_{U,*}^{-1})^* : (V_1 \cup_{\tilde{f}} V_2)^{**} \otimes (V_1 \cup_{\tilde{f}} V_2)^{**} \rightarrow (V_2^* \cup_{\tilde{f}^*} V_1^*)^* \otimes (V_2^* \cup_{\tilde{f}^*} V_1^*)^*$$

and the switch map $\varphi_{X_1 \leftrightarrow X_2} : X_1 \cup_f X_2 \rightarrow X_2 \cup_{f^{-1}} X_1$, we obtain

$$(\Phi_{U,*}^{-1})^* \otimes (\Phi_{U,*}^{-1})^* \circ \tilde{g}^* = \tilde{g}^* \circ (\varphi_{X_1 \leftrightarrow X_2}).$$

The full statement is as follows.

Theorem 5.1. *Let $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$ be two finite-dimensional locally trivial diffeological vector pseudo-bundle, and let (\tilde{f}, f) be a gluing between them such that f and \tilde{f}^* are diffeomorphisms. Assume that there exist pseudo-metrics g_1 and g_2 on V_1 and V_2 respectively, that are compatible with the gluing along (\tilde{f}, f) . Then the following is true:*

$$((\Phi_{U,*}^{-1})^* \otimes (\Phi_{U,*}^{-1})^*) \circ \tilde{g}^* = \tilde{g}^* \circ (\varphi_{X_1 \leftrightarrow X_2}).$$

Notice that the assumptions of the theorem provide for the existence of all the maps that appear in the claim, that is, for the existence of the smooth switch map $\varphi_{X_1 \leftrightarrow X_2}$, that of the gluing-dual commutativity diffeomorphism $\Phi_{U,*}$, and the existence and compatibility of the dual pseudo-metrics g_2^* and g_1^* , through which the pseudo-metrics \tilde{g}^* and \tilde{g}^* are defined.

Proof. The actual meaning of the formula that we wish to prove is as follows: taken an arbitrary $x \in X_1 \cup_f X_2$ and arbitrary $v^*, w^* \in (\pi_2^* \cup_{(\tilde{f}^*, f^{-1})} \pi_1^*)^{-1}(\varphi_{X_1 \leftrightarrow X_2}(x)) \in V_2^* \cup_{\tilde{f}^*} V_1^*$, we should have

$$\tilde{g}^*(\varphi_{X_1 \leftrightarrow X_2}(x))(v^*, w^*) = \tilde{g}^*(x)(\Phi_{U,*}^{-1}(v^*), \Phi_{U,*}^{-1}(w^*)).$$

Since this formula involves the switch map and a gluing-dual commutativity diffeomorphism, both of which are defined separately in three cases, we should check the desired equality in the same three cases as well. These cases are: $x \in i_1^{X_1}(X_1 \setminus Y)$, $x \in i_2^{X_2}(f(Y))$, and $x \in i_2^{X_2}(X_2 \setminus f(Y))$.

Let $x \in i_1^{X_1}(X_1 \setminus Y)$; then $v^*, w^* \in \text{Range}(j_2^{V_1^*})$ and $\varphi_{X_1 \leftrightarrow X_2}(x) = i_2^{X_1}((i_1^{X_1})^{-1}(x))$. Notice that the corresponding v and w that appear in the definition of the pseudo-metric \tilde{g}^* are then

$$v = \left((\Psi_{g_1}^0)^{-1} \circ (j_2^{V_1^*})^{-1} \right) (v^*), \quad w = \left((\Psi_{g_1}^0)^{-1} \circ (j_2^{V_1^*})^{-1} \right) (w^*),$$

therefore we have

$$\begin{aligned} \tilde{g}^*(\varphi_{X_1 \leftrightarrow X_2}(x))(v^*, w^*) &= \tilde{g}^*(i_2^{X_1}((i_1^{X_1})^{-1}(x)))(v^*, w^*) = \\ &= g_1((i_1^{X_1})^{-1}(x)) \left(((\Psi_{g_1}^0)^{-1} \circ (j_2^{V_1^*})^{-1})(v^*), ((\Psi_{g_1}^0)^{-1} \circ (j_2^{V_1^*})^{-1})(w^*) \right). \end{aligned}$$

²⁸As is, in fact, necessary for the two pseudo-metrics just described to be well-defined.

On the other hand, when \tilde{g}^* is applied to two elements v_1^*, w_1^* in the fibre of $(V_1 \cup_{\tilde{f}} V_2)^*$ over a point in $i_1^{X_1}(X_1 \setminus Y)$, its value is $g_1((i_1^{X_1})^{-1}(x))(v_1, w_1)$, where

$$v_1 = \left((j_1^{V_1})^{-1} \circ (\Psi_{\tilde{g}}^0)^{-1} \right) (v_1^*), \quad w_1 = \left((j_1^{V_1})^{-1} \circ (\Psi_{\tilde{g}}^0)^{-1} \right) (w_1^*);$$

in our case we will have $v_1^* := \Phi_{\cup, *}^{-1}(v^*)$ and $w_1^* := \Phi_{\cup, *}^{-1}(w^*)$. Observe now that

$$\Phi_{\cup, *}^{-1}(v^*) = \left(((j_1^{V_1})^*)^{-1} \circ (j_2^{V_1^*})^{-1} \right) (v^*) \quad \text{and} \quad \Phi_{\cup, *}^{-1}(w^*) = \left(((j_1^{V_1})^*)^{-1} \circ (j_2^{V_1^*})^{-1} \right) (w^*),$$

and that in the case we are considering the relation between $(\Psi_{\tilde{g}}^0)^{-1}$ and $(\Psi_{g_1}^0)^{-1}$ is as follows:

$$(\Psi_{\tilde{g}}^0)^{-1} = j_1^{V_1} \circ (\Psi_{g_1}^0)^{-1} \circ (j_1^{V_1})^*.$$

Therefore we have

$$\begin{aligned} v_1 &= \left((j_1^{V_1})^{-1} \circ (\Psi_{\tilde{g}}^0)^{-1} \right) (\Phi_{\cup, *}^{-1}(v^*)) = \left((\Psi_{g_1}^0)^{-1} \circ (j_2^{V_1^*})^{-1} \right) (v^*) \\ w_1 &= \left((j_1^{V_1})^{-1} \circ (\Psi_{\tilde{g}}^0)^{-1} \right) (\Phi_{\cup, *}^{-1}(w^*)) = \left((\Psi_{g_1}^0)^{-1} \circ (j_2^{V_1^*})^{-1} \right) (w^*), \end{aligned}$$

hence

$$\begin{aligned} \tilde{g}^*(x)(\Phi_{\cup, *}^{-1}(v^*), \Phi_{\cup, *}^{-1}(w^*)) &= \\ &= g_1 \left((i_1^{X_1})^{-1}(x) \right) \left(((\Psi_{g_1}^0)^{-1} \circ (j_2^{V_1^*})^{-1})(v^*), ((\Psi_{g_1}^0)^{-1} \circ (j_2^{V_1^*})^{-1})(w^*) \right) = \\ &= \tilde{g}^*(\varphi_{X_1 \leftrightarrow X_2}(x))(v^*, w^*), \end{aligned}$$

as wanted.

Turning to the second case, let $x \in i_2^{X_2}(f(Y))$, so that $v^*, w^* \in \text{Range}(j_2^{V_1^*})$ and $\varphi_{X_1 \leftrightarrow X_2}(x) = (i_2^{X_1} \circ f^{-1} \circ (i_2^{X_2})^{-1})(x)$. To calculate $\tilde{g}^*(\varphi_{X_1 \leftrightarrow X_2}(x))(v^*, w^*)$, write

$$v = \left((\Psi_{g_1}^0)^{-1} \circ (j_2^{V_1^*})^{-1} \right) (v^*), \quad w = \left((\Psi_{g_1}^0)^{-1} \circ (j_2^{V_1^*})^{-1} \right) (w^*);$$

this implies that

$$\begin{aligned} \tilde{g}^*(\varphi_{X_1 \leftrightarrow X_2}(x))(v^*, w^*) &= \\ &= g_1 \left((f^{-1} \circ (i_2^{X_2})^{-1})(x) \right) \left(((\Psi_{g_1}^0)^{-1} \circ (j_2^{V_1^*})^{-1})(v^*), ((\Psi_{g_1}^0)^{-1} \circ (j_2^{V_1^*})^{-1})(w^*) \right) = \\ &= g_2((i_2^{X_2})^{-1})(x) \left((\tilde{f} \circ (\Psi_{g_1}^0)^{-1} \circ (j_2^{V_1^*})^{-1})(v^*), (\tilde{f} \circ (\Psi_{g_1}^0)^{-1} \circ (j_2^{V_1^*})^{-1})(w^*) \right), \end{aligned}$$

where we have used the compatibility of the pseudo-metrics g_1 and g_2 and the fact that v and w belong to the characteristic sub-bundle, on which \tilde{f} is invertible by assumption.

To calculate now the second part of the identity to verify, recall that by definition

$$\tilde{g}^*(x)(v_2^*, w_2^*) = g_2((i_2^{X_2})^{-1}(x))(v_2, w_2),$$

where

$$v_2 := \left((j_2^{V_2})^{-1} \circ (\Psi_{\tilde{g}}^0)^{-1} \right) (v_2^*) \quad \text{and} \quad w_2 := \left((j_2^{V_2})^{-1} \circ (\Psi_{\tilde{g}}^0)^{-1} \right) (w_2^*),$$

with v_2^*, w_2^* denoting for brevity

$$v_2^* := \Phi_{\cup, *}^{-1}(v^*), \quad w_2^* := \Phi_{\cup, *}^{-1}(w^*).$$

We now recall that in the case we are considering,

$$\Phi_{\cup, *}^{-1}(v^*) = \left(((j_1^{V_1})^*)^{-1} \circ \tilde{f}^* \circ (j_2^{V_2^*})^{-1} \right) (v^*) \quad \text{and} \quad \Phi_{\cup, *}^{-1}(w^*) = \left(((j_1^{V_1})^*)^{-1} \circ \tilde{f}^* \circ (j_2^{V_2^*})^{-1} \right) (w^*),$$

and there are the following relations between Ψ_g^0 , $\Psi_{g_1}^0$, and $\Psi_{g_2}^0$ (which we state immediately for their inverses):

$$(\Psi_g^0)^{-1} = j_2^{V_2} \circ (\Psi_{g_2}^0)^{-1} \circ (\tilde{f}^*)^{-1} \circ (j_1^{V_1})^* \quad \text{and} \quad (\Psi_{g_2}^0)^{-1} = \tilde{f} \circ (\Psi_{g_1}^0)^{-1}.$$

Thus, by direct calculation

$$v_2 = \left((\Psi_{g_2}^0)^{-1} \circ (j_2^{V_2^*})^{-1} \right) (v^*) \quad \text{and} \quad w_2 = \left((\Psi_{g_2}^0)^{-1} \circ (j_2^{V_2^*})^{-1} \right) (w^*),$$

which means that

$$\tilde{g}^*(x)(v_2^*, w_2^*) = \tilde{g}^*(\varphi_{X_1 \leftrightarrow X_2}(x))(v^*, w^*),$$

again as wanted.

Finally, let us consider the third case, that of $x \in i_2^{X_2}(X_2 \setminus f(Y))$; then $v^*, w^* \in \text{Range}(j_1^{V_2^*})$ and $\varphi_{X_1 \leftrightarrow X_2}(x) = \left(i_1^{X_2} \circ (i_2^{X_2})^{-1} \right) (x)$. Furthermore,

$$\tilde{g}^*(\varphi_{X_1 \leftrightarrow X_2}(x))(v^*, w^*) = g_2((i_2^{X_2})^{-1}(x))(v, w),$$

where

$$v = \left((\Psi_{g_2}^0)^{-1} \circ (j_1^{V_2^*})^{-1} \right) (v^*) \quad \text{and} \quad w = \left((\Psi_{g_2}^0)^{-1} \circ (j_1^{V_2^*})^{-1} \right) (w^*).$$

On the other side, we have

$$\tilde{g}^*(x)(v_2^*, w_2^*) = g_2((i_2^{X_2})^{-1}(x))(v_2, w_2),$$

where

$$v_2 = \left((j_2^{V_2})^{-1} \circ (\Psi_g^0)^{-1} \right) (v_2^*) \quad \text{and} \quad w_2 = \left((j_2^{V_2})^{-1} \circ (\Psi_g^0)^{-1} \right) (w_2^*),$$

and in turn

$$v_2^* = (\Phi_{\cup, *})^{-1}(v^*) = \left(((j_2^{V_2})^*)^{-1} \circ (j_1^{V_2^*})^{-1} \right) (v^*), \quad w_2^* = (\Phi_{\cup, *})^{-1}(w^*) = \left(((j_2^{V_2})^*)^{-1} \circ (j_1^{V_2^*})^{-1} \right) (w^*).$$

Finally, we observe that there is the following relation between Ψ_g^0 and $\Psi_{g_2}^0$ (stated again for their inverses):

$$(\Psi_g^0)^{-1} = j_2^{V_2} \circ (\Psi_{g_2}^0)^{-1} \circ (j_2^{V_2})^*.$$

Thus, putting together consecutively the expressions for v_2 , v_2^* , and $(\Psi_g^0)^{-1}$ (and likewise, for w_2 , w_2^* , and $(\Psi_g^0)^{-1}$), we obtain

$$v_2 = \left((\Psi_{g_2}^0)^{-1} \circ (j_1^{V_2^*})^{-1} \right) (v^*) = v \quad \text{and} \quad w_2 = \left((\Psi_{g_2}^0)^{-1} \circ (j_1^{V_2^*})^{-1} \right) (w^*) = w,$$

which implies that

$$\tilde{g}^*(x)(v_2^*, w_2^*) = \tilde{g}^*(\varphi_{X_1 \leftrightarrow X_2}(x))(v^*, w^*),$$

and therefore concludes our consideration of the third case. All cases having thus been exhausted, the proof is finished. \square

We can re-phrase our main conclusion by stating the following.

Corollary 5.2. *The gluing-dual commutativity diffeomorphism $\Phi_{\cup, *}$ is a pseudo-bundle isometry between $\left((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^* \right)$ and $\left(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^* \right)$.*

6 The covariant Clifford algebras

We choose this umbrella term to refer to the pseudo-bundles of Clifford algebras that are associated to whatever pseudo-bundles we obtain by interposing the operations of gluing and taking the dual pseudo-bundle. These are, first of all, the pseudo-bundles of Clifford algebras associated to $(V_1 \cup_{\tilde{f}} V_2)^*$ and $V_2^* \cup_{\tilde{f}^*} V_1^*$; as we have seen above, these pseudo-bundles are *a priori* different, but they are naturally identified under appropriate assumptions. Once these assumptions are imposed, we still have another *a priori* difference, that of the two natural pseudo-metrics \tilde{g}^* and \tilde{g}^* on them being different, the possibility treated in the preceding section. Finally, there is a third option, that of the result of gluing of two pseudo-bundles of Clifford algebras, those associated to V_2^* and V_1^* . In this section we complete our consideration of these three cases, and of how they are interrelated.

6.1 The diffeomorphism $\mathcal{C}l((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^*) \cong \mathcal{C}l(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$

The diffeomorphism in question is easily obtained from the gluing-dual commutativity diffeomorphism $\Phi_{\cup,*}$, whose existence we assume and which in this case both guarantees that the pseudo-metrics \tilde{g}^* and \tilde{g}^* exist and are well-defined, and, by Corollary 4.1, is an isometry with respect to them. Extending $\Phi_{\cup,*}$ to a diffeomorphism between $\mathcal{C}l((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^*)$ and $\mathcal{C}l(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$ is then a standard procedure, whose result we denote by

$$\Phi_{\cup,*}^{\mathcal{C}l} : \mathcal{C}l((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^*) \rightarrow \mathcal{C}l(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*).$$

To describe this diffeomorphism, it suffices to recall that for any equivalence class of form

$$[v_1 \otimes \dots \otimes v_k] \in \left((\pi_1 \cup_{(\tilde{f}, f)} \pi_2)^{\mathcal{C}l} \right)^{-1} (Y) \subset \mathcal{C}l((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^*),$$

with a representative $v_1 \otimes \dots \otimes v_k$, its image is the equivalence class in $\mathcal{C}l(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$ of

$$\Phi_{\cup,*}(v'_1) \otimes \dots \otimes \Phi_{\cup,*}(v'_k).$$

In other words, $\Phi_{\cup,*}^{\mathcal{C}l}$ is the pushdown, by the quotient projections

$$\pi^{T((V_1 \cup_{\tilde{f}} V_2)^*)} : T((V_1 \cup_{\tilde{f}} V_2)^*) \rightarrow \mathcal{C}l((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^*) \text{ and } \pi^{T(V_2^* \cup_{\tilde{f}^*} V_1^*)} : T(V_2^* \cup_{\tilde{f}^*} V_1^*) \rightarrow \mathcal{C}l(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*),$$

of the map $\bigoplus_n \Phi_{\cup,*}^{\otimes n}$, so that we have

$$\Phi_{\cup,*}^{\mathcal{C}l} \circ \pi^{T((V_1 \cup_{\tilde{f}} V_2)^*)} = \pi^{T(V_2^* \cup_{\tilde{f}^*} V_1^*)} \circ \left(\bigoplus_n \Phi_{\cup,*}^{\otimes n} \right).$$

Theorem 6.1. *The map $\Phi_{\cup,*}^{\mathcal{C}l} : \mathcal{C}l((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^*) \rightarrow \mathcal{C}l(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$ given by the identity*

$$\Phi_{\cup,*}^{\mathcal{C}l} \circ \pi^{T((V_1 \cup_{\tilde{f}} V_2)^*)} = \pi^{T(V_2^* \cup_{\tilde{f}^*} V_1^*)} \circ \left(\bigoplus_n \Phi_{\cup,*}^{\otimes n} \right)$$

is a well-defined diffeomorphism.

Proof. This is a consequence of Corollary 5.2. □

6.2 The pseudo-bundle $\mathcal{C}l(V_2^*, g_2^*) \cup_{(\tilde{F}^*)^{\mathcal{C}l}} \mathcal{C}l(V_1^*, g_1^*)$

There is a third pseudo-bundle, with all fibres Clifford algebras, that is naturally associated to our data, the pseudo-bundle $\mathcal{C}l(V_2^*, g_2^*) \cup_{(\tilde{F}^*)^{\mathcal{C}l}} \mathcal{C}l(V_1^*, g_1^*)$. It is obtained by gluing together the pseudo-bundles of Clifford algebras relative to, respectively, (V_2^*, g_2^*) and (V_1^*, g_1^*) , along the map $(\tilde{F}^*)^{\mathcal{C}l}$, induced by the map \tilde{f}^* ; this is the construction described in Section 1.5.2. To indicate how this construction works specifically for this case, it suffices to say that, for any given equivalence class in $((\pi_2^*)^{\mathcal{C}l})^{-1}(f(Y)) \subset \mathcal{C}l(V_2^*, g_2^*)$, with an arbitrary representative $v_1^* \otimes \dots \otimes v_k^*$, its image under $(\tilde{F}^*)^{\mathcal{C}l}$ is the equivalence class in $\mathcal{C}l(V_1^*, g_1^*)$ of the element

$$\tilde{f}^*(v_1^*) \otimes \dots \otimes \tilde{f}^*(v_k^*).$$

The main point, however, is that the resulting pseudo-bundle $\mathcal{C}l(V_2^*, g_2^*) \cup_{(\tilde{F}^*)^{\mathcal{C}l}} \mathcal{C}l(V_1^*, g_1^*)$ is naturally diffeomorphic to the pseudo-bundle $\mathcal{C}l(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$ via the diffeomorphism

$$\Phi^{\mathcal{C}l(*)} : \mathcal{C}l(V_2^*, g_2^*) \cup_{(\tilde{F}^*)^{\mathcal{C}l}} \mathcal{C}l(V_1^*, g_1^*) \rightarrow \mathcal{C}l(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$$

that is determined by the following formula:

$$\Phi^{\mathcal{C}l(*)} = \begin{cases} (j_1^{V_2^*})^{\mathcal{C}l} \circ (j_1^{\mathcal{C}l(V_2^*)})^{-1} & \text{on } \left((\pi_2^*)^{\mathcal{C}l} \cup_{((\tilde{F}^*)^{\mathcal{C}l}, f^{-1})} (\pi_1^*)^{\mathcal{C}l} \right)^{-1} (i_1^{X_2}(X_2 \setminus f(Y))) \\ (j_2^{V_1^*})^{\mathcal{C}l} \circ (j_2^{\mathcal{C}l(V_1^*)})^{-1} & \text{on } \left((\pi_2^*)^{\mathcal{C}l} \cup_{((\tilde{F}^*)^{\mathcal{C}l}, f^{-1})} (\pi_1^*)^{\mathcal{C}l} \right)^{-1} (i_2^{X_1}(X_1)), \end{cases}$$

where

$$(j_1^{V_2^*})^{\mathcal{C}\ell} : \mathcal{C}\ell(V_2^*, g_2^*) \setminus ((\pi_2^*)^{\mathcal{C}\ell})^{-1}(f(Y)) \rightarrow \mathcal{C}\ell(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*) \text{ and } (j_2^{V_1^*})^{\mathcal{C}\ell} : \mathcal{C}\ell(V_1^*, g_1^*) \rightarrow \mathcal{C}\ell(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$$

are the fibrewise extensions to $\mathcal{C}\ell(V_2^*, g_2^*) \setminus ((\pi_2^*)^{\mathcal{C}\ell})^{-1}(f(Y))$ and to $\mathcal{C}\ell(V_1^*, g_1^*)$, respectively, of the two natural inclusions

$$V_2^* \setminus (\pi_2^*)^{-1}(f(Y)) \hookrightarrow \mathcal{C}\ell(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*) \text{ and } V_1^* \hookrightarrow \mathcal{C}\ell(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*).$$

The latter inclusions are in turn given by the compositions of either $j_1^{V_2^*} : V_2^* \setminus (\pi_2^*)^{-1}(f(Y)) \rightarrow V_2^* \cup_{\tilde{f}^*} V_1^*$ or $j_2^{V_1^*} : V_1^* \rightarrow V_2^* \cup_{\tilde{f}^*} V_1^*$, with the standard inclusion $V_2^* \cup_{\tilde{f}^*} V_1^* \rightarrow \mathcal{C}\ell(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$.

Theorem 6.2. *The map $\Phi^{\mathcal{C}\ell(*)} : \mathcal{C}\ell(V_2^*, g_2^*) \cup_{(\tilde{F}^*)^{\mathcal{C}\ell}} \mathcal{C}\ell(V_1^*, g_1^*) \rightarrow \mathcal{C}\ell(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$ thus defined is a smooth diffeomorphism.*

Proof. This is a consequence of Theorem 5.5 in [8], applied to V_2^* , V_1^* , and \tilde{f}^* ; notice that \tilde{g}^* is exactly the counterpart of the pseudo-metric \tilde{g} that appears in the statement of the theorem just cited, in that it is obtained from g_2^* and g_1^* in precisely the same way as \tilde{g} is obtained from g_1 and g_2 . \square

6.3 The three diffeomorphisms $\Phi_{\cup, *}^{\mathcal{C}\ell}$, $\Phi^{\mathcal{C}\ell(*)}$, and $\Phi_{\cup}^{\mathcal{C}\ell(*)}$

We now summarize the above by listing the three possible forms of the Clifford algebra pseudo-bundle, together with the assumptions that allow us to identify them to each other, and with the corresponding diffeomorphisms.

The assumptions As (almost) everywhere throughout the paper, we consider two finite-dimensional diffeological vector pseudo-bundles $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$, and a gluing of the former to the latter along the pair (\tilde{f}, f) . In order for the three diffeomorphisms to exist, we must also assume the following:

- the two pseudo-bundles are locally trivial;²⁹
- f and \tilde{f}^* are diffeomorphisms;
- V_1 and V_2 admit compatible pseudo-metrics g_1 and g_2 respectively.

The three shapes of the pseudo-bundle of covariant Clifford algebras, and their equivalences

Under the assumptions just listed, the following three pseudo-bundles are well-defined:

$$\mathcal{C}\ell((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^*), \quad \mathcal{C}\ell(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*), \quad \mathcal{C}\ell(V_2^*, g_2^*) \cup_{(\tilde{F}^*)^{\mathcal{C}\ell}} \mathcal{C}\ell(V_1^*, g_1^*).$$

Although *a priori* these could be three different pseudo-bundles, the same assumptions that guarantee that all three are well-defined at the same time, also guarantee that they are in fact equivalent, via the following diffeomorphisms, already described above:

- $\Phi_{\cup, *}^{\mathcal{C}\ell} : \mathcal{C}\ell((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^*) \rightarrow \mathcal{C}\ell(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$;
- $\Phi^{\mathcal{C}\ell(*)} : \mathcal{C}\ell(V_2^*, g_2^*) \cup_{(\tilde{F}^*)^{\mathcal{C}\ell}} \mathcal{C}\ell(V_1^*, g_1^*) \rightarrow \mathcal{C}\ell(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$; and
- $(\Phi^{\mathcal{C}\ell(*)})^{-1} \circ \Phi_{\cup, *}^{\mathcal{C}\ell} =: \Phi_{\cup}^{\mathcal{C}\ell(*)} : \mathcal{C}\ell((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^*) \rightarrow \mathcal{C}\ell(V_2^*, g_2^*) \cup_{(\tilde{F}^*)^{\mathcal{C}\ell}} \mathcal{C}\ell(V_1^*, g_1^*)$.

7 The pseudo-bundles of exterior algebras, and gluing

We now turn to considering the pseudo-bundles of exterior algebras, first in the contravariant case, and then in the covariant case. We recall their construction, which is standard, and concentrate on the interactions with the operation of gluing.

²⁹This is sufficient but not necessary. What we really need is that the right inverse of the pairing map, that takes values in the characteristic sub-bundle be smooth, and so it suffices that this sub-bundle split off as a smooth direct summand.

7.1 The contravariant case

We first consider the contravariant version of the exterior algebra, by which we mean the following. Let first V be a diffeological vector space; for each tensor power of V consider the alternating operator³⁰

$$\text{Alt} : \underbrace{V \otimes \dots \otimes V}_n \rightarrow \underbrace{V \otimes \dots \otimes V}_n,$$

acting, as usual, by

$$\text{Alt}(v_1 \otimes \dots \otimes v_n) = \frac{1}{n!} \sum_{\sigma} (-1)^{\text{sgn}(\sigma)} v_{\sigma(1)} \otimes \dots \otimes v_{\sigma(n)}$$

and extended by linearity. In this section the n -th exterior power of V is the image

$$\bigwedge_n(V) = \text{Alt}(\underbrace{V \otimes \dots \otimes V}_n);$$

the whole exterior algebra $\bigwedge_*(V)$ is the direct sum of all $\bigwedge_n(V)$. We obtain the pseudo-bundle $\bigwedge_*(V)$ of exterior algebras associated to a given pseudo-bundle $\pi : V \rightarrow X$ by employing the same operations in the pseudo-bundle version, and defining the alternating operator fibrewise.

7.1.1 The induced gluing map \tilde{f}^{\wedge_*}

This map is provided by the universal factorization property for alternating maps. Specifically, let $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$ be two finite-dimensional diffeological vector pseudo-bundles, and let (\tilde{f}, f) be a gluing between them. Then the restriction $\tilde{f}|_{\pi_1^{-1}(y)}$ of \tilde{f} to each fibre in its domain of definition is a smooth linear map between diffeological vector spaces $\pi_1^{-1}(y)$ and $\pi_2^{-1}(f(y))$, and the direct sum of all tensor degrees of $\tilde{f}|_{\pi_1^{-1}(y)}$ is again a smooth linear map between the tensor algebras of these spaces:

$$\bigoplus_n \left(\tilde{f}|_{\pi_1^{-1}(y)} \right)^{\otimes n} : T(\pi_1^{-1}(y)) \rightarrow T(\pi_2^{-1}(f(y))).$$

By construction, this map commutes with the two respective alternating operators, so its restriction, that we denote by $\left(\tilde{f}|_{\pi_1^{-1}(y)} \right)^{\wedge_*}$, to $\bigwedge_*(\pi_1^{-1}(y))$ is a smooth linear map between the exterior algebras of the two fibres:

$$\left(\tilde{f}|_{\pi_1^{-1}(y)} \right)^{\wedge_*} : \bigwedge_* (\pi_1^{-1}(y)) \rightarrow \bigwedge_* (\pi_2^{-1}(f(y))).$$

Finally, the collection

$$\tilde{f}^{\wedge_*} := \bigcup_{y \in Y} \left(\tilde{f}|_{\pi_1^{-1}(y)} \right)^{\wedge_*},$$

where Y is the domain of definition of f , yields a smooth and fibrewise linear map \tilde{f}^{\wedge_*} between the appropriate subsets of $\bigwedge_*(V_1)$ and $\bigwedge_*(V_2)$. Thus, it yields an induced gluing between the corresponding pseudo-bundles of contravariant exterior algebras.

7.1.2 The pseudo-bundles $\bigwedge_*(V_1 \cup_{\tilde{f}} V_2)$ and $\bigwedge_*(V_1) \cup_{\tilde{f}^{\wedge_*}} \bigwedge_*(V_2)$

In a similar manner, for the pseudo-bundle $V_1 \cup_{\tilde{f}} V_2$ there is its own alternating operator Alt , whose image is the pseudo-bundle $\bigwedge_*(V_1 \cup_{\tilde{f}} V_2)$. Since each fibre of $T(V_1 \cup_{\tilde{f}} V_2)$ coincides with either a fibre of $T(V_1)$ or one of $T(V_2)$, and fibrewise each of the three alternating operators under consideration (those relative to V_1 , V_2 , and $V_1 \cup_{\tilde{f}} V_2$) is the usual one of a diffeological vector space, it makes sense to expect the two pseudo-bundles of exterior algebras, $\bigwedge_*(V_1 \cup_{\tilde{f}} V_2)$ and $\bigwedge_*(V_1) \cup_{\tilde{f}^{\wedge_*}} \bigwedge_*(V_2)$, to be diffeomorphic. Indeed, they are, and it is not difficult to describe the natural diffeomorphism between them; it is based on the gluing-tensor product commutativity diffeomorphism $\Phi_{\cup, \otimes}$, as the next construction shows.

³⁰In other terms, the antisymmetrization operator.

A preliminary remark At this moment we explicitly impose the assumption that for each of our pseudo-bundles V_1 and V_2 (and accordingly, for all the results of their gluings, their duals, and any mixture of such), the set of the dimensions of their fibres has a finite upper limit. We denote it by

$$\dim_V = \sup_{x_1 \in X_1, x_2 \in X_2} \{\dim(\pi_1^{-1}(x_1)), \dim(\pi_2^{-1}(x_2))\};$$

we do not go into any detail about how this correlates with any other assumptions of ours, just note that we will need for one of the diffeomorphisms that we define in the next paragraph (specifically, we use to ensure that $\Phi^{\wedge*}$ is indeed onto).

The diffeomorphism $\Phi_{\cup, \otimes}^{(\otimes n)}$: $(V_1 \cup_{\tilde{f}} V_2)^{\otimes n} \rightarrow V_1^{\otimes n} \cup_{\tilde{f}^{\otimes n}} V_2^{\otimes n}$ We first describe the construction of this diffeomorphism, which is by induction on n . The base of the induction is $n = 2$, in which case $\Phi_{\cup, \otimes}^{(\otimes 2)} = \Phi_{\cup, \otimes}$, the already-mentioned gluing-tensor product commutativity diffeomorphism. Suppose that $\Phi_{\cup, \otimes}^{(\otimes(n-1))}$ has already been defined. Then $\Phi_{\cup, \otimes}^{(\otimes n)}$ is obtained as the composition

$$(V_1 \cup_{\tilde{f}} V_2)^{\otimes(n-1)} \otimes (V_1 \cup_{\tilde{f}} V_2) \rightarrow (V_1^{\otimes(n-1)} \cup_{\tilde{f}^{\otimes(n-1)}} V_2^{\otimes(n-1)}) \otimes (V_1 \cup_{\tilde{f}} V_2) \rightarrow V_1^{\otimes n} \cup_{\tilde{f}^{\otimes n}} V_2^{\otimes n},$$

where the first arrow stands for $\Phi_{\cup, \otimes}^{(\otimes(n-1))} \otimes \text{Id}_{V_1 \cup_{\tilde{f}} V_2}$, and the second one, for the version $\Phi_{\cup, \otimes}^{\tilde{f}^{\otimes(n-1)}, \tilde{f}}$ of the gluing-tensor product commutativity diffeomorphism applied to the case of two factors, $V_1^{\otimes(n-1)} \cup_{\tilde{f}^{\otimes(n-1)}} V_2^{\otimes(n-1)}$ and $V_1 \cup_{\tilde{f}} V_2$. We can summarize the whole construction as

$$\Phi_{\cup, \otimes}^{(\otimes n)} = \Phi_{\cup, \otimes}^{\tilde{f}^{\otimes(n-1)}, \tilde{f}} \circ \left(\Phi_{\cup, \otimes}^{(\otimes(n-1))} \otimes \text{Id}_{V_1 \cup_{\tilde{f}} V_2} \right).$$

Note also that we will denote the inverse of $\Phi_{\cup, \otimes}^{(\otimes n)}$ by $\Phi_{\otimes, \cup}^{(\otimes n)}$. Finally, for all k (limited in practice by \dim_V) we define a diffeomorphism

$$\Phi_{\cup, \oplus}^{(k)} : \bigoplus_{n=0}^k (V_1^{\otimes n} \cup_{\tilde{f}^{\otimes n}} V_2^{\otimes n}) \rightarrow (\bigoplus_n V_1^{\otimes n}) \cup_{\bigoplus_n \tilde{f}^{\otimes n}} (\bigoplus_n V_2^{\otimes n});$$

the construction is exactly the same, just using the gluing-direct sum commutativity diffeomorphism $\Phi_{\cup, \oplus}$ in place of $\Phi_{\cup, \otimes}$.

The diffeomorphism $\Phi_{\cup, \otimes}^{(\otimes n)}$, and the alternating operators We write $\text{Alt}^{(n)}$ for the restriction of the alternating operator Alt on $V_1 \cup_{\tilde{f}} V_2$ onto the n -th tensor degree; likewise, $\text{Alt}_i^{(n)}$ stands for the same restriction of the alternating operator on V_i , for $i = 1, 2$. It is then quite trivial to observe that we have

$$\text{Alt}^{(n)} = \Phi_{\otimes, \cup}^{(\otimes n)} \circ \left(\text{Alt}_1^{(n)} \cup_{(\tilde{f}^{\otimes n}, \tilde{f}^{\otimes})} \text{Alt}_2^{(n)} \right) \circ \Phi_{\cup, \otimes}^{(\otimes n)};$$

equivalently,

$$\Phi_{\cup, \otimes}^{(\otimes n)} \circ \text{Alt}^{(n)} = \left(\text{Alt}_1^{(n)} \cup_{(\tilde{f}^{\otimes n}, \tilde{f}^{\otimes})} \text{Alt}_2^{(n)} \right) \circ \Phi_{\cup, \otimes}^{(\otimes n)}.$$

The diffeomorphism $\Phi^{\wedge*}$: $\Lambda_*(V_1 \cup_{\tilde{f}} V_2) \rightarrow \Lambda_*(V_1) \cup_{\tilde{f}^{\wedge*}} \Lambda_*(V_2)$ We can now define the desired diffeomorphism as

$$\Phi^{\wedge*} = \Phi_{\cup, \oplus}^{(\dim_V)} \circ \bigoplus_n \Phi_{\cup, \otimes}^{(\otimes n)} |_{\Lambda_*(V_1 \cup_{\tilde{f}} V_2)}.$$

It remains to observe that $\Phi^{\wedge*}$ is indeed onto $\Lambda_*(V_1) \cup_{\tilde{f}^{\wedge*}} \Lambda_*(V_2)$, as follows from its commutativity (in the sense explained in the previous paragraph) with the alternating operators.

Theorem 7.1. *The map*

$$\Phi^{\wedge*} = \Phi_{\cup, \oplus}^{(\dim_V)} \circ \bigoplus_n \Phi_{\cup, \otimes}^{(\otimes n)} |_{\Lambda_*(V_1 \cup_{\tilde{f}} V_2)}$$

is a pseudo-bundle diffeomorphism $\Lambda_(V_1 \cup_{\tilde{f}} V_2) \rightarrow \Lambda_*(V_1) \cup_{\tilde{f}^{\wedge*}} \Lambda_*(V_2)$ covering the identity map on $X_1 \cup_f X_2$.*

7.2 The pseudo-bundles of covariant exterior algebras

We now consider the covariant case. The basic definition is simple: the **covariant exterior algebra** $\bigwedge(V)$ of a pseudo-bundle V is $\bigwedge_*(V^*)$, the contravariant exterior algebra of its dual pseudo-bundle. So the reason why we consider it separately is to study its behavior with respect to the gluing, which, as we know, is not always well-behaved with respect to duality.

7.2.1 The induced map between $\bigwedge(V_2)$ and $\bigwedge(V_1)$

Indeed, let $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$ be two finite-dimensional diffeological vector pseudo-bundles, and let (\tilde{f}, f) be a gluing between them such that f is invertible. The gluing map between $\bigwedge(V_2)$ and $\bigwedge(V_1)$ is defined exactly as \tilde{f}^\wedge , but it is based on the dual map \tilde{f}^* . This gluing map is denoted by \tilde{f}^\wedge and is in fact

$$\tilde{f}^\wedge := (\tilde{f}^*)^\wedge.$$

7.2.2 The pseudo-bundles $\bigwedge(V_1 \cup_{\tilde{f}} V_2)$ and $\bigwedge(V_2) \cup_{\tilde{f}^\wedge} \bigwedge(V_1)$

As in the contravariant case, there are two natural pseudo-bundles of exterior algebras to consider, namely those mentioned in the title of this section. It is also natural to wonder whether they are diffeomorphic; we show that indeed they are, under the assumption that the gluing-dual commutativity condition is satisfied, by constructing a certain pseudo-bundle diffeomorphism

$$\Phi^\wedge : \bigwedge(V_1 \cup_{\tilde{f}} V_2) \rightarrow \bigwedge(V_2) \cup_{\tilde{f}^\wedge} \bigwedge(V_1)$$

covering the switch map.

The n -th degree component of Φ^\wedge Let $\Phi_{\cup, \otimes}^{(\otimes n)} : (V_2^* \cup_{\tilde{f}^*} V_1^*)^{\otimes n} \rightarrow (V_2^*)^{\otimes n} \cup_{(\tilde{f}^*)^{\otimes n}} (V_1^*)^{\otimes n}$ be the diffeomorphism constructed in the previous section. The n -th tensor degree component of Φ^\wedge is the composition

$$\Phi_{\cup, \otimes}^{(\otimes n)} \circ \Phi_{\cup, *}^{\otimes n} : ((V_1 \cup_{\tilde{f}} V_2)^*)^{\otimes n} \rightarrow (V_2^* \cup_{\tilde{f}^*} V_1^*)^{\otimes n} \rightarrow (V_2^*)^{\otimes n} \cup_{(\tilde{f}^*)^{\otimes n}} (V_1^*)^{\otimes n}.$$

Notice that if

$$\text{Alt}_{\cup, *}: ((V_1 \cup_{\tilde{f}} V_2)^*)^{\otimes n} \rightarrow ((V_1 \cup_{\tilde{f}} V_2)^*)^{\otimes n}, \quad \text{Alt}_2 : (V_2^*)^{\otimes n} \rightarrow (V_2^*)^{\otimes n} \quad \text{and} \quad \text{Alt}_1 : (V_1^*)^{\otimes n} \rightarrow (V_1^*)^{\otimes n}$$

are the n -th degree alternating operators on $(V_1 \cup_{\tilde{f}} V_2)^*$, V_2^* , and V_1^* respectively, then we have

$$\left(\Phi_{\cup, \otimes}^{(\otimes n)} \circ \Phi_{\cup, *}^{\otimes n} \right) \circ \text{Alt}_{\cup, *} = \left(\text{Alt}_2 \cup_{((\tilde{f}^*)^{\otimes n}, (\tilde{f}^*)^{\otimes n})} \text{Alt}_1 \right) \circ \left(\Phi_{\cup, \otimes}^{(\otimes n)} \circ \Phi_{\cup, *}^{\otimes n} \right).$$

The diffeomorphism Φ^\wedge We now employ also the gluing-direct sum commutativity diffeomorphism $\Phi_{\cup, \oplus}^{(\dim_{V^*})}$, also from the previous section, to obtain Φ^\wedge . Indeed, we define

$$\Phi^\wedge = \Phi_{\cup, \oplus}^{(\dim_{V^*})} \circ \bigoplus_{n=0}^{(\dim_{V^*})} \left(\Phi_{\cup, \otimes}^{(\otimes n)} \circ \Phi_{\cup, *}^{\otimes n} \right) \Big|_{\bigwedge(V_1 \cup_{\tilde{f}} V_2)}.$$

This is a well-defined injective, smooth and fibrewise linear map on $\bigwedge(V_1 \cup_{\tilde{f}} V_2)$; that its image is $\bigwedge(V_2) \cup_{\tilde{f}^\wedge} \bigwedge(V_1)$, follows from the commutativity between each $\Phi_{\cup, \otimes}^{(\otimes n)} \circ \Phi_{\cup, *}^{\otimes n}$ and the appropriate alternating operators (see above).

Theorem 7.2. *The map*

$$\Phi^\wedge = \Phi_{\cup, \oplus}^{(\dim_{V^*})} \circ \bigoplus_{n=0}^{(\dim_{V^*})} \left(\Phi_{\cup, \otimes}^{(\otimes n)} \circ \Phi_{\cup, *}^{\otimes n} \right) \Big|_{\bigwedge(V_1 \cup_{\tilde{f}} V_2)}$$

is a pseudo-bundle diffeomorphism $\bigwedge(V_1 \cup_{\tilde{f}} V_2) \rightarrow \bigwedge(V_2) \cup_{\tilde{f}^\wedge} \bigwedge(V_1)$ covering the switch map $\varphi_{X_1 \leftrightarrow X_2}$.

8 The Clifford actions

In this section we consider all possible (shapes of) Clifford actions, first outlining what acts on what, and then establishing the various natural equivalences.

8.1 The outline

As we have seen in the preceding sections, there is a multitude of formally distinct, but (as we are about to see) equivalent with respect to the diffeomorphisms described in the previous two sections, Clifford actions relative to a given gluing of (V_1, g_1) and (V_2, g_2) . In this section we give a list of these actions and their equivalences, with proofs and details appearing in the two sections immediately following. As before, we assume that f and \tilde{f}^* are diffeomorphisms, and g_1 and g_2 are compatible.

8.1.1 The contravariant case

Recall that in this case we have two natural Clifford algebras,³¹ specifically

$$\mathcal{C}l(V_1 \cup_{\tilde{f}} V_2, \tilde{g}) \cong \mathcal{C}l(V_1, g_1) \cup_{\tilde{f}^{\mathcal{C}l}} \mathcal{C}l(V_2, g_2)$$

(recall that the diffeomorphism that we have between them is $\Phi^{\mathcal{C}l} : \mathcal{C}l(V_1, g_1) \cup_{\tilde{f}^{\mathcal{C}l}} \mathcal{C}l(V_2, g_2) \rightarrow \mathcal{C}l(V_1 \cup_{\tilde{f}} V_2, \tilde{g})$) and two natural exterior algebras,

$$\bigwedge_{*} (V_1 \cup_{\tilde{f}} V_2) \cong \bigwedge_{*} (V_1) \cup_{\tilde{f}^{\wedge_*}} \bigwedge_{*} (V_2).$$

Summary of actions The natural actions are, the standard action c_* of $\mathcal{C}l(V_1 \cup_{\tilde{f}} V_2, \tilde{g})$ on $\bigwedge_{*} (V_1 \cup_{\tilde{f}} V_2)$, and the composite action $\tilde{c}_* := c_1 \cup_{(\tilde{f}^{\mathcal{C}l}, \tilde{f}^{\wedge_*})} c_2$ of $\mathcal{C}l(V_1, g_1) \cup_{\tilde{f}^{\mathcal{C}l}} \mathcal{C}l(V_2, g_2)$ on $\bigwedge_{*} (V_1) \cup_{\tilde{f}^{\wedge_*}} \bigwedge_{*} (V_2)$, where c_i is the standard action of $\mathcal{C}l(V_i, g_i)$ on $\bigwedge_{*} (V_i)$. We will show that this is a partial case of the construction considered in [8].

The equivalence of the two actions This is expressed by the formula:

$$\Phi^{\wedge_*} (c_*(v)(e)) = \tilde{c}_* ((\Phi^{\mathcal{C}l})^{-1}(v)) (\Phi^{\wedge_*}(e))$$

for all $v \in \mathcal{C}l(V_1 \cup_{\tilde{f}} V_2, \tilde{g})$ and $e \in \bigwedge_{*} (V_1 \cup_{\tilde{f}} V_2)$ such that $\pi^{\mathcal{C}l}(v) = \pi^{\wedge_*}(e)$. Below we will explain why this relation does hold.

8.1.2 The covariant case

There are three Clifford algebras to consider:

$$\mathcal{C}l((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^*) \cong \mathcal{C}l(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*) \cong \mathcal{C}l(V_2^*, g_2^*) \cup_{(\tilde{f}^*)^{\mathcal{C}l}} \mathcal{C}l(V_1^*, g_1^*),$$

and essentially two exterior algebras:

$$\bigwedge (V_1 \cup_{\tilde{f}} V_2) \cong \bigwedge (V_2) \cup_{\tilde{f}^{\wedge}} \bigwedge (V_1),$$

to which we will also add the contravariant exterior algebra $\bigwedge_{*} (V_2^* \cup_{\tilde{f}^*} V_1^*)$.

³¹A note on slight change in terminology: in the remainder of the paper we will just say *Clifford algebra* instead of a *pseudo-bundle of Clifford algebras*, and *exterior algebra* instead of *pseudo-bundle of exterior algebras*; in the present context this is unlikely to cause confusion.

Summary of Clifford actions We now outline which Clifford algebra (or the result of gluing of such) acts on which pseudo-bundle of exterior algebras:

- $\mathcal{Cl}((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^*)$ acts on $\Lambda(V_1 \cup_{\tilde{f}} V_2)$ via the standard Clifford action c^* ;
- $\mathcal{Cl}(V_2^*, g_2^*) \cup_{(\tilde{F}^*)^{\mathcal{Cl}}} \mathcal{Cl}(V_1^*, g_1^*)$ acts on $\Lambda(V_2) \cup_{\tilde{f}\Lambda} \Lambda(V_1)$ via the action $c_{*,\cup}$ (see Proposition 6.8) induced by the standard Clifford actions c_2^* and c_1^* of $\mathcal{Cl}(V_2^*, g_2^*)$ and $\mathcal{Cl}(V_1^*, g_1^*)$ on $\Lambda(V_2)$ and $\Lambda(V_1)$ respectively;
- $\mathcal{Cl}(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$ has, again, the standard Clifford action, which we have not mentioned yet and which we now denote by $\tilde{c}_{*,\cup}$, on $\Lambda_*(V_2^* \cup_{\tilde{f}^*} V_1^*)$.

The equivalence of actions As in the contravariant case, the actions c^* , $c_{*,\cup}$, and $\tilde{c}_{*,\cup}$ turn out to be equivalent, with the equivalence established via the diffeomorphisms $\Phi^{\mathcal{Cl}(\ast)}$, $\Phi_{\cup, \ast}^{\mathcal{Cl}}$, and $\Phi_{\cup}^{\mathcal{Cl}(\ast)}$, as well as Φ^Λ and $\Phi_{\cup, \ast}^\Lambda : \Lambda(V_1 \cup_{\tilde{f}} V_2) \rightarrow \Lambda_*(V_2^* \cup_{\tilde{f}^*} V_1^*)$. Specifically, we have:

- the action c^* of $\mathcal{Cl}((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^*)$ on $\Lambda(V_1 \cup_{\tilde{f}} V_2)$ is related to the action $\tilde{c}_{*,\cup}$ of $\mathcal{Cl}(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$ on $\Lambda_*(V_2^* \cup_{\tilde{f}^*} V_1^*)$, with respect to the diffeomorphisms $\Phi_{\cup, \ast}^{\mathcal{Cl}}$ and $\Phi_{\cup, \ast}^\Lambda$, via

$$\Phi_{\cup, \ast}^\Lambda(c^*(v)(e)) = \tilde{c}_{*,\cup}(\Phi_{\cup, \ast}^{\mathcal{Cl}}(v))(\Phi_{\cup, \ast}^\Lambda(e)),$$

that holds for all $v \in \mathcal{Cl}((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^*)$ and $e \in \Lambda(V_1 \cup_{\tilde{f}} V_2)$ such that $\pi^\Lambda(e) = \pi^{\mathcal{Cl}}(v)$;

- the action $c_{*,\cup}$ of $\mathcal{Cl}(V_2^*, g_2^*) \cup_{(\tilde{F}^*)^{\mathcal{Cl}}} \mathcal{Cl}(V_1^*, g_1^*)$ on $\Lambda(V_2) \cup_{\tilde{f}\Lambda} \Lambda(V_1)$ is equivalent to the action $\tilde{c}_{*,\cup}$ of $\mathcal{Cl}(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$ on $\Lambda_*(V_2^* \cup_{\tilde{f}^*} V_1^*)$, with respect to the diffeomorphisms $\Phi^{\mathcal{Cl}(\ast)}$ and $\Phi_{\cup}^{\mathcal{Cl}(\ast)}$, via

$$\left(\Phi_{\cup, \ast}^\Lambda \circ (\Phi^\Lambda)^{-1}\right)(c_{*,\cup}(v)(e)) = \tilde{c}_{*,\cup}\left(\Phi^{\mathcal{Cl}(\ast)}(v)\right)\left((\Phi_{\cup, \ast}^\Lambda \circ (\Phi^\Lambda)^{-1})(e)\right)$$

that is true for all $v \in \mathcal{Cl}(V_2^*, g_2^*) \cup_{(\tilde{F}^*)^{\mathcal{Cl}}} \mathcal{Cl}(V_1^*, g_1^*)$ and $e \in \Lambda(V_2) \cup_{\tilde{f}\Lambda} \Lambda(V_1)$ such that $(\pi_2^\Lambda \cup_{(\tilde{f}\Lambda, f^{-1})})\pi_1^\Lambda(e) = (\pi_2^{\mathcal{Cl}} \cup_{((\tilde{F}^*)^{\mathcal{Cl}}, f^{-1})})\pi_1^{\mathcal{Cl}}(v)$.

Notice that these equivalences imply also the equivalence of c^* to $\tilde{c}_{*,\cup}$.

8.2 The standard Clifford action is smooth

The basis for several versions of the Clifford(-type) actions listed above is the usual action of the (contravariant) Clifford algebra on the corresponding (also contravariant) exterior algebra. This means the following.

Let $\pi : V \rightarrow X$ be any locally trivial finite-dimensional diffeological vector pseudo-bundle that admits a pseudo-metric; let g be a fixed choice of a pseudo-metric on it. The **standard Clifford action** of $\mathcal{Cl}(V, g)$ on $\Lambda_*(V)$ is the map $c : \mathcal{Cl}(V, g) \rightarrow \mathcal{L}(\Lambda_*(V), \Lambda_*(V))$ given by

$$c(v)(v_1 \wedge \dots \wedge v_k) = v \wedge v_1 \wedge \dots \wedge v_k - \sum_{j=1}^k (-1)^{j+1} v_1 \wedge \dots \wedge g(\pi(v))(v, v_j) \wedge \dots \wedge v_k.$$

On each fibre $\pi^{-1}(x)$ of V , this is the usual Clifford action of the Clifford algebra relative to the bilinear symmetric form $g(x)$ on the exterior algebra of $\pi^{-1}(x)$.

Proposition 8.1. *The action c is smooth as a map $\mathcal{Cl}(V, g) \rightarrow \mathcal{L}(\Lambda_*(V), \Lambda_*(V))$.*

Proof. Notice first of all that the pseudo-bundle $\Lambda_*(V)$ smoothly splits as the direct sum $\bigoplus_k \Lambda^k V$. It then follows from the above presentation of the action c and the definition of the diffeology of $\mathcal{Cl}(V, g)$, that it suffices to show that the following two maps $c_V, c_j : V \rightarrow \mathcal{L}(\Lambda_*(V), \Lambda_*(V))$ are smooth:

$$c_V(v)(v_1 \wedge \dots \wedge v_k) = v \wedge v_1 \wedge \dots \wedge v_k \quad \text{and} \quad c_j(v)(v_1 \wedge \dots \wedge v_k) = v_1 \wedge \dots \wedge g(\pi(v))(v, v_j) \wedge \dots \wedge v_k.$$

Thus, c_V acts as the exterior product, which is smooth by definition (recall that the diffeology on each exterior product degree can be described as the pushforward of the tensor product diffeology by the alternating operator, which makes it, and the exterior product as a consequence, automatically smooth).

The smoothness of the map c_j follows from the smoothness of the pseudo-metric g . To be slightly more explicit, we note that on a small enough neighborhood U , we can write a plot of the k -th exterior degree as (p_1, \dots, p_k) , where each p_i is a plot of V , acting by $u \mapsto p_1(u) \wedge \dots \wedge p_k(u)$. Therefore the evaluation map that determines the smoothness of c_j is locally of form

$$(u', u) \mapsto p_1(u) \wedge \dots \wedge g(\pi(p(u')))(p(u'), p_j(u)) \wedge \dots \wedge p_k(u)$$

for some other plot $p : U' \rightarrow V$ of V . Since $(u', u) \mapsto g(\pi(p(u')))(p(u'), p_j(u))$ is a smooth function, and the diffeology of $\bigwedge_*(V)$ is a (vector) pseudo-bundle diffeology, we obtain a plot of $\bigwedge_*(V)$, whence the claim. \square

8.3 The compatibility of two standard Clifford actions

Likewise, we can show that under certain assumptions, two standard Clifford actions are compatible with a given gluing; this happens precisely when the gluing itself is commutative. Here is the precise statement.

Proposition 8.2. *Let $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$ be two locally trivial finite-dimensional diffeological vector pseudo-bundles, let (f, \tilde{f}) be a gluing between them such that f and \tilde{f} are diffeomorphisms, and let g_1 and g_2 be compatible pseudo-metrics on V_1 and V_2 respectively. Let c_i for $i = 1, 2$ be the standard Clifford actions of $\mathcal{C}l(V_i, g_i)$ on $\bigwedge_*(V_i)$. Then for all $v, v_1, \dots, v_k \in V_1$ such that $\pi_1(v) = \pi_1(v_1) = \dots = \pi_1(v_k) \in Y$ we have*

$$\tilde{f}^{\bigwedge_*}(c_1(v)(v_1 \wedge \dots \wedge v_k)) = c_2(\tilde{F}^{\mathcal{C}l}(v))(\tilde{f}^{\bigwedge_*}(v_1 \wedge \dots \wedge v_k)).$$

Proof. By the definition of $\tilde{F}^{\mathcal{C}l}$ and that of \tilde{f}^{\bigwedge_*} , we have that

$$c_2(\tilde{F}^{\mathcal{C}l}(v))(\tilde{f}^{\bigwedge_*}(v_1 \wedge \dots \wedge v_k)) = c_2(\tilde{f}(v))(\tilde{f}(v_1) \wedge \dots \wedge \tilde{f}(v_k)).$$

The desired condition easily follows from this. Indeed,

$$\begin{aligned} \tilde{f}^{\bigwedge_*}(c_1(v)(v_1 \wedge \dots \wedge v_k)) &= \\ &= \tilde{f}^{\bigwedge_*}(v \wedge v_1 \wedge \dots \wedge v_k - \sum_{j=1}^k (-1)^{j+1} g_1(\pi_1(v))(v, v_j) v_1 \wedge \dots \wedge v_{j-1} \wedge v_{j+1} \wedge \dots \wedge v_k) = \\ &= \tilde{f}(v) \wedge \tilde{f}(v_1) \wedge \dots \wedge \tilde{f}(v_k) - \\ &\quad - \sum_{j=1}^k (-1)^{j+1} g_1(\pi_1(v))(v, v_j) \tilde{f}(v_1) \wedge \dots \wedge \tilde{f}(v_{j-1}) \wedge \tilde{f}(v_{j+1}) \wedge \dots \wedge \tilde{f}(v_k). \end{aligned}$$

Now, since

$$\begin{aligned} c_2(\tilde{f}(v))(\tilde{f}(v_1) \wedge \dots \wedge \tilde{f}(v_k)) &= \tilde{f}(v) \wedge \tilde{f}(v_1) \wedge \dots \wedge \tilde{f}(v_k) - \\ &\quad - \sum_{j=1}^k (-1)^{j+1} g_2(\pi_2(\tilde{f}(v)))(\tilde{f}(v), \tilde{f}(v_j)) \tilde{f}(v_1) \wedge \dots \wedge \tilde{f}(v_{j-1}) \wedge \tilde{f}(v_{j+1}) \wedge \dots \wedge \tilde{f}(v_k). \end{aligned}$$

The pseudo-metrics g_1 and g_2 being compatible ensures that $g_2(\pi_2(\tilde{f}(v)))(\tilde{f}(v), \tilde{f}(v_j)) = g_1(\pi_1(v))(v, v_j)$, whence the claim. \square

8.4 The contravariant case: the actions on $\bigwedge_*(V_1 \cup_{\tilde{f}} V_2)$ and $\bigwedge_*(V_1) \cup_{\tilde{f}^{\bigwedge_*}} \bigwedge_*(V_2)$

We now describe the action \tilde{c}_* , and prove its equivalence (already stated above) to the action c_* . Notice that c_* is an instance of the standard Clifford action, so it is smooth by Lemma 8.1.

The action $\tilde{c}_* = c_1 \cup_{(\tilde{F}^{Cl}, \tilde{f}\wedge_*)} c_2$ This is a partial case of a more general construction described in [8]. The construction bears some similarity to that of the gluing of smooth maps, although, as mentioned in the same source, it is not quite the same thing. To describe this action, let $v \in cl(V_1, g_1) \cup_{\tilde{F}^{Cl}} Cl(V_2, g_2)$; then $\tilde{c}_*(v)$ is an endomorphism of the fibre of $\wedge_*(V_1) \cup_{\tilde{f}\wedge_*} \wedge_*(V_2)$ over the point $(\pi_1^{Cl} \cup_{(\tilde{F}^{Cl}, f)} \pi_2^{Cl})(v) \in X_1 \cup_f X_2$. Then the action \tilde{c}_* is defined as follows:

$$\tilde{c}_*(v)(e) = \begin{cases} j_1^{\wedge_*(V_1)} \left(c_1((j_1^{Cl(V_1, g_1)})^{-1}(v))((j_1^{\wedge_*(V_1)})^{-1}(e)) \right) & \text{over } i_1^{X_1}(X_1 \setminus Y) \\ j_2^{\wedge_*(V_2)} \left(c_2((j_2^{Cl(V_2, g_2)})^{-1}(v))((j_2^{\wedge_*(V_2)})^{-1}(e)) \right) & \text{over } i_2^{X_2}(X_2). \end{cases}$$

In other words, we just pull back v and e to the respective factors of gluing, apply c_1 or c_2 , as appropriate, and re-insert the result into the pseudo-bundle $\wedge_*(V_1) \cup_{\tilde{f}\wedge_*} \wedge_*(V_2)$. It now suffices to note that by Proposition 8.2 c_1 and c_2 are compatible as Clifford actions, so it follows from [8] that the action $\tilde{c}_{*, \cup}$ is smooth.

The equivalence of c_* to $\tilde{c}_{*, \cup}$ We now prove the already-mentioned statement of equivalence for these actions.

Theorem 8.3. *Let $v \in Cl(V_1 \cup_{\tilde{f}} V_2, \tilde{g})$ and $e \in \wedge_*(V_1 \cup_{\tilde{f}} V_2)$ be such that $\pi^{Cl}(v) = \pi^{\wedge_*}(e)$. Then*

$$\Phi^{\wedge_*}(c_*(v)(e)) = \tilde{c}_*((\Phi^{Cl})^{-1}(v)) \left(\Phi^{\wedge_*}(e) \right).$$

Proof. The proof is almost trivial if we adopt the following viewpoint: since both actions are fibrewise based on the standard Clifford action, it suffices to assume that v is an element of the copy of $V_1 \cup_{\tilde{f}} V_2$ naturally contained in $Cl(V_1 \cup_{\tilde{f}} V_2, \tilde{g})$, and that e belongs to the k -th exterior degree of $V_1 \cup_{\tilde{f}} V_2$, that is, $e = v_1 \wedge \dots \wedge v_k$ for $v_1, \dots, v_k \in V_1 \cup_{\tilde{f}} V_2$. Formally, there are two cases to consider: that of $\pi^{Cl}(v) \in i_1^{X_1}(X_1 \setminus Y)$ and that of $\pi^{Cl}(v) \in i_2^{X_2}(X_2)$.

Thus, suppose that $\pi^{Cl}(v) \in i_1^{X_1}(X_1 \setminus Y)$. Since c_* is the standard action, we have

$$c_*(v)(e) = v \wedge v_1 \wedge \dots \wedge v_k - \sum_{j=1}^k (-1)^{j+1} v_1 \wedge \dots \wedge \tilde{g}(\pi^{Cl}(v))(v, v_j) \wedge \dots \wedge v_k.$$

Therefore

$$\begin{aligned} \Phi^{\wedge_*}(c_*(v)(e)) &= j_1^{\wedge_*(V_1)}((j_1^{V_1})^{-1}(v) \wedge (j_1^{V_1})^{-1}(v_1) \wedge \dots \wedge (j_1^{V_1})^{-1}(v_k)) - \\ &\quad - \sum_{j=1}^k (-1)^{j+1} (j_1^{V_1})^{-1}(v_1) \wedge \dots \wedge g_1(\pi_1((j_1^{V_1})^{-1}(v))((j_1^{V_1})^{-1}(v), (j_1^{V_1})^{-1}(v_j))) \wedge \dots \wedge (j_1^{V_1})^{-1}(v_k). \end{aligned}$$

It therefore suffices to note that $j_1^{Cl(V_1, g_1)}((j_1^{V_1})^{-1}(v)) = (\Phi^{Cl})^{-1}(v)$ and $j_1^{\wedge_*(V_1)}((j_1^{V_1})^{-1}(v_1) \wedge \dots \wedge (j_1^{V_1})^{-1}(v_k)) = \Phi^{\wedge_*}(v_1 \wedge \dots \wedge v_k)$, to draw the desired conclusion. Since the treatment of the case $\pi^{Cl}(v) \in i_2^{X_2}(X_2)$ is exactly the same, the proof is finished. \square

8.5 The covariant case

In this case we have three potential actions, corresponding to the three shapes of the Clifford algebra and those of the three exterior algebras (one of which is actually a contravariant algebra, trivially identified to a covariant one). After a detailed description of the actions, we prove their equivalences, already announced in Section 8.1.2.

8.5.1 The action c^* of $Cl((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^*)$ on $\wedge(V_1 \cup_{\tilde{f}} V_2)$

This is a case of a standard Clifford action, considered in Lemma 8.1; this lemma, in particular, ensures, that c^* is a smooth action. Recall indeed that $\wedge(V_1 \cup_{\tilde{f}} V_2) = \wedge_*((V_1 \cup_{\tilde{f}} V_2)^*)$ by its definition.

8.5.2 The action $c_{*,\cup}$ of $\mathcal{Cl}(V_2^*, g_2^*) \cup_{(\tilde{F}^*)^{\mathcal{Cl}}} \mathcal{Cl}(V_1^*, g_1^*)$ on $\Lambda(V_2) \cup_{\tilde{f}\Lambda} \Lambda(V_1)$

Let $c_2^* : \mathcal{Cl}(V_2^*, g_2^*) \rightarrow \mathcal{L}(\Lambda(V_2), \Lambda(V_2))$ and $c_1^* : \mathcal{Cl}(V_1^*, g_1^*) \rightarrow \mathcal{L}(\Lambda(V_1), \Lambda(V_1))$ be the standard Clifford actions. By Proposition 8.2, they are compatible with the gluings that yield respectively $\mathcal{Cl}(V_2^*, g_2^*) \cup_{(\tilde{F}^*)^{\mathcal{Cl}}} \mathcal{Cl}(V_1^*, g_1^*)$ and $\Lambda(V_2) \cup_{\tilde{f}\Lambda} \Lambda(V_1)$, with respect to the maps $\tilde{f}\Lambda$ and $(\tilde{F}^*)^{\mathcal{Cl}}$. Thus, the procedure described in [8] yields a smooth action $c_{*,\cup} = c_2^* \cup_{((\tilde{F}^*)^{\mathcal{Cl}}, \tilde{f}\Lambda)} c_1^*$. The formula that describes it is as follows.

Let $v \in \mathcal{Cl}(V_2^*, g_2^*) \cup_{(\tilde{F}^*)^{\mathcal{Cl}}} \mathcal{Cl}(V_1^*, g_1^*)$ and $e \in \Lambda(V_2) \cup_{\tilde{f}\Lambda} \Lambda(V_1)$ be such that $(\pi_2^\Lambda \cup_{(\tilde{f}\Lambda, f^{-1})} \pi_1^\Lambda)(e) = (\pi_2^{\mathcal{Cl}} \cup_{((\tilde{F}^*)^{\mathcal{Cl}}, f^{-1})} \pi_1^{\mathcal{Cl}})(v)$. Then we will have

$$c_{*,\cup}(v)(e) = \begin{cases} j_1^{\Lambda(V_2)} \left(c_2^* \left((j_1^{\mathcal{Cl}(V_2^*, g_2^*)})^{-1}(v) \right) \left((j_1^{\Lambda(V_2)})^{-1}(e) \right) \right) & \text{over } i_1^{X_2}(X_2 \setminus f(Y)) \\ j_2^{\Lambda(V_1)} \left(c_1^* \left((j_2^{\mathcal{Cl}(V_1^*, g_1^*)})^{-1}(v) \right) \left((j_2^{\Lambda(V_1)})^{-1}(e) \right) \right) & \text{over } i_2^{X_1}(X_1). \end{cases}$$

8.5.3 The action $\tilde{c}_{*,\cup}$ of $\mathcal{Cl}(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$ on $\Lambda_*(V_2^* \cup_{\tilde{f}^*} V_1^*)$

This is also a standard Clifford action; its smoothness follows, once again, from Lemma 8.1.

8.5.4 The equivalence of c^* to $\tilde{c}_{*,\cup}$, and that of $c_{*,\cup}$ to $\tilde{c}_{*,\cup}$

It now remains to prove the two equivalence formulae for the covariant actions c^* , $c_{*,\cup}$, and $\tilde{c}_{*,\cup}$.

The diffeomorphism $\Phi_{\cup,*}^\Lambda$: $\Lambda(V_1 \cup_{\tilde{f}} V_2) \rightarrow \Lambda_*(V_2^* \cup_{\tilde{f}^*} V_1^*)$ This diffeomorphism is induced by the gluing-dual commutativity diffeomorphism $\Phi_{\cup,*}$ in a completely standard way and is, by definition, the map

$$\Phi_{\cup,*}^\Lambda = \bigoplus_{n=0}^{\dim_{V^*}} \Phi_{\cup,*}^{\otimes n} |_{\Lambda(V_1 \cup_{\tilde{f}} V_2)}.$$

That it has, in particular, the desired range follows from the commutativity of $\Phi_{\cup,*}^{\otimes n}$ with the relevant alternating operators.

The equivalence of c^* to $\tilde{c}_{*,\cup}$ We now show that the action c^* of $\mathcal{Cl}((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^*)$ on $\Lambda(V_1 \cup_{\tilde{f}} V_2)$ is equivalent to the action $\tilde{c}_{*,\cup}$ of $\mathcal{Cl}(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$ on $\Lambda_*(V_2^* \cup_{\tilde{f}^*} V_1^*)$, via the rule described in the following statement.

Theorem 8.4. *Let $v \in \mathcal{Cl}((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^*)$ and $e \in \Lambda(V_1 \cup_{\tilde{f}} V_2)$ be such that $\pi^\Lambda(e) = \pi^{\mathcal{Cl}}(v)$. Then*

$$\Phi_{\cup,*}^\Lambda(c^*(v)(e)) = \tilde{c}_{*,\cup}(\Phi_{\cup,*}^{\mathcal{Cl}}(v))(\Phi_{\cup,*}^\Lambda(e)).$$

Proof. The proof uses the same kind of reasoning as that of Theorem 8.3, in which it suffices observe that both diffeomorphisms $\Phi^{\mathcal{Cl}(\cdot)}$ and $\Phi_{\cup,*}^\Lambda$ are based on the same diffeomorphism $\Phi_{\cup,*}$, and that the latter essentially commutes with the exterior product. \square

The equivalence of $c_{*,\cup}$ to $\tilde{c}_{*,\cup}$ Let us now show that the action $c_{*,\cup}$ of $\mathcal{Cl}(V_2^*, g_2^*) \cup_{(\tilde{F}^*)^{\mathcal{Cl}}} \mathcal{Cl}(V_1^*, g_1^*)$ on $\Lambda(V_2) \cup_{\tilde{f}\Lambda} \Lambda(V_1)$ is equivalent to the action $\tilde{c}_{*,\cup}$ of $\mathcal{Cl}(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$ on $\Lambda_*(V_2^* \cup_{\tilde{f}^*} V_1^*)$. Specifically, we have the following statement.

Theorem 8.5. *Let $v \in \mathcal{Cl}(V_2^*, g_2^*) \cup_{(\tilde{F}^*)^{\mathcal{Cl}}} \mathcal{Cl}(V_1^*, g_1^*)$ and $e \in \Lambda(V_2) \cup_{\tilde{f}\Lambda} \Lambda(V_1)$ be such that $(\pi_2^\Lambda \cup_{(\tilde{f}\Lambda, f^{-1})} \pi_1^\Lambda)(e) = (\pi_2^{\mathcal{Cl}} \cup_{((\tilde{F}^*)^{\mathcal{Cl}}, f^{-1})} \pi_1^{\mathcal{Cl}})(v)$. Then*

$$\left(\Phi_{\cup,*}^\Lambda \circ (\Phi^\Lambda)^{-1} \right) (c_{*,\cup}(v)(e)) = \tilde{c}_{*,\cup} \left(\Phi^{\mathcal{Cl}(\cdot)}(v) \right) \left((\Phi_{\cup,*}^\Lambda \circ (\Phi^\Lambda)^{-1})(e) \right).$$

Proof. This is a direct consequence of Theorem 8.3 applied to (V_2^*, g_2^*) , (V_1^*, g_1^*) , and (\tilde{f}^*, f^{-1}) . \square

9 Examples

The two examples that we describe in this section are chosen with the following considerations in mind. For one thing, even when we start with some usual smooth vector bundles (as in the first example below), the gluing of them may be defined on a non-open set, producing a result which is not a smooth manifold, yet is being treated as if it were one. In the second example we consider a non-diffeomorphic gluing of fibres.

9.1 The wedge of two lines

We start with the two pseudo-bundles $\pi_1 : V_1 \rightarrow X_1$ and $\pi_2 : V_2 \rightarrow X_2$. Let $V_1 = V_2 = \mathbb{R}^2$ with the standard diffeology, let $X_1 = X_2 = \mathbb{R}$, also standard, and let π_1 and π_2 be the two standard projections on the x -axis: $\pi_1 : V_1 \ni (x, y) \mapsto x \in \mathbb{R} = X_1$ and $\pi_2 : V_2 \ni (x, y) \mapsto x \in \mathbb{R} = X_2$. The pseudo-bundle structure is given by imposing on each fibre $(x, y_1) + (x, y_2) \mapsto (x, y_1 + y_2)$ and $\lambda(x, y_1) \mapsto (x, \lambda y_1)$. The gluing of these two pseudo-bundles is given by (\tilde{f}, f) , where $f : X_1 \supset \{0\} \rightarrow \{0\} \subset X_2$ and \tilde{f} is determined by a non-zero constant $a \in \mathbb{R}$ via the rule

$$\tilde{f}(0, 1) = (0, a) \in V_2 = \mathbb{R}^2.$$

Let $f_1, f_2 : \mathbb{R} \rightarrow \mathbb{R}$ be two smooth everywhere positive functions; let g_i be the pseudo-metric on V_i given by

$$g_i(x)(v, w) = f_i(x) \cdot e^2(v) \cdot e^2(w),$$

where e^2 is the second element of the usual dual basis of the canonical basis of \mathbb{R}^2 (relative to the notation used later on it can be written as dy). The compatibility condition is then $f_1(0) = a^2 f_2(0)$.

The result of gluing The pseudo-bundle $V_1 \cup_{\tilde{f}} V_2$ can be described as the union $\{(x, 0, z)\} \cup \{(0, y, z)\}$ of two planes in \mathbb{R}^3 , and, accordingly, $X_1 \cup_f X_2$ as the union $\{(x, 0, 0)\} \cup \{(0, y, 0)\}$ of the two axes, with the projection $\pi_1 \cup_{(\tilde{f}, f)} \pi_2$ acting by $(x, 0, z) \mapsto (x, 0, 0)$, $(0, y, z) \mapsto (0, y, 0)$. The pseudo-metric \tilde{g} is then

$$\tilde{g}(x, y, 0) = \begin{cases} f_1(x) dz^2 & \text{if } y = 0 \text{ and } x \neq 0, \\ f_2(y) dz^2 & \text{if } x = 0. \end{cases}$$

The pairing maps Ψ_{g_1} , Ψ_{g_2} , and $\Psi_{\tilde{g}}$ Since all fibres are standard, the characteristic sub-bundles coincide with the pseudo-bundles themselves, so all three maps are automatically invertible. They act by:

$$\Psi_{g_1}(x, y) = f_1(x) y dy, \quad \Psi_{g_2}(x, y) = f_2(x) y dy,$$

and then, using the just-mentioned presentation of $V_1 \cup_{\tilde{f}} V_2$ as a subset of \mathbb{R}^3 , we have

$$\Psi_{\tilde{g}}(x, y, z) = \begin{cases} f_1(x) z dz & \text{if } y = 0, \\ f_2(y) z dz & \text{if } x = 0 \end{cases}$$

The dual pseudo-metrics The dual pseudo-metrics are therefore described in the same manner as g_1 and g_2 , but the coefficients are inverted:

$$g_2^*(x)(v^*, w^*) = \frac{1}{f_2(x)} \cdot v^*(e_2) \cdot w^*(e_2) \quad \text{and} \quad g_1^*(x)(v^*, w^*) = \frac{1}{f_1(x)} \cdot v^*(e_2) \cdot w^*(e_2).$$

The compatibility condition for g_2^* and g_1^* is thus $\frac{1}{f_2(0)} = \frac{a^2}{f_1(0)}$, and so is equivalent to the one for g_1 and g_2 . The pseudo-metric $\tilde{g}^* \equiv \tilde{g}^*$ is therefore

$$\tilde{g}^*(x', y', 0) = \begin{cases} \frac{1}{f_1(x')} \frac{\partial}{\partial z} \otimes \frac{\partial}{\partial z} & \text{if } y' = 0, \\ \frac{1}{f_2(x')} \frac{\partial}{\partial z} \otimes \frac{\partial}{\partial z} & \text{if } x' = 0. \end{cases}$$

The pseudo-bundles of Clifford algebras All fibres in our case are 1-dimensional, so each of $\mathcal{Cl}(V_1, g_1)$, $\mathcal{Cl}(V_2, g_2)$ is thus a trivial fibering of \mathbb{R}^3 over \mathbb{R} ; the result of their gluing can be described as the subset in \mathbb{R}^4 given by the equation $xy = 0$, so that

$$\mathcal{Cl}(V_1, g_1) \cup_{\tilde{F}^{\mathcal{Cl}}} \mathcal{Cl}(V_2, g_2) = \{(x, 0, z, w), \text{ where } x \neq 0\} \cup \{(0, y, z, w)\},$$

with the Clifford multiplication being defined by

$$\begin{aligned} (x, 0, z_1, w_1) \cdot_{\mathcal{Cl}} (x, 0, z_2, w_2) &= (x, 0, z_1 w_2 + z_2 w_1, -f_1(x) z_1 z_2 + w_1 w_2), \\ (0, y, z_1, w_1) \cdot_{\mathcal{Cl}} (0, y, z_2, w_2) &= (0, y, z_1 w_2 + z_2 w_1, -f_2(y) z_1 z_2 + w_1 w_2). \end{aligned}$$

From this, it is also quite evident that the result trivially coincides with $\mathcal{Cl}(V_1 \cup_{\tilde{F}} V_2, \tilde{g})$, so much in fact, that we can only distinguish between the two by choosing two slightly different forms of designating the same subset in \mathbb{R}^4 . Specifically, in the case of $\mathcal{Cl}(V_1 \cup_{\tilde{F}} V_2, \tilde{g})$ we describe the set of its points as

$$\{(x, y, z, w), \text{ where } xy = 0\}.$$

Obviously, this is the same set as we described as the set of points of $\mathcal{Cl}(V_1, g_1) \cup_{\tilde{F}^{\mathcal{Cl}}} \mathcal{Cl}(V_2, g_2)$; the chosen presentation of the latter emphasizes its structure as the result of a gluing.

The pseudo-bundles of covariant Clifford algebras Consider again the subset in \mathbb{R}^4 given by the equation $xy = 0$. This is the subset that is identified with all three (shapes of) the Clifford algebra. For all three possibilities, we identify the copy of V_1^* contained in either of them with the hyperplane $\{(x, 0, z, 0)\}$, and the copy of V_2^* , with the hyperplane $\{(0, y, z, 0)\}$; the fourth coordinate w corresponds to the scalar part of the Clifford algebra.

The distinction between the various shapes of the Clifford algebra is the following one. When this subset is viewed as $\mathcal{Cl}((V_1 \cup_{\tilde{F}} V_2)^*, \tilde{g}^*)$, we describe the Clifford multiplication as

$$\begin{cases} (x, 0, z_1, w_1) \cdot_{\mathcal{Cl}} (x, 0, z_2, w_2) = (x, 0, z_1 w_2 + z_2 w_1, -\frac{1}{f_1(x)} z_1 z_2 + w_1 w_2) \text{ for } x \neq 0, \\ (0, y, z_1, w_1) \cdot_{\mathcal{Cl}} (0, y, z_2, w_2) = (0, y, z_1 w_2 + z_2 w_1, -\frac{1}{f_2(y)} z_1 z_2 + w_1 w_2) \text{ otherwise.} \end{cases}$$

On the other hand, when we view the same subset as either $\mathcal{Cl}(V_2^*, g_2^*) \cup_{(\tilde{F}^*)^{\mathcal{Cl}}} \mathcal{Cl}(V_1^*, g_1^*)$ or $\mathcal{Cl}(V_2^* \cup_{\tilde{F}^*} V_1^*, \tilde{g}^*)$, we describe the corresponding product by

$$\begin{cases} (x, 0, z_1, w_1) \cdot_{\mathcal{Cl}} (x, 0, z_2, w_2) = (x, 0, z_1 w_2 + z_2 w_1, -\frac{1}{f_1(x)} z_1 z_2 + w_1 w_2) \text{ for all } x, \\ (0, y, z_1, w_1) \cdot_{\mathcal{Cl}} (0, y, z_2, w_2) = (0, y, z_1 w_2 + z_2 w_1, -\frac{1}{f_2(y)} z_1 z_2 + w_1 w_2) \text{ for } y \neq 0. \end{cases}$$

The pseudo-bundles of exterior algebras These can be presented in exactly the same way as those of Clifford algebras. In both the contravariant and the covariant case we have a unique presentation, again as a subset of \mathbb{R}^4 given by the equation $xy = 0$, with the exterior product

$$\begin{cases} (x, 0, z_1, w_1) \wedge (x, 0, z_2, w_2) = (x, 0, z_1 w_2 + z_2 w_1, w_1 w_2), \\ (0, y, z_1, w_1) \wedge (0, y, z_2, w_2) = (0, y, z_1 w_2 + z_2 w_1, w_1 w_2) \end{cases}$$

The Clifford actions In the contravariant case, we have two exterior algebras, $\bigwedge_*(V_1 \cup_{\tilde{F}} V_2)$ and $\bigwedge_*(V_1) \cup_{\tilde{F}^*} \bigwedge_*(V_2)$, with the actions c and \tilde{c} of, respectively, $\mathcal{Cl}(V_1 \cup_{\tilde{F}} V_2, \tilde{g})$ and $\mathcal{Cl}(V_1, g_1) \cup_{\tilde{F}^{\mathcal{Cl}}} \mathcal{Cl}(V_2, g_2)$. In the former case, we have

$$\begin{aligned} c((0, 0, z_2, w_2))(0, 0, z, w) &= (0, 0, z_2, w_2) \wedge (0, 0, z, w) - (0, 0, 0, \tilde{g}(0, 0, 0))((0, 0, z_2), (0, 0, z)) = \\ &= (0, 0, z_2 w + w_2 z, w_2 w + f_2(0) z_2 z); \end{aligned}$$

in the latter case, the only thing that changes with respect to the formula just given, is that the term $\tilde{g}((0, 0, 0)((0, 0, z_2), (0, 0, z)))$ is replaced by the term $g_2((0, 0, 0)((0, z_2), (0, z)))$, whose value however is exactly the same.

The covariant case is analogous, although we have three exterior algebras, $\Lambda(V_1 \cup_{\tilde{f}} V_2)$, $\Lambda_*(V_2^* \cup_{\tilde{f}^*} V_1^*)$, and $\Lambda(V_2) \cup_{(\tilde{f}^*)\wedge} \Lambda(V_1)$, with the actions c , $c_{*,\cup}$, and $c_{\cup,*}$ of, respectively, $Cl((V_1 \cup_{\tilde{f}} V_2)^*, \tilde{g}^*)$, $Cl(V_2^* \cup_{\tilde{f}^*} V_1^*, \tilde{g}^*)$, and $Cl(V_2^*, g_2^*) \cup_{(\tilde{f}^*)c\ell} Cl(V_1^*, g_1^*)$. Once again, these actions have the same form everywhere except over the point of gluing (the origin), where we would formally write the formulae for $c((0, 0, z_2, w_2))(0, 0, z, w)$, $c_{*,\cup}((0, 0, z_2, w_2))(0, 0, z, w)$, and $c_{\cup,*}((0, 0, z_2, w_2))(0, 0, z, w)$ with respect to \tilde{g}^* , g^* , or g_1^* , respectively:

$$\begin{aligned} c((0, 0, z_2, w_2))(0, 0, z, w) &= (0, 0, z_2, w_2) \wedge (0, 0, z, w) - (0, 0, 0, \tilde{g}^*(0, 0, 0)((0, 0, z_2), (0, 0, z))) = \\ &= (0, 0, z_2w + w_2z, w_2w + \frac{1}{f_2(0)}z_2z), \\ c_{*,\cup}((0, 0, z_2, w_2))(0, 0, z, w) &= (0, 0, z_2, w_2) \wedge (0, 0, z, w) - (0, 0, 0, \tilde{g}^*(0, 0, 0)((0, 0, z_2), (0, 0, z))) = \\ &= (0, 0, z_2w + w_2z, w_2w - g_1^*(0, 0)((0, z_2), (0, z))) = (0, 0, z_2w + w_2z, w_2w + \frac{1}{f_1(0)}z_2z), \\ c_{\cup,*}((0, 0, z_2, w_2))(0, 0, z, w) &= (0, 0, z_2, w_2) \wedge (0, 0, z, w) - (0, 0, 0, g_1^*(0, 0)((0, z_2), (0, z))) = \\ &= (0, 0, z_2w + w_2z, w_2w + \frac{1}{f_1(0)}z_2z). \end{aligned}$$

9.2 A non-diffeomorphism \tilde{f} and diffeomorphism \tilde{f}^*

Let $\pi_2 : V_2 \rightarrow X_2$ be the same as in the previous section, *i.e.*, the standard projection $\mathbb{R}^2 \rightarrow \mathbb{R}$; define $\pi_1 : V_1 \rightarrow X_1$ to be the projection of $V_1 = \mathbb{R}^3$ to $X_1 = \mathbb{R}$, where X_1 carries the standard diffeology, and $V_1 = \mathbb{R} \times \mathbb{R} \times \mathbb{R}$ carries the product diffeology relative to the standard diffeologies on the first two factors and the vector space diffeology generated by the plot $\mathbb{R} \ni x \mapsto |x|$ on the third factor.³² The projection π_1 is just the projection onto the first factor. The gluing map \tilde{f} for the bases is the same, $\{0\} \rightarrow \{0\}$, and the one for the total spaces is almost the same, specifically, $\tilde{f}(0, y, z) = (0, ay)$ with $a \neq 0$ (again, notice that zeroing out the third coordinate is necessary for \tilde{f} to be smooth). The pseudo-bundle $\pi_2 : V_2 \rightarrow X_2$ carries the same pseudo-metric g_2 as in the previous example, while the pseudo-metric g_1 on $\pi_1 : V_1 \rightarrow X_1$ extends the previous one in a trivial manner:

$$g_1(x)((x, y_1, z_1), (x, y_2, z_2)) = f_1(x)y_1y_2.$$

The compatibility condition remains the same.

The entire covariant case coincides with that of the example treated in the previous section. We only consider the pseudo-bundle $V_1 \cup_{\tilde{f}} V_2$ and the corresponding contravariant constructions.

The pseudo-bundle $V_1 \cup_{\tilde{f}} V_2$ We represent it as a subset in \mathbb{R}^4 , specifically as the union of the plane given by the equations $x = 0$ and $w = 0$ (the part corresponding to V_2), and of the set $\{y = 0\} \setminus \{x = 0, y = 0, w = 0\}$; this is the part corresponding to V_1 , where excising the line $\{x = 0, y = 0, w = 0\}$ reflects how $V_1 \cup_{\tilde{f}} V_2$ contains $V_1 \setminus \pi_1^{-1}(Y)$, and not the entire V_1 . Thus, the entire set can be described as

$$\begin{cases} (x, 0, z, w) & \text{except the points } (0, 0, z, 0) \\ (0, y, z, 0) & \text{for all } y, z. \end{cases}$$

The two Clifford algebras The Clifford algebra of V_2 is the already seen one; relative to the presentation of $V_1 \cup_{\tilde{f}} V_2$ given above, we could describe it as a subset of \mathbb{R}^5 , adding the 5th coordinate u_1 for the scalar part of $Cl(V_2, g_2) \cong \mathbb{R} \oplus V_2$. Thus,

$$Cl(V_2, g_2) = \{(0, y, z, 0, u_1)\},$$

³²In fact, any non-standard vector space diffeology would be sufficient for our purposes.

with the Clifford multiplication given by

$$(0, y, z', 0, u'_1) \cdot_{Cl} (0, y, z'', 0, u''_1) = (0, y, u'_1 z' + u'_1 z'', 0, u'_1 u''_1 - f_2(y) z' z'').$$

The Clifford algebra $Cl(V_1, g_1)$ is bigger; since the fibres of V_1 have dimension 2, each fibre of $cl(V_1, g_1)$ has dimension 4. Thus, we represent it as a subset in \mathbb{R}^6 , by adding the coordinates u_1, u_2 , where u_1 corresponds to the scalar part and u_2 corresponds to the degree 2 vector part. Thus,

$$Cl(V_1, g_1) = \{(x, 0, z, w, u_1, u_2)\},$$

with the Clifford multiplication given by

$$\begin{aligned} (x, 0, z', w', u'_1, u'_2) \cdot_{Cl} (x, 0, z'', w'', u''_1, u''_2) = \\ = (x, 0, z' u''_1 + z'' u'_1, w' u''_1 + w'' u'_1 + f_1(x) w'', u'_1 u''_1 - f_1(x) z' z'', u'_1 u''_2 + u'_1 u''_2). \end{aligned}$$

Finally, $Cl(V_1 \cup_{\tilde{f}} V_2, \tilde{g})$ can be described as the following subset in \mathbb{R}^6 :

$$\{(x, y, z, w, u_1, u_2) \text{ such that } xy = 0, x = 0 \Rightarrow w = u_2 = 0\},$$

while $Cl(V_1, g_1) \cup_{\tilde{f} \circ \ell} Cl(V_2, g_2)$ is presented as the subset in \mathbb{R}^6 of the following form:

$$\{(x, 0, z, w, u_1, u_2) \text{ such that } x \neq 0\} \cup \{(0, y, z, 0, u_1, 0) \text{ for all } y, z\}.$$

The fibrewise multiplication is described by uniting the two formulae just given.

The contravariant exterior algebras Likewise, the exterior algebras $\bigwedge_*(V_1)$ and $\bigwedge_*(V_2)$ are given by the same sets. Both of these we immediately represent as subsets of \mathbb{R}^6 , with the 5-th coordinate being the scalar part and the 6-th coordinate being the exterior product corresponding to the exterior product relative to the 3-rd and the 4-th coordinates; in the case of V_2 , this part is obviously trivial. Thus, we have

$$\bigwedge_*(V_1) = \{(x, 0, z, w, u_1, u_2)\}, \quad \bigwedge_*(V_2) = \{(0, y, z, 0, u_1, 0)\},$$

with the exterior product given by

$$\begin{aligned} (x, 0, z', w', u'_1, u'_2) \wedge (x, 0, z'', w'', u''_1, u''_2) = \\ = (x, 0, u'_1 z' + u'_1 z'', u''_1 w' + u'_1 w'', u'_1 u''_1, u''_1 u'_2 + u'_1 u''_2 + z' w'' - z'' w'), \end{aligned}$$

$$(0, y, z', 0, u'_1, 0) \wedge (0, y, z'', 0, u''_1, 0) = (0, y, u'_1 z' + u'_1 z'', 0, u'_1 u''_1, 0).$$

The exterior algebras $\bigwedge_*(V_1 \cup_{\tilde{f}} V_2)$ and $\bigwedge_*(V_1) \cup_{\tilde{f} \wedge_*} \bigwedge_*(V_2)$ are then represented respectively by the sets

$$\bigwedge_*(V_1 \cup_{\tilde{f}} V_2) = \{(x, y, z, w, u_1, u_2), \text{ where } xy = 0, x = 0 \Rightarrow w = u_2 = 0\},$$

$$\bigwedge_*(V_1) \cup_{\tilde{f} \wedge_*} \bigwedge_*(V_2) = \{(x, 0, z, w, u_1, u_2) \text{ such that } x \neq 0\} \cup \{(0, y, z, 0, u_1, 0)\}.$$

It is obvious that the two presentations determine the same set, with the second one possibly giving a better idea of the structure of the set, and the first one allowing for the uniform description of the exterior product, in the following way:

$$\begin{aligned} (x, y, z', w', u'_1, u'_2) \wedge (x, y, z'', w'', u''_1, u''_2) = \\ = (x, y, u'_1 z' + u'_1 z'', u''_1 w' + u'_1 w'', u'_1 u''_1, u''_1 u'_2 + u'_1 u''_2 + z' w'' - z'' w'). \end{aligned}$$

The Clifford actions It remains to describe the corresponding Clifford actions. As is standard, in the case of $\mathcal{Cl}(V_1, g_1)$, it suffices to consider the action of elements of form $(x, 0, z, 0, 0, 0)$ and $(x, 0, 0, w, 0, 0)$ on elements of form $(x, 0, z, 0, 0, 0)$, $(x, 0, 0, w, 0, 0)$, $(x, 0, 0, 0, u_1, 0)$, and $(x, 0, 0, 0, 0, u_2)$.

For these elements the multiplication is determined as follows

$$\left\{ \begin{array}{l} c_1(x, 0, z, 0, 0, 0)(x, 0, z', 0, 0, 0) = (x, 0, 0, 0, -f_1(x)z^2, 0) \\ c_1(x, 0, z, 0, 0, 0)(x, 0, 0, w, 0, 0) = (x, 0, 0, 0, 0, zw) \\ c_1(x, 0, z, 0, 0, 0)(x, 0, 0, 0, u_1, 0) = (x, 0, u_1z, 0, 0, 0) \\ c_1(x, 0, z, 0, 0, 0)(x, 0, 0, 0, 0, u_2) = (x, 0, 0, -u_2f_1(x)z, 0, 0) \\ c_1(x, 0, 0, w, 0, 0)(x, 0, z, 0, 0, 0) = (x, 0, 0, 0, 0, -zw) \\ c_1(x, 0, 0, w, 0, 0)(x, 0, 0, w', 0, 0) = (x, 0, 0, 0, -f_1(x)ww', 0) \\ c_1(x, 0, 0, w, 0, 0)(x, 0, 0, 0, u_1, 0) = (x, 0, 0, u_1w, 0, 0) \\ c_1(x, 0, 0, w, 0, 0)(x, 0, 0, 0, 0, u_2) = (x, 0, 0, 0, 0, 0). \end{array} \right.$$

In the case of $\mathcal{Cl}(V_2, g_2)$, it suffices to consider the action of $(0, y, z, 0, 0, 0)$ on elements of form $(0, y, z, 0, 0, 0)$ and $(0, y, 0, 0, u_1, 0)$, and we have

$$\left\{ \begin{array}{l} c_2(0, y, z, 0, 0, 0)(0, y, z', 0, 0, 0) = (0, y, 0, 0, -f_2(y)zz', 0) \\ c_2(0, y, z, 0, 0, 0)(0, y, 0, 0, u_1, 0) = (0, y, u_1z, 0, 0, 0) \end{array} \right.$$

Finally, the Clifford action on both $\bigwedge_*(V_1 \cup_{\tilde{f}} V_2)$ and $\bigwedge_*(V_1) \cup_{\tilde{f} \wedge_*} \bigwedge_*(V_2)$ is obtained by concatenating the two lists; the difference between the two pseudo-bundles is not seen on the level of defining the action, but rather in how we determine the two sets of points (as already been indicated above), underlying the commutativity between the gluing and the exterior product.

References

- [1] K.T. CHEN, *Iterated path integrals*, Bull. Amer. Math. Soc. (5) **1977**, pp. 831-879.
- [2] J.D. CHRISTENSEN – G. SINNAMON – E. WU, *The D-topology for diffeological spaces*, Pacific J. of Mathematics (1) **272** (2014), pp. 87-110.
- [3] J.D. CHRISTENSEN – E. WU, *Tangent spaces and tangent bundles for diffeological spaces*, arXiv:1411.5425v1.
- [4] P. IGLESIAS-ZEMMOUR, *Fibrations difféologiques et homotopie*, Thèse de doctorat d'État, Université de Provence, Marseille, 1985.
- [5] P. IGLESIAS-ZEMMOUR, *Diffeology*, Mathematical Surveys and Monographs, 185, AMS, Providence, 2013.
- [6] Y. KARSHON – J. WATTS, *Basic forms and orbit spaces: a diffeological approach*, SIGMA (2016).
- [7] E. PERVOVA, *Multilinear algebra in the context of diffeology*, arXiv:1504.08186v2.
- [8] E. PERVOVA, *Diffeological Clifford algebras and pseudo-bundles of Clifford modules*, arXiv:1505.06894v2.
- [9] E. PERVOVA, *On the notion of scalar product for finite-dimensional diffeological vector spaces*, arXiv:1507.03787v1.
- [10] E. PERVOVA, *Diffeological vector pseudo-bundles*, Topol. Appl. **202** (2016), pp. 269-300.
- [11] E. PERVOVA, *Diffeologica gluing of vector pseudo-bundles and pseudo-metrics on them*, Topol. Appl. (2017), <http://dx.doi.org/10.1016/j.topol.2017.02.002>
- [12] E. PERVOVA, *Diffeological Dirac operators and diffeological gluing*, arXiv:1701.06785v1.
- [13] I. SATAKE, *The Gauss-Bonnet theorem for V-manifolds*, J. Math. Soc. Japan (4) **9** (1957), pp. 464-492.

- [14] J.M. SOURIAU, *Groups différentiels*, Differential geometrical methods in mathematical physics (Proc. Conf., Aix-en-Provence/Salamanca, 1979), Lecture Notes in Mathematics, 836, Springer, (1980), pp. 91-128.
- [15] J.M. SOURIAU, *Groups différentiels de physique mathématique*, South Rhone seminar on geometry, II (Lyon, 1984), Astérisque 1985, Numéro Hors Série, pp. 341-399.
- [16] A. STACEY, *Comparative smoothology*, Theory Appl. Categ., 25(4) (2011), pp. 64-117.
- [17] M. VINCENT, *Diffeological differential geometry*, Master Thesis, University of Copenhagen, 2008.
- [18] J. WATTS, *Diffeologies, Differential Spaces, and Symplectic Geometry*, PhD Thesis, 2012, University of Toronto, Canada.
- [19] E. WU, *Homological algebra for diffeological vector spaces*, Homology, Homotopy & Applications (1) **17** (2015), pp. 339-376.

University of Pisa
Department of Mathematics
Via F. Buonarroti 1C
56127 PISA – Italy

ekaterina.pervova@unipi.it