

On maximal hypersurfaces in Lorentz manifolds admitting a parallel lightlike vector field

José A. S. Pelegrín^{a*}, Alfonso Romero^{a*} and Rafael M. Rubio^{b*}

^a Departamento de Geometría y Topología,
Universidad de Granada, 18071 Granada, Spain
E-mail: jpelegrin@ugr.es, aromero@ugr.es

^b Departamento de Matemáticas, Campus de Rabanales,
Universidad de Córdoba, 14071 Córdoba, Spain
E-mail: rmrubio@uco.es

Abstract

We study constant mean curvature spacelike hypersurfaces and in particular maximal hypersurfaces immersed in pp-wave spacetimes satisfying the timelike convergence condition. We prove the non-existence of compact spacelike hypersurfaces whose constant mean curvature is non-zero and also that every compact maximal hypersurface is totally geodesic. Moreover, we give an extension of the classical Calabi-Bernstein theorem to this class of pp-wave spacetimes.

PACS number: 02.40.Ma, 02.40.Vh.

2010 MSC: 53C50, 53C42, 53C80.

Keywords: Lorentzian geometry, lightlike parallel vector fields, Calabi-Bernstein theorem.

1 Introduction

The pp-wave type spacetimes are an important family of spacetimes. Considering such a spacetime as an exact solution of Einstein's field equation, it can model radiation (electromagnetic or gravitational) moving at the speed of light. The recent interest on pp-wave type spacetime can be explained, on the one hand, by its classical geometrical properties, on the other by its applications to string theory as well as the possibilities of direct detection of gravitational waves. Historically, the study of gravitational waves goes back to Einstein ([14]) but the standard exact model was already introduced by Brinkmann in order to determine Einstein spaces which can be improperly mapped conformally on some Einstein one (see [4]). The original definition by Brinkmann says that a pp-wave spacetime is any spacetime whose metric tensor can be described, with respect to

suitable coordinates, in the form $ds^2 = H(u, x, y) du^2 + 2 du dv + dx^2 + dy^2$, where H is any smooth function. Nowadays, pp-wave means any spacetime which admits a parallel global light-like vector field [25, 45, p. 383]. Moreover, when the spacetime is Ricci-flat (vacuum solutions) it is called gravitational plane wave. The interest in better understanding the geometry of pp-waves and their potential applications to string theory justifies the study of these spacetimes from a wider perspective.

On the other hand, spacelike hypersurfaces in $(n \geq 3)$ -dimensional Lorentzian spacetimes are geometrical objects of great physical and mathematical interest. Roughly speaking, each of them represents the physical space in a given instant of a time function. In Electromagnetism, a spacelike hypersurface is an initial data set that univocally determines the future of the electromagnetic field which satisfies the Maxwell equations [23, Thm. 3.11.1] and analogously, for the simple matter equations [23, Thm. 3.11.2]. In Causality Theory, the mere existence of a particular spacelike hypersurface implies that the spacetime obeys a certain causal property. Let us remark that the completeness of a spacelike hypersurface is required whenever we study its global properties, and also, from a physical viewpoint, completeness implies that the whole physical space is taken into consideration. To know more details about the relevance of constant mean curvature spacelike hypersurfaces in General Relativity see [17].

From a mathematical point of view, the interest of spacelike hypersurfaces is motivated by their nice Bernstein-type properties. In fact, maximal spacelike hypersurfaces in an $(n \geq 3)$ -dimensional spacetime are critical points of the area functional. In the history of research on maximal hypersurfaces, a striking fact was the discovery of new nonlinear elliptic problems. In fact, the function defining a maximal graph in the $(n + 1)$ -dimensional Lorentz-Minkowski spacetime \mathbb{L}^{n+1} satisfies an elliptic second order PDE similar to the equation of minimal graphs in Euclidean space \mathbb{R}^{n+1} , but with a new and surprising behavior for its entire solutions: The only entire solutions to the maximal hypersurface equation in \mathbb{L}^{n+1} are the affine functions defining spacelike hyperplanes. This result was previously proved by Calabi [6] for $n \leq 4$ and later extended for any n in the seminal paper by S.Y. Cheng and S.T. Yau [9] and it is called Calabi-Bernstein theorem. Recall that the Bernstein theorem for minimal graphs in the Euclidean space \mathbb{R}^{n+1} , holds only for $n \leq 7$, [24]. An important goal in [9] was the introduction of a new tool, the so-called Omori-Yau generalized maximum principle [18], [26]. By means of the use of this technique, many uniqueness results were obtained. For instance, S. Nishikawa [19] proved that a complete maximal hypersurface in a locally symmetric Lorentzian manifold whose Ricci tensor satisfies a natural assumption on timelike tangent vectors must be totally geodesic.

The special case of maximal surfaces in 3-dimensional spacetimes is of mathematical interest and, sometimes, a source of inspiration for higher dimensions. So, for the 3-dimensional Lorentz-Minkowski spacetime several new proofs of the so called classical Calabi-Bernstein theorem have been given by different authors (see [20], [3], [21], [2]). Moreover, several extensions of the classical results in different direction have been proved recently (see [1], [13], [5]).

In this work, we deal with complete maximal hypersurfaces in Lorentz manifolds admitting a parallel lightlike vector field satisfying the timelike convergence condition (and in particular for Ricci-flat case). We show (see Theorem 1) that in these Lorentzian manifolds (in fact, spacetimes)

there are no compact constant mean curvature spacelike hypersurfaces except for the maximal case. Moreover, we prove that every compact maximal hypersurface must be totally geodesic. On the other hand, in section 4 we give a new extension of Calabi-Bernstein theorem for these Lorentzian manifolds.

2 Preliminaries

The notion of symmetry is basic in Physics. In General Relativity, symmetry is usually based on the assumption of the existence of a one-parameter group of transformations generated by a Killing or, more generally, a conformal vector field. In fact, an usual simplification for the search of exact solutions to the Einstein equation is to assume a priori the existence of such an infinitesimal symmetry (see [10], [12] for instance). A complete general approach to symmetries can be found in [27] (see also [11] and references therein).

A vector field V on a Lorentzian manifold (M, g) is called conformal if $\mathcal{L}_V g = 2\lambda g$ for a smooth function λ , where \mathcal{L} denotes the Lie derivative on M . It is easy to see that if V is locally a gradient field (closed), then this condition is equivalent to

$$\nabla_X V = \lambda X, \quad \forall X \in \mathfrak{X}(M) \quad (1)$$

Vector fields satisfying (1) are called closed conformal vector fields. If $V \in \mathfrak{X}(M)$ is lightlike on (M, g) , then from (1) we get

$$Xg(V, V) = 2g(\nabla_X V, V) = 2\lambda g(X, V) = 0$$

for every $X \in \mathfrak{X}(M)$. Thus, V is parallel, since $\lambda V = 0$ and then λ must be identically 0.

Note that if a Lorentzian manifold (M, g) admits a parallel lightlike vector field then it is time-orientable and if we choose a time-orientation and (M, g) is connected then it is a spacetime.

On the other hand, recall that a Lorentzian manifold obeys the timelike convergence condition (TCC) if its Ricci tensor $\overline{\text{Ric}}$ satisfies

$$\overline{\text{Ric}}(Z, Z) \geq 0,$$

for all timelike vector Z . It is normally argued that TCC is the mathematical way to express that gravity, on average, attracts.

3 Principal result

Let $(\overline{M}, \langle, \rangle)$ be a Lorentzian manifold of dimension $n+1$ and let $x : M \rightarrow \overline{M}$ be an isometrically immersed spacelike hypersurface. Let us suppose the existence of a parallel lightlike vector field ξ on \overline{M} , i.e., $\langle \xi, \xi \rangle = 0$, $\xi \neq 0$ such that $\overline{\nabla} \xi = 0$, where $\overline{\nabla}$ denotes the Levi-Civita covariant differential in $(\overline{M}, \langle, \rangle)$. We can decompose ξ into its tangent and normal part, having $\xi = \xi^T +$

$\xi^\perp = \xi^T - \langle N, \xi \rangle N$, being N the timelike unitary normal vector field to M . It is clear that ξ^\perp never vanishes. Observe that the existence of a vector field which never vanishes can give some topological obstructions when the hypersurface M is compact.

Remark 1. Consider the Lorentzian manifold $(\overline{M}, \langle \cdot, \cdot \rangle)$ where \overline{M} is given by the product $\Sigma \times \mathbb{S}^1 \times \mathbb{R}$, endowed with the Lorentzian metric $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_\Sigma + 2d\alpha dv + H(x, \alpha)d\alpha^2$, where $x \in \Sigma$, being $(\Sigma, \langle \cdot, \cdot \rangle_\Sigma)$ a compact Riemannian manifold, \mathbb{S}^1 the unitary sphere, α an angular coordinate on \mathbb{S}^1 and $v \in \mathbb{R}$. Now, it is clear that the vector field $\frac{\partial}{\partial v}$ on \overline{M} is a parallel lightlike vector field and \overline{M} admits a compact spacelike hypersurface, as long as H is positive.

Let us now consider the function $\eta = \langle N, \xi \rangle$ and choose N such that $\langle N, \xi \rangle > 0$. It is not difficult to see that

$$\nabla \eta = -A\xi^T, \quad (2)$$

where ∇ denotes the gradient operator on $(M, \langle \cdot, \cdot \rangle)$ and A the shape operator of the isometric immersion associated with N .

Take a local orthonormal reference frame $\{E_1, \dots, E_n\}$ for $(M, \langle \cdot, \cdot \rangle)$. Since ξ is parallel, taking the tangent component of $\overline{\nabla}_{E_i} \xi$, we obtain

$$\nabla_{E_i} \xi^T = -\langle N, \xi \rangle A E_i. \quad (3)$$

We compute now $\text{div}(A\xi^T)$,

$$\text{div}(A\xi^T) = \sum_{i=1}^n \langle \nabla_{E_i} (A\xi^T), E_i \rangle = \sum_{i=1}^n \langle (\nabla_{E_i} A)\xi^T, E_i \rangle + \sum_{i=1}^n \langle A(\nabla_{E_i} \xi^T), E_i \rangle.$$

Using (3), we have

$$\begin{aligned} \text{div}(A\xi^T) &= \sum_{i=1}^n \langle (\nabla_{E_i} A)\xi^T, E_i \rangle + \sum_{i=1}^n \langle A(-\langle N, \xi \rangle A E_i), E_i \rangle = \\ &= \sum_{i=1}^n \langle (\nabla_{E_i} A)\xi^T, E_i \rangle - \langle N, \xi \rangle \sum_{i=1}^n \langle A^2 E_i, E_i \rangle = \sum_{i=1}^n \langle (\nabla_{E_i} A)\xi^T, E_i \rangle - \langle N, \xi \rangle \text{trace}(A^2). \end{aligned}$$

The Codazzi equation allows us to obtain,

$$\begin{aligned} \text{div}(A\xi^T) &= \sum_{i=1}^n \langle (\nabla_{\xi^T} A)E_i, E_i \rangle + \sum_{i=1}^n \langle \overline{R}(\xi^T, E_i)N, E_i \rangle - \langle N, \xi \rangle \text{trace}(A^2) = \\ &= \sum_{i=1}^n \langle (\nabla_{\xi^T} A)E_i, E_i \rangle - \overline{\text{Ric}}(\xi^T, N) - \langle N, \xi \rangle \text{trace}(A^2), \end{aligned}$$

Where \overline{R} and $\overline{\text{Ric}}$ denote the curvature and Ricci tensors of \overline{M} respectively. Finally, taking into account that tensor derivations commute with contractions we get

$$\text{div}(A\xi^T) = -n\langle \nabla \mathcal{H}, \xi \rangle - \overline{\text{Ric}}(\xi^T, N) - \langle N, \xi \rangle \text{trace}(A^2), \quad (4)$$

where $\mathcal{H} = -\frac{1}{n}\text{trace}(A)$ is the mean curvature function of M . From (2) and (4), the Laplacian of the distinguished function η is given by

$$\Delta \eta = n\langle \nabla \mathcal{H}, \xi \rangle + \overline{\text{Ric}}(\xi^T, N) + \eta \text{trace}(A^2). \quad (5)$$

Moreover, since ξ is parallel, the Ricci tensor can be expressed as $\overline{\text{Ric}}(\xi^T, N) = \overline{\text{Ric}}(\xi, N) + \langle N, \xi \rangle \overline{\text{Ric}}(N, N) = \eta \overline{\text{Ric}}(N, N)$ and (5) leads to

$$\Delta \eta = n\langle \nabla \mathcal{H}, \xi \rangle + \eta\{\overline{\text{Ric}}(N, N) + \text{trace}(A^2)\}. \quad (6)$$

Theorem 1. *Let $(\overline{M}, \langle, \rangle)$ be an $(n+1)$ -dimensional ($n \geq 2$) Lorentzian manifold which admits a parallel global lightlike vector field and suppose that $\overline{\text{Ric}}(Z, Z) \geq 0$, for every timelike vector Z . Consider $x : M \rightarrow \overline{M}$ a compact isometrically immersed spacelike hypersurface. If M has constant mean curvature, then it must be totally geodesic. As a direct consequence, there is no compact constant mean curvature spacelike immersed hypersurface whose mean curvature is different from zero.*

Proof. From (6) we have that the function η is positive and subharmonic on a compact manifold. Thus, η is constant. Again, from (6) we obtain that $\text{trace}(A^2) = 0$. \square

4 A new extension of the Calabi-Bernstein theorem

In this section we consider a $(2+1)$ -dimensional Lorentzian manifold $(\overline{M}, \langle, \rangle)$, which admits a parallel global lightlike vector field. Several examples of such Lorentzian manifolds can be found in [7]. Moreover, the differentiable manifold $\mathbb{R}^3 = \{(u, v, x) : u, v, x \in \mathbb{R}\}$ can be endowed with the family of Lorentzian metrics given by

$$\langle, \rangle = H(u, x) du^2 + 2 du dv + dx^2. \quad (7)$$

When the function H is constant we have a realization of the Lorentz-Minkowski spacetime \mathbb{L}^3 . Nonetheless, taking into account that the Ricci tensor (see [15, Section 2.2]) of this family of metrics is given by

$$\overline{\text{Ric}} = -\frac{\partial^2 H}{\partial x^2} du \otimes du,$$

it is clear that this family admits spacetimes which are not isometric to \mathbb{R}^3 and admit a global parallel lightlike vector field. So, it is enough to consider a function H such that $\frac{\partial^2 H}{\partial x^2}(p) \neq 0$, for some point p .

Futhermore, we can give examples of closed maximal hypersurfaces in spacetimes of the previous family, which are not isometric to \mathbb{L}^3 . Indeed, let $F : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a smooth function and let $c \in \mathbb{R}$ be a regular value of F . It is well-known that the level set $\Sigma = F^{-1}(c)$ is a closed embedded hypersurface in \mathbb{R}^3 . On the other hand, this hypersurface is spacelike with the induced metric, if and only if the vector field $\overline{\nabla}F$ is timelike on Σ . Taking into account the Christoffel symbols of the metric (see [8, Section 2]), it is easy to obtain

$$\overline{\nabla}F = \frac{\partial F}{\partial v} \frac{\partial}{\partial u} + \left(\frac{\partial F}{\partial u} - H \frac{\partial F}{\partial v} \right) \frac{\partial}{\partial v} + \frac{\partial F}{\partial x} \frac{\partial}{\partial x}.$$

Therefore, Σ is spacelike if and only if the inequality

$$2 \frac{\partial F}{\partial u} \frac{\partial F}{\partial v} - H \left(\frac{\partial F}{\partial v} \right)^2 + \left(\frac{\partial F}{\partial x} \right)^2 < 0, \quad (8)$$

holds on Σ .

Now, if we denote by A the shape operator on Σ associated to the unitary normal vector field $\frac{1}{|\overline{\nabla}F|} \overline{\nabla}F$, we have that

$$\langle Aw_1, w_2 \rangle = - \frac{\overline{\text{Hess}}(F)(w_1, w_2)}{|\overline{\nabla}F|}, \quad (9)$$

where $\overline{\text{Hess}}(F)$ denotes the Hessian of the function F in $(\mathbb{R}^3, \langle \cdot, \cdot \rangle)$ and $w_1, w_2 \in T_p \Sigma$.

If we take $W_1, W_2 \in \mathfrak{X}(\mathbb{R}^3)$ and we put

$$W_1 = w_1^u \frac{\partial}{\partial u} + w_1^v \frac{\partial}{\partial v} + w_1^x \frac{\partial}{\partial x}, \quad W_2 = w_2^u \frac{\partial}{\partial u} + w_2^v \frac{\partial}{\partial v} + w_2^x \frac{\partial}{\partial x},$$

then

$$\overline{\text{Hess}}(F)(W_1, W_2) = \langle \overline{\nabla}_{W_1}(\overline{\nabla}F), W_2 \rangle \quad (10)$$

$$\begin{aligned} &= W_1 \left(\frac{\partial F}{\partial v} \right) (H w_2^u + w_2^v) + \frac{\partial F}{\partial v} \left(\frac{1}{2} w_1^u w_2^u \frac{\partial H}{\partial u} - \frac{1}{2} w_1^u w_2^x \frac{\partial H}{\partial x} + \frac{1}{2} w_1^x w_2^u \frac{\partial H}{\partial x} \right) \\ &\quad + W_1 \left(\frac{\partial F}{\partial u} - H \frac{\partial F}{\partial v} \right) w_2^u + W_1 \left(\frac{\partial F}{\partial x} \right) w_2^x + \frac{\partial F}{\partial x} \left(\frac{1}{2} w_1^u w_2^u \frac{\partial H}{\partial x} \right). \end{aligned}$$

Now, consider the spacetime $(\mathbb{R}^3, \langle \cdot, \cdot \rangle)$, whose metric is given in (7), taking H a positive function such that $\frac{\partial H}{\partial u} = 0$ and $\frac{\partial^2 H}{\partial x^2}(p) \neq 0$, for some point $p \in \mathbb{R}^3$. Let $F(u, v, x) = \lambda v$, $\lambda \in \mathbb{R}^+$ be. Taking into account (8) and (10), it is not difficult to see that the hypersurface $F^{-1}(c)$ is spacelike and in particular maximal, since from (9), it is clear that $\text{trace}(A) = 0$.

Consider an immersed maximal surface S in a $(2+1)$ -dimensional Lorentzian manifold $(\overline{M}, \langle, \rangle)$, which admits a parallel global lightlike vector field. Here, taking into account that $\mathcal{H} = 0$, we have from (6),

$$\Delta\eta = \eta\{\overline{\text{Ric}}(N, N) + \text{trace}(A^2)\}. \quad (11)$$

Moreover, in this case, using Cayley-Hamilton theorem we get

$$|\nabla\eta|^2 = \langle A^2\xi^T, \xi^T \rangle = \frac{1}{2}\text{trace}(A^2)|\xi^T|^2 = \frac{1}{2}\text{trace}(A^2)\eta^2. \quad (12)$$

Now, we can state,

Theorem 2. *Let $(\overline{M}, \langle, \rangle)$ be a $(2+1)$ -dimensional Lorentzian manifold $(\overline{M}, \langle, \rangle)$, which admits a parallel global lightlike vector field and suppose that $\overline{\text{Ric}}(Z, Z) \geq 0$, for all timelike vector Z . Let $x : S \rightarrow \overline{M}$ be a complete isometrically immersed maximal surface. Then S admits a positive function η which is constant if and only if S is totally geodesic.*

On the other hand, taking into account the Gauss curvature of the surface, we can enunciate,

Theorem 3. *Let $(\overline{M}, \langle, \rangle)$ be a $(2+1)$ -dimensional manifold which admits a parallel global lightlike vector field. Let $x : S \rightarrow \overline{M}$ be an isometrically immersed maximal surface. Then,*

- (i) *the Gaussian curvature of S satisfies $K \geq \overline{\text{Ric}}(N, N)$, being N the timelike unitary normal vector field to S . Moreover, equality holds if and only if S is totally geodesic.*
- (ii) *if the spacetime satisfies the TCC and the surface is complete, then S is parabolic.*

Proof. Since S is maximal, from the Gauss equation we obtain,

$$\text{Ric}(\xi^T, \xi^T) = \overline{\text{Ric}}(\xi^T, \xi^T) + \langle \overline{\mathbf{R}}(\xi^T, N)N, \xi^T \rangle + \langle A^2\xi^T, \xi^T \rangle. \quad (13)$$

Moreover, being $\dim S = 2$ we have

$$\text{Ric}(\xi^T, \xi^T) = K|\xi^T|^2 = K\eta^2 \quad (14)$$

and as $\overline{\nabla}\xi = 0$, then

$$\overline{\text{Ric}}(\xi^T, \xi^T) = \eta^2\overline{\text{Ric}}(N, N), \quad (15)$$

and

$$\langle \overline{\mathbf{R}}(\xi^T, N)N, \xi^T \rangle = \langle \overline{\mathbf{R}}(\xi, N)N, \xi \rangle + 2\eta\langle \overline{\mathbf{R}}(\xi, N)N, N \rangle + \eta^2\langle \overline{\mathbf{R}}(N, N)N, N \rangle = 0. \quad (16)$$

Therefore, substituting (14), (15), (16) and (12) in (13) we have

$$K = \overline{\text{Ric}}(N, N) + \frac{1}{2}\text{trace}(A^2) \quad (17)$$

and as a direct consequence (i) holds.

Finally, the TCC guarantees $K \geq 0$ and by a classical result of Ahlfors and Blanc-Fiala-Huber (see for instance [16]) a complete Riemannian surface with non-negative Gaussian curvature is parabolic. \square

We can give the following rigidity's result,

Theorem 4. *Let $(\overline{M}, \langle, \rangle)$ be a $(2+1)$ -dimensional Lorentzian manifold $(\overline{M}, \langle, \rangle)$, which admits a parallel global lightlike vector field and suppose that $\overline{\text{Ric}}(Z, Z) \geq 0$, for all timelike vector Z . Let $x : S \rightarrow \overline{M}$ be a complete isometrically immersed maximal surface. Then S is totally geodesic.*

Proof. Making use of (11) and (12),

$$\Delta \left(\frac{1}{\eta} \right) = -\frac{1}{\eta^2} \Delta \eta + \frac{2}{\eta^3} |\nabla \eta|^2 = -\frac{1}{\eta} \{ \overline{\text{Ric}}(N, N) + \text{trace}(A^2) \} + \frac{1}{\eta} \text{trace}(A^2)$$

Thus,

$$\Delta \left(\frac{1}{\eta} \right) = -\frac{1}{\eta} \overline{\text{Ric}}(N, N). \quad (18)$$

Since S is parabolic due to Theorem 3, it is enough to observe that the function η must be constant, and so, Theorem 2 applies. \square

As a corollary of the previous theorem we obtain the classical Calabi-Bernstein theorem.

Corollary 1. *(Classical Calabi-Bernstein theorem) The only complete maximal surfaces in the Lorentz-Minkowski spacetime \mathbb{L}^3 are the spacelike affine planes.*

Acknowledgments

The authors are grateful to the two referees for their deep readings and the suggestions made toward the improvement of this article. The authors are partially supported by the Spanish MICINN Grant with FEDER funds MTM2013-47828-C2-1-P and by the Junta de Andalucía Regional Grant P09-FQM-4496.

References

- [1] A. Albuja and L.J. Alías, Parabolicity of maximal surfaces in Lorentzian product spaces, *Math. Z.* **267** (2011), 453-464.
- [2] J.A. Aledo, A. Romero and R.M. Rubio, The classical Calabi-Bernstein Theorem revisited, *J. Math. Anal. Appl.* **431** (2015), 1172-1177.

- [3] L.J. Alías and B. Palmer, On the Gaussian curvature of maximal surfaces and the Calabi-Bernstein theorem, *Bull. London Math. Soc.* **33** (2001), 454–458.
- [4] H. Brinkmann, Einstein spaces which are mapped conformally on each other, *Math. Ann.*, **94** (1925), 119–145.
- [5] M. Caballero, A. Romero and R.M. Rubio, Uniqueness of maximal surfaces in Generalized Robertson-Walker spacetimes and Calabi-Bernstein type problems, *J. Geom. Phys.*, **60** (2010), 394–402.
- [6] E. Calabi, Examples of Bernstein problems for some nonlinear equations, *Proc. Symp. Pure Math.*, **15** (1970), 223–230.
- [7] M. Chaichi, E. Garcia-Río and M.E. Vázquez Abal Three-dimensional Lorentz manifolds admitting a parallel null vector field, *J. Phys. A*, **3** (2005) 841-850
- [8] A.M. Candela, J.L. Flores and M. Sanchez, On General Plane Fronted Waves. Geodesics, *Gen. Rel. Grav.*, **35** (2003), 631–649.
- [9] S.Y. Cheng and S.T. Yau, Maximal space-like hypersurfaces in the Lorentz-Minkowski spaces, *Ann. Math.*, **104** (1976), 407–419.
- [10] V. Daftardar and N. Dadhich. Gradient conformal Killing vectors and exact solutions. *Gen. Relativity Gravitation*, **26** (1994), 859-868.
- [11] K. Duggal and R. Sharma. Symmetries of spacetimes and Riemannian manifolds, volume 487. Springer Science and Business Media, 2013.
- [12] D. Eardley, J. Isenberg, J. Marsden, and V. Moncrief. Homothetic and conformal symmetries of solutions to Einsteins equations, *Comm. Math. Phys.*, **106** (1996), 137-158.
- [13] F.J.M. Estudillo and A. Romero, Generalized maximal surfaces in Lorentz-Minkowski space \mathbb{L}^3 , *Math. Proc. Camb. Phil. Soc.* **111** (1992), 515–524.
- [14] A. Einstein and N. Rosen, On gravitational waves, *J. Franklin Inst.*, **223** (1937), 4354.
- [15] J.L Flores and M. Sánchez, On the geometry of pp-wave type spacetimes. Analytical and numerical approaches to mathematical relativity, 7998, Lecture Notes in Phys., 692, Springer, Berlin, 2006.
- [16] J.L. Kazdan, Parabolicity and the Liouville property on complete Riemannian manifolds, *Aspects of Math.*, **E10**, Ed. A.J. Tromba, Friedr. Vieweg and Sohn, Bonn (1987), 153–166.
- [17] J.E. Marsden and F.J. Tipler, Maximal hypersurfaces and foliations of constant mean curvature in General Relativity, *Phys. Rep.* **66** (1980), 109–139.

- [18] H. Omori, Isometric immersions of Riemannian manifolds, *J. Math. Soc. Japan*, **19** (1967), 205–214.
- [19] S. Nishikawa, On maximal spacelike hypersurfaces in a Lorentzian manifold, *Nagoya Math. J.*, **95** (1984), 117–124.
- [20] A. Romero, Simple proof of Calabi-Bernstein’s theorem, *Proc. Amer. Math. Soc.* **124** (1996), 1315–1317.
- [21] A. Romero and R.M. Rubio, New proof of the Calabi-Bernstein theorem, *Geom. Dedicata* **147** (2010), 173–176.
- [22] R. M. Rubio and J.J. Salamanca, Maximal surface equation on a Riemannian 2-manifold with finite total curvature, *J. Geom. Phys.*, **92** (2015), 140-146.
- [23] R.K. Sachs and H. Wu, *General Relativity for Mathematicians*, Grad. Texts Math., **48** Springer-Verlag, New York, 1977.
- [24] R. Schoen, L. Simon and S.T. Yau, Curvature estimates for minimal hypersurfaces, *Acta Math.*, **134** (1975), 275–288.
- [25] H. Stephani, D. Kramer, M. MacCallum, C. Hoenselaers and E. Herlt, Exact Solutions of Einsteins Field Equations *Cambridge University Press*, Cambridge 2003.
- [26] S.T. Yau, Harmonic functions on complete Riemannian manifolds, *Comm. Pure Appl. Math.*, **28** (1975), 201–228.
- [27] E. Zafiris. Irreducible decomposition of Einsteins equations in spacetimes with symmetries, *Ann. Physics*, **263** (1998), 155-178.