

Bounds on Multipartite Concurrence and Tangle

Jing Wang^{1,2}, Ming Li¹, Hongfang Li¹, Shao-Ming Fei^{2,3} and Xianqing Li-Jost³

¹ College of the Science, China University of Petroleum, 266580 Qingdao

² Department of Mathematics, Capital Normal University, 100037 Beijing

³ Max-Planck-Institute for Mathematics in the Sciences, 04103 Leipzig

Abstract

We present an analytical lower bound of multipartite concurrence based on the generalized Bloch representations of density matrices. It is shown that the lower bound can be used as an effective entanglement witness of genuine multipartite entanglement. Tight lower and upper bounds for multipartite tangles are also derived. Since the lower bounds depend on just part of the correlation tensors, the result is experimentally feasible.

PACS numbers: 03.67.-a, 02.20.Hj, 03.65.-w

I. INTRODUCTION

Quantum entanglement, as the remarkable nonlocal feature of quantum mechanics, is recognized as a valuable resource in the rapidly expanding field of quantum information science, with various applications [1, 2] such as quantum computation, quantum teleportation, dense coding, quantum cryptographic schemes, quantum radar, entanglement swapping and remote states preparation.

The mixed state $\rho \in \mathcal{H} = \mathcal{H}_1 \otimes \cdots \otimes \mathcal{H}_N$ is said to be a fully separable state, if there exist $|\phi_i^j\rangle \in \mathcal{H}_j, j = 1, \dots, N, p_i > 0, \sum_i p_i = 1$, such that

$$\rho = \sum_i p_i |\phi_i^1\rangle\langle\phi_i^1| \otimes \cdots \otimes |\phi_i^N\rangle\langle\phi_i^N|, \quad (1)$$

otherwise ρ is said to be an entangled state. States that are not biseparable with respect to any partitions are said to be genuinely multipartite entangled. Genuinely multipartite entanglement is a kind of important type of entanglement, which offers significant advantage in quantum information processing tasks [3]. In particular, it is the basic ingredient in

measurement-based quantum computation [4], and is beneficial in various quantum communication protocols [5], including secret sharing [6] (cf.[7]). Despite its importance, characterization and detection of this kind of resource turn out to be rather hard and only a few results have been proposed [8–11].

Quantifying quantum entanglement is a basic and longer standing problem in quantum information theory. A measure of quantum entanglement can be used to detect and classify entanglement of quantum states. In this paper, we use the multipartite concurrence [12] to investigate the multipartite entanglement. Let \mathcal{H}_i , $i = 1, 2, \dots, N$, be d -dimensional vector spaces. The concurrence of an N -partite pure state $|\psi\rangle \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \dots \otimes \mathcal{H}_N$ is defined by

$$\mathcal{C}_N(|\psi\rangle\langle\psi|) = 2^{1-\frac{N}{2}} \sqrt{(2^N - 2) - \sum_{\alpha} \text{Tr}\{\rho_{\alpha}^2\}}, \quad (2)$$

where α labels all the different reduced density matrices. If we list all the $2^N - 2$ reduced matrices in the following way: $\{\rho_1, \rho_2, \dots, \rho_N, \rho_{12}, \rho_{13}, \dots, \rho_{1N}, \rho_{23}, \dots, \rho_{12\dots N-1}, \dots, \rho_{23\dots N}\}$, (2) can be reexpressed as

$$\mathcal{C}_N(|\psi\rangle\langle\psi|) = 2^{1-\frac{N}{2}} \sqrt{(2^N - 2) - 2 \sum_{k=1}^{2^{N-1}-1} \text{Tr}\{\rho_k^2\}}. \quad (3)$$

For a mixed multipartite quantum state, $\rho \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \dots \otimes \mathcal{H}_N$, the corresponding concurrence of (2) is given by the convex roof:

$$\mathcal{C}_N(\rho) = \min_{\{p_i, |\psi_i\rangle\}} \sum_i p_i \mathcal{C}_N(|\psi_i\rangle\langle\psi_i|), \quad (4)$$

where the minimum is taken over all the ensemble decomposition of $\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i|$.

The correlation tensors of the generalized Bloch representation of a quantum state play significant roles in quantum information theory. In [13–16], separable conditions for both bi- and multi-partite quantum states are introduced by studying the norm of the correlation tensors. In [17, 18], the authors present a multipartite entanglement measure for N -qubit and N -qudit pure states, using the norm of the correlation tensors. In [9], the authors have introduced a general framework for detecting genuine multipartite entanglement and non full separability in multipartite quantum systems of arbitrary dimensions based also on the correlation tensors. In [19], we have found that the norms of the correlation tensors are closely related to the maximal violation of a kind of multipartite Bell inequalities.

In the following, we first reform the concurrence for multipartite pure states in terms of the norms of the correlation tensors. The correlation tensors are then used to derive a lower bound of concurrence for mixed multipartite quantum states. The lower bound also provides a fully separable condition for multipartite quantum states. We further show that genuine multipartite entanglement can be detected by the bound. We also investigate the multipartite tangle. Tight lower and upper bounds are derived.

II. LOWER BOUND OF MULTIPARTITE CONCURRENCE

We first consider the concurrence of multipartite pure states $|\psi\rangle \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_N$ in terms of the generalized Bloch representation of $|\psi\rangle\langle\psi|$. Let $\{\lambda_{\alpha_k}\}$ be the $SU(d)$ generators. The generalized Bloch representation for any quantum states $\rho \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_N$ is given by [15],

$$\rho = \frac{1}{d^N} \left(\otimes_{j=1}^N I_d + \cdots + \sum_{M=1}^N \sum_{\{\mu_1 \mu_2 \cdots \mu_M\}} \sum_{\alpha_1 \alpha_2 \cdots \alpha_M} \mathcal{T}_{\alpha_1 \alpha_2 \cdots \alpha_M}^{\{\mu_1 \mu_2 \cdots \mu_M\}} \lambda_{\alpha_1}^{\{\mu_1\}} \lambda_{\alpha_2}^{\{\mu_2\}} \cdots \lambda_{\alpha_M}^{\{\mu_M\}} \right), \quad (5)$$

where $\{\mu_1 \mu_2 \cdots \mu_M\}$ is a subset of $\{1, 2, \dots, N\}$, $\lambda_{\alpha_k}^{\{\mu_k\}} = I_d \otimes I_d \otimes \cdots \otimes \lambda_{\alpha_k} \otimes I_d \otimes \cdots \otimes I_d$ with λ_{α_k} appearing at the μ_k th position and

$$\mathcal{T}_{\alpha_1 \alpha_2 \cdots \alpha_M}^{\{\mu_1 \mu_2 \cdots \mu_M\}} = \frac{d^M}{2^M} \text{Tr}[\rho \lambda_{\alpha_1}^{\{\mu_1\}} \lambda_{\alpha_2}^{\{\mu_2\}} \cdots \lambda_{\alpha_M}^{\{\mu_M\}}], \quad (6)$$

which can be viewed as the entries of the correlation tensors $\mathcal{T}^{\{\mu_1 \mu_2 \cdots \mu_M\}}$.

After computation(detailed processes can be found in the supplementary material), we obtain an alternative representation about the concurrence,

$$\begin{aligned} & 2^{N-2} \mathcal{C}_N^2(|\psi\rangle\langle\psi|) \\ &= [2^N - \frac{(d+1)^N}{d^N} - \frac{1}{d^N} (d^N - 1)(d+1)^{N-1}] \\ & \quad + \sum_{l=2}^N \frac{2^l}{d^{N+l}} [(d+1)^{N-1} - (d+1)^{N-l}] \sum_{k_1 \cdots k_l \subset \{1, 2, \dots, N\}} \|\mathcal{T}^{k_1 \cdots k_l}\|^2. \end{aligned} \quad (7)$$

By noticing that a quantum state ρ is fully separable if and only if $\mathcal{C}_N(\rho) = 0$, one derives a sufficient and necessary condition for the fully separability of multipartite pure states with formula (7). In particular, as the tensors $\mathcal{T}^{\{\mu_1 \mu_2 \cdots \mu_M\}}$ in (6) are mean values of the observables $\lambda_{\alpha_1}^{\{\mu_1\}} \lambda_{\alpha_2}^{\{\mu_2\}} \cdots \lambda_{\alpha_M}^{\{\mu_M\}}$, (7) also gives an experimental way to measure the concurrence of a pure multipartite state. Since we have eliminated the terms containing \mathcal{T}^{k_1} , the measurement can be only operated on the norms of $\mathcal{T}^{\{\mu_1 \mu_2 \cdots \mu_M\}}$ with $M \geq 2$. From (7) we can now derive the lower bound for multipartite concurrence of any mixed states ρ .

Theorem 1: For any mixed quantum state $\rho \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_N$, we have

$$\begin{aligned} \mathcal{C}_N(\rho) &\geq -2^{1-N/2} [-2^N + \frac{(d+1)^N}{d^N} + \frac{1}{d^N} (d^N - 1)(d+1)^{N-1}]^{\frac{1}{2}} \\ & \quad + 2^{1-N/2} \left\{ \sum_{l=2}^N \frac{2^l}{d^{N+l}} [(d+1)^{N-1} - (d+1)^{N-l}] \sum_{k_1 \cdots k_l \subset \{1, 2, \dots, N\}} \|\mathcal{T}^{k_1 \cdots k_l}\|^2 \right\}^{\frac{1}{2}}. \end{aligned} \quad (8)$$

Proof: For simplicity we denote $C = -2^N + \frac{(d+1)^N}{d^N} + \frac{1}{d^N} (d^N - 1)(d+1)^{N-1}$, and C_α the coefficient of $\|\mathcal{T}^\alpha\|^2$ in (7) for $\alpha \in \{k_1 k_2, k_1 k_2 k_3, \dots, 1 \cdots N\}$, which are nonnegative numbers depending only on N and d .

Assume that $\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i|$ is the optimal decomposition such that (4) attains the minimum. We have that

$$\begin{aligned}
\mathcal{C}_N(\rho) &= \sum_i p_i \mathcal{C}_N(|\psi_i\rangle) = 2^{1-N/2} \sum_i p_i \{-C + \sum_\alpha C_\alpha \|\mathcal{T}_i^\alpha\|^2\}^{\frac{1}{2}} \\
&\geq 2^{1-N/2} \left[\sum_i p_i \sqrt{\sum_\alpha C_\alpha \|\mathcal{T}_i^\alpha\|^2} - \sqrt{C} \right] \\
&\geq 2^{1-N/2} \left[\sqrt{\sum_\alpha C_\alpha \left(\sum_i p_i \|\mathcal{T}_i^\alpha\|^2 \right)} - \sqrt{C} \right] \\
&\geq 2^{1-N/2} \left[\sqrt{\sum_\alpha C_\alpha \|\mathcal{T}^\alpha\|^2} - \sqrt{C} \right],
\end{aligned}$$

where we have used the inequalities $\sqrt{a-b} \geq \sqrt{a} - \sqrt{b}$ for $a \geq b \geq 0$ and $\sum_i \sqrt{\sum_j x_{ij}^2} \geq \sqrt{\sum_j (\sum_i x_{ij})^2}$ for real and nonnegative x_{ij} . \blacksquare

The lower bound (8) can be used to estimate the concurrence for multipartite quantum states with arbitrary dimension. It is also a kind of entanglement witness for fully separability. Moreover, this multipartite concurrence can be employed to detect the genuine multipartite entanglement. It has been shown that an N -partite quantum state $\rho \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_N$ is genuine multipartite entangled if [20]

$$\mathcal{C}_N(\rho) > \begin{cases} 2^{1-\frac{N}{2}} \sqrt{2^N - 4 + \frac{2}{d} - 2 \sum_{k=1}^{\frac{N-1}{2}} \frac{\binom{N}{k}}{d^k}}, & \text{for odd } N, \\ 2^{1-\frac{N}{2}} \sqrt{2^N - 4 + \frac{2}{d} - 2 \sum_{k=1}^{\frac{N}{2}-1} \frac{\binom{N}{k}}{d^k} - \frac{\binom{N}{\frac{N}{2}}}{d^{\frac{N}{2}}}}, & \text{for even } N, \end{cases} \quad (9)$$

where $\binom{N}{k} = N!/(k!(N-k)!)$.

Since the concurrence $\mathcal{C}_N(\rho)$ is difficult to compute, our lower bound can be employed to detect the genuine multipartite entanglement.

As an example, let us consider tripartite case. From (9) $\rho \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \mathcal{H}_3$ is genuinely multipartite entangled if $\mathcal{C}_3(\rho) > \sqrt{2 - \frac{2}{d}}$. For a three-qubit GHZ state mixed with noise, $\rho_{GHZ} = \frac{x}{8}I + (1-x)|GHZ\rangle\langle GHZ|$, where $|GHZ\rangle = \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle)$, we have $\mathcal{C}_3(\rho_{GHZ}) \geq \frac{1}{2}\sqrt{6 - 25x + \frac{25}{2}x^2}$ by Theorem 1. Therefore, the lower bound is valid to detect genuinely multipartite entangled for $x < 0.08349$. By $\frac{1}{2}\sqrt{6 - 25x + \frac{25}{2}x^2} > 0$, one can detect general entanglement(not fully separable) of ρ_{GHZ} for $x < 0.27889$. Note that general entanglement will be detected for $x < 0.64645$ in [15].

III. BOUNDS ON MULTIPARTITE TANGLE

We now consider the multipartite tangle that is tightly related with concurrence. By the squared I-concurrence for bipartite quantum systems [21], we introduce the multipartite squared I-concurrence. For a multipartite pure quantum state $|\psi\rangle \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_N$ the multipartite squared I-concurrence is defined by the square of the multi-concurrence,

$$\tau_N(|\psi\rangle\langle\psi|) = \mathcal{C}_N^2(|\psi\rangle\langle\psi|) = 2^{2-N}[(2^N - 2) - \sum_{\alpha} \text{Tr}\{\rho_{\alpha}^2\}], \quad (10)$$

where α labels all the different reduced density matrices. For a mixed multipartite quantum state, $\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i| \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_N$, the corresponding multipartite squared I-concurrence is then given by the convex roof:

$$\tau_N(\rho) = \min_{\{p_i, |\psi_i\rangle\}} \sum_i p_i \tau_N(|\psi_i\rangle\langle\psi_i|). \quad (11)$$

The multipartite squared I-concurrence defined above has the following properties: (i) $\tau_N(\rho) = 0$ if and only if ρ is fully separable; (ii) $\tau_N(\rho)$ is invariant under local unitary transformation of ρ ; (iii) $\tau_N(\rho) \geq \mathcal{C}_N^2(\rho)$. By property (i) above, a multipartite state is not separable if $\tau_N(\rho) > 0$. In the following, we present valid lower and upper bounds for $\tau_N(\rho)$.

Theorem 2: For any mixed quantum state $\rho \in \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_N$, we have

$$\begin{aligned} \tau_N(\rho) \geq & -2^{2-N}[-2^N + \frac{(d+1)^N}{d^N} + \frac{1}{d^N}(d^N - 1)(d+1)^{N-1}] \\ & + 2^{2-N} \left\{ \sum_{l=2}^N \frac{2^l}{d^{N+l}} [(d+1)^{N-1} - (d+1)^{N-l}] \sum_{k_1 \cdots k_l \subset \{1, 2, \dots, N\}} \|\mathcal{T}^{k_1 \cdots k_l}\|^2 \right\}; \end{aligned} \quad (12)$$

$$\tau_N(\rho) \leq 2^{2-N}(2^N - 2 - \sum_{\alpha} \text{Tr}\rho_{\alpha}^2). \quad (13)$$

Proof: We still take the simplified notions C and C_{α} used in the proof of Theorem 1. Assume that $\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i|$ is the optimal decomposition such that (11) attains the minimum. We have that

$$\begin{aligned} \tau_N(\rho) &= \sum_i p_i \mathcal{C}_N^2(|\psi_i\rangle) = 2^{2-N} \sum_i p_i \{-C + \sum_{\alpha} C_{\alpha} \|\mathcal{T}_i^{\alpha}\|^2\} \\ &= 2^{2-N} [\sum_{\alpha} C_{\alpha} (\sum_i p_i \|\mathcal{T}_i^{\alpha}\|)^2 - C] \\ &\geq 2^{2-N} [\sum_{\alpha} C_{\alpha} \|\mathcal{T}^{\alpha}\|^2 - C], \end{aligned}$$

where we have used the triangle inequality for the Hilbert-Schmidt norm.

On the other hand, by the definition of $\tau_N(\rho)$, we have

$$\begin{aligned}\tau_N(\rho) &\leq \sum_i p_i \tau_N(|\psi_i\rangle) = 2^{2-N} (2^N - 2 - \sum_{\alpha,i} p_i \text{Tr}(\rho_\alpha^i)^2) \\ &\leq 2^{2-N} [2^N - 2 - \sum_\alpha \text{Tr}(\sum_i p_i \rho_\alpha^i)^2] \\ &= 2^{2-N} (2^N - 2 - \sum_\alpha \text{Tr} \rho_\alpha^2),\end{aligned}$$

which gives the upper bound. ■

From the proof of Theorem 2, one has that for pure states the lower and upper bounds are exact. Thus the lower and upper bounds (12) for $\tau_N(\rho)$ are tight.

Remark: Our bounds are given by the norms of the correlation tensors. As the Hilbert-Schmidt norm is invariant under local unitary transformation, the bounds give experimentally feasible way in identify both non-separability and genuine multipartite entanglement. Further more, as has been discussed in [9, 15], partial knowledge of the correlation tensors may also allow us to detect entanglement and estimate the degree of entanglement.

IV. CONCLUSIONS AND DISCUSSIONS

It is a basic and fundamental question in quantum entanglement theory to compute the concurrence for multipartite quantum systems. Since the concurrence is defined by taking the optimization over all the ensemble decompositions of a mixed quantum states, it is formidable to derive an analytical formula. We have derived an analytical and experimentally feasible formula for multipartite concurrence of any multipartite pure quantum states by using generalized Bloch representation of density matrices. We have then obtained a lower bound of concurrence for any mixed multipartite quantum states. Genuine multipartite entanglement can be detected by using this bound. We have also investigated the multipartite tangle. Tight lower and upper bounds are obtained. Although the bounds are in general not tighter than that in [22], they are experimentally feasible as the elements in the correlation tensors are just the mean values of the hermitian $SU(d)$ generators. The approach used in this manuscript can also be implemented to investigate the k-separability of multipartite quantum systems. Future research on the construction of genuine multipartite entanglement criteria in terms of the lower bound of multipartite squared I-concurrence and the k norm would be also interesting. The correlation tensors, based on which the results are derived in this paper, would be used further to investigate nonlocality[19] or Measurement-Induced Non locality[23].

Acknowledgments This work is supported by the NSFC 11105226, 11275131; the Fundamental Research Funds for the Central Universities No. 15CX08011A, 15CX05062A,

27R1410016A, 16CX02049A; Qingdao applied basic research program No. 15-9-1-103-jch, and a project sponsored by SRF for ROCS, SEM.

- [1] Nielsen, M.A., and Chuang, I.L.: Quantum computation and quantum information. Cambridge: Cambridge University Press, (2000).
- [2] DiVincenzo, D.P.: Quantum computation. *Science* 270,255-261(1995).
- [3] Horodecki, R., Horodecki, P., Horodecki, M., and Horodecki, K.: Quantum entanglement. *Rev. Mod. Phys.* 81, 865-942(2009).
- [4] Briegel, H.J., Browne, D.E., Dür, W., Raussendorf, R. and Van den Nest, M.: Measurement-based quantum computation. *Nat. Phys.* 5, 19-26(2009).
- [5] Sen(De), A. and Sen, U.: Quantum advantage in communication networks. *Phys. News* 40, 17-32(2010).
- [6] Żukowski, M., Zeilinger, A., Horne, M.A., and Weinfurter, H.: Quest for GHZ states. *Acta Phys. Pol. A* 93, 187-195(1998); Hillery, M., Bužek, V., and Berthiaume, A.: Quantum secret sharing. *Phys. Rev. A* 59, 1829-1834 (1999); Demkowicz-Dobrzanski, R., Sen(De), A., Sen, U., and Lewenstein, M.: Entanglement enhances security in quantum communication. *Phys. Rev. A* 80,012311(2009); Gisin, N., Ribordy, G., Tittel, W., and Zbinden, H.: Quantum cryptography. *Rev. Mod. Phys.* 74, 145-195(2002).
- [7] Cleve, R., Gottesman, D., and Lo, H.K.: How to share a quantum secret. *Phys. Rev. Lett.* 83, 648 (1999); Karlsson, A., Koashi, M., and Imoto, N.: Quantum entanglement for secret sharing and secret splitting. *Phys. Rev. A* 59, 162(1999).
- [8] Huber, M., Mintert, F., Gabriel, A., and Hiesmayr, B.C.: Detection of high-dimensional genuine multipartite entanglement of mixed states. *Phys. Rev. Lett.* 104, 210501(2010).
- [9] de Vicente, J.I., Huber, M.: Multipartite entanglement detection from correlation tensors. *Phys. Rev. A*, 84, 062306(2011).
- [10] Eltschka, C. and Siewert J.: Entanglement of three-qubit Greenberger-Horne-Zeilinger-symmetric states. *Phys. Rev. Lett.* 108, 020502 (2012).
- [11] Jungnitsch, B., Moroder, T., and Gühne, O.: Taming multiparticle entanglement. *Phys. Rev. Lett.* 106, 190502 (2011).
- [12] Aolita, L. and Mintert, F.: Measuring multipartite concurrence with a single factorizable

- observable. Phys. Rev. Lett. 97, 050501(2006); Carvalho, A.R.R., Mintert, F., and Buchleitner, A.: Decoherence and multipartite entanglement. Phys. Rev. Lett. 93, 230501(2004).
- [13] Vicente, J.D.: Separability criteria based on the Bloch representation of density matrices. Quantum Inf. Comput. 7, 624-638(2007).
- [14] Vicente J.D.: Further results on entanglement detection and quantification from the correlation matrix criterion. J. Phys. A: Math. and Theor., 41, 065309(2008).
- [15] Hassan, A.S.M., Joag, P.S.: Separability criterion for multipartite quantum states based on the bloch representation of density matrices. Quantum Inf. Comput. 8, 773-790(2008).
- [16] Li M., Wang, J., Fei, S.M. and Li-Jost, X.Q.: Quantum separability criterions for arbitrary dimensional multipartite states. Phys. Rev. A, 89,022325(2014).
- [17] Hassan, A.S.M., Joag, P.S.: Experimentally accessible geometric measure for entanglement in N-qubit pure states. Phys. Rev. A, 77, 062334 (2008).
- [18] Hassan, A.S.M., Joag, P.S.: Geometric measure for entanglement in N-qudit pure states. Phys. Rev. A, 80, 042302(2009).
- [19] Li, M. and Fei, S.M.: Bell inequality for multipartite qubit quantum system and the maximal violation. Phys. Rev. A, 86, 052119 (2012).
- [20] Li, M., Wang, J., Fei, S.M., Li-Jost, X.Q. and Fan, H.: Genuine multipartite entanglement detection and lower bound of multipartite concurrence. Phys. Rev. A 92, 062338 (2015).
- [21] Coffman, V., Kundu, J., and Wootters, W.K.: Distributed entanglement. Phys. Rev. A 61, 052306 (2000); Osborne, T.J.: Entanglement measure for rank-2 mixed states. Phys. Rev. A 72, 022309(2005); Rungta, P. and Caves, C. M.: Concurrence-based entanglement measures for isotropic states. Phys. Rev. A 67, 012307(2003).
- [22] Li, M., Fei, S.M. and Wang, Z.X.: Lower bound of concurrence for multipartite quantum states. J. Phys. A, Math. Theor. 42, 145303(2009).
- [23] Hassan, A.S.M. and Joag, P.S.: Measurement-Induced Non-locality in an n -partite quantum state. arXiv:1209.1884v2(2012).

Supplemental material for " Bounds on Multipartite Concurrence and Tangle"

A. Equivalent representation of multipartite concurrence for pure states

To rewritten $\mathcal{C}_N(|\psi\rangle\langle\psi|)$, one can represent both $\text{Tr}\rho^2$ and $\text{Tr}\rho_\alpha^2$ with respect to the correlation tensors. By using the orthogonality of the $SU(d)$ generators we have

$$\text{Tr}\rho^2 = \frac{1}{d^{2N}}(d^N + \sum_{M=1}^N 2^M d^{N-M} \sum_{k_1 \cdots k_M \subset \{1,2,\dots,N\}} \|\mathcal{T}^{k_1 \cdots k_M}\|^2) \quad (14)$$

and

$$\text{Tr}\rho_{k_1 \cdots k_M}^2 = \frac{1}{d^{2M}}(d^M + \sum_{r=1}^M 2^r d^{M-r} \sum_{k_{j_1} \cdots k_{j_r} \subset \{k_{j_1} \cdots k_{j_M}\}} \|\mathcal{T}^{k_{j_1} \cdots k_{j_r}}\|^2) \quad (15)$$

for any $1 \leq M \leq N-1$, where $\rho_{k_1 \cdots k_M}$ is the M -partite reduced density matrix supporting on $\mathcal{H}_{k_1} \otimes \mathcal{H}_{k_2} \otimes \cdots \otimes \mathcal{H}_{k_M}$.

For any $1 \leq M \leq N-1$, we get

$$\begin{aligned} & \sum_{k_1, k_2, \dots, k_M} (\text{Tr}\rho^2 - \text{Tr}\rho_{k_1 \cdots k_M}^2) \\ &= \sum_{l=0}^M \left(\binom{N}{M} \frac{2^l d^{N-l}}{d^{2N}} - \binom{N-l}{M-l} \frac{2^l d^{M-l}}{d^{2M}} \right) \sum_{k_1 k_2 \cdots k_l \subset \{1,2,\dots,N\}} \|\mathcal{T}^{k_1 k_2 \cdots k_l}\|^2 \\ &+ \sum_{l=1}^{N-M} \binom{N}{M} \frac{2^{M+l} d^{N-(M+l)}}{d^{2N}} \sum_{k_1 k_2 \cdots k_{M+l} \subset \{1,2,\dots,N\}} \|\mathcal{T}^{k_1 k_2 \cdots k_{M+l}}\|^2. \end{aligned}$$

Notice that for pure state, we have $\text{Tr}\rho^2 = 1$. Thus the concurrence for pure state can be equivalently represented by

$$\mathcal{C}_N(|\psi\rangle\langle\psi|) = 2^{1-\frac{N}{2}} \sqrt{\sum_{\alpha} (\text{Tr}\rho^2 - \text{Tr}\{\rho_{\alpha}^2\})}. \quad (16)$$

By substituting the representation for $\sum_{k_1, k_2, \dots, k_M} (\text{Tr}\rho^2 - \text{Tr}\rho_{k_1 \cdots k_M}^2)$ above into (16), for pure states $|\psi\rangle$ we have

$$\begin{aligned} & 2^{N-2} \mathcal{C}_N^2(|\psi\rangle\langle\psi|) \\ &= \sum_{M=1}^{N-1} \sum_{k_1, k_2, \dots, k_M} (\text{Tr}\rho^2 - \text{Tr}\rho_{k_1 \cdots k_M}^2) \\ &= \frac{1}{d^N} [-(d+1)^N + d^N + 2^N - 1] \\ &+ \sum_{l=1}^N \frac{2^l}{d^{N+l}} [-(d+1)^{N-l} + 2^N - 1] \sum_{k_1 k_2 \cdots k_l \subset \{1,2,\dots,N\}} \|\mathcal{T}^{k_1 k_2 \cdots k_l}\|^2 \quad (17) \end{aligned}$$

Since $\text{Tr}\rho^2 = 1$ for any pure state $\rho = |\psi\rangle\langle\psi|$, from (14) we have

$$\sum_{k_1 \in \{1,2,\dots,N\}} \|\mathcal{T}^{k_1}\|^2 = \frac{d^{2N} - d^N}{2d^{N-1}} - \sum_{l=2}^N \frac{2^l d^{N-l}}{2d^{N-1}} \sum_{k_1 \dots k_l \subset \{1,2,\dots,N\}} \|\mathcal{T}^{k_1 \dots k_l}\|^2. \quad (18)$$

Substituting (18) into (17), we obtain an alternative relation about the the concurrence, i.e. equation (7).