

Gravitational relaxation of electroweak hierarchy problem

Hiroki Matsui^(a,b), Yoshio Matsumoto^(b),

^(a)*KEK Theory Center, IPNS, KEK, Tsukuba, Ibaraki 305-0801, Japan*

^(b)*The Graduate University of Advanced Studies (Sokendai), Tsukuba, Ibaraki 305-0801,
Japan*

Abstract

In the present paper, we discuss gravitational relaxation models for the electroweak hierarchy problem. We show that modified gravity can naturally relax the electroweak hierarchy problem where conformal transformation provides a crucial rule about what modified gravity theories are favored to relax the electroweak hierarchy. The conformal transformation connects different gravitational theories and rescaling the metric changes the dimensional parameters like the Higgs mass or the cosmological constant in different frames drastically. When the electroweak scale is naturally realized by dynamical and running behavior of dilatonic scalar field or scaling parameter, the modified gravity theories can relax the electroweak hierarchy problem. We discuss the theoretical and phenomenological validity of the gravitational relaxation models.

Contents

1	Introduction	1
2	Gravitational relaxation	2
2.1	Sequestered electroweak hierarchy	6
3	Running gravitational relaxation	7
4	Conclusion	9

1 Introduction

The electroweak hierarchy problem has been recognized as the most notorious difficulty for the high-energy physics in the past decades and often rephrased as the naturalness problem [1, 2, 3, 4]. The naturalness that the low-energy effective field theory should not be extremely sensitive to the high-energy theory is a theoretical and reasonable presumption. Actually, dimensional parameters like scalar masses are as large as the ultraviolet (UV) cut-off scale without involving any special fine-tuning of the parameters or any symmetry. For the Standard Model (SM) case, the Higgs boson mass grows up to the UV cut-off scale M_{UV} by the quadratically divergent quantum corrections ¹

$$\delta M_H^2 \simeq \frac{\alpha}{(4\pi)^2} M_{\text{UV}}^2, \quad (1)$$

where δM_H^2 should not be much larger than the observed Higgs boson mass $M_H^{\text{obs}} = 125.09$ GeV [5, 6, 7]. The majority of theoretical efforts to solve the Higgs naturalness or electroweak hierarchy problem assume a TeV-scale new physics and many models has been proposed, e.g. supersymmetry, extra-dimensions and compositeness. However, these prominent proposals has been suffered from the observed Higgs boson mass $M_H^{\text{obs}} = 125.09$ GeV and the current experimental constraints on new physics.

Recently, theoretically different approaches to the electroweak hierarchy problem have been explored in Ref.[8, 9, 10, 11, 12, 13, 14, 15, 16] based on the cosmological relaxation model of Ref.[17] explaining dynamically the smallness of the cosmological constant. This

¹On the other hand, the cosmological constant problem is more serious from the viewpoint of the naturalness or hierarchy problem. The quantum radiative corrections to the vacuum energy density ρ_{vacuum} which is dubbed *zero-point vacuum energy* enlarges up to the cut-off scale M_{UV} as follows:

$$\delta\rho_{\text{vacuum}} = \frac{1}{2} \int^{M_{\text{UV}}} \frac{d^3k}{(2\pi)^3} \sqrt{k^2 + m^2} = \frac{M_{\text{UV}}^4}{16\pi^2} + \frac{m^2 M_{\text{UV}}^2}{16\pi^2} + \frac{m^4}{64\pi^2} \log\left(\frac{m^2}{M_{\text{UV}}^2}\right) + \dots,$$

which is much larger than the dark energy 2.4×10^{-3} eV in the current Universe.

relaxation mechanism is based on the cosmological evolution of the Higgs field and the axion-like field in the inflationary Universe, and can lead to the naturally small electroweak scale against the cut-off scale.

In this paper we discuss other relaxation scenarios to the electroweak hierarchy problem by involving the gravitational modification which has been proposed and discussed in Ref.[18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29]. Especially, we demonstrate that these gravitational relaxation models can be embedded in the framework of modified gravity theory. By using the conformal transformation or rescaling the metric we clearly show that the electroweak hierarchy can be relaxed from the gravity sector. The conformal transformation connects different gravity theories and rescaling the metric drastically changes dimensional parameters like the Higgs boson mass or cosmological constant in the different frame. We consider several gravitational relaxation models, and discuss the theoretical validity and phenomenological constraints.

The layout of this paper is the following: In Section 2 we introduce the basic formulation for the gravitational relaxation scenarios and discuss why the extended gravity theory relax the electroweak hierarchy problem using the conformal transformation. In Subsection 2.1 we apply the vacuum energy sequestering for the electroweak hierarchy problem as an example of the gravitational modification. In Section 3 we consider the running gravitational relaxation scenarios where the quantum equivalence between in the Einstein and Jordan frames is crucial. Finally, in Section 4 we summarize the conclusion of our work.

2 Gravitational relaxation

In this section, we introduce the gravitational relaxation scenario for the electroweak hierarchy problem. We consider modified gravity theory including the Higgs field Φ and the dilatonic scalar field χ non-minimally coupled to the gravity. The classic action is written by

$$S \supset S_{\text{gravity}} + S_{\text{Higgs}}. \quad (2)$$

The action for the gravity sector including the dilatonic scalar field χ is given by [30]

$$S_{\text{gravity}} = \int d^4x \sqrt{-g} \left(F(\chi) R - \frac{G(\chi)}{2} g^{\mu\nu} \nabla_\mu \chi \nabla_\nu \chi - V(\chi) \right), \quad (3)$$

The action for the Higgs sector is given by

$$S_{\text{Higgs}} = - \int d^4x \sqrt{-g} \left(\frac{1}{2} g^{\mu\nu} \nabla_\mu \Phi^\dagger \nabla_\nu \Phi + V(\Phi^\dagger \Phi) \right), \quad (4)$$

where the (bare) Higgs potential can be written by

$$V(\Phi^\dagger \Phi) = \Lambda_b + M_\Phi^2 (\Phi^\dagger \Phi) + \lambda (\Phi^\dagger \Phi)^2, \quad (5)$$

Now, we assume that the Higgs mass M_Φ and the cosmological constant Λ_b are the UV scale to be $\Lambda_b^{1/4} \simeq M_\Phi \simeq M_{\text{UV}}$. If there exists exact supersymmetry or conformal symmetry, such symmetries can force the cosmological constant or the Higgs mass parameter to be smaller than the cut-off scale. However, such symmetries are always broken in real world and the dimensional parameters grow in proportion to the breaking scale. Thus, we eventually encounter the hierarchy problem via the symmetry breaking scale, and that is the situation in the standard SUSY models. Generally, it is difficult to protect the dimensional parameters from both large classical and quantum corrections of the high-energy physics and that is the reason why the hierarchy problem is thought to be serious.

Now, we rescale the metric via the conformal transformations as follows [30]:

$$g_{\mu\nu} \rightarrow \bar{g}_{\mu\nu} = \Omega^2(\chi) g_{\mu\nu}, \quad (6)$$

$$g^{\mu\nu} \rightarrow \bar{g}^{\mu\nu} = \Omega^{-2}(\chi) g^{\mu\nu}, \quad (7)$$

$$\sqrt{-g} \rightarrow \sqrt{-\bar{g}} = \Omega^4(\chi) \sqrt{-g}. \quad (8)$$

The scalar curvature is transformed as follows:

$$\bar{R} = \frac{1}{\Omega^2(\chi)} \left[R - \frac{6\Box\Omega(\chi)}{\Omega(\chi)} \right], \quad (9)$$

where \Box denotes the covariant d'Alembertian operator and satisfies,

$$\Box\Omega = g^{\mu\nu}\nabla_\mu\nabla_\nu\Omega = \frac{1}{\sqrt{-g}}\partial_\mu[\sqrt{-g}g^{\mu\nu}\partial_\nu\Omega]. \quad (10)$$

Thus, the action for the gravity sector can be transformed as follows:

$$S_{\text{gravity}} = \int d^4x \sqrt{-\bar{g}} \left(\frac{M_{\text{pl}}^2}{2} \bar{R} - \frac{\bar{G}(\chi)}{2} \bar{g}^{\mu\nu} \bar{\nabla}_\mu \chi \bar{\nabla}_\nu \chi - \bar{V}(\chi) \right), \quad (11)$$

where we write down the action in Einstein frame where the scalar curvature is not multiplied by the scalar field, and $\Omega(\chi)$, $F(\chi)$, $\bar{V}(\chi)$ and $\bar{G}(\chi)$ are given by

$$\Omega^2(\chi) = \frac{2F(\chi)}{M_{\text{pl}}^2}, \quad \bar{V}(\chi) = \frac{V(\chi)}{\Omega^4(\chi)}, \quad (12)$$

$$\bar{G}(\chi) = \frac{G(\chi)}{\Omega^2(\chi)} + \frac{6M_{\text{pl}}^2}{\Omega^2(\chi)} \frac{\nabla_\mu\Omega\nabla_\nu\Omega}{\nabla_\mu\chi\nabla_\nu\chi} + \dots \quad (13)$$

where $M_{\text{pl}} = 1/(8\pi G)^{1/2} = 2.4 \times 10^{18}$ GeV is the reduced Planck mass and the related Newton's constant has tight constraints from the cosmological observations [31, 32].

For the Higgs sector the action is given by

$$S_{\text{Higgs}} = - \int d^4x \sqrt{-g} \left(\frac{1}{2} g^{\mu\nu} \nabla_\mu \Phi^\dagger \nabla_\nu \Phi + \Lambda_b + M_\Phi^2 (\Phi^\dagger \Phi) + \lambda (\Phi^\dagger \Phi)^2 \right), \quad (14)$$

The Higgs field Φ are transformed as

$$\Phi \rightarrow H = \Omega^{-1}(\chi) \Phi, \quad (15)$$

The Higgs potential is transformed as follows:

$$V(H^\dagger H) = \frac{\Lambda_b}{\Omega^4(\chi)} + \frac{M_\Phi^2}{\Omega^2(\chi)} (H^\dagger H) + \lambda (H^\dagger H)^2. \quad (16)$$

Note that the action in the SM is conformally invariant except for the Higgs potential. On the other hand, the action of the gauge or fermion fields are only rescaled via the conformal transformations and the couplings are not changed. As the mathematical manipulation, there are several metric frames, e.g. Jordan frame (String frame) and Einstein frame. However, we comment that there is no consensus about physical equivalence of these frames over the years [33, 34, 35, 36, 37, 38, 39, 40] and one should not determine a unique physical frame in modified gravity theory.

The modified gravity theory like extra-dimensions or string theory is often written by Jordan frame. The conformal transformation from Jordan frame to Einstein frame suppress the dimensional parameters of the ordinary SM via the scaling parameter. Our set-up is similar to the Randall-Sundrum model [41] where the large hierarchy is suppressed by the exponential warping factor $e^{-kr_c\phi}$ which depends on an addition extra-dimension ². In this scenario, the four-dimensional components of the bulk metric $g^{\mu\nu}$ and the four-dimensional physical metric $\bar{g}^{\mu\nu}$ have the relation $g^{\mu\nu} = e^{-2kr_c\phi} \bar{g}^{\mu\nu}$ where k is a Planck scale constant and ϕ is the extra-dimensional coordinate with the size r_c ³. By using the rescaling physical metric $\bar{g}^{\mu\nu}$ instead of the bulk metric $g^{\mu\nu}$, this action can be written as,

$$\begin{aligned} S_{\text{Higgs}} &= - \int d^4x \sqrt{-\bar{g}} e^{-4kr_c\phi} \left(\frac{e^{2kr_c\phi}}{2} \bar{g}^{\mu\nu} \bar{\nabla}_\mu \Phi \bar{\nabla}_\nu \Phi + \Lambda_b + M_\Phi^2 (\Phi^\dagger \Phi) + \lambda (\Phi^\dagger \Phi)^2 \right) \\ &= - \int d^4x \sqrt{-\bar{g}} \left(\frac{1}{2} \bar{g}^{\mu\nu} \bar{\nabla}_\mu H \bar{\nabla}_\nu H + \Lambda + M_H^2 (H^\dagger H) + \lambda (H^\dagger H)^2 \right), \end{aligned} \quad (17)$$

²The five-dimensional metric in the Randall-Sundrum model takes the form

$$ds^2 = e^{-2kr_c\phi} \eta_{\mu\nu} dx^\mu dx^\nu + r_c^2 d\phi^2,$$

where $\eta_{\mu\nu}$ is the 4D Minkowski metric.

³In the Randall-Sundrum model, the four-dimensional action for the gravity sector is given by

$$S_{\text{gravity}} \supset \int d^4x \int d\phi 2M^3 r_c e^{-2kr_c|\phi|} \sqrt{-\bar{g}} \bar{R},$$

where M is the five-dimensional Planck scale and \bar{R} is constructed by the rescaling metric $\bar{g}_{\mu\nu}$. Ther 4D Planck scale M_{pl} can be determined as follows:

$$M_{\text{pl}}^2 = M^3 r_c \int d\phi e^{-2kr_c|\phi|} = \frac{M^3}{k} \{1 - e^{-2kr_c\pi}\}.$$

where H is the transformed Higgs field satisfying the relation $H = e^{kr_c\phi}\Phi$. The Higgs mass parameter M_Φ in the fundamental higher-dimensional theory can be suppressed when measured with the rescaling physical metric $\bar{g}^{\mu\nu}$ and become the order of the electroweak scale without difficulty as the following,

$$M_H = e^{-kr_c\phi} M_\Phi \quad (18)$$

Let us discuss several gravitational relaxation models. If we take $\Omega^2(\chi) = M_{\text{pl}}^2/\chi^2$ [42, 43] and $F(\chi) = M_{\text{pl}}^4/2\chi^2$, the transformed Higgs potential is given by

$$V(H^\dagger H) = \frac{\Lambda_b}{M_{\text{pl}}^4} \chi^4 + \frac{M_\Phi^2}{M_{\text{pl}}^2} \chi^2 (H^\dagger H) + \lambda (H^\dagger H)^2. \quad (19)$$

where the Higgs mass and cosmological constant are suppressed and screened off from the cut-off scale. Next, we consider $\Omega^2(\chi) = \chi^2/M_{\text{pl}}^2$, $F(\chi) = \chi^2/2$, the transformed Higgs potential can be written as

$$V(H^\dagger H) = \frac{M_{\text{pl}}^4 \Lambda}{\chi^4} + \frac{M_{\text{pl}}^2 M_\Phi^2}{\chi^2} (H^\dagger H) + \lambda (H^\dagger H)^2. \quad (20)$$

which correspond to the screen scenario for the cosmological constant by so-called cosmon field [44, 45, 42, 43]. When the classic scalar field or vacuum expectation value (VEV) become larger and larger $\chi \gg 1$, the Higgs mass or the cosmological constant are sufficiently suppressed and asymptotically vanish. Next we consider the specific dilaton model [22] where the action can be written as follows:

$$S_{\text{gravity+Higgs}} = \int d^4x \sqrt{-g} \left(\frac{M_{\text{UV}}^2}{2} e^{2\chi/\eta} R - \frac{1}{2} g^{\mu\nu} \nabla_\mu \chi \nabla_\nu \chi - \lambda_\chi (\chi^2 - v_\chi^2) - \frac{1}{2} g^{\mu\nu} \nabla_\mu \Phi^\dagger \nabla_\nu \Phi - \Lambda_b - M_\Phi^2 (\Phi^\dagger \Phi) - \lambda (\Phi^\dagger \Phi)^2 \right). \quad (21)$$

where we assume that M is the order of the cut-off scale to be $\Lambda_b^{1/4} \simeq M_\Phi \simeq M_{\text{UV}}$. Let us transform this action into Einstein frame by rescaling the metric,

$$\bar{g}_{\mu\nu} = \frac{M_{\text{UV}}^2}{M_{\text{pl}}^2} e^{2\chi/\eta} g_{\mu\nu} \simeq \frac{M_{\text{UV}}^2}{M_{\text{pl}}^2} e^{2v_\chi/\eta} g_{\mu\nu}. \quad (22)$$

The Higgs mass parameter M_Φ can be exponentially suppressed as the following

$$M_H \simeq \frac{M_{\text{pl}} M_\Phi}{M_{\text{UV}}} e^{-v_\chi/\eta} \quad (23)$$

where we can have two approaches and interpretations. If we regard $g^{\mu\nu}$ as the physical metric, the Planck-mass scale emerges dynamically by the spontaneous symmetry breaking of

the dilaton symmetry as $M_{\text{pl}} \simeq M_{\text{UV}} e^{v_x/\eta}$. We can solve the large hierarchy problem by assuming $\Lambda_{\text{b}}^{1/4} \simeq M_{\Phi} \simeq M_{\text{UV}} \simeq M_{\text{EW}}$ where M_{EW} express the electroweak scale. On the other hand, we can regard $\bar{g}^{\mu\nu}$ as the physical metric and set $M_{\text{EW}} \simeq M_{\text{pl}} e^{-v_x/\eta}$. This approach resembles the Randall-Sundrum model and the simplest possibility of gravitational relaxations although the quantum gravity effects might appear above the electroweak scale [22]. In these models, the dimensional parameters like the Higgs boson mass, or even the cosmological constant are screened off from the cut-off scale. However, we can not solve both the electroweak hierarchy problem and the cosmological constant problem at once. After all we encounter the fine-tuning problem between the TeV scale and the dark energy although the physical cosmological constant to be $\Lambda^{1/4} \simeq \Omega^{-1}(\chi) \Lambda_{\text{b}}^{1/4} \simeq M_{\text{EW}}$ might be more or less relaxed. Thus, we must require another relaxation mechanism for the cosmological constant.

2.1 Sequestered electroweak hierarchy

Hereafter, we will show that the vacuum energy sequestering scenario can also be effective against the electroweak hierarchy problem and discuss the relation between such a model and previous scenarios. The vacuum energy sequestering scenario proposed in Ref.[24, 25, 26, 27, 28, 29] is the simple model to solve the cosmological constant problem, and can relax the large discrepancy between the vacuum energy density from quantum corrections and the current observed value via the scaling parameter η . This scenario assumes a minimal modification of general relativity to make all scales in the matter sector functionals of the 4-volume element of the Universe. In the context of this scenario, the Universe should be finite in space-time and a transient stage with the present epoch of accelerated expansion before the big crunch [46], but it has been shown that these models could be consistent with the cosmological observation (there are similar models in the context of the unimodular gravity and more detailed discussions are given by Ref.[47, 48, 49, 50, 51, 52, 53, 54, 55]). The mechanism of this scenario is almost the same as the previously discussed one and the unknown scaling parameter η can be regarded as the VEV of the dilatonic scalar field. The vacuum energy sequestering scenario is described by the following action [24]

$$S_{\text{gravity+Higgs}} = \int d^4x \sqrt{-g} \left[\frac{M_{\text{pl}}^2}{2} R - \Lambda_{\text{b}} - \eta^4 \mathcal{L}(\eta^{-2} g^{\mu\nu}, \Phi) \right] + \sigma \left(\frac{\Lambda_{\text{b}}}{\eta^4 \mu^4} \right), \quad (24)$$

where Λ_{b} is the bare cosmological constant and η is the scaling parameter relaxing the large hierarchy. The function $\sigma(x)$ is an adequate function to impose the global constraints and μ is a parameter with the mass dimension. The Lagrangian density $\mathcal{L}(\eta^{-2} g^{\mu\nu}, \Phi)$ for the matter sector couples minimally to the rescaled metric $\bar{g}_{\mu\nu} = \eta^2 g_{\mu\nu}$ and includes the Higgs potential as

$$\begin{aligned} \mathcal{L}(\eta^{-2} g^{\mu\nu}, \Phi) &\supset V(\Phi^\dagger \Phi) \\ &= \Lambda_{\text{b}} + M_{\Phi}^2 (\Phi^\dagger \Phi) + \lambda (\Phi^\dagger \Phi)^2. \end{aligned} \quad (25)$$

The parameter η sets the hierarchy between the physical scale (the electroweak scale or the dark energy scale) and the UV cut-off scale M_{UV} . Thus, the Higgs mass parameter of the

order of the UV cut-off scale are sufficiently suppressed as

$$\frac{M_H}{M_{\text{UV}}} = \eta \frac{M_\Phi}{M_{\text{UV}}}, \quad (26)$$

where M_H is the observed Higgs boson mass and $\eta \ll 1$. To show more accurately the heart of this mechanism, we rewrite this action for the Higgs sector by using the rescaled metric $\bar{g}_{\mu\nu}$ as

$$S_{\text{gravity+Higgs}} = \int d^4x \sqrt{-g} \left(\frac{M_{\text{pl}}^2}{2} R - \Lambda_b \right) - \int d^4x \sqrt{-\bar{g}} \mathcal{L}(\bar{g}^{\mu\nu}, \Phi) + \sigma \left(\frac{\Lambda_b}{\eta^4 \mu^4} \right), \quad (27)$$

where

$$\sqrt{-\bar{g}} \mathcal{L}(\bar{g}^{\mu\nu}, \Phi) = \sqrt{-g} \eta^4 \mathcal{L}(\eta^{-2} g^{\mu\nu}, \Phi). \quad (28)$$

The matter Lagrangian is written by the rescaling metric $\bar{g}_{\mu\nu}$ as

$$\sqrt{-\bar{g}} \mathcal{L}(\bar{g}^{\mu\nu}, \Phi) = \sqrt{-\bar{g}} \left[\frac{1}{2} \bar{g}^{\mu\nu} \bar{\nabla}_\mu \Phi^\dagger \bar{\nabla}_\nu \Phi + \Lambda + M_\Phi^2 (\Phi^\dagger \Phi) + \lambda (\Phi^\dagger \Phi)^2 \right].$$

Thus, by using the conformal transformation, the matter Lagrangian can be written as

$$\begin{aligned} \sqrt{-\bar{g}} \mathcal{L}(\bar{g}^{\mu\nu}, \Phi) &= \sqrt{-g} \eta^4 \mathcal{L}(\eta^{-2} g^{\mu\nu}, \Phi) \\ &= \sqrt{-g} \left[\frac{1}{2} g^{\mu\nu} \nabla_\mu H^\dagger \nabla_\nu H + \eta^4 \Lambda + \eta^2 M_\Phi^2 (H^\dagger H) + \lambda (H^\dagger H)^2 \right], \end{aligned}$$

where we assume $H = \eta\Phi$. Therefore, if the Higgs sector is sequestered from the gravitational sector via the scaling parameter $\eta \ll 1$, the large Higgs mass parameter can be sufficiently suppressed. The unknown scaling parameter η can be regarded as the dilatonic scalar field in the scenarios previously discussed. Although the cosmological constraints on the modified gravity theory and the equivalence between the Einstein and Jordan frames should be carefully considered, this mechanisms or scenarios would not significantly change. In this paper we have focused on the possibility relaxing the electroweak hierarchy and left detail discussion on the cosmological constraints for a forthcoming work.

3 Running gravitational relaxation

In the previous section, we have discussed how modified gravity can alleviate the large Higgs mass and the cosmological constant by using the conformal transformation. In this section, we discuss the running gravitational relaxation scenario ⁴ in which we treat the dilatonic

⁴Polyakov proposed that the cosmological constant could be screened by the IR behavior of quantum gravity and the behavior can be translated by the RG running of the auxiliary gravitational field [18, 19]. The electroweak hierarchy is also discussed by Ref. [23].

scalar field χ as the quantum field and consider the renormalisation group (RG) running behaviors

For simplicity, we consider the action for the Higgs sector in Jordan frame as follows:

$$S_{\text{gravity+Higgs}} \supset \int d^4x \sqrt{-g} \left(\frac{M_{\text{pl}}^4}{2\chi^2} R - \frac{1}{2} g^{\mu\nu} \nabla_\mu \Phi^\dagger \nabla_\nu \Phi - \Lambda_b - M_\Phi^2 (\Phi^\dagger \Phi) - \lambda (\Phi^\dagger \Phi)^2 \right), \quad (29)$$

where we assume that the dilatonic scalar χ satisfy

$$G(\chi) = 6M_{\text{pl}}^2/\chi^2 (2 - M_{\text{pl}}^2/\chi^2) + \dots, \quad V(\chi) = 0. \quad (30)$$

This set-up is for simplifying our discussion and this action is consistent with induced gravity theories [56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69]

Let us perform the conformal transformation $\Omega^2(\chi) = M_{\text{pl}}^2/\chi^2$ and consider the action for the Higgs sector in Einstein frame

$$S_{\text{gravity+Higgs}} \supset \int d^4x \sqrt{-\bar{g}} \left(\frac{M_{\text{pl}}^2}{2} \bar{R} - 6\bar{g}^{\mu\nu} \bar{\nabla}_\mu \chi \bar{\nabla}_\nu \chi - \frac{1}{2} \bar{g}^{\mu\nu} \bar{\nabla}_\mu H^\dagger \bar{\nabla}_\nu H - \frac{\Lambda_b}{M_{\text{pl}}^4} \chi^4 - \frac{M_\Phi^2}{M_{\text{pl}}^2} \chi^2 (H^\dagger H) - \lambda (H^\dagger H)^2 \right). \quad (31)$$

Here, we redefine the scalar field as $\chi \rightarrow \phi = \chi\sqrt{12}$ and obtain the action for the Higgs sector as follows:

$$S_{\text{gravity+Higgs}} \supset \int d^4x \sqrt{-\bar{g}} \left(\frac{M_{\text{pl}}^2}{2} \bar{R} - \frac{1}{2} \bar{g}^{\mu\nu} \bar{\nabla}_\mu \phi \bar{\nabla}_\nu \phi - \frac{1}{2} \bar{g}^{\mu\nu} \bar{\nabla}_\mu H^\dagger \bar{\nabla}_\nu H - \frac{\Lambda_b}{144M_{\text{pl}}^4} \phi^4 - \frac{M_\Phi^2}{12M_{\text{pl}}^2} \phi^2 (H^\dagger H) - \lambda (H^\dagger H)^2 \right). \quad (32)$$

The transformed Higgs potential can be given by

$$V(H^\dagger H) = \lambda_\Lambda \phi^4 + \lambda_M \phi^2 (H^\dagger H) + \lambda (H^\dagger H)^2, \quad (33)$$

where the Higgs mass parameter and the cosmological constant become marginal operators with zero scaling dimensions and they are fixed at the UV cut-off scale as

$$\lambda_\Lambda(\mu_{\text{UV}}) = \frac{\Lambda_b}{144M_{\text{pl}}^4}, \quad \lambda_M(\mu_{\text{UV}}) = \frac{M_\Phi^2}{12M_{\text{pl}}^2}. \quad (34)$$

The renormalisation group (RG) runnings of λ_Λ , λ_M and λ are determined by the one-loop β -functions [70, 71]

$$\beta_{\lambda_\Lambda} = \frac{1}{(4\pi)^2} \left[2\lambda_M^2 + 20\lambda_\Lambda^2 \right], \quad (35)$$

$$\beta_{\lambda_M} = \frac{1}{(4\pi)^2} \left[\frac{1}{2} \lambda_M (12y_t^2 - 3g'^2 - 9g^2) + 4\lambda_M (3\lambda + 2\lambda_\Lambda) + 4\lambda_M^2 \right], \quad (36)$$

$$\beta_\lambda = \frac{1}{(4\pi)^2} \left[\lambda (12y_t^2 - 3g'^2 - 9g^2) - 6y_t^4 + \frac{3}{4}g^4 + \frac{3}{8}(g'^2 + g^2)^2 + 24\lambda^2 + \lambda_M^2 \right]. \quad (37)$$

where the one-loop running of the gauge coupling or the Yukawa coupling are not changed by the conformal transformation because the actions of the gauge sectors or the fermion sectors are only rescaled via the conformal transformations. The transformed Higgs potential has classical conformal symmetry and naturally realize the electroweak scale via the RG running behavior, i.e. the radiative symmetry breaking where these theories are free from the electroweak hierarchy problem (see the reference [72, 73, 74, 75] as the more detailed discussion). Now, we found out that the classically conformal theory [76] which solves the electroweak hierarchy problem is closely related with the modified gravity action of Eq. (32).

Following Polyakov's arguments [18] it is found that the RG running effects of the dilaton or graviton would suppress the Higgs boson mass or cosmological constant in Eq. (32). Although the equivalence between the quantum theories in the Einstein and Jordan frames is still under debate, $f(R)$ gravity theories in the Einstein and Jordan frame are equivalent on shell at the quantum level. The (off shell) quantum corrections are ambiguous, but the equivalence of the effective potential or renormalization group equations have been shown (see, e.g. the reference [77, 78]). Thus, the conformal transformation is also effective for the RG running effects and the modified gravity would relax the electroweak hierarchy.

4 Conclusion

In the present paper, we have discussed the gravitational relaxation scenarios for the electroweak hierarchy problem. We have clearly shown that the conformal transformation has an essential role to understand what modified gravity theory solves the hierarchy problem. If the electroweak scale is naturally realized by dynamical and running behavior of dilatonic scalar field or scaling parameter, the modified gravity theory can relax the electroweak hierarchy problem. The running gravitational relaxation models are closely related with the classically conformal theory and are theoretically attractive as suggested by Polyakov's arguments [18]. Also, the vacuum energy sequestering scenario can be recognized as the dilatonic suppression. The modified gravity theory has the possibility of solving the electroweak hierarchy problem, but require specific constraints on the gravity action with dilatonic scalar fields and it might provide non-trivial effects including metric instability and ghost on the observed Universe.

References

- [1] K. G. Wilson and J. B. Kogut, *The Renormalization group and the epsilon expansion*, *Phys. Rept.* **12** (1974) 75–200.
- [2] G. 't Hooft, *Naturalness, chiral symmetry, and spontaneous chiral symmetry breaking*, *NATO Sci. Ser. B* **59** (1980) 135.
- [3] M. Dine, *Naturalness Under Stress*, *Ann. Rev. Nucl. Part. Sci.* **65** (2015) 43–62, [[1501.01035](https://arxiv.org/abs/1501.01035)].

- [4] G. F. Giudice, *Naturalness after LHC8*, *PoS EPS-HEP2013* (2013) 163, [[1307.7879](#)].
- [5] ATLAS, CMS collaboration, G. Aad et al., *Combined Measurement of the Higgs Boson Mass in pp Collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS and CMS Experiments*, *Phys. Rev. Lett.* **114** (2015) 191803, [[1503.07589](#)].
- [6] ATLAS collaboration, G. Aad et al., *Measurements of Higgs boson production and couplings in diboson final states with the ATLAS detector at the LHC*, *Phys. Lett.* **B726** (2013) 88–119, [[1307.1427](#)].
- [7] CMS collaboration, S. Chatrchyan et al., *Measurement of the properties of a Higgs boson in the four-lepton final state*, *Phys. Rev.* **D89** (2014) 092007, [[1312.5353](#)].
- [8] P. W. Graham, D. E. Kaplan and S. Rajendran, *Cosmological Relaxation of the Electroweak Scale*, *Phys. Rev. Lett.* **115** (2015) 221801, [[1504.07551](#)].
- [9] J. R. Espinosa, C. Grojean, G. Panico, A. Pomarol, O. Pujolàs and G. Servant, *Cosmological Higgs-Axion Interplay for a Naturally Small Electroweak Scale*, *Physical Review Letters* **115** (Dec., 2015) 251803, [[1506.09217](#)].
- [10] E. Hardy, *Electroweak relaxation from finite temperature*, *JHEP* **11** (2015) 077, [[1507.07525](#)].
- [11] S. P. Patil and P. Schwaller, *Relaxing the Electroweak Scale: the Role of Broken dS Symmetry*, *JHEP* **02** (2016) 077, [[1507.08649](#)].
- [12] B. Batell, G. F. Giudice and M. McCullough, *Natural Heavy Supersymmetry*, *JHEP* **12** (2015) 162, [[1509.00834](#)].
- [13] S. Di Chiara, K. Kannike, L. Marzola, A. Racioppi, M. Raidal and C. Spethmann, *Relaxion Cosmology and the Price of Fine-Tuning*, *Phys. Rev.* **D93** (2016) 103527, [[1511.02858](#)].
- [14] O. Matsedonskyi, *Mirror Cosmological Relaxation of the Electroweak Scale*, *JHEP* **01** (2016) 063, [[1509.03583](#)].
- [15] J. L. Evans, T. Gherghetta, N. Nagata and Z. Thomas, *Naturalizing Supersymmetry with a Two-Field Relaxion Mechanism*, *JHEP* **09** (2016) 150, [[1602.04812](#)].
- [16] A. Hook and G. Marques-Tavares, *Relaxation from particle production*, *JHEP* **12** (2016) 101, [[1607.01786](#)].
- [17] L. F. Abbott, *A Mechanism for Reducing the Value of the Cosmological Constant*, *Phys. Lett.* **B150** (1985) 427–430.

- [18] A. m. Polyakov, *PHASE TRANSITIONS AND THE UNIVERSE*, *Sov. Phys. Usp.* **25** (1982) 187.
- [19] A. M. Polyakov, *String theory as a universal language*, *Phys. Atom. Nucl.* **64** (2001) 540–547, [[hep-th/0006132](#)].
- [20] R. Jackiw, C. Nunez and S. Y. Pi, *Quantum relaxation of the cosmological constant*, *Phys. Lett.* **A347** (2005) 47–50, [[hep-th/0502215](#)].
- [21] D. A. Demir, *Gravi-Natural Higgs and Conformal New Physics*, [1207.4584](#).
- [22] C.-S. Lin, *Large Hierarchy from Non-Minimal Coupling*, *Commun. Theor. Phys.* **68** (2017) 223, [[1405.4821](#)].
- [23] A. Kobakhidze, *Quantum relaxation of the Higgs mass*, *Eur. Phys. J.* **C75** (2015) 384, [[1506.04840](#)].
- [24] N. Kaloper and A. Padilla, *Sequestering the Standard Model Vacuum Energy*, *Phys. Rev. Lett.* **112** (2014) 091304, [[1309.6562](#)].
- [25] N. Kaloper and A. Padilla, *Vacuum Energy Sequestering: The Framework and Its Cosmological Consequences*, *Phys. Rev.* **D90** (2014) 084023, [[1406.0711](#)].
- [26] N. Kaloper and A. Padilla, *Sequestration of Vacuum Energy and the End of the Universe*, *Phys. Rev. Lett.* **114** (2015) 101302, [[1409.7073](#)].
- [27] N. Kaloper, A. Padilla, D. Stefanyszyn and G. Zahariade, *Manifestly Local Theory of Vacuum Energy Sequestering*, *Phys. Rev. Lett.* **116** (2016) 051302, [[1505.01492](#)].
- [28] N. Kaloper, A. Padilla and D. Stefanyszyn, *Sequestering effects on and of vacuum decay*, *Phys. Rev.* **D94** (2016) 025022, [[1604.04000](#)].
- [29] N. Kaloper and A. Padilla, *Vacuum Energy Sequestering and Graviton Loops*, *Phys. Rev. Lett.* **118** (2017) 061303, [[1606.04958](#)].
- [30] D. I. Kaiser, *Conformal Transformations with Multiple Scalar Fields*, *Phys. Rev.* **D81** (2010) 084044, [[1003.1159](#)].
- [31] K.-i. Umezū, K. Ichiki and M. Yahiro, *Cosmological constraints on Newton’s constant*, *Phys. Rev.* **D72** (2005) 044010, [[astro-ph/0503578](#)].
- [32] S. Galli, A. Melchiorri, G. F. Smoot and O. Zahn, *From Cavendish to PLANCK: Constraining Newton’s Gravitational Constant with CMB Temperature and Polarization Anisotropy*, *Phys. Rev.* **D80** (2009) 023508, [[0905.1808](#)].
- [33] V. Faraoni and E. Gunzig, *Einstein frame or Jordan frame?*, *Int. J. Theor. Phys.* **38** (1999) 217–225, [[astro-ph/9910176](#)].

- [34] S. Capozziello, P. Martin-Moruno and C. Rubano, *Physical non-equivalence of the Jordan and Einstein frames*, *Phys. Lett.* **B689** (2010) 117–121, [[1003.5394](#)].
- [35] S. Capozziello and D. Saez-Gomez, *Conformal frames and the validity of Birkhoff's theorem*, *AIP Conf. Proc.* **1458** (2011) 347–350, [[1202.2540](#)].
- [36] N. Deruelle and M. Sasaki, *Conformal equivalence in classical gravity: the example of 'Veiled' General Relativity*, *Springer Proc. Phys.* **137** (2011) 247–260, [[1007.3563](#)].
- [37] C. F. Steinwachs and A. Y. Kamenshchik, *Non-minimal Higgs Inflation and Frame Dependence in Cosmology*, *AIP Conf. Proc.* **1514** (2013) 161–164, [[1301.5543](#)].
- [38] X. Calmet and T.-C. Yang, *Frame Transformations of Gravitational Theories*, *Int. J. Mod. Phys.* **A28** (2013) 1350042, [[1211.4217](#)].
- [39] G. Domenech and M. Sasaki, *Conformal Frame Dependence of Inflation*, *JCAP* **1504** (2015) 022, [[1501.07699](#)].
- [40] M. Postma and M. Volponi, *Equivalence of the Einstein and Jordan frames*, *Phys. Rev.* **D90** (2014) 103516, [[1407.6874](#)].
- [41] L. Randall and R. Sundrum, *A Large mass hierarchy from a small extra dimension*, *Phys. Rev. Lett.* **83** (1999) 3370–3373, [[hep-ph/9905221](#)].
- [42] C. Wetterich, *Conformal fixed point, cosmological constant and quintessence*, *Phys. Rev. Lett.* **90** (2003) 231302, [[hep-th/0210156](#)].
- [43] C. Wetterich, *Cosmon inflation*, *Phys. Lett.* **B726** (2013) 15–22, [[1303.4700](#)].
- [44] C. Wetterich, *Cosmology and the Fate of Dilatation Symmetry*, *Nucl. Phys.* **B302** (1988) 668–696.
- [45] C. Wetterich, *The Cosmon model for an asymptotically vanishing time dependent cosmological 'constant'*, *Astron. Astrophys.* **301** (1995) 321–328, [[hep-th/9408025](#)].
- [46] P. P. Avelino, *Vacuum energy sequestering and cosmic dynamics*, *Phys. Rev.* **D90** (2014) 103523, [[1410.4555](#)].
- [47] J. L. Anderson and D. Finkelstein, *Cosmological constant and fundamental length*, *Am. J. Phys.* **39** (1971) 901–904.
- [48] W. Buchmuller and N. Dragon, *Einstein Gravity From Restricted Coordinate Invariance*, *Phys. Lett.* **B207** (1988) 292–294.
- [49] W. Buchmuller and N. Dragon, *Gauge Fixing and the Cosmological Constant*, *Phys. Lett.* **B223** (1989) 313–317.

- [50] M. Henneaux and C. Teitelboim, *The Cosmological Constant and General Covariance*, *Phys. Lett.* **B222** (1989) 195–199.
- [51] W. G. Unruh, *A Unimodular Theory of Canonical Quantum Gravity*, *Phys. Rev.* **D40** (1989) 1048.
- [52] S. Nojiri, *Some solutions for one of the cosmological constant problems*, *Mod. Phys. Lett. A* **31** (2016) 1650213, [1601.02203].
- [53] S. Nojiri, S. D. Odintsov and V. K. Oikonomou, *Bounce universe history from unimodular $F(R)$ gravity*, *Phys. Rev.* **D93** (2016) 084050, [1601.04112].
- [54] S. Nojiri, S. D. Odintsov and V. K. Oikonomou, *Unimodular-Mimetic Cosmology*, *Class. Quant. Grav.* **33** (2016) 125017, [1601.07057].
- [55] S. Nojiri, S. D. Odintsov and V. K. Oikonomou, *Newton law in covariant unimodular $F(R)$ gravity*, *Mod. Phys. Lett. A* **31** (2016) 1650172, [1605.00993].
- [56] A. Zee, *Broken-symmetric theory of gravity*, *Phys. Rev. Lett.* **42** (Feb, 1979) 417–421.
- [57] A. Zee, *Horizon problem and the broken-symmetric theory of gravity*, *Phys. Rev. Lett.* **44** (Mar, 1980) 703–706.
- [58] L. Smolin, *Towards a Theory of Space-Time Structure at Very Short Distances*, *Nucl. Phys.* **B160** (1979) 253–268.
- [59] S. L. Adler, *Einstein Gravity as a Symmetry Breaking Effect in Quantum Field Theory*, *Rev. Mod. Phys.* **54** (1982) 729.
- [60] H. Dehnen, H. Frommert and F. Ghaboussi, *Higgs field gravity*, *Int. J. Theor. Phys.* **29** (1990) 537–546.
- [61] H. Dehnen, H. Frommert and F. Ghaboussi, *Higgs field and a new scalar - tensor theory of gravity*, *Int. J. Theor. Phys.* **31** (1992) 109–114.
- [62] B. L. Spokoiny, *INFLATION AND GENERATION OF PERTURBATIONS IN BROKEN SYMMETRIC THEORY OF GRAVITY*, *Phys. Lett.* **B147** (1984) 39–43.
- [63] F. S. Accetta, D. J. Zoller and M. S. Turner, *Induced Gravity Inflation*, *Phys. Rev.* **D31** (1985) 3046.
- [64] F. Lucchin, S. Matarrese and M. D. Pollock, *Inflation With a Nonminimally Coupled Scalar Field*, *Phys. Lett.* **B167** (1986) 163–168.
- [65] R. Fakir and W. G. Unruh, *Induced gravity inflation*, *Phys. Rev.* **D41** (1990) 1792–1795.

- [66] D. I. Kaiser, *Constraints in the context of induced gravity inflation*, *Phys. Rev.* **D49** (1994) 6347–6353, [[astro-ph/9308043](#)].
- [67] D. I. Kaiser, *Induced gravity inflation and the density perturbation spectrum*, *Phys. Lett.* **B340** (1994) 23–28, [[astro-ph/9405029](#)].
- [68] J. L. Cervantes-Cota and H. Dehnen, *Induced gravity inflation in the $SU(5)$ GUT*, *Phys. Rev.* **D51** (1995) 395–404, [[astro-ph/9412032](#)].
- [69] J. L. Cervantes-Cota and H. Dehnen, *Induced gravity inflation in the standard model of particle physics*, *Nucl. Phys.* **B442** (1995) 391–412, [[astro-ph/9505069](#)].
- [70] M. Gonderinger, Y. Li, H. Patel and M. J. Ramsey-Musolf, *Vacuum Stability, Perturbativity, and Scalar Singlet Dark Matter*, *JHEP* **01** (2010) 053, [[0910.3167](#)].
- [71] J. Elias-Miro, J. R. Espinosa, G. F. Giudice, H. M. Lee and A. Strumia, *Stabilization of the Electroweak Vacuum by a Scalar Threshold Effect*, *JHEP* **06** (2012) 031, [[1203.0237](#)].
- [72] S. Iso, N. Okada and Y. Orikasa, *Classically conformal $B-L$ extended Standard Model*, *Phys. Lett.* **B676** (2009) 81–87, [[0902.4050](#)].
- [73] S. Iso, N. Okada and Y. Orikasa, *The minimal $B-L$ model naturally realized at TeV scale*, *Phys. Rev.* **D80** (2009) 115007, [[0909.0128](#)].
- [74] S. Iso and Y. Orikasa, *TeV Scale $B-L$ model with a flat Higgs potential at the Planck scale - in view of the hierarchy problem -*, *PTEP* **2013** (2013) 023B08, [[1210.2848](#)].
- [75] M. Hashimoto, S. Iso and Y. Orikasa, *Radiative symmetry breaking at the Fermi scale and flat potential at the Planck scale*, *Phys. Rev.* **D89** (2014) 016019, [[1310.4304](#)].
- [76] W. A. Bardeen, *On naturalness in the standard model*, in *Ontake Summer Institute on Particle Physics Ontake Mountain, Japan, August 27-September 2, 1995*, 1995.
- [77] M. S. Ruf and C. F. Steinwachs, *Quantum equivalence of $f(R)$ gravity and scalar-tensor theories*, *Phys. Rev.* **D97** (2018) 044050, [[1711.07486](#)].
- [78] N. Ohta, *Quantum equivalence of $f(R)$ gravity and scalar-tensor theories in the Jordan and Einstein frames*, *PTEP* **2018** (2018) 033B02, [[1712.05175](#)].