

A Unified Hidden-Sector Electroweak Model, Para-photons and the X-boson

M. J. Neves ^a and J. A. Helayël-Neto ^b

^a*Departamento de Física,
Universidade Federal Rural do Rio de Janeiro
BR 465-07, 23890-971, Seropédica, Rio de Janeiro, Brazil*

^b*Centro Brasileiro de Pesquisas Físicas (CBPF),
Rua Dr. Xavier Sigaud 150, Urca,
CEP 22290-180, Rio de Janeiro, Brazil*

February 19, 2022

E-mail: mariojr@ufrrj.br, helayel@cbpf.br

ABSTRACT: Our contribution sets out to investigate the phenomenology of a gauge model based on an $SU_L(2) \times U_R(1)_J \times U(1)_K$ -symmetry group. The model can accommodate in its distinct phases - by virtue of different symmetry-breaking patterns - a candidate to a heavy Z' -boson at the TeV-scale and the recently discussed 17 MeV X-boson that may bring about a new physics at the MeV-scale. Furthermore, the para-photon of the dark matter sector and the candidate to the so-called dark photon can also be described in the scenario we are endeavoring to set up in this paper. In the Z' -scenario, the extended Higgs sector introduces a heavy scalar whose mass lies in the region 1.2 – 3.7 TeV, while that the lower energy scale scenario presents a light scalar with mass in the interval 1.9 – 19 GeV. The fermionic sector includes an exotic candidate to dark matter that mixes with the right-neutrino component in the Higgs sector, so that the whole field content yields the cancellation of the U(1)-anomaly. The masses are fixed according to the particular way the symmetry breaking takes place. In view of the possible symmetry breakdown patterns, we study the phenomenological implications in both scenarios of high- and low-energy scales. As examples, we consider the scattering of Standard Model matter into dark sector constituents intermediated by the Z' -boson in the TeV-scale, and the corresponding scattering intermediated by a lightest boson in the lower scale physics. Furthermore, we obtain the magnetic dipole momentum (MDM) of the exotic fermion and the transition MDM due to the mixing with the right-neutrino.

KEYWORDS: Physics beyond Standard Model, Hidden particles, Dark matter.

Contents

1	Introduction	1
2	A review of the Z'-model	4
3	The mixing between right-neutrinos and ζ_i-fermions	11
4	The Z'-phenomenology	15
4.1	The Z' -decay into fermions : $Z' \rightarrow \bar{f} f$	15
4.2	The F -Higgs decays	16
4.3	The quark-quark scattering into the dark sector	18
5	The MDM for the ζ_i-fermions	20
6	The X-boson scenario at the MeV-scale and below	25
7	The X-boson phenomenology	29
7.1	The X-boson decays processes	29
7.2	The X-boson scattering	30
7.3	The electron and muon $g - 2$ factors	32
8	Concluding Comments	34

1 Introduction

The search for new particles and interactions beyond the Standard Model (SM) has been challenging High-Energy Physics, both Theoretical and Experimental, over the recent decades. The results from the LHC's ATLAS- and CMS-Collaborations may point to the existence of a (new) fifth interaction [1, 2]. Analysing data of pp -collisions at a center-of-mass energy scale of $\sqrt{s} = 13$ TeV has revealed peaks of masses that may correspond to the hypothetical heavy W'^- and Z' -bosons [3, 4], which may be an evidence for new Physics communicated from very high energies to the TeV-scale. We should keep in mind that the introduction of an extra Higgs may be needed to explain the heavy mass of the new bosons at the TeV-scale. These masses are associated with a new range of vacuum expectation values (VEVs) of supplementary Higgs scalars.

A well-known model in this direction is based on an $SU_L(2) \times SU_R(2) \times U(1)_{B-L}$ gauge symmetry [6–13]. The extra $SU_R(2)$ -subgroup is introduced to account for the hypothetical W'^- and Z' -bosons, besides the already known W^\pm - and Z^0 -weak mediators of the Glashow-Salam-Weinberg (GSW) model. The right-handed sector also introduces fermion doublets with a charged lepton and its associated neutrino of right chirality. Two

scalar doublets constitute the Higgs sector responsible for the chain of symmetry breakings. The first doublet is introduced to break the $SU_R(2) \times U(1)_{B-L}$ -gauge symmetry at the scale set up by the VEV u , above the SM Electroweak scale $u > v = 246$ GeV. As a consequence, the hypercharge, Y , appears as a combination of generators of $SU_R(2) \times U(1)_{B-L}$. Next, the Standard Model Higgs breaks the remaining gauge symmetry down to the electromagnetic Abelian symmetry, $U(1)_{em}$. Therefore, the recent CMS simulation indicates that the estimate for masses of hypothetical W' and Z' is around the 2 TeV [3, 4].

The well-known model in the literature that describes only the Z' -heavy boson is based on the gauge symmetry $SU_L(2) \times U_Y(1) \times U(1)_{B-L}$ [14–16], which includes a $U(1)$ -extra group in the GSW model. The gauge sector has just one extra boson, while the Higgs is extended to include a doublet and a singlet, or bi-doublets of scalars fields. The fermion sector is enlarged to guarantee anomaly cancellation; for example, the introduction of right-neutrinos components and exotic fermions that could be candidates to dark matter content.

In 2016, in a particular Nuclear Physics experiment, anomalies in the decay of the excited state of 8Be^* to its ground state has suggested the existence of a new neutral boson, called X , through the decay mode $8\text{Be}^* \rightarrow 8\text{Be} + X$ [17]. The X -boson immediately decays into an electron-positron pair $X \rightarrow e^+ + e^-$. It is a vector-type spin-1 neutral particle, with mass around $m_X = 17$ MeV that mixes with the SM photon through a kinetic term. Its origin could, in principle, be traced back to an extra gauge symmetry, $U(1)_X$, in addition to the symmetries of the SM. The discovery of the X -boson may be pointing to the existence of a fifth fundamental interaction in Nature. This is an interesting scenario in which the hypothetical fifth interaction would be accessible at the MeV-scale.

The effective Lagrangian proposed to describe this extra X -boson is given below [20]

$$\mathcal{L}_{eff} = -\frac{1}{4} F_{\mu\nu}^2 - \frac{1}{4} X_{\mu\nu}^2 + \frac{\chi}{2} X_{\mu\nu} F^{\mu\nu} + \frac{1}{2} m_X^2 X_\mu^2 - J_\mu X^\mu, \quad (1.1)$$

where J^μ is the current weakly coupled to X^μ

$$J^\mu = \sum_{\Psi=e,u,d,\dots} e \chi_\Psi \bar{\Psi} \gamma^\mu \Psi, \quad (1.2)$$

with Ψ standing for any fermion of the SM. This vector current exhibits a protophobic character described by the χ_p -parameter whose magnitude for the proton and neutron satisfies the condition $\chi_p \ll \chi_n$, whenever the X -boson interacts with the nucleon. We list some estimates of χ_Ψ for the fermions of the SM, following the phenomenology of the X -boson [18] :

$$\begin{aligned} 2 \times 10^{-4} < |\chi_e| < 1.4 \times 10^{-3}, \\ |\chi_n| < 2.5 \times 10^{-2}, \\ \sqrt{|\chi_n \chi_e|} \lesssim 7 \times 10^{-5}. \end{aligned} \quad (1.3)$$

while that for the u - and d -quarks, the extreme protophobic limit, ($\chi_p = 0$), parameterizes

χ_u and χ_d as it follows below :

$$\begin{aligned}\chi_u &= -\frac{\chi_n}{3} \simeq \pm 3.7 \times 10^{-3} \\ \chi_d &= +\frac{2\chi_n}{3} \simeq \mp 7.4 \times 10^{-3} .\end{aligned}\tag{1.4}$$

The X^μ -boson can also have a chiral interaction with the SM leptons via an axial current, see [19]. For a complete review on the anomaly in Beryllium decays, see [20]. Still in the Lagrangian (1.1), the parameter χ mixes the X^μ -boson with the usual electromagnetic (EM) photon, A^μ , where $X^{\mu\nu} = \partial^\mu X^\nu - \partial^\nu X^\mu$ is the field-strength tensor for X^μ , and $F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu$, the corresponding tensor field for the photon. It is clear that the massive term spoils the $U(1)_X$ -symmetry, and the Lagrangian exhibits only EM gauge symmetry, $U_{em}(1)$. Thus, the presence of the mass in (1.1) motivates us to search for a physical mechanism that underlies its generation. There has emerged an important connection between the $U(1)_X$ -symmetry and dark matter in a model with spontaneous symmetry breaking (SSB); for that, see the recent paper of Ref. [23].

In another context, the search for single photons in 53 fb^{-1} , in the $e^+ e^-$ CM-collision with the production of a spin-1 particle, A' , referred to as dark photon, is associated to the process $e^+ e^- \rightarrow \gamma A'$ [24]. Immediately, the dark photon decays into invisible fermion matter $A \rightarrow \bar{\zeta} \zeta$, that can be a candidate to dark matter constituent. The A' dark-photon has therefore a similar phenomenology to the X -Boson, with a kinetic mixing of order 10^{-3} , with the EM photon. The dominant decay mode for the lowest-mass of the ζ -state fixes the condition $m_{A'} > 2m_\zeta$, considering that the $m_{A'}$ -dark-photon mass is bounded from above by $m_{A'} \leq 8 \text{ GeV}$.

In the literature, there is also a great deal of interest in the activity related to the phenomenology of hidden sector para-photons [25–27]. The para-photon is a neutral vector boson with a sub-eV mass and with the property of electromagnetic interactions with coupling constants referred to as millicharges. The para-photons are characterized by a mixed mass term with the photon and they appear with an extra $U(1)$ gauge factor in the full symmetry group. The particles like the boson- X , dark-photon or para-photon that interact with a current like (1.2) are known in the literature as Weakly Interacting Massive Particles (WIMP). This idea connects the matter content of the SM with possible new dark matter particles via scattering processes.

In the present paper, we re-assess the $SU_L(2) \times U_R(1)_J \times U(1)_K$ model whose spontaneous symmetry breaking (SSB) mechanism can be applied in two scenarios: with a higher (TeV) or a lower (MeV) energy scale; for details, consult [28]. We study its phenomenology such that the new bosons, Z' or X , can connect the world of SM particles with dark matter particles in particular scattering processes. Furthermore, the introduction of a mixing involving a Dirac right-neutrino component with exotic fermions, *i. e.*, heavy and neutral fermions, can unveil new magnetic properties of these new fermions. We pursue this investigation and we obtain the transition Magnetic Dipole Momentum (MDM) and the MDM for the exotic fermion that depend on its mass, as it happens in the neutrino case. Sectors of fermions and scalar bosons are introduced with quantum numbers consistent with the gauge invariance and the chiral anomalies associated to the Abelian sectors cancel out.

The organization of this paper follows the outline below: in Section II, we review the Z' -model based on an $SU_L(2) \times U_R(1)_J \times U(1)_K$ -symmetry presented in details in [28]. In Section III, we study in details the diagonalization of the dark fermion and right-neutrino sector. Section IV is presented in three subsections, where we study the Z' -decay modes into fermions, the decay modes of the extra Higgs and the scattering processes quark-quark and dark fermion mediated by the Z' particle. In the Section V, we obtain the transition MDM and the MDM of the dark fermion. Section VI contains a review of SSB in association with the X -boson and the dark photon scenario of [28]. In Section VII, we study the decay modes of the dark photon (first subsection) and we work out the scattering process for the cascade effect $e^+ e^- \rightarrow X \rightarrow \bar{\zeta} \zeta$ (second subsection). The third subsection is dedicated to form factor calculations in the correction to the QED vertex due to the axial interaction of the X -boson with the leptons of the SM. Finally, our Concluding Comments are cast in Section VIII.

2 A review of the Z' -model

In this Section, we present a short review of the Z' -model; the details may be found in the work of Ref. [28]. Here, we introduce the model in the renormalizable R_ξ -gauge. The sector of fermions and gauge fields of the model $SU_L(2) \times U_R(1)_J \times U(1)_K$ -model is described by the Lagrangian below :

$$\mathcal{L}_f = \bar{\Psi}_L i \not{D} \Psi_L + \bar{\Psi}_R i \not{D} \Psi_R + \bar{\nu}_{iR} i \not{D} \nu_{iR} + \bar{\zeta}_i i \not{D} \zeta_i, \quad (2.1)$$

and

$$\mathcal{L}_{gauge} = -\frac{1}{2} \text{tr} (F_{\mu\nu}^2) - \frac{1}{4} B_{\mu\nu}^2 - \frac{1}{4} C_{\mu\nu}^2. \quad (2.2)$$

The slashed notation corresponds to the contraction of the covariant derivatives with the usual Dirac matrices. The covariant derivatives acting on the fermions of the model are cast according to:

$$\begin{aligned} D_\mu \Psi_L &= \left(\partial_\mu + i g A_\mu^a \frac{\sigma^a}{2} + i J_L g' B_\mu + i K_L g'' C_\mu \right) \Psi_L, \\ D_\mu \Psi_R &= \left(\partial_\mu + i J_R g' B_\mu + i K_R g'' C_\mu \right) \Psi_R, \\ D_\mu \nu_{iR} &= \left(\partial_\mu + i J_{\nu_R} g' B_\mu + i K_{\nu_R} g'' C_\mu \right) \nu_{iR}, \\ D_\mu \zeta_i &= \left(\partial_\mu + i J_\zeta g' B_\mu + i K_\zeta g'' C_\mu \right) \zeta_i, \end{aligned} \quad (2.3)$$

where $A^{\mu a} = \{ A^{\mu 1}, A^{\mu 2}, A^{\mu 3} \}$ are the gauge fields of $SU_L(2)$, B^μ is the Abelian gauge field of $U_R(1)_J$, and C^μ the similar one to $U(1)_K$. Here, we have chosen the symbol J to represent the generator of $U_R(1)_J$, K is the generator of $U(1)_K$, the generators of $SU_L(2)$ are the Pauli matrices $\frac{\sigma^a}{2}$ ($a = 1, 2, 3$), and g, g' and g'' are dimensionless gauge couplings. The fermionic field content is given in the sequel. The notation Ψ_L indicates the usual doublet of neutrinos/leptons $L_i = (\nu_i \ell_i)_L^t$, or quarks Q_{iL} ($i = 1, 2, 3$) left-handed of the

SM, it turn out in the fundamental representation of $SU_L(2)$. The label $\ell_{iL} = (e_L, \mu_L, \tau_L)$ indicates the leptons family displayed in the doublet, and the ν_{iL} -label for neutrinos is defined by $\nu_{iL} = (\nu_{eL}, \nu_{\mu L}, \nu_{\tau L})$. The Ψ_R -fermion is any right-handed of the SM, *i. e.*, it can be the lepton $\ell_{iR} = (e_R, \mu_R, \tau_R)$ or right-handed quarks $Q_{iR} = \{u_R, c_R, t_R\}$ and $q_{iR} = \{d_R, s_R, b_R\}$. The new content of fermions beyond the SM is represented by the Dirac Right-Neutrino $\nu_{iR} = (\nu_{eR}, \nu_{\mu R}, \nu_{\tau R})$ and the exotic neutral ζ_i -fermion ($i = 1, 2, 3$) that we have introduced it associated with the $U(1)_K$ -group:

$$\begin{aligned}
 L_i &= \begin{pmatrix} \nu_e \\ \ell_e \end{pmatrix}_L, \begin{pmatrix} \nu_\mu \\ \ell_\mu \end{pmatrix}_L, \begin{pmatrix} \nu_\tau \\ \ell_\tau \end{pmatrix}_L : \left(\underline{\mathbf{2}}, 0, -\frac{1}{2} \right), \\
 Q_{iL} &= \begin{pmatrix} u \\ d \end{pmatrix}_L, \begin{pmatrix} c \\ s \end{pmatrix}_L, \begin{pmatrix} t \\ b \end{pmatrix}_L : \left(\underline{\mathbf{2}}, 0, +\frac{1}{6} \right), \\
 \ell_{iR} &= \{e_R, \mu_R, \tau_R\} : \left(\underline{\mathbf{1}}, -\frac{1}{2}, -\frac{1}{2} \right), \\
 Q_{iR} &= \{u_R, c_R, t_R\} : \left(\underline{\mathbf{1}}, +\frac{1}{2}, +\frac{1}{6} \right), \\
 q_{iR} &= \{d_R, s_R, b_R\} : \left(\underline{\mathbf{1}}, -\frac{1}{2}, +\frac{1}{6} \right), \\
 \nu_{iR} &= \{\nu_{eR}, \nu_{\mu R}, \nu_{\tau R}\} : \left(\underline{\mathbf{1}}, +\frac{1}{2}, -\frac{1}{2} \right), \\
 \zeta_{iL} &= \{\zeta_{1L}, \zeta_{2L}, \zeta_{3L}\} : \left(\underline{\mathbf{1}}, +\frac{1}{2}, -\frac{1}{2} \right), \\
 \zeta_{iR} &= \{\zeta_{1R}, \zeta_{2R}, \zeta_{3R}\} : \left(\underline{\mathbf{1}}, -\frac{1}{2}, +\frac{1}{2} \right). \tag{2.4}
 \end{aligned}$$

All these fields are singlets that undergo Abelian transformations under the $U_R(1)_J$ - and $U(1)_K$ -groups.

The corresponding field-strength tensors in the sector of gauge fields are defined by

$$\begin{aligned}
 F_{\mu\nu} &= \partial_\mu A_\nu - \partial_\nu A_\mu + ig [A_\mu, A_\nu], \\
 B_{\mu\nu} &= \partial_\mu B_\nu - \partial_\nu B_\mu, C_{\mu\nu} = \partial_\mu C_\nu - \partial_\nu C_\mu, \tag{2.5}
 \end{aligned}$$

The Higgs sector is essential to introduce the masses, the physical fields and the charges for the particle content of the model. The framework of Higgs sector is setting up by two independent scalar fields; the first is singlet scalar, Ξ , that breaks the Abelian subgroup to generate mass to the new gauge boson, that in this scenario we call it Z -boson. The second Higgs field, Φ , is a $SU_L(2)$ - doublet to break the residual electroweak symmetry and, consequently, it yields the known masses for W^\pm and Z^0 . Finally, we end up with the exact electromagnetic symmetry

$$SU_L(2) \times U_R(1)_J \times U(1)_K \xrightarrow{\langle \Xi \rangle_0} SU_L(2) \times U_Y(1) \xrightarrow{\langle \Phi \rangle_0} U_{em}(1), \tag{2.6}$$

where the $U_Y(1)$ -group comes out as the mixing of the $U_R(1)_J$ - and $U(1)_K$ -subgroups. To accomplish this SSB pattern, we start off from the Higgs Lagrangian below:

$$\begin{aligned}
 \mathcal{L}_{Higgs} = & (D_\mu \Xi)^\dagger D^\mu \Xi - \mu_\Xi^2 |\Xi|^2 - \lambda_\Xi |\Xi|^4 \\
 & + (D_\mu \Phi)^\dagger D^\mu \Phi - \mu_\Phi^2 |\Phi|^2 - \lambda_\Phi |\Phi|^4 - \lambda |\Xi|^2 |\Phi|^2, \\
 & - G_{ij}^{(\ell)} \bar{L}_i \Phi \ell_{jR} - G_{ij}^{(\ell)*} \bar{\ell}_{iR} \Phi^\dagger L_j \\
 & - G_{ij}^{(d)} \bar{Q}_{iL} \Phi q_{jR} - G_{ij}^{(d)*} \bar{q}_{iR} \Phi^\dagger Q_{jL} \\
 & - G_{ij}^{(u)} \bar{Q}_{iL} \tilde{\Phi} Q_{jR} - G_{ij}^{(u)*} \bar{Q}_{iR} \tilde{\Phi}^\dagger Q_{jL} \\
 & - X_{ij} \bar{L}_i \tilde{\Phi} \nu_{jR} - X_{ij}^* \bar{\nu}_{iR} \tilde{\Phi}^\dagger L_j \\
 & - Y_{ij} \bar{L}_i \tilde{\Phi} \zeta_{jR} - Y_{ij}^* \bar{\zeta}_{iR} \tilde{\Phi}^\dagger L_j \\
 & - Z_{ij} \bar{\zeta}_{iL} \Xi \nu_{jR} - Z_{ij}^* \bar{\nu}_{iR} \Xi^\dagger \zeta_{jL} \\
 & - W_{ij} \bar{\zeta}_{iL} \Xi \zeta_{jR} - W_{ij}^* \bar{\zeta}_{iR} \Xi^\dagger \zeta_{jL}.
 \end{aligned} \tag{2.7}$$

In (2.7), $\{\mu_\Xi, \mu_\Phi, \lambda_\Xi, \lambda_\Phi, \lambda\}$ are real parameters, $\{G_{ij}^{(\ell)}, G_{ij}^{(u)}, G_{ij}^{(d)}, X_{ij}, Y_{ij}, Z_{ij}, W_{ij}\}$ are Yukawa (complex) coupling parameters needed for the fermions to acquire non-trivial masses. In general, these Yukawa couplings set non-diagonal matrices 3×3 , and as usual, the $\tilde{\Phi}$ -field is defined as $\tilde{\Phi} = i \sigma_2 \Phi^*$ to ensure the gauge invariance.

The covariant derivatives of (2.7) act on the Ξ - and Φ -Higgs as follows:

$$\begin{aligned}
 D_\mu \Xi(x) &= \left(\partial_\mu + i J_\Xi g' B_\mu + i K_\Xi g'' C_\mu \right) \Xi(x) \\
 D_\mu \Phi(x) &= \left(\partial_\mu + ig A_\mu^a \frac{\sigma^a}{2} + ig' J_\Phi B_\mu \right) \Phi(x),
 \end{aligned} \tag{2.8}$$

where the J_Φ is the generator of Φ -Higgs corresponding to the $U_R(1)_J$ -subgroup. The Ξ -field is a scalar singlet of $SU_L(2)$, with transformations under $U_R(1)_J \times U(1)_K$. The second Φ -scalar field is a doublet $\Phi = (\phi^{(+)} \ \phi^{(0)})^t$ that turns out in the fundamental representation of $SU_L(2)$, and it also transforms under $U_R(1)_J$ subgroup. We choose the parametrization of the Ξ - and Φ -complex fields as

$$\Xi(x) = \frac{u + \tilde{F}(x)}{\sqrt{2}} e^{i \frac{\eta(x)}{u}}, \quad \Phi(x) = \exp \left[\frac{i}{v} \begin{pmatrix} \chi^3 & \sqrt{2} \chi^- \\ \sqrt{2} \chi^+ & -\chi^3 \end{pmatrix} \right] \begin{pmatrix} 0 \\ \frac{v + \tilde{H}(x)}{\sqrt{2}} \end{pmatrix}, \tag{2.9}$$

where \tilde{F} , \tilde{H} and $\{\eta, \chi^\pm, \chi_3\}$ (four Goldstone bosons) are real functions. The minima of the Higgs potential are given by the non-trivial VEVs $\{u, v\}$ listed below:

$$\begin{aligned}
 \langle \Xi \rangle_0 &= \frac{u}{\sqrt{2}} \simeq \sqrt{-\frac{\mu_\Xi^2}{2\lambda_\Xi}} \left(1 - \frac{\lambda}{4\lambda_\Phi} \frac{\mu_\Phi^2}{\mu_\Xi^2} \right), \\
 \langle \Phi \rangle_0 &= \frac{v}{\sqrt{2}} \simeq \sqrt{-\frac{\mu_\Phi^2}{2\lambda_\Phi}} \left(1 - \frac{\lambda}{4\lambda_\Xi} \frac{\mu_\Xi^2}{\mu_\Phi^2} \right),
 \end{aligned} \tag{2.10}$$

where the following conditions are satisfied : $\mu_\Xi^2 < 0$, $\mu_\Phi^2 < 0$ and $\lambda \ll 1$. It is important to emphasize that, in this Z' -approach, the necessary condition $u \gg v$ between the VEVs

must be satisfied, such that u -scale generates mass for the heavy Z' -boson, while the $v = 246$ GeV is the electroweak scale of the SM.

After the SSBs, the gauge sector is given by

$$\begin{aligned} \mathcal{L}_{mass} = & m_W^2 W_\mu^+ W^{\mu-} + \frac{u^2}{2} \left(J_\Xi g' B_\mu + K_\Xi g'' C_\mu \right)^2 + \frac{v^2}{2} \left(g' J_\Phi B_\mu - \frac{1}{2} g A_\mu^3 \right)^2 \\ & + u \partial_\mu \eta \left(J_\Xi g' B^\mu + K_\Xi g'' C^\mu \right) + m_W \left(\partial_\mu \chi^+ W^{\mu-} + \partial_\mu \chi^- W^{\mu+} \right) \\ & - v \partial_\mu \chi^3 \left(-\frac{1}{2} g A^{\mu 3} + g' J_\Phi B^\mu \right) . \end{aligned} \quad (2.11)$$

The W^\pm -particles are identified as the combination $\sqrt{2} W_\mu^\pm := A_\mu^1 \mp i A_\mu^2$, and its W^\pm -mass is like in GSW-model $m_W = gv/2$. Thus, the charged Goldstone bosons are $\sqrt{2} \chi^\pm := \chi^1 \mp i \chi^2$. The u^2 - and v^2 -mass terms suggest to introduce the orthogonal $SO(2)$ -transformations :

$$\begin{aligned} B_\mu &= \cos \alpha \tilde{Z}'_\mu + \sin \alpha Y_\mu \\ C_\mu &= -\sin \alpha \tilde{Z}'_\mu + \cos \alpha Y_\mu , \\ A_\mu^3 &= \cos \theta_W \tilde{Z}_\mu + \sin \theta_W A_\mu \\ Y_\mu &= -\sin \theta_W \tilde{Z}_\mu + \cos \theta_W A_\mu , \end{aligned} \quad (2.12)$$

where α is a mixing angle, θ_W is known as the Weinberg's angle $\sin^2 \theta_W = 0.22$ ¹, such that it satisfies the parametrization in terms of the fundamental charge $e^2 = 4\pi/137 \simeq 0.09$

$$\begin{aligned} g_Y &= g' \sin \alpha = g'' \cos \alpha , \\ e &= g \sin \theta_W = g_Y \cos \theta_W . \end{aligned} \quad (2.13)$$

The Y^μ is the massless hypercharge gauge field, and \tilde{Z}'^μ sets a massive boson associated with the VEV u -scale. The \tilde{Z} - and \tilde{Z}' are not the fields that represent the physical Z^0 - and Z -bosons yet due to the mixing they present in the Lagrangian. Another diagonalization must be introduced to obtain the physical masses for the condition $u \gg v$. Therefore, the masses of the W , Z and Z' bosons in terms of the fundamental parameters of the model are given by

$$\begin{aligned} m_W &\simeq 80.2 \text{ GeV} \\ M_{Z^0} &\simeq 91.1 \left(1 - \frac{v^2}{32 u^2} \cos^4 \alpha \right) \text{ GeV} , \\ M_{Z'} &\simeq \frac{0.34 u}{\sin \alpha \cos \alpha} \left(1 + \frac{v^2}{32 u^2} \cos^4 \alpha \right) , \end{aligned} \quad (2.14)$$

in which the experimental data have been accounted for [34]. The parameters u -scale and α -angle were not determined in the previous expressions for the Z - and Z' -masses. The ratio between the masses from (2.14) is given by

$$\frac{M_{Z'}}{M_{Z^0}} \simeq \frac{u}{\sin(2\alpha)} \frac{4 \sin \theta_W}{v} \simeq \frac{u}{\sin(2\alpha)} 0.007 \text{ GeV}^{-1} . \quad (2.15)$$

¹We are considering the Weinberg's angle value of $\sin^2 \theta_W = 0.22$, taking into account the radiative corrections of the Electroweak Theory.

The recent papers of the CMS Collaboration point to the hypothetical Z' upper limits that excludes up to 95% confidence level masses below the 2.0 TeV [4]. Therefore, we fix the Z - and Z' -masses as $M_{Z'} = 2.0$ TeV and $M_{Z^0} = 91$ GeV to estimate the ratio of the u -scale by the α -angle, *i. e.*, $u \simeq 2.8 \times \sin(2\alpha)$ TeV. Thereby, the maximum value for the VEV-scale is $u = 2.8$ TeV, when $\alpha = 45^\circ$.

To eliminate the mixed terms $A^{\mu 3} - \chi^3$, $W^\pm - \chi^\pm$ and $Z' - \eta$ in (2.11), we introduce the R_ξ -gauge fixing Lagrangian

$$\begin{aligned} \mathcal{L}_{gf} = & -\frac{1}{2\alpha} (\partial_\mu A^\mu)^2 - \frac{1}{2\gamma} (\partial_\mu Z^\mu - \gamma M_{Z^0} \chi^3)^2 \\ & - \frac{1}{2\beta} \left(\partial_\mu Z'^\mu + \frac{\beta}{2} M_{Z'} \eta - \beta M_{Z'} \sin \theta_W \chi^3 \right)^2 \\ & - \frac{1}{\delta} (\partial_\mu W^{\mu+} - \delta m_W \chi^+) (\partial_\nu W^{\nu-} - \delta m_W \chi^-), \end{aligned} \quad (2.16)$$

where $\{\alpha, \beta, \gamma, \delta\}$ are real parameters. Then, we obtain all the terms with the gauge fields in the renormalizable R_ξ -gauge :

$$\begin{aligned} \mathcal{L}_{gauge}^0 = & -\frac{1}{4} F_{\mu\nu}^2 - \frac{1}{2\alpha} (\partial_\mu A^\mu)^2 \\ & - \frac{1}{2} W_{\mu\nu}^+ W^{\mu\nu-} - \frac{1}{\delta} \partial_\mu W^{\mu+} \partial_\nu W^{\nu-} + m_W^2 W_\mu^+ W^{\mu-} \\ & - \frac{1}{4} Z_{\mu\nu}^2 - \frac{1}{2\gamma} (\partial_\mu Z^\mu)^2 + \frac{1}{2} M_{Z^0}^2 Z_\mu^2 \\ & - \frac{1}{4} Z'_{\mu\nu}{}^2 - \frac{1}{2\beta} (\partial_\mu Z'^\mu)^2 + \frac{1}{2} M_{Z'}^2 Z'_\mu{}^2. \end{aligned} \quad (2.17)$$

The Higgs sector, after the SSBs, is reduced to the scalar fields \tilde{F} and \tilde{H} , whose diagonalization yields the following masses :

$$\begin{aligned} M_H & \simeq \sqrt{2 \lambda_\Phi v^2} \left(1 - \frac{\lambda^2}{8 \lambda_\Xi \lambda_\Phi} \right) = 125 \text{ GeV} \\ M_F & \simeq \sqrt{2 \lambda_\Xi u^2} \left(1 + \frac{\lambda^2}{8 \lambda_\Xi^2} \frac{v^2}{u^2} \right), \end{aligned} \quad (2.18)$$

where $M_F \gg M_H$ as consequence of the VEV-scales condition. The estimation for M_F is

$$1.24 \text{ TeV} < M_F < 3.7 \text{ TeV}, \quad (2.19)$$

in which the maximum value 3.7 TeV correspond to α -angle of 45° .

The sector of interaction of the Z -, Z' -bosons and photon with fermions of the model is

$$\mathcal{L}^{int} = -e Q_{em} \bar{\Psi} A \Psi - e Q_Z \bar{\Psi} Z \Psi - e Q_{Z'} \bar{\Psi} Z' \Psi. \quad (2.20)$$

The electric charge content of model is

$$Q_{em} = I^3 + J + K, \quad (2.21)$$

and the generators Q_Z and $Q_{Z'}$ are defined by the relation

$$\begin{aligned} Q_Z &:= \frac{1}{\sin \theta_W \cos \theta_W} \left(I^3 - Q_{em} \sin^2 \theta_W \right) , \\ Q_{Z'} &:= \frac{2}{\sin(2\alpha) \cos \theta_W} \left(-J + Y \sin^2 \alpha \right) . \end{aligned} \quad (2.22)$$

The interaction of neutrinos-leptons with the W^\pm are, like in the GSW model, reobtained here :

$$\mathcal{L}^{int} = -\frac{g}{\sqrt{2}} \bar{\nu}_{iL} \mathcal{W}^+ \ell_{iL} - \frac{g}{\sqrt{2}} \bar{\ell}_{iL} \mathcal{W}^- \nu_{iL} . \quad (2.23)$$

The Q_Z -generator emerges as in the usual Electroweak Model, but we have here another charge, $Q_{Z'}$, of the interaction between the fermions and the Z' -boson. The values of Q_{em} , I^3 , Y and the primitive charges J and K are summarized in the table below. These values are due to a possible solution for the Abelian (chiral) anomaly to cancel out. Since the model is based on two Abelian subgroups, there are six triangle graphs of the J - and K -symmetries that contribute for the anomaly : J , K , $J^2 K$, JK^2 , J^3 , K^3 . Therefore, the sum of all the charges J and K in the table (1) yield the cancellation of the six triangle graphs. The necessary condition for an anomaly-free model is that the ζ_i -fermions have no electric charge, *i. e.*, $Y = 0$, with $J = -K = +1/2$ for Left-component and $J = -K = -1/2$ for Right-component. Furthermore, the neutrino right-component must also be added to ensure the model to be free from anomalies. Thereby, the Yukawa interactions introduced in the Higgs sector are gauge invariant and it is satisfied by the J - and K -charges in the table. As consequence, ζ_i -fermions do not interact with Z -boson and photon of the SM, but it do interact with the Z' -boson. All the interactions of the Electroweak Standard Model are reproduced in the model. The right-neutrino components and ζ_i -fermions do not interact with the EW Z -boson, as can be verified by the charges. The new interactions of the leptons, neutrinos and the ζ -fermion with the Z' -boson are displayed in what follows :

$$\mathcal{L}_{Z'}^{int} = -g_{Z'} \bar{f} \mathcal{Z}' \left(g_V^f - g_A^f \gamma_5 \right) f , \quad (2.24)$$

where $g_{Z'} := e \csc(2\alpha) \sec \theta_W$, and we rewrite the Z' -interaction with the Left- and Right-components of fermions Ψ from (2.20). Here, the f -fermions means the the fermion fields with no quiral components. Furthermore, we define the coefficients g_V^f and g_A^f as

$$\begin{aligned} g_V^f &= -J^{fL} - J^{fR} + \left(Y^{fL} + Y^{fR} \right) \sin^2 \alpha \\ g_A^f &= -J^{fL} + J^{fR} + \left(Y^{fL} - Y^{fR} \right) \sin^2 \alpha . \end{aligned} \quad (2.25)$$

The f -sum in (2.24) runs to all fermions (no quiral components) of the model. Thereby,


Fields & particles	Q_{em}	I^3	Y	J	K
lepton-left	-1	-1/2	-1/2	0	-1/2
neutrino-left	0	+1/2	-1/2	0	-1/2
lepton-right	-1	0	-1	-1/2	-1/2
neutrino-right	0	0	0	+1/2	-1/2
ζ_i -fermions left	0	0	0	-1/2	+1/2
ζ_i -fermions right	0	0	0	+1/2	-1/2
u-quark-left	+2/3	+1/2	+1/6	0	+1/6
d-quark-left	-1/3	-1/2	+1/6	0	+1/6
u-quark-right	+2/3	0	+2/3	+1/2	+1/6
d-quark-right	-1/3	0	-1/3	-1/2	+1/6
W^\pm -bosons	± 1	± 1	0	0	0
neutral bosons	0	0	0	0	0
Ξ -Higgs	0	0	0	-1	+1
Φ -Higgs	0	-1/2	+1/2	+1/2	0

Table 1. The particle content for the Z' -model candidate at the TeV-scale physics. The J - and K -charges are such that anomalies cancel out.

we list all values of g_V^f and g_A^f following the charges in the table (1) :

$$\begin{aligned}
 g_V^\ell &= -1 + \frac{3}{2} \cos^2 \alpha \quad , \quad g_A^\ell = -\frac{1}{2} \cos^2 \alpha \\
 g_V^\nu &= -1 + \frac{1}{2} \cos^2 \alpha \quad , \quad g_A^\nu = \frac{1}{2} \cos^2 \alpha \\
 g_V^u &= -\frac{1}{2} + \frac{5}{6} \sin^2 \alpha \quad , \quad g_A^u = \frac{1}{2} \cos^2 \alpha \\
 g_V^d &= \frac{1}{2} - \frac{1}{6} \sin^2 \alpha \quad , \quad g_A^d = -\frac{1}{2} \cos^2 \alpha \\
 g_V^\zeta &= 0 \quad , \quad g_A^\zeta = +1 .
 \end{aligned} \tag{2.26}$$

The Z' -vertex useful to perform loop calculations is reads as



$$\Gamma_{Z'}^\mu = -i g_{Z'} \gamma^\mu \left(g_V^f - g_A^f \gamma_5 \right) .$$

3 The mixing between right-neutrinos and ζ_i -fermions

After the SSB takes place, the neutrinos and the ζ -fermion acquire mass terms as displayed below:

$$\begin{aligned}
 \mathcal{L}_{\nu-\zeta}^0 &= \bar{\nu}_i i \not{\partial} \nu_i + \bar{\zeta}_i i \not{\partial} \zeta_i \\
 &- \frac{v}{\sqrt{2}} (X_{ij} \bar{\nu}_{iL} \nu_{jR} + X_{ij}^* \bar{\nu}_{iR} \nu_{jL}) \\
 &- \frac{v}{\sqrt{2}} (Y_{ij} \bar{\nu}_{iL} \zeta_{jR} + Y_{ij}^* \bar{\zeta}_{iR} \nu_{jL}) \\
 &- \frac{u}{\sqrt{2}} (Z_{ij} \bar{\zeta}_{iL} \nu_{jR} + Z_{ij}^* \bar{\nu}_{iR} \zeta_{jL}) \\
 &- \frac{u}{\sqrt{2}} (W_{ij} \bar{\zeta}_{iL} \zeta_{jR} + W_{ij}^* \bar{\nu}_{iR} \zeta_{jL}) .
 \end{aligned} \tag{3.1}$$

It can be cast in the matrix form

$$\mathcal{L}_{\nu-\zeta}^0 = \bar{\chi}_L i \not{\partial} \chi_L + \bar{\chi}_R i \not{\partial} \chi_R - \bar{\chi}_L M \chi_R + \text{h. c.} , \tag{3.2}$$

where $\chi_{L(R)}^t := (\nu_{L(R)} \ \zeta_{L(R)})$ and the mass matrix is given by

$$M = \frac{1}{\sqrt{2}} \begin{pmatrix} v X & v Y \\ u Z & u W \end{pmatrix} . \tag{3.3}$$

Here, X , Y , Z and W are 3×3 matrices of elements $\{X_{ij}, Y_{ij}, Z_{ij}, W_{ij}\}$, and thus, the mass matrix is 6×6 . We introduce the unitary transformation

$$\chi_{L(R)} \longmapsto \chi'_{L(R)} = S \chi_{L(R)} , \tag{3.4}$$

where S is the unitary matrix, $S^\dagger S = \mathbb{1}$. The sector neutrino- ζ -fermion (3.2) is diagonal such that the mass matrix, after the transformation (3.4), satisfies the relation

$$M_D = S M S^\dagger = \begin{pmatrix} M^{(\nu)} & 0 \\ 0 & M^{(\zeta)} \end{pmatrix} . \tag{3.5}$$

To obtain the form of S , we begin with the most general form of a $U(2)$ -matrix

$$S = \begin{pmatrix} e^{i\alpha} \cos \theta & e^{i\beta} \sin \theta \\ -e^{i(\gamma-\beta)} \sin \theta & e^{i(\gamma-\alpha)} \cos \theta \end{pmatrix} , \tag{3.6}$$

with four independent parameters : θ is mixing angle between the families of Right-Neutrino and ζ -fermion, and three phases $\{\alpha, \beta, \gamma\}$. The M_D -matrix is diagonal if the angles $\{\theta, \alpha, \beta\}$ satisfy the relation

$$(v X - u W) \tan(2\theta) = 2 v Y e^{i(\alpha-\beta)} = 2 u Z e^{-i(\alpha-\beta)} , \tag{3.7}$$

and the diagonal of (3.5) is formed by the two 3×3 matrices. In terms of θ -angle, these matrices are given by

$$\begin{aligned} M^{(\nu)} &= \frac{vX}{\sqrt{2}} + \frac{\tan\theta}{2\sqrt{2}}(vX - uW)\tan(2\theta), \\ M^{(\zeta)} &= \frac{uW}{\sqrt{2}} - \frac{\tan\theta}{2\sqrt{2}}(vX - uW)\tan(2\theta). \end{aligned} \quad (3.8)$$

The diagonalization does not depend on the γ -phase, so we can eliminate it. We can also eliminate the θ -mixing angle in (3.9) in terms of u , v , Z and Y ; after some algebra, we obtain the 3×3 mass matrices

$$\begin{aligned} M^{(\nu)} &= \frac{vX + uW - \sqrt{(vX - uW)^2 + 4vuZY}}{2\sqrt{2}} \\ M^{(\zeta)} &= \frac{vX + uW + \sqrt{(vX - uW)^2 + 4vuZY}}{2\sqrt{2}}. \end{aligned} \quad (3.9)$$

It is immediate that, whenever $Y = Z = 0$, we obtain the mass matrices $M^{(\nu)} = vX/\sqrt{2}$ and $M^{(\zeta)} = uW/\sqrt{2}$, respectively. Other way to obtain the same result is rewrite (3.2) in terms of In terms of the Left- and Right-projectors, the χ -spinor is split as $\chi_i^t = (\nu_i \ \zeta_i)^t$; then, the mass matrix (3.3) is rewritten as

$$M_{\nu-\zeta} = \frac{1}{\sqrt{2}} \begin{pmatrix} vX & YvR + ZuL \\ YvL + ZuR & uW \end{pmatrix}, \quad (3.10)$$

where the eigenvalues have the same results as in (3.9).

Here, we are in the Z' -scenario where the u - VEV scale satisfies the condition $u \gg v = 246 \text{ GeV}$; so, we hope that the set of fermions $\zeta_i = \{\zeta_1, \zeta_2, \zeta_3\}$ should describe three particles heavier than any neutrino of the SM, *i. e.*, it is reasonable to consider that $uW \gg vX$. Furthermore, the elements $\{Y_{ij}, Z_{ij}\}$ can be considered weaker coupling constants due to the mixing of right-neutrinos with the sector of ζ_i -fermions. Under these conditions, the mass matrices (3.9) can be written as corrections of

$$\begin{aligned} M^{(\nu)} &\simeq \frac{vX}{\sqrt{2}} \left(\mathbb{1} - \frac{ZY}{XW} \right) \\ M^{(\zeta)} &\simeq \frac{uW}{\sqrt{2}} \left(\mathbb{1} + \frac{vZY}{uW^2} \right) \simeq \frac{uW}{\sqrt{2}}. \end{aligned} \quad (3.11)$$

The neutrino- ζ -fermion sector is now free from mixed terms $\tilde{\nu} - \tilde{\zeta}$ in the basis $\tilde{\chi}_{L(R)}$:

$$\mathcal{L}_{\nu-\zeta}^0 = \bar{\tilde{\nu}}_i i \not{\partial} \tilde{\nu}_i + \bar{\tilde{\zeta}}_i i \not{\partial} \tilde{\zeta}_i - \bar{\tilde{\nu}}_{iL} M_{ij}^{(\nu)} \tilde{\nu}_{jR} - \bar{\tilde{\zeta}}_{iL} M_{ij}^{(\zeta)} \tilde{\zeta}_{jR} + \text{h. c.} . \quad (3.12)$$

Therefore, the mass matrices in (3.12) can be independently diagonalized. The basis $\chi_{L(R)}$ can be written in terms of $\tilde{\chi}_{L(R)}$ by the inverse transformation of (3.4), *i. e.*, the transformation $(\nu_{L(R)}, \zeta_{L(R)}) \mapsto (\tilde{\nu}_{L(R)}, \tilde{\zeta}_{L(R)})$

$$\begin{aligned} \nu_{iL(R)} &= e^{-i\alpha} \cos\theta \tilde{\nu}_{iL(R)} - e^{-i\beta} \sin\theta \tilde{\zeta}_{iL(R)} \\ \zeta_{iL(R)} &= e^{-i\beta} \sin\theta \tilde{\nu}_{iL(R)} + e^{i\alpha} \cos\theta \tilde{\zeta}_{iL(R)}. \end{aligned} \quad (3.13)$$

The interactions $\nu - \ell - W^\pm$ in (2.23) are written in the new basis $(\tilde{\nu}_{L(R)}, \tilde{\zeta}_{L(R)})$, such that the neutrino-lepton- W^\pm interaction is given by

$$\mathcal{L}_{\tilde{\nu}-\ell-W}^{int} = -\frac{g \cos \theta}{\sqrt{2}} \tilde{\nu}_{iL} \mathbb{W}^+ \ell_{iL} - \frac{g \cos \theta}{\sqrt{2}} \bar{\ell}_{iL} \mathbb{W}^- \tilde{\nu}_{iL}, \quad (3.14)$$

in which the α -phase has been absorbed into the left-neutrino field, and $\cos \theta \simeq 1$ for $\theta \ll 1$. Therefore, this changing of basis does not affect the already-known neutrino-lepton- W^\pm interaction of the GSW model. The mass basis of the Dirac neutrinos is introduced via a unitary transformation of the $\nu_{L(R)}$ -fields, then the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) emerges in the interaction (3.14) with only one Dirac phase for Dirac Right-Neutrinos. Furthermore, there are three independent angles in the PMNS matrix to mix the neutrino fields, as usual. If we introduce the unitary transformation $\tilde{\nu}_{L(R)} \mapsto \nu'_{L(R)} = S_{L(R)} \tilde{\nu}_{L(R)}$, the neutrino mass matrix in (3.12) is diagonal in the mass basis $\nu'_{L(R)}$, in which we have $\tilde{\nu}_L M^{(\nu)} \tilde{\nu}_R = \bar{\nu}'_L M_D^{(\nu)} \nu'_R$, where $M_D^{(\nu)}$ is given by

$$M_D^{(\nu)} = S_L \frac{v X}{\sqrt{2}} \left(\mathbb{1} - \frac{ZY}{XW} \right) S_R^\dagger = \begin{pmatrix} M_{\nu'_e} & 0 & 0 \\ 0 & M_{\nu'_\mu} & 0 \\ 0 & 0 & M_{\nu'_\tau} \end{pmatrix}. \quad (3.15)$$

Since the neutrino masses are measured through their oscillations, the transition probabilities depend on the subtraction of the squared masses. In the case of the electron- and muon-neutrinos, this subtraction is $\Delta M_{\nu'_e-\nu'_\mu}^2 := |M_{\nu'_e}^2 - M_{\nu'_\mu}^2| \simeq (7.53 \pm 0.18) \times 10^{-5} \text{ eV}^2$ [40]. Thus, the subtraction of squared coupling constants are extremely weak, $\Delta X_{\nu'_e-\nu'_\mu}^2 := ||X_{\nu'_e}|^2 - |X_{\nu'_\mu}|^2| \simeq 2.5 \times 10^{-27}$. The mixing between τ - and muon- neutrino yield the squared subtraction $\Delta M_{\nu'_\tau-\nu'_\mu}^2 := |M_{\nu'_\tau}^2 - M_{\nu'_\mu}^2| \simeq (2.44 \pm 0.06) \times 10^{-3} \text{ eV}^2$, then we estimate $\Delta X_{\nu'_\tau-\nu'_\mu}^2 := ||X_{\nu'_\tau}|^2 - |X_{\nu'_\mu}|^2| \simeq 8 \times 10^{-26}$. These estimation helps us to obtain the corresponding values of mixed Yukawa coupling constants Y and Z , but we need define a range for ζ_i -masses.

The ζ_i -fermions interact weakly with the leptons according to

$$\mathcal{L}_{\tilde{\zeta}-\ell-W}^{int} = \frac{g \sin \theta}{\sqrt{2}} \tilde{\zeta}_{iL} \mathbb{W}^+ \ell_{iL} + \frac{g \sin \theta}{\sqrt{2}} \bar{\ell}_{iL} \mathbb{W}^- \tilde{\zeta}_{iL}, \quad (3.16)$$

where the β -phase has been absorbed into the $\tilde{\zeta}_{iL}$ -fields. The leptonic sector is diagonalized like in the SM. The mass term is $-v \bar{\ell}_L G^{(\ell)} \ell_R / \sqrt{2}$, that can be diagonalized by means of the unitary transformations $\ell_{L(R)} \mapsto \ell'_{L(R)} = U_{L(R)} \ell_{L(R)}$, in which $U_L^\dagger U_L = U_R^\dagger U_R = \mathbb{1}$, the mass matrix diagonal is $M_D^{(\ell)} = U_L v G^{(\ell)} / \sqrt{2} U_R^\dagger = \text{diag}(M_e, M_\mu, M_\tau)$. Analogously, the diagonalization of ζ_i - mass matrix is performed by another unitary transformation, that we denote by $\tilde{\zeta}_{L(R)} \mapsto \zeta'_{L(R)} = V_{L(R)} \tilde{\zeta}_{L(R)}$, where $V_L^\dagger V_L = V_R^\dagger V_R = \mathbb{1}$, and we obtain the diagonal mass matrix :

$$M_D^{(\zeta)} = V_L \frac{u W}{\sqrt{2}} V_R^\dagger = \begin{pmatrix} M_{\zeta'_1} & 0 & 0 \\ 0 & M_{\zeta'_2} & 0 \\ 0 & 0 & M_{\zeta'_3} \end{pmatrix}. \quad (3.17)$$

So, we can write the interactions (3.16) in the mass basis $\{\zeta'_L, \ell'_L\}$ as below

$$\mathcal{L}_{\zeta'_L-\ell'_L-W}^{int} = \frac{g \sin \theta}{\sqrt{2}} \bar{\zeta}'_{iL} W^+ V_{ij} \ell'_{jL} + \frac{g \sin \theta}{\sqrt{2}} \bar{\ell}'_{iL} W^- V_{ij}^\dagger \zeta'_{jL}, \quad (3.18)$$

where the most general V -unitary matrix $V := V_L U_L^\dagger$ is parameterized by

$$V = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}c_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix}. \quad (3.19)$$

It has the same structure as the Cabibbo-Kobayashi-Maskawa (CKM) matrix : it displays three mixing angles $\{\theta_{12}, \theta_{13}, \theta_{23}\}$ and only one Dirac δ -phase, since we do not introduce ζ_i -Majorana fermions. In (3.19), we simply the sines and cosines of the angles as : $\cos \theta_{ij} = c_{ij}$ and $\sin \theta_{ij} = s_{ij}$. The ζ'_i -masses depend on the u -VEV scale, so it must exhibit a heavier fermion content in comparison with the SM fermions. The recent simulations of CMS-Collaboration point out to dark matter fermion content with mass of order 0.55 TeV [4]. Therefore, we take here $M_{\zeta'_1} = 0.5$ TeV, and if we use $u = 2.8$ TeV, the estimation for $W_{\zeta'_1}$ -Yukawa constant is $W_{\zeta'_1} \simeq 0.28$. The two others heavy fermions ζ'_2 and ζ'_3 can be particles in the mass range of > 0.5 TeV, so we choose $M_{\zeta'_2} = 0.8$ TeV and $M_{\zeta'_3} = 1$ TeV, so the correspondent coupling constants has the values $W_{\zeta'_2} \simeq 0.4$ and $W_{\zeta'_3} \simeq 0.5$, respectively. Using the uncertainty on $\Delta M_{\nu'_e-\nu'_\mu}^2$, the correction to the neutrinos masses give the upper bound

$$\frac{\Delta Y_{\nu'_e-\nu'_\mu} \Delta Z_{\nu'_e-\nu'_\mu}}{\Delta X_{\nu'_e-\nu'_\mu} W_{\zeta'_1}} \lesssim 0.15, \quad (3.20)$$

and $\Delta Y \simeq \Delta Z$, we obtain $\Delta Y_{\nu'_e-\nu'_\mu} \simeq \Delta Z_{\nu'_e-\nu'_\mu} \simeq 8.3 \times 10^{-8}$. Under these conditions and the previous bounds, the θ -mixing angle in (3.7) turns out to be extremely small : $\tan \theta \simeq -9 \times 10^{-8}$.

The interaction of the ζ_i -fermions with the Z' -boson is not affected by the change of basis dictated by the masses. Other important fact is that the interaction (3.18) connects the fermion sector of the SM with a set of fermions candidate to dark sector via W^\pm -bosons. Since $\theta \ll 1$, the θ -angle rules the magnitude of (3.18), *i. e.*, $\sin \theta \simeq \theta \simeq -9 \times 10^{-8}$. This vertex is represented by the diagram below. Therefore, this vertex yields an important

$$\bar{\zeta}'_i \xrightarrow{\quad} \ell'_j \quad \Gamma_{ij}^\mu = -\frac{i g \theta V_{ij}}{2\sqrt{2}} \gamma^\mu (1 - \gamma_5).$$

contribution to ζ_i -fermions magnetic dipole momentum at the one-loop approximation. The weakly coupling constant that emerges here is $g \theta \simeq -6 \times 10^{-9}$. The interaction (3.18) also violates the CP-symmetry due to the δ -phase in the mixing matrix (3.19).

4 The Z' -phenomenology

4.1 The Z' -decay into fermions : $Z' \rightarrow \bar{f} f$

The recent Z' -phenomenology points to the cascade effects at the tree-level using the CMS data for the pp-collision at $\sqrt{s} = 13$ TeV. We will obtain an expression for the Z' -decay width into the any f -fermion of the model. Then, using the previous rules and quantum field-theoretic results, the decay width of Z' into any f -fermion is given by

$$\Gamma(Z' \rightarrow \bar{f} f) = \frac{g_{Z'}^2 M_{Z'}}{24\pi} \left(|g_V^f|^2 + |g_A^f|^2 \right) \sqrt{1 - \frac{4M_f^2}{M_{Z'}^2}} \left(1 - \frac{3M_f^2}{4M_{Z'}^2} \right), \quad (4.1)$$

where $M_{Z'} > 2M_f$, for $f = \{ \ell_i, \nu_i, \zeta_i \}$ or quarks. The decay width into the lepton pair is

$$\Gamma(Z' \rightarrow \bar{\ell}_i \ell_i) = \frac{g_{Z'}^2 M_{Z'}}{24\pi} \left(\frac{1}{2} - 2\sin^2 \alpha + \frac{5}{2} \sin^4 \alpha \right). \quad (4.2)$$

In the neutrino case, the Left- and Right-components provide the following contributions

$$\Gamma(Z' \rightarrow \bar{\nu}_i \nu_i) = \frac{g_{Z'}^2 M_{Z'}}{48\pi} (1 + \sin^4 \alpha). \quad (4.3)$$

Notice that we have used that $M_{Z'} \gg \{ 2M_\ell, 2M_\nu \}$ for leptons and neutrinos. The processes of the Z' -decay can be useful to search the dark matter through the mono-V jets channels associated with the electroweak bosons W or Z . The observation of these final states could be interpreted as a dark matter particle content, that here we identify as the ζ'_i -fermions. The diagram for this effect is illustrated in the figure (1) :




Figure 1. THE Z' -DECAY INTO ANY PAIR $\bar{\zeta}'_i - \zeta'_i$ OF THE ζ'_i -FERMIONS SET. THE CASCADE EFFECT AS A POSSIBLE DARK MATTER DETECTION VIA W - OR Z -MONOJETS.

Therefore, the result for the total decay width of Z' into the ζ_i -family is given by the sum

$$\Gamma(Z' \rightarrow \bar{\zeta}'_i \zeta'_i) = \sum_{i=1}^3 \Gamma(Z' \rightarrow \bar{\zeta}'_i \zeta'_i), \quad (4.4)$$

in which for a particular ζ'_i , it is shown to be given by

$$\Gamma(Z' \rightarrow \bar{\zeta}'_i \zeta'_i) = \frac{g_{Z'}^2 M_{Z'}}{24\pi} \sqrt{1 - \frac{4M_{\zeta'_i}^2}{M_{Z'}^2}} \left(1 - \frac{3M_{\zeta'_i}^2}{4M_{Z'}^2} \right). \quad (4.5)$$

Here, the condition $M_{Z'} > 2 M_{\zeta'_i}$ must be satisfied for any ζ'_i -fermion. Using the previous values $M_{Z'} = 2 \text{ TeV}$ and $M_{\zeta'_1} = 0.55 \text{ TeV}$, the Z' -width decay rate is

$$\Gamma(Z' \rightarrow \bar{\zeta}'_1 \zeta'_1) \simeq \frac{0.018}{\sin^2(2\alpha)} \text{ TeV} . \quad (4.6)$$

In the case of $\alpha = 45^\circ$, the decay width is $\Gamma(Z' \rightarrow \bar{\zeta}'_1 \zeta'_1) \simeq 0.018 \text{ TeV}$, and the Z' -decay time in this process is estimated by ²

$$\tau(Z' \rightarrow \bar{\zeta}'_1 \zeta'_1) = \frac{1}{\Gamma(Z' \rightarrow \bar{\zeta}'_1 \zeta'_1)} \simeq 3.7 \times 10^{-26} \text{ s} . \quad (4.7)$$

The possible Z' -decays into quarks, *i. e.*, $Z' \rightarrow \bar{q} q$, has also a phenomenological analysis at the CMS Collaboration, see [4]. The Z' -decay cases into the first generation, *i. e.*, $Z' \rightarrow \bar{u} u$ and $Z' \rightarrow \bar{d} d$, have the decays width below :

$$\begin{aligned} \Gamma(Z' \rightarrow \bar{u} u) &= \frac{g_{Z'}^2 M_{Z'}}{24\pi} \left(\frac{1}{2} - \frac{4}{3} \sin^2 \alpha + \frac{17}{18} \sin^4 \alpha \right) , \\ \Gamma(Z' \rightarrow \bar{d} d) &= \frac{g_{Z'}^2 M_{Z'}}{24\pi} \left(\frac{1}{2} - \frac{2}{3} \sin^2 \alpha + \frac{5}{18} \sin^4 \alpha \right) , \end{aligned} \quad (4.8)$$

where we have used that $M_{Z'} \gg m_u$ and $M_{Z'} \gg m_d$. Using the α -angle of $\alpha = 45^\circ$, we obtain the decay widths at the GeV-scale :

$$\Gamma(Z' \rightarrow \bar{u} u) \simeq 2 \text{ GeV} , \quad \Gamma(Z' \rightarrow \bar{d} d) \simeq 7 \text{ GeV} . \quad (4.9)$$

4.2 The F -Higgs decays

The Z' -decay into scalars has a phenomenological interest in the study of Z' -resonance to the final four-lepton state [3]. On the other hand, the F -Higgs decays into the leptons pairs $Z' \rightarrow 4 \ell_i$. The process is illustrated at the tree-level as shown below:




Figure 2. THE LEADING ORDER FEYNMAN DIAGRAM FOR THE CASCADE DECAY OF THE Z' -RESONANCE INTO A FOUR-LEPTON FINAL STATE.

Moreover, the decay process $Z' \rightarrow FF$ can not be described by the model due to definition of covariant derivative in the Higgs sector. For example, after the SSB, the

²We have used the conversion formula $1 \text{ TeV} = 1.52 \times 10^{27} \text{ s}^{-1}$ in the natural units $\hbar = c = 1$.

interaction of F -Scalar field with the Z' -boson is given by

$$\mathcal{L}_{F-Z'}^{int} = \frac{M_{Z'}^2}{u} F Z'_\mu Z'^\mu + \frac{1}{2} \frac{M_{Z'}^2}{u^2} F^2 Z'_\mu Z'^\mu, \quad (4.10)$$

where we have the possible vertex $F Z' Z'$ and $F F Z' Z'$. Using the usual rules of QFT, a process possible described by the $F - Z'$ sector is the decay $F \rightarrow Z' Z'$, that has the following decay width :

$$\Gamma(F \rightarrow Z' Z') = \frac{g_{Z'}^2 M_{Z'} M_{Z'}}{16\pi M_F} \left(1 - \frac{4M_{Z'}^2}{M_F^2}\right)^{1/2} \left(3 - \frac{M_F^2}{M_{Z'}^2} + \frac{M_F^4}{4M_{Z'}^4}\right), \quad (4.11)$$

where it is restricted by the condition $M_F > 2 M_{Z'}$.

However, other cascade effects can be described by the $Z' - F$ -interaction. For example, the decay process in which the Z' decays indirectly into two final states of leptons and others two final states of ζ_i -fermions, *i. e.*, $Z' \rightarrow 2\ell_i + 2\zeta'_i$. This process is illustrated in the figure (3). In this case, we have part of the final state described by the decay width




Figure 3. THE LEADING ORDER FEYNMAN DIAGRAM FOR THE CASCADE DECAY OF Z' RESONANCE TO A FOUR-LEPTON FINAL STATE.

(4.2). The other process in the final state is the F -decay into the ζ'_i -fields, which we denote as $F \rightarrow \bar{\zeta}' \zeta'$. The F -scalar field interacts with the ζ'_i -fields by means of the expression

$$\mathcal{L}_{F-\bar{\zeta}'\zeta'}^{int} = - \sum_{i,j=1}^3 \frac{|W_{\zeta'_i}|}{\sqrt{2}} F \bar{\zeta}'_i \zeta'_j. \quad (4.12)$$

Thus, the total decay width is given by

$$\Gamma(F \rightarrow \bar{\zeta}' \zeta') = \sum_{i=1}^3 \Gamma(F \rightarrow \bar{\zeta}'_i \zeta'_i), \quad (4.13)$$

where we obtain the decay width for

$$\Gamma(F \rightarrow \bar{\zeta}'_i \zeta'_i) = M_F \frac{|W_{\zeta'_i}|^2}{8\pi} \left[\left(1 - \frac{2M_{\zeta'_i}^2}{M_F^2}\right) \sqrt{1 - \frac{4M_{\zeta'_i}^2}{M_F^2} - \frac{2M_{\zeta'_i}^2}{M_F^2} \left(1 - \frac{M_{\zeta'_i}^2}{M_F^2}\right)} \right], \quad (4.14)$$

where $M_F > 2 M_{\zeta'_i}$ ($i = 1, 2, 3$).

4.3 The quark-quark scattering into the dark sector

The cascade effect from (1) has the portal for dark matter scenario following the possible scattering $\bar{u}u \rightarrow Z' \rightarrow \bar{\zeta}'_1 \zeta'_1$, in which ζ'_1 has the mass $M_{\zeta'_1} = 0.5$ TeV. It is illustrated at the tree-level in (4).




Figure 4. THE Z' -SCATTERING $\bar{u}u \rightarrow Z' \rightarrow \bar{\zeta}'_1 \zeta'_1$ THAT CAN CONNECT THE STANDARD MODEL WITH THE DARK MATTER CONTENT.

The scattering amplitude of the diagram (4) reads as follows:

$$i\mathcal{M}_{Z'} = \bar{v}(k, t) i g_{Z'} \gamma^\mu (g_V^u - g_A^u \gamma_5) u(p, s) \frac{-i \eta_{\mu\nu}}{(k+p)^2 - M_{Z'}^2} \times \\ \times \bar{u}(p', s') i g_{Z'} \gamma^\nu (g_V^\zeta - g_A^\zeta \gamma_5) v(k', t'). \quad (4.15)$$

We consider the collision in the center-of-mass frame illustrated in the figure (5).




Figure 5. THE COLLISION IN THE CENTER-OF-MASS FRAME FOR THE PROCESS $\bar{u}u \rightarrow \bar{\zeta}'_1 \zeta'_1$.

The squared amplitude (4.15) is given by

$$\frac{1}{4} \sum_{spins} |\mathcal{M}_{Z'}|^2 = \frac{g_{Z'}^4}{4} \left(1 - \frac{4M_{Z'}^2}{s}\right)^{-2} \left[\cos^4 \alpha (1 + \cos^2 \beta) \left(1 - \frac{4M_{\zeta'_1}^2}{s}\right) + \right. \\ \left. + \left(1 - \frac{5}{3} \sin^2 \alpha\right)^2 \left(1 + \sqrt{1 - \frac{4M_{\zeta'_1}^2}{s}} \cos \beta\right) \right], \quad (4.16)$$

where $s := (p + k)^2 = (p' + k')^2$, $\mathbf{p} + \mathbf{k} = \mathbf{p}' + \mathbf{k}' = \mathbf{0}$ and β is the angle between the 3-momentum \mathbf{p}' and \mathbf{k} , or between \mathbf{p} and \mathbf{k}' , in the CM-collision scheme (5). Thus, we get the relation $s = 4E_u^2 = 4E_{\zeta}^2$ by the energy conservation, and the previous amplitude can be written in terms of s -variable. We fix the condition $s > 4M_{\zeta_1}^2$ to insure that (4.16) is real and positive, and we also use that $s \gg 4m_u^2$. Thus, using the standard rules of QFT, the differential cross section for the CM-collision is

$$\frac{d\sigma}{d\Omega}(\bar{u}u \rightarrow \bar{\zeta}_1' \zeta_1') = \frac{\alpha_e^2}{16s \cos^4 \theta_W \sin^4(2\alpha)} \left(1 - \frac{M_{Z'}^2}{s}\right)^{-2} \times$$

$$\times \left[\cos^4 \alpha (1 + \cos^2 \beta) \left(1 - \frac{4M_{\zeta_1}^2}{s}\right) + \left(1 - \frac{5}{3} \sin^2 \alpha\right)^2 \left(1 + \sqrt{1 - \frac{4M_{\zeta_1}^2}{s}} \cos \beta\right) \right]. \quad (4.17)$$

If we fix $\alpha = 45^\circ$, $\sqrt{s} = 13$ TeV, $M_Z = 2$ TeV and $M_{\zeta_1} = 0.5$ TeV, the result (4.17) appears as a function of β . This differential cross section in terms of the β -angle is plotted in the figure (6).




Figure 6. The differential cross section of $\bar{u}u \rightarrow \bar{\zeta}_1' \zeta_1'$ as function of the β -angle. We use here the values for masses $M_{\zeta_1} = 0.5$ TeV and $M_{Z'} = 2$ TeV, the CM-energy as $\sqrt{s} = 13$ TeV, with a mixing angle of $\alpha = 45^\circ$.

Returning to the expression (4.17), the total cross section as a function of the α -mixing angle is given by

$$\sigma(\bar{u}u \rightarrow \bar{\zeta}_1' \zeta_1') = \frac{\pi \alpha_e^2}{4s \cos^4 \theta_W \sin^4(2\alpha)} \left(1 - \frac{M_{Z'}^2}{s}\right)^{-2} \times$$

$$\times \left[\frac{4}{3} \cos^4 \alpha \left(1 - \frac{4M_{\zeta_1}^2}{s}\right) + \left(1 - \frac{5}{3} \sin^2 \alpha\right)^2 \right], \quad (4.18)$$

in which the σ -function is illustrated in the figure (7). When $\alpha = 45^\circ$, the total cross section assumes the value

$$\sigma(\bar{u}u \rightarrow \bar{\zeta}_1' \zeta_1') \simeq 3.76 \times 10^{-7} \text{ TeV}^{-2}. \quad (4.19)$$




Figure 7. The total cross section of $\bar{u}u \rightarrow \bar{\zeta}'_1 \zeta'_1$ as function of the Z' -mass. Here, we adopt the CM-energy values by $\sqrt{s} = 13$ TeV and $\sqrt{s} = 8$ TeV.

5 The MDM for the ζ_i -fermions

In this Section, we investigate the magnetic properties of ζ_i -fermions through its Magnetic Dipole Momentum (MDM). The mixing of ζ_i with the right-neutrino components motivates us to understand if the ζ_i -MDM depends on its mass, as it happens in the case of neutrinos. For a review on the neutrinos' MDMs, go to the references [29–33]. We start off with the field equations in the mixed basis to obtain the so-called Transition Dipole Momenta (TDM). The Dirac equations for the ν_i -neutrinos and ζ_i -fermions from (3.1) in the momentum space are given by

$$\begin{aligned} (\not{p} - m_{\nu_i}) u_{\nu_i}(p) - M_{ij}^{RL} u_{\zeta_j}(p) &= 0 \\ (\not{p} - m_{\zeta_i}) u_{\zeta_i}(p) - M_{ij}^{LR} u_{\nu_j}(p) &= 0, \end{aligned} \quad (5.1)$$

where m_{ν_i} and m_{ζ_i} are the masses if we make the mixed coupling constants $Y_{ij} = Z_{ij} \rightarrow 0$. The functions $u_{\nu_i}(p)$ and $u_{\zeta_i}(p)$ are the wave plane amplitudes of ν_i and ζ_i in the mixed basis, respectively. For simplicity, we have defined the matrices M_{RL} and M_{LR} as combination of Left- and Right-components :

$$\begin{aligned} M_{ij}^{RL} &:= \frac{Y_{ij} v R + Z_{ij} u L}{\sqrt{2}}, \\ M_{ij}^{LR} &:= \frac{Y_{ij} v L + Z_{ij} u R}{\sqrt{2}}. \end{aligned} \quad (5.2)$$

The hermitian conjugate of (5.1) is written below:

$$\begin{aligned} \bar{u}_{\nu_i}(p) (\not{p} - m_{\nu_i}) - \bar{u}_{\zeta_j}(p) M_{ij}^{LR\dagger} &= 0 \\ \bar{u}_{\zeta_i}(p) (\not{p} - m_{\zeta_i}) - \bar{u}_{\nu_j}(p) M_{ij}^{RL\dagger} &= 0. \end{aligned} \quad (5.3)$$

If we substitute p^μ by p'^μ in (5.3), we can combine the hermitian conjugate equations with the equations (5.1) to obtain the following tree-level Gordon decompositions:

$$\begin{aligned} \bar{u}_{\nu_i}(p') \gamma^\mu u_{\nu_i}(p) &= \bar{u}_{\nu_i}(p') \left(\frac{\ell^\mu}{2m_{\nu_i}} + i \frac{\sigma^{\mu\nu} q_\nu}{2m_{\nu_i}} \right) u_{\nu_i}(p) \\ -\bar{u}_{\nu_i}(p') \left(\mu_{ij} \frac{\ell^\mu}{2m_{\nu_i}} + \mu_{ij} i \frac{\sigma^{\mu\nu} q_\nu}{2m_{\nu_i}} + \eta_{ij} \frac{q^\mu \gamma_5}{2m_{\nu_i}} + \eta_{ij} i \frac{\sigma^{\mu\nu} \ell_\nu}{2m_{\nu_i}} \gamma^5 \right) u_{\zeta_j}(p) &+ \text{h. c.}, \end{aligned} \quad (5.4)$$

and

$$\begin{aligned} \bar{u}_{\zeta_i}(p') \gamma^\mu u_{\zeta_i}(p) &= \bar{u}_{\zeta_i}(p') \left(\frac{\ell^\mu}{2m_{\zeta_i}} + i \frac{\sigma^{\mu\nu} q_\nu}{2m_{\zeta_i}} \right) u_{\zeta_i}(p) \\ -\bar{u}_{\zeta_i}(p') \left(\mu_{ij} \frac{\ell^\mu}{2m_{\zeta_i}} + \mu_{ij} i \frac{\sigma^{\mu\nu} q_\nu}{2m_{\zeta_i}} + \eta_{ij} \frac{q^\mu \gamma_5}{2m_{\zeta_i}} + \eta_{ij} i \frac{\sigma^{\mu\nu} \ell_\nu}{2m_{\zeta_i}} \gamma^5 \right) u_{\nu_j}(p) &+ \text{h. c.}, \end{aligned} \quad (5.5)$$

where $q^\mu = p^\mu - p'^\mu$ is the photon's transfer momentum, and $\ell^\mu := p^\mu + p'^\mu$ is the total 4-momentum. These expressions yield the currents of neutrinos with the ζ_i -fermions of the model, written in momentum space, that we refer to as transition terms. The coefficients μ_{ij} and η_{ij} are matrix elements that depend on the Yukawa complex constant coupling and the u -VEV scale :

$$\begin{aligned} \mu_{ij} &= \frac{1}{2\sqrt{2}} \frac{Z_{ij} u + Y_{ij} v}{m_{\zeta_i} + m_{\nu_i}}, \\ \eta_{ij} &= \frac{1}{2\sqrt{2}} \frac{Z_{ij} u - Y_{ij} v}{m_{\zeta_i} + m_{\nu_i}}. \end{aligned} \quad (5.6)$$

Here, if we use that $m_{\zeta_i} \gg m_{\nu_i}$, and $u \gg v$, so we can approximate $Y_{ij} v + Z_{ij} u \approx Z_{ij} u$, and the coefficients μ_{ij} and η_{ij} are approximately equals, $\mu_{ij} \simeq \eta_{ij}$. We also neglect terms of order $Z_{ij}^2 \approx 0$ in relation to the linear terms of Z_{ij} in (5.4) and (5.5). We observe the emergence of TMDM for neutrinos and ζ_i -fermions in both expressions (5.4) and (5.5). If we multiply these currents by $e A_\mu$, and using the representation $q_\nu \rightarrow i \partial_\nu$ for the photon momentum, the terms $\sigma^{\mu\nu} q_\nu$ have the following TMDM for the ν_i -neutrino :

$$\mu_{ij}^{(\nu_i)} = \frac{e \mu_{ij}}{2 m_{\nu_i}} = \frac{e Z_{ij} u}{4\sqrt{2} m_{\nu_i} m_{\zeta_i}} \simeq \frac{Z_{ij}}{4\sqrt{2}} \left(\frac{2.8 \text{ TeV}}{m_{\zeta_i}} \right) \left(\frac{1 \text{ MeV}}{m_{\nu_i}} \right) \mu_B. \quad (5.7)$$

We have a result at tree-level for Dirac neutrinos that depends on the ν_i -neutrino mass and the ζ_i -mass. This result allows us to fix an small estimate for the Z_{ij} coupling constant using the known result $|\mu^{(\nu)}| \lesssim 8 \times 10^{-15} \mu_B$ in the literature [29]. We consider the mass spectrum of $m_{\nu_i} \sim 1 \text{ eV}$ for neutrinos, the mass of $m_{\zeta_1} = 0.5 \text{ TeV}$ for ζ_1 -fermion, and $u = 2.8 \text{ TeV}$. In so doing, the expression (5.7) yields the Z_{ij} coupling constant below :

$$|Z_{ij}| \lesssim 7.5 \times 10^{-21}. \quad (5.8)$$

The TMDM for ζ_i -hidden fermion is given by expression

$$\mu_{ij}^{(\zeta_i)} = \frac{e \mu_{ij}}{2 m_{\zeta_i}} = \frac{Z_{ij}}{4\sqrt{2}} \left(\frac{2.8 \text{ TeV}}{m_{\zeta_i}} \right) \left(\frac{1 \text{ MeV}}{m_{\zeta_i}} \right) \mu_B. \quad (5.9)$$

Therefore, we can estimate the TMDM for the ζ_1 -hidden fermion :

$$|\mu_{ij}^{(\zeta_1)}| \lesssim 1.6 \times 10^{-26} \mu_B, \quad (5.10)$$

where $\mu_B = 3 \times 10^{-7} \text{ eV}^{-1}$ is the Bohr magneton, in natural units $c = \hbar = 1$.

The important contribution for the MDM of ζ_i combines two external lines of ζ'_i with one external line of photon in a one-loop diagram. This possible loop diagram emerges when we work in the mass basis ζ'_i due to the vertex $W^\pm - \zeta' - \ell'$. This vertex depends on the θ -mixing angle, and we believe that it must be a small effect. Obviously, the vertex goes to zero, when $\theta \rightarrow 0$. The vertex at one loop is illustrated in the figure (8).




Figure 8. THE FIRST CONTRIBUTION AT ONE-LOOP FOR THE MDM OF ζ'_i -FERMION COMBINING THE $W^\pm - \zeta' - \ell'$ VERTEX WITH THE EXTERNAL PHOTON.




Figure 9. THE SECOND CONTRIBUTION AT ONE-LOOP FOR THE MDM OF THE ζ'_i -FERMION WITH THE W^+W^- -PHOTON VERTEX OF THE GSW MODEL.

Following the previous rules, the one-loop vertex (8) is represented by the integral

$$\begin{aligned} \Lambda_{(1)ij}^\mu(p, p') &= -\frac{g^2 \theta^2}{8} V_{ik} V_{kj} \int \frac{d^4 k}{(2\pi)^4} \gamma^\alpha (1 - \gamma_5) \frac{i(\not{k} + \not{p}' + m)}{(k + p')^2 - m^2} \times \\ &\times (ie\gamma^\mu) \frac{i(\not{k} + \not{p} + m)}{(k + p)^2 - m^2} \gamma_\alpha (1 - \gamma_5) \frac{-i}{k^2 - m_W^2}. \end{aligned} \quad (5.11)$$

The second contribution comes from the combination of $W^\pm - \zeta' - \ell'$ interaction with the vertex W^\pm -photon of the GSW model. It is illustrated in the figure (9). The Feynman rules yield the momentum space loop integral below

$$\begin{aligned} \Lambda_{(2)ij}^\mu(p, p') &= -\frac{g^2\theta^2}{8} V_{ik}V_{kj} \int \frac{d^4k}{(2\pi)^4} \gamma_\alpha (1 - \gamma_5) \frac{-i}{(k+p')^2 - m_W^2} \times \\ &\times V^{\mu\alpha\beta}(q, k+p', -k-p) \frac{-i}{(k+p)^2 - m_W^2} \frac{i(\not{k} + m)}{k^2 - m^2} \gamma_\beta (1 - \gamma_5). \end{aligned} \quad (5.12)$$

We have used the W^\pm -propagator in the Feynman gauge in the expressions (5.11) and (5.12), $V^{\mu\alpha\beta}$ sets the $W^\pm - \gamma$ vertex following the GSW model rules. Well-known techniques to deal with Feynman integrals are introduced to calculate the finite part of these integrals and, then, the contributions for the MDM of ζ'_i -fermions. The sum of these two contributions is denoted by $\Gamma^\mu = \Lambda_{(1)}^\mu + \Lambda_{(2)}^\mu$, so the finite part of Γ^μ at one-loop is written into the form

$$J_{\zeta'_i(em)}^\mu = \bar{u}_{\zeta'_i} \Gamma_{ij}^\mu u_{\zeta'_j} = \bar{u}_{\zeta'_i}(p') \left[f_1(q^2) \gamma^\mu + f_2(q^2) \gamma_5 \gamma^\mu + f_3(q^2) i \sigma^{\mu\nu} q_\nu + f_4(q^2) q^\mu \gamma_5 \right]_{ij} u_{\zeta'_j}(p), \quad (5.13)$$

where f_1, f_2, f_3 and f_4 are the form factors of the previous diagrams in this order. The current conservation implies that $q_\mu J_{\zeta'_i(em)}^\mu = 0$, then under this condition we obtain the relation $f_2(q^2) = -q^2 f_4(q^2)/2M_{\zeta'_i}$. Thereby, the EM-current of ζ'_i is reduced to expression

$$J_{\zeta'_i(em)}^\mu = \bar{u}_{\zeta'_i} \Gamma_{ij}^\mu u_{\zeta'_j} = \bar{u}_{\zeta'_i}(p') \left[f_1(q^2) \gamma^\mu + f_A(q^2) \gamma_5 (q^\mu \not{q} - q^2 \gamma^\mu) + f_3(q^2) i \sigma^{\mu\nu} q_\nu \right]_{ij} u_{\zeta'_j}(p). \quad (5.14)$$

We have also used here the mass on-shell conditions for ζ'_i -fermions : $\not{p} u_{\zeta'_i}(p) = M_{\zeta'_i} u_{\zeta'_i}(p)$, $\bar{u}_{\zeta'_i}(p') \not{p}' = \bar{u}_{\zeta'_i}(p') M_{\zeta'_i}$ and $p^2 = p'^2 = M_{\zeta'_i}^2$. The amplitudes $\bar{u}_{\zeta'_i}(p')$ and $u_{\zeta'_i}(p)$ stand for the plane wave solutions in the diagonal basis of ζ'_i -fermions. The f_1 -form factor is the contribution to electric charge given by

$$\begin{aligned} f_1(q^2)_{ij} &= \frac{eg^2\theta^2}{32\pi^2} V_{ik}V_{kj} \int_0^1 dx dy dz \delta(x+y+z-1) \times \\ &\times \left[\frac{M_{\zeta'_i}^2 z(4-z) - m^2}{M_{\zeta'_i}^2 z(1-z) - m_W^2 z - m^2 z(1-z) + q^2 xy} \right. \\ &\left. - \frac{M_{\zeta'_i}^2 (1-z)(2+3z)}{M_{\zeta'_i}^2 z(1-z) - m_W^2 (1-z) - m^2 z + q^2 xy} \right]. \end{aligned} \quad (5.15)$$

The second term in (5.14) is known as the anapole (or toroidal momentum) term with the

f_A -form factor that follows:

$$f_A(q^2)_{ij} = \frac{f_A(q^2)_{ij}}{2M_{\zeta'_i}} = \frac{eg^2\theta^2}{32\pi^2} V_{ik}V_{kj} \int_0^1 dx dy dz \delta(x+y+z-1) \times \left[\frac{2+2z-2(x-y)^2}{M_{\zeta'_i}^2 z(1-z) - m_W^2 z - m^2 z(1-z) + q^2 xy} - \frac{3(1-z) - 2(x-y)^2}{M_{\zeta'_i}^2 z(1-z) - m_W^2(1-z) - m^2 z + q^2 xy} \right]. \quad (5.16)$$

The f_3 -form factor is the contribution to the ζ'_i -fermion MDM :

$$f_3(q^2)_{ij} = -\frac{eg^2\theta^2}{32\pi^2} M_{\zeta'_i} V_{ik}V_{kj} \int_0^1 dx dy dz \delta(x+y+z-1) \times \left[\frac{z(1-z)}{M_{\zeta'_i}^2 z(1-z) - m_W^2 z - m^2 z(1-z) + q^2 xy} - \frac{(1-z)(1/2-z)}{M_{\zeta'_i}^2 z(1-z) - m_W^2(1-z) - m^2 z + q^2 xy} \right]. \quad (5.17)$$

The on-shell condition for the external photon imposes $q^2 = 0$. Therefore, all the form factors depend on the masses of ζ'_i -fermion, W^\pm and the lepton mass, where we take $M_{\zeta'_i}, m_W \gg m$ and also consider $M_{\zeta'_i} > m_W$. Under these conditions, the elements of the f_1 -factor are

$$f_1(0)_{ij} = -\frac{eg^2\theta^2}{16\pi^2} V_{ik}V_{kj} \left(1 + \frac{m_W^2}{M_{\zeta'_i}^2} \right) \left[2 - \left(1 - \frac{m_W^2}{M_{\zeta'_i}^2} \right) \ln \left(\frac{M_{\zeta'_i}^2}{m_W^2} - 1 \right) \right]. \quad (5.18)$$

The interaction of A^μ -field with the ζ'_i -current yields the form factor in the limit $q^2 \rightarrow 0$

$$\bar{u}_{\zeta'_i}(p') \Gamma_{ij}^\mu(q) u_{\zeta'_j}(p) A_\mu \stackrel{q^2 \rightarrow 0}{=} -f_3(0)_{ij} \bar{u}_{\zeta'_i}(p') \vec{\sigma} \cdot \vec{\mathbf{B}} u_{\zeta'_j}(p), \quad (5.19)$$

where the transfer momentum is represented in an operator form, $q_\nu \rightarrow i\partial_\nu$. We have considered the EM tensor as $F^{\mu\nu} = (0, \epsilon^{ijk} B^k)$ with an external magnetic field, and $\sigma^{ij} = \epsilon^{ijk} \sigma^k$. We identify the elements of ζ'_i -MDM as $\mu_{Mij}^{(\zeta'_i)} = f_3(0)_{ij}$, thus, the form factor $f_3(0)_{ij}$ can be written in terms of electron's mass and the Bohr magneton :

$$\mu_{Mij}^{(\zeta'_i)} = -\frac{3G_F m_e M_{\zeta'_i}}{4\sqrt{2}\pi^2} \theta^2 V_{ik}V_{kj} \frac{m_W^2}{M_{\zeta'_i}^2} \left[1 - \frac{1}{3} \ln \left(\frac{M_{\zeta'_i}^2}{m_W^2} - 1 \right) + \frac{m_W^2}{M_{\zeta'_i}^2} \right] \mu_B. \quad (5.20)$$

This depends on the $M_{\zeta'_i}$ -mass and on the ratio $m_W/M_{\zeta'_i}$, if we use $m_W = 80 \text{ GeV}$ and $M_{\zeta'_1} = 0.5 \text{ TeV}$, we have $m_W/M_{\zeta'_1} \simeq 0.14$, and $\theta \simeq -9 \times 10^{-8}$. With these values, the MDM for the ζ'_1 -hidden fermion gets the one-loop contribution

$$\mu_{Mij}^{(\zeta'_1)} \simeq 1.2 \times 10^{-21} V_{ik}V_{kj} \mu_B. \quad (5.21)$$

Therefore, the elements ij of ζ'_1 -MDM depend on the V -matrix elements as given in (3.19). For example, for the diagonal elements, $\mu_{Mii}^{(\zeta'_1)} \simeq 1.2 \times 10^{-21} V_{ik} V_{ki} \mu_B$ with the implicit sum runs on the k -index. For $i = 1$, the element $\mu_{M11}^{(\zeta'_1)}$ depends on the cosines of mixing angles θ_{12} and θ_{13} , so we obtain the upper bound

$$\mu_{M11}^{(\zeta'_1)} \simeq 1.2 \times 10^{-21} c_{12}^2 c_{13}^2 \mu_B \lesssim 1.2 \times 10^{-21} \mu_B. \quad (5.22)$$

6 The X-boson scenario at the MeV-scale and below

Contrary to Section II, the X -boson scenario is introduced through the SSB pattern-II mechanism in which the new VEV-scale is at a lower scale with respect to the EW model, $u \ll v = 246 \text{ GeV}$. This SSB pattern-II can be followed in more details in [28]; the original $SU_L(2) \times U_R(1)_J \times U(1)_K$ -symmetry is broken as below :

$$SU_L(2) \times U_R(1)_J \times U(1)_K \xrightarrow{\langle \Phi \rangle_0} U(1)_G \times U(1)_K \xrightarrow{\langle \Xi \rangle_0} U(1)_{em}, \quad (6.1)$$

where we have the VEV scale- u that defines a lightest massive boson, that can describe the physics of the dark-photon with mass bound of $m_{A'} < 8 \text{ GeV}$ [24], the X -boson of mass 17 MeV , or also can describe the para-photon physics at the Sub-eV-scale. We call the $U(1)_G$ -group as the result from the mixing of $SU_L(2) \times U_R(1)_J$. The generator G of $U(1)_G$ is given by $G = I^3 + J$. In both cases, the extra-sector $U(1)_K$ of C^μ couples kinetically with B^μ of $U_R(1)_J$ by means of mixing parameter χ given by

$$\mathcal{L}_{gauge} = -\frac{1}{2} \text{tr} (F_{\mu\nu}^2) - \frac{1}{4} B_{\mu\nu}^2 - \frac{1}{4} C_{\mu\nu}^2 + \frac{\chi}{2} B_{\mu\nu} C^{\mu\nu}. \quad (6.2)$$

Its currently estimated value is $10^{-6} < \chi < 10^{-3}$ for models that discuss hidden photons as dark matter [27].

The sector of fermions is given by the Lagrangian (2.1), but the covariant derivatives must exhibit the millicharged coupling of extra boson C^μ of $U(1)_K$ with the fermions of the Standard Model. Thus, we modify these couplings introducing the parameters ε_Ψ and ε_ζ

$$\begin{aligned} D_\mu \Psi_L &= \left(\partial_\mu + i g A_\mu^a \frac{\sigma^a}{2} + i J_L g' B_\mu + i K_L \varepsilon_\Psi g'' C_\mu \right) \Psi_L, \\ D_\mu \Psi_R &= \left(\partial_\mu + i J_R g' B_\mu + i K_R \varepsilon_\Psi g'' C_\mu \right) \Psi_R, \\ D_\mu \nu_{iR} &= \left(\partial_\mu + i J_R g' B_\mu + i K_R \varepsilon_\nu g'' C_\mu \right) \nu_{iR}, \\ D_\mu \zeta_i &= \left(\partial_\mu + i J_\zeta g' B_\mu + i K_\zeta \varepsilon_\zeta g'' C_\mu \right) \zeta_i, \end{aligned} \quad (6.3)$$

that represent the magnitude of the weaker interaction with the X -boson mentioned in (1.2). Furthermore, the Ξ -Higgs sector has the modified covariant derivative

$$D_\mu \Xi(x) = \left(\partial_\mu + i J_\Xi g' B_\mu + i K_\Xi \varepsilon_\Xi g'' C_\mu \right) \Xi(x). \quad (6.4)$$

The fermion notation is kept like in the previous Z' -model, but the particle content is modified to insure an anomaly-free model:

$$\begin{aligned}
 L_i &= \begin{pmatrix} \nu_e \\ \ell_e \end{pmatrix}_L, \begin{pmatrix} \nu_\mu \\ \ell_\mu \end{pmatrix}_L, \begin{pmatrix} \nu_\tau \\ \ell_\tau \end{pmatrix}_L : \left(\mathbf{2}, -\frac{1}{2}, -\frac{1}{2} \right), \\
 Q_{iL} &= \begin{pmatrix} u \\ d \end{pmatrix}_L, \begin{pmatrix} c \\ s \end{pmatrix}_L, \begin{pmatrix} t \\ b \end{pmatrix}_L : \left(\mathbf{2}, +\frac{1}{6}, +\frac{1}{6} \right), \\
 \ell_{iR} &= \{ e_R, \mu_R, \tau_R \} : (\mathbf{1}, -1, -1), \\
 Q_{iR} &= \{ u_R, c_R, t_R \} : \left(\mathbf{1}, +\frac{2}{3}, +\frac{2}{3} \right), \\
 q_{iR} &= \{ d_R, s_R, b_R \} : \left(\mathbf{1}, -\frac{1}{3}, -\frac{1}{3} \right), \\
 \nu_{iR} &= \{ \nu_{eR}, \nu_{\mu R}, \nu_{\tau R} \} : (\mathbf{1}, 0, 0), \\
 \zeta_{iL} &= \{ \zeta_{1L}, \zeta_{2L}, \zeta_{3L} \} : \left(\mathbf{1}, 0, +\frac{3}{2} \right), \\
 \zeta_{iR} &= \{ \zeta_{1R}, \zeta_{2R}, \zeta_{3R} \} : \left(\mathbf{1}, 0, -\frac{3}{2} \right).
 \end{aligned} \tag{6.5}$$

After the SSBs, the free Lagrangian for the neutral gauge fields reads as follows :

$$\mathcal{L}_{gauge} = -\frac{1}{4} F_{\mu\nu}^2 - \frac{1}{4} \tilde{Z}_{\mu\nu}^2 - \frac{1}{4} \tilde{X}_{\mu\nu}^2 - \frac{\chi W}{2} \tilde{Z}_{\mu\nu} \tilde{X}^{\mu\nu} + \frac{1}{2} m_{\tilde{Z}}^2 \tilde{Z}_\mu^2 + \frac{1}{2} m_{\tilde{X}}^2 \tilde{X}_\mu^2, \tag{6.6}$$

where we have defined

$$\chi_W := \frac{\chi \sin \theta_W}{\sqrt{1 - \chi^2 \cos^2 \theta_W}}, \tag{6.7}$$

and the masses of \tilde{Z}^μ and \tilde{X}^μ are given by

$$m_{\tilde{Z}} = \frac{ev}{\sin 2\theta_W} \quad \text{and} \quad m_{\tilde{X}} = 3|\varepsilon_\Xi|eu. \tag{6.8}$$

For more details on the diagonalization procedure, see [28]. The sector of neutral gauge bosons is composed by two Proca fields with the mixing parameter χ_W in the kinetic term.

Here, the Weinberg mixing angle θ_W satisfies the parametrization

$$e = g'' = g \sin \theta_W = g' \cos \theta_W. \tag{6.9}$$

In the gauge sector, \tilde{Z}^μ acquires a mass due to VEV-scale of $v = 246$ GeV, and \tilde{X}^μ mass is due to lower VEV-scale u . For convenience, we use here the notation \tilde{Z}^μ and \tilde{X}^μ because they are not the physical Z - and X -bosons yet. The real Z -boson will be mixed with the X -boson, in which a full diagonalization of (6.6) implies to shift fields

$$\begin{aligned}
 \tilde{X}^\mu &= X^\mu + \frac{\chi_W Z^\mu}{\sqrt{1 - \chi_W^2}}, \\
 \tilde{Z}^\mu &= \frac{Z^\mu}{\sqrt{1 - \chi_W^2}},
 \end{aligned} \tag{6.10}$$

Fields	Q_{em}	I^3	$Y = J$	K
leptons-left	-1	-1/2	-1/2	-1/2
leptons-right	-1	0	-1	-1
neutrinos-left	0	+1/2	-1/2	-1/2
neutrinos-right	0	0	0	0
ζ_i -fermions left	0	0	0	+3/2
ζ_i -fermions right	0	0	0	-3/2
u-quark-left	+2/3	+1/2	+1/6	+1/6
d-quark-left	-1/3	-1/2	+1/6	+1/6
s-quark-left	-1/3	-1/2	+1/6	+1/6
u-quark-right	+2/3	0	+2/3	+2/3
d-quark-right	-1/3	0	-1/3	-1/3
s-quark-right	-1/3	0	-1/3	-1/3
W^\pm -bosons	± 1	± 1	0	0
neutral bosons	0	0	0	0
Ξ -Higgs	0	0	0	+3
Φ -Higgs	0	-1/2	+1/2	0

Table 2. The simplest anomaly-free particle content for the $U(1)_K$ -dark photon model.

and it gives the masses of the physical Z - and X -bosons with the correction induced by the χ_W -parameter :

$$\begin{aligned}
 M_Z &\simeq \frac{m_{\tilde{Z}}}{\sqrt{1 - \chi_W^2}} \left(1 + \frac{1}{2} \frac{m_X^2}{m_Z^2} \chi_W^2 \right), \\
 M_X &\simeq \frac{m_{\tilde{X}}}{\sqrt{1 - \chi_W^2}} \left(1 - \frac{\chi_W^2}{2} \right).
 \end{aligned} \tag{6.11}$$

The parametrization (6.9) suggests that the electric charge generator can be defined by $Q_{em} = G = I^3 + J$. For convenience, the new Ξ -Higgs carries the charges $J_\Xi = 0$ and $K_\Xi = +3$ to generate the lightest new fermion and also to avoid stable charged matter, see [35, 36]. Here, the hypercharge Y is not given by the sum of the charges J and K . It is so defined by the proper J -generator, *i. e.*, $J = Y$, and the K -generator has independent values of the Y -charges. The simplest charge values are displayed in the 2, and the model in the X-boson scenario is also anomaly-free. All matter content of the SM has $K = 0$. The ζ_i -fermions do not carry electric charge, such that it just sets the K -charge as an example of a hidden charge. Thus, this fact supports the viewpoint of the ζ_i -fermions as dark matter candidates. Other important point is that the gauge symmetry forbids the mixed Yukawa interactions in (2.7), so we must take the coupling constants, $Y_{ij} = Z_{ij} \rightarrow 0$. Therefore, the gauge symmetry and the anomaly cancellation condition lead us to the values for K_{ζ} -charges : $K_{\zeta_L} = -K_{\zeta_R} = +3/2$. The interactions between Z - and X -bosons and any

quiral fermion Ψ of the model are cast below :

$$\mathcal{L}^{int} = -e Q_{em} \bar{\Psi} A \Psi - e Q_Z \bar{\Psi} Z \Psi - e Q_X \bar{\Psi} X \Psi . \quad (6.12)$$

The charge generator Q_X is defined as follows

$$Q_X := +\tilde{\chi} \cos \theta_W Q_{em} + K_\Psi \tilde{\varepsilon}_\Psi , \quad (6.13)$$


where $\tilde{\chi} := \chi/\sqrt{1-\chi^2}$, $\tilde{\varepsilon}_\Psi := \varepsilon_\Psi/\sqrt{1-\chi^2}$. We observe a resulting millicharged current in the interaction of any fermion with the X -boson, whose coupling constant is given by $e Q_X$. Thereby, the interaction of X -boson with the non-chiral components of f -fermions can be written as

$$\mathcal{L}_X^{int} = -e \bar{f} X \left(c_V^f - c_A^f \gamma_5 \right) f , \quad (6.14)$$

where the coefficients c_V^f and c_A^f are defined by

$$c_V^f = Q_X^{fL} + Q_X^{fR} \quad \text{and} \quad c_A^f = Q_X^{fL} - Q_X^{fR} . \quad (6.15)$$

The X - f interaction is not CP-invariant. This shall have some consequence when we are going to calculate the MDM of the charged leptons. Thereby, we list all expressions of c_V^f



$$\bar{f} \longrightarrow f \quad \Gamma_X^\mu = -i e \gamma^\mu \left(c_V^f - c_A^f \gamma_5 \right) .$$

and c_A^f in terms of ε -parameters, following the charges in the table (2) :

$$\begin{aligned} c_V^\ell &= -\chi \cos \theta_W - \frac{1}{2} \left(\frac{1}{2} \varepsilon_{\ell_L} + \varepsilon_{\ell_R} \right) , \\ c_A^\ell &= +\frac{1}{2} \left(-\frac{1}{2} \varepsilon_{\ell_L} + \varepsilon_{\ell_R} \right) , \\ c_V^\nu &= c_A^\nu = -\frac{1}{4} \varepsilon_{\nu_L} , \\ c_V^u &= +\frac{2}{3} \chi \cos \theta_W + \frac{1}{2} \left(\frac{1}{6} \varepsilon_{u_L} + \frac{2}{3} \varepsilon_{u_R} \right) , \\ c_A^u &= +\frac{1}{2} \left(\frac{1}{6} \varepsilon_{u_L} - \frac{2}{3} \varepsilon_{u_R} \right) , \\ c_V^d &= -\frac{1}{3} \chi \cos \theta_W + \frac{1}{2} \left(\frac{1}{6} \varepsilon_{d_L} - \frac{1}{3} \varepsilon_{d_R} \right) , \\ c_A^d &= +\frac{1}{2} \left(\frac{1}{6} \varepsilon_{d_L} + \frac{1}{3} \varepsilon_{d_R} \right) , \\ c_V^\zeta &= +\frac{3}{2} (\varepsilon_{\zeta_L} - \varepsilon_{\zeta_R}) , \quad c_A^\zeta = +\frac{3}{2} (\varepsilon_{\zeta_L} + \varepsilon_{\zeta_R}) . \end{aligned} \quad (6.16)$$

Here, we have 11 ε -parameters so that we can choose the convenient way to describe the phenomenology of the X -boson interacting with the SM matter. We consider the

simplest case for both vector and axial currents type of coupling with leptons in which the parameters are fixed as : $\varepsilon_{\ell_L} = -2\varepsilon_{\ell_R}$, and consequently, $c_V^\ell = -\chi \cos \theta_W$ and $c_A^\ell = -\varepsilon_{\ell_L}/2$. In the quark sector, the X -boson has only axial coupling to avoid an extra flavour mixing of u and s , so it is convenient to fix $c_V^Q = 0$ that implies in the conditions :

$$\begin{aligned}\varepsilon_{u_L} + 2\varepsilon_{d_L} &= 4(\varepsilon_{d_R} - \varepsilon_{u_R}) \\ \varepsilon_{u_L} + 2\varepsilon_{s_L} &= 4(\varepsilon_{s_R} - \varepsilon_{u_R}) \\ \varepsilon_{d_L} - \varepsilon_{s_L} &= 2(\varepsilon_{d_R} - \varepsilon_{s_R}) .\end{aligned}\tag{6.17}$$

The allowed region to explain the 8 Be-anomaly requires for quarks couplings the conditions $c_A^u < 0$, $c_A^d > 0$ and $c_A^d = c_A^s$ that fix the bounds [37]

$$\begin{aligned}10^{-4} &\lesssim |\varepsilon_{d_L} + 2\varepsilon_{d_R}| \lesssim 10^{-3} \\ 10^{-4} &< |\varepsilon_{u_L} - 4\varepsilon_{u_R}| \lesssim 10^{-3} .\end{aligned}\tag{6.18}$$

So, by adopting the range $10^{-4} < |\varepsilon_{\Xi}| < 10^{-3}$, we fix the value $M_X = 17 \text{ MeV}$, and the VEV-scale u is estimated in the range $u \sim 1.9 - 19 \text{ GeV}$. In the Higgs sector, whenever $u \ll v$, the masses of H - and F -scalars fields are

$$\begin{aligned}M_H &\simeq \sqrt{2\lambda_\Phi v^2} \left(1 + \frac{\lambda^2}{8\lambda_\Phi^2} \frac{u^2}{v^2} \right) = 125 \text{ GeV} , \\ M_F &\simeq \sqrt{2\lambda_\Xi u^2} \left(1 - \frac{\lambda^2}{8\lambda_\Phi\lambda_\Xi} \right) ,\end{aligned}\tag{6.19}$$

and the F -mass is estimated in the range $1.9 \text{ GeV} \lesssim M_F \lesssim 19 \text{ GeV}$. Under these conditions, the neutrino and ζ_i -fermions mass matrices are given by the

$$M_{ij}^{(\nu)} = \frac{v X_{ij}}{\sqrt{2}} \quad \text{and} \quad M_{ij}^{(\zeta)} = \frac{u W_{ij}}{\sqrt{2}} ,\tag{6.20}$$

in which $M_{ij}^{(\zeta)}$ must have lighter elements in the X -Boson scenario if compared with the corresponding case of the Z' scenario. For example, the estimate here satisfies the Treiman-Gunn lower bound of $> 1.5 \text{ KeV}$ for the mass of a dark matter candidate, see [41]. At this stage, the diagonalization procedure is like in the early case of the Z' -scenario. The diagonalization of mass matrix $M^{(\zeta)}$ is carried out by a unitary transformation such that the elements of the diagonal matrix are the masses $M_{\zeta'_i}$ ($i = 1, 2, 3$). If we impose the previous coupling constants $W_{\zeta'_1} = 0.28$, $W_{\zeta'_2} = 0.4$ and $W_{\zeta'_3} = 0.5$, we obtain the following masses for the light-fermions $M_{\zeta'_1} \simeq 0.3 - 3.8 \text{ GeV}$, $M_{\zeta'_2} \simeq 0.5 - 5.3 \text{ GeV}$ and $M_{\zeta'_3} = 0.6 - 6.6 \text{ GeV}$.

7 The X -boson phenomenology

7.1 The X -boson decays processes

The X -boson phenomenology points out to the decay $X \rightarrow e^+ e^-$ as one of main processes. Furthermore, we will obtain here the others possible processes of X -decay into the SM

fermions. Thereby, using the QFT rules for X -interaction in (6.14), the decay width for the process $X \rightarrow \bar{f} f$ is given by

$$\Gamma(X \rightarrow \bar{f} f) = \frac{\alpha M_X}{6} \left(|c_V^f|^2 + |c_A^f|^2 \right) \sqrt{1 - \frac{4M_f^2}{M_X^2}} \left(1 - \frac{3M_f^2}{4M_X^2} \right), \quad (7.1)$$

with the condition $M_X > 2M_f$. It is clear that, for X -boson of $M_X = 17$ MeV some decays processes are forbidden according to the result of (7.1), as for example, the X -decays into the muon and tau-particles in the leptons sector, and X -decays into the heavier quarks.

The decay processes into the $X \rightarrow e^+ e^-$ and $X \rightarrow \bar{\nu}_e \nu_e$ yield the X -decay widths

$$\begin{aligned} \Gamma(X \rightarrow e^+ e^-) &= \frac{\alpha M_X}{6} \left(\chi^2 \cos^2 \theta_W + \frac{\varepsilon_{eL}^2}{4} \right), \\ \Gamma(X \rightarrow \bar{\nu}_e \nu_e) &= \frac{\alpha M_X}{48} \varepsilon_{\nu L}^2, \end{aligned} \quad (7.2)$$

in which we have used $M_X \gg \{M_e, M_\nu\}$. The KLOE-2 collaboration sets the limit $|\varepsilon_{eL}| \lesssim 4 \times 10^{-3}$ [42], and neutrino constraint is in the scattering $\bar{\nu}_e e \rightarrow \bar{\nu}_e e$, that fixes the bound $|\varepsilon_{\nu L}| < 2.2 \times 10^{-5}$ [22]. The decays widths (7.2) are estimated by the results below

$$\begin{aligned} \Gamma(X \rightarrow e^+ e^-) &= 8.2 \times 10^{-8} \text{ MeV}, \\ \Gamma(X \rightarrow \bar{\nu}_e \nu_e) &= 10^{-12} \text{ MeV}. \end{aligned} \quad (7.3)$$

The allowed decay process in the quarks sector has the decay width

$$\Gamma(X \rightarrow \bar{u} u) = \frac{\alpha M_X}{864} |\varepsilon_{uL} - 4\varepsilon_{uR}|^2 \simeq 10^{-12} - 10^{-10} \text{ MeV}. \quad (7.4)$$

7.2 The X -boson scattering

A possible scenario for the X -boson scattering lies in its connection with the ζ_i -fermions. The recent bound for dark photons A' of mass $m_{A'} < 8$ GeV in the process $e^+ e^- \rightarrow \gamma A'$ could establish a connection with a dark matter scenario through the interaction of A' with a dark sector. Therefore, we propose the study of the scattering $e^+ e^- \rightarrow X \rightarrow \bar{\zeta}'_1 \zeta'_1$ in which the X -boson acts like a dark photon, and we choose the mass of ζ'_1 as $M_{\zeta'_1} = 1$ GeV. The process is illustrated in the figure (10).

The amplitude for this scattering is depicted in the figure (11).

$$\begin{aligned} i\mathcal{M}_X &= \bar{v}(k, t) i e \gamma^\mu (c_V^e - c_A^e \gamma_5) u(p, s) \frac{-i \eta_{\mu\nu}}{(k+p)^2 - M_X^2} \times \\ &\quad \times \bar{u}(p', s') i e \gamma^\nu (c_V^\zeta - c_A^\zeta \gamma_5) v(k', t'). \end{aligned} \quad (7.5)$$




Figure 10. THE X -DECAY SCATTERING INTO A POSSIBLE DARK MATTER SCENARIO WITH THE ζ'_1 -FERMION OF MASS $M_{\zeta'_1} = 1$ GeV.




Figure 11. THE ELECTRON-POSITRON SCATTERING $e^+ e^- \rightarrow X \rightarrow \bar{\zeta}'_1 \zeta'_1$ THAT CAN CONNECT THE SECTOR OF LEPTONS WITH THE DARK MATTER CONTENT VIA HIDDEN PHOTON OR X -BOSON.

The differential cross section for the CM collision is

$$\begin{aligned}
 \frac{d\sigma}{d\Omega}(e^+ e^- \rightarrow \bar{\zeta}'_1 \zeta'_1) &= \frac{9\alpha^2}{2s} \left(1 - \frac{M_X^2}{s}\right)^{-2} \times \\
 &\times \left\{ \left[\varepsilon_{\zeta_L} \left(\chi \cos \theta_W + \frac{\varepsilon_{eL}}{2} \right) - \varepsilon_{\zeta_R} \left(\chi \cos \theta_W - \frac{\varepsilon_{eL}}{2} \right) \right]^2 \times \right. \\
 &\times \left[3 + 2 \sqrt{1 - \frac{4M_{\zeta'_1}^2}{s}} \cos \beta + \left(1 - \frac{4M_{\zeta'_1}^2}{s}\right) \cos^2 \beta \right] + \\
 &\left. + \chi^2 \cos^2 \theta_W (\varepsilon_{\zeta_L} - \varepsilon_{\zeta_R})^2 - \frac{\varepsilon_{eL}^2}{4} (\varepsilon_{\zeta_L} + \varepsilon_{\zeta_R})^2 \right\}, \quad (7.6)
 \end{aligned}$$

where $s > 4M_{\zeta'_1}^2$. If we use the ζ'_1 -mass as $M_{\zeta'_1} = 1$ GeV, the condition on the CM-energy is $\sqrt{s} > 2$ GeV. The simplest case of (7.7) assumes that the ε -parameters of the left-components are of the same order as the right-components, *i. e.*, $\varepsilon_{\zeta_L} \simeq \varepsilon_{\zeta_R}$; so, the

previous expression is reduced to

$$\begin{aligned} \frac{d\sigma}{d\Omega}(e^+ e^- \rightarrow \bar{\zeta}'_1 \zeta'_1) &= \frac{9\alpha^2}{s} \left(1 - \frac{M_X^2}{s}\right)^{-2} \varepsilon_{e_L}^2 \varepsilon_{\zeta_L}^2 \times \\ &\times \left[1 + \sqrt{1 - \frac{4M_{\zeta'_1}^2}{s}} \cos \beta + \frac{1}{2} \left(1 - \frac{4M_{\zeta'_1}^2}{s}\right) \cos^2 \beta \right], \end{aligned} \quad (7.7)$$

The differential cross section (7.7) is illustrated in figure (12). The corresponding total




Figure 12. The differential cross section of $e^+ e^- \rightarrow \bar{\zeta}'_1 \zeta'_1$ as function of the β -angle. We use the masses $M_{\zeta'_1} = 1 \text{ GeV}$ and $M_X = 17 \text{ MeV}$, the CM-energy as $\sqrt{s} = 3 \text{ GeV}$.

cross section is obtained upon integration over the solid angle:

$$\sigma(e^+ e^- \rightarrow \bar{\zeta}'_1 \zeta'_1) = \frac{42\pi \alpha^2}{s} \frac{\varepsilon_{e_L}^2 \varepsilon_{\zeta_L}^2}{(1 - M_X^2/s)^2} \left(1 - \frac{4M_{\zeta'_1}^2}{s}\right). \quad (7.8)$$

Using the previous masses, the C.M. energy of $\sqrt{s} = 3 \text{ GeV}$ and $|\varepsilon_{e_L}| \lesssim 4 \times 10^{-3}$, the σ -function has the following estimate

$$\sigma(e^+ e^- \rightarrow \bar{\zeta}'_1 \zeta'_1) = 1.2 \times 10^{-11} \varepsilon_{\zeta_L}^2 \text{ GeV}^{-2}. \quad (7.9)$$

7.3 The electron and muon $g-2$ factors

The $g-2$ factor is an important experimental physical quantity to estimate the ε -parameters of the leptonic sector in interaction with the X -boson. With those, we can obtain the form factors associated to the vertex ℓ^\pm -photon including the correction of the X -boson propagator. This correction is illustrated in the figure (13). With that, we write the QED vertex of lepton-photon corrected by the diagram (13) as the sum : $\Gamma^\mu(p, p') =$




Figure 13. THE ONE-LOOP CORRECTION OF THE X -BOSON TO THE QED VERTEX DIAGRAM.

$\Gamma_{QED}^\mu(p, p') + \Lambda^\mu(p, p')$. The diagram is represented by the loop 4-momentum integral:

$$\begin{aligned} \Lambda^\mu(p, p') = & \int \frac{d^4k}{(2\pi)^4} i e \gamma_\alpha \left(c_V^\ell - c_A^\ell \gamma_5 \right) \frac{i (\not{k} + \not{p}' + m)}{(k + p')^2 - m^2} (i e \gamma^\mu) \times \\ & \times \frac{i (\not{k} + \not{p} + m)}{(k + p)^2 - m^2} i e \gamma^\alpha \left(c_V^\ell - c_A^\ell \gamma_5 \right) \frac{-i}{k^2 - M_X^2}, \end{aligned} \quad (7.10)$$

where m stands for the electron or muon mass. Obviously, the integral is divergent in the ultraviolet regime, but we are interested in its finite part. To simplify the previous integral, we use here that the vector coefficient is of the same order of the axial coefficient for Leptons, *i. e.*, $c_V^\ell \simeq c_A^\ell = -\chi \cos \theta_W = -\varepsilon_{\ell_L}/2 = +\varepsilon_{\ell_R}$. The integral can be calculated by using the known methods of Feynman integrals; then, the finite part of the vertex can be extracted and gives the leptonic current $\bar{u}\Gamma^\mu(q)u$, that can be written as

$$J_{em}^\mu = \bar{u}(p')\Gamma^\mu(q)u(p) = \bar{u}(p') \left[F_1(q^2) \gamma^\mu + F_2(q^2) \gamma^5 \gamma^\mu + F_3(q^2) i \frac{\sigma^{\mu\nu} q_\nu}{2m} + F_4(q^2) \frac{q^\mu \gamma^5}{2m} \right] u(p), \quad (7.11)$$

where q^μ stands for the momentum transfer and $\ell^\mu = p^\mu + p'^\mu$, as before. Here, we obtain the QED form factors with the correction of ε -parameters

$$\begin{aligned} F_1(q^2) = & F_1^{QED}(q^2) + \frac{\alpha \varepsilon_{\ell_L}^2}{4\pi} \int_0^1 dx dy dz \delta(x + y + z - 1) \times \\ & \times \frac{4z - 1 - z^2 - (1-x)(1-y)q^2/m^2}{(1-z)^2 + zM_X^2/m^2 - xyq^2/m^2}, \end{aligned} \quad (7.12)$$

and

$$\begin{aligned} F_3(q^2) = & \frac{\alpha}{2\pi} \int_0^1 dx dy dz \delta(x + y + z - 1) \frac{2z(1-z)}{(1-z)^2 - xyq^2/m^2} \\ & - \frac{\alpha \varepsilon_{\ell_L}^2}{16\pi^2} \int_0^1 dx dy dz \delta(x + y + z - 1) \frac{2z(1-z)}{(1-z)^2 + zM_X^2/m^2 - xyq^2/m^2}. \end{aligned} \quad (7.13)$$

The F_2 form factor is associated with the CP-violation of the X -fermion coupling. The conservation current yields us the condition : $F_2(q^2) = -q^2 F_4(q^2)/4m^2$; in which the

current (7.11) can be rewritten as

$$J_{em}^\mu = \bar{u}(p')\Gamma^\mu(q)u(p) = \bar{u}(p') \left[F_1(q^2) \gamma^\mu + F_4(q^2) \gamma^5 (q^\mu \not{q} - q^2 \gamma^\mu) + F_3(q^2) i \frac{\sigma^{\mu\nu} q_\nu}{2m} \right] u(p), \quad (7.14)$$

Explicitly, we have the anapole term with the F_A -form factor given by

$$F_A(q^2) = \frac{F_4(q^2)}{2m} = \frac{\alpha \varepsilon_{\ell L}^2}{8\pi} \int_0^1 dx dy dz \delta(x+y+z-1) \frac{1+z-(x-y)^2}{(1-z)^2 + zM_X^2/m^2 - xyq^2/m^2}. \quad (7.15)$$

The interesting analysis now is the situation where $q^2 = 0$ for the external photon in the form factor F_3 . The first term yields the $(g-2)_\ell$ factor for the lepton with the correction of the c_V^ℓ coefficient, which in turn, depends on ε_ℓ -parameters. For the electron case, the $F_3(0)$ is given by

$$F_3^e(0) \simeq \frac{\alpha}{2\pi} \left(1 - \frac{\varepsilon_{eL}^2}{12\pi} \frac{m_e^2}{M_X^2} \right), \quad (7.16)$$

where we have used that $M_X \gg m_e$. Using the $(g-2)_e$ uncertainty in the literature, we obtain the upper bound on ε_{eL} -parameter

$$|\varepsilon_{eL}| \lesssim 5.3 \times 10^{-5}, \quad (7.17)$$

and, thus, we estimate the other parameters $|\varepsilon_{eR}| \lesssim 2.6 \times 10^{-5}$ and $|\chi| \lesssim 3 \times 10^{-5}$. For the muon, we obtain

$$F_3^\mu(0) \simeq \frac{\alpha}{2\pi} \left(1 - \frac{\varepsilon_{\mu L}^2}{8\pi} \right), \quad (7.18)$$

and using the corresponding $(g-2)_\mu$ uncertainty, we obtain the following upper bound

$$|\varepsilon_{\mu L}| \lesssim 8.7 \times 10^{-5}. \quad (7.19)$$

8 Concluding Comments

We have made efforts to discuss a model with an extra $U(1)_K$ -factor that may describe a possible scenario of a Particle Physics beyond the Standard Model (SM). The proposal is based on the gauge group $SU_L(2) \times U_R(1)_J \times U(1)_K$, where the extra $U(1)_K$ -factor introduces a (new) massive and neutral vector boson. The mass of the latter is generated upon a spontaneous symmetry breaking mechanism that defines an extra VEV scale, beyond that VEV of 246 GeV associated with the usual Higgs field of SM. In our model, the Higgs sector displays two scalar fields and the gauge symmetry allows interactions between them. This mechanism can be introduced in two ways : in the first case, the SSB pattern-I takes place through a VEV scale- u , where $u > 246$ GeV, and, as consequence, we fix a mass for the hypothetical Z' -boson around 2 TeV. Next, the SM Higgs acquires its VEV of 246 GeV,

breaks the electroweak symmetry to give the masses for W^\pm and Z . The sequence of SSB mechanism is as follows:

$$SU_L \times U_R(1)_J \times U(1)_K \xrightarrow{\langle \Xi \rangle_0} SU_L(2) \times U_Y(1) \xrightarrow{\langle \Phi \rangle_0} U_{em}(1) .$$

The result for the mass of Z' is estimated by the ATLAS and CMS Collaborations as a possible particle at the TeV-scale. The maximum VEV scale for $M_{Z'} = 2 \text{ TeV}$ is $u = 2.8 \text{ TeV}$, and it allows an estimation of the mass of the new Higgs within the range $1.2 \text{ TeV} < M_F < 3.7 \text{ TeV}$. Furthermore, the subgroup $U(1)_K$ also introduces a new family of fermions in the TEV-scale, which we call ζ_i -fermions ($i = 1, 2, 3$), that could be candidates to the dark matter particles. It is a set of neutral heavy fermions associated with the VEV scale of 2.8 TeV and they guarantee that model be free from the chiral anomaly. We use the masses for ζ_i in the range of $0.5 - 1 \text{ TeV}$, motivated by recent simulations of dark matter fermions scenario in the CMS-Collaboration. Furthermore, these new fermions also mix with a Dirac's right-neutrino component. This new mixing motivated us to investigate the MDM of the ζ_i -fermions, and the transition MDMs of the ζ_i mixed with neutrinos. Since ζ_i are neutral fermions, their MDMs depend on their masses and also on the mixing with the right-neutrinos. At one-loop, the diagonal ζ_1 -MDM for mass of 0.5 TeV is estimated by $\mu_M^{(\zeta_1)} = 1.2 \times 10^{-21} \mu_B$.

In the second scenario, the SSB mechanism pattern-II is introduced to break the composite subgroup $SU_L(2) \times U_R(1)_J$ in the VEV scale of the SM, 246 GeV , to yield the usual Z -mass, and so, a further SBB defines the VEV scale- u , $u < 246 \text{ GeV}$. In this second situation, we have the scenario of a lighter extra gauge boson; it may describe a phenomenology of elementary particle physics at a lower energy scale. The scheme for the pattern-II SSB is displayed below:

$$SU_L \times U_R(1)_J \times U(1)_K \xrightarrow{\langle \Phi \rangle_0} U(1)_G \times U(1)_K \xrightarrow{\langle \Xi \rangle_0} U(1)_{em} .$$

This proposal may be discussed in connection with the light X -Boson, dark photon or para-photon phenomenology, at MeV-scale or lower, which can be related to dark matter particles. This context includes the recent description of a new boson needed to explain the excited 8-Beryllium nuclear decay. In this case, the mass of the X -Boson is fixed at the mass scale of $m_X = 17 \text{ MeV}$. Therefore, the results of SSB lead us to the VEV scale of $u \simeq 1.9 - 19 \text{ GeV}$ that must be origin of a hidden Higgs within the range mass $1.9 \text{ GeV} \lesssim M_F \lesssim 19 \text{ GeV}$. In this context, the ζ_i -fermions are interpreted as light particles with masses in the range of $0.3 - 6.6 \text{ GeV}$. It can be a candidate to dark matter constituent. After the SSB pattern-II, we study the phenomenology of the X -Boson through its decay into electron, neutrino and light u -quark, and the scattering $e^+ e^- \rightarrow X \rightarrow \bar{\zeta}'_1 \zeta'_1$, with $M_{\zeta'_1} = 1 \text{ GeV}$; this can connect the SM with the dark sector with a hidden photon mediating the interaction. To conclude, we calculate the correction to the Quantum Electrodynamics (QED) vertex due to axial interaction of X -Boson with the leptons of the SM; in particular, for the electron and muon cases. The results impose the bounds $|\varepsilon_{eL}| \lesssim 5.3 \times 10^{-5}$ and $|\varepsilon_{\mu L}| \lesssim 8.7 \times 10^{-5}$ for the electron and muon, respectively, which measures the magnitude of the weak interaction of the electron/muon with the X -Boson. The Electric Dipole

Momentum of the charged leptons must emerge at higher loop orders, in both cases of the Z' and X -boson scenarios. This is an issue we are now investigating and we shall report on it in a further paper.

A unification scheme including W' - and Z' -bosons is in the framework of an left-right symmetric model $SU_L(2) \times SU_R(2) \times U(1)_J \times U(1)_K$ may also be the subject of a further investigation to be pursued in the presence of dark matter fermion content.

References

- [1] ATLAS Collaboration, *Phys. Rev. D* **90**, no. 5, 052005 (2014).
- [2] The CMS Collaboration, *JHEP* **1504**, 025 (2015).
- [3] The CMS Collaboration, *Phys. Lett. B*, **2017**, 773, 563.
- [4] The CMS Collaboration, *J. High Energy Phys.*, **2017**, 07, 014.
- [5] The CMS Collaboration, *Search for new physics in final states with an energetic jet or a hadronically decaying W or Z boson and transverse momentum imbalance at $\sqrt{s} = 13$ TeV*, Arxiv : 1712.02345.
- [6] Bogdan A. Dobrescu and Zhen Liu, *JHEP* (2015) 2015 : 118.
- [7] Bogdan A. Dobrescu and Zhen Liu, *Phys. Rev. Lett.* **115** (2015) 211802.
- [8] P. Coloma, B. A. Dobrescu and J. Lopez-Pavon, *Phys. Rev. D* **92** (2015) 115023.
- [9] Bogdan A. Dobrescu and Patrick J. Fox, *JHEP* (2016) 2016 : 047.
- [10] P. S. Bhupal Dev and R. N. Mohapatra, *Phys. Rev. Lett.* **115** (2015) 181803.
- [11] S. Patra, W. Rodejohann and C. E. Yaguna *JHEP* (2016) 2016 : 076.
- [12] Pei-Hong Gu and Rabindra N. Mohapatra, *Phys. Rev. D* **96**, 055011 (2017).
- [13] P. S. Bhupal Dev, Rabindra N. Mohapatra and Yongchao Zhang, *Journal of Physics : Conf. Series* **873** (2017) 012029.
- [14] Paul Langacker, *Reviews of Modern Physics*, **81** (2009) 1199-1228.
- [15] Shinya Kanemura, Osamu Seto and Takashi Shimomura, *Phys. Rev. D* **84** (2011) 016004.
- [16] E. Accomano *et al*, *JHEP* (2016) 2016 : 86.
- [17] A. J. Krasznahorkay *et al*, *Phys. Rev. Lett.* **116** (2016) 042501.
- [18] J. L. Feng *et al*, *Phys. Rev. Lett.* **117** (2016) 071803.
- [19] Pei-Hong Gu , Xiao-Gang He, *Nuclear Physics B* **919** (2017) 209-217.
- [20] J. L. Feng *et al*, *Phys. Rev. D* **95**, 035017 (2017).
- [21] Yonathan Kahn, Gordan Krnjaic, Siddharth Mishra-Sharma, Tim M. P. Tait, *J. High Energy Phys.*, **2017**, 05, 002.
- [22] S. Bilmis, I. Turan, T.M. Aliev, M. Deniz, L. Singh, H.T. Wong, *Phys. Rev. D* **2015**, 92, 033009.
- [23] Teppei Kitahara and Yasuhiro Yamamoto, *Phys. Rev. D* **95**, 015008 (2017).
- [24] J. P. Lees *et al.* (BaBar Collaboration), *Phys. Rev. Lett.* **119** (2017) 131804.

- [25] M. Ahlers, H. Gies, J. Jaeckel, J. Redondo, and A. Ringwald, *Phys. Rev. D* **76** (2007) 115005.
- [26] J. Jaeckel, A. Ringwald, *Phys. Lett. B* **659** (2008) 509-514.
- [27] P. Arias, J. Jaeckel, J. Redondo, and A. Ringwald, *Phys. Rev. D* **82** (2016) 115018.
- [28] M. J. Neves and J. A. Helayël-Neto, *Annalen Der Physik* **2017**, 1700112.
- [29] Nicole F. Bell *et. al.*, *Phys. Rev. Lett.* **95** (2005) 151802.
- [30] Nicole F. Bell *et. al.*, *Phys. Lett. B* 642 (2006) 377-383.
- [31] Kaushik Bhattacharya and Palash B Pal, *Proc Indian Natn Sci Acad*, **70**, A No. 1, January 2004, pp. 145161.
- [32] A. B. Balantekin, AIP Conference Proceedings **847**, 128 (2006).
- [33] Alexander Studenikin, *Journal of Physics : Conference Series* **718** (2016) 062076.
- [34] C. Patrignani *et. al.* (Particle Data Group), *Chin. Phys. C* 40 100001 (2016).
- [35] M. Duerr and P. Fileviez Perez, *Phys. Lett. B* **732** 101 (2014).
- [36] M. Duerr and P. Fileviez Perez, *Phys. Rev. D* **91** (2015) 095001.
- [37] Jonathan Kozaczuk, David E. Morrissey, S.R. Stroberg, *Phys. Rev. D* **2017**, *95*, 115024.
- [38] H. Davoudiasl, H. -S. Lee and W. J. Marciano, *Phys. Rev. D* **89**, 095006 (2014).
- [39] M. Raggi *et al.* (NA48/2 Collaboration), NA48/2 studies of rare decays, *Nuovo Cimento Soc. Ital. Fis.* **38**, 132 (2016).
- [40] KamLAND Collaboration, *Phys. Rev. Lett.*, **2005**, *94*, 081801.
- [41] Subinoy Das, Kris Sigurdson, *Phys. Rev. D* **2012**, *85*, 063510.
- [42] KLOE Collaboration, *Phys. Lett. B*, **2016**, *757*, 362.