

A PRIORI ESTIMATES FOR THE FREE-BOUNDARY EULER EQUATIONS WITH SURFACE TENSION IN THREE DIMENSIONS

MARCELO M. DISCONZI
Department of Mathematics
Vanderbilt University
Nashville, TN 37240, USA

IGOR KUKAVICA
Department of Mathematics
University of Southern California
Los Angeles, CA 91107, USA

ABSTRACT. We derive a priori estimates for the incompressible free-boundary Euler equations with surface tension in three spatial dimensions. Working in Lagrangian coordinates, we provide a priori estimates for the local existence when the initial velocity, which is rotational, belongs to H^3 , with some additional regularity on the normal component of the initial velocity. This lowers the requirement on the regularity of initial data in the Lagrangian setting. Our methods are direct and involve three key elements: estimates for the pressure, the boundary regularity provided by the mean curvature, and the Cauchy invariance.

1. INTRODUCTION

In this paper we derive a priori estimates for the incompressible free-boundary Euler equations with surface tension in three space dimensions. While such equations have been extensively studied, estimates with optimal regularity are not available. With this eventual goal in mind, we provide estimates for the Lagrangian formulation of the system that close with the velocity in H^3 , lowering the regularity from previously known results (see the literature review below).

The incompressible free-boundary Euler equations in a domain of \mathbb{R}^3 are given by

$$\frac{\partial u}{\partial t} + \nabla_u u + \nabla p = 0 \quad \text{in } \mathcal{D}, \quad (1.1a)$$

$$\operatorname{div} u = 0 \quad \text{in } \mathcal{D}, \quad (1.1b)$$

$$p = \sigma \mathcal{H} \quad \text{on } \partial \mathcal{D}, \quad (1.1c)$$

$$(\partial_t + u_\alpha \partial_{x_\alpha})|_{\partial \mathcal{D}} \in T \partial \mathcal{D}, \quad (1.1d)$$

$$u(0, \cdot) = u_0, \quad \Omega(0) = \Omega_0, \quad (1.1e)$$

where

$$\mathcal{D} = \bigcup_{0 \leq t < T} \{t\} \times \Omega(t).$$

Above, the quantities $u = u(t, x)$ and $p = p(t, x)$ represent the velocity and the pressure of the fluid, $\Omega(t) \subset \mathbb{R}^3$ is the moving (i.e., changing over time) domain, which may be written as $\Omega(t) = \eta(t)(\Omega_0)$, where η is the flow of u , σ is a non-negative constant known as the coefficient of surface tension, and \mathcal{H} is the

E-mail addresses: marcelo.disconzi@vanderbilt.edu, kukavica@usc.edu.

Marcelo M. Disconzi is partially supported by NSF grant 1305705, NSF grant 1812826, a Sloan Research Fellowship provided by the Alfred P. Sloan foundation, and a Discovery grant administered by Vanderbilt University.

Igor Kukavica is partially supported by NSF grant 1615239 and NSF grant 1907992.

mean curvature of the moving (time-dependent) boundary $\partial\Omega(t)$. Note that since \mathcal{H} is the mean curvature of the embedding of the spatial boundary $\partial\Omega(t)$ into \mathbb{R}^3 , (1.1c) means that for each t one has $p(t, \cdot) = \sigma\mathcal{H}$. Also, $T\partial\mathcal{D}$ is the tangent bundle of $\partial\mathcal{D}$, with ∂ denoting the space-time boundary. The equation (1.1d) means that the boundary $\partial\Omega(t)$ moves at a speed equal to the normal component of u . The quantity u_0 is the velocity at time zero (necessarily divergence-free by (1.1b)) and Ω_0 is the domain at the initial time, assumed with smooth boundary. The symbol ∇_u is the derivative in the direction of u , often written as $u \cdot \nabla$. The unknowns in (1.1) are u , p , and $\Omega(t)$. Note that \mathcal{H} and $T\partial\mathcal{D}$ are functions of the unknowns and thus have to be determined alongside a solution to the problem.

The first existence results for (1.1) (or, equivalently, for the equations in Lagrangian coordinates, see (2.3) below), are those of Nalimov [61] and Yosihara [74], who considered regular irrotational data. In the case of zero surface tension, Ebin has shown in [37] that the problem is ill-posed without the Rayleigh-Taylor stability condition. The problem of well-posedness under this stability condition and in the case of zero surface tension was obtained by Wu [70, 71]. Recently, Wang et al have obtained in [69] the local existence under the sharp Sobolev regularity $H^{2.5+\delta}$ for the zero surface tension case, extending the previous result of Alazard et al [10], who considered irrotational data.

The case with non-zero surface tension has proven to be more difficult to treat in low regularity spaces, at least in the case of rotational fluids. Currently, one does not have estimates that close in spaces near the threshold $H^{2.5+\delta}$. In [66], Schweizer constructed solutions with rotational data in $H^{4.5}$ with an additional vorticity condition at the surface. Coutand and Shkoller [25] used the Lagrangian formulation and constructed solutions with $H^{4.5}$ initial data without this restriction. At the same time, in [67] Shatah and Zeng obtained a priori estimates for H^3 data in Eulerian coordinates using techniques of infinite dimensional geometry in the spirit of Ebin and Marsden [38] (see also [68], where the authors showed how to use their a priori estimates to obtain a local existence result). Ignatova and the second author obtained in [48] a priori estimates with interior regularity in $H^{3.5}$, using the Lagrangian (direct) approach, while Ebin and the first author established a local-existence result in $H^{3.5+\delta}$ using a combination of the Lagrangian approach, infinite-dimensional geometry, and semi-group theory.

For other results on irrotational fluids with surface tension cf. [2, 3, 4, 5, 11, 12, 13, 17, 30, 42, 46, 52, 75]. Further related results with non-zero surface tension, including the case of rotational fluids, vortex sheets, two-phase fluids, and singular limits, are [22, 27, 31, 32, 40, 51, 62, 65]. Free-boundary problems constitute a very active and fast-growing area of research, and a complete, or even thorough review of prior works is beyond the scope of this paper. A partial list of references relevant to the above discussion and the results of this paper is [1, 6, 7, 8, 9, 15, 16, 18, 19, 20, 21, 23, 24, 26, 28, 29, 41, 44, 45, 47, 49, 50, 53, 54, 55, 56, 57, 58, 59, 60, 63, 72, 73].

The aim of the present paper is to lower the known initial interior regularity for the initial data to H^3 using the Lagrangian approach. We believe that the direct nature of the proof which we provide will allow further lowering of the regularity to the optimal one $H^{2.5+\delta}$. So far, this has only been achieved in the irrotational case by Alazard et al in [4].

The paper is organized as follows. Section 2 contains the main setup of the problem and the main statement, Theorem 2.3. Section 3 provides preliminary bounds on the Lagrangian variable and the cofactor matrix. It also provides some basic geometric results that are used throughout (see Lemma 3.5). The next two sections provide the pressure estimates and the boundary regularity of the Lagrangian flow. Section 6 contains the main inequality for system differentiated with respect to time three times. In Section 7, we deduce regularity of time derivatives and the pressure at time zero; this is where some of the technical conditions stated in Theorem 2.3, namely, an extra regularity of some quantities on the boundary, becomes apparent. Section 8 contains the main estimate for the twice time differentiated system. The next section contains the statement on the boundary regularity of the normal component of the velocity. Section 10 provides estimates for the divergence and the curl of the velocity. Finally, in the last section we provide the proof of the main statement.

2. THE SYSTEM IN LAGRANGIAN COORDINATES AND THE MAIN RESULT

We focus on the case $\sigma > 0$ and consider the model case

$$\Omega_0 \equiv \Omega = \mathbb{T}^2 \times (0, 1).$$

Denoting coordinates on Ω by (x^1, x^2, x^3) , set

$$\Gamma_1 = \mathbb{T}^2 \times \{x^3 = 1\}$$

and

$$\Gamma_0 = \mathbb{T}^2 \times \{x^3 = 0\},$$

so that $\partial\Omega = \Gamma_0 \cup \Gamma_1$. Using a change of coordinates, it is easy to adjust the approach to cover the case when the initial domain is a graph (cf. [55]); the estimates remain unchanged modulo lower order terms. When the domain is not a graph, it is possible to use a straightening of the boundary and a partition of unity, as in [25].

We assume that the lower boundary, which corresponds to the rigid bottom of the fluid domain, does not move, and thus $\eta(t)(\Gamma_0) = \Gamma_0$, where η is the flow of the vector field u . We introduce the Lagrangian velocity and the pressure, respectively, by $v(t, x) = u(t, \eta(t, x))$ and $q(t, x) = q(t, \eta(t, x))$. Therefore,

$$\partial_t \eta = v. \quad (2.1)$$

Denoting by ∇ the derivative with respect to the spatial variables x , introduce the matrix

$$a = (\nabla \eta)^{-1},$$

which is well defined for η near the identity. The map η is volume-preserving as a consequence of (1.1b), which in turn implies the Piola identity

$$\partial_\beta a_{\beta\alpha} = 0. \quad (2.2)$$

(The identity (2.2) can be verified by direct computation using (3.13) below, or cf. [39, p. 462].) Above and throughout, we adopt the following agreement.

Notation 2.1. We denote by ∂_α spatial derivatives, i.e., $\partial_\alpha = \partial/\partial x_\alpha$, for $\alpha = 1, 2, 3$. Greek indices (α, β , etc.) range from 1 to 3 and Latin indices (i, j , etc.), range from 1 to 2. Repeated indices are summed over their range.

In terms of v , q , and a , the system (1.1) becomes

$$\partial_t v_\alpha + a_{\mu\alpha} \partial_\mu q = 0 \quad \text{in } [0, T] \times \Omega, \quad (2.3a)$$

$$a_{\alpha\beta} \partial_\alpha v_\beta = 0 \quad \text{in } [0, T] \times \Omega, \quad (2.3b)$$

$$\partial_t a_{\alpha\beta} + a_{\alpha\gamma} \partial_\mu v_\gamma a_{\mu\beta} = 0 \quad \text{in } [0, T] \times \Omega, \quad (2.3c)$$

$$a_{\mu\alpha} N_\mu q + \sigma |a^T N| \Delta_g \eta_\alpha = 0 \quad \text{on } [0, T] \times \Gamma_1, \quad (2.3d)$$

$$v_\mu N_\mu = 0 \quad \text{on } [0, T] \times \Gamma_0, \quad (2.3e)$$

$$\eta(0, \cdot) = \text{id}, \quad v(0, \cdot) = v_0 = (v_{01}, v_{02}, v_{03}) \text{ in } \Omega, \quad (2.3f)$$

where id is the identity diffeomorphism on Ω , N is the unit outer normal to $\partial\Omega$, a^T is the transpose of a , $|\cdot|$ is the Euclidean norm, and Δ_g is the Laplacian on the boundary with respect to the metric g_{ij} induced on $\partial\Omega(t)$ by the embedding η . Explicitly,

$$g_{ij} = \partial_i \eta_\mu \partial_j \eta_\mu \quad (2.4)$$

and

$$\Delta_g(\cdot) = \frac{1}{\sqrt{g}} \partial_i (\sqrt{g} g^{ij} \partial_j(\cdot)), \quad (2.5)$$

where g is the determinant of the matrix (g_{ij}) and (g^{ij}) is the inverse matrix. In (2.3d), $\Delta_g \eta_\alpha$ simply means Δ_g acting on the scalar function η_α , for each $\alpha = 1, 2, 3$. See Lemma 3.5 below for some important identities used to obtain (2.3d).

Since $\eta(0, \cdot) = \text{id}$, the initial Lagrangian and Eulerian velocities agree, i.e., $v_0 = u_0$. Clearly, v_0 is orthogonal to Γ_0 in view of (2.3e). Note that

$$a(0, \cdot) = I,$$

where I is the identity matrix, in light of (2.3f).

Notation 2.2. Sobolev spaces are denoted by $H^s(\Omega)$, or simply by H^s when no confusion can arise, with the corresponding norm $\|\cdot\|_s$; note that $\|\cdot\|_0$ refers to the L^2 norm. We denote by $H^s(\partial\Omega)$ the Sobolev space of maps defined on $\partial\Omega$, with the corresponding norm $\|\cdot\|_{s,\partial}$, and similarly the space $H^s(\Gamma_1)$ with the norm $\|\cdot\|_{s,\Gamma_1}$. The L^p norms on Ω and Γ_1 are denoted by $\|\cdot\|_{L^p(\Omega)}$ (or $\|\cdot\|_{L^p}$ when no confusion can arise) and $\|\cdot\|_{L^p(\Gamma_1)}$, respectively. We use $|$ to denote restriction.

We now state our main result.

Theorem 2.3. *Let Ω be as described above, and let $\sigma > 0$. Let v_0 be a smooth divergence-free vector field on Ω and assume that v_0 is tangent to Γ_0 . Then there exist $T_* > 0$ and a constant C_0 , depending only on $\sigma > 0$, $\|v_0\|_3$, and $\|v_{03}\|_{4,\Gamma_1}$, such that any smooth solution (v, q) to (2.3) with initial condition v_0 and defined on the time interval $[0, T_*)$, satisfies*

$$\|v\|_3 + \|\partial_t v\|_{2.5} + \|\partial_t^2 v\|_{1.5} + \|\partial_t^3 v\|_0 + \|q\|_3 + \|\partial_t q\|_2 + \|\partial_t^2 q\|_1 \leq C_0. \quad (2.6)$$

The rest of the paper is dedicated to proving this statement. We would like to point out that the theorem above constitutes the first step toward establishing a complete proof of local existence with the stated regularity. In order to obtain such a proof, one should also provide the construction of the solution. The main additional difficulty in this regard, which will be addressed in a future work, is to obtain a formulation of the Cauchy invariance that is amenable to perturbations. We would like to add that the adaptation of our results to non-flat domains is standard. However, we believe that the necessary straightening of the boundary and use of a partition of unity would make the paper more difficult to follow.

The dependence of T_* and C_0 on a higher norm of the normal component of the velocity on the boundary Γ_1 comes from the usual difficulties caused by the moving boundary in free-boundary problems. Note that if $v_0 \in H^{4.5}$, as in [25, 66], then this condition is automatically satisfied. Also, if the initial data is irrotational in a neighborhood of the interface Γ_1 , we have $\Delta v_{03} = 0$ there, and the additional condition on the normal component is not necessary; cf. Section 7 below.

For the rest of the paper, let (v, q) be a smooth solution to (2.3) and assume that Ω , Γ_1 , and Γ_0 are as described above.

3. AUXILIARY RESULTS

In this section we state some preliminary results that are employed in the proof of Theorem 2.3.

Lemma 3.1. *Assume that $\|v\|_3 \leq M$. Then, there exists a constant $C > 0$ such that if $T \in [0, 1/CM]$ and (v, q) is defined on $[0, T]$, the following inequalities hold for $t \in [0, T]$:*

- (i) $\|\eta\|_3 \leq C$.
- (ii) $\|a\|_2 \leq C$.
- (iii) $\|\partial_t a\|_{L^p} \leq C\|\nabla v\|_{L^p}$, $1 \leq p \leq \infty$.
- (iv) $\|\partial_\alpha \partial_t a\|_{L^p} \leq C\|\nabla v\|_{L^{p_1}} \|\partial_\alpha a\|_{L^{p_2}} + C\|\partial_\alpha \nabla v\|_{L^p}$, where $1/p = 1/p_1 + 1/p_2$ and $1 \leq p, p_1, p_2 \leq 6$.
- (v) $\|\partial_t a\|_s \leq C\|\nabla v\|_s$, $0 \leq s \leq 2$.
- (vi) $\|\partial_t^2 a\|_s \leq C\|\nabla v\|_s \|\nabla v\|_{L^\infty} + C\|\nabla \partial_t v\|_s$ for $0 \leq s \leq 1$, and $\|\partial_t^2 a\|_1 \leq C\|\nabla v\|_{5/4}^2 + C\|\nabla \partial_t v\|_1$.
- (vii) $\|\partial_t^3 a\|_{L^p} \leq C\|\nabla v\|_{L^p} \|\nabla v\|_{L^\infty}^2 + C\|\nabla \partial_t v\|_{L^p} \|\nabla v\|_{L^\infty} + C\|\nabla \partial_t^2 v\|_{L^p}$, $1 \leq p < \infty$.
- (viii) If $\epsilon \in (0, 1]$ and $T \leq \epsilon/CM^2$ then, for $t \in [0, T]$, we have

$$\|a_{\alpha\beta} - \delta_{\alpha\beta}\|_2 \leq \epsilon$$

and

$$\|a_{\alpha\mu} a_{\beta\mu} - \delta_{\alpha\beta}\|_2 \leq \epsilon.$$

Furthermore, if $\epsilon \in (0, 1/C]$, where C is a sufficiently large constant, then we have the strong ellipticity

$$a_{\alpha\mu} a_{\beta\mu} \xi_\alpha \xi_\beta \geq \frac{1}{C} |\xi|^2.$$

We emphasize that the constant C above, as well as in the rest of the paper, does not depend on M .

Proof of Lemma 3.1. The proofs of these inequalities follow directly from (2.1) and (2.3c) as in any of the references [48, 54, 55] and are thus omitted. \square

We also recall the following statement from [48] on the H^1 regularity of the Neumann problem.

Lemma 3.2 ([48]). *Let f be an H^1 solution to*

$$\begin{aligned} \partial_\alpha(b_{\alpha\beta}\partial_\beta f) &= \partial_\alpha\pi_\alpha \text{ in } \Omega, \\ b_{\alpha\beta}\partial_\alpha f N_\beta &= g \quad \text{on } \partial\Omega, \end{aligned}$$

where $b \in H^2(\Omega)$, $b_{\alpha\beta} = b_{\beta\alpha}$, $\pi, \partial_\alpha\pi_\alpha \in L^2(\Omega)$, and $g \in H^{-\frac{1}{2}}(\partial\Omega)$, with the compatibility condition

$$\int_{\partial\Omega} (\pi_\mu N_\mu - g) = 0.$$

Suppose that $\|b\|_{L^\infty} \leq M$ and $b_{\alpha\beta}\xi_\alpha\xi_\beta \geq |\xi|^2/M$. If

$$\|b - I\|_{L^\infty} \leq \epsilon_0,$$

where I is the identity matrix and ϵ_0 is a sufficiently small constant depending on M , then

$$\|f\|_1 \leq C\|\pi\|_{L^2\Omega} + C\|g - \pi_\mu N_\mu\|_{-\frac{1}{2},\partial} + \|f\|_{L^2(\Gamma_1)}.$$

Notation 3.3. In the rest of the paper, the symbol C denotes a positive sufficiently large constant. It can vary from expression to expression, but it is always independent of $\|v\|_3$, $\|\partial_t v\|_{2.5}$, $\|\partial_t^2 v\|_{1.5}$, $\|\partial_t^3 v\|_0$, $\|q\|_3$, $\|\partial_t q\|_2$, and $\|\partial_t^2 q\|_1$. The a priori estimates require T to be sufficiently small so that it satisfies $TM \leq 1/C$, where M is an upper bound on the norm of the solution (cf. Lemma 3.1 above). In several estimates it suffices to keep track only on the number of derivatives so we write ∂^ℓ to denote any (spatial) derivative of order ℓ and $\bar{\partial}^\ell$ to denote any (spatial) derivative of order ℓ on the boundary, i.e., with respect to x_i . We use upper-case Latin indices to denote x_i or t , so $\bar{\partial}_A$ means ∂_t or ∂_i .

Remark 3.4. Throughout the paper, we have to estimate several rational functions of tangential derivatives of η that come from various combinations of g^{ij} and \sqrt{g} , where by tangential derivatives we mean ∂_1 and ∂_2 . Although there will be several different such expressions, they can all be organized in the same fashion. Therefore, it is beneficial to introduce some general notation and explain how these terms will be handled in a unified manner. We also take this opportunity to introduce some symbolic notation that will help us keep track of the number of derivatives.

Consider a rational function Q of the (tangential) derivatives of η :

$$Q = Q(\partial_1\eta_1, \partial_2\eta_1, \partial_1\eta_2, \partial_2\eta_2, \partial_1\eta_3, \partial_2\eta_3).$$

We assume that Q is defined in an open domain \mathcal{D} of \mathbb{R}^6 . Assume that $0 \notin \bar{\mathcal{D}}$ and that $(1, 0, 0, 1, 0, 0) \in \mathcal{D}$. This last assumption reflects the fact that at $t = 0$ we have $\eta = \text{id}$, so that $\partial_1\eta_1 = \partial_2\eta_2 = 1$ at $t = 0$, while the other derivatives are zero. In what follows it suffices to keep track of the generic form of some expressions so we write Q symbolically as

$$Q = Q(\bar{\partial}\eta).$$

Then

$$\bar{\partial}_A Q(\bar{\partial}\eta) = \tilde{Q}_\alpha^i(\bar{\partial}\eta)\bar{\partial}_A\partial_i\eta_\alpha,$$

where the terms $\tilde{Q}_\alpha^i(\bar{\partial}\eta)$ are also rational function of derivatives of η with respect to x_i . Note that $\tilde{Q}_\alpha^i(\bar{\partial}\eta)$ are simply the partial derivatives of Q evaluated at $\bar{\partial}\eta$. We write the last equality symbolically as

$$\bar{\partial}_A Q(\bar{\partial}\eta) = \tilde{Q}(\bar{\partial}\eta)\bar{\partial}_A\bar{\partial}\eta. \quad (3.1)$$

Regarding \tilde{Q} , we assume that for all $\bar{\partial}\eta$ taking values in some small neighborhood \mathcal{D}' of $(1, 0, 0, 1, 0, 0)$, and for any $1 < s \leq 1.5$, we have

$$\|\tilde{Q}(\bar{\partial}\eta)\|_{s,\Gamma_1} \leq C \quad (3.2)$$

for some constant C depending only on \mathcal{D}' .

For $s > 1$, we have

$$\|\bar{\partial}_A Q(\bar{\partial}\eta)\|_{s,\Gamma_1} \leq C_1 \|\tilde{Q}(\bar{\partial}\eta)\|_{s,\Gamma_1} \|\bar{\partial}_A \bar{\partial}\eta\|_{s,\Gamma_1}$$

where C_1 depends only on s and on the domain Γ_1 .

The term $\|\tilde{Q}(\bar{\partial}\eta)\|_{s,\Gamma_1}$ can be estimated using the Sobolev norm of the map \tilde{Q} , i.e., $\|\tilde{Q}\|_{H^s(\mathcal{D}')}$, and the Sobolev norm of $\bar{\partial}\eta$, i.e., $\|\bar{\partial}\eta\|_{s,\Gamma_1}$. Under the conditions of Lemma 3.1, we have

$$\|\bar{\partial}\eta - \bar{\partial}\eta(0)\|_{L^\infty(\Gamma_1)} \leq \int_0^t \|\partial_t \bar{\partial}\eta\|_{L^\infty(\Gamma_1)} \leq C_2 t \|v\|_3 \leq C_2 M t,$$

where C_2 depends only on the domain Γ_1 and we used that $H^{1.5}(\Gamma_1)$ embeds into $C^0(\Gamma_1)$. Therefore, if $CMT \leq 1$, we can guarantee that

$$\bar{\partial}\eta(\Gamma_1) \subset \mathcal{D}',$$

and thus, shrinking \mathcal{D} if necessary, we can apply the estimate (3.2). Since Lemma 3.1 also provides a bound for $\|\bar{\partial}\eta\|_{s,\Gamma_1}$, with $s \leq 1.5$, we conclude that

$$\|\bar{\partial}_A Q(\bar{\partial}\eta)\|_{s,\Gamma_1} \leq C \|\bar{\partial}_A \bar{\partial}\eta\|_{s,\Gamma_1}, \text{ for } 1 < s \leq 1.5, \quad (3.3)$$

where C depends only on M , s , and Γ_1 , and provided that t is small enough. The above also shows that

$$\|Q(\bar{\partial}\eta)\|_{s,\Gamma_1} \leq C \|\bar{\partial}\eta\|_{s,\Gamma_1}, \text{ for } 1 < s \leq 1.5. \quad (3.4)$$

Above, the restriction $s > 1$ comes from the Sobolev multiplicative theorem, while $s \leq 1.5$ allows us to invoke the estimates of Lemma 3.1.

We also need some geometric identities that may be known to specialists, but we state them below and provide some of the corresponding proofs for the reader's convenience.

Lemma 3.5. *Denote by n the unit outer normal to $\eta(\Gamma_1)$. Then*

$$n \circ \eta = \frac{a^T N}{|a^T N|}. \quad (3.5)$$

Denoting by τ the tangent bundle of $\overline{\eta(\Omega)}$ and by ν the normal bundle of $\eta(\Gamma_1)$, the canonical projection $\Pi: \tau|_{\eta(\Gamma_1)} \rightarrow \nu$ is given by

$$\Pi_{\alpha\beta} = \delta_{\alpha\beta} - g^{kl} \partial_k \eta_\alpha \partial_l \eta_\beta. \quad (3.6)$$

Furthermore, the following identities hold:

$$\Pi_{\alpha\lambda} \Pi_{\lambda\beta} = \Pi_{\alpha\beta}, \quad (3.7)$$

$$|a^T N| = \sqrt{g}, \quad (3.8)$$

$$\sqrt{g} \Delta_g \eta_\alpha = \sqrt{g} g^{ij} \partial_{ij}^2 \eta_\alpha - \sqrt{g} g^{ij} g^{kl} \partial_k \eta_\alpha \partial_l \eta_\mu \partial_{ij}^2 \eta_\mu, \quad (3.9)$$

$$\begin{aligned} \bar{\partial}_A(\sqrt{g} \Delta_g \eta_\alpha) &= \partial_i \left(\sqrt{g} g^{ij} (\delta_{\alpha\lambda} - g^{kl} \partial_k \eta_\alpha \partial_l \eta_\lambda) \bar{\partial}_A \partial_j \eta_\lambda \right. \\ &\quad \left. + \sqrt{g} (g^{ij} g^{kl} - g^{lj} g^{ik}) \partial_j \eta_\alpha \partial_k \eta_\lambda \bar{\partial}_A \partial_l \eta_\lambda \right), \end{aligned} \quad (3.10)$$

$$-\Delta_g(\eta_\alpha|_{\Gamma_1}) = \mathcal{H} \circ \eta n_\alpha \circ \eta. \quad (3.11)$$

Proof of Lemma 3.5. Letting $r = \eta|_{\Gamma_1}$, we know that $n \circ \eta$ is given by

$$n \circ \eta = \frac{\partial_1 r \times \partial_2 r}{|\partial_1 r \times \partial_2 r|} \quad (3.12)$$

(cf. e.g. [43]). By $\det(\nabla \eta) = 1$, we have

$$a = \begin{bmatrix} \partial_2 \eta_2 \partial_3 \eta_3 - \partial_3 \eta_2 \partial_2 \eta_3 & \partial_3 \eta_1 \partial_2 \eta_3 - \partial_2 \eta_1 \partial_3 \eta_3 & \partial_2 \eta_1 \partial_3 \eta_2 - \partial_3 \eta_1 \partial_2 \eta_2 \\ \partial_3 \eta_2 \partial_1 \eta_3 - \partial_1 \eta_2 \partial_3 \eta_3 & \partial_1 \eta_1 \partial_3 \eta_3 - \partial_3 \eta_1 \partial_1 \eta_3 & \partial_3 \eta_1 \partial_1 \eta_2 - \partial_1 \eta_1 \partial_3 \eta_2 \\ \partial_1 \eta_2 \partial_2 \eta_3 - \partial_2 \eta_2 \partial_1 \eta_3 & \partial_2 \eta_1 \partial_1 \eta_3 - \partial_1 \eta_1 \partial_2 \eta_3 & \partial_1 \eta_1 \partial_2 \eta_2 - \partial_2 \eta_1 \partial_1 \eta_2 \end{bmatrix}. \quad (3.13)$$

Using (3.13) to compute $a^T N$ and comparing with $\partial_1 r \times \partial_2 r$, one verifies that $a^T N = \partial_1 r \times \partial_2 r$, and then (3.8) follows from (3.12).

To prove (3.6), we use (3.5) to write

$$(\delta_{\alpha\lambda} - g^{kl} \partial_k \eta_\alpha \partial_l \eta_\lambda) n_\lambda \circ \eta = \frac{a_{\mu\alpha} N_\mu}{|a^T N|} - \frac{g^{kl} \partial_k \eta_\alpha \partial_l \eta_\lambda a_{\mu\lambda} N_\mu}{|a^T N|}.$$

Contracting $g^{kl} \partial_l \eta_\lambda a_{\mu\lambda} N_\mu$ with g_{mk} gives

$$\begin{aligned} g_{mk} g^{kl} \partial_l \eta_\lambda a_{\mu\lambda} N_\mu &= \partial_m \eta_\lambda a_{3\lambda} \\ &= \partial_m \eta_1 (\partial_1 \eta_2 \partial_2 \eta_3 - \partial_2 \eta_2 \partial_1 \eta_3) + \partial_m \eta_2 (\partial_2 \eta_1 \partial_1 \eta_3 - \partial_1 \eta_1 \partial_2 \eta_3) \\ &\quad + \partial_m \eta_3 (\partial_1 \eta_1 \partial_2 \eta_2 - \partial_2 \eta_1 \partial_1 \eta_2) = 0. \end{aligned} \quad (3.14)$$

Above, the first equality follows because $N = (0, 0, 1)$ (and $g_{mk} g^{kl} = \delta_m^l$), the second equality uses (3.13), and the third equality follows upon setting $m = 1$ and then $m = 2$ and observing that in each case all the terms cancel out. Thus, contracting (3.14) with g^{mn} , we obtain $g^{nl} \partial_l \eta_\lambda a_{\mu\lambda} N_\mu = 0$, and hence

$$(\delta_{\alpha\lambda} - g^{kl} \partial_k \eta_\alpha \partial_l \eta_\lambda) n_\lambda \circ \eta = \frac{a_{\mu\alpha} N_\mu}{|a^T N|}.$$

To conclude the proof of (3.6), we need to verify that $\Pi(X) = 0$ if X is tangent to $\eta(\Gamma_1)$. Since the tangent space to $\eta(\Gamma_1)$ is spanned by $\partial_j \eta$, for $j = 1, 2$, it suffices to verify the identity for these vectors. We have

$$\Pi_{\alpha\mu} \partial_j \eta_\mu = (\delta_{\alpha\mu} - g^{kl} \partial_k \eta_\alpha \partial_l \eta_\mu) \partial_j \eta_\mu = \partial_j \eta_\alpha - g^{kl} \partial_k \eta_\alpha g_{lj} = 0,$$

where we used $g_{lj} = \partial_l \eta_\mu \partial_j \eta_\mu$ and $g^{kl} g_{lj} = \delta_j^k$. Thus, (3.6) is proven.

The identity (3.7) follows from the fact that Π is a projection operator or, alternatively, by direct computation using (3.6). The identity (3.8) follows from (3.5), (3.12), and the standard formula

$$\frac{\partial_1 r \times \partial_2 r}{|\partial_1 r \times \partial_2 r|} = \frac{1}{\sqrt{g}} \partial_1 r \times \partial_2 r$$

(see e.g. [43]). In order to prove (3.9), recall that (see e.g. [43])

$$\Delta_g \eta_\alpha = g^{ij} \partial_{ij}^2 \eta_\alpha - g^{ij} \Gamma_{ij}^k \partial_k \eta_\alpha, \quad (3.15)$$

where Γ_{ij}^k are the Christoffel symbols. Recalling (2.4), a direct computation using the definition of the Christoffel symbols gives

$$\Gamma_{ij}^k = g^{kl} \partial_l \eta_\mu \partial_{ij}^2 \eta_\mu, \quad (3.16)$$

and (3.9) follows from (3.15) and (3.16).

Now, we move to establish (3.10). Using (2.5),

$$\bar{\partial}_A(\sqrt{g} \Delta_g \eta_\alpha) = \bar{\partial}_A \partial_i (\sqrt{g} g^{ij} \partial_j \eta_\alpha) = \partial_i (\sqrt{g} g^{ij} \partial_j \bar{\partial}_A \eta_\alpha + \bar{\partial}_A (\sqrt{g} g^{ij}) \partial_j \eta_\alpha). \quad (3.17)$$

Recalling the standard identity (see e.g. [64, p. 51]),

$$\bar{\partial}_A g = g g^{kl} \bar{\partial}_A g_{kl}, \quad (3.18)$$

which is Jacobi's identity connecting the derivative of a determinant with a trace, we find

$$\bar{\partial}_A(\sqrt{g} g^{ij}) = \sqrt{g} \left(\frac{1}{2} g^{ij} g^{kl} - g^{lj} g^{ik} \right) \bar{\partial}_A g_{kl},$$

where we also used

$$\bar{\partial}_A g^{ij} = -g^{lj} g^{ik} \bar{\partial}_A g_{kl} \quad (3.19)$$

that follows from differentiating $g_{kl} g^{lj} = \delta_k^j$ and then contracting with g^{ik} . Computing $\bar{\partial}_A g_{kl}$ directly from (2.4) then leads to

$$\bar{\partial}_A(\sqrt{g} g^{ij}) = \sqrt{g} \left(\frac{1}{2} g^{ij} g^{kl} - g^{lj} g^{ik} \right) (\bar{\partial}_A \partial_k \eta_\lambda \partial_l \eta_\lambda + \partial_k \eta_\lambda \bar{\partial}_A \partial_l \eta_\lambda). \quad (3.20)$$

Using (3.20) in (3.17) then yields (3.10) after some computation.

Finally, the identity (3.11) is a standard formula for the mean curvature of an embedding into \mathbb{R}^3 (see e.g. [43, Section 2.1] or [64, Section I.2]). \square

4. PRESSURE ESTIMATES

Through the rest of the paper, we suppose that the hypotheses of Lemma 3.1 hold. We assume further that T is as in part (viii) of that lemma, and that (v, q) are defined on $[0, T)$.

Proposition 4.1. *We have the estimates*

$$\begin{aligned} \|q\|_3 &\leq C\|\nabla v\|_2\|v\|_2 + C\|\partial_t v\|_{1.5, \Gamma_1} + C, \\ \|\partial_t q\|_2 &\leq C\|\nabla v\|_{1.5+\delta}(\|q\|_{2.5} + \|\partial_t v\|_{1.5}) + C(\|\nabla v\|_{1.5}\|\nabla v\|_{L^\infty} + \|\nabla \partial_t v\|_{1.5})\|v\|_{1.5+\delta} \\ &\quad + C\|\partial_t^2 v\|_{1, \Gamma_1} + C\|v\|_{2.5} + C\|\partial_t v\|_{2.5}, \\ \|\partial_t^2 q\|_1 &\leq C(\|v\|_{1.5}\|\nabla v\|_{L^\infty} + \|\partial_t v\|_{1.5})(\|q\|_2 + \|\partial_t v\|_1) + C\|\nabla v\|_{L^\infty}(\|\partial_t q\|_1 + \|\partial_t^2 v\|_0) \\ &\quad + C(\|v\|_{1.5}\|\nabla v\|_{L^\infty}^2 + \|\partial_t v\|_{1.5}\|\nabla v\|_{L^\infty} + \|\partial_t^2 v\|_{1.5})\|v\|_1 + C\|\partial_t^3 v\|_0 \\ &\quad + C(\|\partial_t v\|_{2.5} + \|v\|_3\|\partial_t q\|_1) + C(1 + \|v\|_3^2 + \|\partial_t v\|_2)(\|\partial_t v\|_2 + \|v\|_3^2), \end{aligned}$$

for $t \in (0, T)$ and with $\delta > 0$ a small number.

The pressure estimates are performed similarly to [48]. The differences are the spaces in which we estimate q and $\partial_t q$. The inequality for $\partial_t^2 q$ is the same as in [48], except for the bound on $\|\partial_t^2 q\|_{0, \Gamma_1}$, which we state in (4.3) below. The adjustment has to be made to account for the evolution of the Riemannian metric on the boundary.

Proof of Proposition 4.1. Contracting $a_{\mu\alpha}\partial_\mu$ with (2.3a) and using (2.3b) we find

$$\Delta q = \partial_\mu((\delta_{\mu\lambda} - a_{\mu\nu}a_{\lambda\nu})\partial_\lambda q) + \partial_\mu(\partial_t a_{\mu\lambda}v_\lambda), \quad (4.1)$$

where we also used (2.2). Contracting (2.3a) with $a_{\mu\alpha}N_\mu$ and restricting to the boundary yields

$$\frac{\partial q}{\partial N} = (\delta_{\mu\lambda} - a_{\mu\nu}a_{\lambda\nu})N_\mu\partial_\lambda q - a_{\mu\nu}N_\mu\partial_t v_\nu \quad \text{on } \partial\Omega. \quad (4.2)$$

Denoting the right-hand sides of (4.1) and (4.2) by f and g , respectively, we have a standard elliptic estimate

$$\|q\|_3 \leq C\|f\|_1 + C\|g\|_{1.5, \partial} + \|q\|_0$$

(see e.g. [21, Theorem 3.6]). With the help of Lemma 3.1, we find

$$\|f\|_1 \leq \epsilon\|q\|_3 + C\|\nabla v\|_2\|v\|_2$$

and

$$\|g\|_{1.5, \partial} \leq \epsilon\|q\|_3 + C\|\partial_t v\|_{1.5, \Gamma_1}.$$

To estimate $\|q\|_0$, we use

$$\left| \int_\Omega q \right| \leq C\|\nabla(q - \bar{q})\|_0 + C\|q\|_{0, \Gamma_1},$$

which follows from the identity

$$\int_\Omega q = \int_\Omega q\Delta h = - \int_\Omega \nabla q \cdot \nabla h + \int_{\partial\Omega} \frac{\partial h}{\partial N} q$$

and the choice $h = x_3^2 - 1$. Using (2.3d) and Lemma 3.1, we obtain $\|q\|_{0, \Gamma_1} \leq C$. Combining the above inequalities gives the estimate for q after a simple application of the ϵ -Cauchy inequality.

The estimate for $\partial_t q$ is obtained similarly after time-differentiating (2.3a) and proceeding as above.

To obtain the estimate for $\partial_t^2 q$, we differentiate (2.3a) in time twice and apply Lemma 3.2. This is done similarly to [48], with an exception of the estimate on $\|\partial_t^2 q\|_{0, \Gamma_1}$ which reads

$$\|\partial_t^2 q\|_{0, \Gamma_1} \leq \epsilon\|\partial_t^2 q\|_1 + C(\|\partial_t v\|_{2.5} + \|v\|_3\|\partial_t q\|_1) + C(1 + \|v\|_3^2 + \|\partial_t v\|_2)(\|\partial_t v\|_2 + \|v\|_3^2) \quad (4.3)$$

leading to the last two terms in the statement. \square

5. REGULARITY ESTIMATE FOR THE FLOW

One of the key features of the free-boundary Euler equations with surface tension is a gain of regularity for the free-boundary [25, 33]. As discussed in [67], this gain is geometric in nature and does not correspond to the regularity of the flow in the interior (see the counter-example in [67], which shows that in the interior of the domain, the flow cannot in general be more regular than the velocity, even if the boundary gains regularity). This gain of regularity is formulated in the next statement.

Proposition 5.1. *We have the estimate*

$$\|\eta\|_{3.5,\Gamma_1} \leq P(\|q\|_{1.5,\Gamma_1}),$$

where P is a polynomial.

Notation 5.2. From here on, we use $P(\cdot)$, with indices attached when appropriate, to denote a generic polynomial expression of its arguments. The polynomial is assumed nonnegative and its value may change from inequality to inequality. Also, we write

$$\mathcal{P} = P(\|v\|_3, \|\partial_t v\|_{2.5}, \|\partial_t^2 v\|_{1.5}, \|\partial_t^3 v\|_0, \|q\|_3, \|\partial_t q\|_2, \|\partial_t^2 q\|_1) \quad (5.1)$$

for a generic polynomial depending on the norms in (2.6). Finally, we denote by \mathcal{P}_0 a generic polynomial of $\|v_0\|_3$ and $\|v_{03}\|_{4,\Gamma_1}$, i.e.,

$$\mathcal{P}_0 = P(\|v_0\|_3, \|v_{03}\|_{4,\Gamma_1}).$$

Throughout the paper, expressions involving the norms in (2.6) evaluated at time zero are replaced by \mathcal{P}_0 . That \mathcal{P}_0 in fact controls all such terms evaluated at time zero is shown in Section 7; cf. (7.5) below.

Proof of Proposition 5.1. We would like to apply elliptic estimates to (2.3d). While we do not know a priori that the coefficients g_{ij} have enough regularity for an application of standard elliptic estimates, we can use improved estimates for coefficients with lower regularity as in [34]. For this, it suffices to check that g_{ij} has small oscillation, in the following sense. Given $r > 0$ and $x \in \Gamma_1$, set

$$\text{osc}_{x,r}(g^{ij}) = \frac{1}{\text{vol}(B_r(x))} \int_{B_r(x)} \left| g^{ij}(y) - \frac{1}{\text{vol}(B_r(x))} \int_{B_r(x)} g^{ij}(z) dz \right| dy$$

and

$$g_R = \sup_{x \in \Gamma_1} \sup_{r \leq R} \text{osc}_{x,r}(g^{ij}).$$

We need to verify that there exists $\tilde{R} \leq 1$ such that

$$g_{\tilde{R}} \leq \rho, \quad (5.2)$$

where ρ is sufficiently small.

Since $g^{ij} \in H^{1.5}(\Gamma_1)$, we have $g^{ij} \in C^{0,\alpha}(\Gamma_1)$ with $0 < \alpha < 0.5$ fixed. Thus, for $y \in B_r(x)$,

$$\begin{aligned} \left| g^{ij}(y) - \frac{1}{\text{vol}(B_r(x))} \int_{B_r(x)} g^{ij}(z) dz \right| &= \left| \frac{1}{\text{vol}(B_r(x))} \int_{B_r(x)} (g^{ij}(y) - g^{ij}(z)) dz \right| \\ &\leq \sup_{y,z \in \Gamma_1} |g^{ij}(y) - g^{ij}(z)| \leq C_\alpha r^\alpha. \end{aligned}$$

Hence,

$$g_{\tilde{R}} \leq C_\alpha \tilde{R}^\alpha,$$

and we can ensure (5.2). Therefore, the results of [34] imply that

$$\|\eta_\alpha\|_{3.5,\Gamma_1} \leq C(\|a_{\mu\alpha} N_\mu q\|_{1.5,\Gamma_1} + \|\eta_\alpha\|_{1.5,\Gamma_1}) \leq C(\|a\|_{1.5,\Gamma_1} \|q\|_{1.5,\Gamma_1} + \|\eta\|_{1.5,\Gamma_1}),$$

where C depends on $\|g_{ij}\|_{1.5,\Gamma_1}$. Or yet,

$$\|\eta_\alpha\|_{3.5,\Gamma_1} \leq C\|q\|_{1.5,\Gamma_1} + C\|\eta\|_3 \leq C\|q\|_{1.5,\Gamma_1} + C \leq P(\|q\|_{1.5,\Gamma_1}).$$

We remark that [34] deals only with Sobolev spaces of integer order, but since the estimates are linear on the norms we can extend them to fractional order Sobolev spaces as well. \square

6. ENERGY ESTIMATE ON THE THREE TIMES DIFFERENTIATED SYSTEM

In the next statement, we provide two important bounds: on the L^2 norm of $\partial_t^3 v$ and on the \dot{H}^1 norm of the projection of $\partial_t^2 v$. In the last section, we show that these bounds lead to an estimate for $\|\partial_t^2 v_3\|_{1,\Gamma_1}$

Lemma 6.1. *We have*

$$\|\partial_t^3 v\|_0^2 + \|\bar{\partial}(\Pi\partial_t^2 v)\|_{0,\Gamma_1}^2 \leq \tilde{\epsilon}(\|\partial_t^2 v\|_{1.5} + \|\partial_t^2 q\|_1) + P(\|v\|_{2.5+\delta}, \|q\|_{1.5,\Gamma_1}) + \mathcal{P}_0 + \int_0^t \mathcal{P}, \quad (6.1)$$

where Π is given by (3.6).

We hereafter adopt the following notations.

Notation 6.2. We use $\tilde{\epsilon}$ to denote a small positive constant which may vary from expression to expression. Typically, $\tilde{\epsilon}$ comes from choosing the time sufficiently small, from Lemma 3.1, or from the Cauchy inequality with epsilon. The important point to keep in mind, which can be easily verified in the expressions containing $\tilde{\epsilon}$, is that once all estimates are obtained, we can fix $\tilde{\epsilon}$ to be sufficiently small universal constant in order to close the estimates.

Notation 6.3. We use $0 < \delta < 0.5$ to denote a number that appears in the Sobolev norms (e.g. $\|v\|_{2.5+\delta}$). The values of $\tilde{\epsilon}$ and δ may vary from expression to expression, but they can be fixed to hold uniformly across all expressions at the end. In Section 11.2 we choose δ appropriately.

Proof of Lemma 6.1. We apply ∂_t^3 on (2.3a), contract the resulting equation with $\partial_t^3 v_\alpha$, and integrate in space and time to find

$$\frac{1}{2}\|\partial_t^3 v\|_0^2 = \frac{1}{2}\|\partial_t^3 v(0)\|_0^2 - \int_0^t \int_\Omega \partial_t^3 \partial_\mu (a_{\mu\alpha} q) \partial_t^3 v_\alpha, \quad (6.2)$$

where we also used (2.2). Integrating by parts and using that on Γ_0 we have

$$\begin{aligned} v_3 &= 0, \text{ so that } \partial_i v_3 = 0 \text{ and } \partial_t^3 v_3 = 0, \\ \text{and } \partial_i \eta_3 &= 0, \text{ so that } a_{31} = a_{32} = 0, \end{aligned} \quad (6.3)$$

we find

$$- \int_0^t \int_\Omega \partial_t^3 \partial_\mu (a_{\mu\alpha} q) \partial_t^3 v_\alpha = I_1 + I_2, \quad (6.4)$$

where

$$I_1 = - \int_0^t \int_{\Gamma_1} \partial_t^3 (a_{3\alpha} q) \partial_t^3 v_\alpha = \sigma \int_0^t \int_{\Gamma_1} \partial_t^3 (\sqrt{g} \Delta_g \eta_\alpha) \partial_t^3 v_\alpha \quad (6.5)$$

and

$$\begin{aligned} I_2 &= \int_0^t \int_\Omega a_{\mu\alpha} \partial_t^3 q \partial_\mu \partial_t^3 v_\alpha + 3 \int_0^t \int_\Omega \partial_t a_{\mu\alpha} \partial_t^2 q \partial_\mu \partial_t^3 v_\alpha \\ &\quad + 3 \int_0^t \int_\Omega \partial_t^2 a_{\mu\alpha} \partial_t q \partial_\mu \partial_t^3 v_\alpha + \int_0^t \int_\Omega \partial_t^3 a_{\mu\alpha} q \partial_\mu \partial_t^3 v_\alpha = I_{21} + I_{22} + I_{23} + I_{24}. \end{aligned} \quad (6.6)$$

Note that we used (3.8) and (2.3d) to rewrite I_1 .

Estimate of I_1 : We first use (3.10) with $\bar{\partial}_A = \partial_t$ and integrate by parts to find

$$\begin{aligned}
\frac{1}{\sigma} I_1 &= \int_0^t \int_{\Gamma_1} \partial_t^3 (\sqrt{g} \Delta_g \eta_\alpha) \partial_t^3 v_\alpha \\
&= - \int_0^t \int_{\Gamma_1} \sqrt{g} g^{ij} (\delta_{\alpha\lambda} - g^{kl} \partial_k \eta_\alpha \partial_l \eta_\lambda) \partial_t^2 \partial_j v_\lambda \partial_t^3 \partial_i v_\alpha \\
&\quad - \int_0^t \int_{\Gamma_1} \sqrt{g} (g^{ij} g^{kl} - g^{lj} g^{ik}) \partial_j \eta_\alpha \partial_k \eta_\lambda \partial_t^2 \partial_l v_\lambda \partial_t^3 \partial_i v_\alpha \\
&\quad - 2 \int_0^t \int_{\Gamma_1} \partial_t (\sqrt{g} g^{ij} (\delta_{\alpha\lambda} - g^{kl} \partial_k \eta_\alpha \partial_l \eta_\lambda)) \partial_t \partial_j v_\lambda \partial_t^3 \partial_i v_\alpha \\
&\quad - 2 \int_0^t \int_{\Gamma_1} \partial_t (\sqrt{g} (g^{ij} g^{kl} - g^{lj} g^{ik}) \partial_j \eta_\alpha \partial_k \eta_\lambda) \partial_t \partial_l v_\lambda \partial_t^3 \partial_i v_\alpha \\
&\quad - \int_0^t \int_{\Gamma_1} \partial_t^2 (\sqrt{g} g^{ij} (\delta_{\alpha\lambda} - g^{kl} \partial_k \eta_\alpha \partial_l \eta_\lambda)) \partial_j v_\lambda \partial_t^3 \partial_i v_\alpha \\
&\quad - \int_0^t \int_{\Gamma_1} \partial_t^2 (\sqrt{g} (g^{ij} g^{kl} - g^{lj} g^{ik}) \partial_j \eta_\alpha \partial_k \eta_\lambda) \partial_l v_\lambda \partial_t^3 \partial_i v_\alpha \\
&= I_{11} + I_{12} + I_{13} + I_{14} + I_{15} + I_{16}.
\end{aligned} \tag{6.7}$$

The main terms are the first two, I_{11} and I_{12} ; the last four terms are treated similarly to each other using integration by parts in time.

We start with I_{11} . Recalling (3.6), we have

$$\begin{aligned}
I_{11} &= -\frac{1}{2} \int_0^t \int_{\Gamma_1} \sqrt{g} g^{ij} \Pi_{\alpha\lambda} \partial_t (\partial_t^2 \partial_j v_\lambda \partial_t^2 \partial_i v_\alpha) \\
&= -\frac{1}{2} \int_{\Gamma_1} \sqrt{g} g^{ij} \Pi_{\alpha\lambda} \partial_t^2 \partial_j v_\lambda \partial_t^2 \partial_i v_\alpha + I_{11,0} + \frac{1}{2} \int_0^t \int_{\Gamma_1} \partial_t (\sqrt{g} g^{ij} \Pi_{\alpha\lambda}) \partial_t^2 \partial_j v_\lambda \partial_t^2 \partial_i v_\alpha,
\end{aligned}$$

where we used the symmetries of g^{-1} and Π in the first equality, integrated by parts in time in the second equality, and denoted

$$I_{11,0} = \frac{1}{2} \int_{\Gamma_1} \sqrt{g} g^{ij} \Pi_{\alpha\lambda} \partial_t^2 \partial_j v_\lambda \partial_t^2 \partial_i v_\alpha \Big|_{t=0}.$$

Using (3.7) to separate $\Pi_{\alpha\lambda} = \Pi_{\alpha\mu} \Pi_{\mu\lambda}$ and writing $\Pi \partial_t^2 \partial_i v = \partial_i (\Pi \partial_t^2 v) - \partial_i \Pi \partial_t^2 v$, we obtain

$$\begin{aligned}
I_{11} &= -\frac{1}{2} \int_{\Gamma_1} \sqrt{g} g^{ij} \partial_i (\Pi_{\alpha\mu} \partial_t^2 v_\alpha) \partial_j (\Pi_{\mu\lambda} \partial_t^2 v_\lambda) + \int_{\Gamma_1} \sqrt{g} g^{ij} \partial_i \Pi_{\alpha\mu} \partial_t^2 v_\alpha \partial_j (\Pi_{\mu\lambda} \partial_t^2 v_\lambda) \\
&\quad - \frac{1}{2} \int_{\Gamma_1} \sqrt{g} g^{ij} \partial_i \Pi_{\alpha\mu} \partial_j \Pi_{\mu\lambda} \partial_t^2 v_\alpha \partial_t^2 v_\lambda + \frac{1}{2} \int_0^t \int_{\Gamma_1} \partial_t (\sqrt{g} g^{ij} \Pi_{\alpha\lambda}) \partial_t^2 \partial_j v_\lambda \partial_t^2 \partial_i v_\alpha + I_{11,0} \\
&= I_{111} + I_{112} + I_{113} + I_{114} + I_{11,0}.
\end{aligned}$$

We proceed to estimate each term. First,

$$\begin{aligned}
I_{111} &= -\frac{1}{2} \int_{\Gamma_1} \delta^{ij} \partial_i (\Pi_{\alpha\mu} \partial_t^2 v_\alpha) \partial_j (\Pi_{\mu\lambda} \partial_t^2 v_\lambda) - \frac{1}{2} \int_{\Gamma_1} (\sqrt{g} g^{ij} - \delta^{ij}) \partial_i (\Pi_{\alpha\mu} \partial_t^2 v_\alpha) \partial_j (\Pi_{\mu\lambda} \partial_t^2 v_\lambda) \\
&= -\frac{1}{2} \|\bar{\partial}(\Pi \partial_t^2 v)\|_{0,\Gamma_1}^2 - \frac{1}{2} \int_{\Gamma_1} (\sqrt{g} g^{ij} - \delta^{ij}) \partial_i (\Pi_{\alpha\mu} \partial_t^2 v_\alpha) \partial_j (\Pi_{\mu\lambda} \partial_t^2 v_\lambda) \\
&\leq -\frac{1}{2} \|\bar{\partial}(\Pi \partial_t^2 v)\|_{0,\Gamma_1}^2 + C \|\sqrt{g} g^{-1} - I\|_{L^\infty(\Gamma_1)} \|\bar{\partial}(\Pi \partial_t^2 v)\|_{0,\Gamma_1}^2 \\
&\leq -\frac{1}{2} \|\bar{\partial}(\Pi \partial_t^2 v)\|_{0,\Gamma_1}^2 + C \|\sqrt{g} g^{-1} - I\|_{1.5,\Gamma_1} \|\bar{\partial}(\Pi \partial_t^2 v)\|_{0,\Gamma_1}^2.
\end{aligned}$$

Writing $\sqrt{g}g^{-1} - I = \int_0^t \partial_t(\sqrt{g}g^{-1} - I)$ and estimating the integrand, we realize that if $T \leq 1/CM$, the integral is bounded by $\tilde{\epsilon}$, and thus we get

$$I_{111} \geq -\frac{1}{4} \|\bar{\partial}(\Pi \partial_t^2 v)\|_{0,\Gamma_1}^2.$$

Next,

$$\begin{aligned} I_{112} &\leq C \|\sqrt{g}g^{-1}\|_{L^\infty(\Gamma_1)} \|\bar{\partial}\Pi\|_{L^\infty(\Gamma_1)} \|\partial_t^2 v\|_{0,\Gamma_1} \|\bar{\partial}(\Pi \partial_t^2 v)\|_{0,\Gamma_1} \\ &\leq C \|\sqrt{g}g^{-1}\|_{1.5,\Gamma_1} \|\bar{\partial}\Pi\|_{1.5,\Gamma_1} \|\partial_t^2 v\|_{0,\Gamma_1} \|\bar{\partial}(\Pi \partial_t^2 v)\|_{0,\Gamma_1} \\ &\leq P(\|q\|_{1.5,\Gamma_1}) \|\partial_t^2 v\|_{0,\Gamma_1} \|\bar{\partial}(\Pi \partial_t^2 v)\|_{0,\Gamma_1} \\ &\leq \tilde{\epsilon} \|\bar{\partial}(\Pi \partial_t^2 v)\|_{0,\Gamma_1}^2 + C_{\tilde{\epsilon}} P(\|q\|_{1.5,\Gamma_1}) \|\partial_t^2 v\|_{0,\Gamma_1}^2, \end{aligned}$$

where in the third inequality we used Remark 3.4 to get

$$\|\sqrt{g}g^{-1}\|_{1.5,\Gamma_1} \leq C$$

and

$$\|\bar{\partial}\Pi\|_{1.5,\Gamma_1} \leq C \|\bar{\partial}\bar{\partial}\eta\|_{1.5,\Gamma_1} \leq C \|\eta\|_{3.5,\Gamma_1}$$

and then invoked Proposition 5.1; also in the fourth line we used the ϵ -Cauchy inequality (so $C_{\tilde{\epsilon}} \rightarrow \infty$ as $\tilde{\epsilon} \rightarrow 0$).

Again, with the help of Remark 3.4, we have

$$I_{113} \leq C \|\bar{\partial}\Pi\|_{L^\infty(\Gamma_1)}^2 \|\partial_t^2 v\|_{0,\Gamma_1}^2 \leq P(\|q\|_{1.5,\Gamma_1}) \|\partial_t^2 v\|_{0,\Gamma_1}^2$$

and

$$I_{114} \leq C \int_0^t \|\partial_t(\sqrt{g}g^{-1}\Pi)\|_{L^\infty(\Gamma_1)} \|\bar{\partial}\partial_t^2 v\|_{0,\Gamma_1}^2 \leq \int_0^t \mathcal{P},$$

recalling the notation (5.1). One also easily obtains

$$I_{11,0} \leq P(\|\bar{\partial}\partial_t^2 v(0)\|_{0,\Gamma_1}) = P(\|\partial_t^2 v(0)\|_{1,\Gamma_1}) \leq \mathcal{P}_0.$$

Hence, choosing $\tilde{\epsilon}$ sufficiently small

$$I_{11} \leq -\frac{1}{C_{11}} \|\bar{\partial}(\Pi \partial_t^2 v)\|_{0,\Gamma_1}^2 + \mathcal{P}_0 + P(\|q\|_{1.5,\Gamma_1}) \|\partial_t^2 v\|_{0,\Gamma_1}^2 + \int_0^t \mathcal{P}, \quad (6.8)$$

for some positive constant C_{11} . In order to remove the term with $\|\partial_t^2 v\|_{0,\Gamma_1}^2$ from the right-hand side of (6.8), we consider the product $\tilde{\mathcal{P}} \|\partial_t^2 v\|_{0,\Gamma_1}^2$, where $\tilde{\mathcal{P}} = \tilde{\mathcal{P}}(t)$ is a polynomial function in the appropriate norms of v and q . Then by

$$\|\partial_t^2 v\|_{0,\Gamma_1} \leq C \|\partial_t^2 v\|_0^{1/2} \|\partial_t^2 v\|_1^{1/2} \leq C \|\partial_t^2 v\|_0^{2/3} \|\partial_t^2 v\|_{1.5}^{1/3} \quad (6.9)$$

we get

$$\begin{aligned} \tilde{\mathcal{P}} \|\partial_t^2 v\|_{0,\Gamma_1}^2 &\leq \tilde{\epsilon} \|\partial_t^2 v\|_{1.5}^2 + \tilde{\mathcal{P}}^3 + C \|\partial_t^2 v\|_0^4 \\ &\leq \tilde{\epsilon} \|\partial_t^2 v\|_{1.5}^2 + \tilde{\mathcal{P}}^3 + C \|\partial_t^2 v_0\|_0^4 + C \left(\int_0^t \|\partial_t^3 v\|_0 \right)^3 \\ &\leq \tilde{\epsilon} \|\partial_t^2 v\|_{1.5}^2 + \tilde{\mathcal{P}}^3 + C \|\partial_t^2 v_0\|_0^4 + CT^2 \int_0^t \|\partial_t^3 v\|_0^3. \end{aligned}$$

By $T \leq C$, we conclude

$$\tilde{\mathcal{P}} \|\partial_t^2 v\|_1^2 \leq \tilde{\epsilon} \|\partial_t^2 v\|_{1.5}^2 + \tilde{\mathcal{P}}^3 + \mathcal{P}_0 + \int_0^t \mathcal{P}. \quad (6.10)$$

Since it is needed below, we note that similar arguments give

$$\tilde{\mathcal{P}} \|\partial_t v\|_{1.5}^2 \leq \tilde{\epsilon} \|\partial_t v\|_{2.5}^2 + \tilde{\mathcal{P}}^a + \mathcal{P}_0 + \int_0^t \mathcal{P}, \quad (6.11)$$

where $a \geq 1$. We also observe that since the power a comes from an application of the Young inequality (e.g. $a = 3$ in (6.10) above), we can choose it so that $\tilde{\mathcal{P}}^a$ is a polynomial if $\tilde{\mathcal{P}}$ is. Finally, we remark that we can obtain similar estimates for other lower order norms of v and q , (e.g. $\|\partial_t v\|_2$ or $\|\partial_t q\|_1$).

Using (6.10), the inequality (6.8) becomes

$$I_{11} \leq -\frac{1}{C_{11}} \|\bar{\partial}(\Pi \partial_t^2 v)\|_{0,\Gamma_1}^2 + \tilde{\epsilon} \|\partial_t^2 v\|_{1.5}^2 + P(\|\partial_t^2 v(0)\|_{1.5}) + P(\|q\|_{1.5,\Gamma_1}) + \int_0^t \mathcal{P}. \quad (6.12)$$

The term I_{12} is estimated with a trick used in [25, p. 868], where the authors observed that we may write

$$I_{12} = \int_0^t \int_{\Gamma_1} \frac{1}{\sqrt{g}} (\partial_t \det A^1 - \det A^2 - \det A^3), \quad (6.13)$$

where

$$A^1 = \begin{bmatrix} \partial_1 \eta_\mu \partial_t^2 \partial_1 v_\mu & \partial_1 \eta_\mu \partial_t^2 \partial_2 v_\mu \\ \partial_2 \eta_\mu \partial_t^2 \partial_1 v_\mu & \partial_2 \eta_\mu \partial_t^2 \partial_2 v_\mu \end{bmatrix}, A^2 = \begin{bmatrix} \partial_1 v_\mu \partial_t^2 \partial_1 v_\mu & \partial_1 \eta_\mu \partial_t^2 \partial_2 v_\mu \\ \partial_2 v_\mu \partial_t^2 \partial_1 v_\mu & \partial_2 \eta_\mu \partial_t^2 \partial_2 v_\mu \end{bmatrix}, A^3 = \begin{bmatrix} \partial_1 \eta_\mu \partial_t^2 \partial_1 v_\mu & \partial_1 v_\mu \partial_t^2 \partial_2 v_\mu \\ \partial_2 \eta_\mu \partial_t^2 \partial_1 v_\mu & \partial_2 v_\mu \partial_t^2 \partial_2 v_\mu \end{bmatrix}.$$

Integrating by parts in time in the first term in (6.13), we get

$$\begin{aligned} I_{12} &= \int_{\Gamma_1} \frac{1}{\sqrt{g}} \det A^1 - \int_{\Gamma_1} \frac{1}{\sqrt{g}} \det A^1 \Big|_{t=0} - \int_0^t \int_{\Gamma_1} \partial_t \left(\frac{1}{\sqrt{g}} \right) \det A^1 - \int_0^t \int_{\Gamma_1} \frac{1}{\sqrt{g}} (\det A^2 + \det A^3) \\ &= I_{121} + I_{12,0} + I_{122} + I_{123}. \end{aligned} \quad (6.14)$$

First,

$$\int_{\Gamma_1} \frac{1}{\sqrt{g}} \det A^1 = \int_{\Gamma_1} Q_{\mu\lambda}^i (\bar{\partial} \eta, \bar{\partial}^2 \eta) \partial_t^2 v_\mu \partial_i \partial_t^2 v_\lambda. \quad (6.15)$$

To see why (6.15) holds, we compute the determinant to find

$$\int_{\Gamma_1} \frac{1}{\sqrt{g}} \det A^1 = \int_{\Gamma_1} \frac{1}{\sqrt{g}} (\partial_1 \eta_\mu \partial_2 \eta_\lambda \partial_1 \partial_t^2 v_\mu \partial_2 \partial_t^2 v_\lambda - \partial_1 \eta_\mu \partial_2 \eta_\lambda \partial_2 \partial_t^2 v_\mu \partial_1 \partial_t^2 v_\lambda).$$

Integrating by parts the ∂_1 derivative in the factor $\partial_1 \partial_t^2 v_\mu$, and the ∂_2 derivative in $\partial_2 \partial_t^2 v_\mu$ produces (6.15) since the terms with four derivatives cancel out. Thus we obtain

$$\begin{aligned} I_{121} &\leq P(\|\bar{\partial}^2 \eta\|_{L^\infty(\Gamma_1)}) \|\partial_t^2 v\|_{0,\Gamma_1} \|\bar{\partial} \partial_t^2 v\|_{0,\Gamma_1} \leq P(\|\eta\|_{3.5,\Gamma_1}) \|\partial_t^2 v\|_{0,\Gamma_1} \|\bar{\partial} \partial_t^2 v\|_{0,\Gamma_1} \\ &\leq \tilde{\epsilon} \|\bar{\partial} \partial_t^2 v\|_{0,\Gamma_1}^2 + C_{\tilde{\epsilon}} P(\|q\|_{1.5,\Gamma_1}) \|\partial_t^2 v\|_{0,\Gamma_1}^2, \end{aligned}$$

where we used Proposition 5.1. Next,

$$\begin{aligned} I_{122} &= - \int_0^t \int_{\Gamma_1} \partial_t \left(\frac{1}{\sqrt{g}} \right) (\partial_1 \eta_\mu \partial_t^2 \partial_1 v_\mu \partial_2 \eta_\lambda \partial_t^2 \partial_2 v_\lambda - \partial_2 \eta_\mu \partial_t^2 \partial_1 v_\mu \partial_1 \eta_\lambda \partial_t^2 \partial_2 v_\lambda) \\ &\leq C \int_0^t \left\| \partial_t \left(\frac{1}{\sqrt{g}} \right) \right\|_{L^\infty(\Gamma_1)} \|\bar{\partial} \eta\|_{L^\infty(\Gamma_1)}^2 \|\bar{\partial} \partial_t^2 v\|_{0,\Gamma_1}^2 \leq \int_0^t \mathcal{P}, \end{aligned}$$

where we used (3.3) to estimate

$$\left\| \partial_t \left(\frac{1}{\sqrt{g}} \right) \right\|_{1.5,\Gamma_1} \leq C \|\partial_t \bar{\partial} \eta\|_{1.5,\Gamma_1} \leq C \|v\|_3.$$

Finally,

$$\begin{aligned} \int_0^t \int_{\Gamma_1} \frac{1}{\sqrt{g}} \det A^2 &= \int_0^t \int_{\Gamma_1} \frac{1}{\sqrt{g}} (\partial_1 v_\mu \partial_t^2 \partial_1 v_\mu \partial_2 \eta_\lambda \partial_t^2 \partial_2 v_\lambda - \partial_2 v_\mu \partial_t^2 \partial_1 v_\mu \partial_1 \eta_\lambda \partial_t^2 \partial_2 v_\lambda) \\ &\leq C \int_0^t \left\| \frac{1}{\sqrt{g}} \right\|_{L^\infty(\Gamma_1)} \|\bar{\partial} v\|_{L^\infty(\Gamma_1)}^2 \|\bar{\partial} \partial_t^2 v\|_{0,\Gamma_1}^2 \leq \int_0^t \mathcal{P} \end{aligned}$$

by (3.4). The term with $\det A^3$ is estimated the same way, and thus $I_{123} \leq \int_0^t \mathcal{P}$. Since also $I_{12,0} \leq P(\|\partial_t^2 v(0)\|_{1.5})$, we conclude that

$$I_{12} \leq \tilde{\epsilon} \|\bar{\partial} \partial_t^2 v\|_{0,\Gamma_1}^2 + \tilde{\epsilon} \|\partial_t^2 v\|_{1.5}^2 + \mathcal{P}_0 + P(\|q\|_{1.5,\Gamma_1}) + \int_0^t \mathcal{P}, \quad (6.16)$$

where we used (6.10).

The terms I_{13} and I_{14} are estimated analogously, so we only show the details for I_{13} . To treat I_{13} (and I_{14}), it suffices to keep track of the general multiplicative structure of the integrands, and for this we use the symbolic notation in Remark 3.4 to write

$$I_{13} = - \int_0^t \int_{\Gamma_1} \partial_t Q(\bar{\partial}\eta) \partial_t \bar{\partial} v \partial_t^3 \bar{\partial} v.$$

From (3.1) we have

$$\partial_t^2 Q(\bar{\partial}\eta) = Q(\bar{\partial}\eta) (\partial_t \bar{\partial}\eta)^2 + Q(\bar{\partial}\eta) \partial_t^2 \bar{\partial}\eta = Q(\bar{\partial}\eta) (\bar{\partial}v)^2 + Q(\bar{\partial}\eta) \partial_t \bar{\partial}v, \quad (6.17)$$

so that integration by parts in t gives

$$\begin{aligned} I_{13} &= - \int_{\Gamma_1} \partial_t Q(\bar{\partial}\eta) \partial_t \bar{\partial} v \partial_t^2 \bar{\partial} v + I_{13,0} \\ &\quad + \int_0^t \int_{\Gamma_1} \partial_t^2 Q(\bar{\partial}\eta) \partial_t \bar{\partial} v \partial_t^2 \bar{\partial} v + \int_0^t \int_{\Gamma_1} \partial_t Q(\bar{\partial}\eta) \partial_t^2 \bar{\partial} v \partial_t^2 \bar{\partial} v, \end{aligned}$$

where

$$I_{13,0} = \int_{\Gamma_1} \partial_t Q(\bar{\partial}\eta) \partial_t \bar{\partial} v \partial_t^2 \bar{\partial} v \Big|_{t=0} \leq \mathcal{P}_0.$$

Thus, recalling (3.3) and using (6.17),

$$\begin{aligned} I_{13} &\leq C \|\partial_t \bar{\partial}\eta\|_{L^\infty(\Gamma_1)} \|\partial_t \bar{\partial} v\|_{0,\Gamma_1} \|\partial_t^2 \bar{\partial} v\|_{0,\Gamma_1} + \mathcal{P}_0 + C \int_0^t \|\bar{\partial} v\|_{L^\infty(\Gamma_1)}^2 \|\partial_t \bar{\partial} v\|_{0,\Gamma_1} \|\partial_t^2 \bar{\partial} v\|_{0,\Gamma_1} \\ &\quad + C \int_0^t \|\partial_t \bar{\partial} v\|_{L^4(\Gamma_1)}^2 \|\partial_t^2 \bar{\partial} v\|_{0,\Gamma_1} + C \int_0^t \|\partial_t \bar{\partial}\eta\|_{L^\infty(\Gamma_1)} \|\partial_t^2 \bar{\partial} v\|_{0,\Gamma_1}^2 \\ &\leq \tilde{\epsilon} \|\partial_t^2 v\|_{1.5}^2 + \mathcal{P}_0 + C_\epsilon \|\partial_t v\|_{1.5}^2 \|v\|_{2.5+\delta}^2 + \int_0^t \mathcal{P}. \end{aligned} \quad (6.18)$$

It follows that the inequality (6.18) becomes

$$I_{13} \leq \tilde{\epsilon} \|\partial_t^2 v\|_{1.5}^2 + \mathcal{P}_0 + P(\|v\|_{2.5+\delta}) + C \int_0^t (1 + \|\partial_t v\|_{1.5}) \|\partial_t v\|_{1.5} \|\partial_t^2 v\|_{1.5} + \int_0^t \mathcal{P}. \quad (6.19)$$

As pointed out above, the same inequality holds for I_{14} as well.

The remaining terms in I_1 , which are I_{15} and I_{16} , are estimated analogously, so we only estimate I_{15} . It again suffices to keep track of the general multiplicative structure of the integrands and we use the symbolic notation of Remark 3.4. From (3.1) and (6.17),

$$\partial_t^3 Q(\bar{\partial}\eta) = Q(\bar{\partial}\eta) (\partial_t \bar{\partial}\eta)^3 + Q(\bar{\partial}\eta) \partial_t \bar{\partial}\eta \partial_t^2 \bar{\partial}\eta + Q(\bar{\partial}\eta) \partial_t^3 \bar{\partial}\eta \quad (6.20)$$

and thus, using integration by parts in time,

$$\begin{aligned} I_{15} &= I_{15,0} - \int_{\Gamma_1} \partial_t^2 Q(\bar{\partial}\eta) \bar{\partial} v \partial_t^2 \bar{\partial} v + \int_0^t \int_{\Gamma_1} \partial_t^3 Q(\bar{\partial}\eta) \bar{\partial} v \partial_t^2 \bar{\partial} v + \int_0^t \int_{\Gamma_1} \partial_t^2 Q(\bar{\partial}\eta) \partial_t \bar{\partial} v \partial_t^2 \bar{\partial} v \\ &= I_{15,0} - \int_{\Gamma_1} Q(\bar{\partial}\eta) (\partial_t \bar{\partial}\eta)^2 \bar{\partial} v \partial_t^2 \bar{\partial} v - \int_{\Gamma_1} Q(\bar{\partial}\eta) \partial_t^2 \bar{\partial}\eta \bar{\partial} v \partial_t^2 \bar{\partial} v + \int_0^t \int_{\Gamma_1} Q(\bar{\partial}\eta) (\partial_t \bar{\partial}\eta)^3 \bar{\partial} v \partial_t^2 \bar{\partial} v \\ &\quad + \int_0^t \int_{\Gamma_1} Q(\bar{\partial}\eta) \partial_t \bar{\partial}\eta \partial_t^2 \bar{\partial}\eta \bar{\partial} v \partial_t^2 \bar{\partial} v + \int_0^t \int_{\Gamma_1} Q(\bar{\partial}\eta) \partial_t^3 \bar{\partial}\eta \bar{\partial} v \partial_t^2 \bar{\partial} v + \int_0^t \int_{\Gamma_1} \partial_t^2 Q(\bar{\partial}\eta) \partial_t \bar{\partial} v \partial_t^2 \bar{\partial} v \\ &= I_{15,0} + I_{151} + I_{152} + I_{153} + I_{154} + I_{155} + I_{156}, \end{aligned}$$

where we set

$$I_{15,0} = \int_{\Gamma_1} \partial_t^2 Q(\bar{\partial}\eta) \bar{\partial}v \partial_t^2 \bar{\partial}v \Big|_{t=0} \leq \mathcal{P}_0.$$

We have

$$I_{151} \leq C \|\bar{\partial}v\|_{L^\infty(\Gamma_1)}^3 \|\partial_t^2 \bar{\partial}v\|_{0,\Gamma_1} \leq \tilde{\epsilon} \|\partial_t^2 v\|_{1.5}^2 + P(\|v\|_{2.5+\delta}),$$

as well as

$$I_{152} \leq C \|\bar{\partial}\partial_t v\|_{0,\Gamma_1} \|\bar{\partial}\partial_t^2 v\|_{0,\Gamma_1} \|\bar{\partial}v\|_{L^\infty(\Gamma_1)} \leq \tilde{\epsilon} \|\partial_t^2 v\|_{1.5}^2 + C \|\partial_t v\|_{1.5}^2 + P(\|v\|_{2.5+\delta}),$$

and

$$I_{153} + I_{154} + I_{155} \leq \int_0^t \mathcal{P}.$$

The term I_{156} has already been dealt with in the estimate of I_{13} and obeys (6.19). Thus, from the above we obtain

$$I_{15} \leq \tilde{\epsilon} \|\partial_t^2 v\|_{1.5}^2 + \mathcal{P}_0 + P(\|v\|_{2.5+\delta}) + C \|\partial_t v\|_{1.5}^2 + \int_0^t \mathcal{P}. \quad (6.21)$$

Invoking (6.11) and (6.21) gives

$$I_{15} + I_{16} \leq \tilde{\epsilon} \|\partial_t^2 v\|_{1.5}^2 + \mathcal{P}_0 + P(\|v\|_{2.5+\delta}) + \int_0^t \mathcal{P}. \quad (6.22)$$

Combining (6.12), (6.16), (6.19), and (6.22), and recalling (6.5) and (6.7), we obtain

$$I_1 \leq -\frac{1}{C_{11}} \|\bar{\partial}(\Pi\partial_t^2 v)\|_{0,\Gamma_1}^2 + \tilde{\epsilon} \|\partial_t^2 v\|_{1.5}^2 + P(\|q\|_{1.5,\Gamma_1}, \|v\|_{2.5+\delta}) + \mathcal{P}_0 + \int_0^t \mathcal{P}. \quad (6.23)$$

Estimate of I_2 : Now, we consider I_2 , rewritten in (6.6). Differentiating (2.3b) in time three times gives

$$a_{\mu\alpha} \partial_\mu \partial_t^3 v_\alpha = -3\partial_t^2 a_{\mu\alpha} \partial_\mu \partial_t v_\alpha - 3\partial_t a_{\mu\alpha} \partial_\mu \partial_t^2 v_\alpha - \partial_t^3 a_{\mu\alpha} \partial_\mu v_\alpha,$$

so that I_{21} becomes

$$\begin{aligned} I_{21} &= -3 \int_0^t \int_\Omega \partial_t^2 a_{\mu\alpha} \partial_\mu \partial_t v_\alpha \partial_t^3 q - 3 \int_0^t \int_\Omega \partial_t a_{\mu\alpha} \partial_\mu \partial_t^2 v_\alpha \partial_t^3 q - \int_0^t \int_\Omega \partial_t^3 a_{\mu\alpha} \partial_\mu v_\alpha \partial_t^3 q \\ &= I_{211} + I_{212} + I_{213}. \end{aligned} \quad (6.24)$$

We have

$$\begin{aligned} I_{211} &= -3 \int_\Omega \partial_t^2 a_{\mu\alpha} \partial_\mu \partial_t v_\alpha \partial_t^2 q \Big|_0^t + 3 \int_0^t \int_\Omega \partial_t (\partial_t^2 a_{\mu\alpha} \partial_\mu \partial_t v_\alpha) \partial_t^2 q \\ &\leq C \|\partial_t^2 a\|_0 \|\partial\partial_t v\|_{L^3(\Omega)} \|\partial_t^2 q\|_{L^6(\Omega)} \Big|_0^t + C \int_0^t \|\partial_t^3 a\|_{L^3(\Omega)} \|\partial\partial_t v\|_{L^6(\Omega)} \|\partial_t^2 q\|_{L^2(\Omega)} \\ &\quad + C \int_0^t \|\partial_t^2 a\|_{L^6(\Omega)} \|\partial\partial_t^2 v\|_{L^3(\Omega)} \|\partial_t^2 q\|_{L^2(\Omega)} \\ &\leq C (\|\partial v\|_0 \|\partial v\|_{L^\infty(\Omega)} + \|\partial\partial_t v\|_0) \|\partial\partial_t v\|_{0.5} \|\partial_t^2 q\|_1 \Big|_0^t \\ &\quad + C \int_0^t (\|\partial v\|_{L^3(\Omega)} \|\partial v\|_{L^\infty(\Omega)}^2 + \|\partial\partial_t v\|_{L^3(\Omega)} \|\partial v\|_{L^\infty(\Omega)} + \|\partial\partial_t^2 v\|_{L^3(\Omega)}) \|\partial_t v\|_2 \|\partial_t^2 q\|_1 \\ &\quad + C \int_0^t (\|\partial v\|_1 \|\partial v\|_{L^\infty(\Omega)} + \|\partial\partial_t v\|_1) \|\partial_t^2 v\|_{1.5} \|\partial_t^2 q\|_1 \\ &\leq \tilde{\epsilon} (\|\partial_t^2 q\|_1^2 + \|\partial_t v\|_{2.5}^2) + P(\|v\|_{2.5+\delta}) + \mathcal{P}_0 + \int_0^t \mathcal{P}, \end{aligned} \quad (6.25)$$

where in the last step we used an argument similar to that for (6.11). Next, we integrate by parts and use (2.2) to find

$$\begin{aligned} I_{212} &= -3 \int_0^t \int_{\partial\Omega} \partial_t a_{3\alpha} \partial_t^2 v_\alpha \partial_t^3 q N_3 + 3 \int_0^t \int_{\Omega} \partial_t a_{\mu\alpha} \partial_t^2 v_\alpha \partial_\mu \partial_t^3 q \\ &= I_{2121} + I_{2122}. \end{aligned} \quad (6.26)$$

By (2.3c) and (2.3e), we have

$$I_{2121} = 3 \int_0^t \int_{\Gamma_1} a_{3\gamma} \partial_\nu v_\gamma a_{\nu\alpha} \partial_t^2 v_\alpha \partial_t^3 q. \quad (6.27)$$

Since in light of (2.3d) and (3.8),

$$a_{3\gamma} \partial_t^3 q = -\sigma \partial_t^3 (\sqrt{g} \Delta_g \eta_\gamma) - 3 \partial_t a_{3\gamma} \partial_t^2 q - 3 \partial_t^2 a_{3\gamma} \partial_t q - \partial_t^3 a_{3\gamma} q,$$

we obtain

$$\begin{aligned} I_{2121} &= -3\sigma \int_0^t \int_{\Gamma_1} \partial_\nu v_\gamma a_{\nu\alpha} \partial_t^2 v_\alpha \partial_t^3 (\sqrt{g} \Delta_g \eta_\gamma) - 9 \int_0^t \int_{\Gamma_1} \partial_\nu v_\gamma a_{\nu\alpha} \partial_t^2 v_\alpha \partial_t a_{3\gamma} \partial_t^2 q \\ &\quad - 9 \int_0^t \int_{\Gamma_1} \partial_\nu v_\gamma a_{\nu\alpha} \partial_t^2 v_\alpha \partial_t^2 a_{3\gamma} \partial_t q - 3 \int_0^t \int_{\Gamma_1} \partial_\nu v_\gamma a_{\nu\alpha} \partial_t^2 v_\alpha \partial_t^3 a_{3\gamma} q \\ &= I_{21211} + I_{21212} + I_{21213} + I_{21214}. \end{aligned} \quad (6.28)$$

In order to estimate I_{21211} , we use (3.10) and $\partial_t \eta = v$ to write

$$\begin{aligned} I_{21211} &= -3\sigma \int_0^t \int_{\Gamma_1} \partial_i (\sqrt{g} g^{ij} (\delta_{\gamma\lambda} - g^{kl} \partial_k \eta_\gamma \partial_l \eta_\lambda)) \partial_t^2 \partial_j v_\lambda \partial_\nu v_\gamma a_{\nu\alpha} \partial_t^2 v_\alpha \\ &\quad - 3\sigma \int_0^t \int_{\Gamma_1} \partial_i (\sqrt{g} (g^{ij} g^{kl} - g^{lj} g^{ik})) \partial_j \eta_\gamma \partial_k \eta_\lambda \partial_t^2 \partial_l v_\lambda \partial_\nu v_\gamma a_{\nu\alpha} \partial_t^2 v_\alpha \\ &\quad - 6\sigma \int_0^t \int_{\Gamma_1} \partial_i (\partial_t (\sqrt{g} g^{ij} (\delta_{\gamma\lambda} - g^{kl} \partial_k \eta_\gamma \partial_l \eta_\lambda))) \partial_t \partial_j v_\lambda \partial_\nu v_\gamma a_{\nu\alpha} \partial_t^2 v_\alpha \\ &\quad - 6\sigma \int_0^t \int_{\Gamma_1} \partial_i (\partial_t (\sqrt{g} (g^{ij} g^{kl} - g^{lj} g^{ik})) \partial_j \eta_\gamma \partial_k \eta_\lambda) \partial_t \partial_l v_\lambda \partial_\nu v_\gamma a_{\nu\alpha} \partial_t^2 v_\alpha \\ &\quad - 3\sigma \int_0^t \int_{\Gamma_1} \partial_i (\partial_t^2 (\sqrt{g} g^{ij} (\delta_{\gamma\lambda} - g^{kl} \partial_k \eta_\gamma \partial_l \eta_\lambda))) \partial_j v_\lambda \partial_\nu v_\gamma a_{\nu\alpha} \partial_t^2 v_\alpha \\ &\quad - 3\sigma \int_0^t \int_{\Gamma_1} \partial_i (\partial_t^2 (\sqrt{g} (g^{ij} g^{kl} - g^{lj} g^{ik})) \partial_j \eta_\gamma \partial_k \eta_\lambda) \partial_l v_\lambda \partial_\nu v_\gamma a_{\nu\alpha} \partial_t^2 v_\alpha. \end{aligned}$$

In each of the above integrals, we integrate by parts the derivative ∂_i . After this, the sum of the first two integrals has the form $3\sigma \int_0^t \int_{\Gamma_1} Q(\bar{\partial}\eta) \partial_t^2 \bar{\partial} v \bar{\partial} (\bar{\partial} v a \partial_t^2 v)$ (cf. Remark 3.4), the sum of the middle terms is of the form $6\sigma \int_0^t \int_{\Gamma_1} \partial_t Q(\bar{\partial}\eta) \partial_t \bar{\partial} v \bar{\partial} (\bar{\partial} v a \partial_t^2 v)$, while the sum of the last two is $3\sigma \int_0^t \int_{\Gamma_1} \partial_t^2 Q(\bar{\partial}\eta) \bar{\partial} v \bar{\partial} (\bar{\partial} v a \partial_t^2 v)$. Each of the three forms is bounded by $\int_0^t \mathcal{P}$, using Hölder and Sobolev inequalities, and we obtain

$$I_{21211} \leq \int_0^t \mathcal{P}. \quad (6.29)$$

The terms I_{21212} and I_{21213} are also bounded using Hölder and Sobolev inequalities, leading to

$$I_{21212}, I_{21213} \leq \int_0^t \mathcal{P}.$$

For instance, we have

$$\begin{aligned} I_{21213} &= -9 \int_0^t \int_{\Gamma_1} \partial_\nu v_\gamma a_{\nu\alpha} \partial_t^2 v_\alpha \partial_t^2 a_{3\gamma} \partial_t q \\ &\leq C \int_0^t \|\partial v\|_{L^6(\Gamma_1)} \|a\|_{L^\infty(\Gamma_1)} \|\partial_t^2 v\|_{L^6(\Gamma_1)} \|\partial_t^2 a\|_{0,\Gamma_1} \|\partial_t q\|_{L^6(\Gamma_1)} \leq \int_0^t \mathcal{P}. \end{aligned} \quad (6.30)$$

The term I_{21214} can not be estimated immediately due to the factor $\partial_t^3 a$. First, we write

$$\begin{aligned} I_{21214} &= -3 \int_0^t \int_{\Gamma_1} \partial_\nu v_\gamma a_{\nu\alpha} \partial_t^2 v_\alpha \partial_t^3 a_{3\gamma} q = -3 \int_0^t \int_{\Gamma_1} \partial_\nu v_\gamma a_{\nu\alpha} \partial_t^2 v_\alpha \partial_t^3 a_{3\gamma} q \\ &\leq C \sum_{\gamma=1}^3 \int_0^t \|\partial v\|_{L^\infty(\Gamma_1)} \|a\|_{L^\infty(\Gamma_1)} \|\partial_t^2 v\|_{0,\Gamma_1} \|\partial_t^3 a_{3\gamma}\|_{0,\Gamma_1} \|q\|_{L^\infty(\Gamma_1)} \\ &\leq C \sum_{\gamma=1}^3 \int_0^t \|v\|_3 \|\partial_t^2 v\|_{1.5} \|\partial_t^3 a_{3\gamma}\|_{0,\Gamma_1} \|q\|_3. \end{aligned}$$

To bound $\partial_t^3 a_{3\gamma}$, we use that $a_{3\gamma}$ contains only tangential derivatives of η , i.e., from (3.13) we see that

$$a_{3\gamma} = \epsilon_{\gamma\alpha\beta} \partial_1 \eta_\alpha \partial_2 \eta_\beta, \quad (6.31)$$

where $\epsilon_{\gamma\alpha\beta}$ is the Levi-Civita symbol, so that

$$\|\partial_t^3 a_{3\gamma}\|_{0,\Gamma_1} \leq C \|\bar{\partial} \partial_t^2 v \bar{\partial} \eta\|_{0,\Gamma_1} + C \|\bar{\partial} \partial_t v \bar{\partial} v\|_{0,\Gamma_1} \leq P(\|\partial_t^2 v\|_{1.5}, \|\partial_t v\|_{1.5}, \|v\|_3).$$

Hence, $I_{21214} \leq \int_0^t \mathcal{P}$, and we obtain

$$I_{2121} \leq \int_0^t \mathcal{P}. \quad (6.32)$$

The term I_{2122} is estimated using integration by parts in t and obeys

$$\begin{aligned} I_{2122} &\leq C \|\partial_t^2 q\|_1 \|\partial_t^2 v\|_{L^3(\Omega)} \|\partial v\|_{L^6(\Omega)} + P(\|\partial_t^2 q(0)\|_1, \|\partial_t^2 v(0)\|_{1.5}, \|v(0)\|_3) + \int_0^t \mathcal{P} \\ &\leq C \|\partial_t^2 q\|_1 \|\partial_t^2 v\|_{0.5} \|v\|_2 + P(\|\partial_t^2 q(0)\|_1, \|\partial_t^2 v(0)\|_{1.5}, \|v(0)\|_3) + \int_0^t \mathcal{P} \\ &\leq \tilde{\epsilon} (\|\partial_t^2 q\|_1^2 + \|\partial_t^2 v\|_{1.5}^2) + \mathcal{P}_0 + P(\|v\|_2) + \int_0^t \mathcal{P}, \end{aligned} \quad (6.33)$$

where in the last line we estimated similarly to (6.10) and (6.11).

Combining (6.26), (6.32), and (6.33), we obtain

$$I_{212} \leq \tilde{\epsilon} (\|\partial_t^2 q\|_1^2 + \|\partial_t^2 v\|_{1.5}^2) + \mathcal{P}_0 + P(\|v\|_2) + \int_0^t \mathcal{P}. \quad (6.34)$$

Next we estimate I_{213} . Using (2.3c), we find

$$I_{213} = - \int_0^t \int_{\Omega} \partial_t^3 a_{\mu\alpha} \partial_\mu v_\alpha \partial_t^3 q = \int_0^t \int_{\Omega} a_{\mu\gamma} \partial_t^2 \partial_\beta v_\gamma a_{\beta\alpha} \partial_\mu v_\alpha \partial_t^3 q + R,$$

where R represents the sum of the lower order terms, which are easily estimated as

$$R \leq \tilde{\epsilon} (\|\partial_t^2 q\|_1^2 + \|\partial_t^2 v\|_{1.5}^2) + \mathcal{P}_0 + P(\|v\|_{2.5+\delta}) + \int_0^t \mathcal{P}. \quad (6.35)$$

Integrating by parts the derivative ∂_β , we have

$$\begin{aligned} \int_0^t \int_{\Omega} a_{\mu\gamma} \partial_t^2 \partial_\beta v_\gamma a_{\beta\alpha} \partial_\mu v_\alpha \partial_t^3 q &= - \int_0^t \int_{\Omega} a_{\mu\gamma} \partial_t^2 v_\gamma a_{\beta\alpha} \partial_\beta \partial_\mu v_\alpha \partial_t^3 q - \int_0^t \int_{\Omega} a_{\mu\gamma} \partial_t^2 v_\gamma a_{\beta\alpha} \partial_\mu v_\alpha \partial_\beta \partial_t^3 q \\ &\quad - \int_0^t \int_{\Omega} \partial_\beta a_{\mu\gamma} \partial_t^2 v_\gamma a_{\beta\alpha} \partial_\mu v_\alpha \partial_t^3 q + \int_0^t \int_{\Gamma_1} a_{\mu\gamma} \partial_t^2 v_\gamma a_{3\alpha} \partial_\mu v_\alpha \partial_t^3 q, \end{aligned}$$

where we used (2.2) and (2.3e). The first three integrals on the right-hand side are estimated using a simple integration by parts in time and are bounded by

$$\tilde{\epsilon}(\|\partial_t^2 q\|_1 + \|\partial_t^2 v\|_{1.5}^2) + \mathcal{P}_0 + P(\|v\|_{2.5+\delta}) + \int_0^t \mathcal{P}, \quad (6.36)$$

whereas the last term is estimated as I_{2121} (see (6.27) and (6.28)). Thus from (6.35) and (6.36) we have

$$I_{213} \leq \tilde{\epsilon}(\|\partial_t^2 q\|_1 + \|\partial_t^2 v\|_{1.5}^2) + \mathcal{P}_0 + P(\|v\|_{2.5+\delta}) + \int_0^t \mathcal{P}. \quad (6.37)$$

Combining (6.24), (6.25), (6.34), and (6.37), we find

$$I_{21} \leq \tilde{\epsilon}(\|\partial_t^2 q\|_1^2 + \|\partial_t^2 v\|_{1.5}^2) + \mathcal{P}_0 + P(\|v\|_{2.5+\delta}) + \int_0^t \mathcal{P}. \quad (6.38)$$

Next, we treat I_{22} , defined in (6.6), which we write as

$$\begin{aligned} I_{22} &= 3 \int_0^t \int_{\Omega} \partial_t a_{\mu\alpha} \partial_t^2 q \partial_{\mu} \partial_t^3 v_{\alpha} \\ &= 3 \int_0^t \int_{\Gamma_1} \partial_t a_{3\alpha} \partial_t^2 q \partial_t^3 v_{\alpha} - 3 \int_0^t \int_{\Omega} \partial_t a_{\mu\alpha} \partial_{\mu} \partial_t^2 q \partial_t^3 v_{\alpha} \\ &= -3 \int_0^t \int_{\Gamma_1} a_{3\gamma} \partial_{\nu} v_{\gamma} a_{\nu\alpha} \partial_t^2 q \partial_t^3 v_{\alpha} - 3 \int_0^t \int_{\Omega} \partial_t a_{\mu\alpha} \partial_{\mu} \partial_t^2 q \partial_t^3 v_{\alpha} \\ &= 3\sigma \int_0^t \int_{\Gamma_1} \partial_{\nu} v_{\gamma} a_{\nu\alpha} \partial_t^2 (\sqrt{g} \Delta_g \eta_{\gamma}) \partial_t^3 v_{\alpha} + 6 \int_0^t \int_{\Gamma_1} \partial_{\nu} v_{\gamma} a_{\nu\alpha} \partial_t a_{3\gamma} \partial_t q \partial_t^3 v_{\alpha} \\ &\quad + 3 \int_0^t \int_{\Gamma_1} \partial_{\nu} v_{\gamma} a_{\nu\alpha} \partial_t^2 a_{3\gamma} q \partial_t^3 v_{\alpha} - 3 \int_0^t \int_{\Omega} \partial_t a_{\mu\alpha} \partial_{\mu} \partial_t^2 q \partial_t^3 v_{\alpha} \\ &= I_{221} + I_{222} + I_{223} + I_{224}, \end{aligned}$$

where in the second equality we integrated by parts in ∂_{μ} , in the third equality we used (2.3c), and in the fourth equality (2.3d). In the second inequality, we also applied (6.3).

To estimate I_{221} we use the identity (3.10). As above, it suffices to consider the general multiplicative structure of the integrands and thus we write the right-hand side of (3.10) as

$$\bar{\partial}(Q(\bar{\partial}\eta)\partial_t\bar{\partial}\eta). \quad (6.39)$$

Integrating by parts the $\bar{\partial}$ derivative in (6.39) and integrating by parts in time one of the derivatives in the $\partial_t^3 v$ term, we find

$$\begin{aligned} I_{221} &= \int_0^t \int_{\Gamma_1} \partial_{\nu} a \partial_t \bar{\partial}(Q(\bar{\partial}\eta)\partial_t\bar{\partial}\eta) \partial_t^3 v \\ &= - \int_0^t \int_{\Gamma_1} \partial_t(Q(\bar{\partial}\eta)\partial_t\bar{\partial}\eta) (\partial_t^3 \bar{\partial} v \partial_{\nu} a + \partial_t^3 v \bar{\partial}(\partial_{\nu} a)) \\ &= - \int_{\Gamma_1} \partial_t(Q(\bar{\partial}\eta)\partial_t\bar{\partial}\eta) \partial_t^2 \bar{\partial} v \partial_{\nu} a \Big|_0^t - \int_{\Gamma_1} \partial_t(Q(\bar{\partial}\eta)\partial_t\bar{\partial}\eta) \partial_t^2 v \bar{\partial}(\partial_{\nu} a) \Big|_0^t \\ &\quad + \int_0^t \int_{\Gamma_1} \partial_t^2(Q(\bar{\partial}\eta)\partial_t\bar{\partial}\eta) \partial_t^2 \bar{\partial} v a \partial_{\nu} + \int_0^t \int_{\Gamma_1} \partial_t(Q(\bar{\partial}\eta)\partial_t\bar{\partial}\eta) \partial_t^2 \bar{\partial} v \partial_t(a \partial_{\nu}) \\ &\quad + \int_0^t \int_{\Gamma_1} \partial_t^2(Q(\bar{\partial}\eta)\partial_t\bar{\partial}\eta) \partial_t^2 v \bar{\partial}(a \partial_{\nu}) + \int_0^t \int_{\Gamma_1} \partial_t(Q(\bar{\partial}\eta)\partial_t\bar{\partial}\eta) \partial_t^2 v \partial_t \bar{\partial}(a \partial_{\nu}). \end{aligned}$$

The estimate of these terms is done in a similar way as other boundary integrals handled above. For example, the first of the two pointwise terms above equals

$$\begin{aligned} & - \int_{\Gamma_1} Q(\bar{\partial}\eta) \partial_t^2 \bar{\partial}\eta \partial_t^2 \bar{\partial}v \partial va \Big|_0^t - \int_{\Gamma_1} Q(\bar{\partial}\eta) (\partial_t \bar{\partial}\eta)^2 \partial_t^2 \bar{\partial}v \partial va \Big|_0^t \\ & = - \int_{\Gamma_1} Q(\bar{\partial}\eta) \partial_t \bar{\partial}v \partial_t^2 \bar{\partial}v \partial va \Big|_0^t - \int_{\Gamma_1} Q(\bar{\partial}\eta) (\bar{\partial}v)^2 \partial_t^2 \bar{\partial}v \partial va \Big|_0^t \\ & \leq \tilde{\epsilon} \|\partial_t^2 v\|_{1.5}^2 + \mathcal{P}_0 + P(\|v\|_{2.5+\delta}) + \int_0^t P(\|v\|_3, \|\partial_t^2 v\|_{1.5}). \end{aligned}$$

Other terms are handled similarly; we do not present the estimates here as they consist of a repetition of ideas used above. We obtain

$$I_{221} \leq \tilde{\epsilon} \|\partial_t^2 v\|_{1.5}^2 + \mathcal{P}_0 + P(\|v\|_{2.5+\delta}) + \int_0^t \mathcal{P}.$$

The terms I_{222} , I_{223} , and I_{224} are also bounded by essentially a repetition of the ideas presented so far and, thus, we again omit the details. We point out that a term in $\partial_t^3 a$ appears in I_{223} after integration by parts in time. But this term is then handled as in I_{21214} , with the help of (6.31). The final estimate for I_{22} is

$$I_{22} \leq \tilde{\epsilon} (\|\partial_t^2 v\|_{1.5}^2 + \|\partial_t^2 q\|_1^2) + P(\|v\|_{2.5+\delta}) + \mathcal{P}_0 + \int_0^t \mathcal{P}. \quad (6.40)$$

Finally, the terms I_{23} and I_{24} are treated similarly to the terms I_{21} with the corresponding estimates similar to the ones that have already been performed above. The final estimate for $I_{23} + I_{24}$ reads

$$I_{23} + I_{24} \leq \tilde{\epsilon} (\|\partial_t^2 v\|_{1.5}^2 + \|\partial_t^2 q\|_1^2) + P(\|v\|_{2.5+\delta}) + \mathcal{P}_0 + \int_0^t \mathcal{P}. \quad (6.41)$$

Collecting all the bounds above then concludes the proof of the lemma. \square

7. ESTIMATES AT $t = 0$

The above estimates involve several quantities evaluated at time zero. In this section we show that all such quantities may be estimated in terms of the initial data.

From (1.1) we have that p satisfies

$$\begin{aligned} \Delta p &= -\operatorname{div}(\nabla_u u) \quad \text{in } \mathcal{D}, \\ \frac{\partial p}{\partial 3} &= 0 \\ p &= \sigma \mathcal{H} \quad \text{on } \Gamma_1(t), \end{aligned}$$

where $\Gamma_1(t) = \eta(\Gamma_1)$. Evaluating at $t = 0$ and denoting $q_0 = q(0)$ gives

$$\begin{aligned} \Delta q_0 &= -\partial_\alpha v_{0\beta} \partial_\beta v_{0\alpha} \quad \text{in } \Omega_0, \\ \partial_3 p &= 0 \quad \text{on } \Gamma_0, \\ p &= 0 \quad \text{on } \Gamma_1, \end{aligned}$$

where we used $\mathcal{H}(0) = 0$ on Γ_1 . We thus obtain the estimate

$$\|q_0\|_4 \leq C \|v_0\|_3^2.$$

Evaluating (2.3a) at $t = 0$ and using $a(0) = I$ produces

$$\|\partial_t v(0)\|_3 \leq C \|q_0\|_4 \leq C \|v_0\|_3^2.$$

From (2.3c) and its time derivative we also obtain

$$\|\partial_t a(0)\|_2 \leq C \|v_0\|_3$$

and

$$\|\partial_t^2 a(0)\|_2 \leq P(\|v_0\|_3),$$

after using the estimate for $\partial_t v(0)$.

From (4.1) and (4.2), and also using (2.2) and (2.3d), we may write

$$\begin{aligned}\partial_\mu(a_{\mu\nu}a_{\lambda\nu}\partial_\lambda q) &= \partial_t a_{\mu\lambda}\partial_\mu v_\lambda && \text{in } \Omega_0, \\ a_{3\nu}a_{\lambda\nu}\partial_\lambda q &= -a_{3\nu}\partial_t v_\nu && \text{on } \Gamma_0, \\ a_{3\alpha}q &= -\sigma\sqrt{g}\Delta_g\eta_\alpha && \text{on } \Gamma_1.\end{aligned}$$

Setting $\alpha = 3$ in the boundary condition on Γ_1 , differentiating in time, and evaluating at $t = 0$ yields

$$\begin{aligned}\Delta\partial_t q(0) &= (-\partial_t a_{\alpha\mu}\partial_{\mu\alpha}^2 q - \partial_t\partial_\alpha a_{\beta\alpha}\partial_\beta q - \partial_t a_{\beta\mu}\partial_{\mu\beta}^2 q \\ &\quad + \partial_t^2 a_{\alpha\beta}\partial_\alpha v_\beta + \partial_t a_{\alpha\beta}\partial_\alpha\partial_t v_\beta)\Big|_{t=0} && \text{in } \Omega_0, \\ \partial_3\partial_t q(0) &= -\partial_t a_{3\beta}\partial_\beta q\Big|_{t=0} && \text{on } \Gamma_0, \\ \partial_t q(0) &= -\partial_t a_{3\alpha}q\Big|_{t=0} - \sigma\overline{\Delta}v_{03} && \text{on } \Gamma_1,\end{aligned}$$

where we used $a(0) = I$, (2.2), and (6.3). Above, $\overline{\Delta}$ is the flat Laplacian on the boundary. Using the estimates we have already obtained at $t = 0$ and invoking the elliptic theory, we obtain

$$\|\partial_t q(0)\|_{2.5} \leq P(\|v_0\|_3, \|v_{03}\|_{4,\Gamma_1})$$

and thus $\|\partial_{tt}v(0)\|_{1.5} \leq P(\|v_0\|_3, \|v_{03}\|_{4,\Gamma_1})$ by differentiating (2.3a) and evaluating at $t = 0$. Observe that if the initial data is irrotational in a neighborhood of the interface Γ_1 , we have $\overline{\Delta}v_{03} = 0$ there, and we simply get $\|\partial_t q(0)\|_{2.5} \leq P(\|v_0\|_3)$ and $\|\partial_t v(0)\|_{1.5} \leq P(\|v_0\|_3)$.

It is now clear how to proceed to bound higher time derivatives at the time zero. We further differentiate the equations in time, evaluate at $t = 0$, and use the estimates derived for lower time derivatives at time zero. We conclude that

$$\begin{aligned}\|\partial_t v(0)\|_{2.5} + \|\partial_t^2 v(0)\|_{1.5} + \|\partial_t^3 v(0)\|_0 \\ + \|q(0)\|_3 + \|\partial_t q(0)\|_{2.5} + \|\partial_t^2 q(0)\|_1 \leq P(\|v_0\|_3, \|v_{03}\|_{4,\Gamma_1}).\end{aligned}\tag{7.5}$$

Combining (6.2), (6.4), (6.6), (6.23), (6.38), (6.40), (6.41), and (7.5), we obtain (6.1).

Remark 7.1. The arguments in this section clarify why our priori bounds also depend on a higher norm of v_{03} on the boundary. We see from the above boundary value problem for $\partial_t q(0)$ that this term is controlled by $\overline{\Delta}v_3 = \overline{\Delta}(v \cdot N)$ on the boundary Γ_1 . Since we want to bound $\partial_t^2 v$ in $H^{1.5}$, we need $\partial_t^2 v$ in $H^1(\Gamma_1)$. By (2.3a) differentiated in time, this requires, considering the tangent component of this equation, that $\partial_t q(0)$ belongs to $H^2(\Gamma_1)$. Thus, one needs $\overline{\Delta}v_3 \in H^2(\Gamma_1)$.

8. ENERGY ESTIMATE ON THE TWO TIMES DIFFERENTIATED SYSTEM

In this section we prove the next lemma, which provides an upper bound on the $\dot{H}^2(\Gamma_1)$ norm of $\Pi\partial_t v$. In the final section, we show that this leads to a bound on $\|\partial_t v\|_{H^2(\Gamma_1)}$.

Lemma 8.1. *We have*

$$\|\overline{\partial}\partial_t^2 v\|_0^2 + \|\overline{\partial}^2(\Pi\partial_t v)\|_{0,\Gamma_1}^2 \leq \tilde{\epsilon}\|\partial_t v\|_{2.5}^2 + \mathcal{P}_0 + P(\|v\|_3, \|q\|_{1.5,\Gamma_1}) + \int_0^t \mathcal{P}.\tag{8.1}$$

Proof of Lemma 8.1. We apply $\partial_m\partial_t^2$ to (2.3a), contract with $\partial_m\partial_t^2 v_\alpha$, integrate in time and space and find

$$\begin{aligned}\frac{1}{2}\int_\Omega \partial_m\partial_t^2 v_\alpha\partial_m\partial_t^2 v_\alpha &= \int_0^t \int_\Omega \partial_\mu\partial_m\partial_t^2 v_\alpha a_{\mu\alpha}\partial_t^2\partial_m q - \int_0^t \int_{\Gamma_1} \partial_m\partial_t^2 v_\alpha a_{3\alpha}\partial_t^2\partial_m q \\ &\quad - \int_0^t \int_\Omega \partial_m\partial_t^2 v_\alpha r_{\alpha m} + \frac{1}{2}\int_\Omega \partial_m\partial_t^2 v_\alpha\partial_m\partial_t^2 v_\alpha\Big|_{t=0} \\ &= J_1 + J_2 + J_3 + J_0,\end{aligned}$$

where in the second equality we integrated by parts in ∂_m and used (2.2) with (6.3); we also denoted

$$r_{\alpha m} = \partial_t^2\partial_m a_{\mu\alpha}\partial_\mu q + 2\partial_t\partial_m a_{\mu\alpha}\partial_t\partial_\mu q + \partial_m a_{\mu\alpha}\partial_t^2\partial_\mu q + \partial_t^2 a_{\mu\alpha}\partial_\mu\partial_m q + 2\partial_t a_{\mu\alpha}\partial_t\partial_\mu\partial_m q.$$

The treatment of the above terms is parallel to the previous proof (the derivatives ∂_t^3 are replaced by $\partial_t^2 \partial_m$), and thus we only sketch the details. Since (2.3b) implies $\partial_m \partial_t^2 (a_{\mu\alpha} \partial_\mu v_\alpha) = 0$, we have

$$\begin{aligned} J_1 &= - \int_0^t \int_\Omega \partial_m a_{\mu\alpha} \partial_\mu \partial_t^2 v_\alpha \partial_t^2 \partial_m q - 2 \int_0^t \int_\Omega \partial_m \partial_t a_{\mu\alpha} \partial_\mu \partial_t v_\alpha \partial_t^2 \partial_m q \\ &\quad - 2 \int_0^t \int_\Omega \partial_t a_{\mu\alpha} \partial_m \partial_\mu \partial_t v_\alpha \partial_t^2 \partial_m q - \int_0^t \int_\Omega \partial_m \partial_t^2 a_{\mu\alpha} \partial_\mu v_\alpha \partial_t^2 \partial_m q \\ &\quad - \int_0^t \int_\Omega \partial_t^2 a_{\mu\alpha} \partial_m \partial_\mu v_\alpha \partial_t^2 \partial_m q. \end{aligned}$$

All the integrals are bounded by $\int_0^t \mathcal{P}$ using Hölder and Sobolev inequalities. Similarly, we have $J_0 \leq \mathcal{P}$. Also, the term J_3 is treated by the same procedure just used for J_1 , yielding $J_3 \leq \int_0^t \mathcal{P}$.

The remaining term $J_2 = - \int_0^t \int_{\Gamma_1} \partial_m \partial_t^2 v_\alpha a_{3\alpha} \partial_t^2 \partial_k q$ may be rewritten as

$$\begin{aligned} J_2 &= - \int_0^t \int_{\Gamma_1} \partial_m \partial_t^2 (a_{3\alpha} q) \partial_m \partial_t^2 v_\alpha + \int_0^t \int_{\Gamma_1} \partial_m (\partial_t^2 a_{3\alpha} q) \partial_m \partial_t^2 v_\alpha \\ &\quad + 2 \int_0^t \int_{\Gamma_1} \partial_m (\partial_t a_{3\alpha} \partial_t q) \partial_m \partial_t^2 v_\alpha + \int_0^t \int_{\Gamma_1} \partial_m a_{3\alpha} \partial_t^2 q \partial_m \partial_t^2 v_\alpha \\ &= J_{21} + J_{22} + J_{23} + J_{24}. \end{aligned} \tag{8.2}$$

To estimate J_{21} , we use (2.3d) and obtain

$$J_{21} = \sigma \int_0^t \int_{\Gamma_1} \partial_m \partial_t^2 (\sqrt{g} \Delta_g \eta_\alpha) \partial_m \partial_t^2 v_\alpha.$$

This term is now bounded in the same fashion as I_1 in the previous proof. Namely, with $\partial_m \partial_t^2$ replacing ∂_t^3 , we use (3.10) and split J_{21} similarly to (6.7), yielding

$$J_{21} = J_{211} + J_{212} + J_{213} + J_{214}, \tag{8.3}$$

where J_{211} is the analogue of I_{11} , J_{212} the analogue of I_{12} , J_{213} the analogue of $I_{13} + I_{14}$, and J_{214} the analogue of $I_{15} + I_{16}$. The terms J_{211} and J_{212} contain the products of top derivatives and are estimated in the same way as I_{11} and I_{12} . The remainders J_{213} and J_{214} need to be treated in a slightly different fashion than $I_{13} + I_{14}$ and $I_{15} + I_{16}$ because of the different ways the time and space derivatives in $\partial_t \bar{\partial}$ may be distributed in (3.10) after taking $\bar{\partial}_A = \partial_t$. We have

$$J_{213} = - \int_0^t \int_{\Gamma_1} \bar{\partial} Q(\bar{\partial} \eta) \partial_t \bar{\partial} v \bar{\partial} \partial_t^2 \bar{\partial} v - \int_0^t \int_{\Gamma_1} \partial_t Q(\bar{\partial} \eta) \bar{\partial}^2 v \bar{\partial} \partial_t^2 \bar{\partial} v = J_{2131} + J_{2132}$$

and

$$J_{214} = - \int_0^t \int_{\Gamma_1} \bar{\partial} \partial_t Q(\bar{\partial} \eta) \bar{\partial} v \bar{\partial} \partial_t^2 \bar{\partial} v,$$

where we followed the symbolic notation in Remark 3.4. Using (3.1) and integrating by parts in space

$$\begin{aligned} J_{2131} &= \int_0^t \int_{\Gamma_1} Q(\bar{\partial} \eta) \bar{\partial}^2 \eta \partial_t \bar{\partial}^2 v \partial_t^2 \bar{\partial} v + \int_0^t \int_{\Gamma_1} Q(\bar{\partial} \eta) \bar{\partial}^3 \eta \partial_t \bar{\partial} v \partial_t^2 \bar{\partial} v \\ &\quad + \int_0^t \int_{\Gamma_1} Q(\bar{\partial} \eta) (\bar{\partial}^2 \eta)^2 \partial_t \bar{\partial} v \partial_t^2 \bar{\partial} v \\ &\leq \int_0^t \|Q(\bar{\partial} \eta) \bar{\partial}^2 \eta\|_{L^\infty(\Gamma_1)} \|\partial_t \bar{\partial}^2 v\|_{0, \Gamma_1} \|\partial_t^2 \bar{\partial} v\|_{0, \Gamma_1} \\ &\quad + \int_0^t \|Q(\bar{\partial} \eta)\|_{L^\infty(\Gamma_1)} \|\bar{\partial}^3 \eta\|_{L^4(\Gamma_1)} \|\partial_t \bar{\partial} v\|_{L^4(\Gamma_1)} \|\partial_t^2 \bar{\partial} v\|_{0, \Gamma_1} \\ &\quad + \int_0^t \|\bar{\partial}^2 \eta\|_{L^\infty(\Gamma_1)}^2 \|\partial_t \bar{\partial} v\|_{0, \Gamma_1} \|\partial_t^2 \bar{\partial} v\|_{0, \Gamma_1} \leq \int_0^t \mathcal{P} \end{aligned}$$

where we used (3.3), (3.4), and Proposition 5.1. Next, integrating by parts in time

$$\begin{aligned}
J_{2132} &= - \int_0^t \int_{\Gamma_1} Q(\bar{\partial}\eta) \bar{\partial}v \bar{\partial}^2 v \bar{\partial} \partial_t^2 \bar{\partial}v \\
&= - \int_{\Gamma_1} Q(\bar{\partial}\eta) \bar{\partial}v \bar{\partial}^2 v \partial_t \bar{\partial}^2 v + J_{2132,0} + \int_0^t \int_{\Gamma_1} \partial_t(Q(\bar{\partial}\eta) \bar{\partial}v \bar{\partial}^2 v) \partial_t \bar{\partial}^2 v \\
&= - \int_{\Gamma_1} Q(\bar{\partial}\eta) \bar{\partial}v \bar{\partial}^2 v \partial_t \bar{\partial}^2 v + J_{2132,0} + \int_0^t \int_{\Gamma_1} Q(\bar{\partial}\eta) (\bar{\partial}v)^2 \bar{\partial}^2 v \partial_t \bar{\partial}^2 v \\
&\quad + \int_0^t \int_{\Gamma_1} Q(\bar{\partial}\eta) \partial_t \bar{\partial}v \bar{\partial}^2 v \partial_t \bar{\partial}^2 v + \int_0^t \int_{\Gamma_1} Q(\bar{\partial}\eta) \bar{\partial}v \partial_t \bar{\partial}^2 v \partial_t \bar{\partial}^2 v,
\end{aligned}$$

where

$$J_{2132,0} = \int_{\Gamma_1} Q(\bar{\partial}\eta) \bar{\partial}v \bar{\partial}^2 v \partial_t \bar{\partial}^2 v \Big|_{t=0} \leq \mathcal{P}_0.$$

A direct estimate of each term in J_{2132} now produces

$$J_{213} \leq \tilde{\epsilon} \|\partial_t v\|_{2.5}^2 + P(\|v\|_{2.5+\delta}) + \mathcal{P}_0 + \int_0^t \mathcal{P}. \quad (8.4)$$

Moving to J_{214} and recalling (3.1),

$$\begin{aligned}
J_{214} &= - \int_0^t \int_{\Gamma_1} \bar{\partial} \partial_t Q(\bar{\partial}\eta) \bar{\partial}v \bar{\partial} \partial_t^2 \bar{\partial}v \\
&= - \int_0^t \int_{\Gamma_1} Q(\bar{\partial}\eta) \bar{\partial}^2 \eta (\bar{\partial}v)^2 \bar{\partial} \partial_t^2 \bar{\partial}v - \int_0^t \int_{\Gamma_1} Q(\bar{\partial}\eta) \bar{\partial}^2 v \bar{\partial}v \bar{\partial} \partial_t^2 \bar{\partial}v = J_{2141} + J_{2142}.
\end{aligned}$$

Integrating by parts in space, we find

$$\begin{aligned}
J_{2141} &= \int_0^t \int_{\Gamma_1} Q(\bar{\partial}\eta) (\bar{\partial}^2 \eta)^2 (\bar{\partial}v)^2 \partial_t^2 \bar{\partial}v + \int_0^t \int_{\Gamma_1} Q(\bar{\partial}\eta) \bar{\partial}^3 \eta (\bar{\partial}v)^2 \partial_t^2 \bar{\partial}v \\
&\quad + \int_0^t \int_{\Gamma_1} Q(\bar{\partial}\eta) \bar{\partial}^2 \eta \bar{\partial}v \bar{\partial}^2 v \partial_t^2 \bar{\partial}v.
\end{aligned}$$

With the help of (3.3), (3.4), and Proposition 5.1, we find

$$\begin{aligned}
J_{2141} &\leq C \int_0^t \|Q(\bar{\partial}\eta)\|_{L^\infty(\Gamma_1)} \|(\bar{\partial}^2 \eta)^2\|_{0,\Gamma_1} \|(\bar{\partial}v)^2\|_{L^\infty(\Gamma_1)} \|\partial_t^2 \bar{\partial}v\|_{0,\Gamma_1} \\
&\quad + C \int_0^t \|Q(\bar{\partial}\eta)\|_{L^\infty(\Gamma_1)} \|\bar{\partial}^3 \eta\|_{0,\Gamma_1} \|(\bar{\partial}v)^2\|_{L^\infty(\Gamma_1)} \|\partial_t^2 \bar{\partial}v\|_{0,\Gamma_1} \\
&\quad + C \int_0^t \|Q(\bar{\partial}\eta)\|_{L^\infty(\Gamma_1)} \|\bar{\partial}^2 \eta\|_{L^4(\Gamma_1)} \|\bar{\partial}v\|_{L^\infty(\Gamma_1)} \|\bar{\partial}^2 v\|_{L^4(\Gamma_1)} \|\partial_t^2 \bar{\partial}v\|_{0,\Gamma_1} \leq \int_0^t \mathcal{P},
\end{aligned}$$

where in the first integral on the right-hand side we used

$$\|(\bar{\partial}^2 \eta)^2\|_{0,\Gamma_1} \leq C \|(\bar{\partial}^2 \eta)^2\|_{1.5,\Gamma_1} \leq C \|\eta\|_{3.5,\Gamma_1}^2.$$

Next, integrating by parts in time and estimating as above, we find

$$\begin{aligned}
J_{2142} &= - \int_{\Gamma_1} Q(\bar{\partial}\eta) \bar{\partial}^2 v \bar{\partial} v \partial_t \bar{\partial}^2 v + \int_{\Gamma_1} Q(\bar{\partial}\eta) \bar{\partial}^2 v \bar{\partial} v \partial_t \bar{\partial}^2 v \Big|_{t=0} \\
&\quad + \int_0^t \int_{\Gamma_1} Q(\bar{\partial}\eta) \bar{\partial} v \bar{\partial}^2 v \bar{\partial} v \partial_t \bar{\partial}^2 v + \int_0^t \int_{\Gamma_1} Q(\bar{\partial}\eta) \partial_t \bar{\partial}^2 v \bar{\partial} v \partial_t \bar{\partial}^2 v \\
&\quad + \int_0^t \int_{\Gamma_1} Q(\bar{\partial}\eta) \bar{\partial}^2 v \partial_t \bar{\partial} v \partial_t \bar{\partial}^2 v \\
&\leq \tilde{\epsilon} \|\partial_t v\|_{2.5}^2 + P(\|v\|_{2.5+\delta}) + \mathcal{P}_0 + \int_0^t \mathcal{P}.
\end{aligned}$$

Therefore,

$$J_{214} \leq \tilde{\epsilon} \|\partial_t v\|_{2.5}^2 + P(\|v\|_{2.5+\delta}) + \mathcal{P}_0 + \int_0^t \mathcal{P}. \quad (8.5)$$

Combining (8.3), (8.4), and (8.5), we find

$$J_{21} \leq -\frac{1}{C_{21}} \|\bar{\partial}^2(\Pi \partial_t v)\|_{0,\Gamma_1}^2 + \tilde{\epsilon} \|\partial_t v\|_{2.5}^2 + \mathcal{P}_0 + P(\|v\|_{2.5+\delta}, \|q\|_{1.5,\Gamma_1}) + \int_0^t \mathcal{P}. \quad (8.6)$$

We now sketch the estimates of the remaining boundary terms J_{22} , J_{23} , and J_{24} in (8.2). The key is to use (6.31) and Proposition 5.1. Recalling that $N = (0, 0, 1)$, we have

$$\begin{aligned}
J_{22} &= \int_0^t \int_{\Gamma_1} \partial_m \partial_t^2 a_{3\alpha} q \partial_m \partial_t^2 v_\alpha + \int_0^t \int_{\Gamma_1} \partial_t^2 a_{3\alpha} \partial_m q \partial_m \partial_t^2 v_\alpha \\
&\leq C \sum_{\alpha=1}^3 \int_0^t \|q\|_{L^\infty(\Gamma_1)} \|\bar{\partial} \partial_t^2 a_{3\alpha}\|_{0,\Gamma_1} \|\bar{\partial} \partial_t^2 v\|_{0,\Gamma_1} + C \sum_{\alpha=1}^3 \int_0^t \|\bar{\partial} q\|_{L^\infty(\Gamma_1)} \|\partial_t^2 a_{3\alpha}\|_{0,\Gamma_1} \|\bar{\partial} \partial_t^2 v\|_{0,\Gamma_1}.
\end{aligned}$$

From (6.31), we have, written symbolically,

$$\partial_t^2 a_{3\alpha} = (\bar{\partial} v)^2 + \bar{\partial} \eta \bar{\partial} \partial_t v \text{ on } \Gamma_1.$$

Differentiating this identity, we get easily $J_{22} \leq \int_0^t \mathcal{P}$, where we used Proposition 5.1. The terms J_{23} and J_{24} are handled similarly leading to

$$J_{22} + J_{23} + J_{24} \leq \int_0^t \mathcal{P}. \quad (8.7)$$

Using (8.2), (8.6), and (8.7), we find

$$J_2 \leq -\frac{1}{C_{21}} \|\bar{\partial}^2(\Pi \partial_t v)\|_{0,\Gamma_1}^2 + \tilde{\epsilon} \|\partial_t v\|_{2.5}^2 + P(\|v\|_{2.5+\delta}, \|q\|_{1.5,\Gamma_1}) + \int_0^t \mathcal{P}. \quad (8.8)$$

Finally, combining all the inequalities concludes the proof. \square

9. BOUNDARY ESTIMATE ON v_3

The purpose of this section is to establish control of v_3 on the free boundary Γ_1 .

Lemma 9.1. *The restriction of the third component of the velocity to Γ_1 satisfies*

$$\|v_3\|_{2.5,\Gamma_1} \leq C \|\partial_t q\|_1 + P(\|v\|_{2.5+\delta}, \|q\|_{1.5}) + \int_0^t \mathcal{P}. \quad (9.1)$$

Remark 9.2. The estimates (6.1) and (8.1) provide bounds for the normal components of $\partial_t^2 v$ and $\partial_t v$. These enter in the div-curl estimates below, thus providing bounds for $\partial_t^2 v$ and $\partial_t v$ in the interior. In order to bound $\|v\|_3$ with the div-curl estimates, we need to estimate $\|v\|_{2.5,\Gamma_1}$ first. For this we use (2.3d) and Proposition 5.1 rather than an energy estimate as in Sections 6 and 8. This is because arguing as in those sections produces $\bar{\partial}^3 v$ on Γ_1 , for which we do not have a good control.

Proof of Lemma 9.1. Differentiating (2.3d) in time and setting $\alpha = 3$ yields

$$\sqrt{g}g^{ij}\partial_{ij}^2v_3 - \sqrt{g}g^{ij}\Gamma_{ij}^k\partial_kv_3 = -\partial_t(\sqrt{g}g^{ij})\partial_{ij}^2\eta_3 - \partial_t(\sqrt{g}g^{ij}\Gamma_{ij}^k)\partial_k\eta_3 - \frac{1}{\sigma}\partial_t a_{33}q - \frac{1}{\sigma}a_{3\sigma}\partial_t q \quad \text{on } \Gamma_1,$$

where we also used (3.15). In light of Proposition 5.1, we have

$$\|g_{ij}\|_{2.5,\Gamma_1}, \|\Gamma_{ij}^k\|_{1.5,\Gamma_1} \leq C. \quad (9.2)$$

Thus, by the elliptic estimates for operators with coefficients bounded in Sobolev norms (see [25, 36]) we get

$$\begin{aligned} \|v_3\|_{2.5,\Gamma_1} &\leq C\|\partial_t(\sqrt{g}g^{ij})\partial_{ij}^2\eta_3\|_{0.5,\Gamma_1} + C\|\partial_t(\sqrt{g}g^{ij}\Gamma_{ij}^k)\partial_k\eta_3\|_{0.5,\Gamma_1} \\ &\quad + C\|\partial_t a_{33}q\|_{0.5,\Gamma_1} + C\|a_{3\sigma}\partial_t q\|_{0.5,\Gamma_1}. \end{aligned}$$

For the first term on the right side,

$$\begin{aligned} \|\partial_t(\sqrt{g}g^{ij})\partial_{ij}^2\eta_3\|_{0.5,\Gamma_1} &\leq C\|\partial_t(\sqrt{g}g^{-1})\|_{1.5,\Gamma_1}\|\bar{\partial}^2\eta_3\|_{0.5,\Gamma_1} \leq C\|Q(\bar{\partial}\eta)\partial_t\bar{\partial}\eta\|_{1.5,\Gamma_1}\|\bar{\partial}^2\eta_3\|_{0.5,\Gamma_1} \\ &\leq C\|v\|_3\|\bar{\partial}^2\eta_3\|_{0.5,\Gamma_1} \leq C\|v\|_3, \end{aligned}$$

where we used (3.3) and Lemma 3.1(i). For the next term, we use (2.4), (3.16), and (3.18) to compute

$$\partial_t(\sqrt{g}g^{ij}\Gamma_{ij}^k) = -\frac{1}{\sqrt{g}}g^{pq}\partial_p\eta_\tau\partial_t\partial_q\eta_\tau g^{ij}g^{kl}\partial_l\eta_\nu\partial_{ij}^2\eta_\nu + \sqrt{g}\partial_t(g^{ij}g^{kl}\partial_l\eta_\nu)\partial_{ij}^2\eta_\nu + \sqrt{g}g^{ij}g^{kl}\partial_l\eta_\nu\partial_{ij}^2\partial_t\eta_\nu.$$

Then

$$\begin{aligned} \left\| \frac{1}{\sqrt{g}}g^{pq}\partial_p\eta_\tau\partial_t\partial_q\eta_\tau g^{ij}g^{kl}\partial_l\eta_\nu\partial_{ij}^2\eta_\nu \right\|_{0.5,\Gamma_1} &\leq \left\| \frac{1}{\sqrt{g}}g^{pq}\partial_p\eta_\tau\partial_t\partial_q\eta_\tau g^{ij}g^{kl}\partial_l\eta_\nu \right\|_{1.5,\Gamma_1} \|\partial_{ij}^2\eta_\nu\|_{0.5,\Gamma_1} \\ &\leq P(\|\bar{\partial}\eta\|_{1.5,\Gamma_1})\|\bar{\partial}v\|_{1.5,\Gamma_1}\|\bar{\partial}^2\eta\|_{0.5,\Gamma_1} \leq C\|v\|_3. \end{aligned}$$

Also,

$$\|\sqrt{g}\partial_t(g^{ij}g^{kl}\partial_l\eta_\nu)\partial_{ij}^2\eta_\nu\|_{0.5,\Gamma_1} \leq P(\|\bar{\partial}\eta\|_{1.5,\Gamma_1})\|\bar{\partial}^2\eta\|_{0.5,\Gamma_1} \leq C\|v\|_3$$

and

$$\|\sqrt{g}g^{ij}g^{kl}\partial_l\eta_\nu\partial_{ij}^2\partial_t\eta_\nu\|_{0.5,\Gamma_1} \leq P(\|\bar{\partial}\eta\|_{1.5,\Gamma_1})\|\bar{\partial}^2v\|_{0.5,\Gamma_1} \leq C\|v\|_3,$$

where we used (3.19) to compute $\partial_t(g^{ij}g^{kl}\partial_l\eta_\nu)$, and (3.3) and (3.4) have also been employed. Therefore,

$$\|\partial_t(\sqrt{g}g^{ij}\Gamma_{ij}^k)\partial_k\eta_3\|_{0.5,\Gamma_1} \leq C\|\partial_t(\sqrt{g}g^{ij}\Gamma_{ij}^k)\|_{0.5,\Gamma_1}\|\partial_k\eta_3\|_{1.5,\Gamma_1} \leq C\sum_{k=1}^2\|v\|_3\|\partial_k\eta_3\|_{1.5,\Gamma_1} \leq C\|v\|_3.$$

The terms containing $\partial_t a$ and $\partial_t q$ are easily estimated, and (9.1) is proven. \square

10. DIV-CURL ESTIMATES

In this section we derive estimates for $\|v\|_3$, $\|\partial_t v\|_{2.5}$, and $\|\partial_t^2 v\|_{1.5}$, which we summarize in the next statement.

Lemma 10.1. *The velocity v satisfies*

$$\|v\|_3 \leq C(\|v_0\|_0 + \|\operatorname{curl} v_0\|_2 + \|v_3\|_{2.5,\Gamma_1}) + \mathcal{P} \int_0^t \mathcal{P}, \quad (10.1)$$

for its time derivative we have

$$\|\partial_t v\|_{2.5} \leq \mathcal{P}_0 + \|\partial_t a\|_2\|v\|_{2.5} + \|\partial_t v_3\|_{2,\Gamma_1} + \mathcal{P} \int_0^t \mathcal{P}, \quad (10.2)$$

while its second time derivative satisfies

$$\|\partial_t^2 v\|_{1.5} \leq \mathcal{P}_0 + \|\partial_t a\|_2\|\partial_t v\|_{1.5} + \|\partial_t^2 a\|_{0.5}\|v\|_{2.5+\delta} + \|\partial_t^2 v_3\|_{1,\Gamma_1} + \|\partial_t v\|_{1.5}\|v\|_{2.5+\delta} + \mathcal{P} \int_0^t \mathcal{P}. \quad (10.3)$$

The arguments below are similar to those in [48, 55].

Proof of Lemma 10.1. First, we recall the inequality

$$\|X\|_s \leq C(\|\operatorname{curl} X\|_{s-1} + \|\operatorname{div} X\|_{s-1} + \|X \cdot N\|_{s-0.5, \partial} + \|X\|_0), \quad (10.4)$$

for $s \geq 1$ and any vector field X on Ω for which the right-hand side is well-defined (cf. [21, 25]), and the Cauchy invariance (see e.g. [14, 55])

$$\epsilon_{\alpha\beta\gamma} \partial_\beta v_\mu \partial_\gamma \eta_\mu = (\operatorname{curl} v)_\alpha, \quad (10.5)$$

where $\epsilon_{\alpha\beta\gamma}$ is the Levi-Civita symbol. First, from (2.3b) we have

$$\|\operatorname{div} v\|_2 = \|(\delta_{\alpha\beta} - a_{\alpha\beta}) \partial_\alpha v_\beta\|_2 \leq C \|\delta_{\alpha\beta} - a_{\alpha\beta}\|_2 \|\partial_\alpha v_\beta\|_2 \leq \tilde{\epsilon} \|v\|_3.$$

Next, using (10.5),

$$\begin{aligned} (\operatorname{curl} v)_\alpha &= \epsilon_{\alpha\beta\gamma} \partial_\beta v_\gamma = \epsilon_{\alpha\beta\gamma} \partial_\beta v_\gamma - \epsilon_{\alpha\beta\gamma} \partial_\beta v_\mu \partial_\gamma \eta_\mu + (\operatorname{curl} v)_\alpha \\ &= \epsilon_{\alpha\beta\gamma} \partial_\beta v_\mu (\delta_{\mu\gamma} - \partial_\gamma \eta_\mu) + (\operatorname{curl} v)_\alpha. \end{aligned}$$

But

$$\delta_{\mu\gamma} - \partial_\gamma \eta_\mu = - \int_0^t \partial_\gamma \partial_t \eta_\mu = - \int_0^t \partial_\gamma v_\mu,$$

so that

$$\|\operatorname{curl} v\|_2 \leq \|\operatorname{curl} v_0\|_2 + C \|v\|_3 \int_0^t \|v\|_3 \leq \|\operatorname{curl} v_0\|_2 + \mathcal{P} \int_0^t \mathcal{P}.$$

Since

$$\|v\|_0 \leq \|v_0\|_0 + C \int_0^t \|\partial_t v\|_0 \leq \|v_0\|_0 + \int_0^t \mathcal{P},$$

we get (10.1) by invoking (10.4). The inequalities (10.2) and (10.3) are obtained analogously. \square

11. CLOSING THE ESTIMATES

In this section we finally collect all our estimates together to obtain Theorem 2.3.

11.1. Comparing ΠX and X_3 . From (10.2) and (10.3), we see that we need to connect $\|X_3\|_{s-0.5, \partial}$ with the norm of ΠX which entered in the energy estimates; cf. the second term on the left side of (6.1). The two necessary inequalities are stated next.

Lemma 11.1. *We have*

$$\|\bar{\partial} X_3\|_{0, \Gamma_1} \leq \tilde{\epsilon} \left(\sum_{\alpha=1}^3 \|\bar{\partial} X_\alpha\|_{0, \Gamma_1} + \sum_{\alpha=1}^3 \|X_\alpha\|_1 \right) + C \|\bar{\partial}(\Pi X)\|_{0, \Gamma_1} \quad (11.1)$$

and

$$\|\bar{\partial}^2 X_3\|_{0, \Gamma_1} \leq \tilde{\epsilon} \|X\|_{2.5} + \tilde{\epsilon} \|\bar{\partial}^2 X\|_{0, \Gamma_1} + C \|\bar{\partial}^2(\Pi X)\|_{0, \Gamma_1} + P(\|q\|_{1.5, \Gamma_1}, \|\partial_t X(0)\|_0) + \int_0^t P(\|\partial_t X\|_0). \quad (11.2)$$

Proof of Lemma 11.1. Recalling (3.6)

$$(\Pi X)_3 = \Pi_{3\lambda} X_\lambda = (\delta_{3\lambda} - g^{kl} \partial_k \eta_3 \partial_l \eta_\lambda) X_\lambda = X_3 - g^{kl} \partial_k \eta_3 \partial_l \eta_\lambda X_\lambda, \quad (11.3)$$

so that

$$\begin{aligned} \|\bar{\partial}((\Pi X)_3 - X_3)\|_{0, \Gamma_1} &= \|\bar{\partial}(g^{kl} \partial_k \eta_3 \partial_l \eta_\lambda X_\lambda)\|_{0, \Gamma_1} \\ &\leq C \|g^{kl} \partial_k \eta_3 \partial_l \eta_\lambda \bar{\partial} X_\lambda\|_{0, \Gamma_1} + C \|\bar{\partial}(g^{kl} \partial_k \eta_3 \partial_l \eta_\lambda) X_\lambda\|_{0, \Gamma_1}. \end{aligned}$$

Since

$$g^{kl} \partial_k \eta_3 \partial_l \eta_\lambda \Big|_{t=0} = 0 \quad (11.4)$$

due to $\eta_3(0) = x_3$, we have

$$\|g^{kl}\partial_k\eta_3\partial_l\eta_\lambda\|_{1.5,\Gamma_1} = \left\| \int_0^t \partial_t(g^{kl}\partial_k\eta_3\partial_l\eta_\lambda) \right\|_{1.5,\Gamma_1} \leq C \int_0^t \|\partial_t\bar{\partial}\eta\|_{1.5,\Gamma_1} \leq C \int_0^t \|v\|_3 \leq CMt.$$

Hence,

$$\|g^{kl}\partial_k\eta_3\partial_l\eta_\lambda\bar{\partial}X_\lambda\|_{0,\Gamma_1} \leq C\|g^{kl}\partial_k\eta_3\partial_l\eta_\lambda\|_{1.5,\Gamma_1}\|\bar{\partial}X_\lambda\|_{0,\Gamma_1} \leq CMt \sum_{\lambda=1}^3 \|\bar{\partial}X_\lambda\|_{0,\Gamma_1}.$$

Next,

$$\begin{aligned} \|\bar{\partial}(g^{kl}\partial_k\eta_3\partial_l\eta_\lambda)X_\lambda\|_{0,\Gamma_1} &\leq C\|\bar{\partial}(g^{kl}\partial_k\eta_3\partial_l\eta_\lambda)\|_{L^4(\Gamma_1)} \sum_{\lambda=1}^3 \|X_\lambda\|_{L^4(\Gamma_1)} \\ &\leq C\|\bar{\partial}(g^{kl}\partial_k\eta_3\partial_l\eta_\lambda)\|_{0.5,\Gamma_1} \sum_{\lambda=1}^3 \|X_\lambda\|_{0.5,\Gamma_1} \leq CMt \sum_{\lambda=1}^3 \|X_\lambda\|_1. \end{aligned}$$

The estimate (11.1) now follows upon choosing t sufficiently small.

We now turn to (11.5). From (11.3), we have

$$\begin{aligned} \bar{\partial}^2(g^{kl}\partial_k\eta_3\partial_l\eta_\lambda X_\lambda) &= \bar{\partial}^2 g^{kl}\partial_k\eta_3\partial_l\eta_\lambda X_\lambda + 2\bar{\partial}g^{kl}\bar{\partial}\partial_k\eta_3\partial_l\eta_\lambda X_\lambda + 2\bar{\partial}g^{kl}\partial_k\eta_3\bar{\partial}\partial_l\eta_\lambda X_\lambda \\ &\quad + 2\bar{\partial}g^{kl}\partial_k\eta_3\partial_l\eta_\lambda\bar{\partial}X_\lambda + g^{kl}\bar{\partial}^2\partial_k\eta_3\partial_l\eta_\lambda X_\lambda + 2g^{kl}\bar{\partial}\partial_k\eta_3\bar{\partial}\partial_l\eta_\lambda X_\lambda \\ &\quad + g^{kl}\bar{\partial}\partial_k\eta_3\partial_l\eta_\lambda\bar{\partial}X_\lambda + g^{kl}\partial_k\eta_3\bar{\partial}^2\partial_l\eta_\lambda X_\lambda + 2g^{kl}\partial_k\eta_3\bar{\partial}\partial_l\eta_\lambda\bar{\partial}X_\lambda \\ &\quad + g^{kl}\bar{\partial}\partial_k\eta_3\partial_l\eta_\lambda\bar{\partial}X_\lambda + g^{kl}\partial_k\eta_3\partial_l\eta_\lambda\bar{\partial}^2 X_\lambda \\ &= A_1 + \dots + A_{11}. \end{aligned}$$

We estimate the terms similarly to those leading to (11.1). We make successive use of the following facts. As before, the terms in $\bar{\partial}\eta_3$ are small (in appropriate norms) in light of (11.4) and what immediately follows. But here also the terms $\bar{\partial}^2\eta_\alpha$ are small (in appropriate norms), for $\alpha = 1, 2, 3$, again because $\eta(0)$ is the identity, so that, as before

$$\bar{\partial}^2\eta = \int_0^t \bar{\partial}^2 v.$$

To proceed, we note that from (2.4) and (3.19) we have

$$\bar{\partial}g^{-1} = Q(\bar{\partial}\eta)\bar{\partial}^2\eta$$

and

$$\bar{\partial}^2g^{-1} = Q(\bar{\partial}\eta)(\bar{\partial}^2\eta)^2 + Q(\bar{\partial}\eta)\bar{\partial}^3\eta,$$

where we adopted a symbolic notation akin to Remark 3.4. We have

$$\begin{aligned} \|A_1\|_{0,\Gamma_1} &\leq C\|\bar{\partial}^2g^{-1}\|_{0,\Gamma_1}\|\bar{\partial}\eta_3\|_{1.5,\Gamma_1}\|\bar{\partial}\eta\|_{1.5,\Gamma_1}\|X\|_{1.5,\Gamma_1} \\ &\leq P(\|Q(\bar{\partial}\eta)\|_{1.5,\Gamma_1})(\|(\bar{\partial}^2\eta)^2\|_{0,\Gamma_1} + \|\bar{\partial}^3\eta\|_{0,\Gamma_1})\tilde{\epsilon}\|X\|_2 \\ &\leq \tilde{\epsilon}P(\|\bar{\partial}^2\eta\|_{L^4(\Gamma_1)}^2 + \|\eta\|_{3,\Gamma_1})\|X\|_2 \\ &\leq \tilde{\epsilon}P(\|q\|_{1.5,\Gamma_1})\|X\|_2 \end{aligned}$$

and

$$\|A_2\|_{0,\Gamma_1} \leq C\|\bar{\partial}^2\eta\|_{1.5,\Gamma_1}\|\bar{\partial}^2\eta_3\|_{0,\Gamma_1}\|X\|_{1.5,\Gamma_1} \leq \tilde{\epsilon}P(\|q\|_{1,\Gamma_1})\|X\|_2.$$

The terms A_3 and A_4 are estimated similarly as

$$\|A_3\|_{0,\Gamma_1} + \|A_4\|_{0,\Gamma_1} \leq \tilde{\epsilon}P(\|q\|_{1,\Gamma_1})\|X\|_2.$$

The term A_5 is different and it gives the term without $\tilde{\epsilon}$ in (11.5),

$$\|A_5\|_{0,\Gamma_1} \leq C\|g^{-1}\bar{\partial}^3\eta\bar{\partial}\eta X\|_{0,\Gamma_1} \leq C\|\eta\|_{3.5,\Gamma_1}\|X\|_{1.5,\Gamma_1} \leq P(\|q\|_{1.5,\Gamma_1})\|X\|_2.$$

Next,

$$\|A_6\|_{0,\Gamma_1} \leq C \|\bar{\partial}^2 \eta_3\|_{L^4(\Gamma_1)} \|\bar{\partial}^2 \eta\|_{L^4(\Gamma_1)} \|X\|_{1.5,\Gamma_1} \leq \tilde{\epsilon} \|X\|_2$$

and

$$\|A_7\|_{0,\Gamma_1} \leq C \|\bar{\partial}^2 \eta_3\|_{L^4(\Gamma_1)} \|\bar{\partial} \eta\|_{L^4(\Gamma_1)} \|\bar{\partial} X\|_{L^4(\Gamma_1)} \leq \tilde{\epsilon} \|X\|_2.$$

The terms A_8 , A_9 and A_{10} are handled similarly. Finally,

$$\|A_{11}\|_{0,\Gamma_1} \leq C \|\bar{\partial} \eta_3\|_{1.5,\Gamma_1} \|\bar{\partial} \eta\|_{1.5,\Gamma_1} \|\bar{\partial}^2 X\|_{0,\Gamma_1} \leq \tilde{\epsilon} \|\bar{\partial}^2 X\|_{0,\Gamma_1}.$$

Combining the above gives

$$\|\bar{\partial}^2 X_3\|_{0,\Gamma_1} \leq \tilde{\epsilon} \sum_{\alpha=1}^3 \|\bar{\partial}^2 X_\alpha\|_{0,\Gamma_1} + P(\|q\|_{1.5,\Gamma_1}) \|X\|_2 + C \|\bar{\partial}^2 (\Pi X)\|_{0,\Gamma_1}. \quad (11.5)$$

Using interpolation, Young's, and Jensen's inequalities as in (6.11), we readily obtain (11.2). \square

11.2. Eliminating the lower order terms. Our estimates so far contain some lower order terms on the right-hand sides that need to be eliminated.

In (9.1), we have $\|\partial_t q\|_1$. Arguing as in (6.11) we obtain

$$\|\partial_t q\|_1^2 \leq \tilde{\epsilon} \|\partial_t q\|_2^2 + \mathcal{P}_0 + \int_0^t \mathcal{P}. \quad (11.6)$$

Next, we look at the term $P(\|v\|_{2.5+\delta}, \|q\|_{1.5,\Gamma_1})$. Up to here, δ was any number between 0 and 0.5. Now we determine it. By the Cauchy-Schwarz inequality, we have

$$P(\|v\|_{2.5+\delta}, \|q\|_{1.5,\Gamma_1}) \leq P(\|v\|_{2.5+\delta}) + P(\|q\|_{1.5,\Gamma_1}),$$

where $P(\|v\|_{2.5+\delta})$ is a linear combination of terms of the form $\|v\|_{2.5+\delta}^a$, where $a \geq 1$ is integer. Since

$$\|v\|_{2.5+\delta} \leq \frac{1}{2} + \frac{1}{2} \|v\|_{2.5+\delta}^2,$$

we may assume $a \geq 2$. The interpolation inequality gives

$$\|v\|_{2.5+\delta}^a \leq C \|v\|_3^{(0.5-\delta)a/3} \|v\|_0^{(2.5+\delta)a/3}. \quad (11.7)$$

Let a_m be the largest of the powers a , and choose $\delta \equiv \delta_m$ sufficiently close to 0.5 so that

$$\frac{6}{a_m(0.5 - \delta_m)} > 1.$$

Set $p_a = 6/a(0.5 - \delta_m)$, and note that $p_a \geq p_{a_m} > 1$, so that the conjugate exponent q_a is well-defined and also greater than 1. Therefore, we may apply Young's inequality with epsilon to the right hand-side of (11.7) to find

$$\|v\|_{2.5+\delta}^a \leq \tilde{\epsilon} \|v\|_3^2 + C \|v\|_0^b,$$

where $b = (2.5 + \delta)aq/3$. Since $a \geq 2$ and $q > 1$, we have $b > 1$ and hence we may use Jensen's inequality to obtain

$$\|v\|_0^b \leq P(\|v_0\|, \|v\|_{3.5,\Gamma_1}) + \int_0^t \mathcal{P}.$$

We therefore conclude that

$$P(\|v\|_{2.5+\delta}) \leq \tilde{\epsilon} \|v\|_3^2 + C + \int_0^t \mathcal{P}.$$

For $P(\|q\|_{1.5,\Gamma_1})$, we simply note that $\|q\|_{1.5,\Gamma_1} \leq C \|q\|_{2.5+\tilde{\delta}}$ for any $0 < \tilde{\delta} < 0.5$. Thus we can proceed exactly as in the estimate for $P(\|v\|_{2.5+\delta})$. We finally conclude

$$P(\|v\|_{2.5+\delta}, \|q\|_{1.5,\Gamma_1}) \leq \tilde{\epsilon} (\|v\|_3^2 + \|q\|_3^2) + C + \int_0^t \mathcal{P}. \quad (11.8)$$

12. PROOF OF THE MAIN THEOREM

We are now ready to combine all our estimates. This consists essentially of keeping track of the energy and pressure estimates and verifying that they always give estimates for higher order terms in terms time integrals of the terms to be bounded. We only sketch the details since the Gronwall argument is fairly standard.

Proof of Theorem 2.3. From (10.2) and (10.3), we see that we need $\|\partial_t v_3\|_{2,\Gamma_1}$ and $\|\partial_t^2 v_3\|_{1,\Gamma_1}$. By interpolation, we have

$$\begin{aligned} \|\partial_t v_3\|_{2,\Gamma_1} &\leq C\|\bar{\partial}^2 \partial_t v_3\|_{0,\Gamma_1} + C\|\partial_t v_3\|_{0,\Gamma_1} \\ &\leq C\|\bar{\partial}^2 \partial_t v_3\|_{0,\Gamma_1} + P(\|v_0\|_3, \|v_0\|_{3.5,\Gamma_1}) + \int_0^t \mathcal{P} \end{aligned}$$

and

$$\begin{aligned} \|\partial_t^2 v_3\|_{1,\Gamma_1} &\leq C\|\bar{\partial} \partial_t^2 v_3\|_{0,\Gamma_1} + C\|\partial_t^2 v_3\|_{0,\Gamma_1} \\ &\leq C\|\bar{\partial} \partial_t^2 v_3\|_{0,\Gamma_1} + \tilde{\epsilon}\|\partial_t^2 v\|_{1.5} + P(\|v_0\|_3, \|v_0\|_{3.5,\Gamma_1}) + \int_0^t \mathcal{P}. \end{aligned}$$

Thus, it suffices to estimate $\|\bar{\partial} \partial_t^2 v_3\|_{0,\Gamma_1}$ and $\|\bar{\partial}^2 \partial_t v_3\|_{0,\Gamma_1}$ which, in light of (11.1) and (11.2), are estimated, up to some harmless lower order terms and terms with $\tilde{\epsilon}$, in terms of $\|\bar{\partial}(\Pi \partial_t^2 v)\|_{0,\Gamma_1}$ and $\|\bar{\partial}^2(\Pi \partial_t v)\|_{0,\Gamma_1}$. An estimate for these last two terms is given in (6.1) and (8.1).

Therefore, combining the estimates of Proposition 4.1 with (10.1), (10.2), (10.3), invoking (6.1), (8.1), and (9.1), eliminating the lower order terms with (11.6) and (11.8), and proceeding as in [48], we obtain a Gronwall-type inequality as in [35] (or cf. [25, 55, 48]), which establishes Theorem 2.3. \square

REFERENCES

- [1] T. Alazard. About global existence and asymptotic behavior for two dimensional gravity water waves. In *Séminaire Laurent Schwartz—Équations aux dérivées partielles et applications. Année 2012–2013*, Sémin. Équ. Dériv. Partielles, pages Exp. No. XVIII, 16. École Polytech., Palaiseau, 2014.
- [2] T. Alazard. Stabilization of the Water-Wave Equations with Surface Tension. *Ann. PDE*, 3(2):3:17, 2017.
- [3] T. Alazard and P. Baldi. Gravity capillary standing water waves. *Arch. Ration. Mech. Anal.*, 217(3):741–830, 2015.
- [4] T. Alazard, N. Burq, and C. Zuily. On the water-wave equations with surface tension. *Duke Math. J.*, 158(3):413–499, 2011.
- [5] T. Alazard, N. Burq, and C. Zuily. Strichartz estimates for water waves. *Ann. Sci. Éc. Norm. Supér. (4)*, 44(5):855–903, 2011.
- [6] T. Alazard, N. Burq, and C. Zuily. The water-wave equations: from Zakharov to Euler. In *Studies in phase space analysis with applications to PDEs*, volume 84 of *Progr. Nonlinear Differential Equations Appl.*, pages 1–20. Birkhäuser/Springer, New York, 2013.
- [7] T. Alazard, N. Burq, and C. Zuily. On the Cauchy problem for gravity water waves. *Invent. Math.*, 198(1):71–163, 2014.
- [8] T. Alazard, N. Burq, and C. Zuily. Cauchy theory for the gravity water waves system with non-localized initial data. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 33(2):337–395, 2016.
- [9] T. Alazard and J.-M. Delort. Global solutions and asymptotic behavior for two dimensional gravity water waves. *Ann. Sci. Éc. Norm. Supér. (4)*, 48(5):1149–1238, 2015.
- [10] T. Alazard and J.-M. Delort. Sobolev estimates for two dimensional gravity water waves. *Astérisque*, (374):viii+241, 2015.
- [11] D. M. Ambrose. Well-posedness of vortex sheets with surface tension. *SIAM J. Math. Anal.*, 35(1):211–244 (electronic), 2003.
- [12] D. M. Ambrose and N. Masmoudi. The zero surface tension limit of two-dimensional water waves. *Comm. Pure Appl. Math.*, 58:1287–1315, 2005.
- [13] J. T. Beale, T. Y. Hou, and J. S. Lowengrub. Growth rates for the linearized motion of fluid interfaces away from equilibrium. *Comm. Pure Appl. Math.*, 46(9):1269–1301, 1993.
- [14] N. Besse and U. Frisch. Geometric formulation of the Cauchy invariants for incompressible Euler flow in flat and curved spaces. *J. Fluid Mech.*, 825:412–478, 2017.
- [15] L. Bieri, S. Miao, S. Shahshahani, and S. Wu. On the Motion of a Self-Gravitating Incompressible Fluid with Free Boundary. *Comm. Math. Phys.*, 355(1):161–243, 2017.
- [16] L. Bieri, S. Miao, S. Shahshahani, and S. Wu. On the motion of a self-gravitating incompressible fluid with free boundary. *Comm. Math. Phys.*, 355(1):161–243, 2017.
- [17] A. Castro, D. Córdoba, C. Fefferman, F. Gancedo, and J. Gómez-Serrano. Finite time singularities for water waves with surface tension. *J. Math. Phys.*, 53(11):115622, 26, 2012.

- [18] A. Castro, D. Córdoba, C. Fefferman, F. Gancedo, and J. Gómez-Serrano. Finite time singularities for the free boundary incompressible Euler equations. *Ann. of Math. (2)*, 178(3):1061–1134, 2013.
- [19] A. Castro, D. Córdoba, C. Fefferman, F. Gancedo, and J. Gómez-Serrano. Structural stability for the splash singularities of the water waves problem. *Discrete Contin. Dyn. Syst.*, 34(12):4997–5043, 2014.
- [20] A. Castro, D. Córdoba, C. Fefferman, F. Gancedo, and M. López-Fernández. Rayleigh-Taylor breakdown for the Muskat problem with applications to water waves. *Ann. of Math. (2)*, 175(2):909–948, 2012.
- [21] C.-H. A. Chen, D. Coutand, and S. Shkoller. Solvability and regularity for an elliptic system prescribing the curl, divergence, and partial trace of a vector field on Sobolev-class domains. *J. Math. Fluid Mech.*, 19(3):375–422, 2017.
- [22] C.-H. Cheng and S. Shkoller. On the motion of vortex sheets with surface tension in three-dimensional Euler equations with vorticity. *Comm. Pure Appl. Math.*, 61(12):1715–1752, 2008.
- [23] D. Christodoulou and H. Lindblad. On the motion of the free surface of a liquid. *Comm. Pure Appl. Math.*, 53(12):1536–1602, 2000.
- [24] D. Coutand. Finite-Time Singularity Formation for Incompressible Euler Moving Interfaces in the Plane. *Arch. Ration. Mech. Anal.*, 232(1):337–387, 2019.
- [25] D. Coutand and S. Shkoller. Well-posedness of the free-surface incompressible Euler equations with or without surface tension. *J. Amer. Math. Soc.*, 20(3):829–930, 2007.
- [26] D. Coutand and S. Shkoller. A simple proof of well-posedness for the free-surface incompressible Euler equations. *Discrete Contin. Dyn. Syst. Ser. S*, 3(3):429–449, 2010.
- [27] D. Coutand and S. Shkoller. On the finite-time splash and splat singularities for the 3-D free-surface Euler equations. *Commun. Math. Phys.*, 325:143–183, 2014.
- [28] W. Craig. On the Hamiltonian for water waves. *arXiv:1612.08971 [math.AP]*, page 10 pages, 2016.
- [29] T. de Poyferré. A priori estimates for water waves with emerging bottom. *Arch. Ration. Mech. Anal.*, 232(2):763–812, 2019.
- [30] Y. Deng, A. D. Ionescu, B. Pausader, and F. Pusateri. Global solutions of the gravity-capillary water-wave system in three dimensions. *Acta Math.*, 219(2):213–402, 2017.
- [31] M. M. Disconzi. On a linear problem arising in dynamic boundaries. *Evolution Equations and Control Theory*, 3(4):627–644, 2014.
- [32] M. M. Disconzi and D. G. Ebin. On the limit of large surface tension for a fluid motion with free boundary. *Comm. Partial Differential Equations*, 39(4):740–779, 2014.
- [33] M. M. Disconzi and D. G. Ebin. The free boundary Euler equations with large surface tension. *Journal of Differential Equations*, 261(2):821–889, 2016.
- [34] H. Dong and D. Kim. On the L_p -solvability of higher order parabolic and elliptic systems with BMO coefficients. *Arch. Ration. Mech. Anal.*, 199(3):889–941, 2011.
- [35] S. S. Dragomir. *Some Gronwall type inequalities and applications*. Nova Science Publishers, Inc., Hauppauge, NY, 2003.
- [36] S. Ebenfeld. L^2 -regularity theory of linear strongly elliptic Dirichlet systems of order $2m$ with minimal regularity in the coefficients. *Quart. Appl. Math.*, 60(3):547–576, 2002.
- [37] D. G. Ebin. The equations of motion of a perfect fluid with free boundary are not well posed. *Comm. Partial Differential Equations*, 12(10):1175–1201, 1987.
- [38] D. G. Ebin and J. Marsden. Groups of diffeomorphisms and the motion of an incompressible fluid. *Annals of Math*, 92:102–163, 1970.
- [39] L. C. Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2010.
- [40] C. Fefferman, A. D. Ionescu, and V. Lie. On the absence of splash singularities in the case of two-fluid interfaces. *Duke Math. J.*, 165(3):417–462, 2016.
- [41] P. Germain, N. Masmoudi, and J. Shatah. Global solutions for the gravity water waves equation in dimension 3. *Ann. of Math. (2)*, 175(2):691–754, 2012.
- [42] P. Germain, N. Masmoudi, and J. Shatah. Global existence for capillary water waves. *Comm. Pure Appl. Math.*, 68(4):625–687, 2015.
- [43] Q. Han and J.-X. Hong. *Isometric embedding of Riemannian manifolds in Euclidean spaces*, volume 130 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2006.
- [44] J. K. Hunter, M. Ifrim, and D. Tataru. Two dimensional water waves in holomorphic coordinates. *Comm. Math. Phys.*, 346(2):483–552, 2016.
- [45] M. Ifrim and D. Tataru. Two dimensional water waves in holomorphic coordinates II: Global solutions. *Bull. Soc. Math. France*, 144(2):369–394, 2016.
- [46] M. Ifrim and D. Tataru. The Lifespan of Small Data Solutions in Two Dimensional Capillary Water Waves. *Arch. Ration. Mech. Anal.*, 225(3):1279–1346, 2017.
- [47] M. Ifrim and D. Tataru. Two-dimensional gravity water waves with constant vorticity I: Cubic lifespan. *Anal. PDE*, 12(4):903–967, 2019.
- [48] M. Ignatova and I. Kukavica. On the local existence of the free-surface Euler equation with surface tension. *Asymptot. Anal.*, 100(1-2):63–86, 2016.
- [49] T. Iguchi, N. Tanaka, and A. Tani. On a free boundary problem for an incompressible ideal fluid in two space dimensions. *Adv. Math. Sci. Appl.*, 9(1):415–472, 1999.

- [50] A. D. Ionescu and F. Pusateri. Global solutions for the gravity water waves system in 2d. *Invent. Math.*, 199(3):653–804, 2015.
- [51] A. D. Ionescu and F. Pusateri. Global analysis of a model for capillary water waves in two dimensions. *Comm. Pure Appl. Math.*, 69(11):2015–2071, 2016.
- [52] A. D. Ionescu and F. Pusateri. Global regularity for 2D water waves with surface tension. *Mem. Amer. Math. Soc.*, 256(1227):v+124, 2018.
- [53] I. Kukavica and A. Tuffaha. On the 2D free boundary Euler equation. *Evol. Equ. Control Theory*, 1(2):297–314, 2012.
- [54] I. Kukavica and A. Tuffaha. A regularity result for the incompressible Euler equation with a free interface. *Appl. Math. Optim.*, 69(3):337–358, 2014.
- [55] I. Kukavica, A. Tuffaha, and V. Vicol. On the local existence and uniqueness for the 3D Euler equation with a free interface. *Appl. Math. Optim.*, 76(3):535–563, 2017.
- [56] D. Lannes. Well-posedness of the water-waves equations. *J. Amer. Math. Soc.*, 18(3):605–654 (electronic), 2005.
- [57] D. Lannes. *The water waves problem*, volume 188 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2013. Mathematical analysis and asymptotics.
- [58] H. Lindblad. The motion of the free surface of a liquid. In *Séminaire: Équations aux Dérivées Partielles, 2000–2001*, Sémin. Équ. Dériv. Partielles, pages Exp. No. VI, 10. École Polytech., Palaiseau, 2001.
- [59] H. Lindblad. Well-posedness for the linearized motion of an incompressible liquid with free surface boundary. *Comm. Pure Appl. Math.*, 56(2):153–197, 2003.
- [60] H. Lindblad and K. H. Nordgren. A priori estimates for the motion of a self-gravitating incompressible liquid with free surface boundary. *J. Hyperbolic Differ. Equ.*, 6(2):407–432, 2009.
- [61] V. I. Nalimov. The Cauchy-Poisson problem. *Dinamika Splošn. Sredy*, (Vyp. 18 Dinamika Zidkost. so Svobod. Granicami):104–210, 254, 1974.
- [62] M. Ogawa and A. Tani. Free boundary problem for an incompressible ideal fluid with surface tension. *Math. Models Methods Appl. Sci.*, 12(12):1725–1740, 2002.
- [63] M. Ogawa and A. Tani. Incompressible perfect fluid motion with free boundary of finite depth. *Adv. Math. Sci. Appl.*, 13(1):201–223, 2003.
- [64] J. Prüss and G. Simonett. *Moving interfaces and quasilinear parabolic evolution equations*, volume 105 of *Monographs in Mathematics*. Birkhäuser/Springer, [Cham], 2016.
- [65] F. Pusateri. On the limit as the surface tension and density ratio tend to zero for the two-phase Euler equations. *J. Hyperbolic Differ. Equ.*, 8(2):347–373, 2011.
- [66] B. Schweizer. On the three-dimensional Euler equations with a free boundary subject to surface tension. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 22(6):753–781, 2005.
- [67] J. Shatah and C. Zeng. Geometry and a priori estimates for free boundary problems of the Euler equation. *Comm. Pure Appl. Math.*, 61(5):698–744, 2008.
- [68] J. Shatah and C. Zeng. Local well-posedness for fluid interface problems. *Arch. Ration. Mech. Anal.*, 199(2):653–705, 2011.
- [69] C. Wang, Z. Zhang, W. Zhao, and Y. Zheng. Local well-posedness and break-down criterion of the incompressible euler equations with free boundary. *arXiv:1507.02478*, 2015.
- [70] S. Wu. Well-posedness in Sobolev spaces of the full water wave problem in 2-D. *Invent. Math.*, 130(1):39–72, 1997.
- [71] S. Wu. Well-posedness in Sobolev spaces of the full water wave problem in 3-D. *J. Amer. Math. Soc.*, 12(2):445–495, 1999.
- [72] S. Wu. Almost global wellposedness of the 2-D full water wave problem. *Invent. Math.*, 177(1):45–135, 2009.
- [73] S. Wu. Global wellposedness of the 3-D full water wave problem. *Invent. Math.*, 184(1):125–220, 2011.
- [74] H. Yosihara. Gravity waves on the free surface of an incompressible perfect fluid of finite depth. *Publ. Res. Inst. Math. Sci.*, 18(1):49–96, 1982.
- [75] H. Yosihara. Capillary-gravity waves for an incompressible ideal fluid. *J. Math. Kyoto Univ.*, 23(4):649–694, 1983.