

Conditional Expectation Bounds with Applications in Cryptography

Kevin J. Compton
Computer Science and Engineering Division
University of Michigan - Ann Arbor
Ann Arbor, MI 48109-2212, USA
kjc@umich.edu

November 15, 2021

Abstract

We derive two conditional expectation bounds, which we use to simplify cryptographic security proofs. The first bound relates the expectation of a bounded random variable and the average of its conditional expectations with respect to a set of i.i.d. random objects. It shows, under certain conditions, that the conditional expectation average has a small tail probability when the expectation of the random variable is sufficiently large. It is used to simplify the proof that the existence of weakly one-way functions implies the existence of strongly one-way functions. The second bound relaxes the independence requirement on the random objects to give a result that has applications to expander graph constructions in cryptography. It is used to simplify the proof that there is a security preserving reduction from weakly one-way functions to strongly one-way functions. To satisfy the hypothesis for this bound, we prove a hitting property for directed graphs that are expander-permutation hybrids.

1. Introduction

Let Z be a random variable on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with $0 \leq Z \leq 1$ and U_i , $0 \leq i < t$, be i.i.d. random objects from the measurable space (Ω, \mathcal{F}) to the measurable space (Ω', \mathcal{F}') . Let W be the average of the conditional expectations $W_i = \mathbb{E}(Z|U_i)$. From the law of iterated expectations and linearity of expectation, we know that $\mathbb{E}(W) = \mathbb{E}(Z)$, but we would expect W to be more concentrated around the mean. Since it is not an average of *independent* random variables, however, this concentration is difficult to quantify. We will derive a tail bound for W which says, roughly, that W has a small tail probability when $\mathbb{E}(Z)$ is sufficiently large. The contrapositive, that $\mathbb{E}(Z)$ is small when W has a large tail probability, may be applied to simplify certain cryptographic security proofs, viz., the proof that the existence of weakly one-way functions implies the existence of strongly one-way functions. We will relax the independence requirement on the random objects U_i to obtain a somewhat weaker inequality that has applications to cryptographic expander graph constructions such as the one used to show that there is a security preserving reduction from weakly one-way functions to strongly one-way functions.

We will assume the reader is familiar with terminology and notation from probability theory found in standard texts, e.g., [2, 20, 29]. Before proceeding, we should point out that

there are two definitions of conditional expectation in the literature. In most cases it does not matter which is used, but it does here. Let Y be a random variable on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and X be a measurable function from measurable space (Ω, \mathcal{F}) to measurable space $(\Omega', \mathcal{F}', \mathbb{P}_X)$. We will take the definition that says $\mathbb{E}(Y|X)$ is a random variable V on $(\Omega', \mathcal{F}', \mathbb{P}_X)$ such that

$$\int_S V d\mathbb{P}_X = \int_{\{X \in S\}} Y d\mathbb{P}$$

for all $S \in \mathcal{F}'$. (Here \mathbb{P}_X is the probability measure induced by X .) This was the first rigorous definition of conditional expectation, formulated by Kolmogorov in his seminal 1933 treatise [27] and found in many current texts (e.g., [2, 20]). Kolmogorov used the Radon-Nikodym Theorem to show that such a V exists whenever $\mathbb{E}(Y)$ exists, and is unique in the sense that any two such V 's differ only on a set of measure 0. Some sources (e.g., [11]) do not define $\mathbb{E}(Y|X)$ as V , but instead as $V \circ X$ (or an equivalent expression), which is a random variable on $(\Omega, \mathcal{F}, \mathbb{P})$. Loève adopts the latter definition, but remarks that the original Komolgorov definition is the “usual interpretation” (cf. [29], p. 343). $V \circ X$ is sometimes denoted $\mathbb{E}(Y|\sigma(X))$, where $\sigma(X)$ is the σ -algebra induced by X .

The situation described in the introductory paragraph arises naturally. Suppose, for example, that we want to approximate the probability of some event $S \subseteq \Omega$, where Ω is the solid unit n -cube $[0, 1]^n$. If n is large, it may not be feasible to generate random points in Ω so instead we try to approximate the probability of a random lower-dimensional cross-section of S . We generate the cross-section by randomly choosing a projection $U_i: \mathbb{R}^n \rightarrow \mathbb{R}^k$, where $k < n$, from some fixed collection U_0, U_1, \dots, U_{t-1} of independent projections, then randomly choosing some $x \in [0, 1]^k$ and intersecting S with the affine subspace $\{U_i = x\}$. If we let Z be the indicator function for S , $W_i = \mathbb{E}(Z|U_i)$, and $W = (W_0 + W_1 + \dots + W_{t-1})/t$, then $W(x)$ is the probability (or expected volume) of this cross-section.

Take a specific example. Let $n = t$ and fix p between 0 and 1. Set $S = [0, p^{1/t}]^t$ so $\mathbb{E}(Z) = p$. The t projection functions U_i given by $U_i(y_0, y_1, \dots, y_{t-1}) = y_i$ are i.i.d. random objects and we compute each $W_i = \mathbb{E}(Z|U_i)$ to be

$$W_i(x) = \begin{cases} p^{1-1/t}, & \text{if } x \leq p^{1/t}, \\ 0, & \text{otherwise.} \end{cases}$$

The random variables W_i are identical, so W , their average, is the same function. Now compare: Z takes the value 1 with probability p ; W takes the value $p^{1-1/t}$ with probability $p^{1/t}$. For large t , W is more concentrated, taking a value closer to the expectation p on an event whose probability is $\mathbb{E}(Z)^{1/t}$. In fact, for any ε such that $0 < \varepsilon < p^{1-1/t}$, $\mathbb{P}_U\{W > \varepsilon\} = \mathbb{E}(Z)^{1/t}$.

Our first conditional expectation bound, Theorem 2.1(i), says that something like this holds in general. Since the random objects U_i are identically distributed we may write $U_i \sim U$ for some fixed U . We show that when $0 \leq Z \leq 1$ and $0 < \varepsilon < 1$,

$$\mathbb{E}(Z) \leq \mathbb{P}_U\{W > \varepsilon\}^t + t\varepsilon. \tag{1.1}$$

This gives a lower bound for $\mathbb{P}_U\{W > \varepsilon\}$ when $t\varepsilon$ is small, and, therefore, a tail bound

$$\mathbb{P}_U\{W \leq \varepsilon\} \leq 1 - (\mathbb{E}(Z) - t\varepsilon)^{1/t}.$$

A further result, Theorem 2.2(i), dispenses with the hypothesis that the random objects U_i are identically distributed. The bound is given in terms of product of tail probabilities $\mathbb{P}_{U_i}\{W_i > \varepsilon_i\}$ plus a correction term $t\varepsilon$, where ε is the average of the values ε_i .

Our second conditional expectation bound, Theorem 2.1(ii), weakens the hypothesis by replacing independence of the random objects U_i with a property, introduced here, called β -independence (where $0 \leq \beta \leq 1$). In the case of identically distributed random objects U_i , β -independence holds if for all events $T_i \in \mathcal{F}'$, setting $\mathbb{P}\{U_i \in T_i\} = 1 - \nu_i$, we have

$$\mathbb{P}\left\{\bigwedge_{0 \leq i < t} U_i \in T_i\right\} \leq \prod_{0 \leq i < t} (1 - \beta \nu_i). \quad (1.2)$$

Under this hypothesis we conclude

$$\mathbb{E}(Z) \leq (\alpha + \beta \mathbb{P}_U\{W > \varepsilon\})^t + t\varepsilon, \quad (1.3)$$

where $\alpha = 1 - \beta$. It is not difficult to see that 1-independence is equivalent to independence, so that part (i) of Theorem 2.1 is a special case of part (ii). We should note that Theorem 2.1(ii) does not require full β -independence. It suffices that (1.2) holds when the events T_i are identical, i.e.,

$$\mathbb{P}\left\{\bigwedge_{0 \leq i < t} U_i \in T\right\} \leq (1 - \beta \nu)^t, \quad (1.4)$$

where $\mathbb{P}\{U_i \in T\} = 1 - \nu$. However, we show a related result, Theorem 2.2, which dispenses with the condition that the random objects U_i be identically distributed, but requires full β -independence given by (1.2).

Inequalities (1.1) and (1.3) are not difficult to prove. We regard them as useful probabilistic tools similar to Chernoff bounds. But whereas we use Chernoff bounds to show that an efficient probabilistic algorithm has a high probability of returning the correct result, we use conditional expectation bounds to show that all efficient probabilistic algorithms have a low probability of breaking a cryptographic construction. One of the principal contributions of this paper is to show how conditional expectation bounds can simplify certain security proofs.

The earliest example of such a security proof deals with Yao's construction [42] of a strongly one-way function as a direct power of a weakly one-way function. (See Section 4 for precise definitions.) The idea is simple. From F , a weakly one-way function, define

$$F'(x_0 x_1 \cdots x_{t-1}) = (F(x_0), F(x_1), \dots, F(x_{t-1})),$$

where $x_0, x_1, \dots, x_{t-1} \in \{0, 1\}^n$ and $t = t(n)$ is suitably chosen polynomial. The proof that F' is strongly one-way is not so simple.

The first proof of this result appeared in an online draft of a text by Goldreich [13]. Let us describe it in terms of a reduction between cryptographic primitives (as found in [30, 8, 28]). For one-way functions, a *reduction* is a pair $(\mathcal{R}, \mathcal{R}^*)$ where \mathcal{R} is an efficient transformation taking F to F' and \mathcal{R}^* is an efficient transformation taking each randomized function G' attempting to invert F' to a randomized function G attempting to invert F . There is also a condition, detailed in section 4, relating the probability that G inverts F to the probability that G' inverts F' . In this situation, we say $(\mathcal{R}, \mathcal{R}^*)$ is a reduction from F to F' (although it would be more accurate to say it is a reduction of the invertibility problem for F to the invertibility problem for F').

Goldreich's proof implicitly gives such a reduction. \mathcal{R} is the the direct power construction. The crux of the proof is to specify \mathcal{R}^* so that if G' is probabilistic polynomial time

(p.p.t.) computable then so is G , and whenever G inverts F with probability significantly less than 1, G' inverts F' with negligible probability. If F is weakly one-way, every p.p.t. G inverts F with probability significantly less than 1, and, hence, every p.p.t. G' inverts F' with negligible probability. Therefore, F' is strongly one-way. The proof of this in [13] relies on a number of technical subsidiary results running to several pages. Using the conditional expectation inequality (1.1), we are able to significantly simplify the proof (cf. Theorem 5.1).

Goldreich et al. [15] later pointed out a drawback to the direct power construction: the associated reduction is not security preserving (see section 6 for the precise definition). The reason is that the forward transformation \mathcal{R} takes F with input size n to F' with input size $nt(n)$; \mathcal{R}^* , then, takes G' with input size $nt(n)$ to G with input size n . It follows that if $S(n)$ is the security of F against G and $S'(n)$ is the security of F' against G' , then $S(n)$ is of the same order as $S'(nt(n))$, which is much larger than $S'(n)$ when S' has superpolynomial growth. They remedied this deficiency by replacing the direct power with a more elaborate construction employing expander graphs to control input size blowup. They give a security preserving reduction in the restricted case where F is a weakly one-way permutation and F' is a strongly one-way permutation, but now rather than $S'(nt(n))$, the security of F' against G' is $S'(n + \omega(\log n))$, which is of the same order as $S'(n)$ (in a sense made precise in section 6). In this case, (1.1) does not suffice, so we use (1.3). Again, the original proof relies on many subsidiary technical results which we bypass by means of the conditional expectation bound (1.3). To apply it, though, we must satisfy the β -independence condition.

The hitting property for expander graphs is closely related to β -independence. The version of hitting property in [41] is equivalent to the statement that if U_i is the i -th vertex along a random walk in an expander graph $\mathcal{G} = (V, E)$ whose spectral gap is β , $T \subseteq V$, and $|T|/|V| = 1 - \nu$, then (1.4) holds. As we noted, (1.4) is weaker than the full β -independence definition (1.2) where we have $T_i \subseteq V$ for each $i < t$, but it is possible to prove this stronger version of the hitting property for expander graphs; this result does not seem to be stated anywhere in the literature (but does follow from Theorem 3.11 of [22] using the Cauchy-Schwarz inequality). In section 3 we will show an even stronger result. We will show that (1.3) holds for random walks on *directed graphs* that are hybrids formed from expander graphs and vertex permutations. This significantly simplifies the proof for the reduction from weakly to strongly one-way permutations in [15, 13].

The outline of the paper is as follows. In Section 2 we prove the conditional expectation bounds. In Section 3 we show that the β -independence property (1.2) holds in hybrid expander-permutation directed graphs. In Section 4 we define and discuss the properties of weakly and strongly one-way functions. In Section 5 we use the conditional expectation inequality (1.1) to simplify Goldreich's proof that existence of weakly one-way functions implies the existence of strongly one-way functions. In Section 6 we use the conditional expectation inequality (1.3) to show that the hybrid expander-permutation directed graph construction yields a strongly one-way permutation when the expander graph is from a certain fully explicit family expander graphs. In section 6 we show that there is a security preserving reduction from weakly to strongly one-way permutations. Our result is stronger than the one in Goldreich et al. [15] because they require that their expander graph families have fully explicit polynomial-time computable edge-colorings. We will show that the edge-coloring assumption is not necessary. In section 7, we conclude with some brief remarks.

2. Conditional Expectation Bounds

Before stating our conditional expectation bounds, we recall some basic terminology and notation from probability theory. A *random object* on $(\Omega, \mathcal{F}, \mathbb{P})$ is a measurable function X from measurable algebra (Ω, \mathcal{F}) to measurable algebra (Ω', \mathcal{F}') . When $\Omega' = \mathbb{R}$ and \mathcal{F}' is the σ -algebra of Borel sets, X is a *random variable*. The *indicator function* for an event S is

$$\mathbb{1}_S(x) = \begin{cases} 1, & \text{if } x \in S, \\ 0, & \text{otherwise,} \end{cases}$$

$\mathbb{E}(Z)$ is the expectation of a random variable Z and $\mathbb{E}_S(Z)$ denotes $\mathbb{E}(\mathbb{1}_S \cdot Z)$. A random object $U: (\Omega, \mathcal{F}) \rightarrow (\Omega', \mathcal{F}')$ induces the probability measure \mathbb{P}_U (also called the *distribution* of U) on (Ω', \mathcal{F}') given by $\mathbb{P}_U(T) = \mathbb{P}\{U \in T\}$.

As we noted earlier, we introduce here a generalization of the notion of independence for random objects. Let $0 \leq \beta \leq 1$ and $\alpha = 1 - \beta$. The random objects $U_i: (\Omega, \mathcal{F}) \rightarrow (\Omega'_i, \mathcal{F}'_i)$, $i < t$, are β -independent if for all events $T_i \in \mathcal{F}'_i$ with $\mathbb{P}_{U_i}(T_i) = \mu_i = 1 - \nu_i$,

$$\mathbb{P}\left\{ \bigwedge_{0 \leq i < t} U_i \in T_i \right\} \leq \prod_{0 \leq i < t} (1 - \beta \nu_i)$$

or, equivalently,

$$\mathbb{P}\left\{ \bigwedge_{0 \leq i < t} U_i \in T_i \right\} \leq \prod_{0 \leq i < t} (\alpha + \beta \mu_i).$$

If, in addition, the random objects U_i are identically distributed, they are said to be β -i.i.d.

We now state the conditional expectation bounds.

Theorem 2.1. *Let Z be a random variable such that $0 \leq Z \leq 1$, and*

$$U_i: (\Omega, \mathcal{F}) \rightarrow (\Omega'_i, \mathcal{F}'_i), i < t,$$

be identically distributed with $U_i \sim U$ for some fixed U . Set $W_i = \mathbb{E}(Z|U_i)$, $W = (W_0 + W_1 + \dots + W_{t-1})/t$ and take any $0 < \varepsilon < 1$.

(i) *If the random objects U_i are independent,*

$$\mathbb{E}(Z) \leq \mathbb{P}_U\{W > \varepsilon\}^t + t\varepsilon.$$

(ii) *If the random objects U_i are β -independent,*

$$\mathbb{E}(Z) \leq (\alpha + \beta \mathbb{P}_U\{W > \varepsilon\})^t + t\varepsilon,$$

where $\alpha = 1 - \beta$.

Proof. Let $T = \{W > \varepsilon\}$, $S_i = \{U_i \in T\}$, and $S = \bigcap_{i < t} S_i$. By the the law of total expectation,

$$\mathbb{E}(Z) = \mathbb{E}(Z|S) \mathbb{P}(S) + \mathbb{E}(Z|\bar{S}) \mathbb{P}(\bar{S}), \quad (2.1)$$

Bound the right side of this equation as follows. $\mathbb{E}(Z|S) \leq 1$ since $Z \leq 1$. Independence of the random objects U_i in part (i) of the theorem implies

$$\mathbb{P}(S) = \mathbb{P}\left(\bigcap_{i < t} S_i\right) = \prod_{i < t} \mathbb{P}_{U_i}(T) = \mathbb{P}_U\{W > \varepsilon\}^t,$$

and β -independence in part (ii) implies

$$\mathbb{P}(S) \leq \prod_{i < t} (\alpha + \beta \mathbb{P}_{U_i}(T)) = (\alpha + \beta \mathbb{P}_U\{W > \varepsilon\})^t.$$

From $\bar{S} = \bigcup_{i < t} \bar{S}_i$ we have

$$\begin{aligned} \mathbb{E}(Z|\bar{S}) \mathbb{P}(\bar{S}) &= \mathbb{E}_{\bar{S}}(Z) \\ &\leq \sum_{i < t} \mathbb{E}_{\bar{S}_i}(Z), \end{aligned}$$

which, by the law of iterated expectations, is equal to

$$\begin{aligned} \sum_{i < t} \mathbb{E}_{\bar{T}}(\mathbb{E}(Z|U_i)) &= \sum_{i < t} \mathbb{E}_{\{W \leq \varepsilon\}}(W_i) \\ &= \mathbb{E}_{\{W \leq \varepsilon\}}\left(\sum_{i < t} W_i\right) \\ &= \mathbb{E}_{\{W \leq \varepsilon\}}(tW) \\ &\leq t\varepsilon, \end{aligned}$$

since $\bar{S}_i = \{U_i \in \bar{T}\}$ and $\bar{T} = \{W \leq \varepsilon\}$.

Substitution into (2.1) completes the proof. \square

Remark. As we noted in the introduction, we may relax the hypotheses of Theorem 2.1 so that rather than independence in part (i), we require only that for every event T ,

$$\mathbb{P}\left\{\bigwedge_{i < t} U_i \in T\right\} = \prod_{i < t} \mathbb{P}_{U_i}(T),$$

and similarly for β -independence in part (ii).

We do not use the following theorem in this paper, but it may prove useful in other contexts where the random variables U_i are not identically distributed.

Theorem 2.2. *Let Z be a random variable such that $0 \leq Z \leq 1$,*

$$U_i: (\Omega, \mathcal{F}) \rightarrow (\Omega'_i, \mathcal{F}'_i), i < t,$$

be random objects with $W_i = \mathbb{E}(Z|U_i)$, $0 < \varepsilon_i < 1$ for $i < t$, and $\varepsilon = (\sum_{i < t} \varepsilon_i)/t$.

(i) *If the random objects U_i are independent,*

$$\mathbb{E}(Z) \leq \prod_{i < t} \mathbb{P}_{U_i}\{W_i > \varepsilon_i\} + t\varepsilon.$$

(ii) *If the random objects U_i are β -independent,*

$$\mathbb{E}(Z) \leq \prod_{i < t} (\alpha + \beta \mathbb{P}_{U_i}\{W_i > \varepsilon_i\}) + t\varepsilon,$$

where $\alpha = 1 - \beta$.

Proof. Let $T_i = \{W_i > \varepsilon_i\}$, $S_i = \{U_i \in T_i\}$, and $S = \bigcap_{i < t} S_i$. As in the previous theorem, we bound the right side of

$$\mathbb{E}(Z) = \mathbb{E}(Z|S)\mathbb{P}(S) + \mathbb{E}(Z|\bar{S})\mathbb{P}(\bar{S}).$$

As before, $\mathbb{E}(Z|S) \leq 1$. Also, for part (i) of the theorem,

$$\mathbb{P}(S) = \mathbb{P}\left(\bigcap_{i < t} S_i\right) = \prod_{i < t} \mathbb{P}_{U_i}(T_i),$$

and for part (ii),

$$\mathbb{P}(S) \leq \prod_{i < t} (\alpha + \beta \mathbb{P}_{U_i}(T_i)).$$

We have

$$\mathbb{E}(Z|\bar{S})\mathbb{P}(\bar{S}) \leq \sum_{i < t} \mathbb{E}_{\bar{S}_i}(Z),$$

This last summation is equal to

$$\begin{aligned} \sum_{i < t} \mathbb{E}_{\bar{T}_i}(\mathbb{E}(Z|U_i)) &= \sum_{i < t} \mathbb{E}_{\{W_i \leq \varepsilon_i\}}(W_i) \\ &\leq \sum_{i < t} \varepsilon_i, \end{aligned}$$

since $\bar{S}_i = \{U_i \in \bar{T}_i\}$ and $\bar{T}_i = \{W_i \leq \varepsilon_i\}$. The theorem follows by substitution. \square

3. Expander-Permutation Graphs and β -Independence.

Theorem 3.5, the main result of this section, provides a natural construction of β -independent random objects based on hybrid expander-permutation directed graphs.

Let \mathbf{A} be a symmetric, real-valued matrix of dimension N . In particular, \mathbf{A} is Hermitian (cf. Horn and Johnson [23] for basic results concerning Hermitian matrices), so it has N real eigenvalues. List them (with repetitions according to multiplicities) in nonincreasing order:

$$\lambda_0(\mathbf{A}) \geq \lambda_1(\mathbf{A}) \cdots \geq \lambda_{N-1}(\mathbf{A}).$$

We write λ_i rather than $\lambda_i(\mathbf{A})$ when matrix \mathbf{A} is clear from context. Because \mathbf{A} is Hermitian, there is an orthonormal basis $\mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{N-1} \in \mathbb{R}^n$, where \mathbf{u}_i is a left (row) eigenvector associated with eigenvalue $\lambda_i(\mathbf{A})$. Since \mathbf{A} is symmetric, the respective right eigenvectors are transposes of the left eigenvectors.

We will consider only real-valued vectors. We write vectors in lower case boldface and denote the transpose of vector \mathbf{v} by \mathbf{v}^T , so the inner product of row vectors \mathbf{v} and \mathbf{w} is $\langle \mathbf{v}, \mathbf{w} \rangle = \mathbf{v} \cdot \mathbf{w}^T$ and $|\mathbf{v}|^2 = \langle \mathbf{v}, \mathbf{v} \rangle$. The Cauchy-Schwarz inequality states that $|\langle \mathbf{v}, \mathbf{w} \rangle| \leq |\mathbf{v}| |\mathbf{w}|$.

Let $\mathcal{G} = (V, E)$ be a d -regular undirected graph and \mathbf{A} be its *transition matrix*; i.e., $\mathbf{A} = (a_{i,j})$, where

$$a_{i,j} = \begin{cases} 1/d, & \text{if there is an edge from vertex } i \text{ to vertex } j, \\ 0, & \text{otherwise.} \end{cases}$$

\mathbf{A} is symmetric, nonnegative, and doubly stochastic. If \mathcal{G} is connected, then \mathbf{A} is irreducible (cf. Seneta [39] for basic results concerning nonnegative matrices). By the Perron-Frobenius Theorem, the largest eigenvalue of \mathbf{A} is $\lambda_0 = 1$, the common row sum. Also, this is a simple eigenvalue so $\lambda_0 > \lambda_1$ and no other eigenvalue is larger in magnitude than λ_0 ; it follows that $\lambda_{N-1} \geq -1$. If \mathcal{G} is not bipartite, then $\lambda_{N-1} > -1$. Under these conditions, define $\alpha = \alpha(\mathbf{A})$ to be the second largest eigenvalue magnitude of \mathbf{A} , and β to be the *spectral gap* of \mathbf{A} , i.e., the difference between the largest and second largest eigenvalue magnitudes. That is, $\alpha = \max(|\lambda_1(\mathbf{A})|, |\lambda_{N-1}(\mathbf{A})|)$ and $\beta = 1 - \alpha$.

Definition. A connected, non-bipartite graph $\mathcal{G} = (V, E)$ is an (N, d, α) -*expander graph* if $|V| = N$, \mathcal{G} is d -regular, and the second largest eigenvalue magnitude of its transition matrix is at most α .

For the remainder of the section, \mathbf{A} is the transition matrix for an (N, d, α) -expander graph \mathcal{G} , $\alpha + \beta = 1$, and vectors $\mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{N-1} \in \mathbb{R}^n$ form an orthonormal basis, where \mathbf{u}_i is a left eigenvector associated with eigenvalue $\lambda_i(\mathbf{A})$. We take \mathbf{u}_0 to be $N^{-1/2}(1, 1, \dots, 1)$. For reference, we collect a few simple facts. Here the *projection matrix* for a set $S \subseteq \{0, 1, \dots, N-1\}$ is the matrix $\mathbf{P} = (p_{i,j})$ where $p_{i,i} = 1$ if $i \in S$ and all other entries $p_{i,j}$ are 0.

Proposition 3.1. *Let V_0 be the subspace of \mathbb{R}^N spanned by \mathbf{u}_0 and V_1 its orthogonal space, the subspace spanned by $\mathbf{u}_1, \dots, \mathbf{u}_{N-1}$.*

- (i) V_0 and V_1 are invariant under the action of \mathbf{A} .
- (ii) Every vector $\mathbf{w} \in \mathbb{R}^N$ can be decomposed as a sum of two vectors $\mathbf{w} = \mathbf{x} + \mathbf{y}$ where $\mathbf{x} \in V_0$ and $\mathbf{y} \in V_1$.
- (iii) If $\mathbf{x} \in V_0$, then $\mathbf{x}\mathbf{A} = \mathbf{x}$. If $\mathbf{y} \in V_1$, then $|\mathbf{y}\mathbf{A}| \leq \alpha|\mathbf{y}|$.
- (iv) If \mathbf{P} is the projection matrix for $T \subseteq \{0, 1, \dots, N-1\}$, then $|\mathbf{u}_0\mathbf{P}|^2 = |T|/N$.

Random walks on expander graphs with spectral gap β give rise to β -i.i.d. random objects. A t -walk on \mathcal{G} is a $(t+1)$ -tuple $\mathbf{y} = (y_0, y_1, \dots, y_t)$ of (not necessarily distinct) vertices such that $\{y_i, y_{i+1}\} \in E$ for $i < t$. Let Ω be the set of all t -walks on \mathcal{G} and \mathbb{P} be the uniform probability measure on Ω . We shall see that the projection functions U_i , defined by $U_i(\mathbf{y}) = y_i$ are β -independent.

Of course, we could just take the projection functions on the Cartesian product V^{t+1} , but many applications require reducing the number of bits needed to represent a sample point. We generate a t -walk in a unique way by picking an initial vertex y_0 and then choosing each successive vertex y_{i+1} by traversing one of the d edges incident with y_i . Representing a t -walk this way requires $\lceil \lg N \rceil + t\lceil \lg d \rceil$ bits while representing a point in V^{t+1} requires $(t+1)\lceil \lg N \rceil$ bits.

We now extend this idea to random walks on directed graphs that are expander-permutation hybrids. Take an (N, d, α) -expander graph $\mathcal{G} = (V, E)$ together with a permutation F on V and define the composition

$$E' = F \circ E = \{(u, F(v)) \mid \{u, v\} \in E\}$$

obtain a directed graph $\mathcal{G}' = (V, E')$ (possibly with loops). (Goldreich et al. implicitly use $E' = E \circ F$, but either construction works.) \mathcal{G}' is a d -regular directed graph in the sense

that every vertex has *both* indegree and outdegree d . Let Ω be the set of *directed t -walks* $\mathbf{y} = (y_0, y_1, \dots, y_t)$, where $(y_i, y_{i+1}) \in E'$ for $i < t$, \mathbb{P} be the uniform probability measure on Ω , and $U_i(\mathbf{y}) = y_i$, as before. Theorem 3.5 below shows that the projection functions U_i are β -independent.¹

\mathcal{G}' has transition matrix $\mathbf{A}' = \mathbf{A}\mathbf{B}$, where \mathbf{B} is the permutation matrix for F (i.e., the (i, j) entry of \mathbf{B} is 1 if $F(i) = j$ and 0 otherwise). \mathbf{A}' represents a step of a random walk on \mathcal{G}' . When it acts on a row vector representing a probability distribution on V , the result is the succeeding probability distribution along the random walk. \mathbf{A}' may also act on an *improper* probability distribution, i.e., one whose coordinates are nonnegative and sum to at most 1. Let Ω be the set of all t -walks on \mathcal{G}' and \mathbb{P} be the uniform probability measure on Ω . The *terminal probability vector* of $B \subseteq \Omega$ is a vector $(p_0, p_1, \dots, p_{N-1})$ where $p_j = \mathbb{P}(B \cap \{U_t = j\})$ for each vertex $j \in V$. This is an improper probability distribution whose coordinates sum to $\mathbb{P}(B)$.

The following result is well known and follows by standard techniques [22].

Proposition 3.2. *For \mathcal{G}' as above, let $T_i \subseteq V$ and \mathbf{P}_i be the projection matrix for T_i for $i \leq t$. Then the terminal probability vector for $\{\bigwedge_{i \leq t} U_i \in T_i\}$ is given by*

$$\mathbf{v} \mathbf{P}_0 \mathbf{A}' \mathbf{P}_1 \cdots \mathbf{P}_{t-1} \mathbf{A}' \mathbf{P}_t,$$

where $\mathbf{v} = N^{-1}(1, 1, \dots, 1)$.

The idea here is clear: the vector \mathbf{v} is the initial probability vector, \mathbf{A}' represents a step along the random walk, and each \mathbf{P}_i eliminates paths whose i -th vertex is not in T_i .

Now we require two lemmas.

Lemma 3.3. *Let \mathbf{A} , N , d , α , and β be as above. Take projection matrices \mathbf{P} and \mathbf{P}' for T and $T' \subseteq \{0, 1, \dots, N-1\}$ and set $\mu = |T|/N$, $\mu' = |T'|/N$. Then for all for all $\mathbf{v} \in \mathbb{R}^N$,*

$$|\mathbf{v} \mathbf{P} \mathbf{A} \mathbf{P}'| \leq (\alpha + \beta\mu)^{1/2} (\alpha + \beta\mu')^{1/2} |\mathbf{v}|$$

Proof. Let \mathbf{z} be a unit vector parallel to $\mathbf{v} \mathbf{P} \mathbf{A} \mathbf{P}'$ and put $\mathbf{w} = \mathbf{v} \mathbf{P}$, $\mathbf{w}' = \mathbf{z} \mathbf{P}'$ so

$$\begin{aligned} |\mathbf{v} \mathbf{P} \mathbf{A} \mathbf{P}'| &= \langle \mathbf{v} \mathbf{P} \mathbf{A} \mathbf{P}', \mathbf{z} \rangle \\ &= \langle \mathbf{v} \mathbf{P} \mathbf{A}, \mathbf{z} \mathbf{P}' \rangle \\ &= \langle \mathbf{w} \mathbf{A}, \mathbf{w}' \rangle \\ &= \langle \mathbf{x} \mathbf{A}, \mathbf{x}' \rangle + \langle \mathbf{x} \mathbf{A}, \mathbf{y}' \rangle + \langle \mathbf{y} \mathbf{A}, \mathbf{x}' \rangle + \langle \mathbf{y} \mathbf{A}, \mathbf{y}' \rangle, \end{aligned}$$

where $\mathbf{w} = \mathbf{x} + \mathbf{y}$ and $\mathbf{w}' = \mathbf{x}' + \mathbf{y}'$ are the decompositions given by Proposition 3.1(ii). Proposition 3.1(i) and (iii) imply $\langle \mathbf{x} \mathbf{A}, \mathbf{y}' \rangle = \langle \mathbf{y} \mathbf{A}, \mathbf{x}' \rangle = 0$, $|\langle \mathbf{x} \mathbf{A}, \mathbf{x}' \rangle| = |\langle \mathbf{x}, \mathbf{x}' \rangle| = |\mathbf{x}| |\mathbf{x}'|$, and $|\langle \mathbf{y} \mathbf{A}, \mathbf{y}' \rangle| \leq |\mathbf{y} \mathbf{A}| |\mathbf{y}'| \leq \alpha |\mathbf{y}| |\mathbf{y}'|$, so

$$|\mathbf{v} \mathbf{P} \mathbf{A} \mathbf{P}'| \leq |\mathbf{x}| |\mathbf{x}'| + \alpha |\mathbf{y}| |\mathbf{y}'|. \tag{3.1}$$

By the Cauchy-Schwarz inequality

$$\begin{aligned} |\mathbf{x}| |\mathbf{x}'| + |\mathbf{y}| |\mathbf{y}'| &\leq (|\mathbf{x}|^2 + |\mathbf{y}|^2)^{1/2} (|\mathbf{x}'|^2 + |\mathbf{y}'|^2)^{1/2} \\ &= |\mathbf{w}| |\mathbf{w}'| \end{aligned}$$

¹Some sources assert that since F is a permutation, \mathcal{G} and \mathcal{G}' have the same mixing properties. This is true, but since the adjacency matrix of \mathcal{G}' is not Hermitian, this assertion does not follow directly.

so $|y| |y'| \leq |w| |w'| - |x| |x'|$. Thus, substituting into (3.1),

$$|\mathbf{vPAP}'| \leq \alpha |w| |w'| + \beta |x| |x'|. \quad (3.2)$$

We must bound $|w|$, $|w'|$, $|x|$, and $|x'|$. First, $|w| = |\mathbf{vP}| \leq |\mathbf{v}|$ and $|w'| = |\mathbf{zP}'| \leq |\mathbf{z}| = 1$. Next, we know $\mathbf{x} + \mathbf{y} = \mathbf{vP}$. Take the inner product of both sides with \mathbf{u}_0 and observe that \mathbf{y} and \mathbf{u}_0 are orthogonal so $\langle \mathbf{x}, \mathbf{u}_0 \rangle = \langle \mathbf{vP}, \mathbf{u}_0 \rangle = \langle \mathbf{v}, \mathbf{u}_0 \mathbf{P} \rangle$. Then $|x| = |\langle \mathbf{v}, \mathbf{u}_0 \mathbf{P} \rangle| \leq |\mathbf{v}| |\mathbf{u}_0 \mathbf{P}| = \mu^{1/2} |\mathbf{v}|$ by Proposition 3.1(iv). Also, we know $\mathbf{x}' + \mathbf{y}' = \mathbf{zP}'$. Take the inner product of both sides with \mathbf{u}_0 as before. This time, $|x'| \leq |\mathbf{z}| |\mathbf{u}_0 \mathbf{P}'| = (\mu')^{1/2}$, again by Proposition 3.1(iv).

Substituting these four bounds into (3.2) gives

$$|\mathbf{vPAP}'| \leq (\alpha + \beta (\mu\mu')^{1/2}) |\mathbf{v}|.$$

By the Cauchy-Schwarz inequality,

$$\begin{aligned} \alpha + \beta (\mu\mu')^{1/2} &= \alpha^{1/2} \alpha^{1/2} + (\beta\mu)^{1/2} (\beta\mu')^{1/2} \\ &\leq (\alpha + \beta\mu)^{1/2} (\alpha + \beta\mu')^{1/2} \end{aligned}$$

which completes the proof. \square

Lemma 3.4. *Let \mathbf{A}' , N , d , α , and β be as above. Take projection matrices \mathbf{P} and \mathbf{P}' for T and $T' \subseteq \{0, 1, \dots, N-1\}$ and set $\mu = |T|/N$, $\mu' = |T'|/N$. Then for all $\mathbf{v} \in \mathbb{R}^N$,*

$$|\mathbf{vPA}'\mathbf{P}'| \leq (\alpha + \beta\mu)^{1/2} (\alpha + \beta\mu')^{1/2} |\mathbf{v}|.$$

Proof. We have

$$\begin{aligned} \mathbf{PA}'\mathbf{P}' &= \mathbf{PABP}' \\ &= \mathbf{PA}(\mathbf{BP}'\mathbf{B}^{-1})\mathbf{B} \\ &= \mathbf{PAP}''\mathbf{B}, \end{aligned}$$

where $\mathbf{P}'' = \mathbf{BP}'\mathbf{B}^{-1}$. It is easy to see that \mathbf{P}'' is the projection matrix for $F^{-1}[T']$. Now $|F^{-1}[T']|/N = |T'|/N = \mu'$, so by Lemma 3.3, $|\mathbf{vPAP}''| \leq (\alpha + \beta\mu)^{1/2} (\alpha + \beta\mu')^{1/2} |\mathbf{v}|$. Thus, since \mathbf{B} is a permutation matrix,

$$\begin{aligned} |\mathbf{vPA}'\mathbf{P}'| &= |\mathbf{vPAP}''\mathbf{B}| \\ &= |\mathbf{vPAP}''| \\ &\leq (\alpha + \beta\mu)^{1/2} (\alpha + \beta\mu')^{1/2} |\mathbf{v}| \end{aligned}$$

by the previous lemma. \square

We come now to the main result of this section giving the construction of β -independent random objects.

Theorem 3.5. *Let \mathcal{G}' be a hybrid expander-permutation directed graph where the expander graph has spectral radius β , Ω be the set of all directed t -walks in \mathcal{G}' , and \mathbb{P} be the uniform probability measure on Ω . Then the projection functions U_i are β -independent random objects.*

Proof. We need to show that for all $T_i \subseteq V$, $i \leq t$,

$$\mathbb{P}\{\bigwedge_{i \leq t} U_i \in T_i\} \leq \prod_{i \leq t} (\alpha + \beta \mu_i), \quad (3.3)$$

where $\mu_i = \mathbb{P}\{U_i \in T_i\}$.

We claim that it is enough to show this in the special case where $T_0 = T_t = V$. In this case $\{U_0 \in T_0\} = \{U_t \in T_t\} = \Omega$, $\mu_0 = \mu_t = 1$, and $\alpha + \beta \mu_0 = \alpha + \beta \mu_t = 1$. In effect, this eliminates constraints on the initial and terminal vertices of walks. Then (assuming $t > 2$) we may delete the initial and terminal vertices to obtain (3.3) for $(t-2)$ -walks rather than t -walks. That is, letting \mathbb{P}' be the uniform measure on the set of $(t-2)$ -walks,

$$\mathbb{P}'\{\bigwedge_{1 \leq i \leq t-1} U_i \in T_i\} = \mathbb{P}\{\bigwedge_{0 \leq i \leq t} U_i \in T_i\} = \prod_{1 \leq i \leq t-1} (\alpha + \beta \mu_i).$$

From Proposition 3.2 we know that the terminal probability vector for $\{\bigwedge_{0 \leq i \leq t} U_i \in T_i\}$ is given by

$$\mathbf{v} \mathbf{P}_0 \mathbf{A}' \mathbf{P}_1 \cdots \mathbf{P}_{t-2} \mathbf{A}' \mathbf{P}_{t-1}. \quad (3.4)$$

where $\mathbf{v} = N^{-1}(1, 1, \dots, 1)$. We may rewrite this as

$$\mathbf{v} (\mathbf{P}_0 \mathbf{A}' \mathbf{P}_1) (\mathbf{P}_1 \mathbf{A}' \mathbf{P}_2) \cdots (\mathbf{P}_{t-1} \mathbf{A}' \mathbf{P}_t)$$

since $\mathbf{P}_i \mathbf{P}_i = \mathbf{P}_i$ for $1 \leq i \leq t-1$. By Lemma 3.4, multiplication by $\mathbf{P}_i \mathbf{A}' \mathbf{P}_{i+1}$ changes the magnitude of a vector by at most a factor of $(\alpha + \beta \mu_i)^{1/2} (\alpha + \beta \mu_{i+1})^{1/2}$, so (3.4) is bounded in magnitude by

$$|\mathbf{v}| \prod_{i \leq t-1} (\alpha + \beta \mu_i)^{1/2} (\alpha + \beta \mu_{i+1})^{1/2}$$

which is equal to

$$|\mathbf{v}| (\alpha + \beta \mu_0)^{1/2} (\alpha + \beta \mu_t)^{1/2} \prod_{1 \leq i \leq t-1} (\alpha + \beta \mu_i).$$

By assumption, $\alpha + \beta \mu_0 = \alpha + \beta \mu_t = 1$, so we can further simplify this to

$$|\mathbf{v}| \prod_{i \leq t} (\alpha + \beta \mu_i).$$

Finally, compute the probability of $\{\bigwedge_{i \leq t} U_i \in T_i\}$ by summing the coordinates of its terminal probability vector (3.4); we do this by taking the inner product with the vector $\mathbf{u} = (1, 1, \dots, 1)^T$. The result is bounded in magnitude by

$$|\mathbf{u}| |\mathbf{v}| \prod_{i \leq t} (\alpha + \beta \mu_i).$$

But $|\mathbf{v}| = n^{-1/2}$ and $|\mathbf{u}| = n^{1/2}$, so we have established (3.3). \square

4. Invertibility and One-Way Functions

In this section we present definitions and notation concerning one-way functions.

For a set S , I_S is the identity function on S . Consider functions $F: S \rightarrow T$ and $G: T \rightarrow S$. We say G is a *right inverse* of F if $F \circ G = I_T$ and is a *left inverse* of F if $G \circ F = I_S$.²

We say F is a *partial function* from S to T (written $F: S \rightarrow T$) if it maps a subset of S (the *domain* of F , denoted $\text{dom}(F)$) onto a subset of T (the *range* of F , denoted $\text{ran}(F)$). If $F: S \rightarrow T$ and $G: T \rightarrow U$, the *composition* of F and G , written $G \circ F$, is a partial function from S to U that maps s to u precisely when there is a t such that $F(s) = t$ and $G(t) = u$. For partial functions $F: S \rightarrow T$ and $G: T \rightarrow S$, G is a *partial right inverse* of F if $F \circ G = I_{\text{ran}(F)}$ and is a *partial left inverse* of F if $G \circ F = I_{\text{dom}(F)}$.

Like functions, partial functions are injective if and only if they have a (partial) left inverse. Unlike functions, which are surjective if and only if they have a left inverse, partial functions always have a partial right inverse.³ In cryptography, invertibility of a partial function refers to existence of an *efficiently computable right inverse* of some kind.

Let $\varphi(n)$ be a proposition concerning the natural numbers $n \in \mathbb{N} = \{0, 1, 2, \dots\}$. We say that $\varphi(n)$ holds *infinitely often*, and write $\varphi(n)$ i.o. or $(\exists^\infty n) \varphi(n)$, if $\forall m (\exists n \geq m) \varphi(n)$. We say that $\varphi(n)$ holds *almost always*, and write $\varphi(n)$ a.a. or $(\forall^\infty n) \varphi(n)$, if $\exists m (\forall n \geq m) \varphi(n)$. \exists^∞ and \forall^∞ are dual quantifiers: $\neg(\exists^\infty n) \varphi(n)$ is equivalent to $(\forall^\infty n) \neg\varphi(n)$ and $\neg(\forall^\infty n) \varphi(n)$ is equivalent to $(\exists^\infty n) \neg\varphi(n)$. It is helpful to keep this in mind when negating statements.

A function $p: \mathbb{N} \rightarrow \mathbb{R}$ is *negligible* if $p(n) = n^{-\omega(1)}$, i.e., $(\forall c > 0) (\forall^\infty n) (|p(n)| \leq n^{-c})$. (Some sources use the term *superpolynomially small* rather than *negligible*.) In contrast, p is *significant* if $p(n) = n^{-O(1)}$, i.e., $(\exists c > 0) (\forall^\infty n) (p(n) \geq n^{-c})$. The term *negligible* is standard in cryptographic theory; the term *significant* is not.

Write $p(n) \approx q(n)$ if $|p(n) - q(n)|$ is negligible and $p(n) \gg q(n)$ if $p(n) - q(n)$ is significant.

Let F be a function from $\{0, 1\}^*$ to $\{0, 1\}^*$. Define the *auxiliary function* of F to be $\bar{F}(x) = (1^{|x|}, F(x))$, where $1^{|x|}$ is the unary representation of the input length. Auxiliary functions are convenient when defining weakly and strongly one-way functions because they inform an adversary attempting to invert F what the length of the preimage is. It is reasonable to suppose that an adversary would have this information.

We will take a probabilistic approach where the arguments of F are uniformly distributed random bit strings $X_n \in \{0, 1\}^n$ and $Y_n = F(X_n)$. Y_n may not be uniformly distributed; indeed, it need not have a fixed length for a given n .

We require another modification: a probabilistic adversary. Thus, the adversary attempting to find X_n such that $\bar{F}(X_n) = (1^n, Y_n)$ is a partial function $\bar{G}(1^n, Y_n, R_n)$ computable in time polynomial in $|1^n, Y_n|$, where R_n is a random bit string independent of X_n . We may assume that R_n is uniformly distributed on $\{0, 1\}^{q(n)}$ for some polynomial $q(n)$. We will say that \bar{G} is a p.p.t. function in this circumstance.

For each $n > 0$ and polynomial $q(n)$, $(1^n, Y_n, R_n)$ is a random vector and $\mathbb{P}_{(1^n, Y_n, R_n)}$, denoted for the sake of simplicity as \mathbb{P}^n , is an induced probability measure on $\Omega' = \{1\}^* \times \{0, 1\}^* \times \{0, 1\}^*$. (Strictly speaking, this notation should specify the particular polynomial

²Other common terms for left inverse are *retract* and *retraction*. Other common terms for right inverse are *coretraction* and *section*.

³The existence of partial right inverses for partial functions is equivalent to the Axiom of Choice. This is almost the same result as Axiom of Choice equivalent AC 5 in Rubin and Rubin [38], which says that the existence of right inverses for onto functions is equivalent to the Axiom of Choice.

$q(n)$ used, but this is cumbersome.) In the definitions below, the function $I(1^n, y, r) = (1^n, y)$ acts as an identity function when the adversary is the p.p.t. function \bar{G} .

Definition. A polynomial time computable function $F : \{0, 1\}^* \rightarrow \{0, 1\}^*$ is a *weakly one-way function* if

$$(\exists \delta \gg 0)(\forall \text{p.p.t. } \bar{G}) (\mathbb{P}^n \{\bar{F} \circ \bar{G} = I\} \leq 1 - \delta \text{ a.a.}),$$

or equivalently, there is a $c > 0$ such that for all p.p.t. \bar{G} , $\mathbb{P}^n \{\bar{F} \circ \bar{G} = I\} \leq 1 - n^{-c}$ a.a.

A polynomial time computable function F is a *strongly one-way function* if

$$(\forall \text{p.p.t. } \bar{G}) (\mathbb{P}^n \{\bar{F} \circ \bar{G} = I\} \approx 0),$$

or equivalently, for all $c > 0$ and all p.p.t. \bar{G} , $\mathbb{P}^n \{\bar{F} \circ \bar{G} = I\} \leq n^{-c}$ a.a. If, in addition to either of the conditions above, F is a length-preserving permutation (i.e., it is a bijection on $\{0, 1\}^n$ when restricted to strings of length n), we say it is a *weakly* or *strongly one-way permutation*.

Remark. The notation used in this paper differs from the notation used in other sources. A typical definition (similar to the one found in [13]) says that F is weakly one-way if there is a $c > 0$ such that for every p.p.t. G and all large n ,

$$\mathbb{P}\{G(1^n, F(x)) \notin F^{-1}F(x)\} > n^{-c}$$

where the probability is taken uniformly over $x \in \{0, 1\}^n$ and the random bits used by G . This is equivalent to the definition above.

Given \bar{F} and \bar{G} as above, let

$$W = \mathbb{P}^n \{\bar{F} \circ \bar{G} = I | I\}.$$

$W(1^n, y)$ is the probability that \bar{G} successfully finds a length n inverse of y . Taking the conditional expectation with respect to I averages over the random bit strings used by \bar{G} . Using this notation, we may prove a standard amplification result which yields a useful tail bound for W when F is a weakly one-way function [15, 13].

Proposition 4.1. *Let \bar{F} be an auxiliary function and \bar{G} be a randomized partial function with $W = \mathbb{P}^n \{\bar{F} \circ \bar{G} = I | I\}$.*

- (i) *Let \bar{G}' be the randomized partial function computed by independently computing \bar{G} k times on a given input $(1^n, y)$ (with fresh random bits each time) and returning the first value x such that $\bar{F}(x) = (1^n, y)$ (if there is such a value). Let $W' = \mathbb{P}^n \{\bar{F} \circ \bar{G}' = I | I\}$. Then for any $0 < \varepsilon < 1$,*

$$\mathbb{P}_I^n \{W > \varepsilon\} = \mathbb{P}_I^n \{W' > 1 - (1 - \varepsilon)^k\}.$$

- (ii) *If F is a weakly one-way function, where for some significant δ , every p.p.t. \bar{G} satisfies $\mathbb{P}^n \{\bar{F} \circ \bar{G} = I\} \leq 1 - \delta$, then for every significant $\varepsilon = \varepsilon(n)$,*

$$\mathbb{P}_I^n \{W > \varepsilon\} \leq 1 - \delta/2 \text{ a.a.}$$

Proof. Fix a value $(1^n, y)$ in the range of \bar{F} . $W(1^n, y) > \varepsilon$ asserts that the probability that $\bar{F} \circ \bar{G}(1^n, y) = (1^n, y)$ is greater than ε or, equivalently, the probability that $\bar{F} \circ \bar{G}(1^n, y) \neq (1^n, y)$ is less than $1 - \varepsilon$. This happens if and only if the probability that k independent computations of $\bar{F} \circ \bar{G}(1^n, y)$ fail to yield $(1^n, y)$ is less than $(1 - \varepsilon)^k$; and this happens if and only if $W'(1^n, y) > 1 - (1 - \varepsilon)^k$. This proves part (i).

Now suppose ε is significant. There is an integer d such that $\varepsilon > n^{-d}$ a.a. In part (i) take $k = n^{d+1}$ so almost always

$$\begin{aligned} \mathbb{P}_I^n\{W > \varepsilon\} &= \mathbb{P}_I^n\{W' > 1 - (1 - \varepsilon)^k\} \\ &\leq \mathbb{P}_I^n\{W' > 1 - (1 - n^{-d})^{n^{d+1}}\} \\ &\leq \mathbb{P}_I^n\{W' > 1 - e^{-n}\}. \end{aligned}$$

By Markov's inequality and the weakly one-way assumption, we have almost always

$$\begin{aligned} \mathbb{P}_I^n\{W' > 1 - e^{-n}\} &\leq \mathbb{E}(W')/(1 - e^{-n}) \\ &\leq (1 - \delta)/(1 - e^{-n}) \\ &= 1 - \delta + e^{-n}(1 - \delta)/(1 - e^{-n}) \\ &\leq 1 - \delta + e^{-n} \\ &\leq 1 - \delta/2, \end{aligned}$$

from which (ii) follows. □

Remark. Rephrased, Proposition 4.1(ii) says that $\mathbb{P}_I^n\{W \leq \varepsilon\} \geq \delta/2$ a.a., i.e., under the hypotheses of the proposition, W has a uniformly significant tail probability.

We often deal with partial functions defined only on arguments of certain prescribed lengths. The following technical result shows that under certain circumstances we can obtain weakly and strongly one-way functions (and permutations) from hard-to-invert partial functions by filling in undefined values.

Proposition 4.2. *Let $F: \{0, 1\}^* \rightarrow \{0, 1\}^*$ be polynomial time computable with domain $\bigcup_{m \geq 0} \{0, 1\}^{\tau(m)}$, where $\tau(m)$ is strictly increasing, computable in time polynomial in m , and for some $k > 0$, $\tau(m+1) \leq \tau(m)^k$. Define $F'(x)$ on strings x of length n as follows. Let m be the largest integer such that $\tau(m) \leq n$, put $x = x'z$ where $|x'| = \tau(m)$, and set $F'(x) = F(x')z$ (the concatenation of $F(x')$ and z).*

- (i) *If there is a $c > 0$ such that for all p.p.t. \bar{G} , $\mathbb{P}^{\tau(m)}\{\bar{F} \circ \bar{G} = I\} \leq 1 - \tau(m)^{-c}$ a.a., then F' is a weakly one-way function.*
- (ii) *If for all $c > 0$ and all p.p.t. \bar{G} , $\mathbb{P}^{\tau(m)}\{\bar{F} \circ \bar{G} = I\} \leq \tau(m)^{-c}$ a.a., then F' is strongly one-way.*
- (iii) *If for every $m \geq 0$, F is a permutation (i.e., its restriction to each domain $\{0, 1\}^{\tau(m)}$ is a bijection), then F' is also a permutation.*

Proof. Part (iii) is obvious.

For parts (i) and (ii) we derive an upper bound for $\mathbb{P}^n\{\bar{F}' \circ \bar{G}' = I\}$, where \bar{G}' is an arbitrary p.p.t. function.

From \bar{G}' construct a p.p.t. function \bar{G} that attempts to invert \bar{F} as follows on input $(1^{\tau(m)}, y)$. It takes successive values of n in the interval $\tau(m) \leq n < \tau(m+1)$, each time choosing a random bit string z of length $n - \tau(m)$ and applying \bar{G}' to $(1^n, yz)$; if the result is of the form xz and $F(x) = y$, it returns the value x and terminates the computation.

In (i) of the proposition, there is a $c > 0$ such that

$$\mathbb{P}^{\tau(m)}\{\bar{F} \circ \bar{G} = I\} \leq 1 - \tau(m)^{-c} \text{ a.a.}$$

\bar{G} finds an inverse image x (with respect to \bar{F}) of $(1^{\tau(m)}, y)$ only if there is an n in the range $\tau(m) \leq n < \tau(m+1)$ such that \bar{G}' finds an inverse image xz of $(1^n, yz)$. But $1 - \tau(m)^{-c} \leq 1 - n^{-c}$, so

$$\mathbb{P}^n\{\bar{F}' \circ \bar{G}' = I\} \leq 1 - n^{-c} \text{ a.a.}$$

Therefore, F' is a weakly one-way function.

In part (ii) of the proposition, for all $c > 0$ and all p.p.t. \bar{G} ,

$$\mathbb{P}^{\tau(m)}\{\bar{F} \circ \bar{G} = I\} \leq \tau(m)^{-c} \text{ a.a.}$$

Again, \bar{G} finds an inverse x of $(1^{\tau(m)}, y)$ only if there is an n in the range $\tau(m) \leq n < \tau(m+1)$ such that \bar{G}' finds an inverse xz of $(1^n, yz)$. But $n < \tau(m+1) \leq \tau(m)^k$ a.a., so

$$\mathbb{P}^n\{\bar{F}' \circ \bar{G}' = I\} \leq n^{-c/k} \text{ a.a.}$$

and, therefore, F' is a strongly one-way function. \square

5. From Weakly to Strongly One-Way.

We now show that the existence of a weakly one-way function implies the existence of a strongly one-way function. As noted in the introduction, the first published proof [13] implicitly defines a reduction $(\mathcal{R}, \mathcal{R}^*)$ between function inversion problems. Let us make this precise.

Recall that from F we define F' by

$$F'(x_0 x_1 \cdots x_{t-1}) = (F(x_0), F(x_1), \dots, F(x_{t-1}))$$

where $t = t(n)$ is a polynomially bounded function and $|x_i| = n$ for all i . This defines F' only for inputs $x' = x_0 x_1 \cdots x_{t-1}$ of length $n t(n)$, but the Proposition 4.2 allows us to extend F' to a total function. It is convenient to view \mathcal{R} as a polynomial time oracle Turing machine that computes the auxiliary function \bar{F}' by making queries to evaluate \bar{F} at the values x_0, x_1, \dots, x_{t-1} . Similarly, \mathcal{R}^* is a probabilistic polynomial time oracle Turing machine that computes \bar{G} by making queries to evaluate the function \bar{F} and the randomized function \bar{G}' at various values.

We now show how the probability inequality in Theorem 2.1(i) figures in Goldreich's proof [13].

Theorem 5.1. *Suppose F is a weakly one-way function, so that for some integer $c > 0$,*

$$(\forall \text{ p.p.t. } \bar{G}) (\mathbb{P}^n\{\bar{F} \circ \bar{G} = I\} \leq 1 - n^{-c} \text{ a.a.}).$$

Then F' , defined as above with $t = n^{c+1}$, is a strongly one-way function.

Proof. Take an arbitrary p.p.t. \bar{G}' . By Proposition 4.2(ii) it is enough to show that for all $d > 0$,

$$\mathbb{P}^{nt}\{\bar{F}' \circ \bar{G}' = I\} \leq (nt)^{-d} \text{ a.a.}$$

Let $Z = \mathbb{P}^{nt}\{\bar{F}' \circ \bar{G}' = I|I\}$ so $\mathbb{E}(Z) = \mathbb{P}^{nt}\{\bar{F}' \circ \bar{G}' = I\}$ by the law of iterated expectations.

\mathcal{R}^* computes a randomized function \bar{G} , which attempts to find the inverse of \bar{F} at $(1^n, y)$, as follows.

1. \mathcal{R}^* forms $\mathbf{y} = (y_0, y_1, \dots, y_{t-1})$ by choosing a random $i < t$ and setting $y_i = y$; then generating random $x'_j \in \{0, 1\}^n$ for each $j \neq i$ and putting $y_j = F(x'_j)$. (Here it queries the \bar{F} oracle).
2. It queries the \bar{G}' oracle on $(1^{nt}, \mathbf{y})$ and receives an answer $x_0 x_1 \cdots x_{t-1}$, where each x_j is of length n .
3. It checks that $F(x_j) = y_j$ for all $j < t$ (again, by querying the \bar{F} oracle) and, if so, returns x_i ; otherwise, the function is undefined.

Clearly, when this procedure returns a value x_i , $(1^n, y) = \bar{F}(x_i)$.

Define $U_i(1^{nt}, \mathbf{y}) = (1^n, y_i)$. The random objects U_i are i.i.d. The probability that \bar{G}' successfully finds an inverse for $(1^{nt}, \mathbf{y})$ given that $y = y_i$ is $W_i(y)$, where $W_i = \mathbb{E}(Z|U_i)$. Therefore, $\mathbb{P}^n\{\bar{F} \circ \bar{G} = I|I\}$ is $W = (\sum_{i < t} W_i)/t$. It is at this point we bypass many of the details in the original proof and invoke Theorem 2.1(i):

$$\mathbb{E}(Z) \leq \mathbb{P}_I^n\{W > \varepsilon\}^t + t\varepsilon.$$

Let d be an arbitrary positive integer and put $\varepsilon = n^{-d}t^{-d-1}/2$. By Proposition 4.1, $\mathbb{P}_I^n\{W > \varepsilon\} \leq 1 - n^{-c}/2$ a.a. Thus, almost always

$$\begin{aligned} \mathbb{P}_I^n\{W > \varepsilon\}^t &\leq (1 - n^{-c}/2)^{n^{c+1}} \\ &\leq (e^{-n^{-c}/2})^{n^{c+1}} \\ &= e^{-n/2} \\ &\leq (nt)^{-d}/2. \end{aligned}$$

Also, $\varepsilon t = (nt)^{-d}/2$. Consequently, $\mathbb{P}^{nt}\{\bar{F}' \circ \bar{G}' = I\} = \mathbb{E}(Z) \leq (nt)^{-d}$ a.a. \square

Using Proposition 4.2, we have the following corollary.

Corollary 5.2.

- (i) *If weakly one-way functions exist, then strongly one-way functions exist.*
- (ii) *If weakly one-way permutations exist, then strongly one-way permutations exist.*

6. A Security Preserving Reduction for One-Way Permutations.

In this section we restrict our attention to one-way permutations. Since $|F(x)| = |x|$, it is not necessary to use an auxiliary function $\bar{F}(x) = (1^{|x|}, F(x))$ – an adversary can infer this information from $|F(x)|$.

Definitions of *security preserving reduction* differ on details [15, 8, 28, 16, 14], but follow the same general pattern. $(\mathcal{R}, \mathcal{R}^*)$ is a reduction from a cryptographic primitive F to a cryptographic primitive G , as in the previous section, but in addition, when \mathcal{R}^* computes F' from G' , the security of F' against G' is at least as great as the security of F against G . For one-way permutations, security means resilience against inversion. To be precise, if G is a randomized function computed in time $T(n)$ (not necessarily a polynomial) and $\varepsilon(n)$ is the probability that G inverts F , the *security* of F against G is $S(n) = T(n)/\varepsilon(n)$. This is essentially the expected time to invert some element with respect to F by applying G independently to random elements in the range of F . We will write $S'(n) \succeq S(n)$ ($S'(n)$ *dominates* $S(n)$) if there are positive constants k and c such that $S'(cn) \geq S(n)/n^k$ a.a., and $S'(n) \asymp S(n)$ ($S'(n)$ and $S(n)$ are *of the same order*) if $S'(n) \succeq S(n)$ and $S(n) \succeq S'(n)$.

The reduction used in Theorem 5.1 is not security preserving because \mathcal{R}^* makes just one query to the G' oracle and it is of length n^{c+2} . If the security of F' against G' is $S'(n)$, then the security of F against G is of the same order as $S'(n^{c+2})$, which is not of the same order as $S'(n)$ in general.

We apply Theorem 2.1(ii) to show that the reduction of Goldreich et al. [15] is security preserving, in fact, that $S(n)$ is essentially $S'(n + \omega(\log n))$, which is of the same order as $S'(n)$. This reduction applies just to one-way permutations rather than arbitrary one-way functions. It uses the set of t -walks in a hybrid expander-permutation directed graph in place of a direct power and β -independence in place of independence.

The expander graphs used for this reduction must be from a fully explicit family, defined as follows.

Definition. Let $\mathcal{G}_m, m \geq 0$, be a family of d -regular graphs where \mathcal{G}_m has vertex set $V_m = \{0, 1, \dots, N_m - 1\}$ with $N_0 < N_1 < N_2 < \dots$. A *rotation function* $R(N, u, j)$ for this family is a partial function satisfying the following conditions.

1. $R(N, u, j)$ is defined if and only if for some m , $N = N_m$, $0 \leq u < N_m$, and $0 \leq j < d$.
2. For each vertex there is a linear order on the d adjacent vertices such that $R(N_m, u, j) = (v, k)$ holds precisely when v is the j -th vertex adjacent to u and u is the k -th vertex adjacent to v .

A family of d -regular graphs is *fully explicit* if it has a polynomial time computable rotation function.

If $\mathcal{G}_m, m \geq 0$, has a rotation function $R(N, u, j)$ such that for every edge $\{u, v\}$ in \mathcal{G}_m there is a j such that $R(N_m, u, j) = (v, j)$, we let $\kappa(u, v) = j$. Thus, κ defines an *edge coloring*: κ assigns a color to each edge such that at each vertex incident edges are uniquely colored. When this occurs for a polynomial time computable $R(N, u, j)$ we have a *fully explicit edge coloring*.

There is an extensive literature on the construction of fully explicit families of (N, d, α) -expander graphs [32, 12, 25, 37, 1, 3].

Remark. Most applications involving fully explicit expander graph families require that the gap between N_m and N_{m+1} not grow too quickly. We require more, viz., that N_0, N_1, N_2, \dots be a smoothly growing sequence of powers of two with $N_m = 2^{cm}$ for some constant c , and that d be a fixed power of two, say 2^e . For the remainder of this section we will assume that $\mathcal{G}_m, m \geq 0$, is a fully explicit (N, d, α) -expander graph family satisfying these conditions,

with d fixed and $\alpha < 1$. Hence, $\mathcal{G}_m = (\{0, 1\}^n, E_m)$ with $n = cm$. As a notational convenience, we will take $E = \bigcup_{m \geq 0} E_m$ and write $\mathcal{G}_m = (\{0, 1\}^n, E)$ rather than $\mathcal{G}_m = (\{0, 1\}^n, E_m)$.

One example of an explicit family of expander graphs satisfying these conditions is the affine torus expander graph family of Margulis [32]. Gabber and Galil [12] established an upper bound for the second largest eigenvalue magnitude of graphs in this family, later improved by Jimbo and Maruoka [25]. Using these results, we may take $n = 2m$ (so $N_m = 2^{2m}$), $d = 8 = 2^3$, and $\alpha = 5\sqrt{2}/8 = 0.88388\dots$. Other constructions may give a better bound for α . The results in [15] require that $\alpha \leq 1/2$, but the approach here based on β -independence works for any fixed bound less than 1.

Goldreich et al. [15] require an expander graph family which has a fully explicit edge coloring (but use different terminology). Many explicit expander graph constructions do, in fact, have a fully explicit edge coloring, but we will extend the proof in [15] so that we may dispense with this assumption.

Let $t = t(n)$ be a polynomially bounded, strictly increasing function. We first describe the transformation \mathcal{R} taking F , a weakly one-way permutation, to F' , a slightly harder to invert permutation.

Take $E' = F \circ E$. This gives a family of directed graphs $\mathcal{G}'_m = (\{0, 1\}^n, E')$. Note that for each directed edge (u, w) in \mathcal{G}'_m , there is a unique v such that $\{u, v\} \in E$ and $F(v) = w$. This suggests two ways to color the directed edges of \mathcal{G}'_m . If $R(N_m, u, j) = (v, k)$, we have the coloring $\kappa(u, w) = j$ and the coloring $\kappa'(u, w) = k$. Thus, κ is an explicit out-edge coloring in the sense that at every vertex u of \mathcal{G}'_m , the d out-edges are uniquely colored; and κ' is an explicit in-edge coloring in the sense that at every vertex w , the d in-edges are uniquely colored.

Let $\mathbf{x} = (x_0, x_1, \dots, x_t)$ be a directed t -walk in \mathcal{G}'_m . The *forward representation* of \mathbf{x} is

$$\varphi(\mathbf{x}) = (x_0, \kappa(x_0, x_1), \kappa(x_1, x_2), \dots, \kappa(x_{t-1}, x_t)).$$

In effect, we regard \mathcal{G}'_m and its out-edge-coloring as a finite automaton with alphabet $\{0, 1, \dots, d-1\}$, and $\kappa(x_0, x_1)\kappa(x_1, x_2)\cdots\kappa(x_{t-1}, x_t)$ as the unique string which, when read, causes this finite automaton to transition through the states x_0, x_1, \dots, x_t . This walk representation uses fewer bits than just listing vertices. Clearly, φ is a bijection from the set of directed t -walks in \mathcal{G}' to $V \times \{0, 1, \dots, d-1\}^t$. We will identify $V \times \{0, 1, \dots, d-1\}^t$ with $\{0, 1\}^{n+te}$. Since R and F are polynomial time computable, so are φ and φ^{-1} .

The *reverse representation* of \mathbf{x} is

$$\rho(\mathbf{x}) = (x_t, \kappa'(x_{t-1}, x_t), \kappa'(x_{t-2}, x_{t-1}), \dots, \kappa'(x_0, x_1)).$$

View this as taking \mathcal{G}'_m together with its in-edge-coloring, reversing the edge directions to form another finite automaton, and specifying a succinct walk representation as before. As with φ , ρ is a bijection from the set of directed t -walks in \mathcal{G}' to $\{0, 1\}^{n+te}$. It is easy to see that ρ is polynomial time computable, but since F is weakly one-way, it does not follow that ρ^{-1} is polynomial time computable.

We now describe how \mathcal{R} computes F' , a permutation on $\{0, 1\}^{n+te}$, from F , a permutation on $\{0, 1\}^n$, where $n = cm$. For each t -walk \mathbf{x} in \mathcal{G}'_m , F' maps $\varphi(\mathbf{x})$ to $\rho(\mathbf{x})$. In other words, $F' = \rho \circ \varphi^{-1}$. To compute $F'(x_0, k_1, \dots, k_t)$, \mathcal{R} begins at vertex x_0 in \mathcal{G}_m , then repeatedly follows the k_i -th edge with respect to the coloring κ from the current vertex x_i and applies F to jump to a new vertex x_{i+1} . In this way, it computes a t -path

$\mathbf{x} = (x_0, x_1, \dots, x_t)$. As it traverses each edge in \mathcal{G}'_m dictated the out-edge coloring k_i (with respect to κ), it computes the corresponding in-edge coloring (with respect to κ'), then at the terminus x_t reverses the in-edge color sequence to output $\rho(\mathbf{x})$. Clearly, F' is a permutation. By this definition, F' is defined only on strings of length $cm + te$ for $m \geq 0$, but we may use Proposition 4.2 to extend F' so that it is defined on strings of any length.

Lemma 6.1. *From a weakly one-way permutation F , construct F' as above with polynomially bounded $t = t(n)$. Suppose $\delta = \delta(n)$ is significant and that for all p.p.t. G ,*

$$\mathbb{P}^n\{F \circ G = I\} \leq 1 - \delta \text{ a.a.}$$

Then the following hold.

(i) For every p.p.t. G' ,

$$\mathbb{P}^{n+te}\{F' \circ G' = I\} \leq (1 - \beta\delta(n)/2)^t \text{ a.a.}$$

(ii) If $t \geq 7/\beta$, then for every p.p.t. G' ,

$$\mathbb{P}^{n+te}\{F' \circ G' = I\} \leq \max(1 - 2\delta(n), 1/2) \text{ a.a.}$$

(iii) If $\delta(n) \geq 1/2$ a.a. and $t = \omega(\log n)$, then for every p.p.t. G' ,

$$\mathbb{P}^n\{F' \circ G' = I\} \approx 0.$$

Proof. (i) Let $Z = \mathbb{P}^{n+te}\{F' \circ G' = I | I\}$ so $\mathbb{E}(Z) = \mathbb{P}^{n+te}\{F' \circ G' = I\}$. As in the proof of Theorem 5.1, we have \mathcal{R} taking F to F' and must specify \mathcal{R}^* taking each p.p.t. function G' , which attempts to invert F' , to another p.p.t. function G , which attempts to invert F . On a given input y , \mathcal{R}^* , querying oracles for G' and F , computes G as follows.

1. \mathcal{R}^* chooses a random i in the interval $1 \leq i < t$, then generates a random sequence of integers k_1, k_2, \dots, k_{t-i} , where each k_j is in the range $0 \leq k_j < d$; $(y, k_1, k_2, \dots, k_{t-i})$ is the forward walk representation of a random $(t-i)$ -walk in \mathcal{G}'_m with initial vertex y . \mathcal{R}^* applies F to obtain the reverse walk representation $(y_t, k'_1, k'_2, \dots, k'_{t-i})$ then generates random integers $k'_{t-i+1}, k'_{t-i+2}, \dots, k'_t$ in the range $0 \leq k'_j < d$ to obtain $\mathbf{y} = (y_t, k'_1, k'_2, \dots, k'_t)$, the reverse walk representation of a t -walk (y_0, y_1, \dots, y_t) chosen randomly from t -walks such that $y_i = y$.
2. \mathcal{R}^* queries the G' oracle on \mathbf{y} and receives an answer

$$\mathbf{x} = (x_0, k''_1, k''_2, \dots, k''_t).$$

3. \mathcal{R}^* applies φ^{-1} to \mathbf{x} to obtain a purported walk (x_0, x_1, \dots, x_t) , and then applies ρ . If the result matches \mathbf{y} , \mathcal{R}^* has verified that (x_0, x_1, \dots, x_t) is indeed a t -walk and that $x_i = y$ and (x_{i-1}, x_i) is an edge in \mathcal{G}'_m . In this case, there is a v such that $\{x_{i-1}, v\}$ is an edge in \mathcal{G}_m and $F(v) = x_i$, so \mathcal{R}^* returns $v = R(2^n, x_{i-1}, k''_i)$; otherwise, the function is undefined.

When this procedure returns a value v , $F(v) = y$.

Define $U_i(\mathbf{y}) = y_i$, where $\rho^{-1}(\mathbf{y}) = (y_0, y_1, \dots, y_t)$. By Theorem 3.5 and the bijectivity of ρ , the random objects U_i are β -i.i.d. The probability that G' successfully finds an inverse of \mathbf{y} , given that $y = y_i$, is $W_i(y)$, where $W_i = \mathbb{E}(Z|U_i)$. Therefore, $\mathbb{P}^n\{F \circ G = I|I\}$ is $W = (\sum W_i)/t$. By Theorem 2.1(ii),

$$\mathbb{E}(Z) \leq (\alpha + \beta \mathbb{P}_I^n\{W > \varepsilon\})^t + t\varepsilon.$$

By Proposition 4.1, $\mathbb{P}_I^n\{W > \varepsilon\} \leq 1 - \delta/2$ a.a. Thus, almost always

$$\begin{aligned} (\alpha + \beta \mathbb{P}_I^n\{W > \varepsilon\})^t &\leq (\alpha + \beta(1 - \delta/2))^t \\ &= (1 - \beta\delta/2)^t. \end{aligned}$$

Hence, for every significant ε , $\mathbb{E}(Z) \leq (1 - \beta\delta/2)^t + t\varepsilon$, which proves (i).

(ii) The function $f(x) = (1 - \beta x)^t$ is convex for $x \geq 0$ and

$$\begin{aligned} f((\beta t)^{-1}) &= (1 - 1/t)^t \\ &< e^{-1} \end{aligned}$$

so it lies below the continuous piecewise linear function

$$g(x) = \begin{cases} 1 - (1 - e^{-1})\beta t x, & \text{if } 0 \leq x \leq (\beta t)^{-1}, \\ e^{-1}, & \text{if } x > (\beta t)^{-1}. \end{cases}$$

In other words, $f(x) \leq g(x) = \max(1 - (1 - e^{-1})\beta t x, e^{-1})$ (see Figure 1). Thus, setting $x = \delta/2$ gives

$$(1 - \beta\delta/2)^t \leq \max(1 - (1 - e^{-1})\beta t\delta/2, e^{-1}).$$

But $\beta t \geq 7$ and $(1 - e^{-1})/2 = 0.31606\dots$, so from part (i) of the theorem

$$\mathbb{P}^{n+te}\{F' \circ G' = I\} \leq \max(1 - 2\delta, 1/2) \text{ a.a.}$$

(iii) This also follows from part (i). Since $\delta \geq 1/2$ a.a., $1 - \beta\delta/2 \leq 1 - \beta/4$ a.a. We know $t = \omega(\log n)$ a.a., so $(1 - \beta\delta/2)^t$ is negligible. \square




Figure 1: Function $f(x)$ lies below $g(x)$.

To conclude, we show that the reduction of Goldreich et al. [15] from a weakly one-way permutation to a strongly one-way permutation is security preserving.

First, consider the reduction $(\mathcal{R}_0, \mathcal{R}_0^*)$ described in the proof of Lemma 6.1(ii) where $t \geq 1/\beta$ is an even integer. (Recall that $n = 2m$ and $e = 3$, so an even t ensures that $n + te$ is even.) If F is a weakly one-way permutation, where there is a significant δ such that for all p.p.t. G $\mathbb{P}^n\{F \circ G = I\} \leq 1 - \delta$ a.a., then applying \mathcal{R}_0 s times to F results in a permutation F' such that for any p.p.t. G' attempting to invert F'

$$\mathbb{P}^{n+рте}\{F' \circ G' = I\} \leq \max(1 - 2^s \delta, 1/2) \text{ a.a.}$$

We know that for some $c > 0$, $\delta(n) < n^{-c}$ a.a., so, setting $s = \lceil c \log n \rceil$, we have that for every p.p.t. G' , $\mathbb{P}^n\{F' \circ G' = I\} \leq 1/2$ a.a.

Next, consider the reduction $(\mathcal{R}_1, \mathcal{R}_1^*)$ described in the proof of Lemma 6.1(iii) where $t = \omega(n)$. Applying \mathcal{R}_1 to F' results in a permutation F'' such that for all p.p.t. G'' attempting to invert F'' ,

$$\mathbb{P}^n\{F'' \circ G'' = I\} \approx 0.$$

Thus, a transformation \mathcal{R} consisting of s applications of \mathcal{R}_0 followed by an application of \mathcal{R}_1 takes F to F'' , thereby increasing input length from n to $n + \omega(\log n)$. Transformation \mathcal{R}^* consisting of an application of \mathcal{R}_1^* followed by s applications of \mathcal{R}_0^* takes p.p.t. function G'' to p.p.t. function G , thereby decreasing input length from $n + \omega(\log n)$ to n . \mathcal{R}^* , computing G , queries the oracle for G'' just once and the probability that G inverts F is precisely the probability that G'' inverts F'' . \mathcal{R}^* runs in polynomial time (assuming constant time to answer a query). Thus, we have the following result.

Theorem 6.2. *For reduction $(\mathcal{R}, \mathcal{R}^*)$ described above, if the security of F'' against G'' is $S''(n)$, then the security of F against G is $S(n) \asymp S''(n + \omega(\log n))$. Thus, $S(n) \asymp S''(n)$. That is, $(\mathcal{R}, \mathcal{R}^*)$ is a security preserving reduction taking weakly to strongly one-way functions.*

Remark. The reduction of Theorem 6.2 takes every weakly one-way permutation to a strongly one-way permutation. In contrast, the proof of Theorem 5.1 shows that for every weakly one-way function there is a reduction to a strongly one-way function. It is not apparent from this proof that there is just one reduction that takes every weakly one-way function to a strongly one-way function. Thus, the reduction of Theorem 6.2 is security preserving but is also uniform in this sense.

7. Conclusion.

One-way functions are not the only cryptographic primitives proved secure using repetition or expander graph constructions to amplify hardness. Others include collision-resistant hash functions [6], encryption schemes [10], weakly verifiable puzzles [5, 24, 26], signature schemes and message authentication codes [9], commitment schemes [19, 7], pseudorandom functions and pseudorandom generators [9, 34], block ciphers [31, 35, 33, 40], and interactive protocols [36, 21, 17]. Another approach to constructing security preserving reductions uses hash functions rather than expander graphs to control input size blowup [8, 18, 4]. Conditional expectation bounds may help simplify proofs in some of these cases and point the way to other cryptographic applications.

References

- [1] Noga Alon, Oded Schwartz, and Asaf Shapira. An elementary construction of constant-degree expanders. *Combinatorics, Probability and Computing*, 17:319–327, 2008.
- [2] Robert B. Ash and Catherine A. Doléans-Dade. *Probability and Measure Theory*. Academic Press, San Diego, CA, USA, second edition, 2000.
- [3] Avraham Ben-Aroya and Amnon Ta-Shma. A combinatorial construction of almost-Ramanujan graphs using the zig-zag product. In *Proc. 40th Annual ACM Symp. on Theory of Computing*, pages 325–334. ACM, 2008.
- [4] Andre Bogdanov and Alon Rosen. Input locality and hardness amplification. *J. Cryptology*, 26:144–171, 2013.
- [5] Ran Canetti, Shai Halevi, and Michael Steiner. Hardness amplification of weakly verifiable puzzles. In Joe Kilian, editor, *2nd Theory of Cryptography Conference - TCC 2005*, volume 3378 of *Lecture Notes in Computer Science*, pages 17–33. Springer, 2005.
- [6] Ran Canetti, Madhu Sudan, Luca Trevisan, Salil Vadhan, and Hoeteck Wee. Amplifying collision resistance: A complexity-theoretic treatment. In Alfred Menezes, editor, *Advances in Cryptology Crypto 2007*, volume 4622 of *Lecture Notes in Computer Science*, pages 264–283. Springer, 2007.
- [7] Kai-Min Chung, Feng-Hao Liu, Chi-Jen Lu, and Bo-Yin Yang. Efficient string-commitment from weak bit-commitment. In Masayuki Abe, editor, *Advances in Cryptology - ASIACRYPT 2010*, volume 6477 of *Lecture Notes in Computer Science*, pages 268–282. Springer, 2010.
- [8] Giovanni Di Crescenzo and Russell Impagliazzo. Security-preserving hardness-amplification for any regular one-way function. In *Proc. 31st ACM Symp. on Theory of Computing*, pages 169–178, New York, 1999. Association for Computing Machinery.
- [9] Yevgeniy Dodis, Russell Impagliazzo, Ragesh Jaiswal, and Valente Kabanets. Security amplification for interactive cryptographic primitives. In Omer Reingold, editor, *6th Theory of Cryptography Conference, TCC 2009*, volume 5444 of *Lecture Notes in Computer Science*, pages 128–145. Springer, 2009.
- [10] Cynthia Dwork, Moni Naor, and Omer Reingold. Immunizing encryption schemes from decryption errors. In Christian Cachin and Jan L. Camenisch, editors, *Advances in Cryptology - EUROCRYPT 2004*, volume 3027 of *Lecture Notes in Computer Science*, pages 342–360. Springer Berlin Heidelberg, 2004.
- [11] William Feller. *Introduction to Probability Theory and its Applications*, volume 2. Wiley, New York, 1971.
- [12] Ofer Gabber and Zvi Galil. Explicit constructions of linear-sized superconcentrators. *Journal of Computer and System Sciences*, 22:407 – 420, 1981.
- [13] Oded Goldreich. *Foundations of Cryptography: Basic Tools*, volume 1. Cambridge, New York, 2001.

- [14] Oded Goldreich. On security preserving reductions – revised terminology. In Oded Goldreich, editor, *Studies in Complexity and Cryptography. Miscellanea on the Interplay between Randomness and Computation*, pages 540–546. Springer, Berlin, Heidelberg, 2011.
- [15] Oded Goldreich, Russell Impagliazzo, Leonid A. Levin, Ramarathnam Venkatesan, and David Zuckerman. Security preserving amplification of hardness. In *Proc. 31st IEEE Symp. on Foundations of Computer Science, FOCS 1990*, pages 318–326. IEEE Computer Society Press, 1990.
- [16] Oded Goldreich, Leonid A. Levin, and Noam Nisan. On constructing 1-1 one-way functions. In Oded Goldreich, editor, *Studies in Complexity and Cryptography. Miscellanea on the Interplay between Randomness and Computation*, pages 13–25. Springer, Berlin, Heidelberg, 2011.
- [17] Iftach Haitner. A parallel repetition theorem for any interactive argument. *54th Annual IEEE Symposium on Foundations of Computer Science, FOCS 2013*, pages 241–250, 2009.
- [18] Iftach Haitner, Danny Harnik, and Omer Reingold. On the power of the randomized iterate. In Cynthia Dwork, editor, *Advances in Cryptology - CRYPTO 2006*, volume 4117 of *Lecture Notes in Computer Science*, pages 22–40. Springer Berlin Heidelberg, 2006.
- [19] Shai Halevi and Tal Rabin. Degradation and amplification of computational hardness. In Ran Canetti, editor, *5th Theory of Cryptography Conference, TCC 2008*, volume 4948 of *Lecture Notes in Computer Science*, pages 626–643. Springer, 2008.
- [20] Paul Halmos. *Measure Theory*. Graduate Texts in Mathematics. Springer, Berlin, 1974.
- [21] Johan Håstad, Rafael Pass, Douglas Wikström, and Krzysztof Pietrzak. An efficient parallel repetition theorem. In Daniele Micciancio, editor, *7th Theory of Cryptography Conference, TCC 2010*, volume 5978 of *Lecture Notes in Computer Science*, pages 1–18. Springer, 2010.
- [22] Shlomo Hoory, Nathan Linial, and Avi Wigderson. Expander graphs and their applications. *Bull. Amer. Math. Soc.*, 43:439–561, 2006.
- [23] Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. Cambridge Univ. Press, New York, 1985.
- [24] Russell Impagliazzo, Ragesh Jaiswal, and Valentine Kabanets. Chernoff-type direct product theorems. *Journal of Cryptology*, 22:75–92, 2009.
- [25] Shuji Jimbo and Akira Maruoka. Expanders obtained from affine transformations. *Combinatorica*, 7:343–355, 1987.
- [26] Charanjit S. Jutla. Almost optimal bounds for direct product threshold theorem. In Daniele Micciancio, editor, *7th Theory of Cryptography Conference, TCC 2010*, volume 5978 of *Lecture Notes in Computer Science*, pages 37–51. Springer, 2010.

- [27] Andrei N. Kolmogorov. *Grundbegriffe der Wahrscheinlichkeitsrechnung*. Ergebnisse der Mathematik und ihrer Grenzgebiete. Springer, Berlin, 1933. Translated by Nathan Morrison as *Foundations of Probability*, Chelsea, New York, 1950.
- [28] Henry C. Lin, Luca Trevisan, and Hoeteck Wee. On hardness amplification of one-way functions. In Joe Kilian, editor, *Second Theory of Cryptography Conference - TCC 2005*, volume 3378 of *Lecture Notes in Computer Science*, pages 34–49. Springer, 2005.
- [29] Michel Loève. *Probability Theory II*. Graduate Texts in Mathematics. Springer, New York, fourth edition, 1978.
- [30] Michael Luby. *Pseudorandomness and Cryptographic Applications*. Princeton University Press, Princeton, NJ, USA, 1996.
- [31] Michael Luby and Charles Rackoff. Pseudo-random permutation generators and cryptographic composition. In Juris Hartmanis, editor, *18th Annual ACM Symposium on Theory of Computing, STOC 1986*, pages 356–363. ACM, 1986.
- [32] G. A. Margulis. Explicit constructions of concentrators. *Probl. Peredachi Inf.*, 9:71–80, 1973.
- [33] Ueli M. Maurer and Stefano Tessaro. Computational indistinguishability amplification: Tight product theorems for system composition. In Shai Halevi, editor, *29th Annual International Conference Advances in Cryptology - CRYPTO 2009*, volume 5677 of *Lecture Notes in Computer Science*, pages 355–373. Springer, 2009.
- [34] Ueli M. Maurer and Stefano Tessaro. A hardcore lemma for computational indistinguishability: Security amplification for arbitrarily weak PRGs with optimal stretch. In Daniele Micciancio, editor, *7th Theory of Cryptography Conference - TCC 2010*, volume 5978 of *Lecture Notes in Computer Science*, pages 237–254. Springer, 2010.
- [35] Moni Naor and Omer Reingold. On the construction of pseudo-random permutations: Luby-Rackoff revisited. *Journal of Cryptology*, 12:29–66, 1997.
- [36] Rafael Pass and Muthuramakrishnan Venkatasubramanian. A parallel repetition theorem for constant-round Arthur-Merlin proofs. *ACM Trans. Comput. Theory*, 4:10:1–10:22, 2012.
- [37] Omer Reingold, Salil Vadhan, and Avi Wigderson. Entropy waves, the zig-zag graph product, and new constant-degree expanders. *Ann. of Math. (2)*, 155:157–187, 2002.
- [38] Herman Rubin and Jean E. Rubin. *Equivalents of the Axiom of Choice*. Studies in Logic and the Foundations of Mathematics. North-Holland, Amsterdam, 1970.
- [39] Eugene Seneta. *Non-negative Matrices and Markov Chains*. Springer Series in Statistics. Springer, New York, second edition, 1981.
- [40] Stefano Tessaro. Security amplification for the cascade of arbitrarily weak PRPs: Tight bounds via the interactive hardcore lemma. In Yuval Ishai, editor, *8th Theory of Cryptography Conference - TCC 2011*, volume 6597 of *Lecture Notes in Computer Science*, pages 37–54. Springer, 2011.

- [41] Salil P. Vadhan. Pseudorandomness. *Foundations and Trends in Theoretical Computer Science*, 7(13):1–336, 2012.
- [42] Andrew Chi-Chih Yao. Theory and applications of trapdoor functions. In *Proc. 23rd Annual IEEE Symp. on Foundations of Computer Science, FOCS 1982*, pages 80–91. IEEE Computer Society Press, 1982.