

# Average degrees of edge-chromatic critical graphs

Yan Cao<sup>a</sup>, Guantao Chen<sup>a</sup>, Suyun Jiang<sup>b\*</sup>, Huiqing Liu<sup>c\*</sup>, Fuliang Lu<sup>d\*</sup>

<sup>a</sup> Department of Mathematics and Statistics, Georgia State University, Atlanta, GA 30303

<sup>b</sup> School of Mathematics, Shandong University, Jinan, 250100

<sup>c</sup> Faculty of Mathematics and Statistics, Hubei University, Wuhan 430062

<sup>d</sup> School of Mathematics and Statistics, Linyi University, Linyi, Shandong 276000

## Abstract

Given a graph  $G$ , denote by  $\Delta$ ,  $\bar{d}$  and  $\chi'$  the maximum degree, the average degree and the chromatic index of  $G$ , respectively. A simple graph  $G$  is called *edge- $\Delta$ -critical* if  $\chi'(G) = \Delta + 1$  and  $\chi'(H) \leq \Delta$  for every proper subgraph  $H$  of  $G$ . Vizing in 1968 conjectured that if  $G$  is edge- $\Delta$ -critical, then  $\bar{d} \geq \Delta - 1 + \frac{3}{n}$ . We show that

$$\bar{d} \geq \begin{cases} 0.69241\Delta - 0.15658 & \text{if } \Delta \geq 66, \\ 0.69392\Delta - 0.20642 & \text{if } \Delta = 65, \text{ and} \\ 0.68706\Delta + 0.19815 & \text{if } 56 \leq \Delta \leq 64. \end{cases}$$

This result improves the best known bound  $\frac{2}{3}(\Delta + 2)$  obtained by Woodall in 2007 for  $\Delta \geq 56$ . Additionally, Woodall constructed an infinite family of graphs showing his result cannot be improved by well-known Vizing's Adjacency Lemma and other known edge-coloring techniques. To overcome the barrier, we follow the recently developed recoloring technique of Tashkinov trees to expand Vizing fans technique to a larger class of trees.

*Keywords:* edge- $k$ -coloring; edge-critical graphs; Vizing's Adjacency Lemma

## 1 Introduction

All graphs in this paper, unless otherwise stated, are simple graphs. Let  $G$  be a graph with vertex set  $V(G)$  and edge set  $E(G)$ . Denote by  $\Delta$  the maximum degree of  $G$ . An edge- $k$ -coloring of a graph  $G$  is a mapping  $\varphi : E(G) \rightarrow \{1, 2, \dots, k\}$  such that  $\varphi(e) \neq \varphi(f)$  for any two adjacent edges  $e$  and  $f$ . We call  $\{1, 2, \dots, k\}$  the color set of  $\varphi$ . Denote by

---

\*Partially supported by NSFC of China (Nos. 11671232, 11571096, 61373019, 11671186).

$\mathcal{C}^k(G)$  the set of all edge- $k$ -colorings of  $G$ . The chromatic index  $\chi'(G)$  is the least integer  $k \geq 0$  such that  $\mathcal{C}^k(G) \neq \emptyset$ . We call  $G$  *class one* if  $\chi'(G) = \Delta$ . Otherwise, Vizing' theorem [12] gives  $\chi'(G) = \Delta + 1$  and  $G$  is said to be of *class two*. An edge  $e$  is called *critical* if  $\chi'(G - e) < \chi'(G)$ , where  $G - e$  is the subgraph obtained from  $G$  by removing the edge  $e$ . A graph  $G$  is called *edge- $\Delta$ -critical* if  $\chi'(G) = \Delta + 1$  and  $\chi'(H) \leq \Delta$  holds for any proper subgraph  $H$  of  $G$ . Clearly, if  $G$  is edge- $\Delta$ -critical, then  $G$  is connected and  $\chi'(G - e) = \Delta$  for any  $e \in E(G)$ . Let  $\bar{d}(G)$  denote the average degree of a graph  $G$ . Vizing [14] made the following conjecture in 1968, which is commonly referred as Vizing's Average Degree Conjecture.

**Conjecture 1.** [Vizing [14]] *If  $G$  is an edge- $\Delta$ -critical graph of  $n$  vertices, then  $\bar{d}(G) \geq \Delta - 1 + \frac{3}{n}$ .*

The conjecture has been verified for  $\Delta \leq 6$ , see [3, 5, 6, 8]. For arbitrary  $\Delta$ , there are a few results on the lower bound for  $\bar{d}(G)$ . Let  $G$  be an edge- $\Delta$ -critical graph. Fiorini [2] showed, for  $\Delta \geq 2$ ,

$$\bar{d}(G) \geq \begin{cases} \frac{1}{2}(\Delta + 1) & \Delta \text{ is odd;} \\ \frac{1}{2}(\Delta + 2) & \Delta \text{ is even.} \end{cases}$$

Haile [4] improved the bounds as follows.

$$\bar{d}(G) \geq \begin{cases} \frac{3}{5}(\Delta + 2) & \Delta = 9, 11, 13; \\ \frac{\Delta+6}{2} - \frac{12}{\Delta+4} & \Delta \geq 10, \Delta \text{ is even;} \\ \frac{15+\sqrt{29}}{2} & \Delta = 15; \\ \frac{\Delta+7}{2} - \frac{16}{\Delta+5} & \Delta \geq 17, \Delta \text{ is odd.} \end{cases}$$

Sanders and Zhao [9] showed  $\bar{d}(G) \geq \frac{1}{2}(\Delta + \sqrt{2\Delta - 1})$  for  $\Delta \geq 2$ . Woodall [16] improved the bound to  $\bar{d}(G) \geq \frac{t(\Delta+t-1)}{2t-1}$ , where  $t = \lceil \sqrt{\Delta/2} \rceil$ . Improving Vizing's Adjacency Lemma, Woodall [15] improved the coefficient of  $\Delta$  from  $\frac{1}{2}$  to  $\frac{2}{3}$  as follows.

$$\bar{d}(G) \geq \begin{cases} \frac{2}{3}(\Delta + 1) & \text{if } \Delta \geq 2; \\ \frac{2}{3}\Delta + 1 & \text{if } \Delta \geq 8; \\ \frac{2}{3}(\Delta + 2) & \text{if } \Delta \geq 15. \end{cases}$$

In the same paper, Woodall provided the following example demonstrating that the above result cannot be improved by the use of his new adjacency Lemmas (see Lemma 2 and Lemma 3) and Vizing's Adjacency Lemma alone.

Let  $G$  be a graph comprising  $k$  vertices of degree 4, all of whose neighbors have degree  $\Delta$ , and  $2k$  vertices of degree  $\Delta$ , each of which is adjacent to two vertices of degree 4 and  $\Delta - 2$  vertices of degree  $\Delta$ . Graph  $G$  can be chosen to be triangle-free, and indeed to have arbitrarily large girth. Then  $G$  may not be edge- $\Delta$ -critical, but it satisfies the conclusions of all the existing lemmas at that time including two mentioned above, and it has average degree  $\frac{2}{3}(\Delta + 2)$ . So, using these known results, it is impossible to prove that the example is not edge- $\Delta$ -critical. On the other hand, we note that using our new result, Claim 3.4 in Section 3, it is readily seen that if  $\Delta \geq 6$  then the above example is not edge- $\Delta$ -critical. By proving a few stronger properties of edge- $\Delta$ -critical graphs, we improve Woodall's result as below for  $\Delta \geq 56$ .

**Theorem 1.** *If  $G$  is an edge- $\Delta$ -critical graph, then*

$$\bar{d}(G) \geq \begin{cases} 0.69241\Delta - 0.15658 & \text{if } \Delta \geq 66, \\ 0.69392\Delta - 0.20642 & \text{if } \Delta = 65, \text{ and} \\ 0.68706\Delta + 0.19815 & \text{if } 56 \leq \Delta \leq 64. \end{cases}$$

We will prove a few technic lemmas in Section 2 and give the proof of Theorem 1 in Section 3. We will use the following terminology and notation. Let  $G$  be a graph. Denote by  $N(x)$  the neighborhood of  $x$  for any  $x \in V(G)$ , and  $d(x)$  the degree of  $x$ , i.e.,  $d(x) = |N(x)|$ . For any nonnegative integer  $m$ , we call a vertex  $x$  an  $m$ -vertex if  $d(x) = m$ , a ( $< m$ )-vertex if  $d(x) < m$ , and ( $> m$ )-vertex if  $d(x) > m$ . Correspondingly, we call a neighbor  $y$  of  $x$  an  $m$ -neighbor if  $d(y) = m$ , etc.. Let  $k$  be a positive integer such that  $\mathcal{C}^k(G - e) \neq \emptyset$ , and let  $\varphi \in \mathcal{C}^k(G - e)$  and  $v \in V(G)$ . Let  $\varphi(v) = \{\varphi(e) : e \text{ is incident with } v\}$  and  $\bar{\varphi}(v) = \{1, \dots, k\} \setminus \varphi(v)$ . We call  $\varphi(v)$  the set of colors seen by  $v$  and  $\bar{\varphi}(v)$  the set of colors missing at  $v$ . A set  $X \subseteq V(G)$  is called *elementary* with respect to  $\varphi$  if  $\bar{\varphi}(u) \cap \bar{\varphi}(v) = \emptyset$  for every two distinct vertices  $u, v \in X$ . For any color  $\alpha$ , let  $E_\alpha$  denote the set of edges assigned color  $\alpha$ . Clearly,  $E_\alpha$  is matching of  $G$ . For any two colors  $\alpha$  and  $\beta$ , the components of induced by edges in  $E_\alpha \cup E_\beta$ , named  $(\alpha, \beta)$ -chains, are even cycles and paths with alternating color  $\alpha$  and  $\beta$ . For a vertex  $v$  of  $G$ , we denote by  $P_v(\alpha, \beta, \varphi)$  the unique  $(\alpha, \beta)$ -chain that contains the vertex  $v$ . Let  $\varphi/P_v(\alpha, \beta, \varphi)$  denote the edge- $k$ -coloring obtain from  $\varphi$  by switching colors  $\alpha$  and  $\beta$  on the edges on  $P_v(\alpha, \beta, \varphi)$ .

## 2 Lemmas

Let  $q$  be a positive number,  $G$  be an edge- $\Delta$ -critical graph and  $x \in V(G)$ . For each  $y \in N(x)$ , let  $\sigma_q(x, y) = |\{z \in N(y) \setminus \{x\} : d(z) \geq q\}|$ , the number of neighbors of

$y$  (except  $x$ ) with degree at least  $q$ . Vizing studied the case  $q = \Delta$  and obtained the following result.

**Lemma 1.** [Vizing's Adjacency Lemma [13]] *Let  $G$  be an edge- $\Delta$ -critical graph. Then  $\sigma_\Delta(x, y) \geq \Delta - d(x) + 1$  holds for every  $xy \in E(G)$ .*

Woodall [15] studied  $\sigma_q(x, y)$  for the case  $q = 2\Delta - d(x) - d(y) + 2$  and obtained the following two results. For convention, we let  $\sigma(x, y) = \sigma_q(x, y)$  when  $q = 2\Delta - d(x) - d(y) + 2$ .

**Lemma 2.** [Woodall [15]] *Let  $xy$  be an edge in an edge- $\Delta$ -critical graph  $G$ . Then there are at least  $\Delta - \sigma(x, y) \geq \Delta - d(y) + 1$  vertices  $z \in N(x) \setminus \{y\}$  such that  $\sigma(x, z) \geq 2\Delta - d(x) - \sigma(x, y)$ .*

Let  $x$  be a vertex in a graph  $G$  and  $y \in N(x)$ . Vizing's Adjacency Lemma shows that  $\sigma_\Delta(x, y) \geq \Delta - d(x) + 1$ . So,  $\sigma(x, y) \geq \Delta - d(x) + 1$ . Woodall studied their difference through the following two parameters.

$$p_{min}(x) := \min_{y \in N(x)} \sigma(x, y) - \Delta + d(x) - 1 \quad \text{and}$$

$$p(x) := \min\left\{ p_{min}(x), \left\lfloor \frac{d(x)}{2} \right\rfloor - 1 \right\}.$$

Clearly,  $p(x) < d(x)/2 - 1$ . As a corollary, the following lemma shows that there are about  $d(x)/2$  neighbors  $y$  of  $x$  such that  $\sigma(x, y) \geq \Delta/2$ . In general, for any positive number  $q$  with  $q \leq \Delta$ , we define the following two parameters.

$$p_{min}(x, q) := \min_{y \in N(x)} \sigma_q(x, y) - \Delta + d(x) - 1 \quad \text{and}$$

$$p(x, q) := \min\left\{ p_{min}(x, q), \left\lfloor \frac{d(x)}{2} \right\rfloor - 3 \right\}.$$

**Lemma 3.** [Woodall [15]] *Every vertex  $x$  in an edge- $\Delta$ -critical graph has at least  $d(x) - p(x) - 1$  neighbors  $y$  for which  $\sigma(x, y) \geq \Delta - p(x) - 1$ .*

When  $d(x) \leq \Delta - 4$ , we generalize the above results by allowing  $q$  taking various values and obtain the following two results, which serve as key ingredients in our proof of Theorem 1.

**Lemma 4.** *Let  $xy$  be an edge in an edge- $\Delta$ -critical graph  $G$  and  $q$  be a positive number. If  $\Delta/2 < q \leq \Delta - d(x)/2 - 2$ , then  $x$  has at least  $\Delta - \sigma_q(x, y) - 2$  vertices  $z \in N(x) \setminus \{y\}$  such that  $\sigma_q(x, z) \geq 2\Delta - d(x) - \sigma_q(x, y) - 4$ .*

Due to its length, the proof of Lemma 4 will be placed at the end of this section. The following is a consequence of it.

**Lemma 5.** *Let  $G$  be an edge- $\Delta$ -critical graph,  $x \in V(G)$  and  $q$  be a positive number. If  $\Delta/2 < q \leq \Delta - d(x)/2 - 2$ , then  $x$  has at least  $d(x) - p(x, q) - 3$  neighbors  $y$  for which  $\sigma_q(x, y) \geq \Delta - p(x, q) - 5$ .*

*Proof.* Let  $y \in N(x)$  such that  $p_{min}(x, q) = \sigma_q(x, y) - \Delta + d(x) - 1$ .

If  $p(x, q) = p_{min}(x, q)$ , by Lemma 4,  $x$  has at least  $\Delta - \sigma_q(x, y) - 2 = \Delta - (\Delta - d(x) + p_{min}(x, q) + 1) - 2 = d(x) - p_{min}(x, q) - 3$  vertices  $z \in N(x) \setminus \{y\}$  such that  $\sigma_q(x, z) \geq 2\Delta - d(x) - 4 - \sigma_q(x, y) = \Delta - p_{min}(x, q) - 5$ .

If  $p(x, q) = \left\lfloor \frac{d(x)}{2} \right\rfloor - 3 < p_{min}(x, q)$ , then for every  $y \in N(x)$ ,  $\sigma_q(x, y) > \Delta - d(x) + 1 + \left\lfloor \frac{d(x)}{2} \right\rfloor - 3 \geq \Delta - \left\lfloor \frac{d(x)}{2} \right\rfloor - 3 = \Delta - p(x, q) - 6$ . So  $\sigma_q(x, y) \geq \Delta - p(x, q) - 5$ .  $\square$

Let  $G$  be a graph (in this paragraph,  $G$  may be a multigraph),  $e_1 = y_0y_1 \in E(G)$  and  $\varphi \in C^k(G - e_1)$ . A Tashkinov tree  $T$  is a sequence  $T = (y_0, e_1, y_1, \dots, e_p, y_p)$  with  $p \geq 1$  consisting of edges  $e_1, e_2, \dots, e_p$  and vertices  $y_0, y_1, \dots, y_p$  such that the following two conditions hold.

- The edges  $e_1, e_2, \dots, e_p$  are distinct and  $e_i = y_r y_i$  for each  $1 \leq i \leq p$ , where  $r < i$ ;
- For every edge  $e_i$  with  $2 \leq i \leq p$ , there is a vertex  $y_h$  with  $0 \leq h < i$  such that  $\varphi(e_i) \in \bar{\varphi}(y_h)$ .

Clearly, a Tashkinov tree is indeed a tree of  $G$ . Tashkinov [11] proved that if  $G$  is edge- $k$ -critical with  $k \geq \Delta + 1$ , then  $V(T)$  is elementary. In the above definition, if  $e_i = y_0 y_i$  for every  $i$  (i.e.,  $T$  is a star with  $y_0$  as the center), then  $T$  is a Vizing fan. The classic result of Vizing [10] show that for every Vizing fan  $T$  the set  $V(T)$  is elementary if  $G$  is edge- $k$ -critical for every  $k \geq \Delta$ , which includes edge- $\Delta$ -critical graphs. In the definition of Tashkinov tree, if  $e_i = y_{i-1} y_i$  for every  $i$  (i.e.  $T$  is a path with end-vertices  $y_0$  and  $y_p$ ), then  $T$  is a Kierstead path, which was introduced by Kierstead [7]. Kierstead proved that for every Kierstead path  $P$  the set  $V(P)$  is elementary if  $G$  is an edge- $k$ -critical with  $k \geq \Delta + 1$ . For simple graphs, following Kierstead's proof, Zhang [17] noticed that for a Kierstead path  $P$  the set  $V(P)$  is elementary if  $G$  is edge- $\Delta$ -critical and  $d(y_i) < \Delta$  for every  $i$  with  $2 \leq i \leq p$ . Clearly, every Kierstead path  $P$  with three vertices is a Vizing fan, so  $V(P)$  is elementary if  $G$  is edge- $\Delta$ -critical.

**Lemma 6.** [Kostochka and Stiebitz [10]] *Let  $G$  be a graph with maximum degree  $\Delta$  and  $\chi'(G) = \Delta + 1$ . Let  $e_1 \in E(G)$  be a critical edge and  $\varphi \in \mathcal{C}^\Delta(G - e_1)$ . If  $K = (y_0, e_1, y_1, e_2, y_2, e_3, y_3)$  is a Kierstead path with respect to  $e_1$  and  $\varphi$ , then the following statements hold:*

1.  $\bar{\varphi}(y_0) \cap \bar{\varphi}(y_1) = \emptyset$ ;
2. if  $d(y_2) < \Delta$ , then  $V(K)$  is elementary with respect to  $\varphi$ ;
3. if  $d(y_1) < \Delta$ , then  $V(K)$  is elementary with respect to  $\varphi$ ;
4. if  $\Gamma = \bar{\varphi}(y_0) \cup \bar{\varphi}(y_1)$ , then  $|\bar{\varphi}(y_3) \cap \Gamma| \leq 1$ .

In the definition of Tashkinov tree  $T = (y_0, e_1, y_1, e_2, y_2, \dots, y_p)$ , we call  $T$  a *broom* if  $e_2 = y_1y_2$  and for each  $i \geq 3$ ,  $e_i = y_2y_i$ , i.e.,  $y_2$  is one of the end-vertices of  $e_i$  for each  $i \geq 3$ . Moreover, we call  $T$  a *simple broom* if  $\varphi(e_i) \in \bar{\varphi}(y_0) \cup \bar{\varphi}(y_1)$  for each  $i \geq 3$ , i.e.,  $(y_0, e_1, y_1, e_2, y_2, e_i, y_i)$  is a Kierstead path.

**Lemma 7.** [Chen, Chen, Zhao [1]] *Let  $G$  be an edge- $\Delta$ -critical graph,  $e_1 = y_0y_1 \in E(G)$  and  $\varphi \in \mathcal{C}^\Delta(G - e_1)$  and  $B = \{y_0, e_1, y_1, e_2, y_2, \dots, e_p, y_p\}$  be a simple broom. If  $|\bar{\varphi}(y_0) \cup \bar{\varphi}(y_1)| \geq 4$  and  $\min\{d(y_1), d(y_2)\} < \Delta$ , then  $V(B)$  is elementary with respect to  $\varphi$ .*

**Lemma 8.** *Let  $G$  be an edge- $\Delta$ -critical graph,  $xy \in E(G)$ , and  $\varphi \in \mathcal{C}^\Delta(G - xy)$ . Let  $q$  be a positive number with  $d(x) < q \leq \Delta - 1$  and  $Z = \{z \in N(x) \setminus \{y\} : d(z) > q, \varphi(xz) \in \bar{\varphi}(y)\}$ . Then for every  $z \in Z$  the following three inequalities hold.*

$$|Z| \geq \Delta - d(y) + 1 - \left\lfloor \frac{d(x) + d(y) - \Delta - 2}{\Delta - q} \right\rfloor \quad (1)$$

$$\sum_{z \in Z} (d(z) - q) \geq (\Delta - d(y) + 1)(\Delta - q) - d(x) - d(y) + \Delta + 2 \quad (2)$$

$$\sigma_q(x, z) \geq 2\Delta - d(x) - d(y) + 1 - \left\lfloor \frac{d(x) + d(y) + d(z) - 2\Delta - 2}{\Delta - q} \right\rfloor \quad (3)$$

*Proof.* Since  $xy$  is a critical edge of  $G$ ,  $\bar{\varphi}(x) \cap \bar{\varphi}(y) = \emptyset$ . Let  $Z_y := \{z \in N(x) \setminus \{y\} : \varphi(xz) \in \bar{\varphi}(y)\}$ . Clearly,  $Z \subseteq Z_y$  and  $|Z_y| = \Delta - d(y) + 1$ . Since  $\{y, x\} \cup Z_y$  forms a Vizing fan with center  $x$ , it is elementary, so  $|\bar{\varphi}(x)| + |\bar{\varphi}(y)| + \sum_{z \in Z_y} |\bar{\varphi}(z)| \leq \Delta$  holds. Since  $|\bar{\varphi}(x)| = \Delta - d(x) + 1$  and  $|\bar{\varphi}(y)| = \Delta - d(y) + 1$ , we have

$$\sum_{z \in Z_y} |\bar{\varphi}(z)| \leq \Delta - |\bar{\varphi}(x)| - |\bar{\varphi}(y)| \leq d(x) + d(y) - \Delta - 2. \quad (4)$$

Since  $d(z) \leq q$  for all  $z \in Z_y - Z$ ,  $\sum_{z \in Z_y} |\bar{\varphi}(z)| \geq (|Z_y| - |Z|)(\Delta - q)$ . Solving for  $|Z|$ , we get  $|Z| \geq |Z_y| - \left\lfloor \frac{d(x)+d(y)-\Delta-2}{\Delta-q} \right\rfloor$ . Since  $|Z_y| = \Delta - d(y) + 1$ , inequality (1) holds.

Plugging  $|\bar{\varphi}(z)| = \Delta - d(z)$  for each  $z \in Z_y$  in inequality (4), we get

$$\sum_{z \in Z_y} d(z) \geq |Z_y| \Delta - (d(x) + d(y) - \Delta - 2).$$

Since  $d(z) \leq q$  for every  $z \in Z_y - Z$ , we have

$$\sum_{z \in Z} (d(z) - q) \geq \sum_{z \in Z_y} (d(z) - q) \geq |Z_y| \Delta - (d(x) + d(y) - \Delta - 2) - |Z_y| q.$$

Plugging  $|Z_y| = \Delta - d(y) + 1$ , we get (2).

For each  $z \in Z$ , let  $U_z^* = \{u \in N(z) \setminus \{x\} : \varphi(zu) \in \bar{\varphi}(x) \cup \bar{\varphi}(y) \setminus \{\varphi(xz)\}\}$  and  $U_z = \{u \in U_z^* : d(u) > q\}$ . Clearly,  $|U_z^*| = 2\Delta - d(x) - d(y) + 1$  and  $\{y, x, z\} \cup U_z^*$  forms a simple broom. Since  $d(x) < q \leq \Delta - 1$ , we have  $d(x) \leq \Delta - 2$ . Thus  $|\bar{\varphi}(x) \cup \bar{\varphi}(y)| \geq 4$  and  $\min\{d(x), d(z)\} = d(x) < \Delta$ . By Lemma 7,  $\{y, x, z\} \cup U_z^*$  is elementary with respect to  $\varphi$ . So

$$\sum_{u \in U_z^*} |\bar{\varphi}(u)| + |\bar{\varphi}(x)| + |\bar{\varphi}(y)| + |\bar{\varphi}(z)| \leq \Delta,$$

which in turn gives  $\sum_{u \in U_z^*} |\bar{\varphi}(u)| \leq d(x) + d(y) + d(z) - 2\Delta - 2$ . Since  $d(u) \leq q$  for every  $u \in U_z^* - U_z$ ,  $\sum_{u \in U_z^*} |\bar{\varphi}(u)| \geq (|U_z^*| - |U_z|)(\Delta - q)$ . So,

$$(|U_z^*| - |U_z|)(\Delta - q) \leq d(x) + d(y) + d(z) - 2\Delta - 2.$$

Solving the above inequality with  $|U_z^*| = 2\Delta - d(x) - d(y) + 1$ , we get

$$|U_z| \geq 2\Delta - d(x) - d(y) + 1 - \left\lfloor \frac{d(x) + d(y) + d(z) - 2\Delta - 2}{\Delta - q} \right\rfloor.$$

Since  $\sigma_q(x, z) \geq |U_z|$ , the inequality (3) holds.  $\square$

## 2.1 Proof of Lemma 4

**Lemma 4.** Let  $xy$  be an edge in an edge- $\Delta$ -critical graph  $G$  and  $q$  be a positive number. If  $\Delta/2 < q \leq \Delta - d(x)/2 - 2$ , then  $x$  has at least  $\Delta - \sigma_q(x, y) - 2$  vertices  $z \in N(x) \setminus \{y\}$  such that  $\sigma_q(x, z) \geq 2\Delta - d(x) - \sigma_q(x, y) - 4$ .

*Proof.* Let graph  $G$ , edge  $xy \in E(G)$  and  $q$  be defined as in Lemma 4. A neighbor  $z \in N(x) \setminus \{y\}$  is called *feasible* if there exists a coloring  $\varphi \in \mathcal{C}^\Delta(G - xy)$  such that  $\varphi(xz) \in \bar{\varphi}(y)$ , and such a coloring  $\varphi$  is called  *$z$ -feasible*. Denote by  $\mathcal{C}_z$  the set of all  $z$ -feasible colorings. For each  $\varphi \in \mathcal{C}_z$ . let

$$\begin{aligned} Z(\varphi) &= \{v \in N(z) \setminus \{x\} : \varphi(vz) \in \bar{\varphi}(x) \cup \bar{\varphi}(y)\}, \\ C_z(\varphi) &= \{\varphi(vz) : v \in Z(\varphi) \text{ and } d(v) < q\}, \\ Y(\varphi) &= \{v \in N(y) \setminus \{x\} : \varphi(vy) \in \bar{\varphi}(x) \cup \bar{\varphi}(z)\}, \text{ and} \\ C_y(\varphi) &= \{\varphi(vy) : v \in Y(\varphi) \text{ and } d(v) < q\}. \end{aligned}$$

Note that  $Z(\varphi)$  and  $Y(\varphi)$  are vertex sets while  $C_z(\varphi)$  and  $C_y(\varphi)$  are color sets. For each color  $k \in \varphi(z)$ , let  $z_k \in N(z)$  such that  $\varphi(zz_k) = k$ . Similarly, we define  $y_k$  for each  $k \in \varphi(y)$ . Let  $T(\varphi) = \{k \in \varphi(x) \cap \varphi(y) \cap \varphi(z) : d(y_k) < q \text{ and } d(z_k) < q\}$ .

Since  $G$  is edge- $\Delta$ -critical,  $\{x, y, z\}$  is elementary with respect to  $\varphi$ . So  $\bar{\varphi}(x)$ ,  $\bar{\varphi}(y)$ ,  $\bar{\varphi}(z)$  and  $\varphi(x) \cap \varphi(y) \cap \varphi(z)$  are mutually exclusive and

$$\bar{\varphi}(x) \cup \bar{\varphi}(y) \cup \bar{\varphi}(z) \cup (\varphi(x) \cap \varphi(y) \cap \varphi(z)) = \{1, 2, \dots, \Delta\}.$$

Recall that  $\sigma_q(x, y)$  and  $\sigma_q(x, z)$  are number of vertices with degree  $\geq q$  in  $N(y) \setminus \{x\}$  and  $N(z) \setminus \{x\}$ , respectively. So, the following inequalities hold.

$$\begin{aligned} &\sigma_q(x, y) + \sigma_q(x, z) \\ &\geq |Y(\varphi)| - |C_y(\varphi)| + |Z(\varphi)| - |C_z(\varphi)| + |\varphi(x) \cap \varphi(y) \cap \varphi(z)| - |T(\varphi)| \\ &= |\bar{\varphi}(x) \cup \bar{\varphi}(z)| + |\bar{\varphi}(x) \cup \bar{\varphi}(y)| - 1 + |\varphi(x) \cap \varphi(y) \cap \varphi(z)| - |C_y(\varphi)| - |C_z(\varphi)| - |T(\varphi)| \\ &= \Delta + |\bar{\varphi}(x)| - |C_y(\varphi)| - |C_z(\varphi)| - |T(\varphi)| - 1 \\ &= 2\Delta - d(x) + 1 - |C_y(\varphi)| - |C_z(\varphi)| - |T(\varphi)| - 1 \end{aligned}$$

So, Lemma 4 follows the three statements below.

- I.** For any  $\varphi \in \mathcal{C}_z$ ,  $|C_z(\varphi)| \leq 1$  and  $|C_y(\varphi)| \leq 1$ ;
- II.** there exists a  $\varphi \in \mathcal{C}_z$  such that  $|T(\varphi)| \leq 2$ ; and
- III.** there are  $\Delta - \sigma_q(x, y) - 2$  feasible vertices  $z \in N(x) \setminus \{y\}$ .

For every  $z$ -feasible coloring  $\varphi \in \mathcal{C}^\Delta(G - xy)$ , let  $\varphi^d \in \mathcal{C}^\Delta(G - xz)$  obtained from  $\varphi$  by assigning  $\varphi^d(xy) = \varphi(xz)$  and keeping all colors on other edges unchange. Clearly,  $\varphi^d$  is a  $y$ -feasible coloring and  $Z(\varphi^d) = Z(\varphi)$ ,  $Y(\varphi^d) = Y(\varphi)$ ,  $C_z(\varphi^d) = C_z(\varphi)$  and



$C_y(\varphi^d) = C_y(\varphi)$ . We call  $\varphi^d$  the *dual coloring* of  $\varphi$ . Considering dual colorings, we see that some properties for vertex  $z$  also hold for vertex  $y$ .

The condition  $q \leq \Delta - d(x)/2 - 2$  implies  $2(\Delta - q) + (\Delta - d(x)) + 1 > \Delta$ . So, for any  $\varphi \in \mathcal{C}^\Delta(G - xy)$ , every elementary set  $X$  with  $x \in X$  contains at most one vertex with degree  $\leq q$ .

Let  $z \in N(x) \setminus \{y\}$  be a feasible vertex and  $\varphi \in \mathcal{C}_z$ . By the definition of  $Z(\varphi)$ ,  $G[\{x, y, z\} \cup Z(\varphi)]$  contains a simple broom, so  $\{x, y, z\} \cup Z(\varphi)$  is elementary with respect to  $\varphi$ . Consequently, it contains at most one vertex other than  $x$  having degree  $< q$ . Thus,  $|C_z(\varphi)| \leq 1$ . By considering its dual  $\varphi^d$ , we have  $|C_y(\varphi)| = |C_y(\varphi^d)| \leq 1$ . Hence, **I** holds. The proofs of **II** and **III** are much more complicated. In the remainder of the proof, we let  $Z = Z(\varphi)$ ,  $Y = Y(\varphi)$ ,  $C_z = C_z(\varphi)$ ,  $C_y = C_y(\varphi)$ , and  $T = T(\varphi)$  if the coloring  $\varphi$  is clearly referred. Let  $R = C_z \cup C_y$  and  $\bar{\varphi}(x, R) = \bar{\varphi}(x) \setminus R$ . A coloring  $\varphi \in \mathcal{C}_z$  is called *optimal* if  $|C_z| + |C_y|$  is maximum over all feasible colorings.

### 2.1.1 Proof of II.

Suppose to the contrary:  $|T| \geq 3$  for every  $\varphi \in \mathcal{C}_z$ . Let  $\varphi$  be an optimal feasible coloring and assume, without loss of generality,  $\varphi(xz) = 1$ .

**Claim A.** For each  $i \in \bar{\varphi}(x, R)$  and  $k \in T$ ,  $P_x(i, k, \varphi)$  contains both  $y$  and  $z$ .

*Proof.* We first show that  $z \in V(P_x(i, k, \varphi))$ . Otherwise,  $P_z(i, k, \varphi)$  is disjoint with  $P_x(i, k, \varphi)$ . Let  $\varphi' = \varphi/P_z(i, k, \varphi)$ . Since  $1 \neq i, k$ ,  $\varphi'$  is also feasible. Since colors in  $R$  are unchanged and  $d(z_k) < q$ ,  $C_z(\varphi') = C_z \cup \{i\}$  and  $C_y(\varphi') \supseteq C_y$ , giving a contradiction to the maximality of  $|C_y| + |C_z|$ . By considering the dual  $\varphi^d$ , we can verify that  $y \in V(P_x(i, k, \varphi))$ .  $\square$

Since  $|T| \geq 3$ , there are three colors  $k_1, k_2, k_3 \in T$ . Let

$$\begin{aligned} V_T &= \{z_{k_1}, z_{k_2}, z_{k_3}\} \cup \{y_{k_1}, y_{k_2}, y_{k_3}\}, \\ W(\varphi) &= \{u \in V_T : \bar{\varphi}(u) \cap \bar{\varphi}(x) \subseteq R\}, \\ M(\varphi) &= V_T - W(\varphi) = \{u \in V_T : \bar{\varphi}(u) \cap \bar{\varphi}(x, R) \neq \emptyset\}, \\ E_T &= \{zz_{k_1}, zz_{k_2}, zz_{k_3}, yy_{k_1}, yy_{k_2}, yy_{k_3}\}, \\ E_W(\varphi) &= \{e \in E_T : e \text{ is incident to a vertex in } W(\varphi)\}, \text{ and} \\ E_M(\varphi) &= E_T - E_W(\varphi) = \{e \in E_T : e \text{ is incident to a vertex in } M(\varphi)\}. \end{aligned}$$

For convenience, we let  $W = W(\varphi)$ ,  $M = M(\varphi)$ ,  $E_W = E_W(\varphi)$  and  $E_M = E_M(\varphi)$ .

We assume that  $|E_W|$  is minimum over all optimal feasible coloring  $\varphi$  and all sets of three colors in  $T(\varphi)$ . For each  $v \in M$ , pick a color  $\alpha_v \in \bar{\varphi}(v) \cap \bar{\varphi}(x, R)$ . Let  $C_M = \{\alpha_v : v \in M\}$ . Clearly,  $|C_M| \leq |M|$ . Note that  $\{z_{k_1}, z_{k_2}, z_{k_3}\} \cap \{y_{k_1}, y_{k_2}, y_{k_3}\}$  may be not empty,  $\frac{|E_W|}{2} \leq |W| \leq |E_W|$  and  $\frac{|E_M|}{2} \leq |M| \leq |E_M|$ .

**Claim B.** If there exist two vertices  $u, v \in V_T$  and a color  $\alpha \in \varphi(x) \setminus R$  such that  $\alpha \in \bar{\varphi}(u) \cap \bar{\varphi}(v)$ , then there is an optimal feasible coloring  $\varphi^*$  such that  $|E_W(\varphi^*)| \leq |E_W|$  and  $\{u, v\} \cap M(\varphi^*) \neq \emptyset$ . Moreover, if  $\bar{\varphi}(x) \setminus (R \cup C_M) \neq \emptyset$ , then  $u$  or  $v \in M$ .

*Proof.* We first note that the condition of  $d(x)$  and  $q$  gives

$$|\bar{\varphi}(x)| = \Delta - d(x) + 1 \geq \Delta - 2(\Delta - q) + 5 > 5. \quad (5)$$

If  $\{u, v\} \cap M \neq \emptyset$ , we are done. Suppose  $u, v \in W$ . Let  $\beta$  be an arbitrary color in  $\bar{\varphi}(x, R)$  with the preference that  $\beta \in \bar{\varphi}(x, R) \setminus C_M$  if the set is not empty. Since  $|R| \leq 2$  and (5), such a color  $\beta$  exists. Since  $u, v \in W$ , we have  $\beta \in \varphi(u) \cap \varphi(v)$ . So, both  $u$  and  $v$  are endvertices of  $(\alpha, \beta)$ -chains. Assume without loss of generality  $P_u(\alpha, \beta, \varphi)$  is disjoint with  $P_x(\alpha, \beta, \varphi)$ . We note that  $\beta \in \varphi(y) \cap \varphi(z)$  since  $\{x, y, z\}$  is an elementary set.

We first consider the case of  $\alpha = 1$ . In this case,  $P_x(\alpha, \beta, \varphi) = P_y(\alpha, \beta, \varphi)$  holds; otherwise,  $\varphi/P_x(\alpha, \beta, \varphi)$  would lead a  $\Delta$ -coloring of  $G$ . Since  $\varphi(xz) = 1$ ,  $z \in P_x(\alpha, \beta, \varphi)$ . So,  $P_u(\alpha, \beta, \varphi) \cap \{x, y, z\} = \emptyset$ . Hence, coloring  $\varphi' = \varphi/P_u(\alpha, \beta, \varphi)$  is feasible,  $C_y(\varphi') = C_y$ ,  $C_z(\varphi') = C_z$  and  $T(\varphi') = T$ . So,  $\varphi'$  is also optimal,  $u \in M(\varphi')$  and  $|E_W(\varphi')| \leq |E_W|$  with that the inequality holds if the other endvertex of  $P_u(\alpha, \beta, \varphi)$  is not in  $M$  or  $\beta \notin R \cup C_M$ .

We now suppose  $\alpha \in \varphi(x) \setminus (R \cup \{1\})$ . So, both  $\alpha$  and  $\beta$  are not in  $R \cup \{1\}$ . Let  $\varphi' = \varphi/P_u(\alpha, \beta, \varphi)$ . Then,  $\varphi'$  is feasible (for  $z$ ),  $C_y(\varphi') = C_y$  and  $C_z(\varphi') = C_z$ . Thus,  $\varphi'$  is still an optimal coloring and  $\beta \in \bar{\varphi}'(u)$ . We have  $|E_W(\varphi')| \leq |E_W|$  and  $u \in M(\varphi')$ . By the minimality of  $|E_W|$ , we have the other endvertex of  $P_u(\alpha, \beta, \varphi)$  must be in  $M$  and  $\beta \in C_M$ , which leads a contradiction to the minimality of  $|E_W|$  if  $\beta \notin R \cup C_M$ .  $\square$

**Claim C.** There exist a color  $k \in \{k_1, k_2, k_3\}$  and three distinct colors  $i, j, \ell$  where  $i, j \in \bar{\varphi}(x, R)$  and  $\ell \in \bar{\varphi}(x, R) \cup \{1\}$  such that  $i \in \bar{\varphi}(z_k)$ ,  $j \in \bar{\varphi}(y_k)$  and  $\ell \in \bar{\varphi}(z_k) \cup \bar{\varphi}(y_k)$ .

*Proof.* We first note that if there exist  $i, j \in \bar{\varphi}(x, R)$  such that  $i \in \bar{\varphi}(z_k)$  and  $j \in \bar{\varphi}(y_k)$ , then  $i \neq j$ ; for otherwise, by Claim A, the path  $P_x(i, k, \varphi)$  contains three endvertices  $x, z_k$  and  $y_k$ , a contradiction.

First we show that there exist  $i, j \in \bar{\varphi}(x, R)$  and  $k \in \{k_1, k_2, k_3\}$  such that  $i \in \bar{\varphi}(z_k)$  and  $j \in \bar{\varphi}(y_k)$ . Suppose not. Then  $|E_M| \leq 3$  and  $|E_W| \geq 3$ , which in turn give  $|W| \geq \lceil \frac{3}{2} \rceil = 2$  and  $|M| \leq |E_M| \leq 3$ . Let  $u, v \in W$ . By (5),  $|\bar{\varphi}(x)| \geq 6 > |R| + |M|$ . There exists a color  $\beta \in \bar{\varphi}(x, R) \setminus C_M$ . Then,  $\beta \in \varphi(u) \cap \varphi(v)$  as  $u, v \in W$ .

Since  $|R| \leq 2$ , we have

$$|\bar{\varphi}(u) \setminus R| + |\bar{\varphi}(v) \setminus R| + |\bar{\varphi}(x)| > 2(\Delta - q - 2) + \Delta - d(x) + 1 \geq \Delta + 1.$$

So, there is a color  $\alpha$  shared by at least two of these three sets. Since  $(\bar{\varphi}(u) \setminus R) \cap \bar{\varphi}(x) = \emptyset$  and  $(\bar{\varphi}(v) \setminus R) \cap \bar{\varphi}(x) = \emptyset$ , we have  $\alpha \in (\bar{\varphi}(u) \setminus R) \cap (\bar{\varphi}(v) \setminus R) \cap \varphi(x)$ . By Claim B, there exists an optimal feasible coloring  $\varphi'$  such that  $|E_W(\varphi')| \leq |E_W|$ . Moreover, since  $\beta \notin R \cup C_M$ , the inequality holds which gives a contradiction to the minimality of  $|E_W|$ .

We now only need to show that additionally there exists another color  $\ell \in \bar{\varphi}(x, R) \cup \{1\}$  such that  $\ell \in \bar{\varphi}(y_k) \cup \bar{\varphi}(z_k)$ . Suppose on the contrary that there is no such a color  $\ell$ . Then the following equalities hold.

$$\bar{\varphi}(z_k) \cap \bar{\varphi}(x, R) = \{i\} \quad \text{and} \quad \bar{\varphi}(y_k) \cap \bar{\varphi}(x, R) = \{j\}$$

Moreover,  $1 \notin \bar{\varphi}(z_k) \cup \bar{\varphi}(y_k)$ . Since  $|R| \leq 2$ , the following inequalities hold.

$$|\bar{\varphi}(z_k) \setminus (R \cup \{i\})| + |\bar{\varphi}(y_k) \setminus (R \cup \{j\})| + |\bar{\varphi}(x) \cup (R \cup \{1\})| > 2(\Delta - q - |R| - 1) + \Delta - d(x) + 2 \geq \Delta$$

So, there is color  $\alpha$  in two of the three sets. Since  $(\bar{\varphi}(z_k) \setminus (R \cup \{i\})) \cap (\bar{\varphi}(x) \cup R \cup \{1\}) = \emptyset$  and  $(\bar{\varphi}(y_k) \setminus (R \cup \{j\})) \cap (\bar{\varphi}(x) \cup R \cup \{1\}) = \emptyset$ ,  $\alpha \in \bar{\varphi}(z_k) \cap \bar{\varphi}(y_k) \cap \varphi(x) \setminus (R \cup \{i, j, 1\})$ .

Since  $|\bar{\varphi}(x)| = \Delta - d(x) + 1 \geq \Delta - 2(\Delta - q) + 5 > 5$ , there exists a color  $\beta \in \bar{\varphi}(x) \setminus (R \cup \{i, j\})$ . Then,  $\beta \notin \bar{\varphi}(z_k) \cup \bar{\varphi}(y_k)$ . So,  $\beta \in (\bar{\varphi}(x) \cap \varphi(z_k) \cap \varphi(y_k)) \setminus (R \cup \{i, j\})$ .

Applying Claim B with color  $\alpha$ , we obtain an optimal coloring  $\varphi'$  and  $|E_W(\varphi')| \leq |E_W|$ , but color  $\beta$  serves as the required color  $\ell$ , giving a contradiction.  $\square$

Let  $k, i, j$  and  $\ell$  be as stated in Claim C. If  $\ell \neq 1$ , we consider coloring obtained from  $\varphi$  by interchange colors 1 and  $\ell$  for edges not on the path  $P_x(1, \ell, \varphi)$ , and rename it as  $\varphi$ . So we may assume  $1 \in \bar{\varphi}(y_k) \cup \bar{\varphi}(z_k)$ .

We first consider the case of  $1 \in \bar{\varphi}(y_k)$ . By Claim A, the paths  $P_x(i, k, \varphi)$  and  $P_x(j, k, \varphi)$  both contain  $y, z$ . Since  $\varphi(y y_k) = \varphi(z z_k) = k$ , these two paths also contain  $y_k, z_k$ . Since  $i \in \bar{\varphi}(z_k)$ , we have  $x$  and  $z_k$  are the two endvertices of  $P_x(i, k, \varphi)$ . So,  $i \in \varphi(y) \cap \varphi(z) \cap \varphi(y_k)$ . Similarly, we have  $j \in \varphi(y) \cap \varphi(z) \cap \varphi(z_k)$ . We now consider the following sequence of colorings of  $G - xy$ .

Let  $\varphi_1$  be obtained from  $\varphi$  by assigning  $\varphi_1(y y_k) = 1$ . Since 1 is missing at both  $y$  and  $y_k$ ,  $\varphi_1$  is an edge- $\Delta$ -coloring of  $G - xy$ . Now  $k$  is missing at  $y$  and  $y_k$ ,  $i$  is still missing at  $z_k$ . Since  $G$  is not  $\Delta$ -colorable,  $P_x(i, k, \varphi) = P_y(i, k, \varphi)$ ; otherwise  $\varphi/P_y(i, k, \varphi)$  can be extended to an edge- $\Delta$ -coloring of  $G$  giving a contradiction. Furthermore,  $z_k, y_k \notin V(P_x(i, k, \varphi_1))$  since either  $i$  or  $k$  is missing at these two vertices, which in turn shows that  $z \notin V(P_x(i, k, \varphi_1))$  since  $\varphi_1(z z_k) = k$ .

Let  $\varphi_2 = \varphi_1/P_x(i, k, \varphi_1)$ . We have  $k \in \bar{\varphi}_2(x)$ ,  $i \in \bar{\varphi}_2(y) \cap \bar{\varphi}_2(z_k)$  and  $j \in \bar{\varphi}_2(x) \cap \bar{\varphi}_2(y_k)$ . Since  $G$  is not edge- $\Delta$ -colorable,  $P_x(i, j, \varphi_2) = P_y(i, j, \varphi_2)$  which contains neither  $y_k$  nor  $z_k$ .

Let  $\varphi_3 = \varphi_2/P_x(i, j, \varphi_2)$ . Then  $k \in \bar{\varphi}_3(x)$  and  $j \in \bar{\varphi}_3(y) \cap \bar{\varphi}_3(y_k)$ .

Let  $\varphi_4$  be obtained from  $\varphi_3$  by recoloring  $yy_k$  by  $j$ . Then  $1 \in \bar{\varphi}_4(y)$ ,  $\varphi_4(xz) = 1$ ,  $k \in \bar{\varphi}_4(x)$ ,  $\varphi_4(zz_k) = k$ . Since  $\varphi_4(xz) = 1 \in \bar{\varphi}_4(y)$ ,  $\varphi_4$  is feasible. Since  $i, j, k \notin R = C_y \cup C_z$ , the colors in  $R$  are unchanged during this sequence of re-colorings, so  $C_y(\varphi_4) \supseteq C_y$  and  $C_z(\varphi_4) \supseteq C_z$ . Since  $\varphi_4(zz_k) = k \in \bar{\varphi}_4(x)$  and  $d(z_k) < q$ , we have  $k = \varphi_4(zz_k) \in C_z(\varphi_4)$ . So,  $C_z(\varphi_4) \supseteq C_z \cup \{k\}$ . We therefore have  $|C_y(\varphi_4)| + |C_z(\varphi_4)| \geq |C_y| + |C_z| + 1$ , giving a contradiction.

For the case of  $1 \in \bar{\varphi}(z_k)$ , we consider the dual coloring  $\varphi^d$  of  $G - xz$  obtained from  $\varphi$  by uncoloring  $xz$  and coloring  $xy$  with color 1. Following the exact same argument above, we can reach a contradiction to the maximum of  $|C_y| + |C_z|$ . This completes the proof of **II**.  $\square$

### 2.1.2 Proof of III.

Denote by  $Z$  the set of all feasible vertices. For a coloring  $\varphi \in \mathcal{C}^\Delta(G - xy)$ , let  $Z(\varphi) = \{z \in N(x) : \varphi(xz) \in \bar{\varphi}(y)\}$  and  $S(\varphi) = \{z \in N(x) \setminus Z(\varphi) : d(y_{\varphi(xz)}) < q\}$ , where  $y_j \in N(y)$  with  $\varphi(yy_j) = j$  for any color  $j$ . We call vertices in  $S(\varphi)$  *semi-feasible* vertices of  $\varphi$ .

**Claim 2.1.** *For any coloring  $\varphi \in \mathcal{C}^\Delta(G - xy)$ , the following two inequalities hold.*

- a.  $|Z(\varphi) \cup S(\varphi)| \geq \Delta - \sigma_q(x, y) - 1$ ;
- b. *With one possible exception, for all  $z \in S(\varphi)$  there exists a coloring  $\varphi^* \in \mathcal{C}^\Delta(G - xy)$  such that  $\varphi^*(xz) \in \bar{\varphi}^*(y)$ .*

*Proof.* Let  $\varphi \in \mathcal{C}^\Delta(G - xy)$ . Since  $G$  is edge- $\Delta$ -critical,  $xy$  is an edge of  $G$ , it is easy to see that  $\bar{\varphi}(y) \subseteq \varphi(x)$  and  $\bar{\varphi}(x) \subseteq \varphi(y)$ . Divide  $\varphi(y)$  into two subsets:

$$\varphi(y, \geq q) = \{i \in \varphi(y) : d(y_i) \geq q\} \quad \text{and} \quad \varphi(y, < q) = \{i \in \varphi(y) : d(y_i) < q\}.$$

Clearly,  $\sigma_q(x, y) = |\varphi(y, \geq q)|$  and  $|\bar{\varphi}(y)| + |\varphi(y, < q)| = \Delta - \sigma_q(x, y)$ . Since  $\bar{\varphi}(y) \subseteq \varphi(x)$ , to prove **a**, we only need to show  $|\bar{\varphi}(x) \cap \varphi(y, < q)| \leq 1$ . Since edge  $xy$  and the edges incident to  $y$  with colors in  $\bar{\varphi}(x)$  form a Vizing fan  $F$ , the vertex set  $V(F)$  is elementary with respect to  $\varphi$ . Since  $d(x) + 2q < 2\Delta$ ,  $V(F) \setminus \{x\}$  contains at most one vertex with degree  $< q$ . So  $|\bar{\varphi}(x) \cap \varphi(y, < q)| \leq 1$  holds.

To prove **b**, we show that for any two distinct vertices  $z_k, z_\ell \in S(\varphi)$ , there is a coloring  $\varphi^* \in \mathcal{C}^\Delta(G - xy)$  such that at least one of  $\varphi^*(xz_k)$  and  $\varphi^*(xz_\ell)$  is in  $\bar{\varphi}^*(y)$ . We assume  $\varphi(xz_k) = k$  and  $\varphi(xz_\ell) = \ell$ . Let  $y_k, y_\ell \in N(y) \setminus \{x\}$  such that  $\varphi(yy_k) = k$  and  $\varphi(yy_\ell) = \ell$ .

By the definition of  $S(\varphi)$ , we have  $d(y_k) < q$  and  $d(y_\ell) < q$ . Since  $\Delta/2 < q \leq \Delta - d(x)/2 - 2$ , the following inequality holds.

$$|\bar{\varphi}(x)| + |\bar{\varphi}(y_k)| + |\bar{\varphi}(y_\ell)| > \Delta \quad (6)$$

We claim that there exists a coloring  $\varphi^* \in \mathcal{C}^\Delta(G - xy)$  such that keeping the property  $\varphi^*(xz) = 1 \in \bar{\varphi}^*(y)$  and having the following property.

$$\bar{\varphi}^*(x) \cap (\bar{\varphi}^*(y_k) \cup \bar{\varphi}^*(y_\ell)) \neq \emptyset \quad (7)$$

Otherwise, by (6), there exists  $r \in \varphi(x) \cap \bar{\varphi}(y_k) \cap \bar{\varphi}(y_\ell)$ . Choose a color  $i \in \bar{\varphi}(x)$ . Since at least one of colors  $i$  and  $r$  is missing at each of  $x, y_k$  and  $y_\ell$ , we may assume  $P_{y_k}(i, r, \varphi)$  is disjoint with  $P_x(i, r, \varphi)$ . Then, in coloring  $\varphi/P_{y_k}(i, r, \varphi)$ , color  $i$  is missing at both  $x$  and  $y_k$ , giving a contradiction.

By (7), we may assume that there exists a color  $i \in \bar{\varphi}(x) \cap \bar{\varphi}(y_k)$ . Since  $G$  is not edge- $\Delta$ -colorable,  $P_x(1, i, \varphi) = P_y(1, i, \varphi)$ . So,  $P_{y_k}(1, i, \varphi)$  is disjoint with  $P_x(1, i, \varphi)$ . If  $1 \in \varphi(y_k)$ , let  $\varphi' = \varphi/P_{y_k}(1, i, \varphi)$ . For coloring  $\varphi'$ , we have  $1 \in \bar{\varphi}'(y_k)$  and  $\varphi'(x) = \varphi(x), \varphi'(y) = \varphi(y)$ . Thus we can assume  $1 \in \bar{\varphi}(y_k)$ . Let  $\varphi^*$  be a coloring obtained from  $\varphi$  by recoloring  $yy_k$  with color 1. Then,  $\varphi^*(xz_k) = k \in \bar{\varphi}^*(y)$ . This completes the proof of **III**.  $\square$

### 3 Proof of Theorem 1

Let  $G$  be an edge- $\Delta$ -critical graph with  $n$  vertices and  $m$  edges. Clearly,  $\bar{d}(G) = 2m/n$ . We assume  $\Delta \geq 56$ . Let  $q := \min\{\frac{2\sqrt{2}(\Delta-1)-2}{2\sqrt{2}+1}, \frac{3}{4}\Delta - 2\}$ , that is,  $q = \frac{2\sqrt{2}(\Delta-1)-2}{2\sqrt{2}+1}$  if  $\Delta \geq 66$  and  $q = \frac{3}{4}\Delta - 2$  if  $56 \leq \Delta \leq 65$ . We initially assign to each vertex  $x$  of  $G$  a charge  $M(x) = d(x)$  and redistribute the charge according to the following rule:

- **Rule of Discharge:** each ( $> q$ )-vertex  $y$  distributes its surplus charge of  $d(y) - q$  equally among all ( $< q$ )-neighbors of  $y$ .

Denote by  $M'(x)$  the resulting charge on each vertex  $x$ . Clearly,  $\sum_{x \in V(G)} M'(x) = \sum_{x \in V(G)} M(x) = 2m$ . Let  $X_1 = \{x \in V(G) : d(x) \leq 3q - 2\Delta\}$ . We show that  $M'(x) \geq 2 + 2(\Delta - q)$  for all vertices in  $X_1$  and  $M'(x) \geq q$  for all other vertices, which gives  $\bar{d}(G) \geq q - (3q - 2\Delta - 2)\frac{|X_1|}{n}$ . We then show that  $|X_1|/n$  is small in order to complete our proof.

Since  $q = \min\{\frac{2\sqrt{2}(\Delta-1)-2}{2\sqrt{2}+1}, \frac{3}{4}\Delta - 2\}$  and  $\Delta \geq 56$ , we have  $\frac{\Delta+2}{2} < q < \frac{3\Delta}{4}$ . Thus  $q > \Delta - q + 2 > 3q - 2\Delta$ .

**Claim 3.1.** *If  $d(x) \leq \Delta - q + 2$ , then  $M'(x) \geq d(x) + 2(\Delta - q)$ . Consequently,  $M'(x) \geq d(x) + 2(\Delta - q)$  for each  $x \in X_1$ .*

*Proof.* Let  $y$  be an arbitrary neighbor of  $x$ . Since  $2\Delta - d(x) - d(y) + 2 \geq \Delta - d(x) + 2 \geq q$ , we have  $\sigma_q(x, y) \geq \sigma(x, y)$ . We will use lower bounds of  $\sigma(x, y)$  to estimate  $\sigma_q(x, y)$ . Following the definition  $p_{min}(x) = \min_{v \in N(x)} \sigma(x, v) - \Delta + d(x) - 1$  and  $p(x) = \min\{p_{min}(x), \lfloor \frac{d(x)}{2} \rfloor - 1\}$ , we have the following inequalities.

$$1 \leq d_{<q}(y) \leq d(y) - \sigma(x, y) \leq d(y) - (\Delta - d(x) + p(x) + 1) \quad (8)$$

By Lemma 3,  $x$  has at least  $d(x) - p(x) - 1$  neighbors  $y$  for which  $\sigma(x, y) \geq \Delta - p(x) - 1$ , so for these neighbors  $y$  the following inequalities hold.

$$1 \leq d_{<q}(y) \leq d(y) - \sigma(x, y) \leq d(y) - (\Delta - p(x) - 1) \quad (9)$$

We first consider the case  $p(x) \geq 1$ . In this case, we have  $q \leq \Delta - d(x) + 2 \leq \Delta - d(x) + p + 1$ . Since  $\frac{d(y)-a}{d(y)-b}$  with  $a \leq b$  is a decreasing function of  $d(y)$ , for each  $y \in N(x)$ ,  $x$  receives charge at least

$$\frac{d(y) - q}{d(y) - (\Delta - d(x) + p(x) + 1)} \geq \frac{\Delta - q}{d(x) - p(x) - 1},$$

And there are at least  $d(x) - p(x) - 1$  neighbors  $y$  of  $x$  giving  $x$  at least

$$\frac{d(y) - q}{d(y) - (\Delta - p(x) - 1)} \geq \frac{\Delta - q}{p(x) + 1},$$

where the inequality holds because  $q \leq \Delta - d(x) + 2 \leq \Delta - p(x) - 1$  as  $1 \leq p(x) \leq \lfloor \frac{d(x)}{2} \rfloor - 1$ . Thus  $x$  receives at least

$$(d(x) - p(x) - 1) \frac{\Delta - q}{p(x) + 1} + (p(x) + 1) \frac{\Delta - q}{d(x) - p(x) - 1} = (\theta + \theta^{-1})(\Delta - q) \geq 2(\Delta - q),$$

where  $\theta = \frac{d(x) - p(x) - 1}{p(x) + 1}$ . It follows that  $M'(x) \geq M(x) + 2(\Delta - q) = d(x) + 2(\Delta - q)$ .

We now consider the case  $p(x) = \min\{p_{min}(x), \lfloor \frac{d(x)}{2} \rfloor - 1\} = 0$ . If  $d(x) = 2$ , then by (8) for every neighbor  $y$  of  $x$  we have  $d_{<q}(y) = 1$  and  $d(y) = \Delta$ , thus  $M'(x) \geq M(x) + 2(\Delta - q) = d(x) + 2(\Delta - q)$ . If  $d(x) \geq 3$ , then by (9) for at least  $d(x) - 1$  neighbors  $y$  of  $x$ , we have  $d_{<q}(y) = 1$  and  $d(y) = \Delta$ . Thus  $M'(x) \geq M(x) + (d(x) - 1)(\Delta - q) \geq d(x) + 2(\Delta - q)$ .  $\square$

**Claim 3.2.** For each  $x \in V(G) - X_1$ ,  $M'(x) \geq q$  holds.

*Proof.* Let  $x \in V(G) - X_1$ , i.e.,  $d(x) > 3q - 2\Delta$ . If  $d(x) \geq q$ , then  $M'(x) = M(x) - \frac{d(x)-q}{d_{<q}(x)}d_{<q}(x) = q$ . If  $3q - 2\Delta < d(x) \leq \Delta - q + 2$ , then by Claim 3.1, we have  $M'(x) \geq d(x) + 2(\Delta - q) > q$ . So we only need to consider the case  $\Delta - q + 2 < d(x) < q$ .

Since  $G$  is edge- $\Delta$ -critical and  $xy \in E(G)$ , there exists a coloring  $\varphi \in \mathcal{C}^\Delta(G - xy)$ . Let  $Z_q = \{z \in N(x) : d(z) > q\}$ ,  $Z_y = \{z \in N(x) \setminus \{y\} : \varphi(xz) \in \bar{\varphi}(y)\}$  and  $Z_q^* = Z_q \cap Z_y$ . Clearly, for each  $z \in Z_q$ ,  $x$  receives at least  $\frac{d(z)-q}{d(z)-\sigma_q(x,z)}$  charge. Thus  $M'(x) \geq d(x) + \sum_{z \in Z_q} \frac{d(z)-q}{d(z)-\sigma_q(x,z)}$ . We consider the following three cases to complete the proof.

**Case 1.**  $\Delta - q + 2 < d(x) < q$  and  $x$  has a neighbor  $y$  such that  $d(y) \leq q$ .

By Lemma 8 (3), for each vertex  $z \in Z_q^*$  we have

$$\sigma_q(x, z) \geq 2\Delta - d(x) - d(y) + 1 - \lfloor \frac{d(x) + d(y) + d(z) - 2\Delta - 2}{\Delta - q} \rfloor \geq 2\Delta - d(x) - d(y),$$

where we used the inequality  $\lfloor \frac{d(x)+d(y)+d(z)-2\Delta-2}{\Delta-q} \rfloor \leq 1$  following  $d(x) < q, d(y) \leq q, d(z) \leq \Delta$  and  $q < \frac{3}{4}\Delta$ . Thus  $\sigma_q(x, z) \geq 2\Delta - d(x) - d(y)$ . So,  $M'(x) \geq M(x) + \sum_{z \in Z_q^*} \frac{d(z)-q}{d(z)-(2\Delta-d(x)-d(y))}$ . By Lemma 8 (2), we have  $\sum_{z \in Z_q^*} (d(z) - q) \geq (\Delta - d(y) + 1)(\Delta - q) - d(x) - d(y) + \Delta + 2$ . Thus

$$\begin{aligned} \sum_{z \in Z_q^*} \frac{d(z) - q}{d(z) - (2\Delta - d(x) - d(y))} &\geq \frac{(\Delta - q)(\Delta - d(y) + 1) - (d(x) + d(y) - \Delta - 2)}{d(x) + d(y) - \Delta} \\ &\geq \frac{(\Delta - q)(\Delta - d(y) + 1) + 2}{d(x) + d(y) - \Delta} - 1. \end{aligned}$$

So, we have the following inequality.

$$\begin{aligned} M'(x) &\geq M(x) + \sum_{z \in Z_q^*} \frac{d(z) - q}{d(z) - (2\Delta - d(x) - d(y))} \\ &\geq d(x) + \frac{(\Delta - q)(\Delta - d(y) + 1) + 2}{d(x) + d(y) - \Delta} - 1 \\ &\geq d(x) + q - \Delta + \frac{(\Delta - q)(\Delta - q + 1) + 2}{d(x) + q - \Delta} - 1 - q + \Delta \\ &\geq 2\sqrt{(\Delta - q)(\Delta - q + 1) + 2} + \Delta - q - 1 \\ &\geq 3(\Delta - q) \geq q. \end{aligned}$$

**Case 2.**  $2(\Delta - q) - 4 < d(x) < q$  and  $d(y) > q$  for every neighbor  $y$  of  $x$ .

Let  $y \in N(x)$  such that  $d(y) := \min\{d(u) : u \in N(x)\}$ . By Lemma 8 (3), for each vertex  $z \in Z_q^*$  we have

$$\begin{aligned}\sigma_q(x, z) &\geq 2\Delta - d(x) - d(y) + 1 - \left\lfloor \frac{d(x) + d(y) + d(z) - 2\Delta - 2}{\Delta - q} \right\rfloor \\ &\geq 2\Delta - d(x) - d(y) - 1,\end{aligned}$$

where we used the inequality  $\lfloor \frac{d(x)+d(y)+d(z)-2\Delta-2}{\Delta-q} \rfloor \leq 2$  when  $d(x) < q$  and  $q < \frac{3}{4}\Delta$ . By Lemma 8 (2), we have

$$\begin{aligned}\sum_{z \in Z_q^*} \frac{d(z) - q}{d(z) - \sigma_q(x, y)} &\geq \frac{(\Delta - d(y) + 1)(\Delta - q) - d(x) - d(y) + \Delta + 2}{d(x) + d(y) - \Delta + 1} \\ &= q - \Delta - 1 + \frac{(\Delta - q)(d(x) + 2) + 3}{d(x) + d(y) - \Delta + 1}.\end{aligned}$$

By Lemma 1, for each neighbor  $u$  of  $x$  we have  $\sigma_\Delta(x, u) \geq \Delta - d(x) + 1$ . Since  $d(u) \geq d(y)$  for each  $u \in N(x)$  and  $q \geq \Delta - d(x) + 1$ , we have

$$\frac{d(u) - q}{d(u) - (\Delta - d(x) + 1)} \geq \frac{d(y) - q}{d(y) - (\Delta - d(x) + 1)}.$$

So,

$$\begin{aligned}\sum_{u \in N(x) \setminus Z_q^*} \frac{d(u) - q}{d(u) - (\Delta - d(x) + 1)} &\geq |N(x) \setminus Z_q^*| \cdot \frac{d(y) - q}{d(y) - (\Delta - d(x) + 1)} \\ &= \frac{(d(x) - (\Delta - d(y) + 1))(d(y) - q)}{d(y) - (\Delta - d(x) + 1)} = d(y) - q.\end{aligned}$$

Thus

$$\begin{aligned}M'(x) &\geq d(x) + q - \Delta - 1 + \frac{(\Delta - q)(d(x) + 2) + 3}{d(x) + d(y) - \Delta + 1} + d(y) - q \\ &= d(x) + d(y) - \Delta + 1 + \frac{(\Delta - q)(d(x) + 2) + 3}{d(x) + d(y) - \Delta + 1} - 2 \\ &\geq 2\sqrt{(\Delta - q)(d(x) + 2) + 3} - 2 \\ &> 2\sqrt{2(\Delta - q)(\Delta - q - 1)} - 2 \geq q.\end{aligned}$$



**Case 3.**  $\Delta - q + 2 < d(x) \leq 2(\Delta - q) - 4$  and  $d(y) > q$  for every neighbor  $y$  of  $x$ .

Since  $\Delta \geq 56$ , we have  $\Delta - q + 2 < 2(\Delta - q) - 4$ , so this case occurs. Since the notation  $p(x, q)$  will be used heavily in this proof, we let  $p' := p(x, q)$  for convenience. So,  $p' = \min\{ p_{\min}(x, q), \lfloor \frac{d(x)}{2} \rfloor - 3 \}$ , where  $p_{\min}(x, q) := \min_{y \in N(x)} \sigma_q(x, y) - \Delta + d(x) - 1$ . Following this definition, for every  $y \in Z_q$ ,  $\sigma_q(x, y) \geq \Delta - d(x) + p' + 1$ , which in turn gives

$$\frac{d(y) - q}{d(y) - \sigma_q(x, y)} \geq \frac{d(y) - q}{d(y) - (\Delta - d(x) + p' + 1)}.$$

So, if  $q \leq \Delta - d(x) + p' + 1$ , then

$$\frac{d(y) - q}{d(y) - \sigma_q(x, y)} \geq \frac{\Delta - q}{d(x) - p' - 1}.$$

By Lemma 5,  $x$  has at least  $d(x) - p' - 3$  neighbors  $y$  for which  $\sigma_q(x, y) \geq \Delta - p' - 5$ . For such neighbors  $y$ , since  $q \leq \Delta - \frac{d(x)}{2} - 2 \leq \Delta - p' - 5$ , we have

$$\frac{d(y) - q}{d(y) - \sigma_q(x, y)} \geq \frac{d(y) - q}{d(y) - (\Delta - p' - 5)} \geq \frac{\Delta - q}{p' + 5}.$$

If  $q \leq \Delta - d(x) + p' + 1$ , then

$$\begin{aligned} M'(x) &\geq d(x) + (d(x) - p' - 3) \frac{\Delta - q}{p' + 5} + (p' + 3) \frac{\Delta - q}{d(x) - p' - 1} \\ &\geq \Delta - q + 2 + (\Delta - q) \left( 2 - \frac{8(d(x) + 2)}{(\Delta - q + 2)(d(x) + 4)} \right) \geq q, \end{aligned}$$

where we used the inequality  $\theta + \theta^{-1} \geq 2$  ( $\theta = \frac{d(x) - p' - 3}{p' + 5}$ ) to show the following

$$\frac{d(x) - p' - 3}{p' + 5} + \frac{p' + 3}{d(x) - p' - 1} > 2 - \frac{2d(x) + 4}{\left(\frac{\Delta - q + 2}{2}\right)\left(\frac{d(x)}{2} + 2\right)} = 2 - \frac{8(d(x) + 2)}{(\Delta - q + 2)(d(x) + 4)}.$$

Suppose  $q > \Delta - d(x) + p' + 1$ , i.e.,  $p' < d(x) + q - \Delta - 1$ . So,  $\frac{d(x) - p' - 3}{p' + 5} > \frac{\Delta - q - 2}{d(x) + q - \Delta + 4}$ , which gives

$$\begin{aligned} M'(x) &\geq d(x) + \frac{\Delta - q - 2}{d(x) + q - \Delta + 4} (\Delta - q) \\ &= (d(x) + q - \Delta + 4) + \frac{(\Delta - q)(\Delta - q - 2)}{d(x) + q - \Delta + 4} - (q - \Delta + 4) \\ &\geq 2\sqrt{(\Delta - q)(\Delta - q - 2)} + \Delta - q - 4 \geq 3(\Delta - q) - 8 \geq q. \end{aligned}$$

□

**Claim 3.3.**  $d(y) > q$  for each  $y \in N(X_1)$  and  $|N(X_1)| \geq 2|X_1|$  where  $N(X_1) = \cup_{x \in X_1} N(x)$ .

*Proof.* Since  $G$  is edge- $\Delta$ -critical, for each edge  $xy \in E(G)$  we have  $d(x) + d(y) \geq \Delta + 2$ . Since  $q < \frac{3}{4}\Delta$  and  $d(x) \leq 3q - 2\Delta$  for each  $x \in X_1$ , we have  $d(y) \geq \Delta + 2 - (3q - 2\Delta) > q$  for each  $y \in N(x)$ . Thus the vertices in  $N(X_1)$  does not receive charges from any other vertices. As the vertices in  $X_1$  receive charges only from the vertices in  $N(X_1)$ , we have

$$\sum_{x \in X_1} M'(x) + \sum_{y \in N(X_1)} M'(y) \leq \sum_{x \in X_1} M(x) + \sum_{y \in N(X_1)} M(y) \leq \sum_{x \in X_1} d(x) + \Delta|N(X_1)|. \quad (10)$$

Also, by Claims 3.1 and 3.2, we have  $M'(x) \geq d(x) + 2(\Delta - q)$  for each  $x \in X_1$  and  $M'(y) \geq q$  for each  $y \in N(X_1)$ . Thus we have

$$\sum_{x \in X_1} M'(x) + \sum_{y \in N(X_1)} M'(y) \geq \sum_{x \in X_1} d(x) + 2(\Delta - q)|X_1| + q|N(X_1)|. \quad (11)$$

Combining (10) with (11), we have  $|N(X_1)| \geq 2|X_1|$ .  $\square$

For each edge  $xy \in E(G)$  and  $\varphi \in \mathcal{C}^\Delta(G - xy)$ , let  $Y(x, \varphi) = \{w \in N(y) \setminus \{x\} : \varphi(yw) \in \bar{\varphi}(x)\}$ ,  $Y^1(x, \varphi) = Y(x, \varphi) \cap N(X_1)$  and  $Y^2(x, \varphi) = Y(x, \varphi) - (X_1 \cup N(X_1))$ . Clearly,  $|Y(x, \varphi)| = \Delta - d(x) + 1$ . Note that with respect to the coloring  $\varphi$ ,  $\{x, y\} \cup Y(x, \varphi)$  forms a Vizing fan, so it is elementary.

**Claim 3.4.** For each  $y \in N(X_1)$  and  $x \in N(y) \cap X_1$ ,  $|Y^2(x, \varphi)| \geq \Delta - 2d(x) + 3$ .

*Proof.* Recall that  $\{x, y\} \cup Y(x, \varphi)$  is elementary. Then for each vertex in  $w \in Y(x, \varphi)$  we have  $|\bar{\varphi}(w)| + |\bar{\varphi}(x)| + |\bar{\varphi}(y)| \leq \Delta$ , it follows that  $d(w) \geq |\bar{\varphi}(x)| + |\bar{\varphi}(y)| > 3q - 2\Delta$ . Thus we have  $Y(x, \varphi) \cap X_1 = \emptyset$ . If  $|Y^1(x, \varphi)| \leq d(x) - 2$ , then  $|Y^2(x, \varphi)| = |Y(x, \varphi) - Y^1(x, \varphi) - (Y(x, \varphi) \cap X_1)| \geq \Delta - d(x) + 1 - (d(x) - 2) \geq \Delta - 2d(x) + 3$ . So the Claim 3.4 is equivalent to show that  $|Y^1(x, \varphi)| \leq d(x) - 2$ .

**Subclaim 3.4.1.** If  $w \in Y^1(x, \varphi)$ , then for each neighbor  $z$  of  $w$  in  $X_1$ , we have  $\varphi(wz) \in \varphi(x) \cap \varphi(y)$ .

*Proof.* If there exists a neighbor of  $w$  in  $X_1$ , say  $z$ , such that  $\varphi(wz) \notin \varphi(x) \cap \varphi(y)$ . Then  $\varphi(wz) \in \bar{\varphi}(x) \cup \bar{\varphi}(y)$ . Thus  $\{x, y, w, z\}$  forms a Kierstead path. By Lemma 6, we have  $|\bar{\varphi}(z) \cap (\bar{\varphi}(x) \cup \bar{\varphi}(y))| \leq 1$ , it follows that  $d(z) \geq (\Delta - d(x) + 1) + (\Delta - d(y) + 1) - 1 > 3q - 2\Delta$ , this contradicts with the fact that  $z \in X_1$ . So Subclaim 3.4.1. holds.  $\square$

For each color  $j \in \varphi(x) \cap \varphi(y)$ , set  $Y_j = \{w \in Y^1(x, \varphi) : j \in \varphi(w)\}$  and  $Z_j = \{z \in X_1 : \text{there exists a vertex } w \in Y_j \text{ such that } \varphi(wz) = j\}$ . Clearly,  $\sum_{j \in \varphi(x) \cap \varphi(y)} |Z_j| \geq |Y^1(x, \varphi)|$ . Since  $|\varphi(x) \cap \varphi(y)| = \Delta - (\Delta - d(x) + 1) - (\Delta - d(y) + 1) \leq d(x) - 2$ , to show that  $|Y^1(x, \varphi)| \leq d(x) - 2$ , we only need to prove that  $|Z_j| \leq 1$  for each  $j$ . Let  $|Z_j| = t$  and  $Z_j = \{z_{\alpha_1}, \dots, z_{\alpha_t}\}$ , where for each  $z_{\alpha_i}$  there exists  $y_{\alpha_i} \in Y_j$  such that  $\varphi(yy_{\alpha_i}) = \alpha_i$  and  $\varphi(y_{\alpha_i}z_{\alpha_i}) = j$ . Clearly,  $\alpha_i \in \bar{\varphi}(x)$  for each  $1 \leq i \leq t$ .

**Subclaim 3.4.2.** Let  $k$  be a color in  $\bar{\varphi}(x)$ . Then the followings hold.

- (1) For each  $k \notin \{\alpha_1, \dots, \alpha_t\}$ , at least  $t - 1$  vertices of  $Z_j$  have the color  $k$ .
- (2) For each  $k \in \{\alpha_1, \dots, \alpha_t\}$ , at least  $t - 2$  vertices of  $Z_j$  have the color  $k$ .

*Proof.* First suppose that  $k \notin \{\alpha_1, \dots, \alpha_t\}$ . We consider the path  $P_x(j, k, \varphi)$ , and  $w$  is the other end vertex of this path. We will show that the color  $k$  seen by each vertex in  $Z_j \setminus \{w\}$ . For otherwise, we assume  $k \notin \varphi(z)$  for some  $z \in Z_j \setminus \{w\}$ , say  $z = z_{\alpha_1}$ , then  $P_{z_{\alpha_1}}(j, k, \varphi)$  is disjoint from  $P_x(j, k, \varphi)$ , thus let  $\varphi' = \varphi / P_{z_{\alpha_1}}(j, k, \varphi)$  be the new coloring which  $\varphi'(yy_{\alpha_1}) = \varphi(yy_{\alpha_1}) \in \bar{\varphi}(x)$  and  $\varphi'(y_{\alpha_1}z_{\alpha_1}) = k \in \bar{\varphi}(x)$ . Thus  $\{x, y, y_{\alpha_1}, z_{\alpha_1}\}$  forms a Kierstead path. So by Lemma 6 we have  $|\bar{\varphi}(z_{\alpha_1}) \cap (\bar{\varphi}(x) \cup \bar{\varphi}(y))| \leq 1$ , it follows that  $d(z_{\alpha_1}) \geq (\Delta - d(x) + 1) + (\Delta - d(y) + 1) - 1 > 3q - 2\Delta$  as  $d(x) \leq 3q - 2\Delta$ , this contradicts with the fact that  $z_{\alpha_1} \in X_1$ .

Then suppose that  $k \in \{\alpha_1, \dots, \alpha_t\}$ . We may assume that  $k = \alpha_t$ . Clearly,  $k \notin \{\alpha_1, \dots, \alpha_{t-1}\}$ . Let  $Z'_j = Z_j - \{z_k\}$ . By (1), we have at least  $|Z'_j| - 1$  vertices of  $Z'_j$  has the color  $k$ , that is, at least  $t - 2$  vertices of  $Z_j$  has the color  $k$ .  $\square$

By Subclaim 3.4.2 and the definition of  $X_1$ , we have

$$(\Delta - d(x) + 1 - t)(t - 1) + t(t - 2) \leq \sum_{z \in Z_j} d(z) \leq t(3q - 2\Delta).$$

Since  $d(x) \leq 3q - 2\Delta$  and  $q < \frac{3}{4}\Delta$ , we have

$$t \leq \frac{\Delta - d(x) + 1}{3\Delta - 3q - d(x)} \leq 1 + \frac{3q - 2\Delta + 1}{5\Delta - 6q} < 2.$$

Since  $t$  is an integer, we have  $t \leq 1$ . Then Claim 3.4 holds.  $\square$

Let  $c$  be a positive integer, set  $Z_1(c) = \{z \in V(G) - (X_1 \cup N(X_1)) : d(z) \geq \Delta - c\}$  and  $Z_2(c) = \{z \in V(G) - (X_1 \cup N(X_1)) : d(z) < \Delta - c\}$ .

**Claim 3.5.**  $|Z_1(c)| \geq \frac{(5c+2)\Delta - (6c+3)q + 3c+2}{c\Delta} |N(X_1)|$ .

*Proof.* For each  $y \in N(X_1)$ ,  $x \in N(y) \cap X_1$  and  $\varphi \in \mathcal{C}^\Delta(G - xy)$ , let  $Y_{<c} = \{z \in Y^2(x, \varphi) : d(z) < \Delta - c\}$ . Since  $\{x, y\} \cup Y(x, \varphi)$  is elementary and  $Y^2(x, \varphi) \subseteq Y(x, \varphi)$ , we have  $\Delta - d(x) + 1 + \Delta - d(y) + 1 + c|Y_{<c}| < \sum_{v \in \{x, y\} \cup Y(x, \varphi)} \bar{\varphi}(v) \leq \Delta$ . Thus  $|Y_{<c}| < \frac{d(x)-2}{c}$ . By Claim 3.4, we have  $|Y^2(x, \varphi) - Y_{<c}| > \Delta - 2d(x) + 3 - \frac{d(x)-2}{c}$ , that is, for each  $y \in N(X_1)$  we have  $d_{Z_1(c)}(y) \geq \Delta - 2d(x) + 3 - \frac{d(x)-2}{c}$ . Hence,

$$(\Delta - 2d(x) + 3 - \frac{d(x)-2}{c})|N(X_1)| \leq |E(N(X_1), Z_1(c))| \leq \Delta|Z_1(c)|,$$

where  $E(N(X_1), Z_1(c))$  are the edges with one vertex in  $N(X_1)$  and the other endvertex in  $Z_1(c)$ . Since  $d(x) \leq 3q - 2\Delta$ , solving the above inequalities we have

$$|Z_1(c)| \geq \frac{(5c+2)\Delta - (6c+3)q + 3c + 2}{c\Delta}|N(X_1)|.$$

□

By Claims 3.1 and 3.2, we have

$$M'(x) \geq \begin{cases} 2 + 2(\Delta - q) & x \in X_1, \\ q & x \in V(G) - X_1. \end{cases}$$

And by the definitions of  $Z_1(c)$  and  $Z_2(c)$ , we get the following two lower bounds of  $\sum_{x \in V(G)} M'(x)$ .

$$\begin{aligned} b_1 &= (2 + 2(\Delta - q))|X_1| + q|N(X_1)| + (\Delta - c)|Z_1(c)| + (3q - 2\Delta)|Z_2(c)| \\ b_2 &= (2 + 2(\Delta - q))|X_1| + (n - |X_1|)q \end{aligned}$$

We now divide into a few cases to estimate the lower bound of  $\max\{b_1, b_2\}$ .

First we consider the case  $\Delta - q - c > 0$ . For fixed value  $|X_1| + |N(X_1)|$ ,  $|Z_1(c)| + |Z_2(c)|$  is a constant. Since  $\Delta - c > q > 3q - 2\Delta$ ,  $\max\{(\Delta - c)|Z_1(c)| + (3q - 2\Delta)|Z_2(c)|, q(|Z_1(c)| + |Z_2(c)|)\}$  takes minimum when  $(\Delta - c)|Z_1(c)| + (3q - 2\Delta)|Z_2(c)| = q(|Z_1(c)| + |Z_2(c)|)$ , that is  $|Z_2(c)| = (\frac{1}{2} - \frac{c}{2\Delta - 2q})|Z_1(c)|$ . So by Claims 3.3 and 3.5,  $|Z_1(c)| + |Z_2(c)| \geq (3 - \frac{c}{\Delta - q})\frac{(5c+2)\Delta - (6c+3)q + 3c + 2}{c\Delta}|X_1|$ . Let  $f(c) = (3 - \frac{c}{\Delta - q})\frac{(5c+2)\Delta - (6c+3)q + 3c + 2}{c\Delta}$ . So  $n = |X_1| + |N(X_1)| + |Z_1(c)| + |Z_2(c)| \geq (3 + f(c))|X_1|$ . Hence,  $\sum_{x \in V(G)} M'(x) \geq \max\{b_1, b_2\} \geq qn + (2 + 2\Delta - 3q)|X_1| \geq (q + \frac{2+2\Delta-3q}{3+f(c)})n$ . So  $\bar{d}(G) \geq q + \frac{2+2\Delta-3q}{3+f(c)}$ .

Let  $q^* = \frac{2\sqrt{2}\Delta}{2\sqrt{2}+1}$  and  $a = 1 + \frac{1}{2\sqrt{2}+1}$  if  $q = \frac{2\sqrt{2}(\Delta-1)-2}{2\sqrt{2}+1}$ , and  $q^* = \frac{3\Delta}{4}$ ,  $a = 2$  if  $q = \frac{3}{4}\Delta - 2$ . So  $q = q^* - a$  and we have

$$\begin{aligned} \frac{2+2\Delta-3q}{3+f(c)} &= \frac{c\Delta(2+2\Delta-3q)}{3c\Delta + (3 - \frac{c}{\Delta-q})((5c+2)\Delta - (6c+3)q + 3c+2)} \\ &= \frac{(2c\Delta - 3cq^*)\Delta + f_1f_2\Delta - f_1f_2\Delta + 2c\Delta + 3ca\Delta}{(18c+6)\Delta - (18c+9)q^* + f_2} \\ &= f_1\Delta + \frac{-f_1f_2\Delta + 2c\Delta + 3ca\Delta}{(18c+6)\Delta - (18c+9)q^* + f_2}, \end{aligned}$$

where  $f_1 = \frac{2c\Delta-3cq^*}{(18c+6)\Delta-(18c+9)q^*}$  and  $f_2 = 9c+6+(18c+9)a - \frac{(5c^2+2c)\Delta-(6c^2+3c)q+3c^2+2c}{\Delta-q}$ .

Clearly,  $f_1 = \frac{2c\Delta-3cq^*}{(18c+6)\Delta-(18c+9)q^*} = \frac{2\Delta-3q^*}{18\Delta-18q^*+\frac{6\Delta-9q^*}{c}}$  is an increasing function of  $c$ . To make  $f_1$  as large as possible when  $\Delta \geq l$ , where  $l$  is a positive integer, we choose  $c$  such that  $c = \min\{[\Delta - q] : \Delta \geq l\}$ . If  $l$  is large enough,  $c$  is large too and we can see that the value of  $f_1$  will approximate to  $\frac{2\Delta-3q^*}{18\Delta-18q^*}$  and  $\bar{d}(G)$  will approximate to  $0.69277\Delta$ . Note that  $q = \frac{2\sqrt{2}(\Delta-1)-2}{2\sqrt{2}+1}$  if  $\Delta \geq 66$ . Let  $l = 66$ . Then we have  $c = 18$ . So we have  $\Delta - q - c > 0$  if  $\Delta \geq 65$ . Plugging  $c = 18$  and the value of  $q^*$  into  $f_2$  and  $\frac{-f_1f_2\Delta+2c\Delta+3ca\Delta}{(18c+6)\Delta-(18c+9)q^*+f_2}$ , we have  $\frac{-f_1f_2\Delta+2c\Delta+3ca\Delta}{(18c+6)\Delta-(18c+9)q^*+f_2} > 0$  and  $f_2 < 0$ . Thus

$$\frac{2+2\Delta-3q}{3+f(c)} \geq f_1\Delta + \frac{2c\Delta + 3ca\Delta - f_2f_1\Delta}{(18c+6)\Delta - (18c+9)q^*}. \quad (12)$$

If  $\Delta \geq 66$ , then  $q = \frac{2\sqrt{2}(\Delta-1)-2}{2\sqrt{2}+1}$ . Plugging  $c = 18$  into the inequality (12), we get  $f_1 \geq -0.04638$ ,  $f_2 \geq -244.43905$  and

$$\frac{2+2\Delta-3q}{3+f(c)} \geq -0.04638\Delta + 1.10463.$$

Thus

$$\bar{d}(G) \geq q - 0.04638\Delta + 1.10463 \geq 0.69241\Delta - 0.15658.$$

If  $\Delta = 65$ , then  $q = \frac{3}{4}\Delta - 2$ . Plugging  $c = 18$  and  $\Delta = 65$  into the inequality (12), we get  $f_1 \geq -0.05608$ ,  $f_2 \geq -1.15069$  and  $\frac{2+2\Delta-3q}{3+f(c)} \geq -0.05608\Delta + 1.79358$ . It follows that

$$\bar{d}(G) \geq q - 0.05608\Delta + 1.79358 \geq 0.69392\Delta - 0.20642.$$

Now we consider the case  $\Delta - q - c \leq 0$ . It is easy to see that  $b_2 > b_1$  and  $\Delta \leq 64$ . Thus  $\sum_{x \in V(G)} M'(x) \geq qn - (3q - 2\Delta - 2)|X_1|$ . So  $\bar{d}(G) \geq q - (3q - 2\Delta - 2)\frac{|X_1|}{n}$ . By Claims 3.3 and 3.5, we have

$$n \geq |X_1| + |N(X_1)| + |Z_1(c)| \geq (3 + f'(c))|X_1|,$$

where  $f'(c) = 2\frac{(5c+2)\Delta - (6c+3)q + 3c+2}{c\Delta}$ .

Plugging  $c = 18$  and  $q = \frac{3}{4}\Delta - 2$  into  $f'(c)$ , we have  $f'(c) = \frac{8.75\Delta + 278}{9\Delta}$ . Since  $\frac{|X_1|}{n} \leq \frac{1}{3+f'(c)}$ , we have  $(3q - 2\Delta - 2)\frac{|X_1|}{n} \leq \frac{2.25\Delta^2 - 72\Delta}{35.75\Delta + 278} = \frac{9\Delta}{143} + \frac{695.94484}{35.75\Delta + 278} - 2.50339$ , thus  $\bar{d}(G) \geq \frac{393\Delta}{572} - \frac{695.94484}{35.75\Delta + 278} + 0.50339$ . It is easy to check that if  $\Delta \geq 56$  then  $\bar{d}(G) \geq \frac{2}{3}(\Delta + 2)$ , which improve Woodall's result in [15]. If  $\Delta \geq 56$ , we have

$$\bar{d}(G) \geq 0.68706\Delta + 0.19815.$$

Hence,

$$\bar{d}(G) \geq \begin{cases} 0.69241\Delta - 0.15658 & \text{if } \Delta \geq 66, \\ 0.69392\Delta - 0.20642 & \text{if } \Delta = 65, \text{ and} \\ 0.68706\Delta + 0.19815 & \text{if } 56 \leq \Delta \leq 64. \end{cases}$$

This completes the proof of Theorem 1.

## References

- [1] G. Chen, X. Chen, and Y. Zhao. Hamiltonianicity of edge chromatic critical graph. Discrete Math., accepted.
- [2] S. Fiorini. Some remarks on a paper by Vizing on critical graphs. Math. Proc. Cambridge Philos. Soc., 77:475-483, 1975.
- [3] S. Fiorini and R. J. Wilson. Edge-colourings of graphs, Research notes in Maths. Pitman, London, 1977.
- [4] D. Haile. Bounds on the size of critical edge-chromatic graphs. Ars Combin., 53:85-96, 1999.
- [5] I. T. Jakobsen. On critical graphs with chromatic index 4. Discrete Math., 9:265-276, 1974.
- [6] K. Kayathri. On the size of edge-chromatic critical graphs. Graphs Combin., 10:139-144, 1994.

- [7] H. A. Kierstead. On the chromatic index of multigraphs without large triangles. *J. Combin. Theory Ser. B*, 36(2):156-160, 1984.
- [8] R. Luo, L. Miao, and Y. Zhao. The size of edge chromatic critical graphs with maximum degree 6. *J. Graph Theory*, 60:149-171, 2009.
- [9] D. P. Sanders and Y. Zhao. On the size of edge chromatic critical graphs. *J. Combin. Theory Ser. B*, 86:408-412, 2002.
- [10] M. Stiebitz, D. Scheide, B. Toft, and L. M. Favrholdt. *Graph edge-coloring: Vizing's theorem and Goldberg's conjecture*. Wiley, 2012.
- [11] V. A. Tashkinov. On an algorithm for the edge coloring of multigraphs. *Diskretn. Anal. Issled. Oper. Ser. 1*, 7(3):72-85, 100, 2000.
- [12] V. G. Vizing. On an estimate of the chromatic class of a p-graph (in Russian). *Diskret. Analiz No.*, 3:25-30, 1964.
- [13] V. G. Vizing. Critical graphs with a given chromatic class (in Russian). *Diskret. Analiz No.*, 5:9-17, 1965.
- [14] V. G. Vizing. Some unsolved problems in graph theory (in Russian). *Uspekhi Mat. Nauk*, 23:117-134, 1968.
- [15] D. R. Woodall. The average degree of an edge-chromatic critical graph. II. *J. Graph Theory*, 56:194-218, 2007.
- [16] D. R. Woodall. The average degree of an edge-chromatic critical graph. *Discrete Math.*, 308:803-819, 2008.
- [17] L. Zhang. Every planar graph with maximum degree 7 is of class 1. *Graphs Combin.*, 16(4):467-495, 2000.