

# Partitioning a graph into monochromatic connected subgraphs

António Girão\*      Shoham Letzter†      Julian Sahasrabudhe‡

March 12, 2022

## Abstract

A well-known result by Haxell and Kohayakawa states that the vertices of an  $r$ -coloured complete graph can be partitioned into  $r$  monochromatic connected subgraphs of distinct colours; this is a slightly weaker variant of a conjecture by Erdős, Pyber and Gyárfás that states that there exists a partition into  $r - 1$  monochromatic connected subgraphs. We consider a variant of this problem, where the complete graph is replaced by a graph with large minimum degree, and prove two conjectures of Bal and DeBiasio, for two and three colours.

## 1 Introduction

An old observation by Erdős and Rado says that when the edges of a complete graph are coloured with two colours, there is a spanning monochromatic component. This simple remark has been the starting point of extensive research. A natural example is the search for large monochromatic components in  $r$ -edge-coloured complete graph (see, for example, [8, 9]). Here we focus on a different direction, namely, the search for covers (or partitions) of the vertices into as few as possible monochromatic connected subgraphs.

A classical example appears in a seminal paper by Erdős, Gyárfás and Pyber [7], who showed that for any  $r$ -colouring of  $K_n$  (the complete graph on  $n$  vertices) the vertices can be partitioned into at most  $O(r^2 \log r)$  monochromatic cycles. We note that throughout the paper, when we say that the vertices of a graph are covered (or partitioned) by a collection of subgraphs, we mean that the vertices are covered by the *vertex sets* of these subgraphs.

Gyárfás, Ruszinkó, Sárközy and Szemerédi [10] improved the above result by showing that if the edges of the complete graph are  $r$ -coloured then the vertices can be partitioned into  $O(r \log r)$  monochromatic

---

\*Department of Pure Mathematics and Mathematical Statistics, University of Cambridge, Cambridge, UK; e-mail: [A.Girao@dpms.cam.ac.uk](mailto:A.Girao@dpms.cam.ac.uk).

†ETH Institute for Theoretical Studies, ETH, 8092 Zurich; e-mail: [shoham.letzter@eth-its.ethz.ch](mailto:shoham.letzter@eth-its.ethz.ch).

‡Department of Mathematics, University of Memphis, Memphis, Tennessee; e-mail: [julian.sahasra@gmail.com](mailto:julian.sahasra@gmail.com).

cycles. In the other direction, Pokrovskiy [15] showed that one needs strictly more than  $r$  cycles, disproving a conjecture of Erdős, Gyárfás and Pyber [7]. Conlon and Stein [4] showed similar results for colourings where every vertex is incident with at most  $r$  distinct colours. The question of whether one can partition an  $r$ -coloured graph into  $O(r)$  monochromatic cycles remains an enticing open problem in this area.

In a slightly different direction, Erdős, Gyárfás and Pyber [7] conjectured that the vertices of an  $r$ -coloured complete graph may be partitioned into at most  $r - 1$  monochromatic connected subgraphs. This conjecture is known to be tight when  $r - 1$  is a prime power and  $n$  is sufficiently large, due to a well-known construction which requires the existence of an affine plane of an appropriate order. Haxell and Kohayakawa [11] proved a slightly weaker result, showing that one can partition an  $r$ -coloured complete graph on  $n$  vertices into  $r$  monochromatic subgraphs, for sufficiently large  $n$ .

Interestingly, this problem is closely related to a well-known conjecture of Ryser on packing and covering edges in  $r$ -partite,  $r$ -uniform hypergraphs. This link was first noted by Gyárfás [8] in 1997 and leads to the following natural formulation of the conjecture of Ryser, published in [12], where  $\alpha(G)$  is the size of the largest independent set in the graph  $G$ .

**Conjecture 1** (Ryser (see [12])). *The vertex set of an  $r$ -coloured graph  $G$  can be covered by at most  $(r - 1)\alpha(G)$  monochromatic connected subgraphs.*

In this form, it is clear that Ryser's conjecture implies the *covering version* of the aforementioned conjecture of Erdős, Gyárfás and Pyber about monochromatic connected subgraphs. Although not much is known about Ryser's conjecture in general, a few special cases are understood. The case  $r = 2$  is equivalent to König's classical theorem (see [5], for example), while the case  $r = 3$  was proved by Aharoni [1] in 2001, who built on the earlier advances of Aharoni and Haxell [2]. The conjecture is also known to hold for  $\alpha(G) = 1$  (i.e.  $G$  is a complete graph) and  $r \leq 5$ , as was proved by Gyárfás [8] ( $r = 3$ ), Duchet [6] and Tuza [17] ( $r = 4$ ), and Tuza [17] ( $r = 5$ ).

Following Schelp [16] who suggested several variants of Ramsey-type problems (e.g. the determining the length of the longest monochromatic path in a 2-coloured graphs), we consider variants of the above problems for graphs with large minimum degree. Our first main result proves a conjecture of Bal and DeBiaso [3] about partitioning the vertices of a 2-coloured graph with large minimum degree; recall that  $\delta(G)$  denotes the minimum degree of the graph  $G$ .

**Theorem 2.** *There exists an integer  $n_0$  such that that every 2-coloured graph  $G$  on  $n \geq n_0$  vertices and with minimum degree at least  $\frac{2n-5}{3}$  can be partitioned into two monochromatic connected subgraphs.*

We note that this is a generalisation of the result by Haxell and Kohayakawa [11] mentioned above for two colours, where the complete graph is replaced by a graph with large minimum degree. This result is seen to be sharp by a construction of Bal and DeBiaso [3]; in Section 4 we describe a more general family of examples which shows, in particular, the sharpness of the minimum degree condition in this result. One can think of this result as saying that  $\frac{2n-5}{3}$  is the minimum degree 'threshold'

that guarantees a partition of every 2-colouring into two monochromatic connected subgraphs. It is therefore natural to ask what minimum degree condition on a graph  $G$  guarantees a partition into  $t$  monochromatic connected subgraphs, no matter how the graph is 2-coloured. We conjecture the following.

**Conjecture 3.** *For every  $t$  there exists  $n_0$ , such that for every 2-colouring of a graph  $G$  on  $n \geq n_0$  vertices with  $\delta(G) \geq \frac{2n-2t-1}{t+1}$  there exists a partition of the vertex set into at most  $t$  monochromatic connected subgraphs.*

We support this conjecture by observing an analogous result for *covers* of the vertices by monochromatic components.

**Proposition 4.** *Let  $t$  be integer and let  $G$  be a 2-coloured graph on  $n$  vertices with  $\delta(G) \geq \frac{2n-2t-1}{t+1}$ . Then vertices of  $G$  can be covered by at most  $t$  monochromatic components.*

We also give a construction, showing that the inequality in this proposition (and therefore the conjecture) cannot be improved.

Bal and DeBiasio [3] also considered the problem of covering coloured graphs with monochromatic components of distinct colours. In particular, they conjectured the following.

**Conjecture 5.** *Let  $G$  be an  $r$ -coloured graph on  $n$  vertices with  $\delta(G) \geq (1 - 1/2^r)n$ . Then the vertices can be covered by monochromatic components of distinct colours.*

Again, Bal and DeBiasio provided examples showing that if true, the bound  $(1 - 2^{-r})n$  is best possible. We shall prove Conjecture 5 for  $r = 2, 3$ . The case  $r = 3$  is the most interesting case but we include a short proof of  $r = 2$  for completeness.

**Theorem 6.** *Let  $G$  be a 3-coloured graph on  $n$  vertices with  $\delta(G) \geq 7n/8$ . Then the vertices of  $G$  can be covered by monochromatic components of distinct colours.*

## 1.1 Structure of the paper

We conclude the introduction with a description of the notation that we shall use in this paper. We prove Theorem 2 in Section 2, and prove Theorem 6 in Section 3. We conclude the paper in Section 4 with some final remarks and open problems and give a proof of Proposition 4.

## 1.2 Notation

By an  $r$ -coloured graph, we mean a graph whose edges are coloured with  $r$  colours. When a graph is 2-coloured we call the colours *red* and *blue*; and when it is 3-coloured, we denote the colours by *red*, *blue* and *yellow*.

For a set of vertices  $W$ , we denote by  $N_r(W)$  the set of vertices in  $V(G) \setminus W$  that are adjacent to a vertex in  $W$  by a red edge. If  $x \in V(G)$  is a vertex, we define  $d_r(x) = |N_r(\{x\})|$  which we refer to as the *red degree* of  $x$ . We say that  $y$  is a *red neighbour* of  $x$  if  $xy$  is a red edge. By a *red component* of a graph  $G$ , we mean the *vertex set* of a component in the graph on vertex set  $V(G)$  whose edges are the red edges of  $G$ . We denote the red component that contains  $x$  by  $C_r(x)$ . A *red set*  $U$  is a set of vertices that is connected in red, i.e. the red edges induced by  $U$  form a connected graph.

All the above definitions and notation, that were defined for red, also works for blue or yellow; e.g.  $d_b(x)$  and  $d_y(x)$  are the blue and yellow degrees of  $x$ , respectively, and a blue set is a set of vertices that is connected in blue.

## 2 Partitioning into monochromatic connected subgraphs

In this section we prove Theorem 2.

**Theorem 2.** *There exists an integer  $n_0$  such that every 2-coloured graph  $G$  on  $n \geq n_0$  vertices and with minimum degree at least  $\frac{2n-5}{3}$  can be partitioned into two monochromatic connected subgraphs.*

We note that the minimum degree condition in this theorem cannot be improved; this can be seen by taking  $t = 2$  in Example 21, which we describe in Section 4.

**Proof of Theorem 2.** Throughout this proof, we assume that the number of vertices  $n$  is sufficiently large. Suppose, for a contradiction, that the vertices of  $G$  cannot be partitioned into two monochromatic sets.

**Claim 7.** *There is a blue component of order at most  $(n + 1)/6$ .*

*Proof of Claim 7.* We may assume that there are at least three red components and at least three blue components, as otherwise the vertices may be partitioned into two red sets or two blue sets (recall that a *red set* is defined to be a set of vertices that is connected in red, and similarly for blue), contradicting our assumption. Let  $R$  be a red component of smallest order; so,  $|R| \leq n/3$ .

Let us assume first that  $|R| \leq (n-5)/3$ . Since every vertex in  $R$  sends at least  $(2n-5)/3 - (|R|-1) > (n-|R|)/2$  blue edges outside of  $R$ , every two vertices in  $R$  have a common blue neighbour outside of  $R$ . Hence,  $R$  is contained in a blue component of order at least  $|R| + (2n-5)/3 - (|R|-1) \geq (2n-1)/3$ . Since there are at least three blue components, there is a blue component of order at most  $(n - (2n-1)/3)/2 = (n+1)/6$ .

We now assume that  $(n-4)/3 \leq |R| \leq n/3$ . If every two vertices in  $R$  have a common blue neighbour, then, again,  $R$  is contained in a blue component of order at least  $(2n-1)/3$  and as before there is a blue component of order at most  $(n+1)/6$ . Otherwise, there exist two vertices  $u, w \in R$  whose blue neighbourhoods do not intersect. But both  $u$  and  $v$  have at least  $(n-5)/3$  blue neighbours outside

of  $R$ , and therefore every vertex in  $R \setminus \{u, v\}$  has a common blue neighbour with either  $u$  or  $w$ . It follows that there are two blue components (namely, the components  $C_b(u)$  and  $C_b(w)$ ) whose union has order at least  $|R| + 2(n-5)/3 > n-5$ , hence there is a blue component of order at most 4.  $\square$

**Claim 8.** *There is a red set  $U$  of size at most  $27 \log n$  such that  $|N_r(U)| \geq 2n/3 - 27 \log n$ .*

*Proof of Claim 8.* By the previous claim, there is a blue component  $B$  of order at most  $(n+1)/6$ . Note that every vertex in  $B$  has at least  $(2n-5)/3 - |B|$  red neighbours in  $V(G) \setminus B$ . Fix a vertex  $u \in B$  and let  $N$  be the set of red neighbours of  $u$  outside  $B$ . Every  $w \in B$  has at least the following number of red neighbours in  $N$ .

$$2 \cdot ((2n-5)/3 - |B|) - (n - |B|) = (n-10)/3 - |B| \geq (n-21)/6.$$

Now let  $U'$  be a random subset of  $N$  where each vertex  $w \in N$  belongs to  $U'$ , independently, with probability  $13 \log n/n$ . Let  $I_w$  be the event that  $w$  (where  $w \in B$ ) does not have a red neighbour in  $U'$ . We bound

$$\mathbb{P}\left(\bigcup_{w \in B} I_w\right) \leq |B| \cdot \mathbb{P}(I_w) \leq n \cdot \left(1 - \frac{13 \log n}{n}\right)^{\frac{n-21}{6}} \leq n \cdot e^{-2 \log n} < 1/2.$$

Note that since  $\mathbb{E}(|U'|) \leq 13 \log n$ , we have  $\mathbb{P}(|U'| \geq 26 \log n) \leq 1/2$ , by Markov's inequality. Therefore, there is a choice of  $U' \subseteq N$  such that  $|U'| \leq 26 \log n$  and every vertex in  $B$  is joined by a red edge to some vertex in  $U'$ . We choose  $U = U' \cup \{u\}$ . Note that

$$\begin{aligned} |N_r(U' \cup \{u\})| &\geq |N \setminus U'| + |B \setminus \{u\}| \\ &\geq ((2n-5)/3 - |B| - 26 \log n) + (|B| - 1) \\ &= 2n/3 - 27 \log n. \end{aligned}$$

Hence, the set  $U = U' \cup \{u\}$  satisfies the requirements of Claim 8.  $\square$

Let  $U$  be a red set as in Claim 8 and let  $N = N_r(U)$ . Now choose a maximal sequence of distinct vertices  $x_1, \dots, x_t \in V \setminus (N \cup U)$  so that  $x_i$  has at least  $\log n$  red neighbours in the set  $N \cup \{x_1, \dots, x_{i-1}\}$ , for every  $i \in [t]$ . Then put  $\overline{N} = N \cup \{x_1, \dots, x_t\}$  and write  $W = V(G) \setminus (U \cup \overline{N})$ . Note that every vertex in  $W$  has at most  $\log n$  red neighbours in  $\overline{N}$ .

**Claim 9.**  $|\overline{N}| \leq 2n/3 + 3 \log n + 4$ .

*Proof of Claim 9.* For a contradiction, suppose that  $|\overline{N}| > 2n/3 + 3 \log n + 4$ . We shall deduce that the vertices can be partitioned into a red set and a blue one, a contradiction.

To define the partition, fix  $w \in W$  and let  $X = N_b(w) \cap \overline{N}$ . Let  $S$  be a random subset of  $X$ , obtained by taking each vertex of  $X$  independently with probability  $1/2$ . We claim that, with positive probability,  $(U \cup \overline{N}) \setminus S$  is red and  $W \cup S$  is blue.

To bound the probability that  $W \cup S$  is blue, we consider the probability that every vertex in  $W$  is joined by a blue edge to  $S$  (an event which would imply that  $W \cup S$  is blue). For every  $x, y \in V$  we have  $|N(x) \cap N(y)| \geq n/3 - 10/3$ , hence  $|N(x) \cap N(y) \cap \overline{N}| \geq 3 \log n$ . Since every vertex in  $W$  has at most  $\log n$  red neighbours in  $\overline{N}$ , we have  $|N_b(x) \cap N_b(y) \cap \overline{N}| \geq \log n$ . Therefore the probability that a given  $x \in W$  has no blue neighbours in  $S$  is at most  $2^{-\log n} = 1/n$ . Thus, the expected number of vertices in  $W$  with no edges to  $S$  is smaller than  $1/2$  (note that  $|W| \leq n/3$ ). Hence,  $\mathbb{P}(W \cup S \text{ is blue}) > 1/2$ .

We now estimate the probability that  $(U \cup \overline{N}) \setminus S$  is red. First note that as  $N = N_r(U)$ , we have that  $U \cup N'$  is red for any subset  $N' \subseteq N$ . So it remains to show that the vertices of  $\{x_1, \dots, x_t\} \setminus S$  can be joined, via a red path, to  $U \cup (N \setminus S)$ , with sufficiently high probability. For  $i \in [t]$ , let  $E_i$  be the event that vertex  $x_i$  is joined by a red edge to  $(N \cup \{x_1, \dots, x_{i-1}\}) \setminus S$ . Note that if the event  $E = \bigcap_i^t E_i$  holds,  $(U \cup \overline{N}) \setminus S$  is red. Now, to estimate  $\mathbb{P}(E_i)$ , for  $i \in [t]$ , note that each vertex  $x_i$  has at least  $\log n$  forward neighbours, and the probability that one of these vertices is deleted is at most  $1/2$ . Thus  $\mathbb{P}(E_i) \geq 1 - 2^{-\log n} = 1 - 1/n$ , therefore  $\mathbb{P}((U \cup \overline{N}) \setminus S \text{ is red}) \geq \mathbb{P}(E) > 1/2$ , where the second inequality holds since  $t < n/2$ .

Thus, with positive probability,  $W \cup S$  is blue and  $(U \cup \overline{N}) \setminus S$  is red. In particular, the vertices can be partitioned into a blue set and a red one, a contradiction.  $\square$

**Claim 10.** *There is a vertex of blue degree at most  $90 \log n$ .*

*Proof of Claim 10.* By definition of  $\overline{N}$  and since  $|\overline{N}| \geq 2n/3 - 27 \log n$ , every vertex in  $W$  has at least  $n/3 - 29 \log n$  blue neighbours in  $\overline{N}$ .

Fix a vertex  $w \in W$ . If there is a vertex  $v \in W$  with  $|N_b(v) \cap N_b(w) \cap \overline{N}| < \log n$ , then the two blue components containing  $v$  and  $w$  cover all vertices of  $W$  and all but at most  $62 \log n$  vertices of  $\overline{N}$  (as  $|\overline{N}| \leq 2n/3 + 3 \log n + 4$ , by the previous claim). Since  $|U| \leq 27 \log n$ , it follows that these two components cover all but at most  $90 \log n$  vertices. Recall that there are at least three blue components, hence there is a component of order at most  $90 \log n$ , and any vertex in that component has blue degree at most  $90 \log n$ .

Otherwise, every vertex  $v \in W$  satisfies  $|N_b(v) \cap N_b(w) \cap \overline{N}| \geq \log n$ . As in Claim 9, let  $S$  be a uniformly random subset of  $N_b(w) \cap \overline{N}$ ; we find that, with positive probability,  $(U \cup \overline{N}) \setminus S$  is red and  $W \cup S$  is blue, so the vertices can be partitioned into a red set and a blue one, a contradiction to our assumption.  $\square$

Let  $r$  be a vertex of blue degree at most  $90 \log n$ , which exists by the previous claim. By symmetry, there is a vertex  $b$  of red degree at most  $90 \log n$ . Then  $d_r(r), d_b(b) \geq 2n/3 - 90 \log n - 2$ . Write  $A_1 = N_b(b) \setminus N_r(r)$ ,  $A_2 = N_b(b) \cap N_r(r)$  and  $A_3 = N_r(r) \setminus N_b(b)$ . Then  $|A_2| \geq n/3 - 180 \log n - 4$  and  $|A_1|, |A_3| \leq n/3 + 90 \log n + 2$ .

**Claim 11.** *There is a vertex with no blue neighbours in  $A_1$ , no red neighbours in  $A_3$ , and at most  $2 \log n$  neighbours in  $A_2$ .*

*Proof of Claim 11.* Suppose that the statement does not hold. Let  $\{B, R\}$  be a random partition of  $A_2$ , obtained by putting vertices in  $B$ , independently, with probability  $1/2$ . Then, with positive probability, every vertex in  $G$  has a blue neighbour in  $A_1 \cup B \subseteq N_b(b)$  or a red neighbour in  $A_3 \cup R \subseteq N_r(r)$ . We thus obtain a partition of the vertices into a red set and a blue one, a contradiction.  $\square$

Let  $x$  be a vertex with no blue neighbours in  $A_1$ , no red neighbours in  $A_3$ , and at most  $2 \log n$  neighbours in  $A_2$  (its existence is guaranteed by the previous claim). Then  $|A_2| \leq n/3 + 3 \log n$ , so  $|A_1|, |A_3| \geq n/3 - 95 \log n$ . Furthermore,  $x$  has at least  $n/3 - 100 \log n$  red neighbours in  $A_1$  and at least  $n/3 - 100 \log n$  blue neighbours in  $A_3$ . Write  $A'_1 = A_1 \cap N_r(x)$ ,  $A'_2 = A_2 \setminus N(x)$ , and  $A'_3 = A_3 \cap N_b(x)$  (so  $|A'_1|, |A'_3| \geq n/3 - 100 \log n$  and  $|A'_2| \geq n/3 - 190 \log n$ ).

**Claim 12.** *The vertices  $x$  and  $b$  are in distinct blue components; similarly,  $x$  and  $r$  are in distinct red components.*

*Proof of Claim 12.* Suppose that  $x$  and  $b$  are in the same blue component. Then there is a blue path  $P$  from  $\{x\} \cup A'_3$  to  $\{b\} \cup A'_1 \cup A'_2$ . We may assume that the inner vertices of  $P$  are outside of  $A'_1 \cup A'_2 \cup A'_3 \cup \{x, b\}$ . Hence,  $|P| \leq 400 \log n$ .

Now, let  $\{B, R\}$  be a random partition of  $(A'_2 \cup A'_3) \setminus V(P)$ , obtained by putting each vertex in  $B$ , independently, with probability  $1/2$ . It is easy to see that, with positive probability, every vertex in  $G$  has a red neighbour in  $R$  or a blue neighbour in  $B$ , from which it can be deduced that there is a partition of the vertices into a red set and a blue one, which is a contradiction. Indeed, note that  $P \cup \{x, b\} \cup B$  is a blue set and  $\{r\} \cup R$  is a red set. Thus, we have that  $b$  and  $x$  are in distinct blue components; by symmetry,  $r$  and  $x$  are in different red components.  $\square$

Note that  $|C_b(b)|, |C_r(r)| \geq 2n/3 - 91 \log n$  and  $|C_b(x)|, |C_r(x)| \geq n/3 - 100 \log n$ . Recall that there are at least three blue components. Hence, there is a vertex  $r_1$  which is not in  $C_b(b)$  or in  $C_b(x)$ . It follows that  $d_b(r_1)$  is at most  $191 \log n$ , hence it has red degree at least  $2n/3 - 192 \log n$ , so  $r_1 \in C_r(r)$ . Similarly, there is a vertex  $b_1$  which is not in  $C_r(r)$  or in  $C_r(x)$ , and therefore it must belong to  $C_b(b)$ . We claim that the set  $X = \{b_1, r_1, x\}$  is independent. We cannot have  $r_1 \tilde{x}$  or  $b_1 \tilde{x}$ , for this either contradicts the choice of  $r_1 \notin C_b(x)$  and  $b_1 \notin C_r(x)$  or it contradicts the statement of Claim 12. If we had  $r_1 \tilde{b}_1$  and this edge was coloured red then  $b_1 \in C_r(r)$  which is a contradiction, by definition of  $b_1$ . If  $r_1 b_1$  is coloured blue then we arrive at the contradiction  $r_1 \in C_b(b)$ . Thus  $X$  is independent. So finally, by the minimum degree condition, there must be a vertex  $w$  that is adjacent to all three vertices in  $X$ . Indeed, if no such  $w$  exists, then the number of edges between  $X$  and  $V(G) \setminus X$  is at most  $2(n-3) < 3(2n-5)/3$ , a contradiction. Without loss of generality,  $w$  sends two red edges into  $X$ , implying that two of these vertices in  $X$  belong to the same red component, a contradiction. This completes our proof of Theorem 2.  $\square$

### 3 Covering with monochromatic components of distinct colours

In this section we verify Conjecture 5 for  $r \in \{2, 3\}$ . Most of the difficulty is in the proof for  $r = 3$ , but we include a short proof for  $r = 2$  for completeness. Actually, the  $r = 2$  case (for  $n$  large) already follows from a difficult result of Letzter [13], who showed that when  $\delta(G) \geq 3n/4$ , the vertices can be partitioned into two monochromatic *cycles* of different colours, for every 2-colouring of  $G$ . Before turning to the proofs, we mention the following construction of Bal and DeBiasio [3], which shows that the minimum degree condition in Conjecture 5 cannot be improved.

**Example 13.** Let  $n \geq 2^r$ ; we shall define a graph on vertex set  $[n]$  as follows. Partition  $[n]$ , as equally as possible, into  $2^r$  sets which are indexed by the sequences  $s \in \{0, 1\}^r$ . We write

$$[n] = \bigcup_{s \in \{0, 1\}^r} A(s)$$

and define the following, where  $\mathbb{1} = (1, \dots, 1)$ .

$$E = [n]^{(2)} \setminus \bigcup_{s \in \{0, 1\}^r} \{xy : x \in A(s), y \in A(\mathbb{1} - s)\}.$$

In other words, we include all edges in the graph except for the edges between parts of the partition corresponding to antipodal elements of  $\{0, 1\}^r$ . Now, colour all edges  $xy$ , where  $x \in A(s), y \in A(s')$ , by the first coordinate on which  $s, s'$  agree; e.g. the edge between  $(0, 1, 0, 0)$  and  $(1, 0, 0, 1)$  is coloured 3.

We now show that  $G$  cannot be covered by components of distinct colours. Suppose that it can, and note that the  $i$ -coloured components are of the form  $\bigcup_{s \in S_i} A(s)$  where  $S_i$  is a set of elements that agree on their  $i$ -th coordinate; denote this coordinate by  $a_i$ . It follows that the vertices of  $A((1 - a_1, \dots, 1 - a_r))$  are not covered by any of these components, a contradiction.

We now prove Conjecture 5 for  $r = 2$ .

**Lemma 14.** Let  $G$  be a 2-coloured graph with  $\delta(G) \geq 3n/4$ . Then the vertices of  $G$  can be covered by a red component and a blue component.

**Proof.** We first show that there is a monochromatic component of order greater than  $n/2$ . If  $G$  is red connected we are done. Hence, there exists a red component  $R$  with  $|R| \leq n/2$ . Then, any two vertices  $u, w \in R$  have a common blue neighbour, as  $|N_b(u) \cap N_b(w) \cap R| \geq 2 \cdot (3n/4 - (|R| - 1)) - (n - |R|) > 0$ . So  $R \subseteq C_b(u)$  and  $C_b(u)$  is a blue component of order at least  $3n/4$ , as required.

Without loss of generality, there is a red component  $R$  of order larger than  $n/2$ . Note that there is a vertex  $x$  which is not in  $R$  (otherwise we are done), and  $|N_b(x) \cap R| = |N(x) \cap R| > n/4$ , as  $x$  does not send red edges to  $R$ . In particular,  $|C_b(x) \cap R| > n/4$ . It follows that every vertex sends at least one edge to  $C_b(x) \cap R$  and thus the components  $R$  and  $C_b(x)$  cover the whole graph.  $\square$

We now turn to prove Theorem 6, which is the case of three colours in Conjecture 5.

**Theorem 6.** *Let  $G$  be a 3-coloured graph on  $n$  vertices with  $\delta(G) \geq 7n/8$ . Then the vertices of  $G$  can be covered by monochromatic components of distinct colours.*

**Proof.** We begin with a series of preparatory claims (Claims 15 to 17).

**Claim 15.** *If there are three monochromatic components of distinct colours whose intersection has order at least  $n/8$ , then the vertices can be covered by monochromatic components of distinct colours.*

*Proof of Claim 15.* Suppose that  $R$ ,  $B$  and  $Y$  are red, blue and yellow components, whose intersection  $U = R \cap B \cap Y$  has size at least  $n/8$ . Then, by the minimum degree condition, every vertex not in  $U$  has a neighbour in  $U$ , implying that every vertex in the graph belongs to at least one of  $R$ ,  $B$  and  $Y$ , as required.  $\square$

**Claim 16.** *If there are two monochromatic components of distinct colours whose intersection has order at least  $n/4$ , then the vertices of  $G$  may be covered by monochromatic components of distinct colours.*

*Proof of Claim 16.* Suppose that  $R$  and  $B$  are red and blue components whose intersection  $U = R \cap B$  has size at least  $n/4$ . We show that one of the following holds.

1.  $R \cup B = V(G)$ ;
2. there is a yellow component whose intersection with  $R \cap B$  has size at least  $n/8$ .

Suppose that the first assertion does not hold. Then there is a vertex  $u \notin R \cup B$ . By the minimum degree condition,  $u$  sends at least  $n/8$  edges to  $R \cap B$ , but these edges cannot be red or blue (because  $u \notin R \cup B$ ), hence they are yellow, so by picking  $Y$  to be the yellow component containing  $u$ , the second assertion holds. If the first assertion holds, we are done immediately; otherwise, we are done by Claim 15.  $\square$

**Claim 17.** *If there is a monochromatic component of order at least  $n/2$ , then the vertices can be covered by three monochromatic components of distinct colours.*

*Proof of Claim 17.* As in the proof of Claim 16, we show that one of the following assertions holds, where  $R$  is a red component of order at least  $n/2$ .

1.  $R = V(G)$ ;
2. there are monochromatic components  $B$  and  $Y$  in colours blue and yellow respectively, such that  $R \cup B \cup Y = V(G)$ ;
3. there are monochromatic components  $B$  and  $Y$  in colours blue and yellow respectively, such that  $|R \cap B \cap Y| \geq n/8$ .

Suppose that  $R \neq V(G)$  and let  $u \notin R$ . Consider the blue and yellow components,  $B$  and  $Y$ , containing  $u$ . By the minimum degree condition,  $u$  sends at least  $|R| - n/8$  edges to  $R$ , none of which are red. So  $|(B \cup Y) \cap R| \geq |R| - n/8$ . Suppose that  $R, B$  and  $Y$  do not cover the whole graph. Let  $w \notin R \cup B \cup Y$ , and denote the blue and yellow components containing  $w$  by  $B'$  and  $Y'$ . By the same argument as before,  $|(B' \cap Y') \cap R| \geq |R| - n/8$ , which implies the following.

$$|(B \cup Y) \cap (B' \cup Y') \cap R| \geq |R| - n/4 \geq n/4.$$

Since  $B \cap B' = \emptyset$  and  $Y \cap Y' = \emptyset$ , either  $|B \cap Y' \cap R| \geq n/8$  or  $|B' \cap Y \cap R| \geq n/8$ . This completes the proof that one of the above assertions holds. If one of the first two assertions holds, we are done immediately; and if the third assertion holds, Claim 17 follows from Claim 15.  $\square$

Henceforth, we assume  $G$  cannot be covered by monochromatic components of distinct colours.

**Claim 18.** *There are two monochromatic components of distinct colours of order at least  $3n/8$ .*

*Proof of Claim 18.* We will show that for every pair of colours there is a monochromatic component of order at least  $3n/8$  in one of the two colours; the claim easily follows from this fact. Let the two colours be red and blue. Since  $G$  is not connected in yellow, we may find a partition  $\{X, Y\}$  of the vertices of  $G$  such that no  $X - Y$  edges are yellow. Without loss of generality, at least half the edges between  $X$  and  $Y$  are red; set  $H = G_r[X, Y]$ , denote  $d(x) = d_H(x)$  for any vertex  $x$ , and given an edge  $xy$  in  $H$ , set  $s(xy) = d(x) + d(y)$ . We will show that there is an edge  $xy$  with  $s(xy) \geq 3n/8$ ; note that this would imply the existence of a red component of order at least  $3n/8$ , as required. Put  $e = e(H)$  and, without loss of generality, we assume that  $|X| \leq |Y|$ . We have

$$\begin{aligned} \frac{1}{e} \sum_{xy \in E(H)} s(xy) &= \frac{1}{e} \sum_{xy \in E(H)} (d(x) + d(y)) \\ &= \frac{1}{e} \left( \sum_{x \in X} d(x)^2 + \sum_{y \in Y} d(y)^2 \right) \\ &\geq \frac{1}{e} \left( \frac{(\sum_{x \in X} d(x))^2}{|X|} + \frac{(\sum_{y \in Y} d(y))^2}{|Y|} \right) \\ &= e \left( \frac{1}{|X|} + \frac{1}{|Y|} \right) \\ &\geq \frac{1}{2}|X|(|Y| - n/8) \left( \frac{1}{|X|} + \frac{1}{|Y|} \right) \\ &= \frac{1}{2} \left( |Y| - n/8 + |X| - \frac{|X| \cdot n/8}{n - |X|} \right) \\ &\geq 3n/8. \end{aligned}$$

Indeed, the first inequality follows from the Cauchy-Schwarz inequality; the second follows from the minimum degree condition and the assumption that red is the majority colour between  $X$  and  $Y$ ; and the last inequality follows since  $|X| + |Y| = n$  and the expression  $\frac{|X|}{n-|X|}$  is maximised when  $|X| = n/2$  (as we have the constraint  $|X| \leq n/2$ ).

This chain of inequalities shows that the average value of  $s(xy)$  is at least  $3n/8$ ; in particular, there is a red component of order at least  $3n/8$ , as required.  $\square$

We remark that the idea of double counting  $s(xy)$  as in the proof of the previous claim originated in a paper by Liu, Morris and Prince [14].

By the previous claim, we may assume that  $R$  and  $B$  are red and blue components of order at least  $3n/8$ .

**Claim 19.** *Either  $|R \setminus B| < n/4$  or  $|B \setminus R| < n/4$ .*

*Proof of Claim 19.* Assume that  $|R \setminus B| \geq n/4$  and  $|B \setminus R| \geq n/4$ . Note that every edge between the disjoint sets  $R \setminus B$  and  $B \setminus R$  is yellow. Furthermore, any two vertices in  $B \setminus R$  has a common neighbour in  $R \setminus B$ , and vice versa. Therefore  $B \triangle R$  is contained in a yellow component; in particular, there is a yellow component of order at least  $n/2$ , a contradiction, by Claim 17.  $\square$

By the previous claim, we may assume that  $|B \setminus R| < n/4$ . Hence,  $|B \cup R| = |R| + |B \setminus R| < n/2 + n/4 = 3n/4$ , by Claim 17. Therefore, the set  $W = V(G) \setminus (R \cup B)$  has size larger than  $n/4$ . Since all edges between  $R \cap B$  and  $W$  are yellow, it follows that every two vertices in  $R \cap B$  have a common yellow neighbour in  $W$  and hence  $R \cap B$  is contained in a yellow component. Thus, Claim 15 implies that  $|R \cap B| < n/8$ . It follows that  $|B| = |B \cap R| + |B \setminus R| < 3n/8$ , in contradiction with the choice of  $B$ . This completes the proof of Theorem 6.  $\square$

## 4 Concluding remarks

As possible lines for future research, we remind the reader of Conjecture 5 by Bal and DeBiasio [3]; in this paper we proved this conjecture for  $r \leq 3$ .

**Conjecture 5.** *Let  $G$  be an  $r$ -coloured graph on  $n$  vertices with  $\delta(G) \geq (1 - 1/2^r)n$ . Then the vertices can be covered by monochromatic components of distinct colours.*

Another conjecture stated by Bal and DeBiasio [3] concerns the minimum degree needed to ensure that an  $r$ -coloured graph can be covered by at most  $r$  monochromatic components, whose colours need not to be distinct.

**Conjecture 20.** *Let  $G$  be an  $r$ -coloured graph on  $n$  vertices with  $\delta(G) \geq \frac{r(n-r-1)+1}{r+1}$ . Then the vertices of  $G$  can be covered by at most  $r$  monochromatic components.*

We further recall our Conjecture 3.

**Conjecture 3.** *For every  $t$  there exists  $n_0$ , such that for every 2-colouring of a graph  $G$  on  $n \geq n_0$  vertices with  $\delta(G) \geq \frac{2n-2t-1}{t+1}$  there exists a partition of the vertex set into at most  $t$  monochromatic connected subgraphs.*

In this conjecture, we attempt to determine the minimum degree condition needed to guarantee the existence of a partition of a 2-coloured graph into  $t$  monochromatic connected subgraphs. This is a generalisation of Theorem 2 which determines this condition for a partition into two monochromatic connected sets. To close the section, we prove Proposition 4, a weaker version of Conjecture 3, where instead of partitioning the vertices, we aim to *cover* the vertices.

*Proof of Proposition 4.* We use the link with König's Theorem first noted by Gyárfás [8]. Let  $G$  be a 2-coloured graph with minimum degree at least  $\frac{2n-2t-1}{t+1}$ . Let  $\mathcal{R}$  be the collection of red components (some of which may be singletons, if there are vertices that are not incident with any red edges), and let  $\mathcal{B}$  be the collection of blue edges. Define an auxiliary bipartite graph  $H = (\mathcal{R}, \mathcal{B}, E)$ , where for  $R \in \mathcal{R}$  and  $B \in \mathcal{B}$ , we have  $RB \in E$  if and only if  $R \cap B \neq \emptyset$ .

We claim that there is no matching of size larger than  $t$ . Indeed, suppose that  $\{R_1B_1, \dots, R_{t+1}B_{t+1}\}$  is a matching of size  $t+1$ . Let  $u_i \in R_i \cap B_i$ , for  $i \in [t+1]$  and  $U = \{u_1, \dots, u_{t+1}\}$ . Then the vertices of  $U$  are in distinct red and blue components. In particular,  $U$  is independent, so the number of edges between  $U$  and  $V(G) \setminus U$  is at least  $2n - 2t - 1$ . On the other hand, no vertex sends more than one red edge into  $U$  (and similarly for blue), so every vertex not in  $U$  sends at most two edges into  $U$ . It follows that the number of edges between  $U$  and  $V(G) \setminus U$  is at most  $2(n - t - 1) < 2n - 2t - 1$ , a contradiction.


By König's theorem, which states that in bipartite graphs, the size of a minimum cover equals the size of a maximum matching, it follows that there is a cover  $W$  of size at most  $t$ ; write  $W = \{C_1, \dots, C_t\}$ . We claim that  $V(G) = C_1 \cup \dots \cup C_t$ . Indeed, consider a vertex  $u$  and denote its red and blue components by  $R$  and  $B$ , respectively. Then  $R \cap B \neq \emptyset$ , hence  $RB$  is an edge in  $H$ , so either  $R$  or  $B$  is in  $W$ , which implies that  $u \in C_1 \cup \dots \cup C_t$ , as required. In other words, the vertices of  $G$  can be covered by at most  $t$  monochromatic components.  $\square$

Finally, we note that the restriction on the minimum degree in Proposition 4 (and therefore Conjecture 3) cannot be improved. The special case of this example, where  $t = 2$ , appears in [3] and shows that the minimum degree condition in Theorem 2 is best possible.

**Example 21.** *Let  $U$  be a set of size  $n \geq t + 1$ , and let  $\{X, A_1, \dots, A_{t+1}\}$  be a partition of  $U$ , where  $|X| = t + 1$  and the sizes of  $A_1, A_2, \dots, A_{t+1}$  are as equal as possible; write  $X = \{x_1, \dots, x_{t+1}\}$ . We define a 2-coloured graph  $G$  on vertex set  $U$  as follows.*

- the sets  $A_i$  are cliques, and we colour them arbitrarily;

- we add all possible edges between  $A_i$  and  $A_{i+1}$ , where  $i \in [t]$ , and colour them red if  $i$  is odd, and blue otherwise;
- we add all edges between  $x_i$  and  $A_i \cup A_{i+1}$ , for  $i \in [t+1]$  (addition is taken modulo  $t+1$ ). We colour these edges red if  $i$  is in  $[t]$  and  $i$  is odd; and blue if  $i$  is in  $[t]$  and  $i$  is even. Finally, we colour the edges from  $x_{t+1}$  to  $A_1$  blue, and colour the edges from  $x_{t+1}$  to  $A_{t+1}$  red if  $t$  is even and blue if  $t$  is odd.



**Figure 1:** an illustration of Example 21 for  $t = 2$  and  $t = 3$  (here grey represents red and black represents blue).

An easy calculation shows that  $G$  has minimum degree<sup>1</sup>  $\lceil (2n - 2t - 1)/(t + 1) \rceil - 1$ , and that no two vertices in  $X$  belong to the same monochromatic component; in particular, the vertices of  $G$  cannot be covered by at most  $t$  monochromatic components.

## Acknowledgements

We would like to thank Béla Bollobás for many valuable comments. We would also like to thank Louis DeBiasio for bringing [17] to our attention. The second author would like to acknowledge the support of Dr. Max Rössler, the Walter Haefner Foundation and the ETH Zurich Foundation. The third author would like to thank Trinity College, Cambridge for support.

## References

- [1] R. Aharoni, *Ryser's conjecture for tripartite 3-graphs*, *Combinatorica* **21** (2001), no. 1, 1–4.
- [2] R. Aharoni and P. Haxell, *Hall's theorem for hypergraphs*, *J. Graph Theory* **35** (2000), no. 2, 83–88.
- [3] D. Bal and L. DeBiasio, *Partitioning random graphs into monochromatic components*, *Electronic Journal of Combinatorics* **24** (2017), #P1.18.
- [4] D. Conlon and M. Stein, *Monochromatic cycle partitions in local edge colourings*, *J. Graph Theory* **81** (2016), 134–145.

<sup>1</sup> In fact, we need to be a bit more careful here. Write  $n = a(t + 1) + r$ , where  $a$  and  $r$  are integers and  $0 \leq r \leq t$ . We consider two cases:  $r < \lceil (t + 1)/2 \rceil$  and  $r \geq \lceil (t + 1)/2 \rceil$ . In the former case, it is easy to see that  $\delta(G) = \lceil (2n - 2t - 1)/(t + 1) \rceil - 1$ . In the latter case, note that exactly  $r$  of the sets  $A_i$  have size  $a$ , and the rest have size  $a - 1$ . Then, again, one can check that  $\delta(G) = \lceil (2n - 2t - 1)/(t + 1) \rceil - 1$  if  $|A_i| = a$  for every odd  $i \in [t + 1]$  (which is possible as  $r \geq (t + 1)/2$ ).

- [5] R. Diestel, *Graph theory*, Graduate Texts in Mathematics, Springer-Verlag, New York, 2016.
- [6] P. Duchet, *Représentations, noyaux en théorie des graphes et hypergraphes*, (1979).
- [7] P. Erdős, A. Gyárfás, and L. Pyber, *Vertex coverings by monochromatic cycles and trees*, J. Combin. Theory Ser. B **51(1)** (1991), 90–95.
- [8] A. Gyárfás, *Partition covers and blocking sets in hypergraphs (in Hungarian)*, MTA SZTAKI Tanulmányok **71** (1977).
- [9] ———, *Large monochromatic components in edge colorings of graphs: a survey*, pp. 77–96, Birkhäuser Boston, Boston, MA, 2011.
- [10] A. Gyárfás, M. Ruszinkó, G. N. Sárközy, and A. Szemerédi, *An improved bound for the monochromatic cycle partition number*, J. Combin. Theory Ser. B **96(6)** (2006), 855–873.
- [11] P. E. Haxell and Y. Kohayakawa, *Partitioning by monochromatic trees*, J. Combin. Theory Ser. B **68** (1996), no. 2, 218–222.
- [12] J. R. Henderson, *Permutation decompositions of  $(0,1)$ -matrices and decomposition transversals*, Ph.D. thesis, California Institute of Technology, 1970.
- [13] S. Letzter, *Monochromatic cycle partitions of 2-coloured graphs with minimum degree  $3n/4$* , preprint, arXiv:1502.07736.
- [14] H. Liu, N. Prince, and R. Morris, *Highly connected monochromatic subgraphs of multicoloured graphs*, Journal of Graph Theory **61** (2009), 22–44.
- [15] A. Pokrovskiy, *Partitioning edge-coloured complete graphs into monochromatic cycles and paths*, J. Combin. Theory Ser. B **106** (2014), 70–97.
- [16] R. Schelp, *Some Ramsey-Turán type problems and related questions*, Discr. Math. **312** (2012), 21582161.
- [17] Sz. Tuza, *Some special cases of Ryser’s conjecture*, (1978), manuscript.