

A Hamilton–Jacobi theory for implicit differential systems

Oğul Esen[†], Manuel de León[‡], Cristina Sardón^{*}

Department of Mathematics[†]
Gebze Technical University
41400 Gebze, Kocaeli, Turkey.

Consejo Superior de Investigaciones Científicas[‡]
C/ Nicolás Cabrera, 13–15, 28049, Madrid. SPAIN

Instituto de Ciencias Matemáticas, Campus Cantoblanco^{*}
Consejo Superior de Investigaciones Científicas

Abstract

In this paper, we propose a geometric Hamilton-Jacobi theory for systems of implicit differential equations.

In particular, we are interested in implicit Hamiltonian systems, described in terms of Lagrangian submanifolds of TT^*Q generated by Morse families. The implicit character implies the nonexistence of a Hamiltonian function describing the dynamics. This fact is here amended by a generating family of Morse functions which plays the role of a Hamiltonian. A Hamilton–Jacobi equation is obtained with the aid of this generating family of functions. To conclude, we apply our results to singular Lagrangians by employing the construction of special symplectic structures.

1 Introduction

Implicit differential equations (IDE) do not only arise in purely mathematical frameworks, as in relation with minimizers of integrals in the calculus of variations, or as intermediate steps for the integration of differential algebraic equations [26], they do appear in many various areas in science as well. Their applications are important in relativity, control theory, chemistry, etc. For example, they describe exchanges of matter, energy, or information that vary in space and time.

Unfortunately, differential equations and, particularly, IDE cannot always be solved analytically. For this matter, different mathematical methods

have been precisely designed. It is desirable that these differential equations are reducible to quadratures, and many attempts have been tried through Fuchsian, Lie’s theory, etc. Nowadays, the methods of numerical integration has been increasingly developed.

The Hamilton–Jacobi theory (HJ theory) has been long known as a powerful problem solving tool [1, 2]. It is particularly useful for identifying conserved quantities for a mechanical system, which may be possible even when the mechanical problem itself cannot be solved completely. Therefore, it constitutes an alternative way of finding solutions of Hamilton’s equations. It is equivalent to other classical formulations of mechanics and its roots in variational calculus. The action functions are solutions of Hamilton–Jacobi equation (HJE). It is important to remark that the classical HJ theory only deals with explicit Hamiltonian systems, but, in the literature there exist tons of physical models governed by IDE. Hence, the necessity of constructing a Hamilton–Jacobi theory for implicit systems.

For example, recall the number of theories described by singular Lagrangians in the sense of Dirac–Bergmann [3, 9, 17], including systems appearing in gauge theories [31]. The Euler–Lagrange equations (EL) give rise to differential equations that are implicit, and because of the degeneracy of the Lagrangian they cannot be put in a normal form. Some authors have introduced a geometric formalism for dealing with dynamical systems in their implicit form [43, 46] and a unified approach for the Lagrangian description of (time-independent) constrained mechanical systems is provided through a technique that generates IDE on T^*Q from one-forms defined on the total space of any fiber bundle over TQ [5]. Other authors have designed algorithms following the Dirac–Bergmann prescription, to be able to deal with singular Hamiltonian and Lagrangian theories, see for example, the geometric Gotay–Nester algorithm [22, 23, 24, 25] (see brief description in our Appendix). In the local coordinate formalism, the classical EL displaying conservative or nonconservative force fields or subject to linear or nonlinear nonholonomic constraints, also arise in implicit form from d’Alambert’s principle of virtual work. In the geometric formalism, the corresponding equations should then be expected to arise in implicit form equivalently from a suitable expression of the above principle.

Our main aim is to generalize the geometric Hamilton–Jacobi explained for explicit systems to the realm of implicit systems on T^*Q . We interpret IDE in terms of arbitrary submanifolds of a higher-order tangent bundle, particularly, Lagrangian submanifolds of TT^*Q in the case of implicit Hamiltonian systems (IHS), not necessarily in the horizontal form. As an application, we shall concentrate the problem of Hamilton–Jacobi theory for

singular Lagrangian theories.

Let us summarize the problem in more technical terms. We consider a first-order IDE as a submanifold E of TT^*Q . We project E to TQ by the tangent mapping $T\pi_Q$ to a submanifold $T\pi_Q(E)$ of TQ , which is another IDE on Q . The philosophy of the geometric Hamilton–Jacobi theory is to retrieve solutions of E , provided the solutions of $T\pi_Q(E)$. In similar fashion as the classical Hamilton–Jacobi theorem, in order to lift the solutions in Q to T^*Q , we are still in need of a closed one-form γ on Q , but two ingredients of the theory are missing. One is that the base manifold $C = \tau_{T^*Q}(E)$ is not necessarily the whole T^*Q , but possibly a proper submanifold of it. The second is the nonexistence of a Hamiltonian vector field due to the implicit character of the equations. In the classical theory, the major role of the Hamiltonian vector field is to connect the image space of γ and the submanifold E . To overcome these two difficulties, we need to introduce a auxiliary section σ of the fibration τ_{T^*Q} defined on $C \cap \text{Im } \gamma$ and taking values in E . If, particularly, the dynamics E is a Lagrangian submanifold then, according to generalized Poincaré theorem [7, 27, 38, 52, 55], there exists a Morse family (a family of generating functions) defined on the total space of a smooth bundle linked to TT^*Q by means of a special symplectic structure. A Morse family also establishes a link from the base space T^*Q to E , so that for this particular case, there is no need for an auxiliary section.

The plan of the manuscript is the following: in section 2 we review the fundamentals of Hamiltonian mechanics, Section 3 develops a geometric interpretation of dynamics as Lagrangian submanifolds and their generation with the aid of Morse families of functions. Section 4 illustrates the geometric HJ theory both for IDE and IHS. In section 5, we contemplate the construction of complete solutions. Section 6 concerns applications of our constructed theory to the case of degenerate Lagrangians.

We assume that functional analytic issues related with the present discussion are satisfied in order to highlight the main aspects of our theory. Accordingly, we assume that all manifolds are connected, all mathematical objects are real, smooth and globally defined.

1.1 Notation chart

Let Q be the configuration space, TQ is the tangent bundle, and T^*Q is the cotangent bundle. Consider the tangent and cotangent bundles of TQ and T^*Q , these are the possibilities: TTQ, T^*TQ, TT^*Q and T^*T^*Q . Here we can establish the canonical projections for the first order tangent and cotangent bundles, denoted as $\pi_Q : T^*Q \rightarrow Q$ and $\tau_Q : TQ \rightarrow Q$. Furthermore,

consider the projections, $\pi_{T^*Q} : T^*T^*Q \rightarrow T^*Q$, $\tau_{T^*Q} : TT^*Q \rightarrow T^*Q$, $T\pi_Q : TT^*Q \rightarrow TQ$ and the two last projections $\pi_{TQ} : T^*TQ \rightarrow TQ$ and $\tau_{TQ} : TTQ \rightarrow TQ$. For the last case there is another possibility $T\tau_Q : TTQ \rightarrow TQ$, and both possibilities are related through a diffeomorphism we shall devise in the following lines.

Table 1. **Canonical coordinates and symplectic forms on second-order tangent and cotangent spaces.** Consider Q a mechanical configuration manifold and note that we are assuming summation over repeated indices.

Space	Coordinates	Symplectic forms
Q	q^i	
TQ	(q^i, \dot{q}^i)	
T^*Q	(q^i, p_i)	$\omega_Q = dq^i \wedge dp_i$
TT^*Q	$(q^i, p_i, \dot{q}^i, \dot{p}_i)$	$\omega_Q^T = d\dot{q}^i \wedge dp_i + dq^i \wedge d\dot{p}_i$
TTQ	$(q^i, \dot{q}^i, \delta q^i, \delta \dot{q}^i)$	
T^*T^*Q	$(q^i, p_i, \alpha_i, \beta^i)$	$\omega_{T^*Q} = dq^i \wedge d\alpha_i + dp_i \wedge d\beta^i$
T^*TQ	$(q^i, \dot{q}^i, a_i, b_i)$	$\omega_{TQ} = dq^i \wedge da_i + d\dot{q}_i \wedge db_i$

Let us recall the definition of the pullback bundle, as we refer to it in forthcoming sections. Let (P, π, M) be a fiber bundle and assume the existence of a differential mapping φ from a manifold P to the base manifold M . Define the following product manifold

$$\varphi^*P = \{(n, p) \in N \times P : \varphi(n) = \pi(p)\}$$

and the surjective submersion $\varphi^*\pi$ which simply projects an element in φ^*P to its first factor in N . The triple $(\varphi^*P, \varphi^*\pi, N)$ is called the pullback bundle of (P, π, M) via the mapping φ . This definition can be summarized within

the following commutative diagram.

$$\begin{array}{ccc}
 \varphi^*P & \xrightarrow{\varepsilon} & P \\
 \downarrow \varphi^*\pi & & \downarrow \pi \\
 N & \xrightarrow{\varphi} & M
 \end{array} \tag{1}$$

Here, ε is the projection which maps an element in φ^*P to its second factor P . Let us point out two particular cases which are important for the present work. If φ is a diffeomorphism, φ^*P and P become diffeomorphic. If M is an embedded submanifold of P , and N is an embedded submanifold of M in (1), then it is evident that φ^*P is an embedded submanifold of P . In this case, the map ε plays the role of an embedding.

Henceforth, we refer to a general, arbitrary manifolds by M or N , we will denote fiber bundles by $\pi : P \rightarrow N$, where P is the complete space and N is its projection by π (also M instead of N indistinctly). IDE will be denoted by E generally, and any general Lagrangian submanifold is denoted by S . We will also refer by E to IHS that are Lagrangian submanifolds generated by a Morse families. By F we denote Morse families. In general, P is the total space of a fiber bundle related with a Morse family F and R is the total space of a special symplectic structure.

2 Fundamentals

This section is dedicated for reviewing fundamentals of Hamiltonian mechanics from a geometric viewpoint, and basics of the Tulczyjew's triple. Here, we are setting the notation we shall be using along the paper.

2.1 Geometry of the cotangent bundle

Consider a manifold Q and a cotangent bundle T^*Q with canonical projection $\pi_Q : T^*Q \rightarrow Q$. We denote by (q^i, p_i) the fibered coordinates in T^*Q such that $\pi_Q(q^i, p_i) = (q^i)$. A cotangent bundle is equipped with a canonical one-form θ_Q defined as follows:

$$\langle X_{\alpha_q}, \theta_Q(\alpha_q) \rangle = \langle T\pi_Q(X_{\alpha_q}), \alpha_q \rangle, \tag{2}$$

where $X_{\alpha_q} \in T_{\alpha_q}(T^*Q)$ and $\alpha_q \in T_q^*Q$. In fibered coordinates, the canonical one-form reads $\theta_Q = p_i dq^i$, which is known as the Liouville one-form on

T^*Q . Consider now the two-form $\omega_Q = -d\theta_Q$, namely, $\omega_Q = dq^i \wedge dp_i$. This two-form has the two properties

1. It has maximal rank $2n$, where n is the dimension of Q .
2. $d\omega_Q = 0$

This two-form is called a symplectic two-form. More generally, a symplectic manifold is a pair (M, ω) such that ω has maximal rank and $d\omega = 0$. Therefore, (T^*Q, ω_Q) is a symplectic manifold and ω_Q is the canonical symplectic form on T^*Q .

Given two symplectic manifolds (M_1, ω_1) and (M_2, ω_2) and a map $F : M_1 \rightarrow M_2$, we say that F is a symplectomorphism if $F^*\omega_2 = \omega_1$.

2.2 Hamiltonian dynamics on the cotangent bundle

A Hamiltonian system on T^*Q is determined by the triple (T^*Q, ω_Q, H) , where H being the Hamiltonian function. Geometrically, Hamilton's equations are defined by

$$\iota_{X_H}\omega_Q = dH, \quad (3)$$

where X_H is the Hamiltonian vector field associated with the Hamiltonian function H , and ι_{X_H} is the inner contraction operator.

In Darboux's coordinates (q^i, p_i) on T^*Q , with $i = 1, \dots, n$ for an n -dimensional configuration manifold Q . The canonical one-form reads $\theta_Q = p_i dq^i$ and the symplectic two-form turns out to be $\omega_Q = dq^i \wedge dp_i$. In this local picture, the Hamiltonian vector field X_H is written as

$$X_H = \frac{\partial H}{\partial p_i} \frac{\partial}{\partial q^i} - \frac{\partial H}{\partial q^i} \frac{\partial}{\partial p_i} \quad (4)$$

whereas the Hamilton's equations (3) turn out to be

$$\dot{q}^i = \frac{\partial H}{\partial p_i}, \quad \dot{p}_i = -\frac{\partial H}{\partial q^i}. \quad (5)$$

Note that, the Hamilton's equations are explicit by definition. Hence, an IDE cannot be recast in the classical Hamiltonian formalism presented in (5). To deal with IHS, we shall redefine Hamiltonian systems in a more abstract framework, as we shall present in the forthcoming sections.

2.3 Geometry of the tangent bundle

Consider the manifold Q and its tangent bundle TQ together with its tangent bundle canonical projection $\tau_Q : TQ \rightarrow Q$. We consider fibered coordinates (q^i, \dot{q}^i) such that $\tau_Q(q^i, \dot{q}^i) = (q^i)$. Given a function $f : Q \rightarrow \mathbb{R}$, we define its complete lift f^T to TQ as the function:

$$f^T(v_q) = df(q)(v_q) \in \mathbb{R}. \quad (6)$$

In local fibered coordinates,

$$f^T(q^i, \dot{q}^i) = \dot{q}^i \frac{\partial f}{\partial q^i}. \quad (7)$$

Given a tangent vector $v_q \in T_q Q$ with components (q^i, v^i) , we define its vertical lift v_q^V for a point $w_q \in T_q Q$ by

$$v_q^V = \left. \frac{d}{dt} \right|_{t=0} (w_q + tv_q) \quad (8)$$

If X is vector field on Q , then its vertical lift to TQ is the vector field

$$X^V(w_q) = (X(q))_{w_q}^V \in T_{w_q}(TQ) \quad (9)$$

for all w_q in TQ . Now, consider the flow $\phi_t : Q \rightarrow Q$ of a vector field X . We define the complete lift X^T of X to TQ as the generator of the tangent flow $T\phi_t : TQ \rightarrow TQ$. We are assuming, for simplicity, that the flow generated by X is complete, but the construction is still valid in general. If $X = X^i \frac{\partial}{\partial q^i}$, a direct computation shows that

$$X^T = X^i \frac{\partial}{\partial q^i} + \dot{q}^i \frac{\partial X^i}{\partial q^i} \frac{\partial}{\partial \dot{q}^i}. \quad (10)$$

Consider now a k -form ω on Q . We define its complete lift ω^T to TQ a k -form characterized by:

$$\omega_Q(X_1^T, \dots, X_n^T) = \omega_Q(X_1, \dots, X_n)^T \quad (11)$$

The following identity follows from a direct computation $d(\omega^T) = (d\omega)^T$. We are particularly interested in the case of lifts of symplectic forms on Q . Therefore, if ω_Q is symplectic on Q , then ω_Q^T is a symplectic form on TQ . Indeed,

$$\text{rank}(\omega_Q^T) = 2 \text{rank}(\omega_Q) \quad (12)$$

and $d(\omega^T) = (d\omega)^T = 0$.

2.4 Submanifolds of symplectic manifolds

Let (M, ω) be a symplectic manifold, and N be a submanifold of M . We define the symplectic orthogonal complement of TN as the set of tangent vectors

$$TN^\perp = \{u \in TM \mid \omega(u, v) = 0, \forall v \in TN\}. \quad (13)$$

Note that, the dimension of the tangent bundle TM is the sum of the tangent bundle TN and its symplectic orthogonal complement TN^\perp . We list some of the important cases.

- N is called an isotropic submanifold of M if $TN \subset TN^\perp$. In this case, the dimension of N is less or equal to the half of the dimension of M .
- N is called a coisotropic submanifold of M if $TN^\perp \subset TN$. In this case, the dimension of N is greater or equal to the half of the dimension of M .
- N is called a Lagrangian submanifold of M if N is a maximal isotropic subspace of (TM, ω) . That is, if $TN = TN^\perp$. In this case, the dimension of N is equal to the half of the dimension of M .
- TN is symplectic if $TN \cap TN^\perp = 0$. In this case, (N, ω_N) is a symplectic manifold.

A diffeomorphism between two symplectic manifolds is called a symplectomorphism if it preserves the symplectic structures. Under a symplectomorphism, the image of a Lagrangian (isotropic, coisotropic, symplectic) submanifold is Lagrangian (resp. isotropic, coisotropic, symplectic) submanifold.

2.5 The complete lift of ω_Q

Consider the canonical symplectic manifold (T^*Q, ω_Q) . The tangent bundle TT^*Q of T^*Q carries a symplectic two-form ω_Q^T that derives from two potential one-forms, denoted by $\theta_1 = i_T \omega_Q$ and $\theta_2 = d_T \theta_Q$. These one-forms are defined by the canonical forms ω_Q and θ_Q , respectively. The definition of the derivation i_T is the manifestation of the double vector bundle structure of TTQ on TQ , and explicitly given by

$$i_T \omega_Q(X) = \omega_Q(\tau_{T^*Q}(X), T\tau_Q(X)).$$

The derivation d_T is the commutator $[d, i_T]$.

Accordingly, in the local coordinate chart $(q^i, p_i; \dot{q}^i, \dot{p}_i)$, the potential one-forms are computed to be

$$\theta_1 = i_T \omega_Q = \dot{p}_i dq^i - \dot{q}^i dp_i, \quad \theta_2 = d_T \theta_Q = \dot{p}_i dq^i + p_i d\dot{q}^i. \quad (14)$$

The exterior derivatives of these one-forms are the same and define the symplectic two-form

$$\omega_Q^T = d\dot{p}_i \wedge dq^i + dp_i \wedge d\dot{q}^i \quad (15)$$

is known as the complete lift to the tangent space of the symplectic two-form [49, 53]. Note that, the difference $\theta_2 - \theta_1$ is an exact one-form. Actually, it is the exterior derivative of coupling function of the Legendre transformation between the Lagrangian and Hamiltonian formalisms. Further, existence of two potential one-forms for ω_Q^T leads to the existence of a Tulczyjew's triple, which is exhibited in the following subsection.

2.6 Tulczyjew's triple

We now consider a particular kind of symplectic manifolds introduced by Tulczyjew in [48, 49, 50, 51, 52, 53].

A special symplectic manifold is a quintuple (R, N, τ, θ, A) where $\tau : R \rightarrow N$ is a fiber bundle, θ is a one-form on R and $A : R \rightarrow T^*N$ is a diffeomorphism such that $\pi = \pi_N \circ A$ and $\theta = A^* \theta_N$. Since $(T^*N, \omega_N = -d\theta_N)$ is a symplectic manifold, then $(R, \omega = -d\theta)$ is symplectic too and $A^* \omega_N = \omega$, therefore, (R, ω) and (T^*N, ω_N) are symplectomorphic. Consider the following diagram.

$$\begin{array}{ccc} R & \xrightarrow{A} & T^*N \\ & \searrow \tau & \swarrow \pi_N \\ & & N \end{array} \quad (16)$$

Tulczyjew's symplectic space (TT^*Q, ω_Q^T) admits two special symplectic structures. Let us study them.

The non-degeneracy of the canonical symplectic structure ω_Q on T^*Q leads to the existence of the following diffeomorphism

$$\beta_Q : TT^*Q \mapsto T^*T^*Q : X \mapsto \iota_X \omega_Q. \quad (17)$$

The mapping β_Q is actually a symplectomorphism if the iterated cotangent bundle T^*T^*Q is equipped with the canonical symplectic two-form ω_{T^*Q} . In coordinates, we have that

$$\beta_Q(q^i, p_i, \dot{q}^i, \dot{p}_i) = (q^i, p_i, -\dot{p}_i, \dot{q}^i). \quad (18)$$

It is a matter of a direct calculation to prove that, the quintuple

$$(TT^*Q, T^*Q, \tau_{T^*Q}, \theta_1, \beta_Q)$$

is a special symplectic manifold. Here, θ_1 is the differential one-form defined in (14).

We define a canonical diffeomorphism $\alpha_Q : TT^*Q \rightarrow T^*TQ$ as follows. First of all, let us recall that there exists a canonical involution $S_Q : TTQ \rightarrow TTQ$ locally given by

$$S_Q(q^i, \dot{q}^i, \delta q^i, \delta \dot{q}^i) = (q^i, \delta q^i, \dot{q}^i, \delta \dot{q}^i) \quad (19)$$

(see [19]). Now, given $v \in TT^*Q$ we define $\alpha_Q(v) \in T^*TQ$ as

$$\langle w, \alpha_Q(v) \rangle = \frac{d}{dt} \langle \gamma, \xi \rangle |_{t=0} \quad (20)$$

where $\gamma : \mathbb{R} \rightarrow TQ$ and $\xi : \mathbb{R} \rightarrow T^*Q$ are curves such that $j^1 \circ \gamma = S_Q(w)$, $j^1 \circ \xi = v$ and $\tau_Q \circ \gamma = \pi_Q \circ \xi$. In local coordinates, we obtain

$$\alpha_Q(q^i, p_i, \dot{q}^i, \dot{p}_i) = (q^i, \dot{q}^i, \dot{p}_i, p_i). \quad (21)$$

The mapping α_Q is a symplectomorphism if the iterated cotangent bundle T^*TQ is equipped with the canonical symplectic two-form ω_{T^*TQ} . Then it becomes immediate to prove that

$$(TT^*Q, TQ, T\pi_Q, \theta_2, \alpha_Q)$$

is a special symplectic manifold. Here, θ_2 is the differential one-form defined in (14).

As a result, we have derived two special symplectic structures for the symplectic manifold (TT^*Q, ω_Q^T) . Tulczyjew's triple is the combination of these two special symplectic structures in one commutative diagram as given below.

$$\begin{array}{ccccc}
 T^*TQ & \xleftarrow{\alpha_Q} & TT^*Q & \xrightarrow{\beta_Q} & T^*T^*Q \\
 \searrow \pi_{TQ} & & \searrow T\pi_Q & & \searrow \pi_{T^*Q} \\
 & & TQ & & T^*Q
 \end{array} \quad (22)$$

3 Lagrangian Submanifolds

In this and following subsections, we will summarize the theory of Morse families and Lagrangian submanifolds generated by them. We refer an incomplete list [8, 38, 50, 51, 52, 55] for more detailed discussions.

Let (M, ω) a symplectic manifold. A sufficient condition for a submanifold $N \subset M$ to be Lagrangian is $TN = TN^\perp$. If N is an isotropic subspace of a symplectic manifold (M, ω) , then N is Lagrangian if and only if $\dim N = \dim M/2$. For different types of manifolds (Poisson, Nambu–Poisson, etc), the definition of a Lagrangian submanifold has been accommodated to its background. See for example [35, 36, 38].

Two principal examples of Lagrangian submanifolds of the symplectic phase space T^*Q are the fibers of the canonical cotangent bundle projection and the image space of closed one-forms $\gamma : Q \rightarrow T^*Q$. The latter case includes the zero section of the projection $T^*Q \mapsto Q$ as well. This statement is the well-known Weinstein tubular neighborhood theorem [55].

3.1 Morse families

Consider a differentiable fibration (P, π, N) . A real valued function F on the total space P can intuitively be understood as a family of functions parameterized by the coordinates of the fibers of π . The critical set of F is defined by

$$Cr(F, \pi) = \{z \in P : \langle dF(z), V \rangle = 0, \forall V \in V_z P\} \quad (23)$$

and is a submanifold of P . The dimension of $Cr(F, \pi)$ is equal to the dimension of N . Here, VP is the vertical bundle on P consisting of vertical vectors projecting to the zero section of TN under the mapping $T\pi$. We define a bilinear mapping

$$\begin{aligned} W(F, z) &: V_z P \times T_z P \rightarrow \mathbb{R} \\ &: (v, w) \rightarrow D^{(1,1)}(F \circ \chi)(0, 0), \end{aligned} \quad (24)$$

where $\chi : \mathbb{R}^2 \rightarrow P$ is the mapping such that the vector v is obtained by taking the derivative of χ with respect to its first entry at $(0, 0)$ and the vector w is obtained by taking the derivative of χ with respect to its second entry at $(0, 0)$.

A family of functions F defined on the total space of the fibration (P, π, N) is said to be regular if the rank of the matrix $W(F, z)$ defined in (24) is the same at each $z \in Cr(F, \pi)$. A family of functions F is said to

be a Morse family (or an energy function) if the rank of $W(F, z)$ is maximal at each $z \in Cr(F, \pi)$.

Let us write the requirement of being a Morse family in terms of local coordinates. Assume that the dimension of the manifold N is n with coordinates (q^i) , the dimension of a fiber is k with coordinates (λ^a) . The function F is called a Morse family if the rank of the $n \times (n + k)$ -matrix

$$\begin{pmatrix} \frac{\partial^2 F}{\partial q^i \partial q^j} & \frac{\partial^2 F}{\partial q^i \partial \lambda^a} \end{pmatrix} \quad (25)$$

is maximal.

3.2 Lagrangian submanifolds generated by Morse families

A Morse family F on the smooth bundle (P, π, N) generates a Lagrangian submanifold

$$S_{T^*N} = \{w \in T^*N : T^*\pi(w) = dF(z)\} \quad (26)$$

of (T^*N, ω_N) . In this case, we say that S_{T^*N} is generated by the Morse family F . Note that, in the definition of S_{T^*N} , there is an intrinsic requirement that $\pi(z) = \pi_{T^*N}(w)$. Here, we are presenting the following diagram in order to summarize the discussion.

$$\begin{array}{ccc} \mathbb{R} & \xleftarrow{F} & P & & T^*N & & \\ & & \downarrow \pi & & \downarrow \pi_N & & \\ & & N & \xlongequal{\quad} & N & & \end{array} \quad (27)$$

In order to exhibit the structure of the submanifold S_{T^*N} , define a fiber preserving mapping κ from the critical set $Cr(F, \pi)$ of the Morse family F to the cotangent bundle T^*N according to the requirement

$$\langle \kappa(z), Z_N \rangle = \langle dF, Z_P \rangle, \quad (28)$$

which is valid for all π -related vector fields $Z_N \in \mathfrak{X}(N)$ and $Z_P \in \mathfrak{X}(P)$. Note that, this mapping is an immersion and that $\dim(S_{T^*N})$ equals to $\dim(Cr(F, \pi)) = n$. A direct calculation shows that the image space of κ is the Lagrangian submanifold S_{T^*N} defined in (26).

Let N be an immersed submanifold of Q , and T_N^*Q denote the inverse image $\pi_Q^{-1}(N)$ in the cotangent bundle T^*Q . We define the mapping

$$\xi : T_N^*Q \rightarrow T^*N \quad (29)$$

by requiring that the equality

$$\langle \xi(p), Z_N(n) \rangle = \langle p, Z_N(n) \rangle$$

holds for each $Z_N \in \mathfrak{X}(N)$, where $n = \pi_Q(p)$ and $p \in T_N^*Q$. So, ξ is the identity if $p \in N$, that is, if $p \in T^*N$. Consider the canonical injection $i : T_N^*Q \rightarrow T^*Q$. If S_{T^*N} is a Lagrangian submanifold of T^*N then the preimage $i \circ \xi^{-1}(S_{T^*N})$ is a Lagrangian submanifold S_{T^*Q} of T^*Q . If, particularly, the Lagrangian submanifold S_{T^*N} is generated by a Morse family F on the fiber bundle (P, π, N) then, using the same terminology, we say that S_{T^*Q} is generated by the Morse family F . We are presenting the following diagram.

$$\begin{array}{ccc} \mathbb{R} & \xleftarrow{F} & P & & T^*Q & \\ & & \downarrow \pi & & \downarrow \pi_Q & \\ & & N & \xrightarrow{\quad} & Q & \end{array} \quad (30)$$

Let us try to see this more explicitly in the following way. For every point $p \in T^*Q$ such that $\pi_N(p) \in N$, we can find a point $z \in P$ satisfying $\pi(z) = \pi_N(p)$. Then, for every vector $Z_P(z)$, we have that $T\pi \circ Z_P$ is a vector field on N , and hence a vector field on Q . Then, the elements of the Lagrangian submanifold S_{T^*Q} are defined by the requirement

$$\langle p, T\pi \circ Z_P(z) \rangle = \langle dF(z), Z_P(z) \rangle,$$

where F is a Morse family on P .

If the local coordinates (q^i, λ^a) are considered, then the Lagrangian submanifold generated by the Morse family F is defined by

$$S_{T^*Q} = \left\{ \left(q^i, \frac{\partial F}{\partial q^i}(q, \lambda) \right) \in T^*Q : \frac{\partial F}{\partial \lambda^a}(q, \lambda) = 0 \right\}. \quad (31)$$

Note that, we are not distinguishing here the base manifold N from Q .

3.3 Lagrangian submanifolds of special symplectic structures

Let $(R, N = Q, \tau, \omega = -d\theta, A)$ be a special symplectic structure for a symplectic manifold (R, ω) with symplectomorphism A . If S_{T^*Q} is a Lagrangian submanifold of T^*Q , then its pre-image $S = A^{-1}(S_{T^*Q})$ under the symplectomorphism A is a Lagrangian submanifold of the symplectic manifold (R, ω) . If the Lagrangian submanifold S_{T^*Q} is generated by the Morse family F as presented in the diagram (30), then we say that S is generated

by the Morse family F as well. To illustrate this, we draw the following diagram.

$$\begin{array}{ccccc}
 \mathbb{R} & \xleftarrow{F} & P & & T^*Q & \xleftarrow{A} & R \\
 & & \downarrow \pi & & \downarrow \pi_Q & \nearrow \tau & \\
 & & N & \xrightarrow{\quad} & Q & &
 \end{array} \tag{32}$$

Let us record here some special cases of the diagram for future reference.

- The simplest case occurs if A is the identity mapping (no special symplectic manifold) on T^*Q , $P = N$ (no Morse family), and the submanifold $N = Q$ (no constraints on Q). Then, we have a function F (not a family) on Q and its exterior derivative is a Lagrangian submanifold of T^*Q . Then, S turns out to be the image space of a closed one-form $dF : Q \rightarrow T^*Q$. This case includes the zero section of the projection $T^*Q \mapsto Q$ as well [55].
- If A is the identity mapping on T^*Q (no special symplectic manifold), $P = N$ (no Morse family). Then we have a function F (not a family) on a submanifold of N of Q , and the Lagrangian submanifold

$$S_{T^*Q} = \{p \in T^*Q : \pi_Q(p) \in N, \langle Z, \theta_Q(p) \rangle = \langle T\pi(Z), dF \rangle\} \tag{33}$$

for any $Z \in T_p T^*N$ such that $T\pi(Z) \in TN \subset TQ$, [31].

- Let $P = N$ (no Morse family) and the submanifold $N = Q$ (no constraints on Q). Instead, consider the existence of a non-trivial special symplectic structure (P, N, π, ω, A) . Then, a Lagrangian submanifold S of (P, ω) is defined by the pre-image of dF under the isomorphism A , that is

$$S = A^{-1}(dF) = \{y \in Y : \langle y, u \rangle = \langle dE, T\tau(u) \rangle, \forall u \in T_y Y\} \tag{34}$$

where E is defined on Q .

- Let $P = N$ (no Morse family), N be a proper submanifold of Q , (R, N, τ, θ, A) be a special symplectic manifold. Then the set

$$S = \{y \in R : \tau(y) \in N, \langle \theta, u \rangle = \langle dF, T\tau(u) \rangle, \forall u \in T_y Y\} \tag{35}$$

is a Lagrangian submanifold of $(R, -d\theta)$, and said to be generated by the function $F : N \rightarrow \mathbb{R}$. We cite chapter 7 of [31] for a proof of this statement in a more general framework.

3.4 Lagrangian submanifolds of Tulczyjew's symplectic space

Assume that a submanifold E of TT^*Q is defined in terms of the constraint functions $\Phi^A : TT^*Q \rightarrow \mathbb{R}$,

$$\Phi^A(q^i, p_i, \dot{q}^i, \dot{p}_i) = 0.$$

If E is an IHS, then it must be a Lagrangian submanifold of the symplectic space (TT^*Q, ω_Q^T) and the number of constraints must be $2n$ assuming that the dimension of Q is n . Note that, in this case, the Poisson brackets of the constraint functions must vanish [38],

$$\{\Phi^A, \Phi^B\} = 0. \quad (36)$$

Here, the Poisson bracket in (36) is the one induced by the Tulczyjew's symplectic two-form ω_Q^T in the subsection (2.5). This Poisson bracket can be computed as

$$\{f, g\} = \omega_Q^T(X_f, X_g) = \frac{\partial f}{\partial \dot{p}_i} \frac{\partial g}{\partial \dot{q}^i} - \frac{\partial g}{\partial \dot{p}_i} \frac{\partial f}{\partial \dot{q}^i} + \frac{\partial f}{\partial p_i} \frac{\partial g}{\partial \dot{q}^i} - \frac{\partial g}{\partial p_i} \frac{\partial f}{\partial \dot{q}^i}. \quad (37)$$

The image of a Hamiltonian vector field is a Lagrangian submanifold of TT^*Q . Conversely, if a Lagrangian submanifold of TT^*Q is the image of a vector field on T^*Q , then this vector field is a (at least locally) Hamiltonian vector field. To see this, we present the following calculations. Assume that E is a Lagrangian submanifold of TT^*Q and there exists a vector field X satisfying $E = \text{Im}(X)$. In Darboux's coordinates, we write X as

$$X = \phi^i(q, p) \frac{\partial}{\partial \dot{q}^i} + \phi_i(q, p) \frac{\partial}{\partial p_i}, \quad (38)$$

where ϕ^i and ϕ_i are arbitrary functions on T^*Q . This local picture enables us to write the fiber variables (\dot{q}, \dot{p}) in terms of the functions of the base variables (q, p) , and the following definition of the Lagrangian submanifold of E given by

$$E = \{(q^i, p_j; \dot{q}^i, \dot{p}_i) \in TT^*Q : \dot{q}^i - \phi^i(q, p) = 0, \dot{p}_i + \phi_i(q, p) = 0\}. \quad (39)$$

Since E is a Lagrangian submanifold, the Poisson brackets of the defining equations in (39) must be identically zero. Here, the Poisson bracket is the one in (37). This requirement dictates n^2 number of conditions

$$\frac{\partial \phi_j}{\partial p_i} - \frac{\partial \phi^i}{\partial q^j} = 0.$$

These conditions are the same with the conditions for closure of the one-form $\phi = \phi_j dq^j + \phi^i dp_i$. Locally, every closed one-form is exact. This implies that there exists a Hamiltonian function H depending on (q, p) satisfying $dH = \phi$. As a result, the system (39) can be written as in form of the Hamilton's equations (4).

A stronger result follows from the generalized Poincaré lemma [7, 27, 38, 52, 55]. The generalized Poincaré guarantees that for a Lagrangian submanifold E of TT^*Q , whether it is explicit or implicit, there exists a Morse family of functions generating E . This theorem is also known as Maslov-Hörmander theorem [6, 8, 11, 12]. We record here a Morse family generating the Lagrangian submanifold E as follows

$$\begin{array}{ccccc}
 \mathbb{R} & \xleftarrow{F} & P & & T^*T^*Q \xleftarrow{\beta_Q} TT^*Q \\
 & & \downarrow \pi & \swarrow \pi_{T^*Q} & \downarrow \\
 & & N & \xrightarrow{\quad} & T^*Q \\
 & & & & \swarrow \tau_{T^*Q}
 \end{array} \quad (40)$$

where N is a submanifold of the cotangent bundle T^*Q , and F is a Morse family defined on the total space P of the smooth bundle (P, π, N) . On the local chart, the Lagrangian submanifold E generated by F is computed to be

$$E = \left\{ \left(q^i, p_i; \frac{\partial F}{\partial p_i}, -\frac{\partial F}{\partial q^i} \right) \in TT^*Q : \frac{\partial F}{\partial \lambda^a}(q, p, \lambda) = 0 \right\}. \quad (41)$$

Note that, if the Morse family F does not depend on the fiber variables (λ) , then E becomes explicit.

Let us comment on a particular case. Consider the following constrained Hamiltonian (Dirac) system

$$\dot{q}^i = \frac{\partial H}{\partial p_i}(q, p), \quad \dot{p}_i = -\frac{\partial H}{\partial q^i}(q, p), \quad \Phi^\alpha(q, p) = 0, \quad (42)$$

where Φ^α , for $\alpha = 1, \dots, k$, are real valued functions defining a constraint submanifold $M \subset T^*Q$ [41]. Note that, we prefer to denote the generator by H to highlight that it is a Hamiltonian function in the classical sense. Diagrammatically, we have the following picture generating the dynamics

$$\begin{array}{ccccc}
 \mathbb{R} & \xleftarrow{F} & P & & T^*T^*Q \xleftarrow{\beta_Q} TT^*Q \\
 & & \downarrow & \swarrow \pi_{T^*Q} & \downarrow \\
 & & T^*Q & \xlongequal{\quad} & T^*Q \\
 & & & & \swarrow \tau_{T^*Q}
 \end{array} \quad (43)$$

where P is the product manifold $T^*Q \times \mathbb{R}^k$ and the Morse family is given by $F(q, p, \lambda) = H(q, p) + \lambda_\alpha \Phi^\alpha(q, p)$.

4 Hamilton–Jacobi theory for implicit systems

4.1 Geometry of the Hamilton–Jacobi equation

A HJE is a partial differential equation for a generating function $S(q^i, t)$ on Q and the time t given by

$$\frac{\partial S}{\partial t} + H\left(q^i, \frac{\partial S}{\partial q^i}\right) = 0. \quad (44)$$

Note that, the generalized momenta do not appear in (44), except as derivatives of S . This equation is a necessary condition describing the external geometry in problems of calculus of variations. Hamilton’s principal function $S = S(q^i, t)$, which is the solution of the HJE, and the classical function H are both closely related to the classical action

$$S = \int L dt.$$

The function S is a generating function for a family of symplectic flows that describes the dynamics of the Hamilton equations. If the generating function is separable in time, then we can make an ansatz $S(q^i, t) = W(q^i) - Et$, where E is the total energy of the system. Then, HJE in (44) reduces to

$$H\left(q^i, \frac{\partial W}{\partial q^i}\right) = E. \quad (45)$$

Physically, this constant is identified with the energy of the mechanical system.

Let us summarize the geometric Hamilton-Jacobi theory. For this, first consider a Hamiltonian vector field X_H on T^*Q , and a one-form section γ on Q . We define a vector field X_H^γ on Q by

$$X_H^\gamma = T\pi \circ X_H \circ \gamma. \quad (46)$$

This definition implies the commutativity of the following diagram.

$$\begin{array}{ccc} T^*Q & \xrightarrow{X_H} & TT^*Q \\ \downarrow \pi_Q & & \downarrow T\pi_Q \\ Q & \xrightarrow{X_H^\gamma} & TQ \end{array} \quad (47)$$

We enunciate the following theorem.

Theorem 1. *The closed one-form $\gamma = dW$ on Q is a solution of the Hamilton–Jacobi equation (45) if the following conditions are satisfied:*

1. *The vector fields X_H and X_H^γ are γ -related, that is*

$$T\gamma(X^\gamma) = X \circ \gamma. \quad (48)$$

2. *Or equivalently, if the following equation is fulfilled*

$$d(H \circ \gamma) = 0.$$

Proof. We refer [13, 32, 33] for proof of this theorem. □

The first item in the theorem says that if $(q^i(t))$ is an integral curve of X_H^γ , then $(q^i(t), \gamma_j(q(t)))$ is an integral curve of the Hamiltonian vector field X_H , hence a solution of the Hamilton's equations (4). Such a solution of the Hamiltonian equations is called horizontal since it is on the image of a one-form on Q . In the local picture, the second condition implies that exterior derivative of the Hamiltonian function on the image of γ is closed, that is, $H \circ \gamma$ is constant

$$H(q^i, \gamma_j(q)) = \text{cst}. \quad (49)$$

Under the assumption that γ is closed, we can find (at least locally) a function W on Q satisfying $dW = \gamma$. After the substitution of this, equation (49) retrieves the HJE (45).

4.2 Implicit differential equations

A first-order differential equation on a manifold M is a submanifold of its tangent bundle. The submanifold is said to be an explicit differential equation (EDE) if it is image of a vector field defined on M , otherwise, it is called implicit (IDE) [27, 39, 41, 45, 47]. Consider a local coordinate system (q^i) on M , and the induced coordinates (q^i, \dot{q}^i) on its tangent bundle TM . An IDE can be written in form

$$\dot{q}^i = g^i(q, \lambda), \quad f^a(q, \lambda) = 0 \quad (50)$$

for i with certain values running from $1, \dots, n$ and $a = 1, \dots, k$. Here, g^i and f^a are real valued differentiable functions on $(q, \lambda) \in M \times \mathbb{R}^k$. Note that f^a 's form a matrix of maximal rank

$$\text{rank} \left(\frac{\partial f^a}{\partial q^i}, \frac{\partial f^a}{\partial \lambda^b} \right) = k. \quad (51)$$

This implies that only certain \dot{q}^i 's are expressible as in (50) depending on certain (q^j, λ^a) for some a 's equal to $k \leq n$.

A solution of an IDE is a curve ϕ on M satisfying that the tangent vectors $\dot{\phi}(t)$ belong to E for all t . The submanifold E is called *integrable* if for all vectors $v \in E$, there exists a solution ϕ satisfying $v = \dot{\phi}(t)$ for some t . Equivalently, we may define ‘‘integrability’’ of an IDE without referring to the solution curves as follows. We say that the IDE is integrable if the restriction of the tangent bundle projection τ_M to E is surjective summersion and if $E \subset T(\tau_M(E))$.

In principle, an IDE is not necessarily integrable. For example, consider the following case [47].

Example 1. Let $E \subset TT^*Q$ be a system of IDE, where Q is coordinated by q and T^*Q by (q, p) , defined by

$$E = \{(q, p, \dot{q}, \dot{p}) \in TT^*Q, q^2 + p^2 + (\dot{q} + 1)^2 + \dot{p} = k\} \quad (52)$$

is not integrable in points $q^2 + p^2 = k$, $\dot{q} = -1$ and $\dot{p} = 0$ with $q \neq 0$.

Nonetheless, we can develop an algorithm to extract its integrable part [47]. This algorithm works as follows. We denote the projection of the submanifold E onto M by C . Each step of the algorithm, there is a fiber bundle (E^k, τ^k, C^k) consisting of two submanifolds $E^k \subset E$ and $C^k \subset C$, and a (surjective summersion) projection $\tau^k : E^k \rightarrow C^k$. The first step is initiated by choosing $(E^0 = E, C^0 = C)$ and, iteratively, the further steps are defined by

$$\left(\begin{array}{l} E^1 := E^0 \cap TC^0 \\ C^1 := \tau_Q(E^1) \\ \tau^1 : E^1 \rightarrow C^1 \end{array} \right) \rightarrow \dots \rightarrow \left(\begin{array}{l} E^k := E^{k-1} \cap TC^{k-1} \\ C^k := \tau_Q(E^k) \\ \tau^k : E^k \rightarrow C^k \end{array} \right) \rightarrow \dots .$$

In finite dimensions, there is an end for the algorithm, that is, there exists a three tuple (E^f, τ^f, C^f) satisfying that $E^{f+1} = E^f$ and $C^{f+1} = C^f$. Note that, the final manifold E^f is integrable. We call E^f the integrable part of E . Constrained

Example 2. Consider the following set $(x, p, r, s, \dot{x}, \dot{p}, \dot{r}, \dot{s})$ on $T\mathbb{R}^4$, with the following equations [47]

$$E = \left\{ r = p, s = 0, \dot{r} = -\frac{\partial H}{\partial x}(x, p), \dot{s} = \dot{x} - \frac{\partial H}{\partial p}(x, p) \right\} \quad (53)$$

Here, $E = E^0$ and $C^0 = \{(x, p, r = p, s = 0)\}$, such that $TC^0 = \{\dot{x}, \dot{p}, \dot{r} = \dot{p}, \dot{s} = 0\}$. So,

$$E^0 \cap TC^0 = \left\{ \dot{x} = \frac{\partial H}{\partial p}(x, p), \dot{r} = \dot{p} = -\frac{\partial H}{\partial x}(x, p), r = p, s = 0 \right\} \quad (54)$$


From here, any other iteration $E^k = E^1$ and $TC^k = TC^1$. Hence, (54) is the integrable part of (53).

4.3 The Hamilton–Jacobi theory for implicit differential equations

In this section we develop a geometric Hamilton–Jacobi theory for IDE. Our problem is that given a set of IDE, we are not necessarily provided with a Hamiltonian vector field as explained in former sections. Here, we propose two methods to construct our theory. The first method consists of a theory which does refer to vector fields. The second is based on the construction of a local vector field defined on the image of a section, but not globally on the phase space.

Let us start with the first method. Consider a submanifold E of TT^*Q . By projecting E by the tangent mapping $T\pi_Q$ onto the tangent bundle TQ , we arrive a submanifold $T\pi_Q(E)$ of TQ . Note that, E refers to an IDE on T^*Q , whereas $T\pi_Q(E)$ refers to an IDE on Q . If E is integrable, then $T\pi_Q(E)$ is integrable too. We see this by considering the projection $T\pi_Q(V) = v \in T\pi_Q(E)$ of an element $V \in E$. Note that, if φ is a curve on T^*Q and it is tangent to $V \in E$, then $\pi_Q \circ \varphi$ is curve on Q which is tangent to $v \in T\pi_Q(E)$. This shows that the projections of the solutions of E are solutions of $T\pi_Q(E)$. Our aim is to discuss the inverse question, starting from the solutions of $T\pi_Q(E)$ construct solutions of E , that is to lift the solutions on Q to the cotangent bundle T^*Q . This is the philosophy of a geometric Hamilton–Jacobi theory. (Recall the geometric Hamilton–Jacobi theory exposed in subsection 4.1.) Furthermore, if E were not integrable, we would have to perform the integrability algorithm explained in subsection 4.2.

To answer the question we have to introduce a section $\gamma : Q \rightarrow T^*Q$. If any solution $\psi : \mathbb{R} \rightarrow Q$ of $T\pi_Q(E)$ is a solution $\gamma \circ \psi : \mathbb{R} \rightarrow T^*Q$ of E , then we denote the submanifold $T\pi_Q(E)$ by E^γ , and say that E and E^γ are γ -related. We illustrate this in a diagram.



In coordinates, a submanifold E of TT^*Q can be given by the set of functions $\Phi^A : TT^*Q \rightarrow \mathbb{R}$,

$$\Phi^A(q^i, p_i, \dot{q}^i, \dot{p}_i) = 0$$

The projection of E onto T^*Q by means of τ_{T^*Q} results with a submanifold C of T^*Q given by the set of functions

$$\Psi^\alpha(q^i, p_i) = 0.$$

Consider the intersection of the projected submanifold C and the image space of the one-form γ . We denote this globally by $C \cap \text{Im}(\gamma)$ and locally by $\Psi^\alpha(q^i, \gamma_j(q)) = 0$. If a solution curve is represented by $\psi^i(t) \subset Q$, the composition $\gamma \circ \psi(t) = (\psi^i(t), \gamma_i(\psi(t)))$ is a curve on T^*Q and the time derivative of the curve is


$$\begin{aligned}
\frac{d}{dt}(\gamma \circ \psi)(0) &= T\gamma(\psi(0)) \cdot \dot{\psi}(0) \\
&= \left(\psi^i(0), \gamma_i(\psi(0)), \dot{\psi}^i(0), \frac{\partial \gamma_j}{\partial q^i} \dot{\psi}^i(0) \right).
\end{aligned}$$

Then, the equations of the submanifold E along γ take the form

$$\Phi^A \left(q^i, \gamma_i(q), \dot{q}^i, \frac{\partial \gamma_j}{\partial q^i} \dot{q}^i \right) = 0, \quad (55)$$

provided that $\Psi^\alpha(q^i, \gamma_i(q)) = 0$ along $\gamma(Q) \subset C$. Here, we assumed that $\psi^i(0) = q^i$.

In the second case, we consider an additional section $\sigma : T^*Q \rightarrow TT^*Q$ such that $\sigma(C) \subset E$.




Since E is implicit, there may exist several vectors in E projecting to the same point, say c , in C . The role of the section σ is to reduce this unknown number to one. We are additionally require that the domain of the section σ be the intersection of $\text{Im}(\gamma)$ and C since, for implicit systems, C may not be the whole of T^*Q . As a result, we arrive at a vector field X_σ . Note that X_σ satisfies

$$\iota_{X_\sigma} \omega_Q = \Theta(\gamma(q)). \quad (56)$$

for an arbitrary one-form Θ defined on $\gamma(q)$.

We define a vector field X_σ^γ on the tangent bundle TQ by the commutation of the following diagram.



Explicitly,

$$X_\sigma^\gamma = T_\pi \circ X_\sigma \circ \gamma. \quad (57)$$

In local coordinates, the vector field X_σ and its projection X_σ^γ can be written as

$$X_\sigma = \sigma^i(q, \gamma(q)) \frac{\partial}{\partial q^i} + \sigma_i(q, \gamma(q)) \frac{\partial}{\partial p_i}, \quad X_\sigma^\gamma = \sigma^i(q, \gamma(q)) \frac{\partial}{\partial q^i}, \quad (58)$$

respectively. Using a one-form section γ on Q , the tangent lift of the projected vector field X_σ^γ is

$$T\gamma(X_\sigma^\gamma) = \sigma^i \left(\frac{\partial}{\partial q^i} + \frac{\partial \gamma_j}{\partial q^i} \frac{\partial}{\partial p_j} \right) \quad (59)$$

Using (57), we find an expression relating the section σ and the vector fields as follows.

$$\sigma^i(q, \gamma(q)) \frac{\partial \gamma_j}{\partial q^i}(q) = \sigma_j(q, \gamma(q)). \quad (60)$$

We are ready now to state the following lemma.

Lemma 2. *Given the conditions above, we say that: the two vector fields X_σ and X_σ^γ are γ -related if and only if (60) is fulfilled.*

Again, this construction can be mimicked for nonintegrable IDE that are submanifolds E of a higher-order bundle, after performing the integrability algorithm given in 4.2.

4.4 A HJ theory for IHS

As we have summarized in subsection (3.4), for every Lagrangian submanifold E of TT^*Q , there exists a Morse family $F : TT^*Q \rightarrow \mathbb{R}$ generating E . This enables us to write E locally as

$$E = \left\{ \left(q^i, p_i; \frac{\partial F}{\partial p_i}, -\frac{\partial F}{\partial q^i} \right) \in TT^*Q : \frac{\partial F}{\partial \lambda^a} = 0 \right\} \quad (61)$$

where $F = F(q, p, \lambda)$. We cite two important studies [7, 42] related with the problem addressed in this subsection.

We introduce a differential one-form γ on the base manifold Q . See that, $\text{Im}(\gamma)$ is a Lagrangian submanifold of T^*Q so that there is an inclusion $\iota : \text{Im}(\gamma) \hookrightarrow T^*Q$. Use the inclusion ι in order to pull the bundle (P, π, T^*Q) back over $\text{Im}(\gamma)$. By this, one arrives at a fiber bundle $(\iota^*P, \iota^*\pi, \text{Im}(\gamma))$. For the present case, the commutative diagram for a generic pullback bundle exhibited in (1) takes the following particular form.

$$\begin{array}{ccc} \iota^*P & \xrightarrow{\varepsilon} & P \\ \downarrow \iota^*\pi & & \downarrow \pi \\ \text{Im}(\gamma) & \xrightarrow{\iota} & T^*Q \end{array} \quad (62)$$

Here, the total space

$$\iota^*P = \{(\gamma(q), z) \in \text{Im}(\gamma) \times P : \pi(z) \in \text{Im}(\gamma)\}$$

of the pull-back bundle is a submanifold of P with ε is the corresponding inclusion. A local coordinate system on i^*P can be taken as $(q, \gamma(q), \lambda)$. Although restriction of the Morse family on i^*P should formally be written as $F \circ \varepsilon$, we will abuse notation by still denoting it by F , but to highlight the difference we shall write the arguments of the function as $F = F(q, \gamma(q), \lambda)$. The submanifold generated by $F = F(q, \gamma(q), \lambda)$ is given by

$$E|_{\text{Im}(\gamma)} = \left\{ \left(q^i, \gamma_i(q); \frac{\partial F}{\partial p_i}, -\frac{\partial F}{\partial q^i} \right) \in TT^*Q : \frac{\partial F}{\partial \lambda^a} = 0 \right\}. \quad (63)$$

Note that, if the Lagrangian submanifold E is the image of a Hamiltonian vector field X_H , then $E|_{\text{Im}(\gamma)}$ reduces to the image space of the composition $X_H \circ \gamma$.

The submanifold $E|_{\text{Im}(\gamma)}$ exhibited in (63) does not depend on the momentum variables. This enables us to project it to a submanifold E^γ of TQ by the tangent mapping $T\pi_Q$ as follows

$$E^\gamma = T\pi_Q \circ E|_{\text{Im}(\gamma)} = \left\{ \left(q^i, \frac{\partial F}{\partial p_i}(q, \gamma(q), \lambda) \right) \in TQ : \frac{\partial F}{\partial \lambda^a} = 0 \right\}. \quad (64)$$

Note that the submanifold E^γ defines an implicit differential equation on Q . We state the generalization of the Hamilton-Jacobi theorem (1) as follows.

Lemma 3. *The following conditions are equivalent for a closed one-form γ :*

1. *The Lagrangian submanifold E in (61) and the submanifold E^γ in (64) are γ -related, that is*

$$T\gamma(E^\gamma) = E|_{\text{Im}(\gamma)}$$

2. *And it is fulfilled that $dF(q, \gamma(q), \lambda) = 0$, where F is the Morse family generating E .*

Proof. The one-form $\gamma = \gamma_i dq^i$ is closed, that is, $\frac{\partial \gamma_i}{\partial q^j} = \frac{\partial \gamma_j}{\partial q^i}$. The first assertion in lemma 3 can be written locally as

$$\frac{\partial \gamma_j}{\partial q^i} \frac{\partial F}{\partial p_j}(q, \gamma(q), \lambda) + \frac{\partial F}{\partial q^i}(q, \gamma(q), \lambda) = 0, \quad (65)$$

with the conditions that $\partial F / \partial \lambda^a = 0$. We make a simple calculation to compute the exterior derivative of the Morse family as follows

$$dF(q, \gamma(q), \lambda) = \frac{\partial F}{\partial q^j} dq^j + \frac{\partial F}{\partial p_i} \gamma_{i,j} dq^j + \frac{\partial F}{\partial \lambda^a} d\lambda^a. \quad (66)$$

Note that, after the substitution of (65) into (66) and by employing the symmetry $\frac{\partial \gamma_i}{\partial q^j} = \frac{\partial \gamma_j}{\partial q^i}$, we arrive at that the exterior derivative of F vanishes when $p = \gamma(q)$. To prove the reverse direction, it is enough to repeat these steps in reverse order. \square

Assume now that the one-form γ is exact so that $\gamma = dW(q)$ for some real valued function W called the characteristic function on the base manifold Q . Then the second condition in lemma 3 gives the implicit Hamilton-Jacobi equation (IHJ equation)

$$F\left(q, \frac{\partial W}{\partial q}, \lambda\right) = \text{cst}, \quad \frac{\partial F}{\partial \lambda^a}\left(q, \frac{\partial W}{\partial q}, \lambda\right) = 0. \quad (67)$$

In the lemma (3), if the Lagrangian submanifold E is the image of a Hamiltonian vector field X_H , then E^γ becomes the image space of the vector field X_H^γ in (46) and we retrieve the classical HJ theory given in (47).

It is possible to generalize Lemma (3) by replacing the image space $\text{Im } \gamma$ by an arbitrary Lagrangian submanifold S of T^*Q . Note that, according to the generalized Poincaré lemma, there exists a Morse family W on the total space of a smooth bundle (R, τ, Q) generating the Lagrangian submanifold S . We equip the total space R with the coordinates (q^i, μ^α) . Then we have that the Lagrangian submanifold S can be written as

$$S = \left\{ \left(q^i, \frac{\partial W}{\partial q^i}(q, \mu) \right) \in T^*Q : \frac{\partial W}{\partial \mu^\beta}(q, \mu) = 0 \right\}. \quad (68)$$

Now, the inclusion in the diagram (62) becomes $\iota : S \mapsto T^*Q$. In this case, the restriction of the Morse family F generating the submanifold E of TT^*Q to the inclusion ϵ defines the following submanifold

$$E|_S = \left\{ \left(q^i, \frac{\partial W}{\partial q^i}; \frac{\partial F}{\partial p_i}, -\frac{\partial F}{\partial q^i} \right) \in TT^*Q : \frac{\partial F}{\partial \lambda^a} = 0, \frac{\partial W}{\partial \mu^\beta}(q, \mu) = 0 \right\}, \quad (69)$$

where $W = W(q, \mu)$ and $F = F(q, \frac{\partial W}{\partial q^i}(q, \mu), \lambda)$. The submanifold $E|_S$ does not depend on the momentum variable p explicitly. So that, its projection E^S to the tangent bundle TQ by means of $T\pi_Q$ is well-defined and given by

$$E^S = \left\{ \left(q^i; \frac{\partial F}{\partial p_i}, \frac{\partial W}{\partial q^i}(q, \mu), \lambda \right) \in TQ : \frac{\partial F}{\partial \lambda^a} = 0, \frac{\partial W}{\partial \mu^\beta} = 0 \right\}. \quad (70)$$

We are now ready to state a generalization of the lemma (3) as follows.

Lemma 4. *Let S be a Lagrangian submanifold of T^*Q generated by Morse family $W = W(q, \mu)$ defined on the total space of a bundle (R, τ, Q) . The following conditions are equivalent*

1. *The Lagrangian submanifold E in (61) and the submanifold E^S in (70) are S -related, that is*

$$T(dW|_\mu)(E^S) = E|_S$$

for every μ , where $E|_S$ is in (69).

2. *$dF\left(q, \frac{\partial W}{\partial q}(q, \mu), \lambda\right) = 0$ for all μ . Here, F is the Morse family generating E .*

We only give some clues instead of writing the whole proof of this lemma since it is very similar to the proof of lemma 3. The closedness of the one-form γ is replaced by the commutativity of the second partial derivatives of the Morse family W with respect to q .

Let us comment on the notation $T(dW|_\mu)$ as well. If the fiber variable μ is frozen, then the exterior derivative $dW|_\mu$ of the Morse family W becomes a differentiable mapping from Q to T^*Q and its tangent mapping $T(dW|_\mu)$ goes from TQ to TT^*Q . We remark that this last comment is a generalization of the Hamilton-Jacobi theory for the Lagrangian submanifolds studied in [4] as well.

Example 3. *Let V be a nonholonomic k -dimensional distribution on Q spanned by the vector fields X_a . We define the following Morse family on the total space of the fiber bundle $T^*Q \times \mathbb{R}^k$*

$$F(q, p, \lambda) = p_i \lambda^a X_a^i(q). \quad (71)$$

*Here F is a Morse family and determines a Lagrangian submanifold of TT^*Q given by*

$$\dot{q}^i = \lambda^a X_a^i(q), \quad \dot{p}_j = -p_i \lambda^a \frac{\partial X_a^i}{\partial q^j}, \quad p_i X_a^i(q) = 0. \quad (72)$$

This system is integrable according to [27]. The corresponding Hamilton-Jacobi equation is computed to be

$$\frac{\partial W}{\partial q^i} \lambda^a X_a^i(q) = cst. \quad (73)$$

5 Complete solutions of the HJ equation for IHS

Before writing the complete solution of a HJ equation of IHS, we first investigate the complete solutions of HJ equations for explicit systems in terms of Morse families and Lagrangian submanifolds.

5.1 Lagrangian submanifolds generated by complete solutions

In the classical sense, a solution W of the HJ equation (45) is called complete if it depends on additional variables that equal in number to the dimension of the base manifold Q [1]. To illustrate this, we start by considering two copies of the configuration manifold and denote them by Q and \bar{Q} . Endow these manifolds with local coordinates (q^i) and (\bar{q}^j) , respectively. The number of arbitrary parameters for the general solution is given by j , which does not necessarily equal i . A complete solution is a real valued function $W = W(\bar{q}, q)$ on the product space $\bar{Q} \times Q$ that resolves the HJ equation (45). This function generates three different Lagrangian submanifolds, let us show them.

Construct the fiber bundle structure $(\bar{Q} \times Q, \rho, Q)$. Here, the bundle projection ρ is assumed to be a projection to the second factor. As discussed previously, a real valued function $W = W(\bar{q}, q)$ on the total space $\bar{Q} \times Q$ is called a Morse family if the matrix $[\partial^2 W / \partial \bar{q}^i \partial q^j]$ is non-degenerate. We draw the following diagram

$$\begin{array}{ccc}
 \mathbb{R} & \xleftarrow{W} & \bar{Q} \times Q & & T^*Q & (74) \\
 & & \downarrow \rho & & \downarrow \pi_Q & \\
 & & Q & \xlongequal{\quad} & Q &
 \end{array}$$

In this case, the Morse family W defines a Lagrangian submanifold of T^*Q given by

$$S = \left\{ \left((q^i, \frac{\partial W}{\partial q^i}(\bar{q}, q)) \in T^*Q : \frac{\partial W}{\partial \bar{q}^i}(\bar{q}, q) = 0 \right) \right\}. \quad (75)$$

Note that, by changing the roles of Q and \bar{Q} , we may define a bundle structure $(\bar{Q} \times Q, \bar{\rho}, \bar{Q})$ over the manifold \bar{Q} and obtain a diagram similar to (74). In this case, W defines a Lagrangian submanifold \bar{S} of $T^*\bar{Q}$ as follows

$$\bar{S} = \left\{ \left((\bar{q}^i, \frac{\partial W}{\partial \bar{q}^i}(\bar{q}, q)) \in T^*\bar{Q} : \frac{\partial W}{\partial q^j}(\bar{q}, q) = 0 \right) \right\}. \quad (76)$$

Another Lagrangian submanifold generated by W is the result of following observation. The cotangent bundle $T^*(\bar{Q} \times Q) = T^*\bar{Q} \times TQ$ of the product space is a symplectic manifold equipped with the symplectic two-form $\omega_{\bar{Q}} \ominus \omega_Q$ [55]. It is evident that image of the exterior derivative dW of a complete solution W is a Lagrangian submanifold

$$\hat{S} = \left\{ \left(\bar{q}, q; \frac{\partial W}{\partial \bar{q}}, \frac{\partial W}{\partial q} \right) \in T^*(\bar{Q} \times Q) \right\}. \quad (77)$$

It is known that a Lagrangian submanifold of $T^*(\bar{Q} \times Q)$ defines a symplectomorphism between $T^*\bar{Q}$ and T^*Q . The Morse family $W = W(\bar{q}, q)$ generates a symplectomorphism according to the following identity

$$\theta_{\bar{Q}} \ominus \theta_Q = \bar{p}_i d\bar{q}^i - p_i dq^i = dW(q, \bar{q}), \quad (78)$$

where we assume the Darboux' coordinates on the cotangent bundles. In the local picture, the induced symplectomorphism is given by

$$\varphi : T^*\bar{Q} \rightarrow T^*Q : \left(\bar{q}^i, \frac{\partial W}{\partial \bar{q}^j} \right) \rightarrow \left(q^i, -\frac{\partial W}{\partial q^j} \right). \quad (79)$$

5.2 Complete solutions for the case of implicit Hamiltonian systems

Let us first try to geometrize the complete solutions of the HJ equation for explicit systems. A function $W = W(\bar{q}, q)$ is a complete solution of the HJ equation (45) if the Hamiltonian function H is constant when it is restricted to S exhibited in (75). That is, a complete solution W is the one satisfying

$$H \left(q, \frac{\partial W}{\partial q}(\bar{q}, q) \right) = \text{cst}, \quad \frac{\partial W}{\partial \bar{q}} = 0. \quad (80)$$

Using the symplectic diffeomorphism φ in (79) generated by the function W , we pull the function H back to $T^*\bar{Q}$ and see that φ^*H is a constant. So that the dynamics generated by φ^*H is trivial.

Now, assume that, we have a Lagrangian submanifold E of TT^*Q . Then there exists a Morse family F generating E . A complete solution of the implicit Hamilton-Jacobi equation (67) is a smooth function W satisfying

$$F \left(q, \frac{\partial W}{\partial q}, \lambda \right) = \text{cst}, \quad \frac{\partial W}{\partial \bar{q}}(\bar{q}, q) = 0, \quad \frac{\partial F}{\partial \lambda} \left(q, \frac{\partial W}{\partial q}, \lambda \right) = 0. \quad (81)$$

We aim to pull the dynamics E or the Morse family F back to $T^*\bar{Q}$. To achieve this goal, we recall the definition of the pullback bundle in (1) and

apply it to the diffeomorphism (79). This way we obtain a fiber bundle structure $(\varphi^*P, \varphi^*\pi, T^*\bar{Q})$ where the total space is defined to be

$$\varphi^*P = \{(\bar{z}, r) \in T^*\bar{Q} \times P : \varphi(z) = \pi(r)\}$$

equipped with the induced coordinates $(\bar{q}^i, \bar{p}^i, \lambda)$, and $\varphi^*\pi$ is the projection to the second factor. We draw the following commutative diagram in order to summarize the discussion.

$$\begin{array}{ccc} \varphi^*P & \xrightarrow{\hat{\varphi}} & P \\ \downarrow \varphi^*\pi & & \downarrow \pi \\ T^*\bar{Q} & \xrightarrow{\varphi} & T^*Q \end{array} \quad (82)$$

Here, $\hat{\varphi}$ is a diffeomorphism and in the local coordinates reads

$$\hat{\varphi} : \varphi^*P \leftrightarrow P : \left(\bar{q}^i, \frac{\partial W}{\partial \bar{q}^i}, \lambda \right) \leftrightarrow \left(q^i, -\frac{\partial W}{\partial q^i}, \lambda \right). \quad (83)$$

The pullback $\bar{F} = F \circ \hat{\varphi}$ of the Morse function F by $\hat{\varphi}$ is a Morse family on the total space of the pullback bundle $(\varphi^*P, \varphi^*\pi, T^*\bar{Q})$. Note that \bar{F} is a constant function and the implicit differential equation generated by \bar{F} renders trivial dynamics.

Generalizing, the most general form of a Lagrangian submanifold of $T^*\bar{Q} \times T^*Q$ is generated by a Morse family W defined on the total space of the fiber bundle (R, τ, M) where the base manifold M is a submanifold of $\bar{Q} \times Q$. Let us depict it in a diagram.

$$\begin{array}{ccc} \mathbb{R} & \xleftarrow{W} & R \\ & & \downarrow \tau \\ & & M \\ & & \xrightarrow{\iota} \bar{Q} \times Q \\ & & \uparrow \pi_{(\bar{Q} \times Q)} \\ & & T^*\bar{Q} \times T^*Q \end{array} \quad (84)$$

A complete solution to the implicit Hamilton Jacobi equation (67) is a Morse family W defined on the total space R . Note that, the Morse family F generating the dynamics on E reduces to a constant function on the Lagrangian submanifold generated by W .

Let us now depict the situation in coordinates. Assume that a submanifold M of $\bar{Q} \times Q$ is defined by a number “ l ” of equations

$$U^a(\bar{q}, q) = 0, \quad a = 1, \dots, l. \quad (85)$$

Consider a real function W' on the submanifold M and define an arbitrary continuation

$$W = W' + \nu_a U^a$$

of W' to the product space $P = \bar{Q} \times Q \times \Lambda$ where (ν_a) 's are the Lagrange multipliers defining a local coordinate system for Λ . This W is a complete solution of the implicit Hamiltonian dynamics (3) generated by F if

$$F\left(q^i, \frac{\partial W'}{\partial q^i}, \lambda\right) = cst, \quad U^a(\bar{q}, q) = 0, \quad a = 1, \dots, l.$$

An implicit description of a Lagrangian submanifold of $T^*\bar{Q} \times T^*Q$ generated by W or equivalently of the corresponding diffeomorphism φ can be computed by

$$\theta_Q \ominus \theta_{\bar{Q}} = p_i dq^i - \bar{p}_i d\bar{q}^i = d(W'(q, \bar{q}) + \nu_a U^a(q, \bar{q})), \quad (86)$$

where the $\theta_{\bar{Q}}$ and θ_Q are the canonical one-forms on \bar{Q} and Q , respectively. In this case, the momenta $p \in T_q^*Q$ and $\bar{p} \in T_{\bar{q}}^*\bar{Q}$ can be explicitly stated as

$$\begin{aligned} \bar{p}_i &= -\frac{\partial W'}{\partial \bar{q}^i} - \nu_a \frac{\partial U^a}{\partial \bar{q}^i} \\ p_i &= \frac{\partial W'}{\partial q^i} + \nu_a \frac{\partial U^a}{\partial q^i} \\ U^a(\bar{q}, q) &= 0, \quad a = 1, \dots, l. \end{aligned} \quad (87)$$

6 Application to degenerate Lagrangian systems

6.1 Lagrangian dynamics

A Lagrangian function L is a real valued function on TQ . Consider the coordinates (q^i, \dot{q}^i) on TQ induced those from Q . We define a vertical endomorphism S given by $S = \partial/\partial \dot{q}^i \otimes dq^i$. Note that, S is a $(1, 1)$ -tensor field on Q . In terms of S the Cartan one-form θ_L is defined to be

$$\theta_L = S^*(dL) = (\partial L / \partial \dot{q}^i) dq^i$$

where dL is the exterior derivative of a Lagrangian density. The Cartan two-form derives from the Cartan one-form $\omega_L = -d\theta_L$. See that ω_L is symplectic if the Hessian matrix

$$(W_{ij}) = \left(\frac{\partial^2 L}{\partial \dot{q}^i \partial \dot{q}^j} \right) \quad (88)$$

is not singular. In this case, the fiber derivative (or the Legendre transformation)

$$\mathbb{F}L : TQ \mapsto T^*Q : (q^i, \dot{q}^j) \mapsto \left(q^i, \frac{\partial L}{\partial \dot{q}^j} \right) \quad (89)$$

becomes a symplectomorphism relating the Cartan two-form ω_L on TQ and the canonical symplectic two-form ω_Q on T^*Q . The Lagrangian is said to be hyperregular if the fiber derivative $\mathbb{F}L$ is a global diffeomorphism.

The energy is defined as $E_L = \Delta(L) - L$, a real valued function on TQ where the Liouville vector field is $\Delta = \dot{q}^i \partial / \partial \dot{q}^i$. The Hamiltonian is retrieved through

$$H(q, p) = E_L \circ \mathbb{F}L^{-1}. \quad (90)$$

If the Lagrangian is regular, or equivalently, if ω_L is symplectic, then the Lagrange equations can be expressed geometrically as

$$\iota_{\xi_L} \omega_L = dE_L, \quad (91)$$

whose solution ξ_L is called a Euler–Lagrange vector field explicitly given by

$$\xi_L = \dot{q}^i \frac{\partial}{\partial q^i} + \xi^i(q, \dot{q}) \frac{\partial}{\partial \dot{q}^i}. \quad (92)$$

The integral curves $(q^i(t), \dot{q}^i(t))$ are lifts of their projections $(q^i(t))$ on Q and are solutions of the system of differential equations

$$\frac{dq^i(t)}{dt} = \dot{q}^i, \quad \frac{d\dot{q}^i(t)}{dt} = \xi^i, \quad (93)$$

which is equivalent to a second-order differential equation

$$\frac{d^2 q^i(t)}{dt^2} = \xi^i. \quad (94)$$

The curves $(q^i(t))$ in Q are called the solutions of ξ_L that correspond with the solutions of the Euler–Lagrange equation

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}^i} \right) = \frac{\partial L}{\partial q^i}. \quad (95)$$

If the Lagrangian is regular, then the fiber derivative (89) has the following geometry.

$$\begin{array}{ccccc}
(TTQ, \omega_L) & \xrightarrow{T\mathbb{F}L} & (TT^*Q, \omega_Q^T) & \xrightarrow{\alpha_Q} & T^*TQ \\
& \searrow \tau_{TQ} & & \nearrow dL & \\
& \searrow \xi_L & TQ & \xrightarrow{L} & \mathbb{R}
\end{array}$$

In this case, the Hamiltonian vector field X_H associated with the Hamiltonian function H in (90) and ξ_L presented in (91) are related as

$$X_H \circ \mathbb{F}L = T\mathbb{F}L \circ \xi_L.$$

The diffeomorphisms α_Q and β_Q maps Lagrangian submanifolds into Lagrangian submanifolds,

$$\alpha_Q(\text{Im}(X_H)) = \text{Im}(dL), \quad \beta_Q \circ \alpha_Q^{-1}(\text{Im}(dL)) = \text{Im}(dH),$$

whereas the Hamiltonian and the Lagrangian vector fields are related to their corresponding Lagrangian submanifolds as

$$\beta_Q \circ X_H = dH, \quad \alpha_Q \circ T\mathbb{F}L \circ \xi_L = dL,$$

respectively.

6.2 Lagrangian dynamics as a Lagrangian submanifold

In this section we depict the geometric interpretation of a HJ theory for Lagrangian dynamics. For it, we present the EL equations in terms of Morse families and special symplectic structures.

Recall the special symplectic structure on the left side of Tulczyjew's triple.

$$\begin{array}{ccc}
T^*TQ & \xleftarrow{\alpha_Q} & TT^*Q \\
\pi_{TQ} \downarrow & \swarrow \tau_{TQ} & \\
\mathbb{R} & \xleftarrow{L} & T^*Q
\end{array} \tag{96}$$

In the induced local picture on TT^*Q , by following the procedure presented in subsection (3.2), we compute the Lagrangian submanifold E generated by the Lagrangian L as

$$E = \left\{ \left(q^i, \frac{\partial L}{\partial \dot{q}^i}; \dot{q}^i, \frac{\partial L}{\partial q^i} \right) \in TT^*Q \right\} \tag{97}$$

which is equivalent to the EL equations (95). We can generate this Lagrangian submanifold from the right wing (the Hamiltonian side) of the

triple (22) by defining a proper Morse family $F^{L \rightarrow H}$ on the Pontryagin bundle $PQ = TQ \times_Q T^*Q$ over T^*Q . On a local chart, the energy function

$$F(q, p, \dot{q}) = p_i \dot{q}^i - L(q, \dot{q}) \quad (98)$$

satisfies the requirements of being a Morse family. Hence, F generates a Lagrangian submanifold of T^*T^*Q as defined in equation (26). In coordinates $(q^i, p_i, \alpha_i, \beta^i)$ of T^*T^*Q , this Lagrangian submanifold is given by

$$\alpha_i = \frac{\partial F}{\partial q^i} = -\frac{\partial L}{\partial q^i}, \quad \beta^i = \frac{\partial F}{\partial p_i} = \dot{q}^i, \quad \frac{\partial F}{\partial \dot{q}^i} = p_i - \frac{\partial L}{\partial \dot{q}^i} = 0 \quad (99)$$

The inverse of the isomorphism β_Q maps this Lagrangian submanifold to the Lagrangian submanifold E presented in (97). Here, we record the following diagram for this.

$$\begin{array}{ccc} \mathbb{R} & \xleftarrow{F} & PQ \\ & & \downarrow \pi \\ & & T^*Q \\ & & \text{=====} \\ & & T^*Q \\ & & \uparrow \pi_{T^*Q} \\ T^*T^*Q & \xleftarrow{\beta_Q} & TT^*Q \\ & \swarrow \tau_{T^*Q} & \\ & & T^*Q \end{array} \quad (100)$$

For regular cases, the Morse family F on PQ can be reduced to a Hamiltonian function H on T^*Q . For degenerate cases, a reduction of the total space PQ to a subbundle larger than T^*Q is possible depending on degeneracy level of Lagrangian function [8]. There exists an intrinsically geometric procedure for dealing with constraints in Hamiltonian and Lagrangian mechanics. It has been available since 1979, with advantages over the Dirac-Bergman algorithm, it is the Gotay-Bergman algorithm [22, 23, 24, 25] (read Appendix for brief description of the method).

6.3 Hamilton-Jacobi theory for degenerate Lagrangians

To write the associated HJ equation of the EL equations generated by (possibly) degenerate Lagrangian densities, we apply lemma 3 to the Lagrangian submanifold presented in (97). Accordingly, we arrive at that the implicit HJ equation

$$F(q, \gamma(q), \dot{q}) = \gamma_i(q) \dot{q}^i - L(q, \dot{q}) = \text{cst}$$

for a closed one-form $\gamma = \gamma_i(q) dq^i$. Taking the exterior derivative of this equation, we arrive at the following local picture of the Hamilton-Jacobi equation for a Lagrangian L

$$\dot{q}^i \frac{\partial \gamma_i}{\partial q^j}(q) - \frac{\partial L}{\partial q^j}(q, \dot{q}) = 0, \quad \gamma_i(q) - \frac{\partial L}{\partial \dot{q}^i}(q, \dot{q}) = 0. \quad (101)$$

To illustrate this, we propose two particular problems [34].

Example 4. Consider the degenerate Lagrangian L on $T\mathbb{R}^3$ given by

$$L(q, \dot{q}) = L(q^1, q^2, q^3, \dot{q}^1, \dot{q}^2, \dot{q}^3) = \frac{1}{2}(\dot{q}^1 + \dot{q}^2)^2,$$

and the Whitney bundle $T\mathbb{R}^3 \oplus T^*\mathbb{R}^3$ fibered on $T^*\mathbb{R}^3$ and parameterized by $(q^1, q^2, q^3, \dot{q}^1, \dot{q}^2, \dot{q}^3, p_1, p_2, p_3)$. The Lagrange multipliers correspond with $(\lambda^i) = (\dot{q}^1, \dot{q}^2, \dot{q}^3)$. Following (98), we define the Morse family

$$F(q, \dot{q}, p) = \dot{q}^1 p_1 + \dot{q}^2 p_2 + \dot{q}^3 p_3 - \frac{1}{2}(\dot{q}^1 + \dot{q}^2)^2.$$

that generates a Lagrangian submanifold E of $TT^*\mathbb{R}^3$. Explicitly,

$$E = \{(q^1, q^2, q^3, p_1, p_2, p_3; \dot{q}^1, \dot{q}^2, \dot{q}^3, 0, 0, 0) \in TT^*\mathbb{R}^3 \\ : p_1 = \dot{q}^1 + \dot{q}^2, p_2 = \dot{q}^1 + \dot{q}^2, p_3 = 0\}. \quad (102)$$

It is evident that the projection of E to $T^*\mathbb{R}^3$ results with the following 4 dimensional submanifold

$$C = \{(q^1, q^2, q^3, p_1, p_2, p_3) \in T^*\mathbb{R}^3 : p_1 = p_2, p_3 = 0\}.$$

Consider now the closed one-form $\gamma : \mathbb{R}^3 \mapsto T^*\mathbb{R}^3$. According to the Lagrangian HJ theorem (3), the system (101) in this particular case takes the form:

$$\begin{cases} [1] & \dot{q}^1 \frac{\partial \gamma_1}{\partial q^1} + \dot{q}^2 \frac{\partial \gamma_2}{\partial q^1} + \dot{q}^3 \frac{\partial \gamma_3}{\partial q^1} = 0, \\ [2] & \dot{q}^1 \frac{\partial \gamma_1}{\partial q^2} + \dot{q}^2 \frac{\partial \gamma_2}{\partial q^2} + \dot{q}^3 \frac{\partial \gamma_3}{\partial q^2} = 0, \\ [3] & \dot{q}^1 \frac{\partial \gamma_1}{\partial q^3} + \dot{q}^2 \frac{\partial \gamma_2}{\partial q^3} + \dot{q}^3 \frac{\partial \gamma_3}{\partial q^3} = 0, \\ [4] & \gamma_1 - \dot{q}^1 - \dot{q}^2 = 0 \\ [5] & \gamma_2 - \dot{q}^1 - \dot{q}^2 = 0 \\ [6] & \gamma_3 = 0. \end{cases} \quad (103)$$

It is immediate to see from equations [4] and [5] that $\gamma_1 = \gamma_2$. If γ is closed and $\gamma_3 = 0$, one obtains that γ_1 and γ_2 are independent of q^3 . Then [3] in (103) is automatically satisfied. The substitution of [4] into [1] and [2] results in

$$\gamma_1 \frac{\partial \gamma_1}{\partial q^1} = 0, \quad \gamma_1 \frac{\partial \gamma_1}{\partial q^2} = 0.$$

A nontrivial solution is possible if γ_1 and γ_2 are constants. Hence, we record the one-form

$$\gamma(q) = (q^1, q^2, q^3, c, c, 0).$$

In terms of submanifolds, that is, according to the first condition in theorem 3, the picture is the following. The constant character of the one-form defines a constraint $\dot{q}^1 + \dot{q}^2 = c$, on the velocity variables. We first restrict the submanifold E to the image space of the γ , we arrive at

$$E|_{\text{Im}(\gamma)} = \{(q^1, q^2, q^3, c, c, 0; \dot{q}^1, \dot{q}^2, \dot{q}^3, 0, 0, 0) \in TT^*\mathbb{R}^3 : c = \dot{q}^1 + \dot{q}^2\}$$

whose generic version is given in (63). The projection of $E|_{\text{Im}(\gamma)}$ to the tangent bundle $T\mathbb{R}^3$ by $T\pi_{\mathbb{R}^3}$ results in a five dimensional submanifold

$$E^\gamma = \{(q^1, q^2, q^3, \dot{q}^1, \dot{q}^2, \dot{q}^3) \in T\mathbb{R}^3 \quad \text{with} \quad \dot{q}^1 + \dot{q}^2 = c\},$$

of the tangent bundle $T\mathbb{R}^3$. The system of equations (103) is equivalent to saying that the tangent lift of E^γ by the tangent mapping $T\gamma$ equals $E|_{\text{Im}(\gamma)}$. It is indeed immediate to see that $T\gamma(E^\gamma) = E|_{\text{Im}(\gamma)}$.

In terms of vector fields, the situation is as follows. Consider a section σ of the tangent bundle τ_{T^*Q} given by

$$\sigma(q^1, q^2, q^3, p_1, p_2, p_3) = (q^1, q^2, q^3, p_1, p_2, p_3; c - \dot{q}^2, \dot{q}^2, \dot{q}^3, 0, 0, 0)$$

on the intersection $C \cap \text{Im}(\gamma)$. Note that, $\text{Im}(\sigma) \subset E$. Accordingly, we write the following vector field

$$X_\sigma = c \frac{\partial}{\partial q^1} + \dot{q}^3 \circ \gamma(q) \frac{\partial}{\partial q^3} + \dot{q}^2 \circ \gamma(q) \left(\frac{\partial}{\partial q^2} - \frac{\partial}{\partial q^1} \right).$$

We project this vector field by $T\pi_Q$ and arrive at the vector field X_σ^γ , which is locally the same as X_σ . Composing with the section γ ,

$$T\gamma(q^1, q^2, q^3, \dot{q}^1, \dot{q}^2, \dot{q}^3) = (q^1, q^2, q^3; c, c, 0, \dot{q}^1, \dot{q}^2, \dot{q}^3, 0, 0, 0)$$

This shows that

$$T\gamma(X_\sigma^\gamma) = X_\sigma \tag{104}$$

is obviously fulfilled.

Example 5. Consider the Lagrangian L on $T\mathbb{R}^2$ given by

$$L(q, \dot{q}) = L(q^1, q^2, \dot{q}^1, \dot{q}^2) = \frac{1}{2}(\dot{q}^1)^2 + q^2(q^1)^2.$$

To recast the EL system generated by this Lagrangian density, we simply define the following Morse family on the Whitney sum $T\mathbb{R}^2 \oplus T^*\mathbb{R}^2$

$$F(q, p, \dot{q}) = p_1 \dot{q}^1 + p_2 \dot{q}^2 - \frac{1}{2}(\dot{q}^1)^2 - q^2(q^1)^2. \tag{105}$$

This family generates the following Lagrangian submanifold

$$E = \{(q^1, q^2, p_1, p_2; \dot{q}^1, \dot{q}^2, 2q^2\dot{q}^1, (q^1)^2) \in TT^*\mathbb{R}^2 : p_1 = \dot{q}^1, p_2 = 0\}. \quad (106)$$

The projection of E onto the cotangent bundle $T^*\mathbb{R}^2$ by the tangent bundle projection $\tau_{T^*\mathbb{Q}}$ results with the following submanifold

$$C = \{(q^1, q^2, p_1, p_2) \in T^*\mathbb{R}^2 : p_2 = 0\}. \quad (107)$$

According to theorem 3, we now introduce a closed one-form γ on \mathbb{R}^2 and require that the Morse family F in (105) is constant on the image space, that is

$$F(q, \gamma(q), \dot{q}) = \gamma_1(q)\dot{q}^1 + \gamma_2(q)\dot{q}^2 - \frac{1}{2}(\dot{q}^1)^2 - q^2(q^1)^2 = cst.$$

For this case, the Hamilton Jacobi equations (101) turn out to be the following set

$$\begin{cases} [1] & \dot{q}^1 \frac{\partial \gamma_1(q)}{\partial q^1} + \dot{q}^2 \frac{\partial \gamma_2(q)}{\partial q^1} - 2q^2\dot{q}^1 = 0 \\ [2] & \dot{q}^1 \frac{\partial \gamma_1(q)}{\partial q^2} + \dot{q}^2 \frac{\partial \gamma_2(q)}{\partial q^2} - (q^1)^2 = 0 \\ [3] & \gamma_1(q) - \dot{q}^1 = 0 \\ [4] & \gamma_2(q) = 0. \end{cases} \quad (108)$$

The closedness of γ , together with equation [4], imply that γ_1 depends only on q^1 . If we substitute this and equations [3] and [4] in [1] and [2], then we arrive at a partial differential equation and a constraint

$$\gamma_1(q) \frac{\partial \gamma_1(q)}{\partial q^1} - 2q^2\dot{q}^1 = 0, \quad q^1 = 0.$$

Note that the constraint $q^1 = 0$ implies that the system is automatically satisfied for any function $\gamma_1 = \gamma_1(q^1)$. Then, the one-form is described by

$$\gamma(q) = (q^1, q^2; q^1, 0). \quad (109)$$

Let us assume that $\gamma_1(q^1) = q^1$ as it is done in [34], and rewrite system (108) in terms of submanifolds. If the Lagrangian submanifold E in (106) is restricted to the image space of γ in (109), then the result becomes

$$E|_{\text{Im}(\gamma)} = \{(q^1, q^2, q^1, 0; \dot{q}^1, \dot{q}^2, 2q^2\dot{q}^1, (q^1)^2) \in TT^*\mathbb{R}^2 : q^1 = 0\}. \quad (110)$$

This is projected to the tangent bundle $T\mathbb{R}^2$ by $T\pi_{\mathbb{R}^2}$ in order to get the reduced dynamics

$$E^\gamma = \{(q^1, q^2; \dot{q}^1, \dot{q}^2) : q^1 = 0, \dot{q}^1 = q^1 = 0\}.$$

The γ -relatedness of E and E^γ can be checked with the following lift

$$T\gamma \circ E^\gamma = (q^1 = 0, q^2, q^1, 0; \dot{q}^1 = 0, \dot{q}^2, \dot{q}^1 = 0, 0).$$

As a result, we have three constraints $q^1 = 0, p_1 = 0$ and $p_2 = 0$. So, the projected submanifold C in (107) must be rectified as

$$C = \{(q^1, q^2, p_1, p_2) : q^1 = 0, p_1 = 0, p_2 = 0\} \subset T^*\mathbb{R}^2.$$

Here, $p_2 = 0$ is called the primary constraint, since it roots in the functional structure of the Lagrangian function, and the other two constraints $q^1 = 0$ and $p_1 = 0$ are called the secondary constraints.

Consider a section σ of the tangent bundle fibration $\tau_{T^*\mathbb{R}^2}$ given by

$$\sigma(q^1, q^2; \dot{q}^1, \dot{q}^2) = (q^1, q^2, \dot{q}^1, \dot{q}^2; \dot{q}^1, \dot{q}^2, 2q^2\dot{q}^1, (q^1)^2).$$

By restricting this section to the intersection $\text{Im}(\gamma) \cap C$, we arrive at the following vector field

$$X_\sigma = \dot{q}^2 \circ \gamma(q) \frac{\partial}{\partial q^2}.$$

The projection of this vector field to $T\mathbb{R}^2$ is the vector field X_σ^γ , and it looks exactly as X_σ , at least locally. X_σ^γ and X_σ are γ related since the tangent lift by γ is given by

$$T\gamma(q, \dot{q}) = (q^1, q^2, q^1, 0; \dot{q}^1, \dot{q}^2, \dot{q}^1, 0)$$

and maps the vector field X_σ^γ into X_σ .

Conclusions and future work

In this work we have presented a geometric Hamilton-Jacobi theory for systems of implicit differential equations. In the general context, due to the implicit character of the equations, the lack of a vector field has been solved by the introduction of a local section σ . In the particular case of the implicit Hamiltonian dynamics, Morse families and special symplectic structures have been employed to derive a Hamilton–Jacobi theory in which the Morse function plays the role of the Hamiltonian. This result has been particularly applied to singular Lagrangians. We expect further applications of the theory in constraint Hamiltonian systems, Dirac systems, and etc.

The obtainance of a Hamilton–Jacobi theorem through reduction is here sketched in terms of coisotropic reduction [37]. As future work, we aim at stating the problem of reduction of the implicit Hamilton-Jacobi theory under the Lie group symmetry of the implicit system of differential equations.

Acknowledgements

This work has been partially supported by MINECO MTM 2013-42-870-P and the ICMAT Severo Ochoa project SEV-2011-0087. One of us (OE) is grateful for ICMAT for warm hospitality where some parts of the works has been done. (OE) is also grateful Prof. Hasan Gümral for valuable discussions in the theory of Tulczyjew triples.

Appendix

The Gotay–Nester algorithm

The Gotay-Nester algorithm is a suitable tool for reducing the dynamics of singular Hamiltonian or Lagrangian systems to a reduced manifold where the motion is well-defined. This algorithm was created as a generalization of the well-known Dirac-Bergman algorithm which has local nature and does not cope with all the singularities appearing in dynamics. Let us briefly describe the Gotay-Nester algorithm [22, 23, 24, 25].

Let us recall that given a singular Lagrangian $L : TQ \rightarrow \mathbb{R}$, the Legendre transform $FL : TQ \rightarrow T^*Q$ and the energy $E_L : TQ \rightarrow \mathbb{R}$, one can define a presymplectic system $(M_1 = FL(TQ), \omega_1)$, where ω_1 is the restriction of the canonical symplectic form on T^*Q to M_1 . We will assume that L is almost regular (the Legendre transformation is a submersion and surjective), then $M_1 = FL(TQ)$ is a submanifold of T^*Q .

The restriction of the Legendre mapping $FL_1 : TQ \rightarrow M_1$ to this submanifold is a submersion with connected fibers. In this case, M_1 is called the submanifold of primary constraints. If L is almost regular, $\ker(TFL) = \ker(\omega_L) \cap V(TQ)$, where $V(TQ)$ denotes the vertical bundle, and the fibers are connected, a direct computation shows that E_L projects onto a function $h_1 : M_1 \rightarrow \mathbb{R}$. The inclusion of this submanifold is denoted by $j_1 : M_1 \rightarrow T^*Q$ and define $\omega_1 = j_1^*(\omega_Q)$. The dynamics in the primary constraint manifold is

$$i_X \omega_1 = dh_1,$$

where $h_1 \in C^\infty(M_1)$ is the projection of the energy $E_L \in C^\infty(TQ)$. Now, there are two possibilities: the solution X defined at all the points of M_1 is such that X defines global dynamics and it is a solution (modulo $\ker \omega_1$), in other words, there are only primary constraints. Or, we are in need of a second submanifold M_2 where $\iota_X \omega_1 = dh_1$ and $X \in TM_1$. But such a solution X is not necessarily tangent to M_2 , so we have to impose an additional tangency condition to M_2 and obtain a new submanifold M_3 along which there exists a solution. Continuing this process, we obtain a sequence of submanifolds

$$\dots M_k \hookrightarrow \dots \hookrightarrow M_2 \hookrightarrow M_1 \hookrightarrow T^*Q$$


where the general description of M_{l+1} is

$$M_{l+1} := \{p \in M_l \text{ such that there exists } X_p \in T_p M_l \text{ satisfying } i_X \omega_1 = dh_1\}.$$

If the algorithm stabilizes at some k , say $M_{k+1} = M_k$, then we say that M_k is the final constraint submanifold which is denoted by M_f , and then there exists a well-defined solution X along M_f . This constraint algorithm produces a solution X of the equation

$$(i_X \omega_1 = dh_1)|_{M_f},$$

where X is a vector field on M_f . We can depict the situation in the following diagram:



References

- [1] V.I. Arnold, *Mathematical methods of classical mechanics*. Graduate texts in mathematics, **60**, 1989.
- [2] R. Abraham and J.E. Marsden, *Foundations of mechanics*. Reading, Massachusetts: Benjamin/Cummings Publishing Company, 1978.
- [3] J.L. Anderson and P.G. Bergmann, *Constraint in covariant field theories*, Phys. Rev **83**, (1951), 1018–1026.
- [4] M. Barbero-Lin, M. de Len and D. M. de Diego, *Lagrangian submanifolds and the Hamilton-Jacobi equation*. Monatshefte fr Mathematik, **171**,(2013), 269-290.
- [5] F. Barone, R. Grassini and G. Mendella, *Unified approach to constrained mechanical systems as implicit differential equations*, Annals de l' I.H.P. Section A **70**, (1999), 515–546.
- [6] O. Bernardi, F. Cardin and A. Siconolfi, *Cauchy problems for stationary Hamilton-Jacobi equations under mild regularity assumptions*. Journal of Geometric Mechanics, **1(3)**, (2009), 271-294.
- [7] S. Benenti and W.M. Tulczyjew, *The geometrical meaning and globalization of the Hamilton-Jacobi method* In *Differential Geometrical Methods in Mathematical Physics* (pp. 9-21). Springer Berlin Heidelberg, 1890.
- [8] S. Benenti, *Hamiltonian structures and generating families*, Springer Science & Business Media, 2011.
- [9] P. G. Bergmann, *Bosonic quantum gravity according to the global one-dimensionality conjecture*, Helv. Phys. Acta, Suppl. **4**, (1956), 79.
- [10] J.F. Cariñena, J. Grabowski, J. de Lucas and C. Sardón, *LieDirac systems and Schwarzian equations*, J. Differential Equations **257**, (2014), 23032340.
- [11] F. Cardin, *Morse families and constrained mechanical systems. Generalized hyperelastic materials*. Meccanica, **26(2-3)**, (1991), 161-167.
- [12] F. Cardin, *Elementary symplectic topology and mechanics*. Springer, 2015.

- [13] J.F. Cariñena, X. Gracia, G. Marmo, E. Martinez, M.C. Muñoz-Lecanda and N. Roman-Roy, *Geometric Hamilton–Jacobi theory*, Int. J. Geom. Methods in Mod. Phys., **3(07)**, (2006), 1417-1458.
- [14] T.J. Courant, *Dirac manifolds*. Transactions of the American Mathematical Society, **319(2)**, (1990), 631-661.
- [15] B. Dacorogna and P. Marcellini, *Progress in Nonlinear differential equations and their applications*, Springer Science+Business Media, LLC, New York, 1999.
- [16] P.A.M. Dirac, *Generalized Hamiltonian dynamics*, Can. J. Math. **2**, (1950), 129–148.
- [17] P.A.M. Dirac, *Lectures in Quantum Mechanics*, Belfer Graduate School of Science, Yeshiva University, New York, 1967.
- [18] D. Dominici, G. Longhi, J. Gomis, J.M. Pons, *HamiltonJacobi theory for constrained systems*. J. Math. Phys., **25(8)**, 2439-2452 (1984).
- [19] C. Godbillion, *Geometrie differentielle et mecanique analytique*, Collection Methodes Hermann, Paris 1969.
- [20] H. Goldstein, *Mecánica Clásica*, 4a Ed. Aguilar SA Madrid 1979.
- [21] M.J. Gotay and J.M. Nester *Presymplectic Lagrangian systems. I. The constraint algorithm and the equivalence theorem*, Ann. Inst. H. Poincar Sect. A (N.S.) **30**, (1979), 129–142.
- [22] M.J. Gotay and J.M. Nester: *Presymplectic Lagrangian systems. II. The second-order equation problem*. Ann. Inst. H. Poincar Sect. A (N.S.), **32**, (1980), 1–13.
- [23] M.J. Gotay and J.M. Nester, *Generalized constraint algorithm and special presymplectic manifolds*, Geom. Methods in Math. Phys. (Proc. NSF-CBMS Conf., Univ. Lowell, Lowell, Mass., 1979), pp. 78-104, Lecture Notes in Math., **775**, Springer, Berlin, 1980.
- [24] M.J. Gotay and J.M. Nester, *Apartheid in the Dirac theory of constraints*, J. Phys. A, **17**, (1984), 3063–3066.
- [25] M.J. Gotay, J.M. Nester and G. Hinds, *Presymplectic manifolds and the Dirac-Bergmann theory of constraints*, J. Math. Phys. **19**, (1978), 2388–2399.

- [26] J. Hoefkens, M. Berz and K. Makino, *Computing validated solutions of implicit differential equations*, Advances in Comp. Math. **19**, (2003), 231–253.
- [27] S. Janeczko, *On implicit Lagrangian differential systems*, In *Annales Polonici Mathematici* **74**, 133-141. Institute of Mathematics Polish Academy of Sciences, 2000.
- [28] S. Janeczko and F. Pelletier, *Singularities of implicit differential systems and Maximum Principle*, BANACH CENTER PUBLICATIONS, **62**, (2004), 117-132.
- [29] T.W. Kibble and F.H. Berkshire, *Classical Mechanics*, Imperial College Press, London, 5th Ed., 2004.
- [30] M. Leok, T. Ohsawa, D. Sosa, *HamiltonJacobi theory for degenerate Lagrangian systems with holonomic and nonholonomic constraints*, J. Math. Phys., **53(7)**, (2012), 072905.
- [31] M. de León and P.R. Rodrigues, *Method of Differential Geometry in Analytical Mechanics*, Mathematical studies **158**, North–Holland, 1989.
- [32] M. de León, D. Iglesias-Ponte and D. Martín de Diego, *Towards a Hamilton-Jacobi theory for nonholonomic mechanical systems*. J. Phys. A: Math. Gen., 015205, (2008), 14 pp.
- [33] M. de León, J.C. Marrero and D. Martín de Diego, *Linear almost Poisson structures and Hamilton-Jacobi equation. Applications to nonholonomic mechanics*, J. Geom. Mech. **2**, (2010), 159198.
- [34] M. de Len, J.C. Marrero, D. M. de Diego and M. Vaquero, *On the Hamilton-Jacobi theory for singular Lagrangian systems*. J. Math. Phys., **54(3)**, (2013), 032902.
- [35] M. de León and C. Sardón, *Cosymplectic and contact structures to resolve time-dependent and dissipative Hamiltonian systems*, Accepted in J. Phys. A
ArXiv:1612.06224 (2017)
- [36] M. de León and C. Sardón, *A geometric Hamilton–Jacobi theory for Nambu–Poisson structures*, J. Math. Phys. **58**, (2017), 033508.
- [37] M. de León, D.M. de Diego and M. Vaquero, *HamiltonJacobi theory, symmetries and coisotropic reduction*. J. Math. Pures et Appliquées, **107(5)**, 591-614.

- [38] P. Libermann and C.M. Marle, *Symplectic geometry and analytical mechanics*, **35**, Springer Science & Business Media, 2012.
- [39] G. Marmo, G. Mendella and W.M. Tulczyjew, *Symmetries and constants of the motion for dynamics in implicit form*, In *Annales de l'IHP Physique théorique*, **57**, (1992), 147-166.
- [40] G. Mendella, G. Marmo and W.M. Tulczyjew, *Integrability of implicit differential equations*, *J. Phys. A: Math. Gen.*, **28(1)**, (1995), 149.
- [41] G. Marmo, G. Mendella and W.M. Tulczyjew, *Constrained Hamiltonian systems as implicit differential equations*. *J. Phys. A: Math. Gen.*, **30(1)**, (1977), 277.
- [42] G. Marmo, G. Morandi and N. Mukunda, *A geometrical approach to the Hamilton-Jacobi form of dynamics and its generalizations*. *La Rivista del Nuovo Cimento*, **13(8)**, (1990), 1-74.
- [43] M.R. Menzio and W.M. Tulczyjew, *Infinitesimal symplectic relations and generalized Hamiltonian dynamics*, *Ann. Inst. H. Poincaré*, **A28**, (1978), 249–367.
- [44] H. T. Davis, *Introduction to nonlinear differential and integral equations*, Dover Books on Mathematics, 2010.
- [45] F. Takens, *Implicit differential equations: Some open problems* In *Singularités d'Applications Différentiables* (pp. 237-253). Springer Berlin Heidelberg, 1976.
- [46] W.M. Tulczyjew, *Hamiltonian systems, Lagrangian systems and the Legendre transformation*, *Sympl. Math.* **16**, (1974) 247–258.
- [47] G. Mendella, G. Marmot, W.M. Tulczyjew, *Integrability of implicit differential equations*. *J. Phys. A: Math. Gen.*, **28(1)**, (1995), 149.
- [48] W.M. Tulczyjew, *A symplectic formulation of relativistic particle dynamics* (No. MPI-PAE/ASTRO-103). Max-Planck-Institut für Physik und Astrophysik, 1976.
- [49] W.M. Tulczyjew, *The legendre transformation*. In *Annales de l'IHP Physique théorique* **27**, (1977), 101-114.
- [50] W.M. Tulczyjew, *A symplectic formulation of particle dynamics*. In *Differential Geometrical Methods in Mathematical Physics* 457-463 Springer Berlin Heidelberg, 1977.

- [51] W.M. Tulczyjew and P. Urbanski, *Homogeneous Lagrangian Systems, Gravitation, Electromagnetism and Geometric Structures*, Pitagora Editrice (1996), 91-136.
- [52] W.M. Tulczyjew, *A slow and careful Legendre transformation for singular Lagrangians*, Acta Phys. In Pol. **B30**, 1999.
- [53] W.M. Tulczyjew, *Geometric Formulations of Physical Theories: Statics and Dynamics of Mechanical Systems* **11**, Amer. Inst. of Phys., 1989.
- [54] P. Urbanski, *Double vector bundles in Classical Mechanics*, Rend. Sem. Matem. Torino, **54**, (1996).
- [55] A. Weinstein, *Lectures on symplectic manifolds*, CBMS Regional Conference Series in Mathematics, **9**, (197y), 48pp.