

Partite Saturation of Complete Graphs

António Girão*

Teeradej Kittipassorn†

Kamil Popielarz‡

November 12, 2021

Abstract

We study the problem of determining $\text{sat}(n, k, r)$, the minimum number of edges in a k -partite graph G with n vertices in each part such that G is K_r -free but the addition of an edge joining any two non-adjacent vertices from different parts creates a K_r . Improving recent results of Ferrara, Jacobson, Pfender and Wenger, and generalizing a recent result of Roberts, we define a function $\alpha(k, r)$ such that $\text{sat}(n, k, r) = \alpha(k, r)n + o(n)$ as $n \rightarrow \infty$. Moreover, we prove that

$$k(2r - 4) \leq \alpha(k, r) \leq \begin{cases} (k - 1)(4r - k - 6) & \text{for } r \leq k \leq 2r - 3, \\ (k - 1)(2r - 3) & \text{for } k \geq 2r - 3, \end{cases}$$

and show that the lower bound is tight for infinitely many values of r and every $k \geq 2r - 1$. This allows us to prove that, for these values, $\text{sat}(n, k, r) = k(2r - 4)n + O(1)$ as $n \rightarrow \infty$. Along the way, we disprove a conjecture and answer a question of the first set of authors mentioned above.

1 Introduction

Given a graph H , the classical Turán-type extremal problem asks for the maximum number of edges in an H -free graph on n vertices. While the corresponding minimization problem is trivial, it is interesting to determine the minimum number of edges in a maximal H -free graph on n vertices. We say that a graph is H -saturated if it is H -free but the addition of an edge joining any two non-adjacent vertices creates a copy of H . The minimum number $\text{sat}(n, H)$ of edges in an H -saturated graph on n vertices was first studied in 1949 by Zykov [17] and independently in 1964 by Erdős, Hajnal, and Moon [5] who proved that $\text{sat}(n, K_r) = (r - 2)(n - 1) - \binom{r-2}{2}$. Soon after this, Bollobás [1] determined exactly $\text{sat}(n, K_r^{(s)})$ where $K_r^{(s)}$ is the complete s -uniform hypergraph on r vertices. Later, in 1986, Kászonyi and Tuza [10] showed that the saturation number $\text{sat}(n, H)$ for a graph H on r vertices is maximized

* Department of Pure Mathematics and Mathematical Statistics, University of Cambridge, Wilberforce Road, Cambridge CB3 0WB, UK; A.Girao@dpms.cam.ac.uk.

† Departamento de Matemática, Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rua Marquês de São Vicente 225, Gávea, Rio de Janeiro, RJ 22451-900, Brazil; ping41@gmail.com.

‡ Department of Mathematical Sciences, University of Memphis, Memphis, TN 38152, USA; kamil.popielarz@gmail.com

at $H = K_r$, and consequently, $\text{sat}(n, H)$ is linear in n for any H . For results on the saturation number, we refer the reader to the survey [6].

This concept can be generalized to the notion of H -saturated subgraphs which are maximal elements of a family of H -free subgraphs of a fixed host graph. A subgraph of a graph G is said to be H -saturated in G if it is H -free but the addition of an edge in $E(G)$ joining any two non-adjacent vertices creates a copy of H . The problem of determining the minimum number $\text{sat}(G, H)$ of edges in an H -saturated subgraph of G was first proposed in the above mentioned paper of Erdős, Hajnal, and Moon. They conjectured a value for the saturation number $\text{sat}(K_{m,n}, K_{r,r})$ which was verified independently by Bollobás [2, 3] and Wessel [15, 16]. Very recently, Sullivan and Wenger [14] studied the analogous saturation numbers for tripartite graphs within tripartite graphs and determined $\text{sat}(K_{n_1, n_2, n_3}, K_{l, l, l})$ for every fixed $l \geq 1$ and every n_1, n_2 and n_3 sufficiently large. Several other host graphs have been considered, including hypercubes [4, 9, 12] and random graphs [11].

In this paper, we are interested in the saturation number $\text{sat}(n, k, r) = \text{sat}(K_{k \times n}, K_r)$ for $k \geq r \geq 3$ where $K_{k \times n}$ is the complete k -partite graph containing n vertices in each of its k parts. This function was first studied recently by Ferrara, Jacobson, Pfender and Wenger [7] who determined $\text{sat}(n, k, 3)$ for $n \geq 100$. Later, Roberts [13] showed that $\text{sat}(n, 4, 4) = 18n - 21$ for sufficiently large n .

For convenience, we say that a k -partite graph with a fixed k -partition is K_r -partite-saturated if it is K_r -free but the addition of an edge joining any two non-adjacent vertices from different parts creates a K_r . Therefore, $\text{sat}(n, k, r)$ is the minimum number of edges in a k -partite graph G with n vertices in each part which is K_r -partite-saturated.

Our first result states that $\text{sat}(n, k, r)$ is linear in n where the constant $\alpha(k, r)$ in front of n is defined as follows. Given $k \geq r \geq 3$, consider a K_r -partite-saturated k -partite graph G containing an independent set X of size k consisting of exactly one vertex from each part of G . We define $\alpha(k, r)$ to be the minimum number of edges between X and X^c taken over all such G and X .

Theorem 1. For $k \geq r \geq 3$,

$$\text{sat}(n, k, r) = \alpha(k, r)n + o(n)$$

as $n \rightarrow \infty$.

Let us shift our focus to the function $\alpha(k, r)$. The next theorem states what we know about it.

Theorem 2. For $k \geq r \geq 3$,

$$(i) \quad k(2r - 4) \leq \alpha(k, r) \leq \begin{cases} (k - 1)(4r - k - 6) & \text{for } r \leq k \leq 2r - 3, \\ (k - 1)(2r - 3) & \text{for } k \geq 2r - 3. \end{cases}$$

$$(ii) \quad \alpha(k, r) = k(2r - 4) \text{ if } \begin{cases} k = 2r - 3, \text{ or} \\ k \geq 2r - 2 \text{ and } r \equiv 0 \pmod{2}, \text{ or} \\ k \geq 2r - 1 \text{ and } r \equiv 2 \pmod{3}. \end{cases}$$

$$(iii) \quad \alpha(k, 3) = 3(k - 1), \alpha(4, 4) = 18 \text{ and } 33 \leq \alpha(5, 5) \leq 36.$$

(iv) $\alpha(r, r) \geq r(2r - 4) + 1$ for $r \geq 4$.

The bounds in (i), together with Theorem 1, imply that $\text{sat}(n, k, r) = O(krn)$, answering a question of Ferrara, Jacobson, Pfender and Wenger [7]. In (ii), we determine exactly $\alpha(k, r)$ for some values of r and every k large enough, allowing us to disprove a conjecture in [7] which states that $\text{sat}(n, k, r) = (k - 1)(2r - 3)n - (2r - 3)(r - 1)$ for $k \geq 2r - 3$ and sufficiently large n . In (iii), we deal with the cases $r = 3, 4, 5$ which have not been covered by (ii). Finally, (iv) shows that the lower bound in (i), which is attained for certain values of r and k mentioned in (ii), is not tight when $k = r$.

Theorem 1 and Theorem 2 imply that $\text{sat}(n, k, r) = k(2r - 4)n + o(n)$ for the values of k and r in (ii). We show that, in this case, the $o(n)$ term can be replaced by a constant.

Theorem 3. For $k \geq r \geq 3$,

$$\text{sat}(n, k, r) = k(2r - 4)n + O(1) \text{ if } \begin{cases} k = 2r - 3, \text{ or} \\ k \geq 2r - 2 \text{ and } r \equiv 0 \pmod{2}, \text{ or} \\ k \geq 2r - 1 \text{ and } r \equiv 2 \pmod{3}, \end{cases}$$

as $n \rightarrow \infty$.

Now we give a summary of the values of $\text{sat}(n, k, r)$ in the case $r = 3, 4, 5$ which are immediate consequences of the first three results.

Corollary 4. (i) $\text{sat}(n, k, 3) = 3(k - 1)n + o(n)$ for $k \geq 3$ and as $n \rightarrow \infty$.

$$(ii) \text{ sat}(n, k, 4) = \begin{cases} 18n + o(n) & \text{for } k = 4, \text{ as } n \rightarrow \infty, \\ 4kn + O(1) & \text{for } k \geq 5, \text{ as } n \rightarrow \infty. \end{cases}$$

$$(iii) \text{ sat}(n, k, 5) \begin{cases} \in [33n + o(n), 36n + o(n)] & \text{for } k = 5, \text{ as } n \rightarrow \infty, \\ \in [36n + o(n), 40n + o(n)] & \text{for } k = 6, \text{ as } n \rightarrow \infty, \\ \in [48n + o(n), 49n + o(n)] & \text{for } k = 8, \text{ as } n \rightarrow \infty, \\ = 6kn + O(1) & \text{for } k = 7 \text{ or } k \geq 9, \text{ as } n \rightarrow \infty. \quad \square \end{cases}$$

We note that (i) and the first half of (ii) are not the best known results. In fact, Ferrara, Jacobson, Pfender and Wenger [7] proved that $\text{sat}(n, k, 3) = 3(k - 1)n - 6$ for sufficiently large n and Roberts [13] proved that $\text{sat}(n, 4, 4) = 18n - 21$ for sufficiently large n .

Let us give some more definitions which will be used throughout the paper. For a k -partite $G = V_1 \cup V_2 \cup \dots \cup V_k$, we refer to each V_i as a *part of G* . We say that an edge (or a non-edge) uv of a k -partite graph is *admissible* if u, v lie in different parts. We say that a non-edge uv of a K_r -free graph is *K_r -saturated* if adding uv to the graph completes a K_r . In other words, a k -partite graph is *K_r -partite-saturated* if it is K_r -free and every admissible non-edge is K_r -saturated.

The rest of this paper is organized as follows. Section 2 is devoted to the proof of Theorem 1. In Section 3, we study the function $\alpha(k, r)$ and prove Theorem 2(i). In Section 4, we prove Theorem 2(ii)

by describing constructions matching the lower bound $\alpha(k, r) \geq k(2r - 4)$ in Theorem 2(i). We prove Theorem 2(iii), Theorem 2(iv) and Theorem 3 in Section 5, Section 6 and Section 7 respectively. Finally, we conclude the paper in Section 8 with some open problems.

2 Proof of Theorem 1

First we show that the upper bound follows easily from the definition of $\alpha(k, r)$.

Proposition 5. *For every $k \geq r \geq 3$ and any integer $n \geq \alpha(k, r) + 1$, we have $\text{sat}(n, k, r) \leq \alpha(k, r)n + \alpha(k, r)^2$.*

Proof. Let G be a K_r -partite-saturated k -partite graph containing an independent set X of size k consisting of exactly one vertex from each part of G with $e(X, X^c) = \alpha(k, r)$. We may assume that $|X^c| \leq \alpha(k, r)$. Indeed, since there are $\alpha(k, r)$ edges between X and X^c , deleting all the vertices in X^c with no neighbors in X leaves at most $\alpha(k, r)$ vertices in X^c . Note that any admissible non-edge with at least one endpoint in X is still K_r -saturated. We finish by keeping adding admissible edges inside X^c until every admissible non-edge inside X^c is K_r -saturated.

Let V_1, V_2, \dots, V_k be the parts of G . It follows that $|V_i| = |V_i \cap X| + |V_i \cap X^c| \leq 1 + \alpha(k, r) \leq n$, and so we can modify G to have exactly n vertices in each part by blowing up the vertex of X in V_i to a class of size $n - |V_i \cap X^c|$ for each i . The resulting graph is K_r -partite-saturated and has exactly n vertices in each of its k parts. Moreover, the number of edges is at most $\alpha(k, r)n + e(G[X^c]) \leq \alpha(k, r)n + \alpha(k, r)^2$. \square

Now we prove the lower bound $\text{sat}(n, k, r) \geq \alpha(k, r)n + o(n)$.

Let $\varepsilon > 0$ and let $G = V_1 \cup V_2 \cup \dots \cup V_k$ be a K_r -partite-saturated k -partite graph with $|V_i| = n$ for all $i \in [k]$. We shall show that $e(G) \geq \alpha(k, r)n - \varepsilon n$ for all sufficiently large n . Let d be a large natural number to be chosen later. For each i , we partition V_i into $V_i^+ = \{v \in V_i : d(x) \geq d\}$ and $V_i^- = \{v \in V_i : d(x) < d\}$. First we show that V_i^+ is small. Since $e(G) \geq \frac{d}{2}|V_i^+|$, we are done unless $|V_i^+| \leq \frac{2\alpha(k, r)}{d}n$. Now we show that we can delete a constant number of vertices from $\bigcup_{i=1}^k V_i^-$ to make it independent.

Lemma 6. *There exists a subset $U \subset \bigcup_{i=1}^k V_i^-$ of size $C_{k, d}$ such that $(\bigcup_{i=1}^k V_i^-) \setminus U$ forms an independent set in G for some constant $C_{k, d}$.*

Let us first show how to finish the proof of Proposition 5 using the lemma. For each $1 \leq i \leq k$, let v_i be a vertex of smallest degree in $V_i^- \setminus U$. Since G is a K_r -partite-saturated k -partite graph and $X = \{v_1, v_2, \dots, v_k\}$ is an independent set with exactly one vertex in each part of G , we have $\sum_{i=1}^k d(v_i) \geq \alpha(k, r)$ by the definition of $\alpha(k, r)$. Since $(\bigcup_{i=1}^k V_i^-) \setminus U$ forms an independent set,

$$\begin{aligned} e(G) &\geq \sum_{i=1}^k \sum_{v \in V_i^- \setminus U} d(v) \geq \sum_{i=1}^k |V_i^- \setminus U| d(v_i) \geq (n - |V_i^+| - |U|) \sum_{i=1}^k d(v_i) \\ &\geq \alpha(k, r) \left(n - \frac{2\alpha(k, r)}{d}n - C_{k, d} \right) = \alpha(k, r)n - \left(\frac{2\alpha(k, r)^2}{d} + \frac{\alpha(k, r)C_{k, d}}{n} \right) n \geq \alpha(k, r)n - \varepsilon n \end{aligned}$$

by taking d and n sufficiently large. It remains to prove the lemma.

Proof of Lemma 6. It is sufficient to show that any matching between V_i^- and V_j^- has size less than 4^{d^2} for all $i \neq j$. Indeed, we can take U to be the endpoints of maximal matchings between V_i^- and V_j^- for all $i \neq j$ and $|U| < 4^{d^2} \binom{k}{2}$.

Suppose for contradiction that $\{x_1y_1, x_2y_2, \dots, x_{4^{d^2}}y_{4^{d^2}}\}$ is a matching of size 4^{d^2} where $X = \{x_1, x_2, \dots, x_{4^{d^2}}\} \subset V_1^-$ and $Y = \{y_1, y_2, \dots, y_{4^{d^2}}\} \subset V_2^-$. The strategy of the proof is to iteratively find vertices $x_{t_1}, x_{t_2}, \dots, x_{t_d}$ of X such that $d(x_{t_i}) \geq i$ for all $1 \leq i \leq d$, which would contradict the fact that $x_{t_d} \in V_1^-$. In fact, we shall find vertices $x_{t_1}, x_{t_2}, \dots, x_{t_d}$ of X such that

- (i) there exists a common neighbor of x_{t_i} and y_{t_j} which is not a neighbor of $y_{t_1}, y_{t_2}, \dots, y_{t_{j-1}}$ for all $i > j$.

Clearly, this implies that $d(x_{t_i}) \geq i$ for all $1 \leq i \leq d$. To find such vertices, it is sufficient to find vertices $x_{t_1}, x_{t_2}, \dots, x_{t_d}$ of X satisfying

- (ii) x_{t_i} and y_{t_j} are not neighbors for all $i > j$, and
- (iii) $N(x_{t_i}) \cap N(y_{t_l}) = N(x_{t_j}) \cap N(y_{t_l})$ for all $i > j > l$.

First we show that (ii) and (iii) imply (i). Let $i > j$. By (ii), $x_{t_i}y_{t_j}$ is a non-edge. Since G is K_r -partite-saturated, there exists a clique W of size $r - 2$ in the common neighborhood of x_{t_i} and y_{t_j} . Since $r \geq 3$, we are done by picking a required vertex from W unless each vertex in W is joined to some y_{t_l} with $l < j$. In this case, $W \cup \{x_{t_j}, y_{t_j}\}$ forms a clique of size r , contradicting the fact that G is K_r -free. Indeed, each $w \in W$ belongs to some $N(y_{t_l})$ with $l < j$, and since $w \in N(x_{t_i})$, we must have $w \in N(x_{t_j})$, by (iii).

Now, we find vertices $x_{t_1}, x_{t_2}, \dots, x_{t_d}$ of X satisfying (ii) and (iii). To help us do so, we shall iteratively construct a nested sequence of sets $X \supset X_1 \supset X_2 \supset \dots \supset X_d$ with $x_{t_i} \in X_i$ for all $2 \leq i \leq d$, satisfying

- (iv) x and $y_{t_{i-1}}$ are not neighbors for all $x \in X_i$, and
- (v) $N(x) \cap N(y_{t_{i-1}}) = N(x') \cap N(y_{t_{i-1}})$ for all $x, x' \in X_i$.

Clearly, such vertices $x_{t_1}, x_{t_2}, \dots, x_{t_d}$ satisfy (ii) and (iii). Start with $x_{t_1} = x_1$ and $X_1 = X$. Let $i \leq d$ and suppose that we have found vertices $x_{t_1}, x_{t_2}, \dots, x_{t_{i-1}}$ and sets $X_1 \supset X_2 \supset \dots \supset X_{i-1}$ with $x_{t_j} \in X_j$ for all $j < i$, satisfying (iv) and (v). We delete the neighbors of $y_{t_{i-1}}$ from X_{i-1} and partition the remaining vertices into $2^{d(y_{t_{i-1}})} \leq 2^d$ subsets according to their common neighborhood with $y_{t_{i-1}}$. In other words, $X_{i-1} \setminus N(y_{t_{i-1}})$ is partitioned into subsets $\{x : N(x) \cap N(y_{t_{i-1}}) = S\}$ for $S \subset N(y_{t_{i-1}})$. We choose X_i to be such subset of maximum size, i.e. $|X_i| \geq \frac{|X_{i-1}| - d}{2^d}$. Clearly, X_i satisfies (iv) and (v). We then choose x_{t_i} be any vertex in X_i . It remains to prove that $|X_i| > 0$. Recall that $|X_1| = |X| = 4^{d^2}$, and we can see, by induction, that $|X_i| \geq 4^{d(d-i)}$ for $i \leq d$. Indeed,

$$|X_i| \geq \frac{|X_{i-1}| - d}{2^d} \geq \frac{|X_{i-1}|}{4^d} \geq \frac{4^{d(d-i+1)}}{4^d} \geq 4^{d(d-i)}$$

as required. □

3 Bounding $\alpha(k, r)$

In this section, we establish a number of results that will help us prove Theorem 2. We shall deduce Theorem 2(i) at the end of the section.

For $k \geq r \geq 2$ and $1 \leq i \leq k - r + 1$, let $\beta_i(k, r)$ be the minimum number of vertices in a K_r -free k -partite graph such that the subgraph induced by any $k - i$ parts contains a K_{r-1} , i.e. the deletion of any i parts does not destroy all the K_{r-1} .

We observe that β_1 and β_2 are useful for bounding α .

Proposition 7. *For $k \geq r \geq 3$,*

$$k\beta_1(k-1, r-1) \leq \alpha(k, r) \leq (k-1)\beta_2(k, r-1).$$

Proof. To prove the lower bound, let G be a K_r -partite-saturated k -partite graph containing an independent set X of size k consisting of exactly one vertex from each part of G . We shall show that $e(X, X^c) \geq k\beta_1(k-1, r-1)$. It is sufficient to show that each vertex in X has degree at least $\beta_1(k-1, r-1)$. Let $x \in X$ and consider the $(k-1)$ -partite graph $H = G[N(x)]$. Clearly, it is K_{r-1} -free since G is K_r -free. It remains to show that, for each part U of H , $H \setminus U$ contains a K_{r-2} . If x' is a vertex of X in the corresponding part of U in G then, since the non-edge xx' is K_r -saturated in G , $H \setminus U$ must contain a K_{r-2} . Hence, $|N(x)| = |H| \geq \beta_1(k-1, r-1)$.

For the upper bound, let G_1 be a K_{r-1} -free k -partite graph on $\beta_2(k, r-1)$ vertices such that the subgraph induced by any $k-2$ parts contains a K_{r-2} . Let G_2 be the graph obtained from G_1 by adding one vertex of $X = \{x_1, x_2, \dots, x_k\}$ to each part of G_1 and joining each x_i to every vertex of G_1 outside its part. By construction, X forms an independent set and $e(X, X^c) = (k-1)\beta_2(k, r-1)$ edges. Note that G_2 is K_r -free since a clique in G_2 contains at most one vertex from X and G_1 is K_{r-1} -free. Now, let G be the graph obtained from G_2 by adding admissible edges inside X^c , until every admissible non-edge inside X^c is K_r -saturated. To conclude that G is K_r -partite-saturated, we need to show that every admissible non-edge inside X is K_r -saturated. Note that, for every pair of distinct vertices $x, x' \in X$, G_1 contains a K_{r-2} not using vertices from the parts containing x and x' . Since x and x' are joined to every vertex outside their parts, the addition of the edge xx' completes a K_r . Hence, $\alpha(k, r) \leq e(X, X^c) = (k-1)\beta_2(k, r-1)$. \square

In the next sections, the argument above used in the proof of the lower bound will be used several times. Let us state it as a lemma.

Lemma 8. *Let G be a k -partite K_r -free graph containing an independent set X of size k consisting of exactly one vertex from each part of G such that the non-edges inside X are K_r -saturated. Then, for each $x \in X$, $G[N(x)]$ is a K_{r-1} -free $(k-1)$ -partite graph such that the subgraph induced by any $k-2$ parts contains a K_{r-2} . In particular, $d(x) \geq \beta_1(k-1, r-1)$ for all $x \in X$.*

In the next two subsections, we shall bound β_1 from below and β_2 from above.

3.1 Upper bounds for β_i

We start with an easy observation which helps us bound β_i from above.

Lemma 9. *For $k \geq r \geq 3$ and $1 \leq i \leq k - r + 1$, $\beta_i(k, r) \leq \beta_i(k - 1, r - 1) + i + 1$.*

Proof. Let $H = U_1 \cup U_2 \cup \dots \cup U_{k-1}$ be a K_{r-1} -free $(k-1)$ -partite graph on $\beta_i(k-1, r-1)$ vertices such that the subgraph induced by any $k-i-1$ parts contains a K_{r-2} . We shall construct a K_r -free k -partite graph $G = V_1 \cup V_2 \cup \dots \cup V_k$ from H with $|G| = |H| + (i+1)$ as follows. First, add new vertices v_1 to U_1 , v_2 to U_2 , \dots , v_i to U_i and v_{i+1} to the new part V_k . This is possible since $k \geq i+2$. Now, join v_{i+1} to every vertex in H and, for every $1 \leq j \leq i$, join v_j to every vertex in $H \setminus U_j$. Clearly, G is K_r -free since H is K_{r-1} -free.

Let \mathcal{C} be a collection of $k-i$ parts of G . It remains to check that the subgraph of G induced by \mathcal{C} contains a K_{r-1} . First, suppose that $V_k \in \mathcal{C}$. By the induction hypothesis, the other $(k-1)-i$ parts $\mathcal{C} \setminus \{V_k\}$ induce a subgraph of H containing a K_{r-2} . Together with $v_{i+1} \in V_k$, they form a K_{r-1} in the subgraph of G induced by \mathcal{C} as required. Now, let us suppose that $V_k \notin \mathcal{C}$. Then \mathcal{C} must contain at least one of V_1, V_2, \dots, V_i . Without loss of generality, we may assume that \mathcal{C} contains V_1 . By the induction hypothesis, the other $(k-1)-i$ parts $\mathcal{C} \setminus \{V_1\}$ induce a subgraph of H containing a K_{r-2} . Together with $v_1 \in V_1$, they form a K_{r-1} in the subgraph of G induced by \mathcal{C} as required. \square

Lemma 9 immediately implies the following upper bound on β_i .

Corollary 10. $\beta_i(k, r) \leq (i+1)(r-1)$ for $k \geq r \geq 2$ and $1 \leq i \leq k - r + 1$.

Proof. It is clear that $\beta_i(k, 2) = i + 1$ for $k \geq i + 1$ by considering the empty graph on $i + 1$ vertices where each vertex is in a different part and the remaining $k - i - 1$ parts are empty.

By induction on r and applying Lemma 9, $\beta_i(k, r) \leq \beta_i(k - 1, r - 1) + i + 1 \leq (i + 1)(r - 2) + i + 1 = (i + 1)(r - 1)$ as required. \square

We remark that there is a straightforward construction proving Corollary 10 for the case $k \geq (i + 1)(r - 1)$, namely, a disjoint union of $i + 1$ cliques of size $r - 1$ where each vertex is in a different part and the remaining $k - (i + 1)(r - 1)$ parts are empty. Clearly, the deletion of any i parts does not destroy all the K_{r-1} .

Now we prove a better upper bound for $\beta_i(k, r)$ in the case when $i \geq 2$ and $k \geq i(r - 1) + 1$ by considering the $(r - 2)$ th power of the cycle $C_{i(r-1)+1}$.

Proposition 11. $\beta_i(k, r) \leq i(r - 1) + 1$ for $k \geq i(r - 1) + 1$ and $r, i \geq 2$.

Proof. Since $\beta_i(k, r)$ is decreasing in k (by adding empty parts), it is enough to show that $\beta_i(k, r) \leq i(r - 1) + 1$ for $k = i(r - 1) + 1$. Let G be the $(r - 2)$ th power of the cycle $C_{i(r-1)+1}$, i.e. G is a graph on $\mathbb{Z}_{i(r-1)+1}$ where u, v are neighbors if $u - v = 1, 2, \dots, r - 2$. We view G as a $(i(r - 1) + 1)$ -partite graph with one vertex in each part. Clearly, G is K_r -free if $i \geq 2$. Note that, after deleting any i vertices of G , there are at least $r - 1$ consecutive vertices remaining in $\mathbb{Z}_{i(r-1)+1}$, which form a K_{r-1} as required. \square

Proposition 11 together with Lemma 9 imply a better upper bound than that in Corollary 10 for $\beta_2(k, r)$ in the remaining cases, i.e when $k < 2r - 1$.

Proposition 12. $\beta_2(k, r) \leq 4r - k - 2$ for $2 \leq r < k < 2r - 1$.

Proof. We proceed by induction on $2r - k$. The base case when $2r - k = 1$ follows from Proposition 11. Now, suppose that $2r - k \geq 2$. Applying Lemma 9,

$$\beta_2(k, r) \leq \beta_2(k - 1, r - 1) + 3 \leq (4(r - 1) - (k - 1) - 2) + 3 = 4r - k - 2,$$

by the induction hypothesis, since $2r - k > 2(r - 1) - (k - 1) \geq 1$, □

Let us remark that a similar upper bound for general β_i can be obtained by the same method. We believe that the bound in Proposition 12 is, in fact, an equality.

Conjecture 13. $\beta_2(k, r) = 4r - k - 2$ for $2 \leq r < k < 2r - 1$.

For the remaining values of k , we shall see in the next subsection that $\beta_2(k, r) = 2r - 1$ for $k \geq 2r - 1$.

3.2 Determining β_1

We shall show that the upper bound for β_1 given by Corollary 10 is an equality. Recall that the clique number of a graph is the order of a maximum clique.

Proposition 14. $\beta_1(k, r) = 2(r - 1)$ for $k \geq r \geq 2$.

The lower bound, is a consequence of the following observation.

Proposition 15. *Let G be a graph on at most $2s - 1$ vertices with clique number s . Then there is a vertex which lies in every K_s of G .*

Proof of Proposition 14. The upper bound follows from Corollary 10. To prove the lower bound, suppose for contradiction that G is a K_r -free k -partite graph on at most $2r - 3$ vertices such that the subgraph induced by any $k - 1$ parts contains a K_{r-1} . Applying Proposition 15 with $s = r - 1$, there is a vertex v which lies in every K_{r-1} . In particular, the deletion of the part containing v destroys all the K_{r-1} . Hence, $\beta_1(k, r) \geq 2r - 2$. □

Let us remark that Proposition 15 is a consequence of the clique collection lemma of Hajnal [8] which states that the sum of the number of vertices in the union and the intersection of a collection of maximum cliques is at least twice the clique number. Our argument below can also be used to give a new proof of Hajnal's clique collection lemma.

Proof of Proposition 15. Let $V_1, V_2, \dots, V_m \subset V(G)$ be the vertex sets of the copies of K_s in G . For a vertex $v \in V(G)$, let $I_v = \{i \in [m] : v \in V_i\}$ be the set of K_s containing v . For a collection $\mathcal{C} \subset \mathcal{P}([m])$ of subsets of $[m]$, let $V_{\mathcal{C}} = \{v \in V(G) : I_v \in \mathcal{C}\}$. Observe that if $\mathcal{C} \subset \mathcal{P}([m])$ is intersecting then $V_{\mathcal{C}}$ induces a clique in G . Indeed, $u, v \in V_{\mathcal{C}}$ are neighbors since $I_u \cap I_v \neq \emptyset$, i.e. there is a clique containing both u and v . Therefore, $|V_{\mathcal{C}}| \leq s$ since G is K_{s+1} -free. The following lemma implies the result.

Lemma 16. For $m \geq 3$, there exist intersecting families $\mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_{m-2} \subset \mathcal{P}([m])$ such that, for

$$I \subset [m], \text{ the number of } \mathcal{C}_j \text{ containing } I \text{ is } \begin{cases} 0 & \text{if } I = \emptyset \\ |I| - 1 & \text{if } I \neq \emptyset, [m] \\ m - 2 & \text{if } I = [m]. \end{cases}$$

Proof. The proof is by induction on m . For $m = 3$, $\mathcal{C}_1 = \{\{1, 2\}, \{2, 3\}, \{3, 1\}, \{1, 2, 3\}\}$ satisfies the required property. For $m \geq 4$, suppose by induction that there exist intersecting families $\mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_{m-3} \subset \mathcal{P}([m-1])$ satisfying the property. We define $\mathcal{D}_1, \mathcal{D}_2, \dots, \mathcal{D}_{m-2} \subset \mathcal{P}([m])$ as follows. For $1 \leq j \leq m-3$, let

$$\mathcal{D}_j = \mathcal{C}_j \cup \{I \cup \{m\} : I \in \mathcal{C}_j\}$$

and

$$\mathcal{D}_{m-2} = \{I \subset [m] : m \in I \text{ and } |I| \geq 2\} \cup \{[m-1]\}.$$

It is easy to check that $\mathcal{D}_1, \mathcal{D}_2, \dots, \mathcal{D}_{m-2}$ satisfy the required property. \square

Let us deduce the result. This is trivial when $m = 1, 2$ so we may assume that $m \geq 3$. Observe that

$$\sum_{i=1}^m |V_i| = \left| \bigcup_{i=1}^m V_i \right| + \sum_{j=1}^{m-2} |V_{\mathcal{C}_j}| + \left| \bigcap_{i=1}^m V_i \right|.$$

Indeed, a vertex v is counted on both sides $|I_v|$ times by the lemma. Using $|V_i| = s$, $|\bigcup_{i=1}^m V_i| \leq 2s - 1$ and $|V_{\mathcal{C}_j}| \leq s$, we have

$$ms \leq (2s - 1) + (m - 2)s + \left| \bigcap_{i=1}^m V_i \right|$$

i.e. $|\bigcap_{i=1}^m V_i| \geq 1$ as required. \square

We remark that the fact that $\beta_1(k, r) = 2(r - 1)$ allows us to show that the upper bound for $\beta_2(k, r)$ when $k \geq 2r - 1$ in Proposition 11 is an equality.

Corollary 17. $\beta_2(k, r) = 2r - 1$ for $k \geq 2r - 1$ and $r \geq 2$.

Proof. Observe that $\beta_i(k, r) \geq \beta_{i-1}(k - 1, r) + 1$. Indeed, if G is a K_r -free k -partite graph on $\beta_i(k, r)$ vertices such that the subgraph induced by any $k - i$ parts contains a K_{r-1} , then, by deleting a non-empty part of G , we obtain a K_r -free $(k - 1)$ -partite graph such that the subgraph induced by any $(k - 1) - (i - 1)$ parts contains a K_{r-1} . This graph must contain at least $\beta_{i-1}(k - 1, r)$ vertices and therefore, $|G| - 1 \geq \beta_{i-1}(k - 1, r)$.

Hence, $\beta_2(k, r) \geq \beta_1(k - 1, r) + 1 = 2(r - 1) + 1 = 2r - 1$ by Proposition 14. \square

3.3 Proof of Theorem 2(i)

The lower bound follows from Proposition 7 and Proposition 14. The upper bound follows from Proposition 7, Proposition 12 and Corollary 17. \square

4 Proof of Theorem 2(ii)

For $k = 2r - 3$, we are done since the lower and upper bounds in Theorem 2(i) match, i.e. $\alpha(k, r) = k(2r - 4) = (k - 1)(2k - 3)$.

Now we shall describe constructions that match the lower bound $\alpha(k, r) \geq k(2r - 4)$ in Theorem 2(i) for the cases when ($k \geq 2r - 2$ and r is even) and ($k \geq 2r - 1$ and $r \equiv 2 \pmod{3}$), i.e. a K_r -partite-saturated k -partite graph G containing an independent set X of size k consisting of exactly one vertex from each part of G with $e(X, X^c) = k(2r - 4)$. Lemma 8 tells us that such graph must satisfy $d(x) = 2r - 4$, for all $x \in X$.

Note that we do not have to worry about making the admissible non-edges inside X^c , K_r -saturated since we can keep adding admissible edges inside X^c until every admissible non-edge inside X^c is K_r -saturated.

Let $p \in \{2, 3\}$ be a divisor of $r - 2$. First we shall construct such k -partite graph G , for $k = 2r - 4 + p$. We define $X = \{x_1, x_2, \dots, x_k\}$ and $X^c = \{y_1, y_2, \dots, y_k\}$, where the parts of G are $\{x_i, y_i\}$, for $i = 1, 2, \dots, k$. There are no edges inside X . Let $y_i y_j$ be an edge iff i, j are not consecutive elements of the circle \mathbb{Z}_k , and so $G[X^c]$ is the graph K_k minus a cycle C_k . Let $x_i y_j$ is an edge iff $i \not\equiv j \pmod{\frac{k}{p}}$, i.e. x_i is joined to all but p equally spaced y_j . We claim that G satisfies the required properties.

Clearly, we have $d(x) = k - p = 2r - 4$ for all $x \in X$ and $e(X, X^c) = k(2r - 4)$. Let us verify that G is K_r -free. A clique inside X^c is a set of non-consecutive elements of \mathbb{Z}_k , and so a largest clique inside X^c has size $\lfloor \frac{k}{2} \rfloor = r - 1$ for $p \in \{2, 3\}$. Since a clique which is not inside X^c can contain at most one vertex of X , it remains to check that the neighborhood of each x_i does not contain a clique of size $r - 1$. Viewing X^c as a circle, $N(x_i)$ consists of p segments of the circle, each of size $\frac{2r-4}{p}$, separated by gaps of size one. Since $\frac{2r-4}{p}$ is even, a largest clique in $N(x_i)$ has size $\frac{p(2r-4)}{2p} = r - 2$.

It remains to show that the admissible non-edges inside X , and those between X and X^c are K_r -saturated. Let $x_i y_j$ be an admissible non-edge, and so $j = i \pm \frac{k}{p}$ in \mathbb{Z}_k . Clearly, $N(x_i)$ contains $r - 2$ vertices which form a non-consecutive set of the circle with y_j . Therefore, there exists a K_{r-2} in the common neighborhood of x_i and y_j as required. Now let $x_i x_j$ be an admissible non-edge. Then the common neighborhood of x_i and x_j consists of $2p$ segments of the circle separated by gaps of size one such that they form p pairs where the sum of the sizes of each pair is $\frac{2r-4}{p} - 1$, and so each pair consists of a segment of even size and a segment of odd size. Therefore, a largest non-consecutive set in $N(x_i) \cap N(x_j)$ has size $\frac{p(2r-4)}{2p} = r - 2$. Hence, there exists a K_{r-2} in $N(x_i) \cap N(x_j)$ as required.

We have constructed such k -partite graph G_k for $k = 2r - 4 + p$ with . Let us obtain G_k for $k > 2r - 4 + p$ from G_{2r-4+p} by blowing up x_1 to a class $\{x_1\} \cup \{x_i : 2r - 3 + p \leq i \leq k\}$ of size $k - (2r - 4 + p) + 1$ where each copy of x_1 (not including itself) forms a part of G_k of size one. Clearly, we have $d(x) = 2r - 4$ for all $x \in X = \{x_1, x_2, \dots, x_k\}$ and $e(X, X^c) = k(2r - 4)$. Since G_{2r-4+p} is K_r -free, so is G_k .

It remains to check that the admissible non-edges inside X , and those between X and X^c are K_r -saturated. Any admissible non-edge inside X which is not inside the blow up class of x_1 is K_r -saturated by the same property of G_{2r-4+p} . Any admissible non-edge inside the blow up class of x_1 is K_r -saturated since $N(x_1)$ contains a K_{r-2} by the construction of G_{2r-4+p} . Any admissible non-edge $x_i y_j$

where $j \neq 1$ or ($j = 1$ and $i \leq 2r - 4 + p$), is K_r -saturated by the same property of G_{2r-4+p} . Any admissible non-edge $x_i y_j$ where $j = 1$ and $2r - 3 + p \leq i \leq k$, is K_r -saturated since $N(x_1) \cap N(y_1)$ contains a K_{r-2} by the construction of G_{2r-4+p} . \square

5 Proof of Theorem 2(*iii*)

In this section, we study $\alpha(k, r)$ for $r = 3, 4, 5$. The values of $\alpha(k, 3)$ and $\alpha(k, 4)$ are completely determined while the values of $\alpha(k, 5)$ are unknown for $k = 5, 6, 8$.

5.1 The function $\alpha(k, 3)$

We shall prove that $\alpha(k, 3) = 3(k - 1)$ for $k \geq 3$. The upper bound follows from Theorem 2(*i*). Let us prove the lower bound.

Let $G = V_1 \cup V_2 \cup \dots \cup V_k$ be a K_3 -partite-saturated k -partite graph G containing an independent set $X = \{x_1, x_2, \dots, x_k\}$ with $x_i \in V_i$ for all i . By Lemma 8, the deletion of any part of G does not destroy all vertices of $N(x_i)$ for all i , i.e. x_i is joined to at least two parts of G . Suppose for contradiction that $e(X, X^c) < 3(k - 1)$, i.e. X contains at least four vertices of degree 2, say x_1, x_2, x_3, x_4 . Let $y_i \in V_i$ and $y_j \in V_j$ with $1 < i < j \leq k$ be the neighbors of x_1 , and so y_i and y_j are not neighbors otherwise $x_1 y_i y_j$ forms a triangle. Since $\{2, 3, 4\} \setminus \{i, j\} \neq \emptyset$, we may assume that $i, j \neq 2$, i.e. x_1, x_2, y_i, y_j are from different parts of G . Since any pair in X forms a K_3 -saturated non-edge in G , they have a common neighbor. So x_1 and x_2 have a common neighbor, say y_i .

First we suppose that $x_2 y_j$ is a non-edge. Then x_2 and y_j have a common neighbor $y_l \in V_l$. Since y_i and y_j are not neighbors, $l \neq i$. We obtain a contradiction by observing that $x_i y_j y_l$ forms a triangle. We observe that $x_i y_j$ are neighbors since x_1 and x_i have a common neighbor and $N(x_1) = \{y_i, y_j\}$. Similarly, $x_i y_l$ are neighbors since x_2 and x_i have a common neighbor and $N(x_2) = \{y_i, y_l\}$.

Now, suppose that $x_2 y_j$ is an edge, and so $N(x_1) = N(x_2) = \{y_i, y_j\}$. Then $x_i y_j$ are neighbors since x_1 and x_i have a common neighbor. Similarly, $x_j y_i$ are neighbors. We know that x_i and x_j have a common neighbor y_l with $l \neq i, j$. Then either $l \neq 1$ or $l \neq 2$, say $l \neq 1$. Since the non-edge $x_1 y_l$ is K_3 -saturated, y_l is joined to either y_i or y_j . This implies a contradiction that either $x_j y_i y_l$ or $x_i y_j y_l$ forms a triangle. \square

5.2 The function $\alpha(k, 4)$

As a consequence of Theorem 2(*ii*), we obtain that $\alpha(k, 4) = 4k$ for $k \geq 5$. For the remaining case $k = 4$, we have the bounds $16 \leq \alpha(4, 4) \leq 18$ from Theorem 2(*i*). We shall show that $\alpha(4, 4) = 18$.

Consider the family of graphs appearing in the definition of $\alpha(r, r)$. Let $G = V_1 \cup V_2 \cup \dots \cup V_r$ be an K_r -partite-saturated r -partite graph G containing an independent set $X = \{x_1, x_2, \dots, x_r\}$ with $x_i \in V_i$ for all i . We shall establish some properties of G which will be useful in this subsection, the next subsection and Section 6.

We say that a vertex $y \in X^c$ is *i-special* if y is the only neighbor of x_i in the part of G containing y . The *special degree* of a vertex $y \in X^c$ is the number of $i \in [r]$ such that y is *i-special*. We say that a vertex $y \in X^c$ is *special* if the special degree of y is at least one. Let us make some easy observations regarding the special vertices.

Lemma 18. *Let $G = V_1 \cup V_2 \cup \dots \cup V_r$ be an K_r -partite-saturated r -partite graph G containing an independent set $X = \{x_1, x_2, \dots, x_r\}$ with $x_i \in V_i$ for all i . The following hold for $r \geq 4$.*

- (i) *A special vertex $y_i \in V_i$ is joined to every vertex of X except x_i .*
- (ii) *Each V_i contains at most one special vertex.*
- (iii) *If $y_i \in V_i$ is i' -special and $y_j \in V_j$ is j' -special with $i' \neq j$ and $j' \neq i$ then $y_i y_j$ is an edge.*
- (iv) *The number of vertices of special degree at least 2 is at most $r - 2$.*
- (v) *If $y_i \in V_i$ is i' -special and $y_j \in V_j$ with $j \neq i, i'$ then y_j is joined to either y_i or $x_{i'}$.*
- (vi) *For a special vertex $y_i \in V_i$, there exist parts V_j and V_l where i, j, l are distinct such that $N(x_i) \cap V_j$ and $N(x_i) \cap V_l$ both contain a non-neighbor of y_i .*

Proof. (i) Let $y_i \in V_i$ be i' -special and let $j \neq i, i'$. Since the non-edge $x_{i'} x_j$ is K_r -saturated, the common neighborhood of $x_{i'}$ and x_j contains a K_{r-2} consisting of one vertex from each part of $G \setminus (V_{i'} \cup V_j)$. Then y_i is in this K_{r-2} since y_i is the only neighbor of $x_{i'}$ in V_i , and so y_i is joined to x_j .

(ii) Suppose for contradiction that V_i contains two special vertices y_i and z_i where y_i is i' -special. Then, by (i), $x_{i'}$ is joined to both y_i and z_i contradicting the fact that y_i is the only neighbor of $x_{i'}$ in V_i .

(iii) First, suppose that $i' \neq j'$. Since the non-edge $x_{i'} x_{j'}$ is K_r -saturated, the common neighborhood of $x_{i'}$ and $x_{j'}$ contains a K_{r-2} consisting of one vertex from each part $G \setminus (V_{i'} \cup V_{j'})$. Since y_i is the only neighbor of $x_{i'}$ in V_i and y_j is the only neighbor of $x_{j'}$ in V_j , both y_i and y_j lie in this K_{r-2} . In particular, $y_i y_j$ is an edge.

Now, suppose that $i' = j'$. We can pick $l \neq i, j, i'$ because $r \geq 4$. Since the non-edge $x_{i'} x_l$ is K_r -saturated, the common neighborhood of $x_{i'}$ and x_l contains a K_{r-2} consisting of one vertex from each part of $G \setminus (V_{i'} \cup V_l)$. Since y_i is the only neighbor of $x_{i'}$ in V_i and y_j is the only neighbor of $x_{i'}$ in V_j , both y_i and y_j lie in this K_{r-2} . In particular, $y_i y_j$ is an edge.

(iv) Suppose for contradiction that there exist vertices y_1, y_2, \dots, y_{r-1} of special degree at least 2. By (ii), they lie in different parts of G , say $y_i \in V_i$ for $1 \leq i \leq r-1$. We claim that they form a K_{r-1} which would be a contradiction since, together with x_r , they form a K_r by (i). Now we show that any $y_i y_j$ is an edge. Since y_i and y_j have special degree at least 2, there exist $i' \neq j$ and $j' \neq i$ such that y_i is i' -special and y_j is j' -special. Therefore, $y_i y_j$ is an edge by (iii).

(v) Suppose that $x_{i'} y_j$ is a non-edge. Then the common neighborhood of $x_{i'}$ and y_j contains a K_{r-2} consisting of one vertex from each part of $G \setminus (V_{i'} \cup V_j)$. Then y_i is in this K_{r-2} since y_i is the only neighbor of $x_{i'}$ in V_i , and so y_i is joined to y_j .

(vi) Suppose for contradiction that there exists $j \in [r] \setminus \{i\}$ such that $y_i \in V_i$ is joined to every vertex in $N(x_i) \cap V_l$ for all $l \neq i, j$. Since the non-edge $x_i x_j$ is K_r -saturated, the common neighborhood of x_i and x_j contains a K_{r-2} consisting of one vertex from each part of $(G \setminus X) \setminus (V_i \cup V_j)$. We obtain a contradiction by observing that this K_{r-2} , together with x_j and y_i , form a K_r . Indeed, by assumption, this K_{r-2} is also in the neighborhood of y_i and $x_j y_i$ is an edge by (i). \square

Now we are ready to show that $\alpha(4, 4) \geq 18$. Suppose for contradiction that $\alpha(4, 4) \leq 17$, i.e. there exists a K_4 -partite-saturated 4-partite graph $G = V_1 \cup V_2 \cup V_3 \cup V_4$ containing an independent set $X = \{x_1, x_2, x_3, x_4\}$ with $x_i \in V_i$ for all i such that $\sum_{i=1}^4 d(x_i) \leq 17$. By Lemma 8, $d(x_i) \geq \beta_1(3, 3) = 4$ and each x_i has some neighbor in V_j for $j \neq i$. Therefore, there are at least three vertices of degree 4 and possibly one of degree 5. Since a vertex of degree 4 in X creates at least two special vertices and a vertex of degree 5 in X creates at least one special vertex, the sum of the special degrees of the vertices in X^c is at least $2 + 2 + 2 + 1 = 7$. By Lemma 18(iv), there is a vertex of special degree 3, say $y_1 \in V_1$.

For $i = 2, 3, 4$, since y_1 is i -special, x_i has at least three neighbors in $N(y_1) \cup \{y_1\}$, each in a different part of G , by Lemma 8. On the other hand, y_1 has at least two non-neighbors, say $y_2 \in V_2$ and $y_3 \in V_3$, by Lemma 18(vi). By Lemma 18(v), $x_i y_2$ is an edge for $i \neq 2$ and $x_i y_3$ is an edge for $i \neq 3$. So x_4 has five neighbors, i.e. y_2, y_3 and three vertices in $N(y_1) \cup \{y_1\}$, and $d(x_1) = d(x_2) = d(x_3) = 4$. Since x_2 has four neighbors including y_3 and it has some neighbor in $(N(y_1) \cup \{y_1\}) \cap V_j$ for each $j = 1, 3, 4$, it has exactly one neighbor in V_4 , say y_4 . Similarly, x_3 has exactly one neighbor in V_4 which has to be the same vertex y_4 by Lemma 18(ii).

We obtain a contradiction by observing that $x_1 y_2 y_3 y_4$ forms a K_4 . First, note that $x_1 y_4$ is an edge by Lemma 18(i). Now y_4 is not 1-special otherwise y_4 would have special degree 3 and by repeating the argument above with y_1 replaced by y_4 , we could deduce that x_1, x_2 , or x_3 had degree 5. Therefore, the neighbors of x_1 are y_2, y_3, y_4 and a vertex in V_4 . Since y_2, y_3 are both 1-special and y_4 is 2, 3-special, $y_2 y_3 y_4$ forms a triangle by Lemma 18(iii). \square

5.3 The function $\alpha(k, 5)$

As a consequence of Theorem 2(i) and (ii), we obtain that

$$\begin{aligned} \alpha(k, 5) &= 6k \quad \text{for } k = 7 \text{ or } k \geq 9, \\ 30 &\leq \alpha(5, 5) \leq 36, \\ 36 &\leq \alpha(6, 5) \leq 40, \\ 48 &\leq \alpha(8, 5) \leq 49. \end{aligned}$$

We shall improve the lower bound for $\alpha(5, 5)$ to 33.

Suppose for contradiction that $\alpha(5, 5) \leq 32$, i.e. there exists a K_5 -partite-saturated 5-partite graph $G = V_1 \cup V_2 \cup V_3 \cup V_4 \cup V_5$ containing an independent set $X = \{x_1, x_2, x_3, x_4, x_5\}$ with $x_i \in V_i$ for all i such that $\sum_{i=1}^5 d(x_i) \leq 32$. Write Y_i for $V_i \setminus \{x_i\}$. By Lemma 8, $d(x_i) \geq \beta_1(4, 4) = 6$ and each x_i has some neighbor in V_j for $j \neq i$. Therefore, there are either four vertices in X of degree 6 or there

are three vertices of degree 6 and two of degree 7. Since a vertex of degree 6 in X creates at least two special vertices and a vertex of degree 7 in X creates at least one special vertex, the sum of the special degrees of the vertices in X^c is at least 8, and hence, there exists a vertex of special degree at least two. Let i be such that there is a special vertex $y \in Y_i$ with special degree $d_s(y)$ at least two where $(d(x_i), d_s(y))$ is maximum in lexicographical order¹. Without loss of generality we can assume that $i = 1$. Let $N = N(y) \setminus X$. By Lemma 18(vi), x_1 has two neighbours, say y_2, y_3 , belonging to two distinct parts of G , different from V_1 , which are non-neighbours of y . Without loss of generality, we can assume that $y_2 \in Y_2$ and $y_3 \in Y_3$.

For a pair of non-adjacent vertices $u, v \in G$ and $S \subset G$, we say that S is uv -saturating if adding the edge of uv to G creates a copy K of K_5 such that $S \subseteq K$. If $S = \{z\}$ then we simply say that z is uv -saturating. Notice that if S is uv -saturating then S induces a clique.

In the rest of the proof, we shall repeatedly use the following lemma.

Lemma 19. *Given $i \in \{2, 3, 4, 5\}$ the following hold.*

- (i) *If $j \in \{2, 3, 4, 5\} \setminus \{i\}$ then x_i has a neighbour in $V_j \cap N$. In particular, $d_N(x_i) \geq 3$.*
- (ii) *If y is i -special then x_i is adjacent to y_j for every $j \in \{2, 3\} \setminus \{i\}$.*
- (iii) *If y is $x_i x_j$ -saturating, for every $j \in \{2, 3, 4, 5\} \setminus \{i\}$, then $d_N(x_i) \geq 4$.*
- (iv) *If y is i -special or $d_s(y) \geq 3$ then $d_N(x_i) \geq 4$.*
- (v) *If y is 2, 3-special and $i \in \{4, 5\}$, then $d(x_i) \geq 7$.*
- (vi) *If $i \in \{2, 3\}$ and there are p vertices in $X \setminus \{x_1\}$ all of which have neighbours in $Y_i \setminus N$ then there is no vertex in V_i with special degree bigger than $\max\{1, 3 - p\}$.*
- (vii) $Y_3 \cup Y_4 \subset N$.

Proof. (i) Observe that we can choose $k \in \{2, 3, 4, 5\} \setminus \{i, j\}$ such that y is either i -special or k -special. Since there must be a triangle in the common neighbourhood of x_i and x_j which uses y , we have that the remaining two vertices belong to N . Hence x_i has a neighbour in $N \cap x_j$.

(ii) This follows directly from Lemma 18(v).

(iii) We shall show that $d_N(x_i) \geq \beta_1(3, 3) = 4$. Take any $j \in \{2, 3, 4, 5\} \setminus \{i\}$. Since y is $x_i x_j$ -saturating then there is an edge in the common neighbourhood of x_i and x_j in $N \setminus (V_i \cup V_j)$. Observe that the common neighbourhood of x_i and y cannot contain a K_3 , hence $d_N(x_i) \geq \beta_1(3, 3) = 4$.

(iv) Take any $j \in \{2, 3, 4, 5\} \setminus \{i\}$. Since y is either i - or j -special, it follows that y is $x_i x_j$ -saturating. Hence, by (ii), $d_N(x_i) \geq 4$.

(v) Without loss of generality we can assume that $i = 4$. If y is also 4-special then it follows from (ii) and (iv) that $d_N(x_4) \geq 4$ and x_4 is adjacent to y, y_2, y_3 , therefore $d(x_4) \geq 7$. Hence we can assume that y is not 4-special. Suppose for contradiction that $d(x_4) = 6$. From (i), we have that $d_N(x_4) \geq 3$

¹We say that $(a, b) \preceq (c, d)$ if $a < c$ or $a = c$ and $b \leq d$, where \preceq denotes the lexicographical order relation.

and since y is not 4-special we have that $d_{Y_1}(x_4) \geq 2$. Moreover, x_4 has to have at least one neighbour not in $Y_1 \cup N$ as otherwise there would be a copy of K_5 in G , as seen by considering the non-edge x_1x_4 . Therefore, $d(x_4) = d_{Y_1}(x_4) + d_N(x_4) + |N(x_4) \setminus (Y_1 \cup N)| \geq 3 + 2 + 1 = 6 = d(x_4)$. Hence, $d_{Y_1}(x_4) = 3$, $d_N(x_4) = 4$ and $|N(x_4) \setminus (Y_1 \cup N)| = 1$. We shall obtain a contradiction by finding a copy of K_5 in the graph G .

Suppose $\{z_1, z_2, z_3\}$ is x_4x_5 saturating, with $z_i \in V_i$. We claim that $y \neq z_1$ and $\{z_2, z_3\} \not\subseteq N$. Suppose for contradiction that it is not the case. If y is x_4x_5 -saturating then from (iii) we have that $d_N(x_4) \geq 4$ hence we obtain a contradiction. We can therefore assume that y is not x_4x_5 -saturating and hence $z_1 \neq y$. Whence $z_2, z_3 \in N$. Recall that $\{z_1, z_2, z_3\}$ form a triangle and therefore there is an edge between z_2, z_3 . By assumption z_2 and z_3 are neighbours of y , hence y, z_2, z_3 form a triangle, and therefore y is x_4x_5 -saturating since y, z_2, z_3 belong to the common neighbourhood of x_4 and x_5 , which contradicts the assumption that y is not x_4x_5 -saturating.

Without loss of generality we can assume that $z_2 \notin N$. Using (i), we can therefore suppose that $N(x_4) \cap Y_1 = \{y, z_1\}$, $N(x_4) \cap Y_2 = \{w, z_2\}$, $N(x_4) \cap Y_3 = \{z_3\}$ and $N(x_4) \cap Y_5 = \{z_5\}$, for some $w, z_3, z_5 \in N$. We shall obtain a contradiction by observing that z_1, z_2, z_3, x_4, z_5 form a copy of K_5 . First we claim that $\{z_2, z_3, z_5\}$ is x_1x_4 -saturating. Indeed, there must be a triangle in the common neighbourhood of x_1 and x_4 , with one vertex in each V_3, V_4, V_5 . There are only two candidates for the triangle: z_2, z_3, z_5 or w, z_3, z_5 . It cannot be w, z_3, z_5 since they are all neighbours of y , hence y, w, z_3, x_4, z_5 would form a copy of K_5 . Hence we must have that the set $\{z_2, z_3, z_5\}$ is x_1x_4 -saturating. Now, since x_4 is not adjacent to y_3 , and y_3 is not adjacent to y we must have an edge between z_1 and z_5 . Indeed, there must be a triangle in the common neighbourhood of x_4 and y_3 with a vertex in each V_1, V_2, V_3 . Since x_4 has only one neighbour in V_5 , i.e. z_5 , and x_4 and y_3 have only one common neighbour in V_1 , i.e. z_1 , we must have an edge between z_1 and z_5 .

Therefore we have that z_1, z_2, z_3 form a triangle, z_2, z_3, z_5 form a triangle, and z_1, z_5 are adjacent. It easy to see now that z_1, z_2, z_3, x_4, z_5 form a copy of K_5 .

(vi) Let v be a special vertex in $V_2 \cup V_3$, say in V_2 . First observe that if v is 1-special then x_3, x_4, x_5 are all adjacent to $y_2 \in Y_2 \setminus N$. On the other hand, it follows from (i) that x_3, x_4, x_5 all have neighbours in $N \cap Y_2$ hence they all have degree at least 2 in Y_2 . It follows that v has special degree 1. If we assume that v is not 1-special then v has special degree at most $3 - p$, since p of the vertices x_3, x_4, x_5 have degree 2 in Y_2 .

(vii) Assume for contradiction that there is v , say in $Y_4 \setminus N$. Observe that if y is i -special then it follows from (ii) and (iv) that $d(x_i) \geq 7$, hence if $d_s(y) \geq 3$ we obtain contradiction by finding three vertices in X of degree at least 7. Therefore we can assume that $d_s(y) = 2$.

If y is 5, i -special, then from (ii) and (iv) we have that $d(x_5) \geq 8$ and $d(x_i) \geq 7$ hence again we obtain a contradiction. Therefore we can assume that y is not 5-special. If y is 2, 3-special then $d(x_2), d(x_3) \geq 7$ and from (iv) we have that $d(x_4), d(x_5) \geq 7$. Hence we can assume that y is 2, 4-special or 3, 4-special. Suppose that the former is the case. Then $d(x_2), d(x_4) \geq 7$. It follows that $d(x_1) = 6$. Therefore by maximality (x_1, y) and from (v) we have that every vertex in $Y_2 \cup Y_3 \cup Y_4$ has special degree at most 1 and no vertex in Y_5 has special degree bigger than 2. Which gives a contradiction since the sum of special degree is then at most 7. \square

We are now ready to finish showing that $\alpha(5, 5) \geq 33$. We consider several cases depending on the special degree of y .

Case 1. $d_s(y) = 4$

Consider the 4-partite graph $H = G[N(y)]$ with an independent set $X' = \{x_2, x_3, x_4, x_5\}$. Clearly, H is K_4 -free since G is K_5 -free. We modify H by keeping adding admissible edges inside $H \setminus X'$ until every admissible non-edge inside $H \setminus X'$ is K_4 -saturated. We claim that H is K_4 -partite-saturated, which would imply that $e(X', H \setminus X') \geq \alpha(4, 4) = 18$ by the previous subsection. It remains to show that the admissible non-edges with at least one endpoint in X' are K_4 -saturated.

Consider the non-edge $x_i y_j$ with $y_j \in V_j \cap H$ (possibly $y_j = x_j$) and distinct $2 \leq i, j \leq 5$. Since the non-edge $x_i y_j$ is K_5 -saturated in G , the common neighborhood in G of x_i and y_j contains a K_3 consisting of one vertex from each part of $G \setminus (V_i \cup V_j)$. Since y is i -special, this K_3 must contain y , and so the common neighborhood in H of x_i and y_j contains a K_2 , i.e. $x_i y_j$ is K_4 -saturated in H as required.

Recall that y has two non-neighbors, $y_2 \in V_2$ and $y_3 \in V_3$. By Lemma 18(v), $x_i y_2$ is an edge for $i \neq 2$ and $x_i y_3$ is an edge for $i \neq 3$. We shall partition the edges between X and X^c as follows:

$$\begin{aligned} e(X, X^c) &\geq e(X', H \setminus X') + d(x_1) + e(X, y) + e(X', y_2) + e(X', y_3) \\ &\geq 18 + 6 + 4 + 3 + 3 = 34, \end{aligned}$$

contradicting the assumption.

Case 2. $d_s(y) = 3$

If y is 4, 5-special then from Lemma 19(ii) and 19(iii) we have that $d(x_4), d(x_5) \geq 7$. Otherwise y is 2, 3-special and hence it follows from Lemma 19(v) that $d(x_4), d(x_5) \geq 7$. We shall obtain a contradiction by showing that $d(x_1) \geq 7$, hence showing that there are three vertices in X with degrees at least 7, which is against an assumption made in the beginning of the subsection. It follows from Lemma 19(vi) with $p \geq 2$, that the sum of special degrees in $Y_2 \cup Y_3$ is at most 2. Since the sum of special degrees is at least 8, it follows that there is a special vertex in $Y_4 \cup Y_5$ with special degree at least 2. Therefore from the maximality of $d(x_1)$ we have that $d(x_1) \geq 7$.

Case 3. $d_s(y) = 2$

We split this case into three subcases.

Case 3.1. y is 2, 3-special

It follows from Lemma 19(v) that $d(x_4), d(x_5) \geq 7$. We shall obtain a contradiction by showing that $d(x_1) \geq 7$, hence showing that there are three vertices in X with degrees at least 7, which is against an assumption made in the beginning of the subsection. It follows from Lemma 19(vi) that the sum of special degrees in $Y_2 \cup Y_3$ is at most 2. Since the sum of special degrees is at least 8, it follows that there is a special vertex in $Y_4 \cup Y_5$ with special degree at least 2. Therefore from the maximality of $d(x_1)$ we have that $d(x_1) \geq 7$.

Case 3.2. y is 4, 5-special

It follows from Lemma 19(ii) and 19(iv) that $d(x_4), d(x_5) \geq 7$. We shall obtain a contradiction by showing that $d(x_1) \geq 7$, hence showing that there are three vertices in X with degrees at least 7, which is against an assumption made in the beginning of the subsection. It follows from Lemma 19(vi) that the sum of special degrees in $Y_2 \cup Y_3$ is at most 3. Since the sum of special degrees is at least 8, it follows that there is a special vertex in $Y_4 \cup Y_5$ with special degree at least 2. Therefore from the maximality of $d(x_1)$ we have that $d(x_1) \geq 7$.

Case 3.3. y is neither 2, 3-special nor 4, 5-special

Without loss of generality we can assume that y is 2, 4-special. It follows from Lemma 19(ii) and 19(iv) that $d(x_4) \geq 7$ and from Lemma 19(vi) with $p \geq 2$ that there is no special vertex in $Y_2 \cup Y_3$ with special degree bigger than 1. Hence there is either a vertex in Y_4 with special degree at least 2 or a vertex in Y_5 with special degree at least 3. Therefore we can assume that $d(x_1) = 7$ as otherwise we obtain a contradiction to the maximality of $(d(x_1), d_s(y))$.

We shall obtain a contradiction by showing that at least one of x_2, x_3 or x_5 has degree at least 7, thus finding three vertices with degree at least 7. Suppose $d(x_2) = d(x_3) = d(x_5) = 6$. Observe that if there is a vertex in X_4 of special degree bigger than 2 then we obtain a contradiction to the maximality of $(d(x_1), d_s(y))$. Therefore there are two vertices in X with at least two neighbours in X_4 . Suppose that $i \in \{3, 5\}$ and x_i has at least two neighbours in X_4 . Then it follows from Lemma 19(i) that $d_N(x_i) \geq 4$, and hence x_i has degree at least 7 as x_i has at least three neighbours outside N . We can therefore assume that x_3 and x_5 have only one neighbour in X_4 . For the same reason we can assume that x_3 has only one neighbour in Y_5 . If x_2 has two neighbours in Y_5 then $d_N(x_2) \geq 5$ and therefore $d(x_2) \geq 7$. Hence we can assume that there is $z_5 \in Y_5$ which is 2, 3-special.

Suppose $\{z_1, z_2, z_4\}$ is x_3x_5 -saturating, with $z_i \in V_i$. We claim that $y \neq z_1$ and $z_2 \notin N$. Suppose for contradiction that it is not the case. If y is x_3x_5 -saturating then from (iii) we have that $d_N(x_3) \geq 4$ hence we obtain a contradiction. We can therefore assume that y is not x_3x_5 -saturating and hence $z_1 \neq y$. Whence $z_2 \in N$. Observe that by Lemma 19(vii) we have $z_4 \in N$. Recall that $\{z_1, z_2, z_4\}$ form a triangle and therefore there is an edge between z_2, z_4 . By assumption z_2 and z_4 are neighbours of y , hence y, z_2, z_4 form a triangle, and therefore y is x_3x_5 -saturating since y, z_2, z_4 belong to the common neighbourhood of x_3 and x_5 , which contradicts the assumption that y is not x_3x_5 -saturating.

We shall obtain a contradiction by showing that z_1, z_2, x_3, z_4, z_5 form a copy of K_5 . Indeed, by assumption $\{z_1, z_2, z_4\}$ is x_3x_5 -saturating and similar analysis to the one made in the proof of Lemma 19(v) shows that $\{z_2, z_4, z_5\}$ is x_1x_3 -saturating. Since y is 2-special it follows that x_2 is not adjacent to z_1 , and moreover z_5 , as the only neighbour of x_2 in Y_5 , is x_2z_1 -saturating, and therefore there is an edge between x_2 and z_5 . Hence we have that z_2, z_4, z_5 form a triangle, z_1, z_2, z_4 form a triangle, and z_1, z_5 are adjacent. It easy to see now that z_1, z_2, x_3, z_4, z_5 form a copy of K_5 .

□

6 The diagonal case $\alpha(r, r)$

6.1 Proof of Theorem 2(iv)

We have seen that the lower bound $\alpha(k, r) \geq k(2r - 4)$ in Theorem 2(i) is attained for some k . In this subsection, we show that this is not the truth for the diagonal case $k = r \geq 4$, i.e. $\alpha(r, r) \geq r(2r - 4) + 1$. We shall again use the concept of special vertices introduced in Section 5.

Suppose for contradiction that for some $r \geq 4$, $\alpha(r, r) = r(2r - 4)$, i.e. there exists a K_r -partite-saturated r -partite graph $G = V_1 \cup V_2 \cup \dots \cup V_r$ containing an independent set $X = \{x_1, x_2, \dots, x_r\}$ with $x_i \in V_i$ for all i such that $\sum_{i=1}^r d(x_i) = r(2r - 4)$. Lemma 8 tells us that we must have $d(x_i) = 2r - 4$ for all i and each x_i has some neighbor in V_j for $j \neq i$. Therefore, each x_i creates at least two special vertices, and so the sum of the special degrees of the vertices in X^c is at least $2r$. By Lemma 18(iv), there is a vertex of special degree at least 3, say $y_1 \in V_1$.

We observe that y_1 has at least two non-neighbors, say $y_2 \in V_2$ and $y_3 \in V_3$ by Lemma 18(vi). Since y_1 has special degree at least 3, we can pick $i \geq 4$ such that y_1 is i -special. By Lemma 18(v), y_2 and y_3 are neighbors of x_i . Therefore,

$$|N(x_i) \cap N(y_1)| = d(x_i) - |N(x_i) \setminus N(y_1)| \leq (2r - 4) - 3 = 2r - 7.$$

On the other hand, we shall obtain a contradiction by showing that the graph $H = G[N(x_i) \cap N(y_1)]$ contains at least $\beta_1(r - 2, r - 2) = 2(r - 3)$ vertices. It is sufficient to prove that H is an $(r - 2)$ -partite K_{r-2} -free graph such that the subgraph induced by any $k - 3$ parts contains a K_{r-3} . Clearly, H is K_{r-2} -free since G is K_r -free. The parts of H are $N(x_i) \cap N(y_1) \cap V_j$ for $j \notin [r] \setminus \{1, i\}$. It remains to verify that the deletion of the part $N(x_i) \cap N(y_1) \cap V_j$ does not destroy all the K_{r-3} . Since the non-edge $x_i x_j$ is K_r -saturated in G , the common neighborhood in G of x_i and x_j contains a K_{r-2} consisting of one vertex from each part of $G \setminus (V_i \cup V_j)$. Since y_1 is i -special, this K_{r-2} must contain y_1 , and so the common neighborhood $N(x_i) \cap N(y_1) \cap N(x_j) \subset H$ contains a K_{r-3} not using the vertices of V_j as required. \square

6.2 Remark on $\beta_2(r, r - 1)$

Recall from Proposition 7 that $\alpha(r, r) \leq (r - 1)\beta_2(r, r - 1)$. Thus, a better estimate on β_2 would translate to a better understanding of the saturation numbers. While we could not find the exact value of $\beta_2(r, r - 1)$, we suspect that $\beta_2(r, r - 1) = 3r - 6$ as mentioned in Conjecture 13. In this subsection, we make an observation about $\beta_2(r, r - 1)$ which can be viewed as a first step towards determining its exact value. For simplicity of notation, let us write $\beta_2(r) = \beta_2(r, r - 1)$.

Proposition 20. *Either*

- $\beta_2(r) = 3r - 6$ for all $r \geq 3$, or
- $\beta_2(r) \leq (c + o(1))r$ for some constant $c < 3$, as $r \rightarrow \infty$.

Proof. The result is an immediate consequence of the following lemma.

Lemma 21. $\beta_2(r_1 + r_2) \leq \beta_2(r_1) + \beta_2(r_2) + 6$ for $r_1, r_2 \geq 3$.

Proof. For $i \in \{1, 2\}$, let $G_i = V_{i,1} \cup V_{i,2} \cup \dots \cup V_{i,r_i}$ be a K_{r_i-1} -free r_i -partite graph on $\beta_2(r_i)$ vertices such that the subgraph induced by any $r_i - 2$ parts contains a K_{r_i-2} . We shall construct a $K_{r_1+r_2-1}$ -free $(r_1 + r_2)$ -partite graph G from G_1 and G_2 with $|G| = |G_1| + |G_2| + 6$ by starting with the disjoint union of G_1 and G_2 and then adding six new vertices $U = \{x_1, x_2, y_1, y_2, z_1, z_2\}$ as follows: add x_i, y_i to $V_{i,1}$ and add z_i to $V_{i,2}$ for $i \in \{1, 2\}$. Now, join all admissible pairs between U and $V(G) \setminus U$, and add the edges $x_1z_1, x_2z_2, y_1y_2, z_1z_2, y_1z_2, z_1y_2$ inside U .

First, we show that G is $K_{r_1+r_2-1}$ -free. Suppose otherwise. Since G_i is K_{r_i-1} -free for $i \in \{1, 2\}$, this $K_{r_1+r_2-1}$ must contain at least three vertices forming a triangle in U , contradicting the fact that $G[U]$ is triangle-free. It remains to show that the deletion of any two parts does not destroy all the $K_{r_1+r_2-2}$. Suppose first that both deleted parts are from G_1 . Since G_1 contains a K_{r_1-2} not using these two parts and G_2 contains a K_{r_2-2} not using $V_{2,1}$ and $V_{2,2}$, we obtain a $K_{r_1+r_2-2}$ not using the deleted parts, formed by these two cliques and x_2, z_2 . Now suppose that one of the deleted parts is from G_1 and the other is from G_2 . For $i \in \{1, 2\}$, let V_i be a part in $\{V_{i,1}, V_{i,2}\}$ which was not deleted. By construction, $G[U]$ contains an edge between $V_{1,j}$ and $V_{1,l}$ for all $j, l \in \{1, 2\}$ and so there exists an edge in $G[U]$ between V_1 and V_2 , say e . Since G_1 contains a K_{r_1-2} not using the deleted part in G_1 and V_1 , and G_2 contains a K_{r_2-2} not using the deleted part in G_2 and V_2 , we obtain a $K_{r_1+r_2-2}$ not using the deleted parts, formed by these two cliques and the endpoints of e . \square

Suppose that $\beta_2(s) < 3s - 6$ for some $s \geq 3$. We shall show that $\beta_2(r) \leq (c + o(1))r$ with $c = \frac{\beta_2(s) + 6}{s} < 3$. Applying the lemma and induction on m , we deduce that $\beta_2(ms) \leq cms - 6$ for all positive integer m . Hence, writing $r = ms + t$ with $3 \leq t \leq s + 2$ and applying the lemma again,

$$\beta_2(r) \leq \beta_2(ms) + \beta_2(t) + 6 \leq cms + d \leq \left(c + \frac{d}{r}\right)r = (c + o(1))r$$

where $d = \max\{\beta_2(t) : 3 \leq t \leq s + 2\}$. \square

7 Proof of Theorem 3

Theorems 1 and Theorem 2(ii) imply that

$$\text{sat}(n, k, r) = k(2r - 4)n + o(n) \text{ if } \begin{cases} k = 2r - 3, \text{ or} \\ k \geq 2r - 2 \text{ and } r \equiv 0 \pmod{2}, \text{ or} \\ k \geq 2r - 1 \text{ and } r \equiv 2 \pmod{3}. \end{cases}$$

In this section, we shall show that the $o(n)$ term can be replaced with $O(1)$. The upper bound follows from Proposition 5 and Theorem 2(ii). We prove that the lower bound holds for any $k \geq r \geq 3$ using the fact that $\beta_1(k - 1, r - 1) = 2r - 4$.

Proposition 22. For $k \geq r \geq 3$, there is an integer $C_{k,r}$ such that $\text{sat}(n, k, r) \geq k(2r - 4)n + C_{k,r}$, for every integer $n \geq 0$.

Proof. Suppose, as we may, that n is sufficiently large. Let $G = V_1 \cup V_2 \cup \dots \cup V_k$ be a K_r -partite-saturated k -partite graph with $|V_i| = n$ for all i . We shall find a subset U of $V(G)$ of constant size such that every vertex in U^c has at least $2r - 4$ neighbors in U . Then we would be done since $e(G) \geq e(U, U^c) \geq (2r - 4)(kn - |U|)$. Let v_1 be a vertex of smallest degree in V_1 . Having defined v_1, v_2, \dots, v_{i-1} , let $v_i \in V_i$ be a vertex of smallest degree in $V_i \setminus (N(v_1) \cup N(v_2) \cup \dots \cup N(v_{i-1}))$. We shall take U to be $N(v_1) \cup N(v_2) \cup \dots \cup N(v_k)$. Now we may assume that $d(v_i) < 2k(2r - 4)$ for all $1 \leq i \leq k$. Indeed, if v_i is the first vertex in the sequence such that $d(v_i) \geq 2k(2r - 4)$ then we are done since

$$e(G) \geq e(V_i, V_i^c) \geq d(v_i) \left(n - \sum_{j < i} d(v_j) \right) \geq 2k(2r - 4) \left(n - 2k(2r - 4)(i - 1) \right) \geq k(2r - 4)n$$

for sufficiently large n . Therefore, U has size bounded by a function of k and r . It remains to show that every vertex $v \in U^c$ has at least $2r - 4$ neighbors in U . We shall prove that $H = G[N(v) \cap U]$ contains at least $\beta_1(k - 1, r - 1) = 2r - 4$ vertices by showing that H is a K_{r-1} -free $(k - 1)$ -partite graph such that the subgraph induced by any $k - 2$ parts contains a K_{r-2} . Clearly, H is K_{r-1} -free since G is K_r -free. Without loss of generality, $v \in V_1$. The parts of H are $N(v) \cap U \cap V_i$ for $2 \leq i \leq k$. The deletion of the part $N(v) \cap U \cap V_i$ does not destroy all the K_{r-2} since the non-edge vv_i is K_{r-1} -saturated in G , i.e. $N(v) \cap N(v_i) \subset H$ contains a K_{r-2} not using the vertices of V_i . \square

8 Concluding remarks

We have reduced the problem of determining $\text{sat}(n, k, r)$ for large n to that of $\alpha(k, r)$. Although, we have determined $\alpha(k, r)$ for some values of k and r , a large number of cases remain unknown. In particular, the seemingly easiest case when r is fixed and k is large, is still open.

Problem 23. Determine $\alpha(k, r)$ for $k \geq 2r - 2$ and $r \equiv 1, 3 \pmod{6}$.

For $k \geq 2r - 2$ and $r \equiv 0, 2, 4, 5 \pmod{6}$, we have determined $\alpha(k, r)$ except one missing case when 3 is the smallest divisor of $r - 2$ and $k = 2r - 2$. Theorem 2(i) implies that $\alpha(2r - 2, r) \in \{(2r - 3)^2, (2r - 3)^2 - 1\}$ and we suspect that $\alpha(2r - 2, r) = (2r - 3)^2$.

Not only we believe that $\beta_2(k, r) = 4r - k - 2$ for $r < k \leq 2r - 1$ (see Conjecture 13) but we also think that the upper bound $\alpha(k, r) \leq (k - 1)\beta_2(k, r - 1) \leq (k - 1)(4r - k - 6)$ in Theorem 2(i) is the correct value for $\alpha(k, r)$ in this case.

Conjecture 24. $\alpha(k, r) = (k - 1)(4r - k - 6)$ for $5 \leq r \leq k \leq 2r - 4$.

We have shown that $33 \leq \alpha(5, 5) \leq 36$. This is the smallest case for which the value of α is not yet known.

Problem 25. Find $\alpha(5, 5)$.

To prove the lower and upper bounds on $\alpha(k, r)$, we extensively used the bounds on $\beta_1(k, r)$ and $\beta_2(k, r)$. We believe that determining the values of $\beta_i(k, r)$ is an interesting problem on its own.

Problem 26. Determine $\beta_i(k, r)$ for $k \geq r \geq 2$ and $2 \leq i \leq k - r + 1$.

We end the paper with a remark on a related problem. Recall that $\text{sat}(n, K_r)$ is the minimum number of edges in a K_r -free graph on n vertices but the addition of an edge joining any two non-adjacent vertices creates a K_r . In the pioneer paper of Erdős, Hajnal, and Moon [5], they determined $\text{sat}(n, K_r)$ by considering a more general problem where the graphs were not required to be K_r -free. Interestingly, the two problems have the same answer since the extremal graph is K_r -free. We remark that this phenomenon does not happen for partite saturation. Roberts [13] studied the corresponding more general problem for $\text{sat}(K_{r \times n}, K_r)$ and showed that the minimum number of edges in a K_r -saturated subgraph of $K_{r \times n}$ where the subgraph is allowed to contain K_r is $\binom{r}{2}(2n - 1)$ for $r \geq 4$ and sufficiently large n . On the other hand, Theorem 1 and Theorem 2 imply that $\text{sat}(K_{r \times n}, K_r) \geq r(2r - 4)n + o(n) > \binom{r}{2}(2n - 1)$ for sufficiently large n .

References

- [1] B. BOLLOBÁS, *On generalized graphs*, Acta Math. Acad. Sci. Hungar., 16 (1965), pp. 447–452.
- [2] ———, *Determination of extremal graphs by using weights*, Wiss. Z. Techn. Hochsch. Ilmenau, 13 (1967), pp. 419–421.
- [3] ———, *On a conjecture of Erdős, Hajnal and Moon*, Amer. Math. Monthly, 74 (1967), pp. 178–179.
- [4] S.-Y. CHOI AND P. GUAN, *Minimum critical squarefree subgraph of a hypercube*, in Proceedings of the Thirty-Ninth Southeastern International Conference on Combinatorics, Graph Theory and Computing, vol. 189, 2008, pp. 57–64.
- [5] P. ERDŐS, A. HAJNAL, AND J. W. MOON, *A problem in graph theory*, Amer. Math. Monthly, 71 (1964), pp. 1107–1110.
- [6] J. R. FAUDREE, R. J. FAUDREE, AND J. R. SCHMITT, *A survey of minimum saturated graphs.*, Electron. J. Comb., DS19 (2011), p. 36.
- [7] M. FERRARA, M. S. JACOBSON, F. PFENDER, AND P. S. WENGER, *Graph saturation in multipartite graphs*, J. Comb., 7 (2016), pp. 1–19.
- [8] A. HAJNAL, *A theorem on k -saturated graphs*, Canad. J. Math., 17 (1965), pp. 720–724.
- [9] J. R. JOHNSON AND T. PINTO, *Saturated subgraphs of the hypercube*, Combin. Probab. Comput., 26 (2017), pp. 52–67.
- [10] L. KÁSZONYI AND Z. TUZA, *Saturated graphs with minimal number of edges*, J. Graph Theory, 10 (1986), pp. 203–210.

- [11] D. KORÁNDI AND B. SUDAKOV, *Saturation in random graphs*, ArXiv e-prints, (2015).
- [12] N. MORRISON, J. A. NOEL, AND A. SCOTT, *Saturation in the hypercube and bootstrap percolation*, *Combin. Probab. Comput.*, 26 (2017), pp. 78–98.
- [13] B. ROBERTS, *Partite saturation problems*, *Journal of Graph Theory*, 85 (2017), pp. 429–445.
- [14] E. SULLIVAN AND P. S. WENGER, *Saturation numbers in tripartite graphs*, *Journal of Graph Theory*, 84 (2017), pp. 428–442.
- [15] W. WESSEL, *über eine Klasse paarer Graphen. I. Beweis einer Vermutung von Erdős, Hajnal und Moon*, *Wiss. Z. Techn. Hochsch. Ilmenau*, 12 (1966), pp. 253–256.
- [16] ———, *über eine Klasse paarer Graphen. II. Bestimmung der Minimalgraphen*, *Wiss. Z. Techn. Hochsch. Ilmenau*, 13 (1967), pp. 423–426.
- [17] A. A. ZYKOV, *On some properties of linear complexes*, *Mat. Sbornik N.S.*, 24(66) (1949), pp. 163–188.