

A note on the intersection property for flat boxes and boxicity in \mathbb{R}^d

Hector Baños*

University of Alaska, Fairbanks

Deborah Oliveros†

Instituto de Matemáticas

Universidad Nacional Autónoma de México, México

March 1, 2022

Celebrating the 70th birthday of T. Bisztriczky, G. Fejes Toth, and A. Makai

Abstract

By extending the definition of boxicity, we extend a Helly-type result given by Danzer and Grünbaum on 2-piercings of families of boxes in d -dimensional Euclidean space by lowering the dimension of the boxes in the ambient space.

Key words. Boxicity, Helly-type results, intersection graph of boxes, p -boxicity.

1 Introduction

A p -box in \mathbb{R}^d is a rectangular p dimensional parallelotope whose edges are parallel to the coordinate axes in \mathbb{R}^d , where $d \geq p$. A family of boxes is called n -pierceable if there exists a set of n points such that each box contains at least one of these points.

For positive integers d and n , what is the smallest number $h = h(d, n)$ such that the following property holds:

“Every finite family \mathcal{F} of d -boxes in \mathbb{R}^d is n -pierceable if and only if, every subfamily of cardinality h is n -pierceable.”

This problem was originally studied by Danzer and Grünbaum in [1], showing in particular that the following theorem holds for piercing number 2:

Theorem 1. [1] *For d odd, $h(d, 2) = 3d$ and for d even $h(d, 2) = 3d - 1$.*

Results of the type “if every subset of cardinality μ of \mathcal{F} is n -pierceable, then the entire family \mathcal{F} is n -pierceable” are called Helly–Gallai type theorems. Clearly Helly–Gallai type theorems are a natural generalization of Helly’s classical theorem for $n = 1$

*hdbanos cervantes@alaska.edu

†dolivero@matem.unam.mx

when \mathcal{F} is a family of convex sets in \mathbb{R}^d and $\mu = d + 1$. Results of this type have been widely studied in different settings (see, for instance, surveys such as [1], [5]). However, in the same paper [1] of Danzer and Grünbaum they show that such theorems do not exist in general, even for the case of families of d -boxes in \mathbb{R}^d .

In [10] the authors give an alternative proof of Theorem 1 using intersection graphs of families of boxes, the fact that for the case $n = 1$, $h(d, 1) = 2$ in any dimension, analyzing the structure of the complement of odd cycles and the chromatic number $\chi(G)$ of a graph G ; see Proposition 13 in this paper.

In 1969 F.S. Roberts [7] extended the definition of interval graph to higher dimensions by considering the intersection graph of a family of d -boxes in d -dimensional Euclidean space \mathbb{R}^d by defining the boxicity of a graph G , denoted $Box(G)$, as the smallest positive integer d for which a graph G is the intersection graph of a family of d -boxes in \mathbb{R}^d . For further reference in this topic and part of its state of art, see [12], [13], [14], and [15].

In his work, Roberts gave a characterization of the boxicity of a noncomplete graph G in terms of interval graphs by showing the following theorem:

Theorem 2. [7] *The boxicity of a noncomplete graph G is the minimum positive integer k such that there exists interval graphs F_1, F_2, \dots, F_k such that $G = F_1 \cap F_2 \cap \dots \cap F_k$.*

In this paper, we introduce the definition of p -boxicity, $Box_p(G)$, as the minimum dimension d such that a graph G is realizable as the intersection graph of p -boxes in \mathbb{R}^d (boxes of dimension p in \mathbb{R}^d), and give a generalization of Roberts' result (Theorem 2) in terms of p -boxes. We believe this definition is interesting in its own right and may yield to further research activity. Furthermore, we extend Danzer's and Grünbaum's theorem for piercing number 2 to the family of flat (that is, not necessarily full dimensional) boxes using this new definition. In particular, we prove the following theorem:

Theorem 3. *Given any family \mathcal{F} of m -boxes in \mathbb{R}^d , $m \leq d$, \mathcal{F} is 2-pierceable if and only if every subfamily of cardinality $h = h(d, m, 2)$ is 2-pierceable provided*

$$h(d, m, 2) = \begin{cases} 5 & \text{for } m = 1 \\ 7 & \text{for } m = 2 \\ 3m & \text{for } m \neq 1 \text{ odd} \\ 3m - 1 & \text{for } m \neq 2 \text{ even} \end{cases}$$

2 Boxicity and p -boxicity

Given a finite family $\mathcal{F} = \{X_1, X_2, \dots, X_k\}$ of p -boxes (boxes of dimension p) in \mathbb{R}^d , a graph $G = (V(G), E(G))$ is the intersection graph of \mathcal{F} if $V(G) := \{x_1, x_2, \dots, x_k\}$ and $(x_i, x_j) \in E(G)$ if and only if $X_i \cap X_j \neq \emptyset$.

Recall that given a graph G , its p -boxicity $Box_p(G)$ is the minimum dimension d such that G is the intersection graph of family of p -boxes in \mathbb{R}^d . Observe that $p \leq d$ and if $p = d$ then $Box_p(G) = Box(G)$. Note that for any graph G and $p \in \mathbb{N}$, if $p \geq Box(G)$




Figure 1: Graph G and its realization as 2-boxes in \mathbb{R}^2 .

then $\text{Box}_p(G) = p$. In general $\text{Box}(G) \leq \text{Box}_p(G)$. We will say that a family \mathcal{F} of p -boxes is a *realization* of G if G is the intersection graph of \mathcal{F} .

To illustrate some of these statements, observe that for a cycle C_s , $s \geq 4$, we have $\text{Box}_1(C_s) = 2$ and $\text{Box}_2(C_s) = \text{Box}(C_s)$ (see figure 6). If, however, G is the graph shown on the left of Figure 1, then $\text{Box}_1(G) = \infty$ but $\text{Box}(G) = 2$. Figure 2 shows a graph G where $\text{Box}(G) = 2$ and $\text{Box}_1(G) = 3$; in this case the realization of G as 1-boxes in \mathbb{R}^3 can be thought of as the edges of a cube.

For us, all graphs $G = (V(G), E(G))$ are finite and simple, i.e. with no loops or multiple edges. Recall that a subgraph $H \subset G$ is an induced subgraph of G if $V(H) \subset V(G)$ and if $x_i, x_j \in V(H)$ and $(x_i, x_j) \in E(G)$ then $(x_i, x_j) \in E(H)$. The neighborhood of a vertex $v \in V(G)$ is the set of vertices adjacent to v , $N(v) := \{w \in V(G) \mid (v, w) \in E(G)\}$. We will define the closure of the neighborhood of v as the set of all possible edges in $N_G(v)$, $[N_G(v)] := \{(u, w) \mid u, w \in N(v)\}$. When it is clear from context, the graph induced by the neighborhood of a vertex will be also referred as the neighborhood of a vertex. As usual K_t will denote the complete graph with t vertices.

One consequence of the following definition and lemma is that we can determine when a graph G that satisfies $\text{Box}(G) = d$ also satisfies $\text{Box}_p(G) = d$.

Definition 4. We will say that a graph G satisfies the *p-slim box property* in \mathbb{R}^d if the following two conditions hold:

- i) G is an intersection graph of d -boxes, i.e. $G = \cap_{i \in I} F_i$ where every F_i is an interval graph and $I := \{1, 2, \dots, d\}$.
- ii) For all $v \in V(G)$, there is a subset $J_v \subset I := \{1, 2, \dots, d\}$ of $d - p$ indices such that the neighborhood of v is a complete subgraph, $[N_{F_i}(v)] \subset E(F_i)$, $i \in J_v$.

Lemma 5. A graph G is realizable as p -boxes in \mathbb{R}^d ($p \leq d$) if and only if G has the *p-slim property* in \mathbb{R}^d .




Figure 2: Top: a graph G where $\text{Box}(G) = 2$ and $\text{Box}_1(G) = 3$. Bottom left: the realization of G as 1-boxes in \mathbb{R}^3 ; each 1-box is an edge of the cube. Bottom right: the realization of G as 2-boxes in \mathbb{R}^2 .

Proof.

Suppose that G is realizable as p -boxes in \mathbb{R}^d . Let F_i be the intersection graph of the projection of the realization of G on the i th coordinate axis, and observe that $G = F_1 \cap F_2 \cap \dots \cap F_d$. Let V be a p -box in the realization of G and let $v \in V(G)$ be its representation in the graph. Since V is a p -box there are $d - p$ coordinate axes where the projection of V is a point; hence, in any of these $d - p$ axes the corresponding F_j satisfies $[N_{F_j}(v)] \subset E(F_j)$. Therefore G has the p -slim property in \mathbb{R}^d .

Suppose now that G has the p -slim property in \mathbb{R}^d . Then there exist interval graphs F_1, \dots, F_k such that $G = \bigcap_{i=1}^d F_i$, and for any $v \in V(G)$ there exists a subset $J_v \subset I$ of $d - p$ indices such that $[N_{F_i}(v)] \subset E(F_i)$, $i \in J_v$. Let I_j be the realization of F_j as intervals. Observe that for any $i \in J_v$ we can reduce v in I_i to a point since $[N_{F_i}(v)] \subset E(F_i)$. We observe that the intersection of the projection of all I_j as the j -th axis is a family of p -boxes in \mathbb{R}^d with intersection graph G . ■

3 Flat boxes and Piercing numbers

For the remainder of this paper C_s will denote the cycle of length s , with $V(C_s) = \{v_1, v_2, \dots, v_s\}$ and $E(C_s) = \{(v_1, v_2), (v_2, v_3), \dots, (v_{s-1}, v_s), (v_s, v_1)\}$. We will denote by $P_{\{v_1, v_2, \dots, v_l\}}$ the path of length $l - 1$ with vertices $V(P_{\{v_1, v_2, \dots, v_l\}}) = \{v_1, v_2, \dots, v_l\}$ and edges $E(P_{\{v_1, v_2, \dots, v_l\}}) = \{(v_1, v_2), \dots, (v_{l-1}, v_l)\}$. We denote the complement of C_s by C_s^c . One graph of particular interest to us is the path $P_{v_k}(C_s) := P_{\{v_{(k+2 \bmod s)}, v_{(k+3 \bmod s)}, \dots, v_{(k+s-2 \bmod s)}\}}$ as a subgraph of C_s with respect to some vertex v_k of C_s (see Figure 3).

We will say that a graph G is m -forbidden in \mathbb{R}^d if G^c is not realizable as m -boxes in \mathbb{R}^d .




Figure 3: The path $P_{v_k}(C_s)$ colored in green.

The following observation is well known.

Observation 6. *Since interval graphs do not contain C_4 as an induced subgraph, then $2K_2$ (two disjoint edges) and a path of length greater or equal to 4, are 1-forbidden in \mathbb{R}^1 .*

We will prove some lemmas that will help us to prove Theorem 3. The following observation will be really helpful to prove such lemmas.

Observation 7. *Suppose that $G = C_s^c$ satisfies the p -slim box property in \mathbb{R}^d for some $p \leq d$ and some s . By Definition 4 we know that $G = \bigcap_{i \in I} F_i$ where F_i is an interval graph, $I = \{1, \dots, d\}$, and for every $v_k \in V(G)$ there exists $J_{v_k} \subset I$, with $|J_{v_k}| = d - p$, such that $[N_{v_k}] \subset F_j$ for all $j \in J_{v_k}$. Then $P_{v_k}(C_s) \subset \bigcap_{j \in J_{v_k}} F_j$.*

The following lemmas will allow us to prove the main theorem.

Lemma 8. *The cycle C_s with $s \geq 7$ is 1-forbidden in \mathbb{R}^d for any $d > 1$.*

Proof. We need to show that $G := C_s^c$, $s \geq 7$, is not realizable as 1-boxes in \mathbb{R}^d . Suppose that G is realizable as 1-boxes in \mathbb{R}^d . Then by Lemma 5, G satisfies the 1-slim box property. Thus $G = \cap_{i \in I} F_i$, $I = \{1, \dots, d\}$, where F_i are interval graphs and for any vertex $v \in V(G)$ there is a subset $J_v \subset I$ with $|J_v| = d - 1$, such that $[N_{F_i}(v)] \subset E(F_i)$, $i \in J_v$.

Without loss of generality assume that $J_{v_1} := \{1, 2, \dots, d - 1\}$. By Observation 7, $P_{v_1}(C_s) \subset \cap_{j=1}^{d-1} F_j$ and thus $(v_3, v_4), (v_4, v_5), \dots, (v_{s-2}, v_{s-1}) \notin E(F_d)$ otherwise $G \neq \cap_{i \in I} F_i$ (see Figure 4).

If $s > 7$, then there would be at least four edges missing in F_d . This contradicts Observation 6. If $s = 7$, by Observation 7 on v_6 the path $P_{v_6}(C_s)$ should belong to $d - 1$ of the F_i interval graphs, but since $(v_3, v_4) \notin E(F_d)$, $J_{v_1} = J_{v_6} = \{1, \dots, d - 1\}$. Then $P_{v_6}(C_s) \in \cap_{i=1}^{d-1} F_i$. This implies $(v_2, v_3) \notin E(F_d)$, otherwise (v_2, v_3) belongs to every F_i contradicting $G = C_7^c$. Then the edges $(v_2, v_3), (v_3, v_4), (v_4, v_5), (v_5, v_6)$ are not in $E(F_d)$, contradicting Observation 6. ■




Figure 4: The d interval graphs whose intersection is G . Until this point of the proof we do not know anything about the path $P_{s-1, s, 1, 2, 3}$ in each of the graphs, so a dashed edge means that such an edge may or may not be in the projection. Observe that the first $d - 1$ graphs contain $[N(v_1)]$ and therefore $P_{v_1}(C_s)$, but the last graph does not contain $P_{v_1}(C_s)$, otherwise the intersection would not be G .

Lemma 9. *The cycle C_s , $s \geq 9$, is 2-forbidden in \mathbb{R}^d for any $d > 2$.*

Proof. Again we need to show that $G := C_s^c$ is not realizable as 2-boxes in \mathbb{R}^d .

Suppose that G is realizable as 2-boxes in \mathbb{R}^d . Then by Lemma 5, G satisfies the 2-slim box property in \mathbb{R}^d . Thus $G = \cap_{i \in I} F_i$, $I = \{1, \dots, d\}$, where F_i are interval graphs and for any vertex $v \in V(G)$ there is a subset $J_v \subset I$, $|J_v| = d - 2$, such that $[N_{F_i}(v)] \subset E(F_i)$, $i \in J_v$. Again without loss of generality suppose $J_{v_1} := \{1, 2, \dots, d - 2\}$. By Observation 7, $P_{v_1}(C_s) \subset \cap_{j=1}^{d-2} F_j$, and thus $(v_3, v_4), (v_4, v_5), \dots, (v_{s-2}, v_{s-1}) \notin E(F_d) \cap E(F_{d-1})$.

Without loss of generality suppose $(v_{s-2}, v_{s-1}) \notin E(F_{d-1})$. Since $s \geq 9$, by Observation 6 $(v_3, v_4), (v_4, v_5) \notin E(F_d)$ and therefore $(v_{s-2}, v_{s-3}) \notin E(F_{d-1})$.

Note that if $(v_{s-3}, v_{s-4}) \notin E(F_{d-1})$, $J_{v_{s-1}} = \{1, 2, \dots, d-2\}$. This is because $(v_3, v_4) \notin E(F_d)$ and $(v_{s-3}, v_{s-4}), (v_3, v_4) \in [N_{v_{s-1}}]$; by the p -slim property there exists $J_{v_{s-1}} \subset I$ where $[N_{v_{s-1}}] \subset F_j$ for all $j \in J_{v_{s-1}}$.

Therefore by Observation 7 any edge of the path $P_{v_{s-1}}(C_s)$ is not in $E(F_d) \cap E(F_{d-1})$ (in particular this includes the edge (v_1, v_2)). This is a contradiction, since by Observation 6 the edge (v_1, v_2) must be in $E(F_d) \cap E(F_{d-1})$ otherwise we obtain either an empty path of size at least 4 or a disjoint path in any of $E(F_d), E(F_{d-1})$. We proceed with an analogous argument if $(v_{s-3}, v_{s-4}) \notin E(F_d)$. ■

Let $p, d \in \mathbb{N}$ with $2 < p < d$. Suppose that $G := C_s^c$ is realizable as p -boxes in \mathbb{R}^d for some $s \geq 3p + 1$. Then by Lemma 5, G satisfies the p -slim box property in \mathbb{R}^d . Thus there exists d interval graphs F_1, F_2, \dots, F_d such that $G = \bigcap_{i \in I} F_i$, $I = \{1, \dots, d\}$ and for any vertex $v \in V(G)$, there is a subset $J_v \subset I$, $|J_v| = d - p$, such that $[N_{F_i}(v)] \subset E(F_i)$, $i \in J_v$. Without loss of generality suppose that $J_{v_1} := \{1, 2, \dots, d - p\}$. Denote the rest of the vertices by $J = \{d - p + 1, \dots, d\}$ and observe that $|J| = p$. Since $d > p$, J_{v_1} is not empty.

By Observation 7 the path of length $s - 4$ is contained in F_i with $i \in J_{v_1}$, i.e. $P_{v_1}(C_s) \subset \bigcap_{j=1}^{d-p} F_j$. Since $G = \bigcap_{i \in I} F_i$, then for every edge e of the path $P_{v_1}(C_s)$ there exists at least one $i \in J$ such that $e \notin E(F_i)$, in this case we say that e is a *missing edge* of F_i . Similarly a *missing path*, is a path of missing edges in F_i .

We also say that F_j , with $j \in J$, satisfies the *missing property* if there is a missing edge of F_j , $e \in E(P_{v_1}(C_s)) = E(P_{v_3, \dots, v_{s-1}})$, such that $e \in E(F_i)$ for all $i \in J \setminus \{j\}$. For example, in Figure 5 F_3 has the missing property since $e := (v_9, v_{10}) \notin E(F_3)$ but $e \in E(F_4)$ and $e \in E(F_5)$.

If there are $r, t \in J$ such that either the path P_{v_3, v_4, v_5, v_6} is a missing path of F_r and the path $P_{v_{s-3}, v_{s-2}, v_{s-1}}$ is a missing path of F_t or the path P_{v_3, v_4, v_5} is a missing path of F_r and the path $P_{v_{s-4}, v_{s-3}, v_{s-2}, v_{s-1}}$ is a missing path of F_t , we say that J satisfies the *extreme condition*. If there exists u such that $6 \leq u \leq s - 4$ and $r, t \in J$ such that $P_{v_{u-3}, v_{u-2}, \dots, v_{u+2}, v_{u+3}}$ is missing in $F_t \cup F_r$, we say that J satisfies the *contiguous condition*. The following two technical lemmas imply that if J satisfies the missing property for all $j \in J$, then neither the contiguous nor the extreme condition holds.

Lemma 10. *Suppose that for all $j \in J$, F_j satisfies the missing property (where J and F_j are defined as above). Then J does not satisfy the contiguous condition.*

Proof. Suppose the opposite. Since the edges $(v_u, v_{u-2}), (v_u, v_{u-3})$ and (v_{u-2}, v_{u-3}) are missing in, say, F_r , then $r \notin J_{v_u}$. Analogously $(v_u, v_{u+2}), (v_u, v_{u+3}) \in E(G)$ and (v_{u+2}, v_{u+3}) is missing in $E(F_t)$, so $t \notin J_{v_u}$. For any other $j \in J \setminus \{r, t\}$, the missing property implies there exists a missing edge (x, y) of F_j such that $e \in F_r$ and $e \in F_t$. This implies $x, y \notin \{v_{u-2}, v_{u-1}, v_u, v_{u+2}, v_{u+3}\}$. Hence $(v_u, x), (v_u, y) \in E(G)$ and thus $j \notin J_{v_u}$. Therefore $J_{v_u} = J_{v_1} = \{1, \dots, d - p\}$. From $6 \leq u \leq s - 4$ and Observation 7,

we note that P_{v_u} is in F_k for all $k \in J_{v_1}$. That is, F_k is a complete graph for all $k \in J_{v_1}$. Thus for any edge e of the cycle C_s , $e \notin \cap_{j \in J} F_j$.

Given $s \geq 3p + 1$ and $|J| = p$, there is some F_j with $j \in J$ missing more than 3 edges, which contradicts Observation 6. ■

Lemma 11. *Suppose that for all $j \in J$, F_j satisfies the missing property. Then J does not satisfy the extreme condition.*

Proof. Suppose that there are $r, t \in J$ such that P_{v_3, v_4, v_5, v_6} is a missing path of F_r and $P_{v_{s-3}, v_{s-2}, v_{s-1}}$ is a missing path of F_t .

Since $(v_3, v_5), (v_3, v_6) \in E(G)$ but (v_6, v_5) is a missing edge of F_r , we have $r \notin J_{v_3}$. For any other $j \in J$, by the missing property there exists a missing edge (x, y) of F_j such that $e \in F_r$. In particular we observe that $(v_3, x), (v_3, y) \in E(G)$, and thus $j \notin J_{v_3}$. Therefore $J_{v_3} = J_{v_1} = \{1, 2, \dots, d - p\}$ and (v_{s-1}, v_s) is a missing edge of $\cap_{j \in J} F_j$.

By Observation 6 and the missing property, the edge (v_{s-1}, v_s) is only a missing edge of F_t . Therefore $P_{v_{s-3}, v_{s-2}, v_{s-1}, v_s}$ is a missing path of F_t . Since $(v_s, v_{s-3}), (v_s, v_{s-4}) \in E(G)$, but (v_{s-3}, v_{s-4}) is a missing edge of F_t , it follows $t \notin J_{v_s}$. For any other $j \in J$, by the missing property there exists a missing edge (x, y) of F_j such that $e \in F_t$. In particular $(v_s, x), (v_s, y) \in E(G)$, so $j \notin J_{v_s}$. Therefore $J_{v_s} = J_{v_1} = \{1, \dots, d - p\}$. By Observation 7 $P_{v_s}(C_s)$ is in F_k for all $k \in J_{v_1}$, which implies that the edge (v_2, v_3) is a missing edge of $\cap_{j \in J} F_j$. By Observation 6 and the fact that for any $j \in J$, F_j satisfies the missing property, there cannot be some F_j , $j \in J$, with such a missing edge. This is a contradiction.

Note that an analogous argument can be used if the path P_{v_3, v_4, v_5} is a missing path of F_r and the path $P_{v_{s-4}, v_{s-3}, v_{s-2}, v_{s-1}}$ is a missing path of F_t , for some $r, t \in J$. ■

We apply Lemmas 10 and 11 to show the following useful lemma.

Lemma 12. *The cycle C_s is p -forbidden in \mathbb{R}^d for any $d > p > 2$ when the following two conditions hold*

- i) $s \geq 3p + 1$ for p even,*
- ii) $s \geq 3p + 2$ for p odd.*

Proof.

We will show that $G := C_s^c$ for the corresponding values of s is not realizable as p -boxes in \mathbb{R}^d . Suppose that G is realizable as p -boxes in \mathbb{R}^d . By Lemma 5, G has the p -slim property. Then there exists d interval graphs F_1, F_2, \dots, F_d such that $G = \cap_{i \in I} F_i$, $I = \{1, \dots, d\}$ and, as before, we assume that for v_1 , $J_{v_1} := \{1, 2, \dots, d - p\}$ and $J = \{d - p + 1, \dots, d\}$ are the rest of indices (observe that $|J| = p$). Since $d > p$, J_{v_1} is not empty. So $P_{v_1}(C_s)$ has length $s - 4$, and satisfies $P_{v_1}(C_s) \subset \cap_{i \in J_{v_1}} F_i$ and each edge of $P_{v_1}(C_s)$ is missing in $\cap_{j \in J} F_j$.

Observe first that if $s - 4 \geq 3p + 1$, at least one of the F_j , $j \in J$, will contain a path with 4 or more missing edges yielding a contradiction of Observation 6. Thus we may assume

$$3p + 1 \leq s < 3p + 5 \text{ for any } d > p > 2.$$

Furthermore, by Observation 6 each F_i , $i \in J$, can only have a missing subpath of P_{v_1} of length 3, 2 or 1.

Since $s < 3p + 5$ and for any edge $e \in E(P_{v_1})$ there exists $j \in J$ such that e is missing in F_j , it is possible to construct a partition (not necessarily unique) of $P_{v_1}(C_s)$ into disjoint connected subpaths of length 1, 2 and 3, with the following two properties:

1. If Q is an element of such a partition then Q is a missing path of F_j for some $j \in J$. In this case we may say that F_j represents Q .
2. If Q and P are different elements of the partition then $i \neq j$ for its corresponding representative F_i and F_j .

For example, in Figure 5 we see a particular case when $s = 11$, $p = 2$ and $d = 5$. Here $J_{v_1} = \{1, 2\}$ and $J = \{3, 4, 5\}$. The dashed edges may or may not be on the graph. Observe that $P_{v_1}(C_{11}) = P_{v_3, \dots, v_{10}}$ is in F_1, F_2 . Consider the following partition K of $P_{v_1}(C_{11})$: $Q_3 := P_{v_7, v_8, v_9, v_{10}}$ represented by F_3 , $Q_4 := P_{v_6, v_7}$ represented by F_4 and $Q_5 := P_{v_3, v_4, v_5, v_6}$ represented by F_5 . Clearly we may have also chosen the partition K' as $Q_3 := P_{v_8, v_9, v_{10}}$ represented by F_3 , $Q_4 := P_{v_6, v_7, v_8}$ represented by F_4 and $Q_5 := P_{v_3, v_4, v_5, v_6}$ represented by F_5 .

For a given partition, let $J_i \subset J$ for $i = 1, 2, 3$, be the set of indices $j \in J_i$ such that F_j represents an element of the partition of size i . We observe that for any partition, $\bigcap_{i=1}^3 J_i = \emptyset$ and $\bigcup_{i=1}^3 J_i \subseteq J$. Furthermore,

$$\begin{aligned} s - 4 &= |J_1| + 2|J_2| + 3|J_3| \\ p &\geq |J_1| + |J_2| + |J_3| \end{aligned} \tag{1}$$

In the example in Figure 5, for the partition K , we obtain $J_1 = \{4\}$, $J_2 = \emptyset$ and $J_3 = \{3, 5\}$. For the partition K' we obtain $J_1 = \emptyset$, $J_2 = \{3, 4\}$ and $J_3 = \{5\}$.

In general, observe that if for all possible partitions we have

$$|J_1| + |J_2| + |J_3| = p, \tag{2}$$

then the missing property holds for all $j \in J$. To establish this, assume that there is $j \in J$ such that for any edge e missing in F_j , there exists $i_e \in J$ such that e is missing in F_{i_e} . We can construct a partition with $j \notin J_1 \cup J_2 \cup J_3$, and thus this partition of $P_{v_1}(C_s)$ satisfies $|J_1| + |J_2| + |J_3| = p - 1$ and $s - 4 = |J_1| + 2|J_2| + 3|J_3|$. This is a contradiction.




Figure 5: In this figure we see a particular example of the partition K when $s = 11, p = 2, d = 5$ and, $J = \{3, 4, 5\}$. The red paths represent the partition K formed by Q_3, Q_4 and Q_5 .

Next we will show that in most of the cases for p and s , Equation (2) holds and therefore the missing property holds for all $j \in J$. Moreover, with appropriate values of s , this yields a contradiction of Lemmas 10 and 11.

Observe that if $s = 3p + 4$, the only solution for System (1) is $|J_3| = p, |J_2| = 0$ and $|J_1| = 0$. This implies the missing property holds for all $j \in J$. Note also that since all elements in the partition have length 3, and since each element of the partition is uniquely represented by some F_j with $j \in J$, J satisfies the contiguous condition and thus contradicts Lemma 10. Therefore we may assume that $s < 3p + 4$.

For $s = 3p + 3$, there are at least $s - 4 = 3p - 1$ missing edges in $\cap_{j \in J} F_j$ (one per edge in P_{v_1}). By solving System (1) we find that $|J_3| = p - 1, |J_2| = 1$ and $|J_1| = 0$, which implies that the missing property holds.

If $p > 3$, then $p - 1 > 2$. By the pigeon hole principle and the fact that each element of the partition is uniquely represented by some F_j with $j \in J$, J satisfies the contiguous condition and contradicts Lemma 10. If $p = 3$, $|J| = 3, |J_3| = 2$ and $|J_2| = 1$. Thus J satisfies either the contiguous condition or the extreme condition, contradicting either Lemma 10 or Lemma 11 (again, this is also using the fact that each element of the partition is uniquely represented by an F_j with $j \in J$). Therefore we may assume $s \leq 3p + 2$.

For $s = 3p + 2$, there are at least $s - 4 = 3p - 2$ missing edges in $\cap_{j \in J} F_j$. Solving System (1) we find that either $|J_3| = p - 1, |J_2| = 0$ and $|J_1| = 1$ or $|J_3| = p - 2, |J_2| = 2$ and $|J_1| = 0$, which implies that the missing property holds for all $j \in J$. If $p > 5$, then $p - 2 > 3$. As before, by the pigeon hole principle J satisfies the contiguous condition and contradicts Lemma 10. Note that if $p = 5$, either $|J_3| = 4$ and $|J_1| = 1$, or $|J_3| = 3$ and $|J_2| = 2$. In either case, the pigeon hole principle implies J satisfies either the contiguous condition or the extreme condition and thereby contradicts either Lemma 10 or Lemma 11.

If $p = 4$, solving System (1) gives $|J_3| = 3$, $|J_2| = 0$, and $|J_1| = 1$ or $|J_3| = 2$, $|J_2| = 2$, and $|J_1| = 0$. In the first case, by the pigeon hole principle J satisfies the contiguous condition and contradicts Lemma 10. In the second case, again by the pigeon hole principle J satisfies the extreme condition and contradicts Lemma 11.

If $p = 3$, solving System (1) gives either $|J_3| = 2$, $|J_2| = 0$, and $|J_1| = 1$ or $|J_3| = 1$, $|J_2| = 2$, and $|J_1| = 0$. In the first case, by the pigeon hole principle J satisfies the contiguous condition and contradicts Lemma 10. For the second case we observe that the only way J does not satisfy the extreme condition (which would contradict Lemma 11) is if there is an F_j missing no more than P_{v_3, v_4, v_5} from $P_{v_1}(C_{11})$, F_i missing no more than P_{v_5, v_6, v_7, v_8} from $P_{v_1}(C_{12})$, and F_k missing no more than $P_{v_8, v_9, v_{10}}$ from $P_{v_1}(C_{12})$ for $i, j, k \in J$. Without loss of generality suppose that $i = d - 2$, $j = d - 1$ and $k = d$. Since we know that the path $P_{v_{11}}(C_{11})$ must exist in at least $d - 3$ projections, we have $J_{v_{11}} = \{1, 2, \dots, d - 3\}$. Therefore $(v_2, v_3) \in E(F_i)$ for all $i \in \{1, 2, \dots, d - 3\}$. This implies (v_2, v_3) is a missing edge of $F_{d-2} \cap F_{d-1} \cap F_d$. By Observation 6 (v_2, v_3) can be only missing in F_{d-2} . Analogously we observe that $J_{v_2} = \{1, 2, \dots, d - 3\}$ and thus (v_{10}, v_{11}) is a missing edge of F_d . This implies $J_{v_5} = \{1, 2, \dots, d - 3\}$, since the path $P_{v_5}(C_{11})$ is in $d - 3$ of the F_i 's. Therefore F_1, \dots, F_{d-3} are complete graphs K_{11} . But then (v_1, v_2) is a missing edge of $F_{d-2} \cap F_{d-1} \cap F_d$, which is a contradiction of Observation 6 since there cannot be more than three edges missing in any of F_{d-2}, F_{d-1} and F_d .

We have shown that $s \geq 3p + 2$ is p -forbidden for odd p . Thus we may assume that $s \leq 3p + 1$.

Let $s = 3p + 1$. Then there are $3p - 3$ edges of P_{v_1} missing in $\cap_{j \in J} F_j$. Solving System (1) gives

$$\begin{aligned} |J_3| &= p - 1, |J_2| = 0 \text{ and, } |J_1| = 0, \\ |J_3| &= p - 2, |J_2| = 1 \text{ and, } |J_1| = 1, \text{ or} \\ |J_3| &= p - 3, |J_2| = 3 \text{ and, } |J_1| = 0. \end{aligned}$$

Observe that the case where $|J_3| = p - 1$, $|J_2| = 0$, and $|J_1| = 0$ is the only case where the missing property does not hold for all $j \in J$. Without loss of generality suppose $J_3 = \{d - p + 1, \dots, d - 1\}$. Since $G = \cap_{i=1}^d F_i$, there exists $k \in \{1, \dots, d\}$ such that (v_{s-1}, v_s) is a missing edge of F_k . By Observation 6, $k \notin \{d - p + 1, \dots, d - 1\}$. Similarly there exists $r \in \{1, \dots, d\}$ such that the edge (v_2, v_3) is missing in F_r . By Observation 6, $r \notin \{k, d - p + 1, \dots, d - 1\}$. Since $|J_3| = p - 1$, $|J_2| = 0$, and $|J_1| = 0$, we have $(v_3, v_4), (v_4, v_5)$ and (v_5, v_6) are missing in some F_t with $t \in J_3$. Similarly $(v_6, v_7), (v_7, v_8)$ and (v_8, v_9) are missing in some F_q for some $q \in J_3$. Thus there is no $J_{v_6} \subset \{1, \dots, d\}$ such that $|J_{v_6}| = d - p$ and such that the neighborhood of v_6 is a complete subgraph, given that we know $|\{r, k, d - p + 1, \dots, d - 1\}| = p + 1$ and in any of F_i , $i \in \{r, k, d - p + 1, \dots, d - 1\}$, the neighborhood of v_6 is not a complete subgraph. This yields a contradiction.

Now suppose that $|J_3| = p - 2$, $|J_2| = 1$ and $|J_1| = 1$. Note that the missing property holds for every $j \in J$. Let $p > 3$. Then Lemma 10 or Lemma 11 are contradicted since

J would satisfy either the contiguous or the extreme condition unless there are $r, t \in J$ where F_r is missing in at most the edge (v_{s-1}, v_{s-2}) from $P_{v_1}(C_s)$ and F_t is missing in at most P_{v_3, v_4, v_5, v_6} from $P_{v_1}(C_s)$ (or similarly, F_r is at most missing $P_{v_{s-1}, v_{s-2}, v_{s-3}, v_{s-4}}$ from $P_{v_1}(C_s)$ and F_t with missing only (v_3, v_4) from $P_{v_1}(C_s)$). Without loss of generality suppose that F_r is missing at most (v_{s-1}, v_{s-2}) from $P_{v_1}(C_s)$ and F_t is missing at most P_{v_3, v_4, v_5, v_6} from $P_{v_1}(C_s)$. Since $p > 3$, $p-2 > 1$. This implies that there is a $g \in J$ where F_g is missing P_{v_6, v_7, v_8, v_9} or $P_{v_{s-2}, v_{s-3}, v_{s-4}, v_{s-5}}$. If F_g is missing P_{v_6, v_7, v_8, v_9} , then $J_{v_6} = \{1, 2, \dots, d-p\}$ and therefore (v_1, v_2) is missing in $\bigcap_{j \in \{d-p+1, \dots, d\}} F_j$ which contradicts Observation 6. If F_g is missing $P_{v_{s-2}, v_{s-3}, v_{s-4}, v_{s-5}}$, then $J_{v_{s-2}} = \{1, 2, \dots, d-p\}$ and therefore (v_1, v_2) is missing in $\bigcap_{j \in \{d-p+1, \dots, d\}} F_j$ which is a contradiction of Observation 6.

Now suppose that $|J_3| = p-3$, $|J_2| = 3$, and $|J_1| = 0$. Again the missing property holds. For $p \geq 6$, by the pigeon hole principle either Lemma 10 or Lemma 11 is contradicted (given that J would satisfy either the contiguous or the extreme condition). For $p = 4$ ($s = 3p + 1 = 13$) we observe that $|J_3| = 1$, $|J_2| = 3$, and $|J_1| = 0$. If there exists $j \in J$ where F_j is missing $P_{v_{12}, v_{11}, v_{10}, v_9}$ or P_{v_3, v_4, v_5, v_6} , Lemma 11 is contradicted (given that J would satisfy the extreme condition).

Therefore there are $r, t \in J$ with F_r at most missing P_{v_3, v_4, v_5} from $P_{v_1}(C_{13})$ and F_t with at most missing $P_{v_{12}, v_{11}, v_{10}}$ from $P_{v_1}(C_{13})$. Therefore $J_{v_{12}} = \{1, 2, \dots, d-4\} = J_{v_1}$. This implies that (v_{13}, v_{12}) and (v_2, v_3) are in $E(F_k)$ for all $k \in \{1, 2, \dots, d-4\}$ and thus they are missing in $\bigcap_{j \in \{d-3, d-2, d-1, d\}} F_j$. By Observation 6, (v_{13}, v_{12}) is missing in F_t and (v_2, v_3) is missing in F_r . Observe then that either there is an F_l missing the 3-path $P_{v_{10}, v_9, v_8, v_7}$ or P_{v_5, v_6, v_7, v_8} . Without loss of generality suppose that F_l is missing $P_{v_{10}, v_9, v_8, v_7}$. We observe that $J_{v_{10}} = \{1, 2, \dots, d-4\}$. Therefore (v_1, v_2) is missing in $\bigcap_{j \in \{d-3, d-2, d-1, d\}} F_j$ which contradicts Observation 6. Therefore the cycle C_s with $s \geq 3p + 1$ is p -forbidden for even p . ■

4 Piercing two for families of flat boxes

In this section we prove Theorem 3. To do so, we apply the following proposition that seems to be widely known but for which we did not find a precise reference. In any case it is easy to show and the proof is omitted.

Proposition 13. *The piercing number of a family of boxes \mathcal{F} is n if and only if $\chi(G_{\mathcal{F}}^c) = n$, where $\chi(G_{\mathcal{F}}^c)$ denotes the chromatic number of the complement of the intersection graph of \mathcal{F} .*

Observation 14. *Let G be a graph such that $\chi(G) > 2$. If for any $v \in V(G)$ we have $\chi(G \setminus \{v\}) = 2$, then $\chi(G) = 3$ and G is an odd cycle. This is easy to see: if $\chi(G \setminus \{v\}) = 2$, then $G \setminus \{v\}$ is bipartite and thus by coloring v with a third color, we obtain $\chi(G) = 3$. It is well known that odd cycles are the only 3-critical chromatic graphs, i.e. they are the only family of graphs with chromatic number 3 such that when any vertex is removed the chromatic number decreases.*




Figure 6: Realization of C_5^c as 1-boxes in \mathbb{R}^2

Observation 15. *By Observation 14 and Proposition 13, if there is a family of p -boxes \mathcal{F} in \mathbb{R}^d such that any subset of \mathcal{F} has piercing number 2, but \mathcal{F} has a greater piercing number, then the piercing number of \mathcal{F} is 3, and $G_{\mathcal{F}}^c$ is an odd cycle and vice versa.*

Now we are ready to prove Theorem 3.

Proof. (Theorem 3) We assume $V(K_m) := \{v_1, v_2, \dots, v_m\}$.

For $m = 5$ define F_1 and F_2 to be two graphs with the same vertices as K_5 and with $E(F_1) := E(K_5) \setminus \{(v_3, v_4), (v_4, v_5), (v_5, v_1)\}$ and $E(F_2) := E(K_5) \setminus \{(v_1, v_2), (v_2, v_3)\}$. Observe that $F_1 \cap F_2 = C_5^c$, and since F_1, F_2 are chordal they are interval graphs. We observe that C_5^c has the 1-slim property in \mathbb{R}^2 and therefore is realizable as 1-boxes in \mathbb{R}^2 (see the realization of this family as 1-boxes in Figure 6). So there exists a family of 1-boxes in \mathbb{R}^d , $d > 2$, such that any 4 elements in the family have piercing 2 but the whole family has piercing 3. Hence $h(d, 1, 2) \geq 5$.

By Lemma 8, the cycle C_s with $s \geq 7$ is 1-forbidden in \mathbb{R}^d for any $d > m$. Thus there cannot be a family of intervals in \mathbb{R}^d such that its intersection graph C_s^c has the 1-slim property. It follows from this and Observation 15 that $h(d, 1, 2) \leq 5$.

For $n = 7$, let F_1, F_2 and F_3 be three graphs with vertices $V(F_i) = V(K_7)$ for $i := \{1, 2, 3\}$ and edges $E(F_1) = E(K_7) \setminus \{(v_1, v_2), (v_2, v_3)\}$, $E(F_2) = E(K_7) \setminus \{(v_3, v_4), (v_4, v_5), (v_5, v_6)\}$, and $E(F_3) = E(K_7) \setminus \{(v_6, v_7), (v_7, v_1)\}$. Observe that $F_1 \cap F_2 \cap F_3 = C_7^c$ and since F_1, F_2 and F_3 are chordal, they are interval graphs and thus C_7^c satisfies the 2-slim box property in \mathbb{R}^3 . Therefore C_7^c is realizable as 1-boxes in \mathbb{R}^3 . So there exists a family of 2-boxes in \mathbb{R}^d , $d > 3$, such that any 6 elements in the family have piercing 2 but the whole family has piercing 3 (see the realization of this family as 2-boxes in Figure 7). Hence $h(d, 2, 2) \geq 7$.

By Lemma 9, the cycle C_s with $s \geq 9$ is 2-forbidden in \mathbb{R}^d for any $d > m$. Thus there cannot be a family of 2-boxes in \mathbb{R}^d such that its intersection graph is C_s^c , and along with Observation 15 this yields $h(d, 2, 2) \leq 7$.

By Theorem 1 it is clear that $h(m, m, 2) \geq 3m$ for m odd and $h(m, m, 2) \geq 3m - 1$




Figure 7: Realization of C_7^c as 1-boxes in \mathbb{R}^3 , where all the facets of the cube are elements of the family except for the gray area that is missing in each of the facets. Each facet is labeled with the corresponding capital letter of the graph at the right. The facet on the xz -axes corresponds to V_1 , the facet parallel to it to the right V_2 , the top V_4 , front V_6 , and back V_7 . There are two elements on the bottom (xy -axes), V_3 drawn in the cube and V_5 slightly below for clarification.

for even m . By Lemma 12, the cycle C_s with $s \geq 3m + 2$ is m -forbidden in \mathbb{R}^d for any m odd, $d > m > 2$, and the cycle C_s with $s \geq 3m + 1$ is m -forbidden in \mathbb{R}^d for any m even, $d > m > 2$. Thus there cannot be a family of p -boxes in \mathbb{R}^d such that its intersection graph is C_s^c . Applying Observation 15, the proof is complete. ■

Acknowledgements.

The first author wish to acknowledge partial support by a research assistantship, funded by the National Institutes of Health grant R01 GM117590. The second author appreciate the hospitality of the department of Mathematics at U.C. Davis, support from scholarship PASPA (UNAM) and CONACYT during her sabbatical visit, as well as acknowledge support by proyecto PAPIIT 104915, 106318 and CONACYT Ciencia Básica 282280.

References

- [1] L. Danzer and B. Grünbaum. *Intersection properties of boxes in \mathbb{R}^d* . *Combinatorica*, (2), 237–246, (1982).

- [2] L. Danzer, B. Grünbaum, V. Klee. *Helly's theorem and its relatives*. Convexity, American Mathematical Society, (7), 101–179, (1963).
- [3] V. Dol'nikov, *Some Problems for Mathematical Olympiad Combinatorial Geometry*. Sociedad Matemática Mexicana–Centro de Investigación en Matemáticas, A.C. (2004).
- [4] J. Eckhoff, Discrete and Computational Geometry, The Goodman-Pollack Festschrift, *Israel J. Math.* **3**, (1965), 187–198; 2), *Algorithms and Combinatorics*, Springer-Verlag, Berlin Heidelberg New York, **Vol. 25** (2003), 347–377
- [5] J. Eckhoff. *Helly, Radon, and Caratheodory type theorems*. Handbook of Convex Geometry, 389–448, (1993).
- [6] P. Erdős, A. Goodman, L. Pósa, *The representation of a graph by set intersections*, Canadian Journal of Mathematics, (18), 106–112, (1966).
- [7] F.S. Roberts, *On the boxicity and cubicity of a graph*, in Tutte, W. T., Recent Progress in Combinatorics, Academic Press, 301–310, (1969).
- [8] B.V. Subramanya Bharadwaj, H. Rao Chintan, P. Ashok., S. Govindarajan. *On Piercing (Pseudo)Lines and Boxes*, CCCG 2012, Charlottetown, P.E.I., 8–10, (2012).
- [9] W. T. Trotter, *A characterization of Roberts' inequality for boxicity*, Discrete Mathematics, (28), 303–313, (1979).
- [10] G. Woeginger, Q. Puite, R. Pendavingh, *2-piercing via graph theory*, Discrete Applied Mathematics, (156), 3510–3512, (2008).
- [11] D.B. West, *Introduction to Graph Theory*, Prentice Hall (1996).
- [12] L. Sunil Chandran, N. Sivadasan. *Boxicity and treewidth*. Journal of Combinatorial Theory. Series B, (5), 733–744, (2007).
- [13] L. Sunil Chandran, M. Francis, N. Sivadasan. *Boxicity and maximum degree*. Journal of Combinatorial Theory. Series B, (2), 443–445, (2008).
- [14] L. Esperet, G. Joret. *Boxicity of Graphs on Surfaces*. Graphs and Combinatorics, 417–427, (2013).
- [15] A. Adiga, L. Sunil Chandran, N. Sivadasan. *Lower bounds for boxicity*. Combinatorica, (6), 631–655, (2014).