

Forbidden subgraphs for graphs of bounded spectral radius, with applications to equiangular lines

Zilin Jiang^{*,†}Alexandr Polyanskii^{†,‡}

Abstract

The spectral radius of a graph is the largest eigenvalue of its adjacency matrix. Let $\mathcal{F}(\lambda)$ be the family of connected graphs of spectral radius $\leq \lambda$. We show that $\mathcal{F}(\lambda)$ can be defined by a finite set of forbidden subgraphs if and only if $\lambda < \lambda^* := \sqrt{2 + \sqrt{5}} \approx 2.058$ and $\lambda \notin \{\alpha_2, \alpha_3, \dots\}$, where $\alpha_m = \beta_m^{1/2} + \beta_m^{-1/2}$ and β_m is the largest root of $x^{m+1} = 1 + x + \dots + x^{m-1}$. The study of forbidden subgraphs characterization for $\mathcal{F}(\lambda)$ is motivated by the problem of estimating the maximum cardinality of equiangular lines in the n -dimensional Euclidean space \mathbb{R}^n — a family of lines through the origin such that the angle between any pair of them is the same. Denote by $N_\alpha(n)$ the maximum number of equiangular lines in \mathbb{R}^n with angle $\arccos \alpha$. We establish the asymptotic formula $N_\alpha(n) = c_\alpha n + O_\alpha(1)$ for every $\alpha \geq \frac{1}{1+2\lambda^*}$. In particular, $N_{1/3}(n) = 2n + O(1)$ and $N_{1/5}(n), N_{1/(1+2\sqrt{2})}(n) = \frac{3}{2}n + O(1)$. Besides we show that $N_\alpha(n) \leq 1.49n + O_\alpha(1)$ for every $\alpha \neq \frac{1}{3}, \frac{1}{5}, \frac{1}{1+2\sqrt{2}}$, which improves a recent result of Balla, Dr axler, Keevash and Sudakov.

1 Introduction

The *spectral radius* of a graph G , denoted by $\lambda_1(G)$, is the largest eigenvalue of its adjacency matrix. Let $\mathcal{F}(\lambda)$ be the family of connected graphs of spectral radius $\leq \lambda$. It is well known that λ_1 is monotone in the sense that $\lambda_1(G_1) \leq \lambda_1(G_2)$ if G_1 is a subgraph of G_2 , moreover $\lambda_1(G_1) < \lambda_1(G_2)$ if G_1 is a proper subgraph of a connected graph G_2 . This implies that $\mathcal{F}(\lambda)$ is closed under taking connected subgraphs. It is natural to ask if $\mathcal{F}(\lambda)$ can be defined by a finite set of forbidden subgraphs. We determine the set of λ for which the answer is yes.

Theorem 1. *For $m = 1, 2, \dots$, let β_m be the largest root of $x^{m+1} = 1 + x + \dots + x^{m-1}$, and let $\alpha_m := \beta_m^{1/2} + \beta_m^{-1/2}$. For every $\lambda < \lambda^* := \sqrt{2 + \sqrt{5}}$ such that $\lambda \notin \{\alpha_2, \alpha_3, \dots\}$, there exist finitely*

^{*}Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA. Email: zilin@mit.edu. Supported in part by Israel Science Foundation (ISF) grant nos 1162/15, 936/16.

[†]Moscow Institute of Physics and Technology, Institute for Information Transmission Problems RAS, and Caucasus Mathematical Center, Adyghe State University. Email: alexander.polyanskii@yandex.ru. Supported in part by ISF grant no. 409/16, by the Russian Foundation for Basic Research through grant no. 15-01-03530 A, and by the Leading Scientific Schools of Russia through grant no. NSh-6760.2018.1.

[‡]The work was done when the authors were postdoctoral fellows at Technion – Israel Institute of Technology.

many graphs G_1, G_2, \dots, G_n such that $\mathcal{F}(\lambda)$ consists exactly of the connected graphs which do not contain any of G_1, G_2, \dots, G_n as a subgraph. However, the same conclusion does not hold for any $\lambda \in \{\alpha_2, \alpha_3, \dots\} \cup [\lambda^*, \infty)$.

The mysterious number λ^* has appeared before in the study of $\mathcal{F}(\lambda)$. After Smith [Smi70] precisely determined the graphs in $\mathcal{F}(2)$, Cvetković, Doob and Gutman [CDG82] classified the graphs in $\mathcal{F}(\lambda^*)$, which were later completely described by Brouwer and Neumaier [BN89]. On the face of it, for $\lambda < \lambda^*$, Theorem 1 may appear to be an immediate consequence of Brouwer and Neumaier’s theorem. However, Theorem 1 is genuinely about the minimal graphs outside $\mathcal{F}(\lambda)$.

Our motivation to understand the forbidden subgraphs characterization for $\mathcal{F}(\lambda)$ comes from the problem of estimating the maximum cardinality $N(n)$ of *equiangular lines* in the n -dimensional Euclidean space \mathbb{R}^n — a family of lines through the origin such that the angle between any pair of them is the same. It is considered to be one of the founding problems of algebraic graph theory to determine $N(n)$. The “absolute bound” $N(n) \leq \binom{n+1}{2}$ was established by Gerzon (see [LS73, Theorem 3.5]). For a long time, it was an open problem to determine whether n^2 is the correct order of magnitude until a remarkable construction of de Caen [dC00] shows that $N(n) \geq \frac{2}{9}(n+1)^2$ for n of the form $n = 6 \cdot 4^k - 1$ (see [GKMS16] for a generalization and [JW15] for an alternative constructions). In these constructions, the common angle tends to $\pi/2$ as dimension grows.

The question of determining the maximum number $N_\alpha(n)$ of equiangular lines in \mathbb{R}^n with common angle $\arccos \alpha$ was first raised by Lemmens and Seidel [LS73] in 1973, who showed that $N_{1/3}(n) = 2n - 2$ for $n \geq 15$ and also conjectured that $N_{1/5}(n)$ equals $\lfloor 3(n-1)/2 \rfloor$ for n sufficiently large. This conjecture was later confirmed by Neumaier [Neu89] (see also [GKMS16]).

In the absence of exact formulas for $N_\alpha(n)$ except when $\alpha = 1/3, 1/5$, various upper bounds were established. The “relative bound” $N_\alpha(n) \leq \frac{1-\alpha^2}{1-n\alpha^2} \cdot n$ is valid in small dimensions $n < 1/\alpha^2$ (see [vLS66, Lemma 6.1] and [LS73, Theorem 3.6]), and a general bound attributed to Neumann [LS73, Theorem 3.4] states that $N_\alpha(n) \leq 2n$ unless $1/\alpha$ is an odd integer. For many years, a linear upper bound for $N_\alpha(n)$ was not known when $1/\alpha$ is an odd integer bigger than 5. An important progress was recently made by Bukh [Buk16], who proved that $N_\alpha(n) \leq c_\alpha n$ for some $c_\alpha = 2^{O(1/\alpha^2)}$. Subsequently, Balla, Dräxler, Keevash and Sudakov [BDKS18] drastically improved the upper bound to $N_\alpha(n) \leq 1.93n$ for sufficiently large n relative to α whenever $\alpha \neq \frac{1}{3}$. A universal upper bound $N_\alpha(n) \leq (2/(3\alpha^2) + 4/7)n + 2$ for all $n \in \mathbb{N}$ when $1/\alpha$ is an odd integer was later found by Glazyrin and Yu [GY18].


In this paper, we combine Theorem 1 and the framework developed in [BDKS18] to establish a result of the form $N_\alpha(n) = c_\alpha n + O(1)$ for all $\alpha \geq \frac{1}{1+2\lambda^*}$. To describe the coefficient c_α , we introduce the following notion.

Definition 1. Given $\lambda > 0$, the *spectral radius order* $k(\lambda)$ of λ is the smallest order of a graph with spectral radius λ . In case λ is not the spectral radius of any graph, we write $k(\lambda) = \infty$.

Theorem 2. Given $\alpha \in (0, 1)$ such that $\lambda := \frac{1-\alpha}{2\alpha} \leq \lambda^* := \sqrt{2 + \sqrt{5}}$, the maximum number $N_\alpha(n)$ of equiangular lines in \mathbb{R}^n with angle $\arccos \alpha$ equals $\frac{k(\lambda)}{k(\lambda)-1} \cdot n + O(1)$, where $k(\lambda)$ is the spectral radius order of λ ; in case $k(\lambda) = \infty$, $N_\alpha(n) = n + O(1)$.

Remark 1. Throughout, as all the big-O notations depend on α , we suppress the subscript in $O_\alpha(\cdot)$.

Applying the above theorem to the connected graphs with 2 or 3 vertices,



whose spectral radii are 1, $\sqrt{2}$ and 2 respectively, we obtain the known results $N_{1/3}(n) = 2n + O(1)$, $N_{1/5}(n) = \frac{3}{2}n + O(1)$ and surprisingly a new result $N_{1/(1+2\sqrt{2})}(n) = \frac{3}{2}n + O(1)$. This refutes the second part of Conjecture 6.1 in [BDKS18]. The first part of the conjecture, which was also raised by Bukh [Buk16, Conjecture 8], says the following.

Conjecture A. The maximum number $N_\alpha(n)$ of equiangular lines in \mathbb{R}^n with angle $\arccos \alpha$ equals $\frac{1+\alpha}{1-\alpha} \cdot n + O(1)$ if $1/\alpha$ is an odd natural number.

When $1/\alpha$ is an odd natural number, or equivalently $\lambda = \frac{1-\alpha}{2\alpha}$ is a natural number, the complete graph on $\lambda + 1$ vertices is the smallest graph with spectral radius λ , hence $k(\lambda) = \lambda + 1$. In this case, the linear coefficient $\frac{k(\lambda)}{k(\lambda)-1}$ in Theorem 2 matches the linear coefficient $\frac{1+\alpha}{1-\alpha}$ in Conjecture A. This phenomenon motivates the following stronger conjecture.

Conjecture B. The maximum number $N_\alpha(n)$ of equiangular lines in \mathbb{R}^n with angle $\arccos \alpha$ equals $\frac{k(\lambda)}{k(\lambda)-1} \cdot n + O(1)$, where $k(\lambda)$ is the spectral radius order of $\lambda := \frac{1-\alpha}{2\alpha}$; in case $k(\lambda) = \infty$, $N_\alpha(n) = n + O(1)$.

One of the unknown cases that are not addressed by Theorem 2 is $\alpha = 1/7$. Conjecture A asserts that $N_{1/7}(n) = \frac{4}{3}n + O(1)$. For such cases, we derive an asymptotic upper bound on $N_\alpha(n)$ (see Theorem 20), from which it follows that $N_{1/7}(n) \leq (\frac{4}{3} + \frac{1}{36} + o(1))n$, and $N_\alpha(n) \leq 1.49n + O(1)$ for $\alpha \neq 1/3, 1/5, 1/(1 + 2\sqrt{2})$.

The rest of the paper is organized as follows. In Section 2, we prove Theorem 1. In Section 3.1, we adapt the framework developed by Balla et al. In Section 3.2, we apply Theorem 1 to estimate $N_\alpha(n)$ when $\lambda = \frac{1-\alpha}{2\alpha} \in (0, \lambda^*) \setminus \{\alpha_2, \alpha_3, \dots\}$. In Section 3.3 we complete the proof of Theorem 2. In Section 3.4, we extrapolate our method to obtain an upper bound on $N_\alpha(n)$ when $\lambda > \lambda^*$. In the concluding section we discuss evidences supporting Conjecture B and a possible extension of our method.

2 Forbidden subgraphs of $\mathcal{F}(\lambda)$

Suppose there is a finite forbidden subgraphs characterization, say G_1, G_2, \dots, G_n , for $\mathcal{F}(\lambda)$. The monotonicity of spectral radius implies that no connected graph has spectral radius in the open

interval $(\lambda, \min \{\lambda_1(G_i) : i \in [n]\})$. Let Λ_1 consist of the spectral radii of all connected graphs of all orders. The contrapositive of the above observation says the following.


Proposition 3. *Let $\lim_+ \Lambda_1 := \{\lambda \in \mathbb{R} : (\lambda, \lambda + \varepsilon) \cap \Lambda_1 \neq \emptyset \text{ for all } \varepsilon > 0\}$ be the set of right-sided limit points of Λ_1 . The family $\mathcal{F}(\lambda)$ does not have a finite forbidden subgraphs characterization for all $\lambda \in \lim_+ \Lambda_1$.*

Hoffman was interested in a related set R consisting of the largest eigenvalues of all symmetric matrices of all orders with non-negative integer entries, and he proved the following theorem on the limit points of R .

Theorem 4 (Hoffman [Hof72]). *Let α_m be defined as in Theorem 1. Then $2 = \alpha_1 < \alpha_2 < \dots$ are all the limit points of R smaller than $\lim_m \alpha_m = \lambda^*$.*

In fact, Hoffman proved the above theorem by first showing [Hof72, Proposition 2.1] that $\Lambda_1 = R$. He also computed the limit of spectral radii for several families of graphs. We compile some of his computation and other relevant results in the following lemma, the proof of which is presented in Appendix A. We use the notation $\alpha_n \nearrow \alpha$ if $\alpha_1 < \alpha_2 < \dots$ and $\lim_n \alpha_n = \alpha$, and $\alpha_n \searrow \alpha$ if $\alpha_1 > \alpha_2 > \dots$ and $\lim_n \alpha_n = \alpha$.

Lemma 5. *Let α_m be defined as in Theorem 1. Denote by C_n the cycle with n vertices, P_n the path with n vertices and S_n the star with n leaves. Define $A_n, B_{m_1, n, m_2}, D_n, E_{m, n}, F_n$ as below. The spectral radii of these graphs satisfy: (a) $\lambda_1(A_n) \nearrow 3/\sqrt{2}$ ¹, (b₁) $\lambda_1(B_{1, n, 1}) = 2$, (b₂) $\lambda_1(B_{m_1, n, m_2}) \searrow \max(\alpha_{m_1}, \alpha_{m_2})$ for fixed $(m_1, m_2) \neq (1, 1)$, (c) $\lambda_1(C_n) = 2$, (d) $\lambda_1(D_n) \searrow \lambda^*$, (e) $\lambda_1(E_{m, n}) \nearrow \alpha_m$ for fixed m , (f) $\lambda_1(F_n) \nearrow \lambda^*$, (p) $\lambda_1(P_n) = 2 \cos(\frac{\pi}{n+1}) \nearrow 2$, (s) $\lambda_1(S_n) = \sqrt{n}$.*



The work of finding all the limit points of Λ_1 was completed by Shearer.

Theorem 6 (Shearer [She89]). *For any $\lambda \geq \lambda^*$, there exists a sequence of distinct graphs G_1, G_2, \dots such that $\lim \lambda_1(G_i) = \lambda$.*

Equipped with these facts from spectral graph theory, we are in the position to prove Theorem 1.

¹It was mistakenly asserted that $\lambda_1(A_n) \nearrow 4/\sqrt{3}$ in [Hof72]. However it will not affect the main result of [Hof72] as the correct limit $3/\sqrt{2}$ is still $> \lambda^*$.

Proof of Theorem 1. Note that Lemma 5(b₂) and Theorem 6 imply that $\{\alpha_2, \alpha_3, \dots\} \cup [\lambda^*, \infty) \subseteq \lim_+ \Lambda_1$. Thus by Proposition 3, the second half of Theorem 1 is proved. For the first half of Theorem 1, we break the proof into two cases.

Case 1: $\lambda < 2$. By Lemma 5, we have $S_4 \notin \mathcal{F}(\lambda)$ and $P_n \notin \mathcal{F}(\lambda)$ for some n . Clearly, a connected graph G that contains neither S_4 nor P_n has maximum degree ≤ 3 and diameter $< n$. Therefore

$$\{S_4, P_n\} \cup \{\text{connected graph } G \notin \mathcal{F}(\lambda) \text{ with maximum degree } \leq 3 \text{ and diameter } < n\}$$

is a finite forbidden subgraphs characterization for $\mathcal{F}(\lambda)$.

Case 2: $\lambda \in [2, \lambda^*) \setminus \{\alpha_2, \alpha_3, \dots\}$. Choose the smallest m such that $\alpha_m > \lambda$. Observe that $\lambda \in [2, \alpha_2)$ when $m = 2$, and $\lambda \in (\alpha_{m-1}, \alpha_m)$ when $m \geq 3$. Then choose, in view of Lemma 5, $n > m$ such that $A_n, E_{m,n}, F_n \notin \mathcal{F}(\lambda)$ and $B_{m_1, n', m_2} \in \mathcal{F}(\lambda)$ for all $m_1, m_2 < m$ and $n' > n$. In fact, when $m = 2$, the requirement that $B_{m_1, n', m_2} \in \mathcal{F}(\lambda)$ for all $m_1, m_2 < m$ and $n' > n$ is redundant because of Lemma 5(b₁). By Lemma 5, we also have $S_5 \notin \mathcal{F}(\lambda)$, $B_{m,0,1}, B_{m,1,1}, \dots, B_{m,n,1} \notin \mathcal{F}(\lambda)$ and $D_2, D_3, \dots, D_{m+n} \notin \mathcal{F}(\lambda)$.

We claim that if a connected graph G contains none of

$$\mathcal{G}_0 := \{S_5, A_n, B_{m,0,1}, B_{m,1,1}, \dots, B_{m,n,1}, D_2, D_3, \dots, D_{m+n}, E_{m,n}, F_n\},$$

then G is a path, or a cycle, or E_{m_0, n_0} for some $m_0 < m$ and some n_0 , or B_{m_1, n', m_2} for some $m_1, m_2 < m$ and some $n' > n$, or G has maximum degree ≤ 4 and radius $< m + n$. Recall that the eccentricity of a vertex v in G is the maximum distance between v and any other vertex of G , and the radius of G is the minimum eccentricity of any vertex in G .

Notice that, by Lemma 5, the spectral radii of paths and cycles are at most 2, the spectral radius of E_{m_0, n_0} is $< \alpha_{m_0} \leq \lambda$ for all $m_0 < m$ and n_0 by the choice of m , and the spectral radius of B_{m_1, n', m_2} is $\leq \lambda$ for all $m_1, m_2 < m$ and $n' > n$ by the choice of n . The claim thus implies that

$$\mathcal{G}_0 \cup \{\text{connected graph } G \notin \mathcal{F}(\lambda) \text{ with maximum degree } \leq 4 \text{ and radius } < m + n\}$$

is a finite forbidden subgraphs characterization for $\mathcal{F}(\lambda)$.

Finally, it suffices to prove the claim. Because G does not contain $D_2, \dots, D_{m+n}, E_{m,n}$ and $D_{m+n+1}, D_{m+n+2}, \dots$ contain $E_{m,n}$, the graph G does not contain D_2, D_3, \dots , and so G must be a tree or a cycle. Suppose that G is not a path or a cycle. As G does not contain S_5 , G is a tree of maximum degree 3 or 4. If the degree of a vertex v is 4, then as G does not contain A_n , the eccentricity of v is $< n$ and so the radius of G is $< n$. If a vertex v of degree 3 is not adjacent to any leaf, then as G does not contain F_n , the eccentricity of v is $< n$ and so the radius of G is $< n$.

Hereafter, we may assume that the maximum degree of G is 3 and that every vertex of degree 3 is adjacent to a leaf, in other words, G is a caterpillar² of maximum degree 3. Let v_1, \dots, v_ℓ be the central path of G . Because G does not contain $B_{m,0,1}, B_{m,1,1}, \dots, B_{m,n,1}, E_{m,n}$ and $B_{m,n+1,1}, B_{m,n+2,1}, \dots$ contain $E_{m,n}$, the graph G does not contain $B_{m,0,1}, B_{m,1,1}, B_{m,2,1}, \dots$, and so for every pair of vertices

²A caterpillar is a tree in which all the vertices are within distance 1 of a central path.

v_i, v_j of degree 3 with $i < j$ the distances between v_1, v_i and between v_j, v_ℓ are $< m$. We finish the proof of the claim by checking the following three cases.

1. If G has only one vertex of degree 3, then as G does not contain $E_{m,n}$, the graph G must be E_{m_0, n_0} for some $m_0 < m$ and some n_0 .
2. If G has exactly two vertices of degree 3, then the graph G must be B_{m_1, n', m_2} for some $m_1, m_2 < m$ and some n' . In this case, either $n' > n$ or the radius of G is $\leq \frac{1}{2}(2m + n) < m + n$.
3. If G has at least 3 vertices, say v_i, v_j, v_k with $i < j < k$, of degree 3, then the distances from v_j to v_1 and to v_ℓ are $< m$, hence the radius of G is $< m$. \square

3 Equiangular lines

3.1 The framework to estimate $N_\alpha(n)$

We shall set the stage for the proof of Theorem 2 by adapting the framework developed in Section 2.1 of [BDKS18] to estimate $N_\alpha(n)$.

Definition 2. Let L be a subset of the interval $[-1, 1)$. A finite set C of unit vectors in \mathbb{R}^n is called a *spherical L -code* if $\langle v_1, v_2 \rangle \in L$ for any pair of distinct vectors v_1, v_2 in C . The *Gram matrix* M is given by $M_{ij} = \langle v_i, v_j \rangle$. The *underlying graph* G is defined as follows: let C be its vertex set, and for any distinct $v_i, v_j \in C$, we put the edge (v_i, v_j) if and only if $\langle v_i, v_j \rangle < 0$.

By choosing a unit vector in the direction of each line in a family of equiangular lines with angle $\arccos \alpha$ in \mathbb{R}^n , we obtain a spherical $\{\pm\alpha\}$ -code C_α in \mathbb{R}^n . The key insight of Lemma 2.8 of Balla et al. [BDKS18] is that one can obtain a spherical $L(\alpha, t)$ -code C in \mathbb{R}^n , where

$$L(\alpha, t) := \left\{ -\frac{1}{\lambda} \cdot \left(1 - \frac{1}{t + \alpha^{-1}} \right) + \frac{1}{t + \alpha^{-1}}, \frac{1}{t + \alpha^{-1}} \right\}, \lambda := \frac{1 - \alpha}{2\alpha} \text{ and } t = \lfloor \log \log n \rfloor,$$

from a spherical $\{\pm\alpha\}$ -code C_α such that $|C_\alpha| \leq |C| + o(n)$, and moreover the size of C is much easier to bound from above than C_α . However, to estimate $N_\alpha(n)$ up to a constant error relative to α , we need instead $|C_\alpha| \leq |C| + O(1)$, which requires t be independent from n . Although Lemma 7 is stated differently from Lemma 2.8 of Balla et al. [BDKS18], the same proof goes through without alternation. For the convenience of the readers, we include here a streamlined proof.

Lemma 7. *Let $\alpha \in (0, 1)$ and $t \in \mathbb{N}$ be fixed such that $t > \left(\frac{1-\alpha}{2\alpha}\right)^2 + 1$. For any $n \in \mathbb{N}$ and any spherical $\{\pm\alpha\}$ -code C_α in \mathbb{R}^n , there exists a spherical $L(\alpha, t)$ -code C in \mathbb{R}^n such that $|C_\alpha| \leq |C| + O_{\alpha, t}(1)$.*

Proof. Denote by $R(\cdot, \cdot)$ the Ramsey number. Let G be the underlying graph of C_α . The clique number of the underlying graph G is $\leq 1 + \lfloor 1/\alpha \rfloor$. In fact, if K is a clique in G , then

$$0 \leq \left\| \sum_{v \in K} v \right\|^2 = \sum_{v \in K} \|v\|^2 + \sum_{\substack{v_1, v_2 \in K \\ v_1 \neq v_2}} \langle v_1, v_2 \rangle = |K| - |K|(|K| - 1)\alpha \implies |K| \leq 1 + 1/\alpha.$$

If $|C_\alpha| < R(2 + \lfloor 1/\alpha \rfloor, t)$, then we are done by taking $C = \emptyset$. Otherwise, by Ramsey's theorem, G contains an independent set I of size t . For every $v \in C_\alpha \setminus I$, we may opt to switch v to $-v$ to ensure that the degree of v to I is at most $t/2$. We partition $C_\alpha \setminus I$ by how the vertices are connected to I . In particular, each part in the partition is indexed by some $I_1 \subset I$ with $|I_1| \leq t/2$, and is defined by

$$C_\alpha(I_1) := \{v \in C_\alpha \setminus I : \forall u \in I \setminus I_1 \ u \not\sim v \text{ and } \forall u \in I_1 \ u \sim v\}.$$

For every $I_1 \subset I$ such that $I_1 \neq \emptyset$ and $|I_1| \leq t/2$, we bound the independent number of the subgraph $G[C_\alpha(I_1)]$ induced on $C_\alpha(I_1)$ as follows. Suppose that $J \subseteq C_\alpha(I_1)$ is an independent set of $G[C_\alpha(I_1)]$. Let $I_0 := I \setminus I_1$ and set

$$u_0 := \sum_{v \in I_0} v, \quad u_1 := \sum_{v \in I_1} v, \quad u_2 := \sum_{v \in J} v.$$

One can work out the Gram matrix of u_0, u_1, u_2 :

$$\begin{pmatrix} |I_0| + |I_0|(|I_0| - 1)\alpha & |I_0||I_1|\alpha & |I_0||J|\alpha \\ |I_0||I_1|\alpha & |I_1| + |I_1|(|I_1| - 1)\alpha & -|I_1||J|\alpha \\ |I_0||J|\alpha & -|I_1||J|\alpha & |J| + |J|(|J| - 1)\alpha \end{pmatrix}.$$

Because the Gram matrix is positive semidefinite, its determinant

$$|I_0||I_1||J| \left((1 - \alpha)^2(1 - \alpha + \alpha(|I_0| + |I_1|)) - 4\alpha^3 \left(|I_0||I_1| - \left(\frac{1 - \alpha}{2\alpha} \right)^2 \right) |J| \right)$$

is non-negative. Together with $|I_0| + |I_1| = |I| = t$ and $|I_0||I_1| \geq t - 1 > \left(\frac{1 - \alpha}{2\alpha} \right)^2$, we obtain a bound on the size of the independent set J :

$$|J| \leq \frac{(1 - \alpha)^2(1 - \alpha + \alpha(|I_0| + |I_1|))}{4\alpha^3 \left(|I_0||I_1| - \left(\frac{1 - \alpha}{2\alpha} \right)^2 \right)} = O_{\alpha, t}(1).$$

Again, since the clique number of $G[C_\alpha(I_1)]$ is $\leq 1 + \lfloor 1/\alpha \rfloor$, by Ramsey's theorem, $|C_\alpha(I_1)| = O_{\alpha, t}(1)$. We thus have the estimation

$$|C_\alpha| = |I| + |C_\alpha(\emptyset)| + \sum_{\substack{\emptyset \neq I_1 \subset I \\ |I_1| \leq t/2}} |C_\alpha(I_1)| = |C_\alpha(\emptyset)| + O_{\alpha, t}(1).$$

Pick any $v \in C_\alpha(\emptyset)$ and consider the projection $p(v)$ of v onto the orthogonal complement of $\text{span}(I)$. The vector $p(v)$ is given by $v - \sum_{u \in I} c_u u$ for some $\mathbf{c} = \{c_u : u \in I\}$ that minimizes $m(\mathbf{c}) := \|v - \sum_{u \in I} c_u u\|^2$. Since $\{v\} \cup I$ is an independent set, the quadratic function $m(\mathbf{c})$ is symmetric with respect to $\{c_u : u \in I\}$, and so it is minimized at $\mathbf{c} = c\mathbf{1}$ for some $c \in \mathbb{R}$. Note that

$$m(c\mathbf{1}) = \left\| v - c \sum_{u \in I} u \right\|^2 = \|v\|^2 - 2c \sum_{u \in I} \langle v, u \rangle + c^2 \left\| \sum_{u \in I} u \right\|^2 = 1 - 2ct\alpha + c^2(t + t(t - 1)\alpha),$$

which is minimized at

$$c := \frac{\alpha}{1 + (t-1)\alpha}. \quad (1)$$

Therefore $p(v) = v - c \sum_{u \in I} u$.

Denote by $\hat{p}(v) := p(v)/\|p(v)\|$ the normalized vector of $p(v)$. Lastly we show that

$$C := \{\hat{p}(v) : v \in C_\alpha(\emptyset)\}$$

is a spherical $L(\alpha, t)$ -code. For any $v_1, v_2 \in C_\alpha(\emptyset)$, we compute $\langle p(v_1), p(v_2) \rangle$ in terms of $\langle v_1, v_2 \rangle$:

$$\begin{aligned} \langle p(v_1), p(v_2) \rangle &= \left\langle v_1 - c \sum_{u \in I} u, v_2 - c \sum_{u \in I} u \right\rangle = \langle v_1, v_2 \rangle - c \sum_{u \in I} \langle v_1, u \rangle - c \sum_{u \in I} \langle v_2, u \rangle + c^2 \left\| \sum_{u \in I} u \right\|^2 \\ &= \langle v_1, v_2 \rangle - ct\alpha - ct\alpha + c^2(t + t(t-1)\alpha) \stackrel{(1)}{=} \langle v_1, v_2 \rangle - 2ct\alpha + ct\alpha = \langle v_1, v_2 \rangle - ct\alpha. \end{aligned} \quad (2)$$

For any $v_1, v_2 \in C_\alpha(\emptyset)$ such that $v_1 \neq v_2$, we get

$$\langle \hat{p}(v_1), \hat{p}(v_2) \rangle = \frac{\langle p(v_1), p(v_2) \rangle}{\|p(v_1)\| \|p(v_2)\|} \stackrel{(2)}{=} \frac{\langle v_1, v_2 \rangle - ct\alpha}{\sqrt{\langle v_1, v_1 \rangle - ct\alpha} \sqrt{\langle v_2, v_2 \rangle - ct\alpha}} = \frac{\langle v_1, v_2 \rangle - ct\alpha}{1 - ct\alpha},$$

which takes its value in $\left\{ \frac{\pm\alpha - ct\alpha}{1 - ct\alpha} \right\} \stackrel{(1)}{=} L(\alpha, t)$. \square

The following lemma, which will be applied in conjunction with Lemma 7, imposes structural restriction on the underlying graph of a spherical $L(\alpha, t)$ -code.

Lemma 8. *Given $\alpha \in (0, 1)$ and a finite family \mathcal{G} of graphs with spectral radius $> \lambda := \frac{1-\alpha}{2\alpha}$, there exists $t \in \mathbb{N}$ such that the underlying graph of any spherical $L(\alpha, t)$ -code does not contain any graph in \mathcal{G} as a subgraph.*

We recall a necessary condition on eigenvalues of the sum of two matrices.

Theorem 9 (Weyl's inequality [Wey12]). *Given two $n \times n$ Hermitian matrices A and B . Denote the eigenvalues of A as $\lambda_1(A) \geq \lambda_2(A) \geq \dots \geq \lambda_n(A)$, and similarly denote the eigenvalues of B and $A + B$. Whenever $0 \leq i, j, i + j < n$, $\lambda_{i+j+1}(A + B) \leq \lambda_{i+1}(A) + \lambda_{j+1}(B)$.*

We shall denote by I_n the identity matrix of order n and J_n the all-ones matrix of order n , and we suppress the subscripts when the order of the matrices could be inferred from the context.

Proof of Lemma 8. We first choose $t \in \mathbb{N}$ depending only on α and \mathcal{G} . Because $\lambda_1(G_0) > \lambda$ for every $G_0 \in \mathcal{G}$, we thus choose $t \in \mathbb{N}$ large enough so that

$$1 - \frac{\lambda_1(G_0)}{\lambda} + \frac{v(G_0)}{t + \alpha^{-1} - 1} < 0 \text{ for all } G_0 \in \mathcal{G}, \quad (3)$$

where $v(G_0)$ is the order in G_0 .

We claim that the underlying graph G of any spherical $L(\alpha, t)$ -code does not contain any graph in \mathcal{G} as a subgraph. Suppose on the contrary that G contains some graph $G_0 \in \mathcal{G}$. Let G'_0 be the minimal induced subgraph of G that contains G_0 as a subgraph, and C'_0 the vertex set of G'_0 . Note that $v(G'_0) = v(G_0)$. By the monotonicity of λ_1 and the choice of t in (3), we obtain

$$1 - \frac{\lambda_1(G'_0)}{\lambda} + \frac{v(G'_0)}{t + \alpha^{-1} - 1} < 0. \quad (4)$$

Let M'_0 be the Gram matrix of C'_0 and let A'_0 be the adjacency matrix of G'_0 . Since C'_0 is still a spherical $L(\alpha, t)$ -code, the two matrices M'_0 and A'_0 are related by the equation

$$\left(1 + \frac{1}{t + \alpha^{-1} - 1}\right) M'_0 = I - \frac{A'_0}{\lambda} + \frac{J}{t + \alpha^{-1} - 1}.$$

Using the fact that $\lambda_1(J) = v(G'_0)$ and (4), we know from Weyl's inequality that the least eigenvalue of M'_0 is negative. This contradicts with the fact that a Gram matrix is positive semidefinite. \square

3.2 Application of Theorem 1

Our application of Theorem 1 addresses the upper bound on $N_\alpha(n)$ in Theorem 2 when $\mathcal{F}(\lambda)$ has a finite forbidden subgraphs characterization.

Corollary 10. *Define $\lambda^*, \alpha_2, \alpha_3, \dots$ as in Theorem 1. Given $\alpha \in (0, 1)$ such that $\lambda := \frac{1-\alpha}{2\alpha} < \lambda^*$ and $\lambda \notin \{\alpha_2, \alpha_3, \dots\}$, the maximum number $N_\alpha(n)$ of equiangular lines in \mathbb{R}^n with angle $\arccos \alpha$ is at most $\frac{k(\lambda)}{k(\lambda)-1} \cdot n + O(1)$, where $k(\lambda)$ is the spectral radius order of λ ; in case $k(\lambda) = \infty$, $N_\alpha(n) \leq n + O(1)$.*

We recall a classical fact about the spectral radius of a connected graph and prove an immediate consequence.

Theorem 11 (Corollary of the Perron–Frobenius theorem [Fro12, Per07]). *For every connected graph G , the largest eigenvalue $\lambda_1(G)$ has multiplicity 1, with an eigenvector whose components are all positive.*

Corollary 12. *For every $\lambda > 0$ and every connected graph G with spectral radius $\leq \lambda$,*

$$v(G) \leq \frac{k(\lambda)}{k(\lambda) - 1} \cdot \text{rank} \left(I - \frac{A}{\lambda} \right),$$

where $v(G)$ is the order of G , $k(\lambda)$ is the spectral radius order of λ , and A is the adjacency matrix of G ; in case $k(\lambda) = \infty$, $v(G) \leq \text{rank} (I - A/\lambda)$.

Proof. If $\lambda_1(G) = \lambda$, the definition of $k(\lambda)$ implies that $v(G) \geq k(\lambda)$, and so the Perron–Frobenius theorem implies that $\text{rank} (I - A/\lambda) = v(G) - 1 \geq (1 - 1/k(\lambda))v(G)$; otherwise $\lambda_1(G) \neq \lambda$, or equivalently $\lambda_1(G) < \lambda$, and so $\text{rank} (I - A/\lambda) = v(G)$. Therefore when $k(\lambda) < \infty$ we always have $\text{rank} (I - A/\lambda) \geq (1 - 1/k(\lambda))v(G)$; in case $k(\lambda) = \infty$, the definition of $k(\lambda)$ implies that $\lambda_1(G) \neq \lambda$, and so $\text{rank} (I - A/\lambda) \geq v(G)$. \square

The last preparation for the proof of Corollary 10 links $\text{rank}(I - A/\lambda)$ with the dimension n .

Proposition 13. *Given $\alpha \in (0, 1)$, $t \in \mathbb{N}$, for any spherical $L(\alpha, t)$ -code C in \mathbb{R}^n , the adjacency matrix A of the underlying graph of C satisfies $\text{rank}(I - A/\lambda) \leq n + 1$, where $\lambda := \frac{1-\alpha}{2\alpha}$.*

Proof. Let M be the Gram matrix of C . Since C is a spherical $L(\alpha, t)$ -code, the matrices M and A are related by

$$\left(1 + \frac{1}{t + \alpha^{-1} - 1}\right)M = I - \frac{A}{\lambda} + \frac{J}{t + \alpha^{-1} - 1}.$$

Therefore $\text{rank}(I - A/\lambda) \leq \text{rank}(M) + \text{rank}(J) \leq n + 1$. \square

Proof of Corollary 10. Since $\lambda = \frac{1-\alpha}{2\alpha} \in (0, \lambda^*) \setminus \{\alpha_2, \alpha_3, \dots\}$, by Theorem 1, there exists a finite family \mathcal{G} of graphs such that $\mathcal{F}(\lambda)$ consists exactly of the connected graphs which do not contain any graph in \mathcal{G} as a subgraph.

Suppose C_α is a spherical $\{\pm\alpha\}$ -code in \mathbb{R}^n . By Lemma 7 and Lemma 8, we can choose $t \in \mathbb{N}$ depending only on α such that there exists a spherical $L(\alpha, t)$ -code C in \mathbb{R}^n with $|C_\alpha| \leq |C| + O(1)$ whose underlying graph G does not contain any graph in \mathcal{G} .

Let G_1, \dots, G_m be the connected components of G . By the claim that G does not contain any graph in \mathcal{G} , neither does G_i for every $i \in [m]$. Because \mathcal{G} is a forbidden subgraphs characterization for $\mathcal{F}(\lambda)$, the connected graph $G_i \in \mathcal{F}(\lambda)$, or in other words, $\lambda_1(G_i) \leq \lambda$. Let $v(G_i)$ be the order of G_i and A_i the adjacency matrix of G_i . By Corollary 12 and Proposition 13, we have

$$|C| = \sum_{i=1}^m v(G_i) \leq \frac{k(\lambda)}{k(\lambda) - 1} \sum_{i=1}^m \text{rank}\left(I - \frac{A_i}{\lambda}\right) = \frac{k(\lambda)}{k(\lambda) - 1} \cdot \text{rank}\left(I - \frac{A}{\lambda}\right) \leq \frac{k(\lambda)}{k(\lambda) - 1} \cdot (n + 1).$$

In case $k(\lambda) = \infty$, by Corollary 12 and Proposition 13, we have

$$|C| = \sum_{i=1}^m v(G_i) \leq \sum_{i=1}^m \text{rank}\left(I - \frac{A_i}{\lambda}\right) = \text{rank}\left(I - \frac{A}{\lambda}\right) \leq n + 1. \quad \square$$

3.3 Proof of Theorem 2

The following lemma connects a spherical $\{\pm\alpha\}$ -code with a spherical $\{-1/\lambda, 0\}$ -code, and it provides the lower bound on $N_\alpha(n)$ in Theorem 2.

Lemma 14. *Given $\alpha \in (0, 1)$, let $\lambda := \frac{1-\alpha}{2\alpha}$. For any spherical $\{-1/\lambda, 0\}$ -code C_0 in \mathbb{R}^k , there exists a spherical $\{\pm\alpha\}$ -code in \mathbb{R}^n of size $\lfloor \frac{n-1}{k} \rfloor |C_0|$. In particular, the maximum number $N_\alpha(n)$ of equiangular lines in \mathbb{R}^n with angle $\arccos \alpha$ is at least $\lfloor \frac{n-1}{k(\lambda)-1} \rfloor k(\lambda)$, where $k(\lambda)$ is the spectral radius order of λ ; in case $k(\lambda) = \infty$, we have $N_\alpha(n) \geq n$.*

Proof. Given a spherical $\{-1/\lambda, 0\}$ -code C_0 in \mathbb{R}^k . Let $m := \lfloor \frac{n-1}{k} \rfloor$ and M_0 be the Gram matrix of C_0 . Consider the matrix $M := (1 - \alpha)M_0 \otimes I_m + \alpha J$ of order $m|C_0|$. For both M_0 and J are positive semidefinite, so is M . The rank of M is at most $m \cdot \text{rank}(M_0) + 1 \leq mk + 1 \leq n$. Moreover, the

diagonal entries of M are ones and its off-diagonal entries are either $-\alpha$ or α . Therefore M can be realized as the Gram matrix of a spherical $\{\pm\alpha\}$ -code of size $m|C_0|$ in \mathbb{R}^n .

It suffices to construct a spherical $\{-1/\lambda, 0\}$ -code C_0 in $\mathbb{R}^{k(\lambda)-1}$ of size $k(\lambda)$. Because λ is the spectral radius of a graph G on $k(\lambda)$ vertices, $I - A/\lambda$ is positive semidefinite, where A is the adjacency matrix of G , and $\text{rank}(I - A/\lambda) \leq k(\lambda) - 1$. Clearly, $I - A/\lambda$ can be realized as the Gram matrix of a spherical $\{-1/\lambda, 0\}$ -code of size $k(\lambda)$ in $\mathbb{R}^{k(\lambda)-1}$.

In case $k(\lambda) = \infty$, $(1 - \alpha)I_n + \alpha J_n$ can be realized as the Gram matrix of a spherical $\{\pm\alpha\}$ -code of size n in \mathbb{R}^n . \square

In view of Corollary 10 and Lemma 14, we are left to prove Theorem 2 for the exceptional $\lambda \in \{\alpha_2, \alpha_3, \dots\} \cup \{\lambda^*\}$. Clearly, if λ is an eigenvalue of the adjacency matrix of a graph, then

1. λ is an algebraic integer — it is a root of some monic polynomial with coefficients in \mathbb{Z} ,
2. λ is totally real — its conjugate elements are in \mathbb{R} .

On the converse, it follows from Estes [Est92, Theorem 1] that any totally real algebraic integer occurs as an eigenvalue of the adjacency matrix of a graph.

When λ is not a totally real algebraic integer, the spectral radius order $k(\lambda) = \infty$, and Conjecture B predicts that $N_\alpha(n) = n + O(1)$, which we prove in the affirmative.

Proposition 15. *Given $\alpha \in (0, 1)$, let $\lambda := \frac{1-\alpha}{2\alpha}$. If λ is not a totally real algebraic integer, then $n \leq N_\alpha(n) \leq n + 1$.*

Proof. Let C be a spherical $\{\pm\alpha\}$ -code in \mathbb{R}^n . Let M be its Gram matrix, and A the adjacency matrix of the underlying graph G . We know that $M = (1 - \alpha)(I - A/\lambda) + \alpha J$. Since λ is not an eigenvalue of A , $\text{rank}(I - A/\lambda) = |C|$ and so $n \geq \text{rank}(M) \geq \text{rank}(I - A/\lambda) - \text{rank}(J) = |C| - 1$. Together with Lemma 14, we have $n \leq N_\alpha(n) \leq n + 1$. \square

Observe that one of the conjugate elements $\sqrt{2 - \sqrt{5}}$ of $\lambda^* = \sqrt{2 + \sqrt{5}}$ is not real. Proposition 15 readily gives the proof of Theorem 2 for $\lambda = \lambda^*$. The rest of the cases $\lambda \in \{\alpha_2, \alpha_3, \dots\}$ in Theorem 2 follow similarly from the following proposition, the proof of which is due in Appendix B.

Proposition 16. *The algebraic integers $\alpha_2, \alpha_3, \dots$, defined as in Theorem 1, are not totally real.*

3.4 An improved upper bound on $N_\alpha(n)$

In this section, we prove $N_\alpha(n) \leq 1.49n + O(1)$ for every $\alpha \neq 1/3, 1/5, 1/(1 + 2\sqrt{2})$. We recall the following spectral tool bounding the rank of a symmetric matrix in terms of its trace and the trace of its square.

Lemma 17. *The rank of a symmetric matrix A is greater than or equal to $(\text{tr}(A))^2/\text{tr}(A^2)$.*

This lemma appeared as Inequality 14 of Bellman [Bel97, page 137] in the form of an exercise. We refer the readers to [BDKS18, Lemma 2.9] for a short proof of Lemma 17. Additionally, we develop a spectral result, which can be seen as an averaged version of the following lemma.

Lemma 18 (Lemma 2.13 of Balla et al. [BDKS18]). *Let G be a graph with minimum degree $\delta \geq 2$. Let v_0 be a vertex of G and let H be the subgraph consisting of all vertices within distance k of v_0 . Then $\lambda_1(H) \geq \frac{2k}{k+1} \sqrt{\delta - 1}$.*

Lemma 19. *Let G be a graph with average degree $d \geq 2$. There exists a vertex v_0 of G such that $\lambda_1(H) \geq 2 \cos(\frac{\pi}{k+2}) \sqrt{d - 1}$, where H is the subgraph consisting of all vertices within distance k of v_0 .*

By Lemma 5(p), the coefficient $2 \cos(\frac{\pi}{k+2})$ in Lemma 19 equals $\lambda_1(P_{k+1})$, where P_{k+1} is the path with k edges. Notice that $\lambda_1(P_{k+1})$ is greater than the average degree $\frac{2k}{k+1}$ of P_{k+1} , which is the coefficient in Lemma 18. This comparison between Lemma 18 and Lemma 19 is reminiscent of that between Nilli's [Nil91] and Friedman's [Fri93, Corollary 3.7] (see also [Nil04]) proofs of the Alon–Boppana bound on the second largest eigenvalue of a regular graph. In fact, Lemma 19 was recently applied [Jia19, Theorem 8] to improve Hoory's bound [Hoo05, Theorem 3] on the second largest eigenvalue for a class of graphs that are not necessarily regular.

Proof. Since removing leaf vertices from a graph of average degree $d \geq 2$ cannot decrease its average degree, without loss of generality, we may assume that the minimum degree of G is ≥ 2 . A walk $(v_{-1}, v_0, v_1, \dots)$ on G is non-backtracking if $v_i \neq v_{i+2}$ for all i . For all $i \geq 0$, define W_i to be the set of all non-backtracking walks $(v_{-1}, v_0, v_1, \dots, v_i)$ on G of length $i + 1$. Define the forest T as follows: the vertex set is $\cup_{i=0}^k W_i$ and two vertices are adjacent if and only if one is a simple extension of the other. For every $e = (v_{-1}, v_0) \in W_0$, denote by T_e the connected component of T containing e . We also denote by G_{v_0} the subgraph of G consisting of all vertices within distance k of v_0 .

We claim that $\lambda_1(T_e) \leq \lambda_1(G_{v_0})$ for every $e = (v_{-1}, v_0) \in W_0$. Let s_i and t_i be the number of closed walks of length i starting respectively at e in T_e and v_0 in G_{v_0} . It is well known that $\lambda_1(T_e) = \limsup \sqrt[i]{s_i}$ and $\lambda_1(G_{v_0}) = \limsup \sqrt[i]{t_i}$. We naturally map a closed walk $e = e_0, e_1, \dots, e_i = e$ in T_e to a closed walk $v_0, v_1, \dots, v_i = v_0$ in G_{v_0} , where v_j is the terminal vertex of the non-backtracking walk e_j for $j = 0, 1, \dots, i$. One can show that this map is injective, and so $s_i \leq t_i$ for all i , from which the claim follows.

Because $\lambda_1(T) = \max \{\lambda_1(T_e) : e \in W_0\}$, it suffices to prove $\lambda_1(T) \geq \lambda \sqrt{d - 1}$, where $\lambda := \lambda_1(P_{k+1}) = 2 \cos(\frac{\pi}{k+2})$. Consider the non-backtracking random walk on T , where the start vertex $w_0 = (v_{-1}, v_0)$ is chosen uniformly at random from W_0 and, for $i \in [k]$, at i th step the next vertex $w_i = (v_{-1}, v_0, \dots, v_i)$ is chosen uniformly at random among the available choices in W_i . The transition matrix of this walk is

$$P_{(v_{-1}, v_0, \dots, v_i), (v_{-1}, v_0, \dots, v_{i+1})} = \frac{1}{d(v_i) - 1},$$

where $d(v)$ denotes the degree of v in G . Clearly $|W_0| = d|V(G)|$. Since W_i is a finite set and $w_i = (v_{-1}, v_0, \dots, v_i)$ is a random element of W_i with distribution

$$p(w_i) := \frac{1}{d|V(G)|} \prod_{j=0}^{i-1} \frac{1}{d(v_j) - 1},$$

for any $c: W_i \rightarrow \mathbb{R}$, the basic identity of importance sampling allows us to represent $\sum_{w \in W_i} c(w)$ as follows:

$$\sum_{w \in W_i} c(w) = \sum_{w \in W_i} p(w) \frac{c(w)}{p(w)} = \mathbb{E}[c(w_i)/p(w_i)]. \quad (5)$$

Let $(x_0, x_1, \dots, x_k) \in \mathbb{R}^{k+1}$ be an eigenvector of P_{k+1} such that $x_0, x_1, \dots, x_k > 0$. Define the vector $f: V(T) \rightarrow \mathbb{R}$ by $f(w) = x_i \sqrt{p(w)}$ for $w \in W_i$, and define the matrix A to be the adjacency matrix of the forest T . For $w = (u_{-1}, u_0, \dots, u_i)$, denote by $w^- = (u_{-1}, u_0, \dots, u_{i-1})$. By the importance sampling identity (5), we observe that

$$\begin{aligned} \langle f, f \rangle &= \sum_{i=0}^k \sum_{w \in W_i} f(w)^2 = \sum_{i=0}^k \mathbb{E}[x_i^2] = \sum_{i=0}^k x_i^2, \\ \frac{1}{2} \langle f, Af \rangle &= \sum_{i=1}^k \sum_{w \in W_i} f(w^-) f(w) = \sum_{i=1}^k \mathbb{E} \left[x_{i-1} x_i \sqrt{p(w^-)/p(w)} \right] = \sum_{i=1}^k x_{i-1} x_i \mathbb{E} \left[\sqrt{d(v_{i-1}) - 1} \right]. \end{aligned}$$

It can be verified by induction that (v_{i-1}, v_i) is uniformly distributed on W_0 for all $i = 0, 1, \dots, k$. Thus $\Pr(v_{i-1} = v) = \frac{d(v)}{d|V(G)|} =: \pi(v)$ for all $v \in V(G)$ and $i = 1, 2, \dots, k$. Since each v_{i-1} has distribution π and the function $x \mapsto x\sqrt{x-1}$ is convex on $[2, \infty]$, Jensen's inequality gives

$$\frac{1}{2} \langle f, Af \rangle = \sum_{i=1}^k x_{i-1} x_i \sum_{v \in V(G)} \frac{d(v)}{d|V(G)|} \sqrt{d(v) - 1} \geq \sqrt{d-1} \sum_{i=1}^k x_{i-1} x_i.$$

Finally we invoke the Rayleigh principle $2 \sum_{i=1}^k x_{i-1} x_i = \lambda \sum_{i=0}^k x_i^2$ and $\lambda_1(T) \geq \langle f, Af \rangle / \langle f, f \rangle$. \square

Remark 2. Lemma 19 is asymptotically tight when d is a prime plus one due to the existence of regular graphs of high girth. The Ramanujan graphs constructed independently by Margulis [Mar82] and Lubotzky, Phillips and Sarnak [LPS88] are d -regular graphs on n vertices of girth $\Omega_d(\log n)$. In these Ramanujan graphs, every subgraph induced on the vertices within a bounded distance of a given vertex looks like a d -regular tree, whose spectral radius is bounded from above by the spectral radius $2\sqrt{d-1}$ of the infinite d -regular tree.

Remark 3. See the expository note by Levin and Peres [LP17] for other applications of Markov chains and importance sampling.

Theorem 20. *Given $\alpha \in (0, 1)$, let $\lambda := \frac{1-\alpha}{2\alpha}$. If $\lambda \geq 2$, then the maximum number $N_\alpha(n)$ of equiangular lines in \mathbb{R}^n is at most $(1 + \frac{1}{4} + \frac{1}{\lambda^2} + o_\alpha(1))n$. In particular, $N_{1/7}(n) \leq (\frac{4}{3} + \frac{1}{36} + o(1))n$.*

Proof. For a fixed $\varepsilon > 0$, we shall prove that the size a spherical $\{\pm\alpha\}$ -code C_α in \mathbb{R}^n is at most $(1 + \frac{1}{4} + \frac{1}{\lambda^2} + \varepsilon)n + O(1)$. Choose $k \in \mathbb{N}$ so that

$$\lambda' := 2 \cos \left(\frac{\pi}{k+2} \right) > \frac{2}{\sqrt{1+4\varepsilon}}. \quad (6)$$

We first find a finite family \mathcal{G}_0 of graphs of spectral radius $> \lambda$ that approximates a forbidden subgraphs characterization of $\mathcal{F}(\lambda)$ in the following sense: if a connected graph G does not contain any graph in \mathcal{G}_0 , then either $G \in \mathcal{F}(\lambda)$ or the average degree of G is at most

$$d := \left(\frac{\lambda}{\lambda'}\right)^2 + 1. \quad (7)$$

Choose $D \in \mathbb{N}$ such that the star $S_D \notin \mathcal{F}(\lambda)$ in view of Lemma 5(s). Suppose a connected graph G does not contain S_D and it has average degree $d_G > d$. Because $\lambda \geq 2$ hence $d_G > d \geq (2/\lambda')^2 + 1 > 2$, Lemma 19 implies that there exists a vertex v_0 of G such that $\lambda_1(H) \geq \lambda' \sqrt{d_G - 1} > \lambda' \sqrt{d - 1} = \lambda$, where H is the subgraph consisting of all vertices within distance k of v_0 . This means that G contains a subgraph $H \notin \mathcal{F}(\lambda)$ with radius $\leq k$. Therefore we can approximate a forbidden subgraphs characterization by

$$\mathcal{G}_0 := \{S_D\} \cup \{\text{connected graph } H \notin \mathcal{F}(\lambda) \text{ with maximum degree } < D \text{ and radius } \leq k\}.$$

By Lemma 7 and Lemma 8, we can choose $t \in \mathbb{N}$ depending only on α such that there exists a spherical $L(\alpha, t)$ -code C in \mathbb{R}^n with $|C_\alpha| \leq |C| + O(1)$ whose underlying graph G does not contain any graph in \mathcal{G}_0 . Let G_1, \dots, G_m be the connected components of G . As G does not contain any graph of \mathcal{G}_0 , neither does G_i for all $i \in [m]$. By our choice of \mathcal{G}_0 , we know that either $\lambda_1(G_i) \leq \lambda$ or the average degree of G_i is $\leq d$.

In the former case that $\lambda_1(G_i) \leq \lambda$, Corollary 12 gives

$$v(G_i) \leq \frac{k(\lambda)}{k(\lambda) - 1} \cdot \text{rank} \left(I - \frac{A_i}{\lambda} \right),$$

where $v(G_i)$ is the order of G_i , $k(\lambda)$ is the spectral radius order of λ , and A_i is the adjacency matrix of G_i . Because the complete graph on $k(\lambda)$ vertices has the largest spectral radius $k(\lambda) - 1$ among all graphs on $k(\lambda)$ vertices, we know that λ , the spectral radius of some graph on $k(\lambda)$ vertices, is at most $k(\lambda) - 1$. Using the fact that $1/\lambda \leq 1/4 + 1/\lambda^2$, we obtain

$$v(G_i) \leq \frac{k(\lambda)}{k(\lambda) - 1} \cdot \text{rank} \left(I - \frac{A_i}{\lambda} \right) \leq \left(1 + \frac{1}{\lambda} \right) \text{rank} \left(I - \frac{A_i}{\lambda} \right) \leq \left(1 + \frac{1}{4} + \frac{1}{\lambda^2} \right) \text{rank} \left(I - \frac{A_i}{\lambda} \right).$$

In case $k(\lambda) = \infty$, by Corollary 12, $v(G_i) \leq \text{rank} (I - A_i/\lambda)$, so the same estimation holds trivially.

In the latter case that the average degree $d(G_i)$ of G_i is $\leq d$, we can apply Lemma 17 to the matrix $I - A_i/\lambda$ and get

$$\text{rank} \left(I - \frac{A_i}{\lambda} \right) \geq \frac{(\text{tr} (I - A_i/\lambda))^2}{\text{tr} ((I - A_i/\lambda)^2)} = \frac{v(G_i)^2}{v(G_i) + v(G_i)d(G_i)/\lambda^2} \geq \frac{v(G_i)}{1 + d/\lambda^2}.$$

Thus we obtain the estimation

$$\begin{aligned} v(G_i) &\leq \left(1 + \frac{d}{\lambda^2} \right) \text{rank} \left(I - \frac{A_i}{\lambda} \right) \stackrel{(7)}{=} \left(1 + \frac{1}{\lambda'^2} + \frac{1}{\lambda^2} \right) \text{rank} \left(I - \frac{A_i}{\lambda} \right) \\ &\stackrel{(6)}{\leq} \left(1 + \frac{1}{4} + \frac{1}{\lambda^2} + \varepsilon \right) \text{rank} \left(I - \frac{A_i}{\lambda} \right). \end{aligned}$$

Summing up these estimations for all the G_i 's and using Proposition 13, we get

$$\begin{aligned} |C| &= \sum_{i=1}^m v(G_i) \leq \left(1 + \frac{1}{4} + \frac{1}{\lambda^2} + \varepsilon\right) \sum_{i=1}^m \text{rank} \left(I - \frac{A_i}{\lambda}\right) \\ &\leq \left(1 + \frac{1}{4} + \frac{1}{\lambda^2} + \varepsilon\right) \text{rank} \left(I - \frac{A}{\lambda}\right) \leq \left(1 + \frac{1}{4} + \frac{1}{\lambda^2} + \varepsilon\right) (n+1). \end{aligned}$$

Recalling that $|C_\alpha| \leq |C| + O(1)$, we get $|C_\alpha| \leq \left(1 + \frac{1}{4} + \frac{1}{\lambda^2} + \varepsilon\right) n + O(1)$. \square

Corollary 21. *Given $\alpha \in (0, 1) \setminus \{1/3, 1/5, 1/(1 + 2\sqrt{2})\}$, the maximum number $N_\alpha(n)$ of equiangular lines in \mathbb{R}^n with angle $\arccos \alpha$ is at most $1.49n + O(1)$.*

Proof. Recall that $\lambda^* = \sqrt{2 + \sqrt{5}} \approx 2.058$. On the one hand, Theorem 2 implies that $N_\alpha(n) \leq \frac{4}{3}n + O(1)$ for $\lambda := \frac{1-\alpha}{2\alpha} \in (0, \lambda^*) \setminus \{1, \sqrt{2}, 2\}$. On the other hand, because $\varepsilon := 0.24 - 1/(\lambda^*)^2 > 0$, Theorem 20 implies that for $\lambda > \lambda^*$, $N_\alpha(n) \leq \left(1 + \frac{1}{4} + \frac{1}{\lambda^2} + \varepsilon\right) n + O(1) \leq 1.49n + O(1)$. \square

4 Concluding remarks

Besides Theorem 2, Lemma 14 and Proposition 15, we discuss two other evidences supporting Conjecture B. Notice that the spectral radius order of λ is at least the algebraic degree $\deg(\lambda)$ of λ . Conjecture B predicts that $N_\alpha(n) \leq \frac{\deg(\lambda)}{\deg(\lambda)-1} \cdot n + O(1)$. This is indeed a cheap upper bound on $N_\alpha(n)$.

Proposition 22. *Given $\alpha \in (0, 1)$, if $\lambda := \frac{1-\alpha}{2\alpha}$ is a totally real algebraic integer, then $N_\alpha(n) \leq \frac{\deg(\lambda)}{\deg(\lambda)-1} \cdot (n+1)$, where $\deg(\lambda)$ is the algebraic degree of λ .*

Proof. Let C be a spherical $\{\pm\alpha\}$ -code in \mathbb{R}^n . Let M be its Gram matrix, and A the adjacency matrix of the underlying graph G . We know that $M = (1 - \alpha)(I - A/\lambda) + \alpha J$. If λ is a totally real algebraic number, then the multiplicity of λ as an eigenvalue of A is $\leq \frac{1}{\deg(\lambda)} |C|$. Thus $\text{rank}(I - A/\lambda) \geq \left(1 - \frac{1}{\deg(\lambda)}\right) |C|$ and so $n \geq \text{rank}(M) \geq \text{rank}(I - A/\lambda) - \text{rank}(J) \geq \left(1 - \frac{1}{\deg(\lambda)}\right) |C| - 1$. \square

Lemma 14 and Proposition 22 would imply Conjecture B in the equality case $k(\lambda) = \deg(\lambda)$. Note that $k(\lambda) = \deg(\lambda)$ if and only if λ is the spectral radius of a graph with irreducible characteristic polynomial. A result of Mowshowitz [Mow71] (see [GM81, Theorem 3.8] for a generalization) states that a graph with irreducible characteristic polynomial has trivial automorphism group. Such graphs are known as asymmetric graphs. Erdős and Rényi [ER63] showed that asymmetric graphs have at least 6 vertices and there are 8 asymmetric graphs on 6 vertices. Interestingly, all these 8 graphs indeed have irreducible characteristic polynomial, and their spectral radii are $> \lambda^*$.

Clearly, if λ is the spectral radius of a graph, then λ is a totally real algebraic integer, and λ is the largest among its conjugate elements. It would be interesting to study a complete set of necessary conditions for the spectral radius of a graph.

When λ is a totally real algebraic integer but not the largest among its conjugate elements, the spectral radius order $k(\lambda) = \infty$, and Conjecture B asserts that $N_\alpha(n) = n + O(1)$. This is indeed the case.

Proposition 23. *Given $\alpha \in (0, 1)$, if $\lambda := \frac{1-\alpha}{2\alpha}$ is a totally real algebraic integer, but λ is not the largest among its conjugate elements, then $n \leq N_\alpha(n) \leq n + 2$.*

Proof. We denote by $\lambda_{-i}(\cdot)$ and $\lambda_i(\cdot)$ respectively the i th smallest eigenvalue and the i th largest eigenvalue of a matrix. Let $\lambda' > \lambda$ be a conjugate element of λ . Let C be a spherical $\{\pm\alpha\}$ -code in \mathbb{R}^n . Let M be its Gram matrix, and A the adjacency matrix of the underlying graph G . We know that $M = (1 - \alpha)(I - A/\lambda) + \alpha J$.

Assume for the sake of contradiction that $\text{rank}(I - A/\lambda) \leq |C| - 2$, that is, λ is an eigenvalue of A with multiplicity ≥ 2 , then $1 - \lambda'/\lambda < 0$ is an eigenvalue of $I - A/\lambda$ with multiplicity ≥ 2 , hence $\lambda_{-2}(I - A/\lambda) < 0$. By Weyl's inequality, $\lambda_{-1}(M) \leq (1 - \alpha)\lambda_{-2}(I - A/\lambda) + \alpha\lambda_2(J) < 0$. This contradicts with the fact that M is positive semidefinite.

Therefore $\text{rank}(I - A/\lambda) \geq |C| - 1$ and so $n \geq \text{rank}(M) \geq \text{rank}(I - A/\lambda) - \text{rank}(J) \geq |C| - 2$. Together with Lemma 14, we have $n \leq N_\alpha(n) \leq n + 2$. \square

Lastly, we remark on a possible extension of our method. Our proof strategy would resolve Conjecture B provided the following connection between forbidden subgraphs characterization and multiplicity of the second largest eigenvalue.

Conjecture C. *For every $\lambda > 0$, there exist graphs G_1, G_2, \dots, G_n of spectral radius $> \lambda$ such that for every connected graph G that does not contain any of G_1, G_2, \dots, G_n as a subgraph, if λ is the second largest eigenvalue of G then the multiplicity of λ is $\leq v(G)/k(\lambda)$, where $v(G)$ is the order of G and $k(\lambda)$ is the spectral radius order of λ .*

In this direction, Woo and Neumaier [WN07] investigated the structure of graphs whose spectral radius is in $(2, 3/\sqrt{2}]$. In particular, such a graph is either an open quipu³, a closed quipu⁴ or a dagger⁵, for which we assert that the multiplicity of any eigenvalue larger than 2 is at most 2.

Acknowledgements

Thanks to Boris Bukh for introducing equiangular lines to the first author, and to Jun Su and Sebastian Cioabă for useful correspondence. We wish to express our deep appreciation to the referee for meticulous reading, and for pointing out a mistake in Theorem 1 and many other inaccuracies in an earlier version of the manuscript. All remaining errors are ours.

³An *open quipu* is a tree of maximum degree 3 such that all vertices of degree 3 lie on a path.

⁴A *closed quipu* is a connected graph of maximum degree 3 such that all vertices of degree 3 lie on a unique cycle.

⁵A *dagger* is A_n defined in Lemma 5.

References

- [BDKS18] Igor Balla, Felix Dräxler, Peter Keevash, and Benny Sudakov. Equiangular lines and spherical codes in Euclidean space. *Invent. Math.*, 211(1):179–212, 2018. [arXiv:math/1606.06620](#) [math.CO].
- [Bel97] Richard Bellman. *Introduction to matrix analysis*, volume 19 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1997. Reprint of the second (1970) edition, With a foreword by Gene Golub.
- [BN89] A. E. Brouwer and A. Neumaier. The graphs with spectral radius between 2 and $\sqrt{2 + \sqrt{5}}$. *Linear Algebra Appl.*, 114/115:273–276, 1989.
- [Buk16] Boris Bukh. Bounds on equiangular lines and on related spherical codes. *SIAM J. Discrete Math.*, 30(1):549–554, 2016. [arXiv:math/1508.00136](#) [math.CO].
- [CDG82] Dragoš Cvetković, Michael Doob, and Ivan Gutman. On graphs whose spectral radius does not exceed $(2 + \sqrt{5})^{1/2}$. *Ars Combin.*, 14:225–239, 1982.
- [dC00] D. de Caen. Large equiangular sets of lines in Euclidean space. *Electron. J. Combin.*, 7:Research Paper 55, 3, 2000.
- [ER63] P. Erdős and A. Rényi. Asymmetric graphs. *Acta Math. Acad. Sci. Hungar.*, 14:295–315, 1963.
- [Est92] Dennis R. Estes. Eigenvalues of symmetric integer matrices. *J. Number Theory*, 42(3):292–296, 1992.
- [Fri93] Joel Friedman. Some geometric aspects of graphs and their eigenfunctions. *Duke Math. J.*, 69(3):487–525, 1993.
- [Fro12] Ferdinand Georg Frobenius. Über Matrizen aus nicht negativen Elementen. *S.-B. Deutsch. Akad. Wiss. Berlin*, 1912:456–477, 1912.
- [GKMS16] Gary Greaves, Jacobus H. Koolen, Akihiro Munemasa, and Ferenc Szöllösi. Equiangular lines in Euclidean spaces. *J. Combin. Theory Ser. A*, 138:208–235, 2016. [arXiv:math/1403.2155](#) [math.CO].
- [GM81] C. D. Godsil and B. D. McKay. Spectral conditions for the reconstructibility of a graph. *J. Combin. Theory Ser. B*, 30(3):285–289, 1981.
- [GY18] Alexey Glazyrin and Wei-Hsuan Yu. Upper bounds for s -distance sets and equiangular lines. *Adv. Math.*, 330:810–833, 2018. [arXiv:math/1611.09479](#) [math.MG].

- [Hof72] Alan J. Hoffman. On limit points of spectral radii of non-negative symmetric integral matrices. pages 165–172. *Lecture Notes in Math.*, Vol. 303, 1972.
- [Hoo05] Shlomo Hoory. A lower bound on the spectral radius of the universal cover of a graph. *J. Combin. Theory Ser. B*, 93(1):33–43, 2005.
- [Jia19] Zilin Jiang. On spectral radii of unraveled balls. *J. Combin. Theory Ser. B*, 136:72–80, 2019. [arXiv:1710.06719](https://arxiv.org/abs/1710.06719) [math.CO].
- [JW15] Jonathan Jedwab and Amy Wiebe. Large sets of complex and real equiangular lines. *J. Combin. Theory Ser. A*, 134:98–102, 2015. [arXiv:math/1501.05395](https://arxiv.org/abs/math/1501.05395) [math.CO].
- [LP17] David A. Levin and Yuval Peres. Counting walks and graph homomorphisms via Markov chains and importance sampling. *Amer. Math. Monthly*, 124(7):637–641, 2017.
- [LPS88] A. Lubotzky, R. Phillips, and P. Sarnak. Ramanujan graphs. *Combinatorica*, 8(3):261–277, 1988.
- [LS73] P. W. H. Lemmens and J. J. Seidel. Equiangular lines. *J. Algebra*, 24:494–512, 1973.
- [Mar82] G. A. Margulis. Explicit constructions of graphs without short cycles and low density codes. *Combinatorica*, 2(1):71–78, 1982.
- [Mow71] Abbe Mowshowitz. Graphs, groups and matrices. pages 509–522, 1971.
- [Neu89] A. Neumaier. Graph representations, two-distance sets, and equiangular lines. *Linear Algebra Appl.*, 114/115:141–156, 1989.
- [Nil91] A. Nilli. On the second eigenvalue of a graph. *Discrete Math.*, 91(2):207–210, 1991.
- [Nil04] A. Nilli. Tight estimates for eigenvalues of regular graphs. *Electron. J. Combin.*, 11(1):Note 9, 4, 2004.
- [Per07] Oskar Perron. Zur Theorie der Matrices. *Math. Ann.*, 64(2):248–263, 1907.
- [She89] James B. Shearer. On the distribution of the maximum eigenvalue of graphs. *Linear Algebra Appl.*, 114/115:17–20, 1989.
- [Smi70] John H. Smith. Some properties of the spectrum of a graph. In *Combinatorial Structures and their Applications (Proc. Calgary Internat. Conf., Calgary, Alta., 1969)*, pages 403–406. Gordon and Breach, New York, 1970.
- [vLS66] J. H. van Lint and J. J. Seidel. Equilateral point sets in elliptic geometry. *Indag. Math.*, 28(3):335–348, 1966.

- [Wey12] Hermann Weyl. Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen (mit einer Anwendung auf die Theorie der Hohlraumstrahlung). *Math. Ann.*, 71(4):441–479, 1912.
- [WN07] Renee Woo and Arnold Neumaier. On graphs whose spectral radius is bounded by $\frac{3}{2}\sqrt{2}$. *Graphs Combin.*, 23(6):713–726, 2007.

A Proof of Lemma 5

Given a connected graph G , let A be its adjacency matrix. The Perron–Frobenius theorem asserts that the eigenvector $f: V(G) \rightarrow \mathbb{R}$ corresponding to the unique largest eigenvalue of A can be chosen so that all of its components are positive. As an eigenvector f' corresponding to any other eigenvalue is orthogonal to f , f' must have at least one negative component. Therefore to compute $\lambda_1(G)$ it suffices to demonstrate an eigenvector whose components are positive.

Proof of Lemma 5(b₁). The eigenvector that maps the leaves to 1 and the rest of the vertices to 2 gives $\lambda_1(B_{1,n,1}) = 2$. \square

Proof of Lemma 5(c). The constant eigenvector that maps every vertex to 1 gives $\lambda_1(C_n) = 2$. \square

Proof of Lemma 5(p). Let v_1, \dots, v_n be the path P_n . The eigenvector that maps v_i to $\sin(\frac{\pi i}{n+1})$ gives $\lambda_1(P_n) = 2 \cos(\frac{\pi}{n+1})$. \square

Proof of Lemma 5(s). The eigenvector that maps the leaves to 1 and the vertex of degree n to \sqrt{n} gives $\lambda_1(S_n) = \sqrt{n}$. \square

For the proof of other facts in Lemma 5, we shall use the following lemmas due to Hoffman.

Lemma 24 (Lemma 3.4 of Hoffman [Hof72]). *Let A_{-1} be a principal submatrix of order $n - 1$ of a symmetric matrix A_0 of order n with non-negative entries. Define A_{i+1} recursively by*


$$A_{i+1} = \begin{pmatrix} A_i & e_i^T \\ e_i & 0 \end{pmatrix}, \text{ where } e_i = \begin{pmatrix} 0 & 0 & \dots & 0 & 1 \end{pmatrix}.$$

Assume further that $\lim_{i \rightarrow \infty} \lambda_1(A_i) > 2$. Then $\lim_{i \rightarrow \infty} \lambda_1(A_i)$ is the largest positive root of

$$\left(\frac{x + \sqrt{x^2 - 4}}{2} \right) p_0(x) = p_{-1}(x), \tag{8}$$

where p_i is the characteristic polynomial of A_i for $i = -1, 0$.

Definition 3. Let G be a connected graph, and let v be a vertex of G . Denote (G, v, n) the graph obtained from G by appending a path of n vertices to G at v . Let G_1, G_2 be disjoint connected graphs, and let v_1, v_2 be vertices of G_1, G_2 respectively. Define (G_1, v_1, n, v_2, G_2) to be the graph obtained from G_1 and G_2 by joining them by a path of n vertices connecting v_1 and v_2 .



Remark 4. When we apply Lemma 24 to the adjacency matrix of a graph, we get the following interpretation. Let G be a connected graph, and let v be a vertex of G . Assume further that $\lambda_1(G, v, n) \geq 2$ for some n . Then $\lim \lambda_1(A_n)$ is the largest positive root of (8), where p_{-1}, p_0 are the characteristic polynomials of $G \setminus \{v\}$ and G respectively.

Lemma 25 (Proposition 4.2 of Hoffman [Hof72]). *Let G_1, G_2 be disjoint connected graphs, v_1, v_2 vertices of degree ≥ 2 of G_1, G_2 respectively. Then*

$$\lim_{n \rightarrow \infty} \lambda_1(G_1, v_1, n, v_2, G_2) = \max \left\{ \lim_{n \rightarrow \infty} \lambda_1(G_1, v_1, n), \lim_{n \rightarrow \infty} \lambda_1(G_2, v_2, n) \right\}.$$

Definition 4. Let e be an edge of a graph G . If there exists a path in G , x_1, x_2, \dots, x_k where x_{k-1} and x_k are the end vertices of e , and the degrees of x_1, x_2, \dots, x_{k-1} are respectively $1, 2, 2, \dots, 2$, then e is said to be on an *end path* of G .

Lemma 26 (Proposition 4.1 of Hoffman [Hof72]). *Let G be a connected graph with $\lambda_1(G) > 2$, $e = (x, y)$ an edge of G not on an end path of G . Let G_e^+ be the graph obtained from G by deleting edge e , and adding a vertex z adjacent to x and y only. Then $\lambda_1(G_e^+) < \lambda_1(G)$.*

The monotonicity of the spectral radii of each family of graphs in Lemma 5(a,e,f) follows immediately from the monotonicity of the spectral radii. The monotonicity in Lemma 5(b₂,d) follows from Lemma 26 and the facts that $\lambda_1(B_{m_1,0,m_2}) > \lambda_1(B_{1,0,1}) = 2$ for $(m_1, m_2) \neq (1, 1)$ and $\lambda_1(D_2) > 2$.

We are left to compute the limits.

Proof of Lemma 5(a). Note that $A_n = (S_3, v, n)$, where v is the vertex of degree 3 in S_3 . Note that $\lambda_1(A_1) = 2$. By Remark 4, $\lim \lambda_1(A_n)$ is the largest positive root of

$$\left(\frac{x + \sqrt{x^2 - 4}}{2} \right) x^2(x^2 - 3) = x^3,$$

which turns out to be $3/\sqrt{2}$. □

Proof of Lemma 5(f). Note that $F_n = (P_5, v, n)$, where v is the third vertex of P_5 . Observe that $\lambda_1(F_2) = 2$. By Remark 4, $\lim \lambda_1(F_n)$ is the largest positive root of

$$\left(\frac{x + \sqrt{x^2 - 4}}{2} \right) (x^5 - 4x^3 + 3x) = (x^2 - 1)^2,$$

which turns out to be λ^* . □

We need the characteristic polynomials of paths and cycles for Lemma 5(d, e). The readers are invited to derive them by reduction and induction.

Lemma 27. *Denote p_n and q_n the characteristic polynomials of P_n and C_n respectively. Then*

$$\begin{aligned} p_0(x) &= 1, & p_1(x) &= x, & p_n &= xp_{n-1}(x) - p_{n-2}(x), \\ q_{n+1}(x) &= p_{n+1}(x) - p_{n-1}(x) - 2, & & & & \text{for all } n \geq 2. \end{aligned}$$

Moreover, the recursions give

$$p_n(x) = \frac{\theta^n}{1 - \theta^{-2}} + \frac{\theta^{-n}}{1 - \theta^2}, \quad q_n(x) = \theta^n + \theta^{-n} - 2, \quad (9)$$

where $\theta = \theta(x) := \frac{x + \sqrt{x^2 - 4}}{2}$.

Proof of Lemma 5(e). For the $m = 1$ case, $\lambda_1(E_{1,n})$ is at least the average degree $2 - 2/(n + 3)$ and so $\lim \lambda_1(E_{1,n}) \geq 2$. On the other hand, assume for the sake of contradiction that $\lim \lambda_1(E_{1,n}) > 2$. By Remark 4, $\lim \lambda_1(E_{1,n})$ is the largest positive root of

$$\left(\frac{x + \sqrt{x^2 - 4}}{2} \right) x (x^2 - 2) = x^2,$$

which turns out to be 2 contradicting the assumption $\lim \lambda_1(E_{1,n}) > 2$.

For the $m \geq 2$ case, because $\lambda_1(E_{m,8}) \geq \lambda_1(E_{2,8}) = 2$ and $E_{m,n} = (P_{m+2}, v, n)$, where v is the second vertex of P_{m+2} , Remark 4 gives that $\lim_{n \rightarrow \infty} E_{m,n}$ is the largest positive root of

$$\theta p_{m+2}(x) = xp_m(x),$$

where θ and p_i are defined as in Lemma 27. From this and (9), using $x = \theta + 1/\theta$ and $z = \theta^2$, we seek the largest root z_m of

$$z^{m+1} = 1 + z + \dots + z^{m-1}.$$

By the definitions of β_m and α_m in Theorem 1, this proves that $z_m = \beta_m$ and $\lim_n \lambda_1(E_{m,n}) = \alpha_m$. \square

Proof of Lemma 5(b₂). Let v_1 and v_2 be the second vertices in P_{m_1+2} and P_{m_2+2} respectively. By Lemma 25,

$$\begin{aligned} \lim_{n \rightarrow \infty} \lambda_1(B_{m_1, n, m_2}) &= \max \left\{ \lim_{n \rightarrow \infty} \lambda_1(P_{m_1+2}, v_1, n), \lim_{n \rightarrow \infty} \lambda_1(P_{m_2+2}, v_2, n) \right\} \\ &= \max \left\{ \lim_{n \rightarrow \infty} \lambda_1(E_{m_1, n}), \lim_{n \rightarrow \infty} \lambda_1(E_{m_2, n}) \right\} = \max(\alpha_{m_1}, \alpha_{m_2}). \quad \square \end{aligned}$$

Proof of Lemma 5(d). Let M_n be the adjacency matrix of D_n and let r_n be its characteristic polynomial. By expanding the determinant of $xI - M_n$ along the row indexed by the leaf of D_n , one can obtain that $\lambda_1(D_n)$ is the largest root of

$$r_n(x) = xq_{n+1}(x) - p_n(x),$$

where p_n and q_{n+1} are defined as in Lemma 27. From this and (9), using $x = \theta + 1/\theta$ and $z = \theta^2$, we seek the largest root z_n of

$$(z^2 - z - 1) \left(1 - z^{-(n+1)}\right) = 2z^{-n/2} \left(z^{1/2} + z^{-1/2}\right).$$

As $z_n > 1$, we get $z_n^2 - z_n - 1 > 0$, hence $z_n > \phi$, where $\phi = \frac{1+\sqrt{5}}{2}$ is the golden ratio. As $n \rightarrow \infty$, the largest root z_n tends to ϕ , and so $\lim \lambda_1(D_n) = \phi^{1/2} + \phi^{-1/2} = \lambda^*$. \square

B Proof of Proposition 16

Fix $m \geq 2$ and let β be the largest root of the equation $x^{m+1} = 1 + x + \dots + x^{m-1}$. We shall prove that $\alpha := \beta^{1/2} + \beta^{-1/2}$ is an algebraic integer but not totally real. We need the following lemma to characterize the conjugate elements of α .

Lemma 28. *Suppose that the polynomial $p(x) \in \mathbb{Q}[x]$ of degree n has n complex roots, say $\gamma_1, \gamma_2, \dots, \gamma_n$, counted with multiplicity. If these roots are not the poles of a rational function $r(x) \in \mathbb{Q}(x)$, then $q(x) := \prod_{i=1}^n (x - r(\gamma_i))$ is a polynomial in $\mathbb{Q}[x]$.*

Proof. For every $d \in [n]$, let $e_d \in \mathbb{Z}[x_1, x_2, \dots, x_n]$ be the elementary symmetric polynomial of degree d in n variables. Note that $f_d(x_1, x_2, \dots, x_n) := e_d(r(x_1), r(x_2), \dots, r(x_n))$ is a symmetric rational function with rational coefficients. Thus by the fundamental theorem of symmetric functions, f_d can be written as a rational function of the elementary symmetric functions e_1, e_2, \dots, e_n with rational coefficients. Note that the coefficient of x^d in the polynomial $q(x)$ is precisely $(-1)^{n-d} f_{n-d}(\gamma_1, \gamma_2, \dots, \gamma_n)$. Since $p(x) \in \mathbb{Q}[x]$, Vieta's formulas tell us $e_d(\gamma_1, \gamma_2, \dots, \gamma_n)$ is a rational for all $d \in [n]$, hence the coefficients of $q(x)$ are rational. \square

Proof of Proposition 16. We first prove the existence and the uniqueness of the positive root of $p(x) := x^{m+1} - (1 + x + \dots + x^{m-1})$. Since $p(1) = 1 - m < 0$ and $p(\infty) = \infty$, by the intermediate value theorem, p has a positive root. Suppose for a moment that β is just a positive root of p . Then $\beta^{m+1} = 1 + \beta + \dots + \beta^{m-1} > 1$, and so $\beta > 1$, hence $\beta^{m+1} = 1 + \beta + \dots + \beta^{m-1} > m$. Notice that

$$\beta^{m+1} = 1 + \beta + \dots + \beta^{m-1} = \frac{\beta^m - 1}{\beta - 1} \implies \beta - 1 = \frac{1}{\beta} - \frac{1}{\beta^{m+1}}. \quad (10)$$

This means that β is a zero of the function $f(x) := x - 1 - \frac{1}{x} + \frac{1}{x^{m+1}}$ in $(m^{1/(m+1)}, \infty)$. Note that $f'(x) = 1 + \frac{1}{x^2} - \frac{m+1}{x^{m+2}}$, and when $x > m^{1/(m+1)}$, the derivative

$$f'(x) \geq \frac{2}{x} - \frac{m+1}{x^{m+2}} = \frac{2x^{m+1} - (m+1)}{x^{m+2}} \geq \frac{2m - (m+1)}{x^{m+2}} > 0,$$

which shows the uniqueness of the positive root of p .

Next we show that α is an algebraic integer. Let $\beta = \beta_0, \beta_1, \dots, \beta_m$ be the $m+1$ complex roots of p counted with multiplicity. Let $\gamma_i := \sqrt{\beta_i}$ be the principal square root of β_i for $i \in$

$\{0, 1, \dots, m\}$. Clearly, $\pm\gamma_0, \pm\gamma_1, \dots, \pm\gamma_m$ are the roots of the monic polynomial $p(x^2) \in \mathbb{Z}[x]$, and $\pm 1/\gamma_0, \pm 1/\gamma_1, \dots, \pm 1/\gamma_m$ are the roots of the monic polynomial $-x^{2(m+1)}p(1/x^2) \in \mathbb{Z}[x]$. Therefore both γ_0 and $1/\gamma_0$ are algebraic integers, and so is $\alpha = \gamma_0 + 1/\gamma_0$.

Suppose α' is a conjugate element of α . Applying Lemma 28 to $\pm\gamma_0, \dots, \pm\gamma_m$ and the rational function $x \mapsto x + 1/x$, we know that α' must be of the form $\pm(\gamma_i + 1/\gamma_i)$, for some $i \in \{0, 1, \dots, m\}$. Assume for the sake of contradiction that $\alpha' = \pm(\gamma_i + 1/\gamma_i)$ is real for some $i \neq 0$. Without loss of generality, we may assume that $i = 1$. We can solve the quadratic equation $\gamma_1^2 \mp \alpha'\gamma_1 + 1 = 0$ and get $\gamma_1 = \frac{\pm\alpha' \pm \sqrt{\alpha'^2 - 4}}{2}$, where the plus-minus signs are independent. As $\beta = \beta_0$ is the only positive root of p , $\gamma_1 = \sqrt{\beta_1}$ is not real. Since $\alpha' \in \mathbb{R}$ but $\gamma_1 \notin \mathbb{R}$, it must be the case that $-2 < \alpha' < 2$ and $\gamma_1 = \frac{\pm\alpha' \pm i\sqrt{4 - \alpha'^2}}{2}$. Therefore we have

$$|\beta_1| = |\gamma_1|^2 = \left(\frac{\alpha'}{2}\right)^2 + \left(\frac{\sqrt{4 - \alpha'^2}}{2}\right)^2 = 1.$$

Let θ be the argument of β_1 , that is $\beta_1 = \cos \theta + i \sin \theta$. Since β_1 is a root of p , (10) holds for β_1 , that is, $\beta_1 - 1 = \beta_1^{-1} - \beta_1^{-(m+1)}$, or $\beta_1^{-(m+1)} = 1 - (\beta_1 - \beta_1^{-1}) = 1 - 2i \sin \theta$ after rearrangement. Using $|\beta_1| = 1$, we get $\sin \theta = 0$, hence $\beta_1 = \pm 1$. As $p(1) < 0$, $\beta_1 = -1$ is the only possibility, in which case $\alpha' = \pm(\sqrt{\beta_1} + 1/\sqrt{\beta_1}) = 0$ and it cannot be a conjugate element of α .

The contradiction shows that any real conjugate element of α is of the form $\pm(\gamma_0 + 1/\gamma_0)$, that is, $\pm\alpha$. If α were a totally algebraic integer, its degree would be either 1 or 2. In the former case, α would be an integer, which contradicts with $2 < \alpha < \lambda^* < 3$. In the latter case, α^2 would be an integer, which contradicts with $2 < \alpha < \lambda^* < \sqrt{5}$. \square