

Harmonic Index of Total Graphs of Some Graphs

V. Anandkumar^{1*}, Radha Rajamani Iyer²

Department of Mathematics,
EASA College of Engineering and Technology,
Coimbatore -641 112,INDIA.
Department of Mathematics,
Amrita Vishwa Vidyapeetham,
Coimbatore -641 112,INDIA.

Abstract

Let $G = G(V, E)$ be a graph with $|V|$ vertices and $|E|$ edges and total graph, $T(G)$ is obtained from G . In This paper we have study the Harmonic index of total graph for standards graphs, bipartite graph of particular type, regular graph, Grid graph $G_{m,n}$ and Complete Binary tree.

2010 Mathematics Subject Classification: Primary 92E10, Secondary 94C15

Keywords:Total graph, Harmonic Index, Grid graph, Binary tree.

1.Introduction

Let $G(V, E)$ be a simple connected undirected graph with $|V| = n$ vertices, $|E| = m$ edges, $uv \in E(G)$ if u and v are adjacent in G . The *Randić* index of graph G is denoted by $R(G)$ and it is defined as $R(G) = \sum_{uv \in E(G)} (d_u d_v)^{\frac{-1}{2}}$ [11], where d_u is the degree of the vertex u in G . Few related variants of the *Randić* index has been introduced in last few years. Sum-connectivity index is one among them introduced by Zhou and Trinajstić in [14,15] and it is defined as $X(G) = \sum_{uv \in E(G)} (d_u + d_v)^{\frac{-1}{2}}$ [11]. $X_\alpha(G) = \sum_{uv \in E(G)} (d(u) + d(v))^\alpha$, where α is the real number. It has been found that $X_\alpha(G)$ and $R(G)$ correlate well between themselves and with Π -electron energy of benzenoid hydrocarbons. In this paper we study the another related variant of the index namely the *Randić* Harmonic index of a graph G and is denoted by $H(G)$ and defined as $H(G) = \sum_{uv \in E(G)} \frac{2}{d(u)+d(v)}$.

This Index was first appeared in [5]. Zhong found the minimum and maximum values of the harmonic index for simple connected graph and trees, and characterized the corresponding extremal graph.

2.Harmonic Index of Total Graphs of Some Graphs

For every graph G there exist a graph called the *line-graph* of G and is denoted by $L(G)$, whose points correspond in a one-to-one manner with the lines of G in such a way that two points in the $L(G)$ are adjacent if and only if the corresponding lines are adjacent in G . Vertices and edges are elements of a graph G . For any graph G , *total graph*, denoted by $T(G)$, is obtained from G whose

vertex set $V(T(G)) = V(G) \cup E(G)$ and two vertices are adjacent if and only if the corresponding elements are adjacent or incident in G . Total graph is introduced by Mehdi Behzad and Gary Chartrand in the year 1966. They have presented the characterization of graphs hamiltonian total graphs and of graph whose total graphs are Eulerian.

Observation

If v' is the point in the total graph of G corresponding to point v of G , then the degree of v' , denoted by $\deg v'$, equals $2 \deg v$. Where $x = v_1v_2$ is a line of G and x' is the corresponding point of $T(G)$, then $\deg x' = \deg v_1 + \deg v_2$.

In this paper Exact values of Harmonic index of total graph of standard graphs such as Path, Cycle, Gird Graph, Bipartite graph of a particular type and complete binary tree are obtained.

Proposition 2.1. For any graph P_n of order $n \geq 4$, $H[T(P_n)] = n - \frac{269}{420}$.

Proposition 2.2. For any star graph S_n , $H[T(S_n)] = (n - 1)[\frac{2}{n+2} + \frac{3(n+1)}{2(3n-1)}]$.

Theorem 2.3. Let G be a graph with $|V| = n$ and $|E| = m$ then

$$H(T(G)) = \frac{1}{2}H(G) + \sum_{\substack{uv \in E(G) \\ vw \in E(G)}} \frac{2}{d(u)+2d(v)+d(w)} + 8 \sum_{uv \in E(G)} \frac{[d(u)+d(v)]}{3(d(u)+d(v))^2+4d(u)d(v)}. \text{ proof: Total}$$

graph of a graph G has $|E(T(G))| = 3m + \frac{1}{2} \sum_{v_i} d(v_i)^2 - m$

$$= 2m + \frac{1}{2} \sum_{v_i} d(v_i)^2$$

each edge belongs any one of the following category.

(i) edges between u' and v' , where u' and v' are vertices corresponding to the vertices u and v in G .

(ii) edges between X' and Y' , where X' and Y' are vertices corresponding to the edges X and Y in G .

(iii) edges between X' and u' , where X' and u' are vertices corresponding to the edge X and the vertex u respectively, $X = uv \in E(G)$.

$$\begin{aligned} \therefore H[T(G)] &= \sum_{u'v' \in E(T(G))} \frac{2m + \frac{1}{2} \sum_{v_i \in G} d(v_i)^2}{d(u')+d(v')} \\ &= \sum_{u'v' \in E(T(G))} \frac{2}{d(u')+d(v')} + \sum_{X'Y' \in E(T(G))} \frac{\frac{1}{2} \sum_{v_i \in G} d(v_i)^2 - m}{d(X')+d(Y')} + \sum_{X'u' \in E(T(G))} \frac{2m}{d(X')+d(u')}. \\ &= A + B + C. \end{aligned}$$

$$\begin{aligned} \text{where } A &= \sum_{u'v' \in E(T(G))} \frac{2}{d(u')+d(v')} = \sum_{uv \in E(G)} \frac{2}{2(d(u)+d(v))} \\ &= \frac{1}{2}H(G). \end{aligned} \tag{1}$$

$$\begin{aligned}
\text{where } B &= \frac{1}{2} \sum_{v_i \in G}^{n} d(v_i^2) - m \\
&= \sum_{X'Y' \in E(T(G))} \frac{2}{d(X') + d(Y')} = \frac{1}{2} M_1(G) - m \sum_{XY \in E(L(G))} \frac{2}{d(X) + d(Y) + 4}. \\
&= \frac{1}{2} M_1(G) - m \sum_{\substack{uv \in E(G) \\ vw \in E(G)}} \frac{2}{d(u) + 2d(v) + d(w)}. \tag{2}
\end{aligned}$$

$$\begin{aligned}
\text{where } C &= \sum_{X'u' \in E(T(G))}^{2m} \frac{2}{d(X') + d(u')}. \\
&= \sum_{X=uv \in E(G)}^{2m} \frac{2}{2d(u) + d(X) + 2} \\
&= \sum_{uv \in E(G)}^m \left[\frac{2}{2d(u) + d(u) + d(v)} + \frac{2}{d(u) + 2d(v) + d(v)} \right] \\
&= \sum_{uv \in E(G)}^m \left[\frac{2}{3d(u) + d(v)} + \frac{2}{d(u) + 3d(v)} \right] \\
&= \sum_{uv \in E(G)}^m \left[\frac{8d(u) + 8d(v)}{3d(u)^2 + 9d(u)d(v) + d(u)d(v) + 3d(v)^2} \right] \\
&= \sum_{uv \in E(G)}^m \left[\frac{8[d(u) + d(v)]}{3d(u)^2 + 10d(u)d(v) + 3d(v)^2} \right] \\
&= 8 \sum_{uv \in E(G)}^m \left[\frac{[d(u) + d(v)]}{3(d(u) + d(v))^2 + 4d(u)d(v)} \right]. \tag{3}
\end{aligned}$$

From (1),(2) and (3),

$$H(T(G)) = \frac{1}{2} H(G) + \sum_{\substack{uv \in E(G) \\ vw \in E(G)}}^{M_1(G) - m} \frac{2}{d(u) + 2d(v) + d(w)} + 8 \sum_{uv \in E(G)}^m \left[\frac{[d(u) + d(v)]}{3(d(u) + d(v))^2 + 4d(u)d(v)} \right].$$

Theorem 2.4. If G is regular graph then $H[T(G)] = \frac{n(k+2)}{4}$.

proof: If G is a regular then $T(G)$ is also regular and it has $\frac{n}{2}(2+k)$ vertices and degree of all the vertices are the same. Then $|E(T(G))| = \frac{nk}{2}(k+2)$.

$$\text{Hence, } H[T(G)] = \sum_{u'v' \in E(T(G))}^{\frac{nk}{2}(k+2)} \frac{2}{d(u') + d(v')}$$

here degree of every vertex in $T(G)$ is $2k$.

therefore,

$$= \frac{nk}{2}(k+2) \frac{2}{2k+2k}$$

$$H[T(G)] = \frac{n(k+2)}{4}.$$

Corollary 2.4.1. Let C_n be a cycle with n vertices then $H[T(C_n)] = n$.

proof: Total graph $T(C_n)$ is the 4-regular graph of the size $4n$.

$$\begin{aligned}
\therefore H[T(C_n)] &= \sum_{u'v' \in E[T(C_n)]}^{4n} \frac{2}{d(u') + d(v')} \\
&= 4n \frac{2}{4+4}
\end{aligned}$$

$$H[T(C_n)] = n.$$

Corollary 2.4.2. Let K_n be a complete graph then $H[T(K_n)] = \frac{n(n+1)}{4}$.

proof: $T(K_n)$ is the $(2(n-1))$ -regular graph of the size $\frac{n(n-1)(n+1)}{2}$.

$$\begin{aligned} \therefore H[T(K_n)] &= \sum_{uv \in E[T(K_n)]} \frac{2}{d(u)+d(v)} \\ &= \frac{n(n-1)(n+1)}{2} \left(\frac{2}{2(n-1)+2(n-1)} \right) \end{aligned}$$

$$H[T(K_n)] = \frac{n(n+1)}{4}.$$

Theorem 2.5. Let graph G be a bipartite graph with bipartition X and Y for all i . $\deg v_i = \begin{cases} k_1, & \text{if } v_i \in X \\ k_2, & \text{if } v_i \in Y \end{cases}$,

$$\text{then } H[T(G)] = \frac{mk_1^2+nk_2^2}{2(k_1+k_2)} + \frac{4(mk_1+nk_2)(k_1+k_2)}{(3k_1+k_2)(k_1+3k_2)}.$$

proof: Let G be a bipartite graph with partition $|X| = m$ and $|Y| = n$.

number of edges in G is $\frac{mk_1+nk_2}{2}$

In the Total graph of G , $|V(T(G))| = m + n + \frac{mk_1+nk_2}{2}$

$$|E(T(G))| = \frac{mk_1+nk_2}{2} + \frac{nk_2(k_2-1)}{2} + \frac{mk_1(k_1-1)}{2} + 2\left(\frac{mk_1+nk_2}{2}\right)$$

$$|E(T(G))| = \frac{1}{2}[nk_2(k_2+2) + mk_1(k_1+2)]$$

$$\begin{aligned} H[T(G)] &= \sum_{u'v' \in E[T(G)]} \frac{2}{d(u')+d(v')} \\ &= \frac{mk_1+nk_2}{2} \left(\frac{2}{2(k_1+k_2)} \right) + \frac{mk_1(k_1-1)}{2} \left(\frac{2}{2(k_1+k_2)} \right) + \frac{nk_2(k_2-1)}{2} \left(\frac{2}{2(k_1+k_2)} \right) + \\ &\quad \frac{mk_1+nk_2}{2} \left(\frac{2}{3k_1+k_2} \right) + \frac{mk_1+nk_2}{2} \left(\frac{2}{k_1+3k_2} \right) \\ &= \frac{mk_1^2+nk_2^2}{2(k_1+k_2)} + (mk_1+nk_2) \left(\frac{1}{3k_1+k_2} + \frac{1}{k_1+3k_2} \right) \\ &= \frac{mk_1^2+nk_2^2}{2(k_1+k_2)} + (mk_1+nk_2) \left(\frac{4k_1+4k_2}{(3k_1+k_2)(k_1+3k_2)} \right) \end{aligned}$$

$$\therefore H[T(G)] = \frac{mk_1^2+nk_2^2}{2(k_1+k_2)} + \frac{4(mk_1+nk_2)(k_1+k_2)}{(3k_1+k_2)(k_1+3k_2)}.$$

Corollary 2.5.1. let $K_{m,n}$ be a complete bipartite graph then

$$H[T(K_{m,n})] = \frac{5mn}{6(m+n)^2+8mn}.$$

proof: $H[T(K_{m,n})] = mn \frac{2}{2n+2m} + \frac{mn(m+n-2)}{2} \frac{2}{(m+n+m+n)} + mn \left(\frac{2}{2n+m+n} \right)$

$$\begin{aligned} &+ mn \frac{2}{2m+m+n} \\ &= \frac{mn}{m+n} + \frac{mn(m+n-2)}{2(m+n)} + \frac{2mn}{2n+m+n} + \frac{2mn}{2m+m+n} \end{aligned}$$

$$\begin{aligned}
&= \frac{2mn+mn(m+n)-2mn}{2(m+n)} + \frac{2mn}{3n+m} + \frac{2mn}{3m+n} \\
&= \frac{mn}{2} + \frac{2mn}{3n+m} + \frac{2mn}{3m+n}
\end{aligned}$$

$$H[T(K_{m,n})] = \frac{5mn}{6(m+n)^2+8mn}.$$

Theorem 2.6. Let G be a gird graph then

$$H[T(G_{m,n})] = \frac{M_1(G)+mn}{2} + (m+n)\frac{99}{260} - \frac{58237}{12870} \text{ where } m, n \geq 5.$$

proof: The total graph of gird graph has

$$\begin{aligned}
|V| &= 3mn - m - n \text{ and} \\
|E| &= \frac{1}{2}M_1(G) + 4mn - 2(m+n).
\end{aligned}$$

The degree pairs of the number of edges 8, 8, 4, 16, 8, $4(2m+2n-13)$, $4(3m+3n-16)$, $2(m+n-4)$, $4, 8(m+n-5)$, $\frac{1}{2}M_1(G) + 4mn - 26(m+n) + 84$ in the $T(G_{m,n})$ are (4,5), (4,6), (5,5), (5,6), (5,7), (6,6), (6,7), (6,8), (7,7), (7,8), (8,8) respectively.

$$\begin{aligned}
\therefore H[T(G_{m,n})] &= 8\left(\frac{2}{4+5}\right) + 8\left(\frac{2}{4+6}\right) + 4\left(\frac{2}{5+5}\right) + 16\left(\frac{2}{5+6}\right) + 8\left(\frac{2}{5+7}\right) + 4(2m+2n-13)\left(\frac{2}{6+6}\right) + 2(3m+3n-16)\left(\frac{2}{6+7}\right) \\
&\quad + 2(m+n-4)\left(\frac{2}{6+8}\right) + 4\left(\frac{2}{7+7}\right) + 8(m+n-5)\left(\frac{2}{7+8}\right) + \left(\frac{1}{2}M_1(G) + 4mn - 26(m+n) + 84\right)\left(\frac{2}{8+8}\right) \\
&= \frac{1}{16}M_1(G) + \frac{mn}{2} + \left(\frac{16}{9} + \frac{12}{5} + \frac{32}{11} - \frac{38}{3} - \frac{64}{13} - \frac{4}{7} + \frac{21}{2}\right) + (m+n)\left(\frac{4}{3} + \frac{16}{13} + \frac{16}{15} - \frac{52}{16}\right) \\
&= \frac{1}{16}M_1(G) + \frac{mn}{2} + \frac{1188}{3120}(m+n) - \left(\frac{287539}{90090}\right) \\
H[T(G_{m,n})] &= \frac{1}{16}M_1(G) + \frac{mn}{2} + \frac{99}{260}(m+n) - \left(\frac{41077}{12870}\right).
\end{aligned}$$

Theorem 2.7. For any complete binary tree T of level $l \geq 3$, $H(T(T)) = 2^l\left(\frac{223}{120}\right) - \frac{2274}{1485}$.

proof: Let T be a binary tree with level l . number of vertices and number of edges in the total graph $T(T)$ are $2^{l+2} - 3$ and $9 \cdot 2^l - 11$ respectively.

ie., $|V[T(T)]| = 2^{l+2} - 3$ and $|E[T(T)]| = 9 \cdot 2^l - 11$

the number of edges, $2^l, 2^l, 2^{l-1}, 2, (2^{l+1} + 2), 1, 6, (9 \cdot 2^{l-1} - 22)$ have the degree pairs (ie., end degree vertices) in the total graph are (2,4), (2,6), (4,4), (4,5), (4,6), (5,5), (5,6) and (6,6) respectively.

$$\begin{aligned}
H[T(T)] &= \sum_{u'v' \in E(T(G))}^{9 \cdot 2^l - 11} \frac{2}{d(u')+d(v')} \\
&= 2^l\left(\frac{2}{2+4}\right) + 2^l\left(\frac{2}{2+6}\right) + 2^{l-1}\left(\frac{2}{4+4}\right) + 2\left(\frac{2}{4+5}\right) + (2^{l+1} + 2)\left(\frac{2}{4+6}\right) + \\
&\quad \left(\frac{2}{5+5}\right) + 6\left(\frac{2}{5+6}\right) + (9 \cdot 2^{l-1} - 22)\left(\frac{2}{6+6}\right) \\
&= 2^l\left(\frac{2}{6}\right) + 2^l\left(\frac{2}{8}\right) + 2^{l-1}\left(\frac{2}{8}\right) + 2\left(\frac{2}{9}\right) + 2^{l+1}\left(\frac{2}{10}\right) + 2\left(\frac{2}{10}\right) + \frac{2}{10} + 6\left(\frac{2}{11}\right) \\
&\quad + 9 \cdot 2^{l-1}\left(\frac{2}{12}\right) - 22\left(\frac{2}{12}\right) \\
&= 2^l\left(\frac{2}{6} + \frac{2}{8} + \frac{2}{16} + \frac{4}{10} + \frac{18}{24}\right) + 4\left(\frac{1}{9} + \frac{1}{10} + \frac{1}{20} + \frac{3}{11} - \frac{11}{12}\right) \\
&= 2^{l-1}\left(\frac{40+30+15+48+90}{240}\right) - \frac{2274}{1485} \\
&= 2^{l+1}\left(\frac{223}{240}\right) - \frac{2274}{1485}
\end{aligned}$$

$$\therefore H(T(T)) = 2^l\left(\frac{223}{120}\right) - \frac{2274}{1485}.$$

Acknowledgement. The authors are thankful to the anonymous referees for their useful com-

ments.

References

1. G.H. Fath-Tabar, A.Hamzeh, and S.Hosseini-Zadeh, GA_2 index of some graph operation, *Filomat*, 24(1):21-28, 2010.
2. Xinli Xu, Relationships between Harmonic index and other topological indices, *applied mathematical Sciences*, 6(41):2013-2018, 2012.
3. Ismael G.Yero and Juan A.Rodriguez-Velazquez, On the Randic index of coronoid product graph, *International Scholarly Research notices*, 2011.
4. M.Randic, On characterization of molecular branching, *J.Am.Chem.Soc.*97(1975)6609-6615.
5. L.Pogliani, From molecular connectivity indices to semi empirical connectivity terms: recent trends in graph theoretical descriptors, *Chem.Rev.*100(2000)3827-3858.
6. I. Gutman, B.Furtula(Eds.), *Recent Results in the Theory of Randic Index*, Univ.Kragujevac,2008.
7. B.Zhou, N.Trinajstić, On general sum-connectivity index, *J.Math.Chem.*47(2010)210-218.
8. B.Lucic, N.Trinajstić, B.Zhou, Comparison between the sum-connectivity index and product-connectivity index for benzenoid hydrocarbons, *Chem.Phys.Lett.*475(2009)146-148.
9. Z.Du, B.Zhou, N.Trinajstić, On the general sum-connectivity index of trees, *appl.Math.Lett.*24(2011)402-405.
10. Lingping Zhong, The harmonic index for graphs, *applied Mathematics Letters* 25(2012)561-566.
11. Hanyuan Deng, S.Balachandran, S.K.Ayyaswamy, Y.B.VenkataKrishnan, On the harmonic index and the chromatic number of a graph, *Discrete applied Mathematics* 161(2013)2740-2744.
12. Z.Du, B.Zhou, On sum-connectivity index of bicyclic graphs, *Bull.Math.Sci.Soc.*(2)35(1)(2012)101-117.
13. Z.Du, B.Zhou, N.Trinajstić, minimum sum-connectivity indices of trees and unicyclic graphs of a given matching number, *J.Math.Chem.*47(2010)842-855.
14. R.Xing, B.Zhou, N.Trinajstić, sum-connectivity index of molecular trees, *J.Math.Chem.*48(2010)583-591.
15. L.Zhong, The harmonic index for graphs, *appl.Math.Lett.*25(2012)561-566.
16. B.Shwetha Shetty, V.Lokesh and P.S.Ranjini, On the harmonic index of graph operations, *Transactions on combinatorics*,4(4)(2015),5-14.