

LOCAL LARGE DEVIATIONS FOR EMPIRICAL LOCALITY MEASURE OF TYPED RANDOM GRAPH MODELS

BY KWABENA DOKU-AMPONSAH

University of Ghana

Abstract. In this article, we prove a local large deviation principle (LLDP) for the empirical locality measure of typed random networks on n nodes conditioned to have a given *empirical type measure* and *empirical link measure*. From the LLDP, we deduce a full large deviation principle for the typed random graph, and the classical Erdos-Renyi graphs, where $nc/2$ links are inserted at random among n nodes. No topological restrictions are required for these results.

1. INTRODUCTION AND BACKGROUND

1.1 Introduction. The Erdos-Renyi graph $\mathcal{G}(n, nc/2)$ is the simplest imaginary random, which arises by taking n nodes and inserting a fixed number $nc/2$ links at random among the n nodes. See, Van Hofstad [10]. Some large deviation principles for this random graph have been found. See, Bordenave & Caputo [1], Doku-Amponsah [5], and Doku-Amponsah and Moerters [8]

Doku-Amponsah and Moerters [8] provided LDPs for the near-critical or sparse typed random graphs with this model as a special case. Bordenave and Caputo [1] obtained large deviation principle for the empirical neighbourhood measure of the model $\mathcal{G}(n, nc/2)$. Large deviation for the empirical distribution was presented in Doku-Amponsah [5] using the method of types and the coupling argument presented by Boucheron et. al [2].

In this paper we present a Local Large deviation principle for the empirical locality measure of the conditional typed random graph model. See Bakhtin [3], Doku-Amponsah [6], Doku-Amponsah [7] for similar results for the empirical measure of iid random variables, the empirical offspring measure of multitype Galton-Watson trees and the empirical locality measure of the sparse typed random graphs, in which links appear in the graph with fix probability depending on the type of the links.

To state our main results in Section 2 we need some notions, concepts and notations from [8] and [4] in the background section below.

1.2 Background. Denote by \mathcal{Z} a finite alphabet or type set. Let $[n]$ be a fixed set of n nodes, say $[n] = \{1, \dots, n\}$. Denote by \mathcal{G}_n the set of all (simple) graphs with node set $[n] = \{1, \dots, n\}$ and link set \mathcal{E} . By $\mathcal{L}(\mathcal{Z})$ we denote the set of all probability distributions on \mathcal{Z} , $\tilde{\mathcal{L}}(\mathcal{Z})$ we denote the set of all finite distributions on \mathcal{Z} , and by $\mathcal{M}(\mathcal{Z})$ we denote the set of counting measures on \mathcal{Z} .

For any typed graph Z (with n nodes), recall from [5] the definitions of the *empirical type distribution* $P^1 \in \mathcal{L}(\mathcal{Z})$, the *empirical link distribution* $P^2 \in \tilde{\mathcal{L}}(\mathcal{Z} \times \mathcal{Z})$ and the *empirical locality distribution*

Mathematics Subject Classification : 94A15, 94A24, 60F10, 05C80

Keywords: Local large deviation, conditional typed random graphs, relative entropy, method of types, coupling, classical Erdos Renyi graph

Address: Statistics Department, University of Ghana, Box LG 115, Legon, Ghana. *E-mail*: kdoku@ug.edu.gh.

$P \in \mathcal{L}(\mathcal{Z} \times \mathcal{N}(\mathcal{Z}))$. For every $p \in \mathcal{L}(\mathcal{Z} \times \mathcal{M}(\mathcal{Z}))$ let p_1, p_2 be the \mathcal{Z} -marginal, respectively the $\mathcal{M}(\mathcal{Z})$ -marginal, of the measure p . Moreover, we define a measure $\langle p(\cdot, e), e(\cdot) \rangle \in \tilde{\mathcal{L}}(\mathcal{Z} \times \mathcal{Z})$ by

$$\langle p(\cdot, e), e(\cdot) \rangle(a, b) := \sum_{e \in \mathcal{M}(\mathcal{Z})} p(a, e)e(b), \quad \text{for } a, b \in \mathcal{Z}.$$

Define the function $\Psi: \mathcal{L}(\mathcal{Z} \times \mathcal{M}(\mathcal{Z})) \rightarrow \mathcal{L}(\mathcal{Z}) \times \tilde{\mathcal{L}}(\mathcal{Z} \times \mathcal{Z})$ by $\Psi(p) = (p_1, \langle p(\cdot, e), e(\cdot) \rangle)$. Note that $\Psi(P) = (P^1, P^2)$, and if these quantities are defined as empirical locality, type, and link measures of a type graph.

The remaining part of the paper is arranged in the following way: Section 2 starts with statement of the main Theorems of the paper; Theorem 2.1 and 2.2 and Corollary 2.3. This results are proved in Section 3. See subsections 3.1, 3.2 and 3.3.

2. STATEMENT OF MAIN RESULTS

To present the LDP, for a pair of distributions $(\pi, p) \in \tilde{\mathcal{L}}(\mathcal{Z} \times \mathcal{Z}) \times \mathcal{L}(\mathcal{Z} \times \mathcal{M}(\mathcal{Z}))$ we recall from [8] the concept of *sub-consistency* and *consistency* of measures. We shall throughout this section assume that $\eta(a) > 0$ for all $a \in \mathcal{Z}$. We write $\langle f, \sigma \rangle := \sum_{y \in \mathcal{Y}} \sigma(y)f(y)$ and denote by

$$Q_{(\eta_n, \pi_n)}(z) = \mathbb{Q}_{(\eta_n, \pi_n)}\{Z = z\} = \mathbb{Q}\{Z = z | \Psi(P) = (\eta_n, \pi_n)\}$$

the distribution of the typed random graph y conditioned to have typed law and link law (η_n, π_n) respectively. We define the rate function $J_{(\eta, \pi)}$ by

$$J_{(\eta, \pi)}(p) = \begin{cases} H(p \| q) & \text{if } (\pi, p) \text{ is sub-consistent and } p_1 = \eta \\ \infty & \text{otherwise.} \end{cases} \quad (2.1)$$

where

$$q(a, e) = \eta(a) \prod_{b \in \mathcal{Z}} \frac{e^{-\pi(a, b)/\eta(a)} [\pi(a, b)/\eta(a)]^{e(b)}}{e(b)!}, \quad \text{for } e \in \mathcal{M}(\mathcal{Z}).$$

The first results, see Theorem 2.1 below is the local large deviation principle for typed random graphs.

Theorem 2.1 (LLDP). *Suppose the sequence (η_n, π_n) converges to $(\pi, \eta) \in \mathcal{L}(\mathcal{Z}) \times \mathcal{L}_*(\mathcal{Z} \times \mathcal{Z})$. Let $z = ((z(i), i \in [n]), \mathcal{E})$ be coloured random graph conditioned on the event $\{\Psi(P) = (\eta_n, \pi_n)\}$. Then, we have*

- (i) *for any functional $p \in \mathcal{L}(\mathcal{Z} \times \mathcal{M}(\mathcal{Z}))$ and a number $\varepsilon > 0$, there exists a weak neighborhood B_p such that*

$$Q_{(\eta_n, \pi_n)}\left\{z \in \mathcal{G}([n], (\eta_n, \pi_n)) \mid P_z \in B_p\right\} \leq e^{-nJ_{(\pi, \eta)}(p) - n\varepsilon + o(n)}.$$

- (ii) *for any $p \in \mathcal{L}_*(\mathcal{Z} \times \mathcal{Z}_*)$, a number $\varepsilon > 0$ and a fine neighborhood B_p we have the asymptotic estimate:*

$$Q_{(\eta_n, \pi_n)}\left\{z \in \mathcal{G}([n], (\eta_n, \pi_n)) \mid P_z \in B_p\right\} \geq e^{-nJ_{(\pi, \eta)}(p) + n\varepsilon - o(n)}.$$

We state in Theorem 2.2 the full LDP for the typed random graph.

Theorem 2.2 (LDP). *Suppose the sequence (η_n, π_n) converges to $(\pi, \eta) \in \mathcal{L}(\mathcal{Z}) \times \mathcal{L}_*(\mathcal{Z} \times \mathcal{Z})$. Let $z = \left((z(i), i \in [n]), \mathcal{E} \right)$ be coloured random graph conditioned on the event $\left\{ \Psi(P) = (\eta_n, \pi_n) \right\}$.*

(i) *Let F be open subset of $\mathcal{L}(\mathcal{Z} \times \mathcal{Z}_*)$. Then we have*

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log Q_{(\eta_n, \pi_n)} \left\{ z \in \mathcal{G}([n], (\eta_n, \pi_n)) \mid P_z \in F \right\} \leq - \inf_{p \in F} J_{(\eta, \pi)}(p).$$

(ii) *Let Γ be closed subset of $\mathcal{L}(\mathcal{Z} \times \mathcal{Z}_*)$. Then we have*

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \log Q_{(\eta_n, \pi_n)} \left\{ y \in \mathcal{G}([n], (\eta_n, \pi_n)) \mid P_z \in \Gamma \right\} \geq - \inf_{p \in \Gamma} J_{(\eta, \pi)}(p).$$

Finally we state from Theorem 2.2 a corollary for the case of the classical Erdos-Renyi graph $\mathcal{G}([n], nc/2)$.

Corollary 2.3 (LDP). *Suppose D_z is the degree distribution of $z \in \mathcal{G}([n], nc/2)$ and let Γ be subset of $\mathcal{L}(\mathcal{Z} \times \mathcal{Z}_*)$. Then we have*

$$\begin{aligned} - \inf_{p \in \text{int}(\Gamma)} I_c(p) &\leq \liminf_{n \rightarrow \infty} \frac{1}{n} \log Q_{(\eta_n, \pi_n)} \left\{ z \in \mathcal{G}([n], nc/2) \mid D_z \in \Gamma \right\} \\ &\leq \limsup_{n \rightarrow \infty} \frac{1}{n} \log Q_{(\eta_n, \pi_n)} \left\{ z \in \mathcal{G}([n], nc/2) \mid D_z \in \Gamma \right\} \leq - \inf_{p \in \text{cl}(\Gamma)} I_c(p), \end{aligned}$$

where q_c is the poisson distribution with mean c , $I_c(p) = H(p \mid q_c)$ and ∞ if otherwise.

3. PROOF OF MAIN RESULTS

3.1 Proof of Theorem 2.1.

We denote by \tilde{z} the multitype random allocation process, see example [5] or [4] and note that for any consistent triple (η_n, π_n, p_n) we have

$$\frac{Q_{(\eta_n, \pi_n)} \left\{ Z = z \mid P_z \in B_p \right\}}{\tilde{Q}_{(\eta_n, \pi_n)} \left\{ Z = z \mid P_z \in B_{\tilde{p}} \right\}} = \frac{\#\{y: P_z = p_n\}}{\#\{z: P_z = \tilde{p}_n\}} = \frac{\#\{z: P_z = p_n\}}{\#\{y: (P^1(y), P^2(y)) = (\eta_n, \pi_n)\}} \times \frac{\#\{z: (P^1(z), P^2(z)) = (\eta_n, \pi_n)\}}{\#\{z: P_z = \tilde{p}_n\}}. \quad (3.1)$$

Now we define a locality of the functional ρ as follows:

$$B_p = \left\{ \mu \in \mathcal{L}(\mathcal{Z} \times \mathcal{Z}_*) : \left\langle \mu, \log \frac{\mu}{q} \right\rangle > \left\langle p, \log \frac{p}{q} \right\rangle - \frac{\varepsilon}{2} \right\}.$$

Using Doku-Amponsah [5], under the condition $P_z \in B_p$ and $\tilde{p} = q$ we have that

$$e^{-nH(p_n \parallel q_n) + nH(q \parallel q_n) - o(n)} \leq \frac{dQ_{(\eta_n, \pi_n)}(z)}{d\tilde{Q}_{(\eta_n, \pi_n)}(z)} \leq e^{-nH(p_n \parallel q_n) + nH(q \parallel q_n) + o(n)}, \quad (3.2)$$

Fix $\varepsilon > 0$, and note that by 3.2 we have

$$\begin{aligned} Q_{(\eta_n, \pi_n)} \left\{ z \in \mathcal{G}([n], (\eta_n, \pi_n)) \mid P_z \in B_p \right\} &\leq \int_{P_z \in B_p} e^{-nH(p_n \parallel q_n) + nH(q \parallel q_n) + o(n)} d\tilde{Q}_{(\eta_n, \pi_n)}(z) \\ &\leq e^{-nJ_{(\eta, \pi)}(p) - n\varepsilon + o(n)}. \end{aligned}$$

Also we have

$$\begin{aligned} Q_{(\eta_n, \pi_n)} \left\{ z \in \mathcal{G}([n], (\eta_n, \pi_n)) \mid P_z \in B_p \right\} &\geq \int_{P_z \in B_p} e^{-nH(p_n \| q_n) + nH(q \| q_n) - o(n)} d\tilde{Q}_{(\eta_n, \pi_n)}(z) \\ &\geq e^{-nJ_{(\eta, \pi)}(p) - n\varepsilon - o(n)}, \end{aligned}$$

which completes the proof of the Theorem. The proof of Theorem 2.2 below, follows from Theorem 2.1 above using similar arguments as in [3, p. 544].

3.2 Proof of Theorem 2.2.

Proof. We observe that the empirical locality distribution is a probability distribution and so it is contained in the unit ball in $\tilde{\mathcal{L}}(\mathcal{Z} \times \mathcal{Z}^*)$. Henceforth, without loss of generality we can assume that the set Γ in Theorem 2.2(ii) above is relatively compact. If we choose any $\varepsilon > 0$, then for each functional $p \in \Gamma$ we can find a weak neighbourhood such that the estimate of Theorem 2.1(i) above holds. From all these neighbourhoods, we choose a finite cover of $\mathcal{G}([n], (\eta_n, \pi_n))$ and sum up over the estimate in Theorem 2.1(i) above to obtain

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log Q_{(\eta_n, \pi_n)} \left\{ z \in \mathcal{G}([n], (\eta_n, \pi_n)) \mid P_z \in \Gamma \right\} \leq - \inf_{p \in \Gamma} J_{(\eta, \pi)}(p) + \varepsilon.$$

Since ε was arbitrarily chosen and the lower bound in Theorem 2.1(ii) implies the lower bound in Theorem 2.2(i) we have the required results which completes the proof. □

3.3 Proof of Corollary 2.3.

Proof. The proof of Corollary 2.3 follows from Theorem 2.2 if we take $P = D_z$ then $\Psi(D_z) = 2|\mathcal{E}|/n = c$. Hence, in the case of the classical Erdos-Renyi graph $\mathcal{G}([n], nc/2)$, we have the convex rate function

$$I_c(p) = J_{(\eta, \pi)}(p)$$

which reduces to $H(p \| q_c)$ if the probability measure p has mean c and ∞ if otherwise. □

Acknowledgement

This article was finalized at the Carnegie Banga-Africa, June 27-July 2017 writeshop, in Koforidua.

REFERENCES

- [1] C. BORDENAVE AND P. CAPUTO Large deviations of empirical neighbourhood distribution in sparse random graphs. *arxiv:1308.5725(2013)*.
- [2] S. BOUCHERON, F. GAMBOA and C. LEONARD. Bins and balls: Large deviations of the empirical occupancy process. *Ann. Appl. Probab.* 12 607-636 (2002).
- [3] I.V. BAKHTIN. Spectral Potential, Kullback Action, and Large deviations of empirical measures on measurable spaces. *Theory of Probability and application.* Vol. 50, No.4. (2015) pp.535-544.
- [4] K. DOKU-AMPONSAH. Large deviations and basic information theory for hierarchical and networked data structures. *Bath Thesis(2006)*.
- [5] K. DOKU-AMPONSAH. Exponential Approximation of empirical neighbourhood measures of random graphs by random allocation. *International J. of Statistics and Probability,* Vol. 3, No 2; 2014. pp 110-120.

- [6] K. DOKU-AMPONSAH. Local Large deviations: McMillian Theorem for multitype Galton-Watson Processes. *Far East J. of Mathematical Sciences* 102(10),pp. 313-337
- [7] K. DOKU-AMPONSAH. Local Large deviations: a McMillian Theorem for Typed Random Graph Processes. *J. of Mathematics and Statistics* 13(4), pp.325-329
- [8] K. DOKU-AMPONSAH and P. MÖRTERS. Large deviation principle for empirical measures of coloured random graphs. *The Annals of Applied Probability*, 20(6), 1989-2021.
- [9] A. DEMBO and O. ZEITOUNI. Large deviations techniques and applications. Springer, New York, (1998).
- [10] R. VAN DER HOFSTAD Random Graphs and Complex Networks. *Eindhoven University of Technology*. Unpublished Manuscript.