

Weight hierarchy of a class of linear codes relating to non-degenerate quadratic forms

Fei Li

Received: date / Accepted: date

Abstract In this paper, we discuss the generalized Hamming weights of a class of linear codes associated with non-degenerate quadratic forms. In order to do so, we study the quadratic forms over subspaces of finite field and obtain some interesting results about subspaces and their dual spaces. On this basis, we solve all the generalized Hamming weights of these linear codes.

Keywords Generalized Hamming weight · Linear code · Quadratic form · Weight hierarchy · Dual space

Mathematics Subject Classification (2010) 94B05 · 11E04 · 11T71

1 Introduction

Let p^m be a power of an odd prime p . Denote F_{p^m} the finite field with p^m elements and $F_{p^m}^*$ the multiplicative group of F_{p^m} .

If C is a k -dimensional F_p -vector subspace of F_{p^n} , then it is called an $[n, k, d]$ p -ary linear code with length n and minimum Hamming distance d [10]. For linear code C , the concept of generalized Hamming weights (GHW) $d_r(C)$ ($0 < r \leq k$) can be viewed as the extension of Hamming weight (see [13, 18]). Let $[C, r]_p$ be the set of all r -dimensional F_p -vector subspaces of C . For $V \in [C, r]_p$, define $Supp(V) = \{i | x_i \neq 0 \text{ for some } x = (x_1, x_2, \dots, x_n) \in V\}$. Then we define the r -th generalized Hamming weight (GHW) $d_r(C)$ of linear code C by

$$d_r(C) = \min\{|Supp(V)| | V \in [C, r]_p\},$$

This research is supported in part by National Natural Science Foundation of China (Grant Nos.61602342).

Fei Li
E-mail: cczxf@163.com

Faculty of School of Statistics and Applied Mathematics, Anhui University of Finance and Economics, Bengbu, Anhui Province, 233041, P.R.China

In particular, $d_1(C) = d$. And $\{d_i(C) | 1 \leq i \leq k\}$ is called the weight hierarchy of C .

Generalized Hamming weight of linear codes has been an interesting topic in both theory and practice for many years. In 1991, Wei in the paper [18] presented his classic results, in which GHW was shown that it can characterise the cryptography performance of linear codes used on the wire-tap channel of type II. From then on, much more attention was paid to the generalized Hamming weight. A detailed survey on the results up to 1995 about GHW can be found In [17]. Afterwards lots of authors devoted themselves to generalized Hamming weight about particular classes of codes [1, 2, 3, 6, 9, 11, 12, 19, 20]. Recently, Minghui Yang et al. in their work [20] gave a very constructive method for the GHWs of irreducible cyclic codes. Generally, it is hard to settle the weight hierarchy of a linear code.

Ding et al. proposed a generic construction of linear code as below ([4, 5]). Let Tr be the trace function from F_{p^m} to F_p and $D = \{d_1, d_2, \dots, d_n\}$ be contained in F_{p^m} . Define a p -ary linear code C_D with length n as following.

$$C_D = \{(Tr(xd_1), Tr(xd_2), \dots, Tr(xd_n)) : x \in F_{p^m}\} \quad (1)$$

and D is called the defining set. Many classes of linear codes with few weights were obtained by choosing properly defining sets [8, 15, 21, 22, 24]. In this paper, we discuss the generalized Hamming weights of a class of p -ary linear codes C_D , whose defining set was chosen to be

$$D = D_a = \{x \in F_{p^m} | f(x) = a\}, \quad a \in F_p. \quad (2)$$

Here f is a non-degenerate quadratic form over F_{p^m} .

And the weight hierarchy of C_{D_0} can be deduced by Theorem 9 in [25], in which Mr. Wan used the theory of finite projective geometry for the purpose.

In reference [23], Kelan Ding and Cunsheng Ding presented the linear codes C_{D_a} of the case $a = 0$ for the special quadratic form $Tr(X^2)$ and determined their weight distributions. In [22] and [7], the weight distributions of C_{D_a} for general quadratic forms were settled. By the main results of them, we know that C_{D_a} is an m -dimensional linear code. So we can employ a general formula for linear codes defined in (1) to calculate $d_r(C_{D_a})$. We give it in the following lemma.

Lemma 1.(Theorem 1, [16]) For each $r(1 \leq r \leq m)$, if the dimension of C_D is m , then $d_r(C_D) = n - \max\{|D \cap H| : H \in [F_{p^m}, m - r]_p\}$.

The rest of this paper is organised as follows: in Section 2, we present some basic definitions and results on quadratic form over subspace of finite fields and dual spaces; in Section 3, using the results in Section 2, we give all the generalized Hamming weights of linear codes defined in (2); in Section 4, we give the conclusion of this paper.

2 Quadratic Form over Subspace of Finite Fields

We may view F_{p^m} as an m -dimensional vector space over F_p . Namely, F_{p^m} is F_p -linearly isomorphic to F_p^m . We fix m elements $v_1, v_2, \dots, v_m \in F_{p^m}$ as a basis and express each vector $X \in F_{p^m}$ in the unique form $X = x_1v_1 + x_2v_2 + \dots + x_mv_m$, for some $x_1, x_2, \dots, x_m \in F_p$. By the isomorphism, we can write $X = (x_1, x_2, \dots, x_m)^T$, where T represents the transpose of a matrix.

Now we fix a basis $v_1, v_2, \dots, v_m \in F_{p^m}$. Recall that a map $f : F_{p^m} \rightarrow F_p$ is called a quadratic form over F_{p^m} with values in F_p if

$$f(X) = f(x_1, x_2, \dots, x_m) = \sum_{1 \leq i, j \leq m} a_{ij}x_ix_j, a_{ij} \in F_p,$$

for each element $X = x_1v_1 + x_2v_2 + \dots + x_mv_m$. Denote $F(X, Y) = \frac{1}{2}[f(X+Y) - f(X) - f(Y)]$, so $f(X) = F(X, X)$. If $a_{ij} = a_{ji}$, we can write $f(X) = X^TAX$, where A is the symmetric matrix $(a_{ij})_{1 \leq i, j \leq m}$. The rank R_f of quadratic form f is defined to be the rank of matrix A . It is well-known that $a_{ii} = f(v_i)$, $a_{ij} = F(v_i, v_j)$.

For a subspace $H \subseteq F_{p^m}$, the dual space H^\perp of H under quadratic form f is defined by

$$H^\perp = \{x \in F_{p^m} | f(x+y) = f(x) + f(y) \text{ for each } y \in H\}.$$

Let G be another subspace of F_{p^m} . If $f(x+y) = f(x) + f(y)$ for each $x \in H$ and $y \in G$, we say that H and G are perpendicular to each other under quadratic form f . It is denoted by $H \perp G$. It is well-known that $R_f + \dim(F_{p^m}^\perp) = m$. If $R_f = m$, we say that the quadratic form f is non-degenerate. And the discriminant $\Delta(f)$ of the quadratic form f is defined as the determinant $\det(A)$ of A . We can find an invertible matrix M such that M^TAM is a diagonal matrix $\Lambda = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_{R_f}, 0, \dots, 0)$. When $0 < R_f < m$, we also can define the discriminant $\Delta(f) = \lambda_1 \cdot \lambda_2 \cdot \dots \cdot \lambda_{R_f}$. When $R_f = 0$, we define $\Delta(f) = 1$. It is easy to see that $\Delta(f)$ varies with the change of the basis of F_{p^m} . But $\bar{\eta}(\Delta(f))$ is an invariant, which is called the sign of the quadratic form f . We denote it by ϵ_f . Here $\bar{\eta}$ is the quadratic character of F_p . Meanwhile, $\bar{\eta}(0)$ is defined to be zero.

Let H be a d -dimensional subspace of F_{p^m} . Restricting the quadratic form f on H , we get a quadratic form over H in d variables. It is denoted by $f|_H$. Similarly, we define the dual space $H_{f|_H}^\perp$ of H under $f|_H$ in itself by

$$H_{f|_H}^\perp = \{x \in H | f(x+y) = f(x) + f(y) \text{ for each } y \in H\}.$$

Let $R_{f|_H}, \Delta(f|_H)$ be the rank and discriminant of $f|_H$ over H , respectively. Obviously, $H_{f|_H}^\perp = H \cap H^\perp$ and $R_{f|_H} = d - \dim(H_{f|_H}^\perp)$.

For k elements $\beta_1, \beta_2, \dots, \beta_k \in F_{p^m}$, the discriminant $\Delta(\beta_1, \beta_2, \dots, \beta_k)$ of them is defined as the determinant of the matrix $M = (F(\beta_i, \beta_j))_{1 \leq i, j \leq k}$.

From now on, we discuss only the non-degenerate quadratic forms. Its symmetric matrix is $(F(v_i, v_j))_{1 \leq i, j \leq m}$ if we fix an F_p -basis v_1, v_2, \dots, v_m

of F_{p^m} . Thus we have that the discriminant $\Delta(f) = \Delta(v_1, v_2 \cdots v_m)$. For a subspace H of F_{p^m} , we know that $\dim(H) + \dim(H^\perp) = m$ and denote $R_H = R_{f|_H}$, $\Delta_H = \Delta(f|_H)$. If we fix an F_p -basis $\mu_1, \mu_2 \cdots \mu_d$ of H , then $\Delta_H = \Delta(\mu_1, \mu_2 \cdots \mu_d)$. Let $v(x)$ be a function over F_p defined by $v(x) = p-1$ if $x = 0$, otherwise $v(x) = -1$. For $a \in F_p$, set $\overline{D}_a = \{x \in F_{p^m} | f(x) = a\}$. Then we have following results.

Proposition 1. Let H be a d -dimensional ($d > 0$) subspace of F_{p^m} , then

$$|H \cap \overline{D}_a| = \begin{cases} p^{d-1} + v(a)\overline{\eta}((-1)^{\frac{R_H}{2}} \Delta_H) p^{d-\frac{R_H+2}{2}}, & \text{if } R_H \equiv 0 \pmod{2}, \\ p^{d-1} + \overline{\eta}((-1)^{\frac{R_H-1}{2}} a \Delta_H) p^{d-\frac{R_H+1}{2}}, & \text{if } R_H \equiv 1 \pmod{2}. \end{cases}$$

Proof. It is the natural result of quadratic forms from Theorem 6.26 and 6.27 in [14]. We omit the details.

Proposition 2. For each r ($0 < 2r < m$), there exist an r -dimensional subspace $H \subseteq F_{p^m}$ ($m > 2$) such that $H \subseteq H^\perp$, i.e., $H \perp H$.

Proof. It suffices to prove there exist r elements $\alpha_1, \alpha_2, \dots, \alpha_r \in F_{p^m}$ satisfying the following two conditions:

1. $F(\alpha_i, \alpha_j) = 0, 1 \leq i, j \leq r$;
2. $\alpha_1, \alpha_2, \dots, \alpha_r$ are linearly independent over F_p .

If $r = 1$, by Lemma 1, $\overline{D}_0 \neq \{0\}$. So it is true for $r = 1$. Suppose we have k elements $\alpha_1, \alpha_2, \dots, \alpha_k$, which satisfy the two conditions. It might as well assume $2k+2 < m$. Otherwise, the proof is finished. Let $H_k = \langle \alpha_1, \alpha_2, \dots, \alpha_k \rangle$. So H_k^\perp is an $(m-k)$ -dimension subspace. By Proposition 1, $|\overline{D}_0 \cap H_k^\perp| = p^{m-k-1} \pm (p-1)p^{m-k-1-\frac{k}{2}}$ if t is even, otherwise $|\overline{D}_0 \cap H_k^\perp| = p^{m-k-1}$. Here $t = R_{H_k^\perp}$. Let $V_k = \{x \in H_k^\perp | F(x, y) = 0, y \in H_k^\perp\}$. So $\dim(V_k) = m-k-t$. Since $V_k \subset H_k, m-k-t \leq k$. Hence $t \geq m-2k > 2$. Therefore $|\overline{D}_0 \cap H_k^\perp| > p^{m-k-1} - (p-1)p^{m-k-2} = p^{m-k-2} > p^k$. We obtain that there is an element $\alpha_{k+1} \in (\overline{D}_0 \cap H_k^\perp) \setminus H_k$. So $\alpha_1, \alpha_2, \dots, \alpha_k, \alpha_{k+1}$ are $(k+1)$ elements that match the conditions as above. By mathematical induction, we finish the proof.

Proposition 3. Let $m = 2s > 2$. There exists an s -dimensional subspace $H_s \subset F_{p^m}$ such that $H_s = H_s^\perp$ if and only if $\epsilon_f = (-1)^{\frac{m(p-1)}{4}}$.

Proof. By Proposition 2, there exist $(s-1)$ elements $\alpha_1, \alpha_2, \dots, \alpha_{s-1}$ that match the following conditions:

1. $F(\alpha_i, \alpha_j) = 0, 1 \leq i, j \leq s-1$;
2. $\alpha_1, \alpha_2, \dots, \alpha_{s-1}$ are linearly independent over F_p .

Let $H_{s-1} = \langle \alpha_1, \alpha_2, \dots, \alpha_{s-1} \rangle$. So H_{s-1}^\perp is an $(s+1)$ -dimension subspace and the rank of quadratic form $f(x)$ over H_{s-1}^\perp is 2. Let

$$H_{s-1}^\perp = \langle \alpha_1, \alpha_2, \dots, \alpha_{s-1}, \alpha_s, \alpha_{s+1} \rangle$$

and $H_2 = \langle \alpha_s, \alpha_{s+1} \rangle$. It is easy to see $\Delta_{H_{s-1}^\perp} = \Delta(\alpha_s, \alpha_{s+1})$ and $H_2 \cap H_2^\perp = \{0\}$. Suppose

$$H_2^\perp = \langle \alpha_1, \alpha_2, \dots, \alpha_{s-1}, \beta_1, \beta_2, \dots, \beta_{s-1} \rangle$$

So $\Delta_m = \Delta(\alpha_1, \dots, \alpha_{s-1}, \alpha_s, \alpha_{s+1}, \beta_1, \dots, \beta_{s-1}) = (-1)^{s+1} \det(M^2) \Delta_{H_{s-1}^\perp}$. Here M is the square matrix $(F(\alpha_i, \beta_j))_{1 \leq i, j \leq s-1}$. By Proposition 1, we have $|H_{s-1}^\perp \cap \overline{D_0}| = p^s + \overline{\eta}((-1)^s \Delta(f))(p-1)p^{s-1}$. If $\overline{\eta}(\Delta(f)) = (-1)^{\frac{m(p-1)}{4}}$, then $|H_{s-1}^\perp \cap \overline{D_0}| = 2p^s - p^{s-1} > p^{s-1}$. So we can choose one element $\alpha_s \in (H_{s-1}^\perp \cap \overline{D_0}) \setminus H_{s-1}$ and set $H_s = \langle \alpha_1, \alpha_2, \dots, \alpha_{s-1}, \alpha_s \rangle$. It is easy to check $H_s^\perp = H_s$. If $\overline{\eta}(\Delta(f)) = -(-1)^{\frac{m(p-1)}{4}}$, then $|H_{s-1}^\perp \cap \overline{D_0}| = p^{s-1}$. So $H_{s-1}^\perp \cap \overline{D_0}$ must be H_{s-1} . In this case, the conclusion of the proposition is untenable. We complete the proof.

3 Weight Hierarchy of Linear Codes of (2)

In this section, we settle the weight hierarchy of the linear code defined in (2).

3.1 Case 1: $a \neq 0$

Theorem 1. If m is even, then for the linear codes defined in (2) we have

$$d_r(C_{D_a}) = \begin{cases} p^{m-1} - p^{m-r-1} - ((-1)^{\frac{m(p-1)}{4}} \epsilon_f + 1)p^{\frac{m-2}{2}}, & \text{if } 1 \leq r \leq \frac{m}{2}, \\ p^{m-1} - 2p^{m-r-1} - (-1)^{\frac{m(p-1)}{4}} \epsilon_f p^{\frac{m-2}{2}}, & \text{if } \frac{m}{2} \leq r < m, \\ p^{m-1} - (-1)^{\frac{m(p-1)}{4}} \epsilon_f p^{\frac{m-2}{2}}, & \text{if } r = m. \end{cases}$$

Proof. We only give the proof of the case: $\epsilon_f = -(-1)^{\frac{m(p-1)}{4}}$. The proof of the other case: $\epsilon_f = (-1)^{\frac{m(p-1)}{4}}$ is similar.

For $1 \leq r < \frac{m}{2}$, let $t = R_{H_{m-r}}$, the rank of $f(x)$ over some $(m-r)$ -dimensional subspace H_{m-r} . By Proposition 2,3, $0 \leq \dim(H_{m-r} \cap H_{m-r}^\perp) \leq r$. So $0 \leq m-r-t \leq r$. Hence $m-2r \leq t \leq m-r$.

If $t = m-2r$, we may construct

$$H_{m-r} = \langle \alpha_1, \alpha_2, \dots, \alpha_{m-2r}, \beta_1, \beta_2, \dots, \beta_r \rangle, H_r = H_{m-r}^\perp = \langle \beta_1, \beta_2, \dots, \beta_r \rangle,$$

$$H_{m-2r} = \langle \alpha_1, \alpha_2, \dots, \alpha_{m-2r} \rangle, H_{2r} = H_{m-2r}^\perp = \langle \beta_1, \beta_2, \dots, \beta_r, \beta_{r+1}, \dots, \beta_{2r} \rangle.$$

Because $H_{m-2r} \cap H_{m-2r}^\perp = \{0\}$,

$$\begin{aligned} \Delta_m &= \Delta(\alpha_1, \alpha_2, \dots, \alpha_{m-2r}, \beta_1, \beta_2, \dots, \beta_r, \beta_{r+1}, \dots, \beta_{2r}) \\ &= \Delta_{H_{m-r}} \Delta(\beta_1, \beta_2, \dots, \beta_r, \beta_{r+1}, \dots, \beta_{2r}) = (-1)^r \Delta_{H_{m-r}} \det(M^2). \end{aligned}$$

Here M is the square matrix $M = (F(\beta_i, \beta_j))_{1 \leq i \leq r, r+1 \leq j \leq 2r}$. Hence for $t = m-2r$, by Proposition 1, $|D_a \cap H_{m-r}| = p^{m-r-1} - \overline{\eta}((-1)^{\frac{m}{2}}) \epsilon_f p^{\frac{m-2}{2}} =$

$p^{m-r-1} + p^{\frac{m-2}{2}}$. So by Proposition 1, we have that $\max\{|D_a \cap H| : H \in [F_{p^m}, m-r]_p\} = p^{m-r-1} + p^{\frac{m-2}{2}}$.

For $\frac{m}{2} \leq r < m$, by Proposition 2, there is such an $(m-r-1)$ -dimensional subspace $H_{m-r-1} = \langle \alpha_1, \alpha_2, \dots, \alpha_{m-r-1} \rangle$ that $H_{m-r-1} \subseteq H_{m-r-1}^\perp$. By Proposition 1, $|D_b \cap H_{m-r-1}^\perp| > 0$ with $b \in F_p^*$. We choose an element $\alpha_{m-r} \in (D_b \cap H_{m-r-1}^\perp)$ and get an $(m-r)$ -dimensional subspace

$$H_{m-r} = \langle \alpha_1, \alpha_2, \dots, \alpha_{m-r-1}, \alpha_{m-r} \rangle.$$

By the construction of H_{m-r} , we know that the rank of $f(x)$ over H_{m-r} is 1 and $\Delta_{H_{m-r}} = b$. So by Proposition 1, $|D_a \cap H_{m-r}| = p^{m-r-1} + \bar{\eta}(ab)p^{m-r-1}$. Also by Proposition 1, we have that $\max\{|D_a \cap H| : H \in [F_{p^m}, m-r]_p\} = 2p^{m-r-1}$. By Proposition 1, we have $|D_a| = p^{m-1} - \epsilon_f(-1)^{\frac{m(p-1)}{4}} p^{\frac{m-2}{2}}$. Then the desired results follow directly from Lemma 1. And we complete the proof.

Theorem 2. If m is odd and $\bar{\eta}(a) = (-1)^{\frac{(m-1)(p-1)}{4}} \epsilon_f$, then for the linear codes defined in (2) we have

$$d_r(C_{D_a}) = \begin{cases} p^{m-1} - p^{m-r-1}, & \text{if } 1 \leq r < \frac{m}{2}, \\ p^{m-1} + p^{\frac{m-1}{2}} - 2p^{m-r-1}, & \text{if } \frac{m}{2} < r < m, \\ p^{m-1} + p^{\frac{m-1}{2}}, & \text{if } r = m. \end{cases}$$

Proof. For $1 \leq r < \frac{m}{2}$, we may construct subspaces

$$H_{m-r} = \langle \alpha_1, \alpha_2, \dots, \alpha_{m-2r}, \beta_1, \beta_2, \dots, \beta_r \rangle, H_r = H_{m-r}^\perp = \langle \beta_1, \beta_2, \dots, \beta_r \rangle,$$

$$H_{m-2r} = \langle \alpha_1, \alpha_2, \dots, \alpha_{m-2r} \rangle, H_{2r} = H_{m-2r}^\perp = \langle \beta_1, \beta_2, \dots, \beta_r, \beta_{r+1}, \dots, \beta_{2r} \rangle.$$

Because $H_{m-2r} \cap H_{m-2r}^\perp = \{0\}$,

$$\begin{aligned} \Delta_m &= \Delta(\alpha_1, \alpha_2, \dots, \alpha_{m-2r}, \beta_1, \beta_2, \dots, \beta_r, \beta_{r+1}, \dots, \beta_{2r}) \\ &= \Delta_{H_{m-r}} \Delta(\beta_1, \beta_2, \dots, \beta_r, \beta_{r+1}, \dots, \beta_{2r}) = (-1)^r \Delta_{H_{m-r}} \det(M^2). \end{aligned}$$

Here M is the square matrix $M = (F(\beta_i, \beta_j))_{1 \leq i \leq r, r+1 \leq j \leq 2r}$. Note that $\bar{\eta}(a) = (-1)^{\frac{(m-1)(p-1)}{4}} \epsilon_f$. So $|D_a \cap H_{m-r}| = p^{m-r-1} + \bar{\eta}((-1)^{\frac{m}{2}} a) \epsilon_f p^{\frac{m-1}{2}} = p^{m-r-1} + p^{\frac{m-1}{2}}$. By Proposition 1, we have that $\max\{|D_a \cap H| : H \in [F_{p^m}, m-r]_p\} = p^{m-r-1} + p^{\frac{m-1}{2}}$.

For the case: $\frac{m}{2} < r < m$, the proof is the same as that of Theorem 1. By Proposition 1, we have $|D_a| = p^{m-1} + \epsilon_f(-1)^{\frac{m(p-1)}{4}} \bar{\eta}(a) p^{\frac{m-1}{2}}$. Then the desired conclusions follow from Lemma 1. And the proof is completed.

Theorem 3. If m is odd and $\bar{\eta}(a) = -(-1)^{\frac{(m-1)(p-1)}{4}} \epsilon_f$, then for the linear codes defined in (2) we have

$$d_r(C_{D_a}) = \begin{cases} p^{m-1} - p^{m-r-1} - p^{\frac{m-1}{2}} - p^{\frac{m-3}{2}}, & \text{if } 1 \leq r < \frac{m}{2}, \\ p^{m-1} - p^{\frac{m-1}{2}} - 2p^{m-r-1}, & \text{if } \frac{m}{2} < r < m, \\ p^{m-1} - p^{\frac{m-1}{2}}, & \text{if } r = m. \end{cases}$$

Proof. The proof is similar to those of Theorem 1 and Theorem 2. We omit the details.

3.2 Case 2: $a = 0$

Theorem 4. If m is even and $\epsilon_f = (-1)^{\frac{m(p-1)}{4}}$, then for the linear codes defined in (2) we have

$$d_r(C_{D_0}) = \begin{cases} p^{m-1} - p^{m-r-1}, & \text{if } 1 \leq r \leq \frac{m}{2}, \\ p^{m-1} + (p-1)p^{\frac{m-2}{2}} - p^{m-r}, & \text{if } \frac{m}{2} \leq r \leq m. \end{cases}$$

Proof. Denote by H_k a k -dimensional subspace in F_{p^m} . By Lemma 1 and Proposition 2 and 3 as above, it is easy to know $d_r(C_{D_0}) = |\overline{D}_0| - p^{m-r}$, if $\frac{m}{2} \leq r \leq m$.

For $1 \leq r < \frac{m}{2}$, let $t = R_{H_{m-r}}$, the rank of quadratic form $f(x)$ over some $(m-r)$ -dimensional subspace H_{m-r} . For $0 \leq \dim(H_{m-r} \cap H_{m-r}^\perp) \leq r$, so $0 \leq m-r-t \leq r$. Hence $m-2r \leq t \leq m-r$.

If $r = 1$, then $d_r(C_{D_0})$ is the minimum Hamming distance of C_{D_0} . The result for this case can be obtained in reference [22].

For the case $\frac{m}{2} > r \geq 2$, if $t = m-2r$, we may construct

$$H_{m-r} = \langle \alpha_1, \alpha_2, \dots, \alpha_{m-2r}, \beta_1, \beta_2, \dots, \beta_r \rangle, H_r = H_{m-r}^\perp = \langle \beta_1, \beta_2, \dots, \beta_r \rangle,$$

$$H_{m-2r} = \langle \alpha_1, \alpha_2, \dots, \alpha_{m-2r} \rangle, H_{2r} = H_{m-2r}^\perp = \langle \beta_1, \beta_2, \dots, \beta_r, \beta_{r+1}, \dots, \beta_{2r} \rangle.$$

Because $H_{m-2r} \cap H_{m-2r}^\perp = \{0\}$,

$$\Delta_m = \Delta(\alpha_1, \alpha_2, \dots, \alpha_{m-2r}, \beta_1, \beta_2, \dots, \beta_r, \beta_{r+1}, \dots, \beta_{2r})$$

$$= \Delta_{H_{m-r}} \Delta(\beta_1, \beta_2, \dots, \beta_r, \beta_{r+1}, \dots, \beta_{2r}) = (-1)^r \Delta_{H_{m-r}} \det(M^2).$$

Here M is the square matrix $M = (F(\beta_i, \beta_j))_{1 \leq i \leq r, r+1 \leq j \leq 2r}$. Notice that $\epsilon_f = (-1)^{\frac{m(p-1)}{4}}$. Hence for $t = m-2r$, by Proposition 1, $|\overline{D}_0 \cap H_{m-r}| = p^{m-r-1} + \eta((-1)^{\frac{m-2r}{2}} \Delta_{H_{m-r}})(p-1)p^{m-r-1-\frac{m-2r}{2}} = p^{m-r-1} + (p-1)p^{\frac{m-2}{2}}$. So by Proposition 1, we have that $\max\{|\overline{D}_0 \cap H| : H \in [F_{p^m}, m-r]_p\} = p^{m-r-1} + (p-1)p^{\frac{m-2}{2}}$. Then the desired results follow directly from Lemma 1. And we complete the proof.

Theorem 5. If $m = 2s > 2$ is even and $\epsilon_f = -(-1)^{\frac{m(p-1)}{4}}$, then for the linear codes defined in (2) we have

$$d_r(C_{D_0}) = \begin{cases} (p-1)(p^{m-2} - p^{\frac{m-2}{2}}), & \text{if } r = 1, \\ p^{m-1} - p^{m-r-1} - (p-1)(p^{\frac{m-2}{2}} + p^{\frac{m-4}{2}}), & \text{if } 2 \leq r \leq \frac{m}{2}, \\ p^{m-1} - p^{m-r} - (p-1)p^{\frac{m-2}{2}}, & \text{if } \frac{m}{2} < r \leq m. \end{cases}$$

Proof. We only give the proof of the two cases: $2 \leq r < \frac{m}{2}$ and $r = \frac{m}{2}$. The proofs of the remaining cases are the same as that of Theorem 1.

For $2 \leq r < \frac{m}{2}$, let the symbols be as in the proof of Theorem 1. By the proof of Theorem 1, we have $|\overline{D}_0 \cap H_{m-r}| = p^{m-r-1} - (p-1)p^{\frac{m-2}{2}}$ if $t = m-2r$.

If $t = m - 2r + 2$, by Proposition 3, we can choose such a subspace $G_{r-2} = \langle \varepsilon_1, \varepsilon_2, \dots, \varepsilon_{r-2} \rangle$ that $G_{r-2} \subset G_{r-2}^\perp$. So the rank of $f(x)$ over G_{r-2}^\perp is $m - 2r + 4$. By Proposition 1, $|\overline{D}_a \cap G_{r-2}^\perp| = p^{m-r+2-1} \pm p^{m-r+2-1-\frac{m-2r+4}{2}} > 0$ with $a \in F_p^*$. So we can choose a nonzero element $\varepsilon_{r-1} \in \overline{D}_a \cap G_{r-2}^\perp$ and get an $(r-1)$ -dimensional subspace $G_{r-1} = \langle \varepsilon_1, \varepsilon_2, \dots, \varepsilon_{r-2}, \varepsilon_{r-1} \rangle$. It is easy to see that the rank of $f(x)$ over G_{r-1} is 1 and $G_{r-1} \cap G_{r-1}^\perp = G_{r-2}$. So the rank of $f(x)$ over G_{r-1}^\perp is $m - 2r + 3$. By Proposition 1, $|\overline{D}_b \cap G_{r-1}^\perp| = p^{m-r} \pm p^{m-r+1-\frac{m-2r+4}{2}} > p^{r-1}$ with $b \in F_p^*$. So we obtain an r -dimensional subspace $H_r = \langle \varepsilon_1, \varepsilon_2, \dots, \varepsilon_{r-1}, \varepsilon_r \rangle$ by choosing $\varepsilon_r \in (\overline{D}_b \cap G_{r-1}^\perp) \setminus G_{r-1}$ and the rank of $f(x)$ over H_r is 2. So we can construct another $(m-r)$ -dimensional subspace $H_{m-r} = H_r^\perp$ and set

$$H_{m-r} = \langle \gamma_1, \gamma_2, \dots, \gamma_{m-2r+2}, \varepsilon_1, \varepsilon_2, \dots, \varepsilon_{r-2} \rangle.$$

Let $H_{m-2r+2} = \langle \gamma_1, \gamma_2, \dots, \gamma_{m-2r+2} \rangle$ and set

$$H_{m-2r+2}^\perp = \langle \varepsilon_1, \dots, \varepsilon_{r-1}, \varepsilon_r, \varepsilon_{r+1}, \dots, \varepsilon_{2r-2} \rangle.$$

Because $R_{H_{m-r}} = m - 2r + 2$, we get $H_{m-2r+2}^\perp \cap H_{m-2r+2} = \{0\}$ and

$$\begin{aligned} \Delta_m &= \Delta(\gamma_1, \gamma_2, \dots, \gamma_{m-2r+2}, \varepsilon_1, \dots, \varepsilon_{r-1}, \varepsilon_r, \varepsilon_{r+1}, \dots, \varepsilon_{2r-2}) \\ &= \Delta_{H_{m-r}} \cdot \Delta(\varepsilon_1, \dots, \varepsilon_{r-1}, \varepsilon_r, \varepsilon_{r+1}, \dots, \varepsilon_{2r-2}) \\ &= (-1)^r \Delta_{H_{m-r}} ab \det(M^2). \end{aligned}$$

Here M is the square matrix $M = (F(\varepsilon_i, \varepsilon_j))_{1 \leq i \leq r-2, r+1 \leq j \leq 2r-2}$. According to the selection of $\varepsilon_{r-1}, \varepsilon_r$, we know that $\overline{\eta}(\Delta_{H_{m-r}})$ may take 1 or -1 . Hence for $t = m - 2r + 2$, by Proposition 1, $|\overline{D}_0 \cap H_{m-r}| = p^{m-r-1} \pm (p-1)p^{\frac{m}{2}-2}$.

For $r = \frac{m}{2}$, let H_s be an s -dimensional subspace. We have $1 \leq R_{H_s} \leq s$ since $\epsilon_f = -(-1)^{\frac{m(p-1)}{4}}$ by Proposition 3. If $R_{H_s} = 1$, then $|\overline{D}_0 \cap H_s| = p^{s-1}$. If $R_{H_s} = 2$, then $|\overline{D}_0 \cap H_s| = p^{s-1} + \overline{\eta}(-\Delta_{H_s})(p-1)p^{s-2}$. By Proposition 2 and Proposition 3, there exists an $(s-2)$ -dimensional subspace H_{s-2} such that $H_{s-2} \subset H_{s-2}^\perp$. Set $H_{s-2} = \langle \alpha_1, \alpha_2, \dots, \alpha_{s-2} \rangle$. So the rank of quadratic form f over H_{s-2}^\perp is 4. By Proposition 1, $|\overline{D}_a \cap H_{s-2}^\perp| = p^{s+1} \pm p^{s-1} > 0$ with $a \in F_p^*$. We choose an element $\alpha_{s-1} \in \overline{D}_a \cap H_{s-2}^\perp$ and get an $(s-1)$ -dimensional subspace $H_{s-1} = \langle \alpha_1, \alpha_2, \dots, \alpha_{s-2}, \alpha_{s-1} \rangle$. The rank of quadratic form f over H_{s-1}^\perp is 3. Also by Proposition 1, $|\overline{D}_b \cap H_{s-1}^\perp| = p^s \pm p^{s-1} > p^{s-1}$ with $b \in F_p^*$. So we choose an element $\alpha_s \in (\overline{D}_b \cap H_{s-1}^\perp) \setminus H_{s-1}$ and get an s -dimensional subspace $H_s = \langle \alpha_1, \alpha_2, \dots, \alpha_{s-2}, \alpha_{s-1}, \alpha_s \rangle$. By the construction of H_s , we know that the rank of f over H_s is 2 and $\overline{\eta}(\Delta_{H_s}) = \overline{\eta}(\Delta(\alpha_{s-1}, \alpha_s))$ may take -1 or 1. So by Proposition 1, $|\overline{D}_0 \cap H_s| = p^{s-1} \pm (p-1)p^{s-2}$.

In a word, we have that when $2 \leq r \leq \frac{m}{2}$, $\max\{|\overline{D}_0 \cap H| : H \in [F_p^m, m-r]_p\} = p^{m-r-1} + (p-1)p^{\frac{m-4}{2}}$. Then we arrive at the conclusion by Lemma 1. And the proof is finished.

Theorem 6. If m is odd, then for the linear codes defined in (2) we have

$$d_r(C_{D_0}) = \begin{cases} p^{m-1} - p^{m-r-1} - (p-1)p^{\frac{m-3}{2}}, & \text{if } 1 \leq r < \frac{m}{2}, \\ p^{m-1} - p^{m-r}, & \text{if } \frac{m}{2} < r \leq m. \end{cases}$$

Proof. We only give the proof of the case: $2 \leq r < \frac{m}{2}$. For an $(m-r)$ -dimensional subspace H_{m-r} , let $t = R_{H_{m-r}}$. We have $m-2r \leq t \leq m-r$. By Proposition 2, we can choose such an $(r-1)$ -dimensional subspace $G_{r-1} = \langle \beta_1, \beta_2, \dots, \beta_{r-1} \rangle$ that $G_{r-1} \subset G_{r-1}^\perp$. So the rank of f over G_{r-1}^\perp is $m-2r+2$. By Proposition 1, $|D_a \cap G_{r-1}^\perp| = p^{m-r} \pm p^{\frac{m-1}{2}} > 0$ with $a \in F_p^*$. So we can choose a nonzero element $\beta_r \in D_a \cap G_{r-1}^\perp$ and get an r -dimensional subspace $H_r = \langle \beta_1, \beta_2, \dots, \beta_{r-1}, \beta_r \rangle$. It is easy to see that the rank of f over H_r is 1 and $H_r \cap H_r^\perp = G_{r-1}$. So the rank of f over H_r^\perp is $m-2r+1$. We construct an $(m-r)$ -dimensional subspace $H_{m-r} = H_r^\perp$ and set

$$H_{m-r} = \langle \alpha_1, \alpha_2, \dots, \alpha_{m-2r+1}, \beta_1, \beta_2, \dots, \beta_{r-1} \rangle.$$

Let $H_{m-2r+1} = \langle \alpha_1, \alpha_2, \dots, \alpha_{m-2r+1} \rangle$ and set

$$H_{m-2r+1}^\perp = \langle \beta_1, \beta_2, \dots, \beta_{r-1}, \beta_r, \beta_{r+1}, \dots, \beta_{2r-1} \rangle.$$

Because $R_{H_{m-r}} = m-2r+1$, we get $H_{m-2r+1}^\perp \cap H_{m-2r+1} = \{0\}$ and

$$\begin{aligned} \Delta_m &= \Delta(\alpha_1, \alpha_2, \dots, \alpha_{m-2r+1}, \beta_1, \beta_2, \dots, \beta_{r-1}, \beta_r, \beta_{r+1}, \dots, \beta_{2r-1}) \\ &= \Delta_{H_{m-r}} \cdot \Delta(\beta_1, \beta_2, \dots, \beta_{r-1}, \beta_r, \beta_{r+1}, \dots, \beta_{2r-1}) \\ &= (-1)^r \Delta_{H_{m-r}} a \det(M^2). \end{aligned}$$

Here M is the square matrix $M = (F(\beta_i, \beta_j))_{1 \leq i \leq r-1, r+1 \leq j \leq 2r-1}$. According to the selection of β_r , we know that $\bar{\eta}(\Delta_{H_{m-r}})$ may take 1 or -1 . Hence for an $(m-r)$ -dimensional subspace H_{m-r} with $R_{H_{m-r}} = m-2r+1$, by Proposition 1, we have $|\overline{D}_0 \cap H_{m-r}| = p^{m-r-1} \pm (p-1)p^{\frac{m-3}{2}}$. Applying Proposition 1 once again, we have that when $2 \leq r < \frac{m}{2}$, $\max\{|\overline{D}_0 \cap H| : H \in [F_p^m, m-r]_p\} = p^{m-r-1} + (p-1)p^{\frac{m-3}{2}}$. Using Lemma 1, we can obtain the results in the theorem. The proof is finished.

4 Concluding Remarks

Quadratic forms are ones of the well-known polynomials in the theory of algebra. In the paper, we restrict quadratic forms over finite fields of odd characteristic on their subspaces and prove the existence of some special subspaces and their dual spaces related to non-degenerate quadratic forms. Then we determine completely the weight hierarchy of a class of linear codes from non-degenerate quadratic forms. We hope that our results may help to research the generalized Hamming weight of other linear codes.

References

1. M. Bras-Amors, K. Lee, and A. Vico-Oton, New lower bounds on the generalized Hamming weights of AG codes, *IEEE Trans. Inf. Theory*, 60(10), 5930-5937(2014).
2. A. I. Barbero and C. Munuera, The weight hierarchy of Hermitian codes, *SIAM J. Discrete Math.*, 13(1), 79-104(2000).
3. J. Cheng and C.-C. Chao, On generalized Hamming weights of binary primitive BCH codes with minimum distance one less than a power of two, *IEEE Trans. Inf. Theory*, 43(1), 294-298(1997).
4. C. Ding, Linear codes from some 2-designs, *IEEE Trans. Inf. Theory*, 61(6), 3265-3275(2015).
5. K. Ding, C. Ding, Binary linear codes with three weights, *IEEE Communication Letters*, 18(11), 1879-1882(2014).
6. M. Delgado, J. I. Farnn, P. A. Garca-Snchez, and D. Llena, On the weight hierarchy of codes coming from semigroups with two generators, *IEEE Trans. Inf. Theory*, 60(1), 282-295(2014).
7. Xiaoni Du and Yunqi Wan, Linear codes from quadratic forms, *Applicable Algebra in Engineering, Communication and Computing*, doi:10.1007/s00200-017-0319-x(2017).
8. C. Ding, C. Li, N.Li, Z. Zhou, Three-weight cyclic codes and their weight distributions. *Discret. Math.*, 339(2), 415-427 (2016).
9. P. Heijnen and R. Pellikaan, Generalized Hamming weights of q-ary ReedCMuller codes, *IEEE Trans. Inf. Theory*, 44(1), 181-196(1998).
10. W. C. Huffman and V. Pless, *Fundamentals of error-correcting codes*, Cambridge University Press, Cambridge(2003).
11. G. Jian, R. Feng and H. Wu, Generalized Hamming weights of three classes of linear codes, *Finite Fields and Their Applications*, 45, 341-354(2017).
12. H. Janwa and A. K. Lal, On the generalized Hamming weights of cyclic codes, *IEEE Trans. Inf. Theory*, 43(1), 299-308(1997).
13. T. Kløve, The weight distribution of linear codes over $GF(q^l)$ having generator matrix over $GF(q >)$, *Discrete Math.*, 23(2), 159-168(1978).
14. R. Lidl, H. Niederreiter, *Finite fields*, Cambridge University Press, New York(1997).
15. C. Li, Q. Yue, and F. Li, Hamming weights of the duals of cyclic codes with two zeros, *IEEE Trans. Inf. Theory*, 60(7), 3895-3902(2014).
16. F. Li, A class of cyclotomic linear codes and their generalized Hamming weights, arXiv:1708.04415(2017).
17. M. A. Tsfasman, S. G. Vladut, Geometric approach to higher weights, *IEEE Trans. Inf. Theory*, 41(6), 1564-1588(1995).
18. V.K.Weil, Generalized Hamming weights for linear codes, *IEEE Trans. Inf. Theory*, 37(5), 1412-1418(1991).
19. M. Xiong, S. Li, and G. Ge. The weight hierarchy of some reducible cyclic codes. *IEEE Trans. Inf. Theory*, 62(7), 4071-4080(2016).
20. M. Yang, J. Li, K. Feng and D. Lin, Generalized Hamming weights of irreducible cyclic codes, *IEEE Trans. Inf. Theory*, 61(9), 4905-4913(2015).
21. S.Yang, Z.Yao, Complete weight enumerators of a family of three-weight linear codes. *Des. Codes Cryptogr.* 82(3), 663-674(2017).
22. Z. Zhou, N. Li, C. Fan and T. Hellesteth, Linear codes with two or three weights from quadratic bent functions, *Des. Codes Cryptogr.*, 81(2), 283-295(2016).
23. K. Ding, C. Ding, A class of two-weight and three-weight codes and their applications in secret sharing, *IEEE Trans. Inf. Theory*, 61(11), 5835-5842(2015).
24. C. Tang, C. Xiang and K. Feng, Linear codes with few weights from inhomogeneous quadratic functions, *Des. Codes Cryptogr.*, 83(3), 691-714(2017).
25. Z. Wan, The weight hierarchies of the projective codes from nondegenerate quadrics, *Des. Codes Cryptogr.*, 4(4), 283-300(1994).