

Standard Bases for Linear Codes over Prime Fields

Jean Jacques Ferdinand RANDRIAMIARAMPANAHY⁽¹⁾

e-mail : randriamiferdinand@gmail.com

Harinaivo ANDRIATAHINY⁽²⁾

e-mail : hariandriatahiny@gmail.com

Toussaint Joseph RABEHERIMANANA⁽³⁾

e-mail : rabeherimanana.toussaint@yahoo.fr

^{1,2,3}Mention : Mathematics and Computer Science,
Domain : Sciences and Technologies,
University of Antananarivo, Madagascar

January 29, 2022

Abstract

It is known that a linear code can be represented by a binomial ideal. In this paper, we give standard bases for the ideals in a localization of the multivariate polynomial ring in the case of the linear codes over prime fields.

Keywords : Linear code, semigroup order, Groebner basis, local ring, standard basis

MSC 2010 : 13P10, 94B05, 12E20

1 Introduction

Coding theory is the mathematical basis for data transmission through noisy communication channels. It contains two main parts. The first part is to encode the message to reduce its sensitivity to noise during transmission. The second part is to decode the received message by detecting and correcting the errors.

Bruno Buchberger introduced the theory of Groebner bases for polynomial ideals in 1965. The Groebner bases theory can be used to solve some problems concerning the ideals by developing computations in multivariate polynomial rings. In 1964, Hironaka [12] introduced the analogues of Groebner bases called standard bases for ideals in the localization of the polynomial ring at the origin. In [6], standard bases for ideals generated by polynomials in local rings can be determined by using the same method as Groebner bases.

Connection between linear codes and ideals in polynomial rings was presented in [2]. And it was proved that a Groebner basis of the ideal associated to a binary linear code can be used for determining the minimum distance . A generalization to linear codes over prime fields can be found in [15, 16]. In [15], it has been proved that a linear code can be described by a binomial ideal, and a Groebner basis with respect to lexicographic order for the binomial ideal is determined .

The aim of this paper is to present the standard basis of the ideal of a linear code over a prime field in the local ring of rational functions that are regular at a point of the affine variety associated to the ideal. The idea is to generalize the method developed by N. Dück and K. H. Zimmermann in [9].

This paper is organized as follows. The second section presents a background for Groebner bases. The third section contains the division algorithm in a local ring. In section 4, the notion of linear codes and their connections with binomial ideals are presented. The main result is contained in section 5.

2 Preliminaries

Throughout this paper, n denotes a positive integer, \mathbb{K} a commutative field and $\mathbb{K}[X] := \mathbb{K}[X_1, \dots, X_n]$ the polynomial ring in n variables over \mathbb{K} . We denote by 0 the zero element of \mathbb{N}^n where \mathbb{N} is the set of non negative integers.

A monomial in $\mathbb{K}[X]$ is an algebraic expression of the form $X_1^{\alpha_1} \dots X_n^{\alpha_n}$ which is denoted by X^α where $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$. The monomial $X^\alpha = X_1^{\alpha_1} \dots X_n^{\alpha_n}$ can be identified with the n -tuple of exponents $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ and vice versa, thus there exists a one-to-one correspondence between the monomials in $\mathbb{K}[X]$ and the elements of \mathbb{N}^n . The degree of the monomial X^α with $\alpha = (\alpha_1, \dots, \alpha_n)$ is $|\alpha| := \alpha_1 + \alpha_2 + \dots + \alpha_n$. Any order $>$ we establish on \mathbb{N}^n will give us an order on the set of monomials in $\mathbb{K}[X]$: if $\alpha > \beta$ according to this order, we have $X^\alpha > X^\beta$. An order $>$ in $\mathbb{K}[X]$ is compatible with multiplication if for all X^α, X^β and X^γ in $\mathbb{K}[X]$ with $X^\alpha > X^\beta$ then $X^\alpha X^\gamma > X^\beta X^\gamma$. Now let $>$ be an order on the set of monomials X^u where $u \in \mathbb{N}^n$. We say that $>$ is a semigroup order in $\mathbb{K}[X]$ if $>$ is a total order and it is compatible with the multiplication of monomials. A monomial order on $\mathbb{K}[X]$ is a semigroup order such that $1 < X_i$ for $i = 1, \dots, n$. Usual monomial orders on $\mathbb{K}[X]$ are the lexicographic order, the degree lexicographic order and the degree reverse lexicographic order.

Let f be a non-zero polynomial of $\mathbb{K}[X]$ such that $f = \sum_{i=1}^k c_i X^{\alpha_i}$ where $c_i \in \mathbb{K}$ and $\alpha_i \in \mathbb{N}^n$. Let

us fix a monomial order $>$ on $\mathbb{K}[X]$. A term in $\mathbb{K}[X]$ is a scalar times a monomial. The leading term of f , denoted by $\text{lt}_>(f)$, is the largest involved term with respect to $>$. If $\text{lt}_>(f) = c_m X^{\alpha_m}$ where $1 \leq m \leq k$, then c_m is called the leading coefficient of f ($\text{lc}_>(f)$) and X^{α_m} is the leading monomial of f ($\text{lm}_>(f)$). We denote by $\text{deg}(f)$ the maximal degree of all monomials occurring in f . The reduction of a polynomial f by a polynomial g , denoted by $\text{Red}(f, g)$ is defined by

$$\text{Red}(f, g) := f - q.g$$

where $\text{lt}_>(f) = q.\text{lt}_>(g)$, for some term $q = cX^\alpha$. Let $\mathcal{F} = (f_1, \dots, f_s)$ be a s -tuple of polynomials in $\mathbb{K}[X]$. Each polynomial $f \in \mathbb{K}[X]$ can be written in the form :

$$f = a_1 f_1 + a_2 f_2 + \dots + a_s f_s + r$$

where $a_1, \dots, a_s, r \in \mathbb{K}[X]$ and either $r = 0$ or r is a \mathbb{K} -linear combination of monomials, none of which is divisible by any of $\text{lt}_>(f_1), \dots, \text{lt}_>(f_s)$. Moreover, if $a_i f_i \neq 0$, then $\text{lt}_>(f) \geq \text{lt}_>(a_i f_i)$, $1 \leq i \leq s$. The polynomial r is called the remainder of f on division by \mathcal{F} . The remainder r is produced by the following algorithm called division algorithm in $\mathbb{K}[X]$ (see [5]).

```

Input :  $f_1, \dots, f_s, f$ 
Output :  $r$ 
 $r := 0$ 
 $p := f$ 
WHILE  $p \neq 0$  DO

```

```

i := 1
divisionoccurred:= false
WHILE i ≤ s AND divisionoccurred = false DO
    IF lt>(fi) divides lt>(p) THEN
        p := Red(p , fi)
        divisionoccurred:= true
    ELSE
        i := i + 1
IF divisionoccurred = false THEN
    r := r + lt>(p)
    p := p - lt>(p)

```

The division algorithm terminates after a finite number of steps.

Let $I \subset \mathbb{K}[X]$ be a non-zero ideal and $>$ a monomial order. The ideal generated by the set of $\text{lt}_>(f)$ where $f \in I$ is called the leading ideal of I , denoted by $\text{lt}_>(I)$, i.e

$$\text{lt}_>(I) := \langle \text{lt}_>(f) / f \in I \rangle.$$

For a finite subset $G = \{g_1, \dots, g_s\}$ of the ideal I , we denote by $\text{lt}_>(G)$ the ideal generated by the $\text{lt}_>(g_i)$, for $i = 1, \dots, s$, i.e

$$\text{lt}_>(G) := \langle \text{lt}_>(g_1), \dots, \text{lt}_>(g_s) \rangle.$$

A finite subset $G = \{g_1, \dots, g_s\}$ of an ideal I is called a Groebner basis of I if $\text{lt}_>(G) = \text{lt}_>(I)$. A Groebner basis G for the polynomial ideal I is called a minimal Groebner basis of I if for all $g \in G$, $\text{lc}_>(g) = 1$ and $\text{lt}_>(g) \notin \text{lt}_>(G - \{g\})$. The reduced Groebner basis for the ideal I is a Groebner basis G satisfying :

- (1) $\text{lc}_>(g) = 1$ for all $g \in G$,
- (2) no monomial of g lies in $\text{lt}_>(G - \{g\})$, for all $g \in G$.

A Groebner basis can be determined by using Buchberger's algorithm. Let $f, g \in \mathbb{K}[X]$ be non-zero polynomials. Fix a monomial order $>$ and let X^γ be the least common multiple of the leading monomial of f and the leading monomial of g . The S-polynomial of f and g , denoted by $\text{spoly}(f, g)$ is

$$\text{spoly}(f, g) := \frac{X^\gamma}{\text{lt}_>(f)} \cdot f - \frac{X^\gamma}{\text{lt}_>(g)} \cdot g$$

If I is a polynomial ideal, then a basis $G = \{g_1, \dots, g_s\}$ for I is a Groebner basis for I if and only if for all pairs $i \neq j$, the remainder on division of $\text{spoly}(g_i, g_j)$ by G is zero.

3 Localization

Let \mathbb{K} be a commutative field and $p = (p_1, \dots, p_n)$ an n -tuple of \mathbb{K}^n . $\mathbb{K}(X) := \mathbb{K}(X_1, \dots, X_n)$ denotes the field of rational functions containing $\mathbb{K}[X]$. We define a local ring in $\mathbb{K}(X)$ by the set $\mathcal{O}_p := \left\{ \frac{f}{g} / f, g \in \mathbb{K}[X], g(p) \neq 0 \right\}$. We say also that we localize at the maximal ideal $\langle X_1 - p_1, \dots, X_n - p_n \rangle \subset \mathbb{K}[X]$. Let m_p be the ideal generated by $X_1 - p_1, \dots, X_n - p_n$ in \mathcal{O}_p . Then each element in $\mathcal{O}_p \setminus m_p$ is a unit in \mathcal{O}_p .

A local order in $\mathbb{K}[X] = \mathbb{K}[X_1, \dots, X_n]$ is a semigroup order such that $1 > X_i$ for all $1 \leq i \leq n$. For instance, for two n -tuples of \mathbb{N}^n $\alpha = (\alpha_1, \dots, \alpha_n)$ and $\beta = (\beta_1, \dots, \beta_n)$, we define the negative degree lexicographic order by $\alpha > \beta$ if $|\alpha| < |\beta|$ or $|\alpha| = |\beta|$ and there exists an integer $i \in \{1, \dots, n\}$ such that $\alpha_1 = \beta_1, \dots, \alpha_{i-1} = \beta_{i-1}, \alpha_i > \beta_i$.

Let $>$ be a local order on the set of monomials of $\mathbb{K}[X]$ and let $S_{>} := \{1 + g \in \mathbb{K}[X] / g = 0 \text{ or } \text{lt}_{>}(g) < 1\}$. $S_{>}$ is a multiplicative part of $\mathbb{K}[X]$ and we remark that $S_{>} = \mathbb{K}[X] \setminus \langle X_1, \dots, X_n \rangle$. Define the localization of $\mathbb{K}[X]$ in view of the order $>$ by the ring ([6, 13])

$$\text{Loc}_{>}(\mathbb{K}[X]) := S_{>}^{-1}\mathbb{K}[X] = \left\{ \frac{f}{(1+g)} / f \in \mathbb{K}[X], 1+g \in S_{>} \right\}. \quad (1)$$

We have $\text{Loc}_{>}(\mathbb{K}[X]) = \mathcal{O}_{p=0}$. Under the local order, there is a difficulty for the successive reductions, because we may have an infinite strictly decreasing sequence of terms. For example, consider the polynomials of one variable X , $f = X$ and $g = X - X^2$, and we divide f by g by using the division algorithm, so that we successively reduce by $X - X^2$. This gives the reductions:

$$\begin{aligned} f_1 &:= \text{Red}(f, g) = X^2 \\ f_2 &:= \text{Red}(f_1, g) = X^3 \\ &\vdots \\ f_n &:= \text{Red}(f_{n-1}, g) = X^{n+1}, \end{aligned}$$

and so on.

Mora introduced a method to solve this problem. The following result can be found in [6].

Theorem 3.1 (Mora normal form algorithm).

Given non-zero polynomials $f, f_1, \dots, f_s \in \mathbb{K}[X]$ and let $>$ be a local order. There is an algorithm which gives the polynomials $u, a_1, \dots, a_s, h \in \mathbb{K}[X]$ such that

$$uf = a_1f_1 + \dots + a_sf_s + h \quad (2)$$

where $\text{lt}_{>}(u) = 1$ ($u = 1 + g$ is unit in $\text{Loc}_{>}(\mathbb{K}[X])$), $\text{lt}_{>}(f) \geq \text{lt}_{>}(a_i)\text{lt}_{>}(f_i)$ for all i with $a_i \neq 0$, and $h = 0$ or $\text{lt}_{>}(h)$ is not divisible by any of $\text{lt}_{>}(f_i)$. We denote $NF(f | G) := h$ with $G = \{f_1, \dots, f_s\}$, and we say that h is the weak normal form of f on division by G .

For $f \in \mathbb{K}[X]$, we define $\text{ecart}(f) := \deg(f) - \deg(\text{lt}_{>}(f))$. The remainder h in (2) is produced by the following algorithm called Mora's division algorithm

```

h := f;  L := {f_1, ..., f_s};  M := {g \in L : lt_{>}(g) | lt_{>}(h)}
WHILE (h \neq 0 AND M \neq \emptyset) DO
SELECT  g \in M  with ecart(g)  minimal
IF ecart(g) > ecart(h)  THEN
    L := L \cup {h}
h := Red(h, g)
IF h \neq 0  THEN
    M := {g \in L : lt_{>}(g) | lt_{>}(h)}

```

4 Linear codes and binomial ideals

Let \mathbb{F}_p be the finite field with p elements where p is a prime number. A linear code \mathcal{C} of length n and dimension k over \mathbb{F}_p is the image of a linear (injective) mapping

$$\psi : \mathbb{F}_p^k \longrightarrow \mathbb{F}_p^n$$

where $k \leq n$. The elements $x = (x_1, \dots, x_n) \in \mathcal{C}$ are called the codewords. The weight of a word $x = (x_1, \dots, x_n) \in \mathbb{F}_p^n$ is defined by $w_t(x) := \text{card}\{i/x_i \neq 0, 1 \leq i \leq n\}$. The minimum distance of the linear code \mathcal{C} is $d := \min\{d(x, y)/x, y \in \mathcal{C}, x \neq y\}$ or $d := \min\{w_t(x)/x \in \mathcal{C}, x \neq 0\}$ where $d(x, y) := \text{card}\{i/x_i \neq y_i\}$. We define the support of an element $x \in \mathcal{C}$ by $\text{supp}(x) := \{i/x_i \neq 0\}$. A linear code \mathcal{C} of length n and dimension k is called an $[n, k]$ -code. Moreover, if the minimum distance is d , we say that \mathcal{C} is an $[n, k, d]$ -code.

Let \mathcal{C} be an $[n, k]$ -code, $e_i = (\zeta_{i1}, \dots, \zeta_{ik})$ where $i = 1, \dots, k$ the canonical basis of \mathbb{F}_p^k and $\psi(e_i) = (g_{i1}, \dots, g_{in})$. The generating matrix of \mathcal{C} is the matrix of dimension $k \times n$ defined by $G = (g_{ij})$ where $g_{ij} \in \mathbb{F}_p$. The linear code \mathcal{C} is represented as follows $\mathcal{C} = \{xG \mid x \in \mathbb{F}_p^k\}$. We will say that G is in standard form if $G = (I_k \mid M)$ where I_k is the $k \times k$ identity matrix. Let \mathcal{C} be an $[n, k]$ -code over \mathbb{F}_p . Define the ideal associated with \mathcal{C} as ([2, 8, 10, 16])

$$I_{\mathcal{C}} := \langle X^c - X^{c'} \mid c - c' \in \mathcal{C} \rangle + \langle X_i^p - 1 \mid 1 \leq i \leq n \rangle \quad (3)$$

where each word $c \in \mathbb{F}_p^n$ is considered as an integral vector in the monomial X^c . Let \mathcal{C} be an $[n, k]$ -code over \mathbb{F}_p and

$$G = (g_{ij}) = (I_k \mid M) \quad (4)$$

a generating matrix in standard form. Let m_i be the vector of length n over \mathbb{F}_p defined by

$$m_i = (0, \dots, 0, p - g_{i,k+1}, \dots, p - g_{i,n}) \quad (5)$$

for $1 \leq i \leq k$. We have $X^{m_i} = X_{k+1}^{p-g_{i,k+1}} X_{k+2}^{p-g_{i,k+2}} \dots X_n^{p-g_{i,n}} = \prod_{j \in \text{supp}(m_i)} X_j^{p-g_{i,j}}$. In particular, if $\text{supp}(m_i) = \emptyset$, then $X^{m_i} = 1$.

Theorem 4.1. *Let us take the lexicographic order on $\mathbb{K}[X]$ with $X_1 > X_2 > \dots > X_n$. The code ideal $I_{\mathcal{C}}$ has the reduced Groebner basis*

$$\mathcal{G} = \{X_i - X^{m_i} \mid 1 \leq i \leq k\} \cup \{X_i^p - 1 \mid k+1 \leq i \leq n\}. \quad (6)$$

Proof. A proof can be found in [15].

5 Standard bases

In this section, we will describe the analogues of Groebner bases called standard bases for the ideals in local rings by Mora's division algorithm. Given any local order $>$ on monomials in $\mathbb{K}[X]$, there is a natural extension of $>$ to $\text{Loc}_{>}(\mathbb{K}[X])$, which we will also denote by $>$. For any $h = \frac{f}{1+g} \in \text{Loc}_{>}(\mathbb{K}[X])$ as in (1), we define $\text{lm}_{>}(h) := \text{lm}_{>}(f)$; $\text{lc}_{>}(h) := \text{lc}_{>}(f)$ and $\text{lt}_{>}(h) := \text{lt}_{>}(f)$. We fix a local order $>$ on $\text{Loc}_{>}(\mathbb{K}[X])$ and let I be an ideal in $\text{Loc}_{>}(\mathbb{K}[X])$. A standard basis of I is a subset $\{f_1, \dots, f_r\}$ of I such that $\text{lt}_{>}(I) = \langle \text{lt}_{>}(f_1), \dots, \text{lt}_{>}(f_r) \rangle$ where $\text{lt}_{>}(I)$ is the ideal generated by the set of $\text{lt}_{>}(f)$ with $f \in I$.

Proposition 5.1 (Product criterion). [11]

Let $f, g \in \mathbb{K}[X_1, \dots, X_n]$ be polynomials such that $\text{lcm}(\text{lm}_{>}(f), \text{lm}_{>}(g)) = \text{lm}_{>}(f) \cdot \text{lm}_{>}(g)$, then

$$NF\left(\text{spoly}(f, g) \mid \{f, g\}\right) = 0 \quad (7)$$

where $NF(- \mid -)$ is defined as in theorem(3.1).

We will consider the ideals of the local ring $\text{Loc}_{>}(\mathbb{K}[X])$ which are generated by polynomials of $\mathbb{K}[X]$. A more general result of the following theorem can be found in [11].

Theorem 5.2 (Buchberger criterion).

Let $I \subset \text{Loc}_>(\mathbb{K}[X])$ be an ideal, $G = \{g_1, \dots, g_s\}$ a set of polynomials of I and $>$ a local order. Let $NF(- | -)$ be the weak normal form as in theorem(3.1). Then the following are equivalent:

- i) G is a standard basis of I .
- ii) G generates I and $NF(\text{spoly}(g_i, g_j) | G) = 0$ for $i, j = 1, \dots, s$.
- iii) G generates I and $NF(\text{spoly}(g_i, g_j) | G_{ij}) = 0$ for a suitable subset $G_{ij} \subset G$ and $i, j = 1, \dots, s$.

Let \mathcal{C} be an $[n, k]$ -code, the point $(1, \dots, 1)$ is a zero of the code ideal $I_{\mathcal{C}}$ in the affine space over \mathbb{F}_p . Rather than localizing at the maximal ideal $\langle X_1 - 1, \dots, X_n - 1 \rangle$, we change coordinates to translate the point to the origin. Denote $I'_{\mathcal{C}}$ the corresponding ideal, and $I := I'_{\mathcal{C}} \text{Loc}_>(\mathbb{F}_p[X])$ the ideal of $\text{Loc}_>(\mathbb{F}_p[X])$ generated by $I'_{\mathcal{C}}$.

Lemma 5.3. Let p a prime number, we have

$$I'_{\mathcal{C}} = \left\langle (X_i + 1) + (p - 1) \prod_{j \in \text{supp}(m_i)} (X_j + 1)^{p - g_{i,j}} / 1 \leq i \leq k \right\rangle + \left\langle (X_i + 1)^{p + p - 1 / k + 1} \leq i \leq n \right\rangle$$

where $g_{i,j}$ is defined in (4) and m_i in (5).

Proof. The ideal $I_{\mathcal{C}}$ defined in (3) has the reduced Groebner basis (6) by the theorem(4.1) with respect to the lexicographic order on $\mathbb{K}[X]$. This is an ideal basis of $I_{\mathcal{C}}$ in $\mathbb{K}[X]$. The translation is made via the ring map $X_i \mapsto X_i + 1$. Since $\mathbb{K}[X] \subset \text{Loc}_>(\mathbb{K}[X])$, then the claim for the translated ideal follows. □

Now we present our main result

Theorem 5.4. Let \mathcal{C} be an $[n, k]$ -code over \mathbb{F}_p with p a prime number. Under the negative degree lexicographic order $>$ on $\mathbb{F}_p[X]$, the ideal $I = I'_{\mathcal{C}} \text{Loc}_>(\mathbb{F}_p[X])$ in $\text{Loc}_>(\mathbb{F}_p[X])$ has the standard basis

$$S = \left\{ X_i - \sum_{\substack{0 \leq t_l \leq p - g_{i,j_l} (1 \leq l \leq \sigma_i) \\ (t_1, \dots, t_{\sigma_i}) \neq (0, \dots, 0)}} \prod_{h=1}^{\sigma_i} \binom{p - g_{i,j_h}}{t_h} X_{j_h}^{t_h} / 1 \leq i \leq k \right\} \cup \left\{ X_i^p / k + 1 \leq i \leq n \right\} \quad (8)$$

where $\sigma_i := \text{card}(\text{supp}(m_i))$.

Proof. We will show that S generates $I'_{\mathcal{C}}$, then we prove that S is a standard basis.

For $k + 1 \leq i \leq n$

Let $X_i^p \in S$ and we will prove that $X_i^p \in I'_{\mathcal{C}}$. For $j \in \{1, 2, \dots, p - 1\}$, the number $\binom{p}{j}$ is a

multiple of p because p is prime. Since we work over a field of characteristic p , we have

$$\begin{aligned}
X_i^p &= \sum_{j=1}^{p-1} \binom{p}{j} X_i^j + X_i^p \\
&= 1 + \sum_{j=1}^{p-1} \binom{p}{j} X_i^j + X_i^p - 1 \\
&= \binom{p}{0} X_i^0 + \sum_{j=1}^{p-1} \binom{p}{j} X_i^j + \binom{p}{p} X_i^p - 1 \\
&= \sum_{j=0}^p \binom{p}{j} X_i^j - 1 \\
&= \sum_{j=0}^p \binom{p}{j} X_i^j + p - 1 \\
&= (X_i + 1)^p + p - 1 \in I'_C.
\end{aligned}$$

For $1 \leq i \leq k$

Let $X_i - \sum_{\substack{0 \leq t_l \leq p - g_{i,j_l} (1 \leq l \leq \sigma_i) \\ (t_1, \dots, t_{\sigma_i}) \neq (0, \dots, 0)}} \prod_{h=1}^{\sigma_i} \binom{p - g_{i,j_h}}{t_h} X_{j_h}^{t_h} \in S$. Suppose that $\text{supp}(m_i) = \{j_1, j_2, \dots, j_{\sigma_i}\}$

with $j_1 < j_2 < \dots < j_{\sigma_i}$ and denote $A := X_i - \sum_{\substack{0 \leq t_l \leq p - g_{i,j_l} (1 \leq l \leq \sigma_i) \\ (t_1, \dots, t_{\sigma_i}) \neq (0, \dots, 0)}} \prod_{h=1}^{\sigma_i} \binom{p - g_{i,j_h}}{t_h} X_{j_h}^{t_h}$. We have

$$\begin{aligned}
A &= X_i + 1 - \left[\sum_{\substack{0 \leq t_l \leq p - g_{i,j_l} (1 \leq l \leq \sigma_i) \\ (t_1, \dots, t_{\sigma_i}) \neq (0, \dots, 0)}} \prod_{h=1}^{\sigma_i} \binom{p - g_{i,j_h}}{t_h} X_{j_h}^{t_h} + 1 \right] \\
&= X_i + 1 - \sum_{0 \leq t_l \leq p - g_{i,j_l} (1 \leq l \leq \sigma_i)} \prod_{h=1}^{\sigma_i} \binom{p - g_{i,j_h}}{t_h} X_{j_h}^{t_h} \\
&= X_i + 1 - \sum_{t_1=0}^{p - g_{i,j_1}} \sum_{t_2=0}^{p - g_{i,j_2}} \dots \sum_{t_{\sigma_i}=0}^{p - g_{i,j_{\sigma_i}}} \binom{p - g_{i,j_1}}{t_1} \binom{p - g_{i,j_2}}{t_2} \dots \binom{p - g_{i,j_{\sigma_i}}}{t_{\sigma_i}} X_{j_1}^{t_1} X_{j_2}^{t_2} \dots X_{j_{\sigma_i}}^{t_{\sigma_i}} \\
&= X_i + 1 - \left[\sum_{t_1=0}^{p - g_{i,j_1}} \binom{p - g_{i,j_1}}{t_1} X_{j_1}^{t_1} \right] \left[\sum_{t_2=0}^{p - g_{i,j_2}} \binom{p - g_{i,j_2}}{t_2} X_{j_2}^{t_2} \right] \dots \left[\sum_{t_{\sigma_i}=0}^{p - g_{i,j_{\sigma_i}}} \binom{p - g_{i,j_{\sigma_i}}}{t_{\sigma_i}} X_{j_{\sigma_i}}^{t_{\sigma_i}} \right] \\
&= X_i + 1 - \left[(X_{j_1} + 1)^{p - g_{i,j_1}} \right] \left[(X_{j_2} + 1)^{p - g_{i,j_2}} \right] \dots \left[(X_{j_{\sigma_i}} + 1)^{p - g_{i,j_{\sigma_i}}} \right] \\
&= X_i + 1 - \prod_{j \in \text{supp}(m_i)} (X_j + 1)^{p - g_{i,j}} \\
&= X_i + 1 + p \left[\prod_{j \in \text{supp}(m_i)} (X_j + 1)^{p - g_{i,j}} \right] - \prod_{j \in \text{supp}(m_i)} (X_j + 1)^{p - g_{i,j}} \\
&= X_i + 1 + (p - 1) \left[\prod_{j \in \text{supp}(m_i)} (X_j + 1)^{p - g_{i,j}} \right] \in I'_C
\end{aligned}$$

By lemma (5.3), S is a generating set for I'_C .

Let us now show that S is a standard basis over \mathbb{F}_p

* Let the pair (i, j) such that $k + 1 \leq i < j \leq n$.

We have $\text{spoly}(X_i^p, X_j^p) = \frac{X_i^p X_j^p}{X_i^p} X_i^p - \frac{X_i^p X_j^p}{X_j^p} X_j^p = X_i^p X_j^p - X_i^p X_j^p = 0$, and $NF(0 \mid S) = 0$.

* Let the pair (i, j) such that $1 \leq i \leq k$ and $k + 1 \leq j \leq n$.

Denote $f_i := X_i - \sum_{\substack{0 \leq t_l \leq p - g_{i,j_l} (1 \leq l \leq \sigma_i) \\ (t_1, \dots, t_{\sigma_i}) \neq (0, \dots, 0)}} \prod_{h=1}^{\sigma_i} \binom{p - g_{i,j_h}}{t_h} X_{j_h}^{t_h}$. We have

$$\begin{aligned} \text{spoly}(f_i, X_j^p) &= \frac{X_i X_j^p}{X_i} \left[X_i - \sum_{\substack{0 \leq t_l \leq p - g_{i,j_l} (1 \leq l \leq \sigma_i) \\ (t_1, \dots, t_{\sigma_i}) \neq (0, \dots, 0)}} \prod_{h=1}^{\sigma_i} \binom{p - g_{i,j_h}}{t_h} X_{j_h}^{t_h} \right] - \frac{X_i X_j^p}{X_j^p} X_j^p \\ &= X_i X_j^p - X_j^p \left[\sum_{\substack{0 \leq t_l \leq p - g_{i,j_l} (1 \leq l \leq \sigma_i) \\ (t_1, \dots, t_{\sigma_i}) \neq (0, \dots, 0)}} \prod_{h=1}^{\sigma_i} \binom{p - g_{i,j_h}}{t_h} X_{j_h}^{t_h} \right] - X_i X_j^p \\ &= -X_j^p \left[\sum_{\substack{0 \leq t_l \leq p - g_{i,j_l} (1 \leq l \leq \sigma_i) \\ (t_1, \dots, t_{\sigma_i}) \neq (0, \dots, 0)}} \prod_{h=1}^{\sigma_i} \binom{p - g_{i,j_h}}{t_h} X_{j_h}^{t_h} \right] \end{aligned}$$

In the last expression, all these monomials are multiple of $X_j^p \in S$. Therefore the remainder of the division of $\text{spoly}(f_i, X_j^p)$ by $\{X_j^p\}$ is zero, i.e

$$NF\left(\text{spoly}(f_i, X_j^p) \mid \{X_j^p\}\right) = 0.$$

* Finally, let $1 \leq i < j \leq k$.

Let $g_i := X_i - \sum_{\substack{0 \leq t_l \leq p - g_{i,r_l} (1 \leq l \leq \sigma_i) \\ (t_1, \dots, t_{\sigma_i}) \neq (0, \dots, 0)}} \prod_{h=1}^{\sigma_i} \binom{p - g_{i,r_h}}{t_h} X_{r_h}^{t_h}$

and $g_j := X_j - \sum_{\substack{0 \leq t'_u \leq p - g_{j,s_u} (1 \leq u \leq \sigma_j) \\ (t'_1, \dots, t'_{\sigma_j}) \neq (0, \dots, 0)}} \prod_{u=1}^{\sigma_j} \binom{p - g_{j,s_u}}{t'_u} X_{s_u}^{t'_u}$.

We have $\text{lt}_>(g_i) = X_i$ and $\text{lt}_>(g_j) = X_j$,

then $\text{lcm}(\text{lm}_>(g_i), \text{lm}_>(g_j)) = \text{lm}_>(g_i) \cdot \text{lm}_>(g_j)$. According to the Product Criterion in Proposition (5.1), we obtain

$NF\left(\text{spoly}(g_i, g_j) \mid \{g_i, g_j\}\right) = 0$. And by the Buchberger's criterion in theorem (5.2), the assertion follows. □

Example 5.5. Consider the generator matrix $G = (g_{ij})$ defined by

$$G = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 2 & 1 & 0 \\ 0 & 0 & 1 & 2 & 2 & 1 \end{pmatrix}$$

Under the negative degree lexicographic order $>$ on $\mathbb{F}_3[X]$, the ideal $I = I'_C \text{Loc}_>(\mathbb{F}_3[X])$ where I'_C is defined in lemma (5.3) with $p = 3$, $n = 6$ and $k = 3$ has the standard basis

$$S = \left\{ g_1, g_2, g_3, X_4^3, X_5^3, X_6^3 \right\}$$

where

$$g_1 = X_1 + X_4 + X_6 + 2X_4^2 + 2X_4X_6 + 2X_6^2 + X_4^2X_6 + X_4X_6^2 + 2X_4^2X_6^2,$$

$$g_2 = X_2 + 2X_4 + X_5 + X_4X_5 + 2X_5^2 + 2X_4X_5^2,$$

$$g_3 = X_3 + 2X_4 + 2X_5 + X_6 + 2X_4X_5 + X_4X_6 + X_5X_6 + 2X_6^2 + X_4X_5X_6 + 2X_4X_6^2 + 2X_5X_6^2 + 2X_4X_5X_6^2.$$

An immediate consequence is the result in [9] (for $p = 2$)

Theorem 5.6. *In view of the negative degree lexicographic order $>$ on $\mathbb{F}_2[X]$, the ideal $I = I'_C \text{Loc}_{>}(\mathbb{F}_2[X])$ in $\text{Loc}_{>}(\mathbb{F}_2[X])$ where $I'_C = \langle (X_i + 1) + \prod_{j \in \text{supp}(m_i)} (X_j + 1) / 1 \leq i \leq k \rangle + \langle (X_i + 1)^2 + 1/k + 1 \leq i \leq n \rangle$ has the standard basis :*

$$S = \left\{ X_i - \sum_{\substack{J \subseteq \text{supp}(m_i) \\ J \neq \emptyset}} X_J / 1 \leq i \leq k \right\} \cup \left\{ X_i^2 / k + 1 \leq i \leq n \right\}.$$

References

- [1] W. Adams and P. Loustanaun, *An Introduction to Groebner Bases*, American Mathematical Society, Vol.3, 1994.
- [2] M. Borges-quintana, M. Borges-trenard, P. Fitzpatrick and E. Martinez-moro, *Groebner bases and combinatorics for binary codes*, Applicable Algebra in Engineering Communication and Computing - AAEECC, Vol.19, 2008, pp. 393-411.
- [3] B. Buchberger, *An Algorithm for Finding the Basis Elements of the Residue Class Ring Modulo a Zero Dimensional Polynomial Ideal*, PhD thesis, University of Innsbruck, 1965.
- [4] A.Cooper, *Towards a new method of decoding Algebraic codes using Groebner bases*, Transactions 10th Army Conf. Appl. Math. Comp., Vol.93, 1992, pp. 293-297.
- [5] D. Cox, J. Little and D. O'Shea, *Ideals, Varieties, and Algorithms*, Springer, 1996.
- [6] D. Cox, J. Little and D. O'Shea, *Using Algebraic Geometry*, Springer, 1998.
- [7] N. Dück and K.-H. Zimmermann, *Graver Bases and Universal Gröbner Bases for Linear Codes*, May 2014.
- [8] N. Dück and K.-H. Zimmermann, *Gröbner bases for perfect binary linear codes*. International Journal of Pure and Applied Mathematics, Vol 91, 2014, 155-167.
- [9] N. Dück and K.-H. Zimmermann, *Standard Bases for binary Linear Codes*, International Journal of Pure and Applied Mathematics Volume 80, 2012.
- [10] N. Dück and K.-H. Zimmermann, *Universal Groebner bases for Binary Linear Code*, International Journal of Pure and Applied Mathematics, Appr 2013.
- [11] G.-M. Greuel and G. Pfister. *A Singular Introduction to Commutative Algebra*, Springer-Verlag, Berlin, 2002.

- [12] H.Hironaka, *Resolution of singularities of an algebraic variety over a field of characteristic zero*, Ann.Math.79(1964),109-326
- [13] T. Mora, G.Pfister and C. Traverso, *An introduction to the tangent cone algorithm*, Advances in Computing Research, 6 (1992), pp. 199-270.
- [14] M. Sala, T. Mora, L. Perret, S. Sakata, and C. Traverso, *Groebner Bases, Coding, and Cryptography*, Springer, Berlin 2009.
- [15] M. Saleemi and K.-H. Zimmermann, *Groebner Bases for Linear Codes*, International journal of Pure and Applied Mathematics, 2010, 62: 481-491.
- [16] M. Saleemi and K.-H. Zimmermann, *Linear Codes as Binomial Ideals*, International Journal of Pure and Applied Mathematics, Vol.61, June 2010, pp. 147-156.