

# Constructions of Coded Caching Schemes with Flexible Memory Size

Minquan Cheng, Jing Jiang, Qifa Yan, Xiaohu Tang, *Member, IEEE*

## Abstract

Coded caching scheme recently has become quite popular in the wireless network due to its effectively reducing the transmission amount (denote such an amount by  $R$ ) during peak traffic times. However to realize a coded caching scheme, each file must be divided into  $F$  packets which usually increases the computation complexity of a coded caching scheme. So we prefer to construct a caching scheme that decreases the order of  $F$  for practical implementations.

In this paper, we construct four classes of new schemes where two classes can significantly reduce the value of  $F$  by increasing a little  $R$  comparing with the well known scheme proposed by Maddah-Ali and Niesen, and  $F$  in the other two classes grows sub-exponentially with  $K$  by sacrificing more  $R$ . It is worth noting that a tradeoff between  $R$  and  $F$ , which is a hot topic in the field of caching scheme, is proposed by our constructions. In addition, our constructions include all the results constructed by Yan et al., (IEEE Trans. Inf. Theory 63, 5821-5833, 2017) and some main results obtained by Shangquan et al., (arXiv preprint arXiv:1608.03989v1) as the special cases.

## Index Terms

Coded caching scheme, placement delivery array, rate, packet number

## I. INTRODUCTION

Recently, the explosive increasing mobile services, especially applications such as video streaming, have imposed a tremendous pressure on the data transmission over the core network [1]. As a result, during the peak-traffic times, the communication systems are usually congested. Coded caching scheme, which was proposed by Maddah-Ali and Niesen in [12], can effectively reduce congestion during the peak-traffic times, and now is a hot topic in both industrial and academic fields (see [6]–[10], [12]–[14], [25], [28], and references therein).

The benchmark work in [12] focused on the centralized caching system where a single server containing  $N$  files with the same length connects to  $K$  users over a shared link and each user has a cache memory of size  $M$  files. A coded caching scheme consists of two phases: a placement phase during off-peak times and a delivery phase during peak times. In the placement phase, the user caches are populated. This phase does not depend on the user demands which are assumed to be arbitrary. In delivery phase, each user requires a file from server. Then server sends a coded signal of at most  $R$  files to the users such that various user demands are satisfied with the help the local caches. It is meaningful to minimize the load  $R$  files in the delivery phase. Here  $R$  is always called the rate. A coded caching scheme is called  $F$ -division scheme if each file is split into  $F$  packets. If the packets of all files are directly cached in the placement phase, we call it uncoded placement. Otherwise we call it coded placement. Through an elaborate uncoded placement and a coded delivery, the first determined scheme for an  $F$ -division  $(K, M, N)$  coded caching system with  $F = \binom{K}{KM/N}$ , when  $\frac{KM}{N}$  is an integer, was proposed by Maddah-Ali and Niesen in [12]. Such a scheme is referred to as MN scheme in this paper. By means of graph theory, reference [22] showed that MN scheme has minimum rate under the constraint of uncoded cache placement when  $K \leq N$ . So far, many results have been obtained following MN schemes, for instances, [3]–[5], [15], [21], [23], [26] etc.

M. Cheng and J. Jiang are with Guangxi Key Lab of Multi-source Information Mining & Security, Guangxi Normal University, Guilin 541004, China (e-mail: {chengqinshi, jjiang2008}@hotmail.com).

Q. Yan and X. Tang are with the Information Security and National Computing Grid Laboratory, Southwest Jiaotong University, Chengdu, 610031, China (e-mail: qifa@my.swjtu.edu.cn, xhutang@swjtu.edu.cn).

However it is easy to check that  $F$  in MN scheme increases very quickly with the number of users  $K$ . This would become infeasible when  $K$  becomes large. Furthermore, the complexity of a coded caching scheme increases with the parameter  $F$ . Shanmugam et al. in [17] first discussed such subpacketization problem for coded caching schemes. Very recently, Yan et al. in [24] characterized an  $F$ -division  $(K, M, N)$  caching scheme by a simple array which is called  $(K, F, Z, S)$  placement delivery array (PDA), where  $\frac{M}{N} = \frac{Z}{F}$  and  $R = \frac{S}{F}$ . Then they proved that MN scheme is equivalent to a special PDA which is referred to as MN PDA. Furthermore, comparing with MN scheme they obtained two classes of coded caching schemes by means of constructing two infinite classes of PDAs such that  $F$  reduces significantly by increasing the rate very little. Inspired by the concept of a PDA, we construct some coded caching schemes with lower level subpacketization were proposed in [16], [18], [20], [24] from the view point of hypergraphs, resolvable combinatorial designs, Ruzsa-Szemerédi graphs, bipartite graphs respectively. For ease of exposition, the main previously known determined schemes are listed in Table I, which was summarised by Shangguang et al. in [16].

TABLE I: Summary of some known results

References	$K$	$M/N$	$R$	$F$
[12]	$K$	$\frac{z}{q}, z = 1, \dots, q-1$	$\frac{K(q-z)}{zK+q}$	$\binom{K}{Kz/q}$
[20], [24]	$(m+1)q$	$\frac{1}{q}$	$q-1$	$q^m$
	$(m+1)q$	$\frac{q-1}{q}$	$\frac{1}{q-1}$	$(q-1)q^m$
[16]	$\binom{m}{t}q^t$	$1 - (\frac{q-1}{q})^t$	$(q-1)^t$	$q^m$
	$\binom{m}{t}q^t$	$1 - \frac{1}{q^t}$	$\frac{1}{(q-1)^t}$	$(q-1)^t q^m$
[18]	$F^{\Omega(\frac{1}{\log \log F})}$	$\frac{2}{3} + o(1)$	1	$F$

In this paper, we first propose two classes of schemes by directly constructing PDAs, i.e., Theorems 2 and 3. Based on these two classes, another two classes of extended schemes, i.e., Theorems 4 and 5, are obtained. We list these four classes of schemes in Table II.

TABLE II: Main constructions in this paper, where  $z = 1, 2, \dots, q-1$ 

Main results	$K$	$M/N$	$R$	$F$
Theorem 2	$\binom{m}{t}q^t$	$1 - (\frac{q-z}{q})^t$	$(\frac{q-z}{\lfloor \frac{q-1}{q-z} \rfloor})^t$	$\lfloor \frac{q-1}{q-z} \rfloor^t q^m$
Theorem 3	$(m+1)q$	$\frac{z}{q}$	$\frac{q-z}{\lfloor \frac{q-1}{q-z} \rfloor}$	$\lfloor \frac{q-1}{q-z} \rfloor q^m$
Theorem 4	$\binom{m}{t} \lfloor \frac{q-1}{q-z} \rfloor^t q^t$	$1 - (\frac{q-z}{q})^t$	$(q-z)^t$	$q^m$
Theorem 5	$(m \lfloor \frac{q-1}{q-z} \rfloor + 1)q$	$\frac{z}{q}$	$q-z$	$q^m$

From Table II, it is not difficult to check that our schemes have more flexible  $\frac{M}{N}$  than that of the results in [16], [18], [24] and include all the results of [16], [24] listed in Table I as special cases. Because of the flexible memory size, our new schemes have the following two important advantages.

- For the fixed  $K$ ,  $\frac{M}{N}$  we can obtain more different types of schemes directly. For example, given  $K = 405$  and  $\frac{M}{N} = \frac{2}{3}$ . Then from the results in [16] and [18], we have two types of schemes directly, while according to the results in Table II, the following 14 schemes can be obtained.

TABLE III: PDAs obtained by choosing different values  $q$  and  $z$  in Table II

$q$	$z$	$m$	$R$	$\ln F$
3	2	134	0.50	147.9072
3	2	67	1	73.6070
15	10	26	2.50	71.1025
9	6	22	3	48.3389
27	18	14	4.50	46.8349
15	10	13	5	35.2047
45	30	8	7.50	31.1464
27	18	7	9	23.0709
81	54	4	13.5	18.2709
45	30	4	15	15.2266
135	90	2	22.5	10.5037
81	54	2	27	8.7889
135	90	1	45	4.9053

- For many values of  $\frac{M}{N}$ , the schemes in Theorem 2 have both smaller  $F$  and  $R$  than those of the schemes generated by memory sharing based on [16] listed in Table I, and the schemes in Theorems 3 have both smaller  $F$  and  $R$  than those of the schemes generated by memory sharing based on [24] listed in Table I. The detailed discusses are proposed in Subsections III-A and III-B.

The rest of this paper is organized as follows. Section II introduces caching system model, placement delivery array and main result in this paper. In Section III, the detailed performance analyses of Theorems 2 and 3 are proposed. In IV, we first propose the constructions for Theorems 2 and 3 respectively. Based on these two constructions, two extended constructions are proposed for Theorem 4 and 5. Finally conclusion is drawn in Section V.

## II. SYSTEM MODEL AND PLACEMENT DELIVERY ARRAY

In this paper, we use bold capital letter, bold lower case letter and curlicue letter to denote array, vector and set respectively. We use  $[a, b] = \{a, a+1, \dots, b\}$  and  $[a, b) = \{a, a+1, \dots, b-1\}$  for intervals of integers for any integers  $a$  and  $b$  with  $a \leq b$ .

### A. System model

Denote  $N$  files by  $\mathcal{W} = \{W_1, W_2, \dots, W_N\}$  and  $K$  users by  $\mathcal{K} = \{1, 2, \dots, K\}$ . An  $F$ -division  $(K, M, N)$  coded caching scheme proposed by Maddah-Ali and Niesen in [12] operates in two separated phases:

- 1) **Placement Phase:** Each file is sub-divided into  $F$  equal packets<sup>1</sup>, i.e.,  $W_i = \{W_{i,j} : j = 1, 2, \dots, F\}$ . Each user is accessible to the files set  $\mathcal{W}$ . Denote  $\mathcal{Z}_k$  the packets subset of  $\mathcal{W}$  cached by user  $k$ .
- 2) **Delivery Phase:** Each user requests one file from  $\mathcal{W}$  randomly. Denote the request by  $\mathbf{d} = (d_1, d_2, \dots, d_K)$ , i.e., user  $k$  requests file  $W_{d_k}$ , where  $k \in \mathcal{K}, d_k \in \{1, 2, \dots, N\}$ . Once the server receives the request  $\mathbf{d}$ , it broadcasts XOR of packets with size of at most  $R_{\mathbf{d}}F$  to users such that each user is able to recover its requested file.

Denote the maximum transmission amount among all the request during the delivery phase by  $R$ , i.e.,

$$R = \sup_{\substack{\mathbf{d} = (d_1, \dots, d_K) \\ d_k \in [1, N], \forall k \in [1, K]}} \{R_{\mathbf{d}}\}.$$

$R$  is always called the rate of a coded caching scheme.

<sup>1</sup>Memory sharing technique may lead to non equally divided packets [12], in this paper, we will not discuss this case.

## B. Placement delivery array

In this paper we focus on the caching system in the above subsection. Yan et al., in [24] proposed an interesting and simple combinatorial structure, called placement delivery array, which can characterize the placement phase and delivery phase simultaneously.

**Definition 1:** ([24]) For positive integers  $K, F, Z$  and  $S$ , an  $F \times K$  array  $\mathbf{P} = (p_{j,k})$ ,  $1 \leq j \leq F, 1 \leq k \leq K$ , composed of a specific symbol “\*” and  $S$  positive integers  $1, 2, \dots, S$ , is called a  $(K, F, Z, S)$  placement delivery array (PDA) if it satisfies the following conditions:

- C1. The symbol “\*” appears  $Z$  times in each column;
- C2. Each integer occurs at least once in the array;
- C3. For any two distinct entries  $p_{j_1, k_1}$  and  $p_{j_2, k_2}$ ,  $p_{j_1, k_1} = p_{j_2, k_2} = s$  is an integer only if
  - a.  $j_1 \neq j_2, k_1 \neq k_2$ , i.e., they lie in distinct rows and distinct columns; and
  - b.  $p_{j_1, k_2} = p_{j_2, k_1} = *$ , i.e., the corresponding  $2 \times 2$  subarray formed by rows  $j_1, j_2$  and columns  $k_1, k_2$  must be of the following form

$$\begin{pmatrix} s & * \\ * & s \end{pmatrix} \text{ or } \begin{pmatrix} * & s \\ s & * \end{pmatrix}.$$

**Theorem 1:** ([24]) Using Algorithm 1, an  $F$ -division caching scheme for a  $(K, M, N)$  caching system can be realized by a  $(K, F, Z, S)$  PDA with  $\frac{M}{N} = \frac{Z}{F}$ . Each user can decode his requested file correctly for any request  $\mathbf{d}$  at the rate  $R = \frac{S}{F}$ .

---

**Algorithm 1** caching scheme based on PDA in [24]

---

- 1: **procedure** PLACEMENT( $\mathbf{P}, \mathcal{W}$ )
  - 2:   Split each file  $W_i \in \mathcal{W}$  into  $F$  packets, i.e.,  $W_i = \{W_{i,j} \mid j = 1, 2, \dots, F\}$ .
  - 3:   **for**  $k \in \mathcal{K}$  **do**
  - 4:      $\mathcal{Z}_k \leftarrow \{W_{i,j} \mid p_{j,k} = *, \forall i = 1, 2, \dots, N\}$
  - 5:   **end for**
  - 6: **end procedure**
  - 7: **procedure** DELIVERY( $\mathbf{P}, \mathcal{W}, \mathbf{d}$ )
  - 8:   **for**  $s = 1, 2, \dots, S$  **do**
  - 9:     Server sends  $\bigoplus_{p_{j,k}=s, 1 \leq j \leq F, 1 \leq k \leq K} W_{d_k, j}$ .
  - 10:   **end for**
  - 11: **end procedure**
- 

**Example 1:** It is easy to verify that the following array is a  $(4, 6, 3, 4)$  PDA:

$$\mathbf{P}_{6 \times 4} = \begin{pmatrix} * & * & 1 & 2 \\ * & 1 & * & 3 \\ * & 2 & 3 & * \\ 1 & * & * & 4 \\ 2 & * & 4 & * \\ 3 & 4 & * & * \end{pmatrix}.$$

Using Algorithm 1, one can obtain a 6-division  $(4, 3, 6)$  coded caching scheme in the following way.

- **Placement Phase:** From Line 2 we have  $W_i = \{W_{i,1}, W_{i,2}, W_{i,3}, W_{i,4}, W_{i,5}, W_{i,6}\}$ ,  $i \in [1, 6]$ . Then by Lines 3-5, the contents in each user are

$$\begin{aligned} \mathcal{Z}_1 &= \{W_{i,1}, W_{i,2}, W_{i,3} : i \in [1, 6]\} & \mathcal{Z}_2 &= \{W_{i,1}, W_{i,4}, W_{i,5} : i \in [1, 6]\} \\ \mathcal{Z}_3 &= \{W_{i,2}, W_{i,4}, W_{i,6} : i \in [1, 6]\} & \mathcal{Z}_4 &= \{W_{i,3}, W_{i,5}, W_{i,6} : i \in [1, 6]\} \end{aligned}$$

- **Delivery Phase:** Assume that the request vector is  $\mathbf{d} = (1, 2, 3, 4, 5, 6)$ . Table IV shows the transmitting process by Lines 8-10.

Time Slot	Transmitted Signal
1	$W_{1,4} \oplus W_{2,2} \oplus W_{3,1}$
2	$W_{1,5} \oplus W_{2,3} \oplus W_{4,1}$
3	$W_{1,6} \oplus W_{3,3} \oplus W_{4,2}$
4	$W_{2,6} \oplus W_{3,5} \oplus W_{4,4}$

TABLE IV: Delivery steps in Example 1

From Theorem 1, an  $F$ -division  $(K, M, N)$  coded caching scheme can be obtained by constructing an appropriate PDA. In this paper, we only consider the caching schemes with  $K \leq N$ .

### C. Main Results

In this paper, we directly construct the following two classes of PDAs.

**Theorem 2:** For any positive integers  $q, z, m$  and  $t$  with  $q \geq 2, z < q$  and  $t < m$ , the array  $\mathbf{P}$  given in Construction 1 is an  $((\binom{m}{t}q^t, \lfloor \frac{q-1}{q-z} \rfloor^t q^m, \lfloor \frac{q-1}{q-z} \rfloor^t (q^m - q^{m-t}(q-z)^t), (q-z)^t q^m)$  PDA with  $\frac{M}{N} = 1 - (\frac{q-z}{q})^t$  and rate  $R = (q-z)/\lfloor \frac{q-1}{q-z} \rfloor^t$ .

**Theorem 3:** For any positive integers  $q, z, m$  with  $q \geq 2$  and  $z < q$ , the array  $\mathbf{H}$  generated by Construction 2 is an  $((m+1)q, \lfloor \frac{q-1}{q-z} \rfloor q^m, z \lfloor \frac{q-1}{q-z} \rfloor q^{m-1}, (q-z)q^m)$  PDA with  $\frac{M}{N} = \frac{z}{q}$  and rate  $R = (q-z)/\lfloor \frac{q-1}{q-z} \rfloor$ .

Based on the above two results, the following extended results can be obtained.

**Theorem 4:** For any positive integers  $q, z, m$  and  $t$  with  $q \geq 2, z < q$  and  $t < m$ , there exists an  $((\binom{m}{t}q^t \lfloor \frac{q-1}{q-z} \rfloor^t, q^m, q^m - (q-z)^t q^{m-t}, (q-z)^t q^m)$  PDA with  $\frac{M}{N} = 1 - (\frac{q-z}{q})^t$  and rate  $R = (q-z)^t$ .

**Theorem 5:** For any positive integers  $q, z, m$  with  $q \geq 2$  and  $z < q$ , the array  $(\mathbf{P}, \mathbf{C})$  generated by Construction 4 is an  $((m \lfloor \frac{q-1}{q-z} \rfloor + 1)q, q^m, zq^{m-1}, (q-z)q^m)$  PDA with  $\frac{M}{N} = \frac{z}{q}$  and rate  $R = q-z$ .

## III. PERFORMANCE ANALYSES

### A. Performance analyses of Theorem 2

First we claim that  $F$  of the PDA in Theorem 2 grows sub-exponentially with  $K$  when  $t \geq 2$ . Let  $q, z$  and  $t$  be fixed and let  $m$  approximate infinity, it holds that  $R$  and  $\frac{M}{N}$  are constants independent of  $K$  for all PDAs in Theorem 2. Similar to the discussion in [16], by means of the inequality  $(m/t)^t < \binom{m}{t} < (em/t)^t$ , we can estimate the value of  $m$  by  $K$ , i.e.,

$$\frac{tK^{\frac{1}{t}}}{eq} < m < \frac{tK^{\frac{1}{t}}}{q}. \quad (1)$$

So we have  $F = \mathcal{O}(\lfloor \frac{q-1}{q-z} \rfloor^t q^{\frac{tK^{1/t}}{q}})$ . This implies that  $F$  grows sub-exponentially with  $K$  if  $t \geq 2$ .

Secondly let us consider the performance in Theorem 2 comparing the result in [16] listed in Table I. When  $\frac{M}{N} = 1 - (\frac{q-1}{q})^t$ ,  $1 - \frac{1}{q^t}$ , from Theorem 2 we have

- a  $((\binom{m}{t}q^t, q^m, q^m - q^{m-t}(q-1)^t, (q-1)^t q^m)$  PDA with  $\frac{M}{N} = 1 - (\frac{q-1}{q})^t$  and rate  $R = (q-1)^t$  and
- a  $((\binom{m}{t}q^t, (q-1)^t q^m, (q-1)^t (q^m - q^{m-t}), q^m)$  PDA with  $\frac{M}{N} = 1 - \frac{1}{q^t}$  and rate  $R = \frac{1}{(q-1)^t}$ .

Clearly these two PDAs are exactly the results in [16] listed in Table I. So we only need to consider the positive real number  $1 - (\frac{q-1}{q})^t < \frac{M}{N} < 1 - \frac{1}{q^t}$ . Let  $K = \binom{m}{t}q^t$ . Based on the schemes in [16], we can obtain the schemes with  $K$  users and memory ratio  $\frac{M}{N}$  by memory sharing. Now let us introduce memory sharing method first. Assume that there exist  $r$  schemes with  $(\frac{M_i}{N}, R_i, F_i), i = 1, 2, \dots, r$ . [12] showed that a scheme with

$$\begin{aligned} \frac{M}{N} &= \lambda_1 \frac{M_1}{N} + \lambda_2 \frac{M_2}{N} + \dots + \lambda_r \frac{M_r}{N} \\ R_M &= \lambda_1 R_1 + \lambda_2 R_2 + \dots + \lambda_r R_r \\ F_M &= F_1 + F_2 + \dots + F_r \end{aligned} \quad (2)$$

can be obtained where  $M_1 \leq M \leq M_r$ ,  $0 < \lambda_i \leq 1$  and  $\sum_{i=1}^r \lambda_i = 1$ . From the results in [16] of Table I, there exist two schemes with

$$\left(\frac{M_1}{N}, R_1, F_1\right) = \left(1 - \left(\frac{q-1}{q}\right)^t, (q-1)^t, q^m\right) \quad \text{and} \quad \left(\frac{M_2}{N}, R_2, F_2\right) = \left(1 - \frac{1}{q^t}, \frac{1}{(q-1)^t}, (q-1)^t q^m\right). \quad (3)$$

So the scheme in (2) based on the schemes in (3) has the parameters

$$\begin{aligned} \frac{M}{N} &= \lambda \left(1 - \left(\frac{q-1}{q}\right)^t\right) + (1-\lambda) \left(1 - \frac{1}{q^t}\right) \\ R_{M-S} &= \lambda(q-1)^t + (1-\lambda) \frac{1}{(q-1)^t} \\ F_{M-S} &= (q-1)^t q^m + q^m \end{aligned} \quad (4)$$

for some positive real number  $0 < \lambda < 1$ . From Theorem 2, there exist  $q-1$  schemes with

$$\left(\frac{M_z}{N}, R_z, F_z\right) = \left(1 - \left(\frac{q-z}{q}\right)^t, \left(\frac{q-z}{\lfloor \frac{q-1}{q-z} \rfloor}\right)^t, \lfloor \frac{q-1}{q-z} \rfloor^t q^m\right), \quad z = 1, 2, \dots, q-1. \quad (5)$$

If there exists  $z$ ,  $1 \leq z \leq q-1$ , satisfying  $\frac{M}{N} = 1 - \left(\frac{q-z}{q}\right)^t$ , by (4) and (5) we have

$$\begin{aligned} \frac{R_z}{R_{M-S}} &= \frac{\left(\frac{q-z}{\lfloor \frac{q-1}{q-z} \rfloor}\right)^t}{\lambda(q-1)^t + (1-\lambda) \frac{1}{(q-1)^t}} = \frac{1}{\left(\lambda \left(\frac{q-1}{q-z}\right)^t + \frac{1-\lambda}{(q-1)^t (q-z)^t}\right) \lfloor \frac{q-1}{q-z} \rfloor^t} \\ &< \frac{1}{\lambda \left(\frac{q-1}{q-z}\right)^t \lfloor \frac{q-1}{q-z} \rfloor^t} < \frac{1}{\lambda \lfloor \frac{q-1}{q-z} \rfloor^{2t}} \end{aligned} \quad (6)$$

and

$$\begin{aligned} \frac{F_z}{F_{M-S}} &= \frac{\lfloor \frac{q-1}{q-z} \rfloor^t q^m}{(q-1)^t q^m + q^m} = \frac{\lfloor \frac{q-1}{q-z} \rfloor^t}{(q-1)^t + 1} = \frac{1}{\left(\frac{q-1}{\lfloor \frac{q-1}{q-z} \rfloor}\right)^t + \frac{1}{\lfloor \frac{q-1}{q-z} \rfloor^t}} \\ &< \frac{1}{(q-z)^t + \frac{1}{\lfloor \frac{q-1}{q-z} \rfloor^t}} < \frac{1}{(q-z)^t}. \end{aligned} \quad (7)$$

**Remark 1:** From (6) and (7), when  $\lambda \lfloor \frac{q-1}{q-z} \rfloor^{2t} > 1$ ,  $R_z$  and  $F_z$  are at least  $\frac{1}{\lambda \lfloor \frac{q-1}{q-z} \rfloor^{2t}}$  and  $\frac{1}{(q-z)^t}$  times smaller than  $R_{M-S}$  and  $F_{M-S}$  respectively.

**Example 2:** When  $t = 3$ ,  $q = 20$  and  $\lambda = 0.1$ , the following table can be obtained by (6) and (7).

TABLE V: Comparisons of the performances between schemes from [16] and Theorem 2

$z$	$\frac{R_z}{R_{M-S}} <$	$\frac{F_z}{F_{M-S}}$
11	0.15625	0.00137174211248285
12	0.15625	0.001953125
13	0.15625	0.00291545189504373
14	0.0137174211248285	0.00462962962962963
15	0.0137174211248285	0.008
16	0.00244140625	0.015625
17	0.000214334705075446	0.037037037037037
18	1.88167642315892e-05	0.125

If there is no integer  $z$  satisfying  $\frac{M}{N} = 1 - \left(\frac{q-z}{q}\right)^t$ , a scheme in (2) based on the schemes in (5) has the parameters

$$\begin{aligned} \frac{M}{N} &= \lambda' \left(1 - \left(\frac{q-z}{q}\right)^t\right) + (1-\lambda') \left(1 - \left(\frac{q-z-1}{q}\right)^t\right) \\ R_{M-C_1} &= \lambda' \left(\frac{q-z}{\lfloor \frac{q-1}{q-z} \rfloor}\right)^t + (1-\lambda') \left(\frac{q-z-1}{\lfloor \frac{q-1}{q-z-1} \rfloor}\right)^t < \left(\frac{q-z}{\lfloor \frac{q-1}{q-z} \rfloor}\right)^t \\ F_{M-C_1} &= \lfloor \frac{q-1}{q-z} \rfloor^t q^m + \lfloor \frac{q-1}{q-z-1} \rfloor^t q^m \end{aligned} \quad (8)$$

for some positive integer  $z$  with  $1 - \left(\frac{q-z}{q}\right)^t < \frac{M}{N} < 1 - \left(\frac{q-z-1}{q}\right)^t$  and a real number  $0 < \lambda' < 1$ . Here the last item in the second formula of (8) holds since  $\left(\frac{q-z}{\lfloor \frac{q-1}{q-z} \rfloor}\right)^x > \left(\frac{q-z-1}{\lfloor \frac{q-1}{q-z-1} \rfloor}\right)^x$  always holds for any positive integer  $x$ . Similar to (6) and (7), we have

$$\frac{R_{M-C_1}}{R_{M-S}} < \frac{\left(\frac{q-z}{\lfloor \frac{q-1}{q-z} \rfloor}\right)^t}{\lambda(q-1)^t + (1-\lambda)\frac{1}{(q-1)^t}} < \frac{1}{\lambda \lfloor \frac{q-1}{q-z} \rfloor^{2t}} \quad (9)$$

and

$$\begin{aligned} \frac{F_{M-C_1}}{F_{M-S}} &= \frac{\lfloor \frac{q-1}{q-z} \rfloor^t q^m + \lfloor \frac{q-1}{q-z-1} \rfloor^t q^m}{(q-1)^t q^m + q^m} = \frac{\lfloor \frac{q-1}{q-z} \rfloor^t + \lfloor \frac{q-1}{q-z-1} \rfloor^t}{(q-1)^t + 1} = \frac{\lfloor \frac{q-1}{q-z} \rfloor^t}{(q-1)^t + 1} + \frac{\lfloor \frac{q-1}{q-z-1} \rfloor^t}{(q-1)^t + 1} \\ &< \frac{1}{(q-z)^t} + \frac{1}{(q-z-1)^t}. \end{aligned} \quad (10)$$

Similar to the Remark 1, we have that when  $\lambda \lfloor \frac{q-1}{q-z} \rfloor^{2t} > 1$ ,  $R_{M-C_1}$  and  $F_{M-C_1}$  are at least  $\frac{1}{\lambda \lfloor \frac{q-1}{q-z} \rfloor^{2t}}$  and  $\frac{1}{(q-z)^t} + \frac{1}{(q-z-1)^t}$  times smaller than  $R_{M-S}$  and  $F_{M-S}$  respectively.

### B. Performance analyses of Theorem 3

For any positive integers  $m$  and  $q$ , assume that  $K = (m+1)q$ . Similar to the discussion in Subsection III-A, we also first consider the case  $\frac{M}{N} = \frac{1}{q}$  and  $\frac{q-1}{q}$ . From Theorem 3 we have

- an  $((m+1)q, q^m, q^{m-1}, (q-1)q^m)$  PDA with  $\frac{M}{N} = \frac{1}{q}$  and  $R = q-1$ , and
- an  $((m+1)q, (q-1)q^m, (q-1)^2 q^{m-1}, q^m)$  PDA with  $\frac{M}{N} = 1 - \frac{1}{q}$  and rate  $R = \frac{1}{q-1}$ .

Clearly these two PDAs are exactly the results in [24] listed in Table I. It is worth to note that the authors in [24] showed that comparing with MN PDA, the packet number  $F$  of these two PDAs in Theorem 3 reduces significantly while the rate  $R$  increases little. Now let us consider the case  $\frac{1}{q} < \frac{M}{N} < \frac{q-1}{q}$ .

From the results in [24] of Table I, there exist two schemes with

$$\left(\frac{M_1}{N}, R_1, F_1\right) = \left(\frac{1}{q}, q-1, q^m\right) \quad \text{and} \quad \left(\frac{M_2}{N}, R_2, F_2\right) = \left(1 - \frac{1}{q}, \frac{1}{q-1}, (q-1)q^m\right). \quad (11)$$

So the scheme in (2) based on the schemes in (3) has the parameters

$$\begin{aligned} \frac{M}{N} &= \lambda \frac{1}{q} + (1-\lambda) \left(1 - \frac{1}{q}\right) \\ R_{M-Y} &= \lambda(q-1) + (1-\lambda) \frac{1}{q-1} \\ F_{M-Y} &= (q-1)q^m + q^m \end{aligned} \quad (12)$$

for some positive real number  $0 < \lambda < 1$ . From Theorem 2, there exist  $q-1$  schemes with

$$\left(\frac{M_z}{N}, R_z, F_z\right) = \left(\frac{z}{q}, \frac{q-z}{\lfloor \frac{q-1}{q-z} \rfloor}, \lfloor \frac{q-1}{q-z} \rfloor q^m\right), \quad z = 1, 2, \dots, q-1. \quad (13)$$

If there exists  $z$ ,  $1 \leq z \leq q-1$ , satisfying  $\frac{M}{N} = \frac{z}{q}$ , by (12) and (13) we have

$$\begin{aligned} \frac{R_z}{R_{M-Y}} &= \frac{\frac{q-z}{\lfloor \frac{q-1}{q-z} \rfloor}}{\lambda(q-1) + (1-\lambda)\frac{1}{(q-1)}} = \frac{1}{\left(\frac{\lambda(q-1)}{q-z} + \frac{1-\lambda}{(q-1)(q-z)}\right) \lfloor \frac{q-1}{q-z} \rfloor} \\ &< \frac{1}{\lambda \left(\frac{q-1}{q-z}\right) \lfloor \frac{q-1}{q-z} \rfloor} < \frac{1}{\lambda \lfloor \frac{q-1}{q-z} \rfloor^2} \end{aligned} \quad (14)$$

and

$$\frac{F_z}{F_{M-Y}} = \frac{\lfloor \frac{q-1}{q-z} \rfloor q^m}{(q-1)q^m + q^m} = \frac{\lfloor \frac{q-1}{q-z} \rfloor}{(q-1)+1} = \frac{\lfloor \frac{q-1}{q-z} \rfloor}{q}. \quad (15)$$

**Remark 2:** From (14) and (15), when  $\lambda \lfloor \frac{q-1}{q-z} \rfloor^2 > 1$ ,  $R_z$  and  $F_z$  are at least  $\frac{1}{\lambda \lfloor \frac{q-1}{q-z} \rfloor^2}$  and  $\frac{\lfloor \frac{q-1}{q-z} \rfloor}{q}$  times smaller than  $R_{M-Y}$  and  $F_{M-Y}$  respectively.

**Example 3:** When  $q = 20$  and  $\lambda = 0.5$ , the following table can be obtained by (14) and (15).

TABLE VI: Comparisons of the performances between scheme from [24] and Theorem 3

$z$	$\frac{R_z}{R_{M-S}} <$	$\frac{F_z}{F_{M-S}}$
11	0.5	0.1
12	0.5	0.1
13	0.5	0.1
14	0.2222222222222222	0.15
15	0.2222222222222222	0.15
16	0.125	0.2
17	0.0555555555555556	0.3
18	0.0246913580246914	0.45

If there is no integer  $z$  satisfying  $\frac{M}{N} = \frac{z}{q}$ , a scheme in (2) based on the schemes in (13) has the parameters

$$\begin{aligned} \frac{M}{N} &= \lambda' \frac{z}{q} + (1 - \lambda') \frac{z+1}{q} \\ R_{M-C_2} &= \lambda' \frac{q-z}{\lfloor \frac{q-1}{q-z} \rfloor} + (1 - \lambda') \frac{q-z-1}{\lfloor \frac{q-1}{q-z-1} \rfloor} < \frac{q-z}{\lfloor \frac{q-1}{q-z} \rfloor} \\ F_{M-C_2} &= \lfloor \frac{q-1}{q-z} \rfloor q^m + \lfloor \frac{q-1}{q-z-1} \rfloor q^m \end{aligned} \quad (16)$$

for some positive integer  $z$  with  $\frac{z}{q} < \frac{M}{N} < \frac{z+1}{q}$  and a real number  $0 < \lambda < 1$ . Similar to (14) and (15), we have

$$\frac{R_{M-C_2}}{R_{M-S}} < \frac{\frac{q-z}{\lfloor \frac{q-1}{q-z} \rfloor}}{\lambda(q-1) + (1-\lambda)\frac{1}{q-1}} < \frac{1}{\lambda \lfloor \frac{q-1}{q-z} \rfloor^2} \quad (17)$$

and

$$\begin{aligned} \frac{F_{M-C_2}}{F_{M-S}} &= \frac{\lfloor \frac{q-1}{q-z} \rfloor q^m + \lfloor \frac{q-1}{q-z-1} \rfloor q^m}{(q-1)q^m + q^m} = \frac{\lfloor \frac{q-1}{q-z} \rfloor + \lfloor \frac{q-1}{q-z-1} \rfloor}{(q-1) + 1} = \frac{\lfloor \frac{q-1}{q-z} \rfloor}{(q-1) + 1} + \frac{\lfloor \frac{q-1}{q-z-1} \rfloor}{(q-1) + 1} \\ &< \frac{1}{q-z} + \frac{1}{q-z-1}. \end{aligned} \quad (18)$$

Similar to the Remark 2, we have that when  $\lambda \lfloor \frac{q-1}{q-z} \rfloor^2 > 1$  and  $q-z > 1$ ,  $R_{M-C_2}$  and  $F_{M-C_2}$  are at least  $\frac{1}{\lambda \lfloor \frac{q-1}{q-z} \rfloor^2}$  and  $\frac{1}{q-z} + \frac{1}{q-z-1}$  times smaller than  $R_{M-Y}$  and  $F_{M-Y}$  respectively.

#### IV. EXPLICIT CONSTRUCTIONS

In this section, four classes of PDAs in Table II are constructed.

##### A. The first construction

First let us consider an example. When  $m = 2$ ,  $t = 1$  and  $q = 3$ , from the construction in [16], the following  $(6, 9, 3, 9)$  PDA can be obtained.

$$\mathbf{P} = \left( \begin{array}{ccc|ccc} * & 11 & 3 & * & 13 & 7 \\ 1 & * & 12 & * & 14 & 8 \\ 10 & 2 & * & * & 15 & 9 \\ * & 14 & 6 & 1 & * & 16 \\ 4 & * & 15 & 2 & * & 17 \\ 13 & 5 & * & 3 & * & 18 \\ * & 17 & 9 & 10 & 4 & * \\ 7 & * & 18 & 11 & 5 & * \\ 16 & 8 & * & 12 & 6 & * \end{array} \right) \quad (19)$$



For the detailed constructions, the interested readers can be referred to [16]. Replacing three integer entries of each column of  $\mathbf{P}$  in (19) by “\*”s elaborately, we have  $\mathbf{P}_1$  in the following.

$$\mathbf{P}_1 = \left( \begin{array}{ccc|ccc} * & * & 3 & * & * & 7 \\ 1 & * & * & * & * & 8 \\ * & 2 & * & * & * & 9 \\ * & * & 6 & 1 & * & * \\ 4 & * & * & 2 & * & * \\ * & 5 & * & 3 & * & * \\ * & * & 9 & * & 4 & * \\ 7 & * & * & * & 5 & * \\ * & 8 & * & * & 6 & * \end{array} \right)$$

Then changing the integer in each entry, the following array  $\mathbf{P}_2$  can be obtained.

$$\mathbf{P}_2 = \left( \begin{array}{ccc|ccc} * & * & 2 & * & * & 4 \\ 3 & * & * & * & * & 5 \\ * & 1 & * & * & * & 6 \\ * & * & 5 & 7 & * & * \\ 6 & * & * & 8 & * & * \\ * & 4 & * & 9 & * & * \\ * & * & 8 & * & 1 & * \\ 9 & * & * & * & 2 & * \\ * & 7 & * & * & 3 & * \end{array} \right)$$

It is interesting that the following array  $\mathbf{P}' = \begin{pmatrix} \mathbf{P}_1 \\ \mathbf{P}_2 \end{pmatrix}$  is a  $(6, 18, 12, 9)$  PDA too.

$$\mathbf{P}' = \left( \begin{array}{ccc|ccc} * & * & 3 & * & * & 7 \\ 1 & * & * & * & * & 8 \\ * & 2 & * & * & * & 9 \\ * & * & 6 & 1 & * & * \\ 4 & * & * & 2 & * & * \\ * & 5 & * & 3 & * & * \\ * & * & 9 & * & 4 & * \\ 7 & * & * & * & 5 & * \\ * & 8 & * & * & 6 & * \\ \hline * & * & 2 & * & * & 4 \\ 3 & * & * & * & * & 5 \\ * & 1 & * & * & * & 6 \\ * & * & 5 & 7 & * & * \\ 6 & * & * & 8 & * & * \\ * & 4 & * & 9 & * & * \\ * & * & 8 & * & 1 & * \\ 9 & * & * & * & 2 & * \\ * & 7 & * & * & 3 & * \end{array} \right) \quad (20)$$

Inspired by replacing method in the above example, we can propose our first construction. For ease of understanding, we will use  $q$ -ary sequences to represent the parameters  $K$ ,  $F$ ,  $Z$  and  $S$  of a PDA unless otherwise stated.

**Construction 1:** For any positive integers  $q$ ,  $z$ ,  $m$  and  $t$  with  $0 < z < q$  and  $0 < t < m$ , let

$$\begin{aligned} \mathcal{F} &= \{ \mathbf{a} = (a_0, a_1, \dots, a_{m-1}, \varepsilon_0, \dots, \varepsilon_{t-1}) \mid a_0, \dots, a_{m-1} \in \mathbb{Z}_q, \varepsilon_0, \dots, \varepsilon_{t-1} \in [0, \lfloor \frac{q-1}{q-z} \rfloor] \}, \\ \mathcal{K} &= \{ \mathbf{b} = (b_0, b_1, \dots, b_{t-1}, \delta_0, \delta_1, \dots, \delta_{t-1}) \mid b_0, \dots, b_{t-1} \in \mathbb{Z}_q, 0 \leq \delta_0 < \dots < \delta_{t-1} < m, \}. \end{aligned} \quad (21)$$

Then a  $\lfloor \frac{q-1}{q-z} \rfloor^t q^m \times \binom{m}{t} q^t$  array  $\mathbf{P} = (p_{\mathbf{a}, \mathbf{b}})$  can be defined in the following way

$$p_{\mathbf{a}, \mathbf{b}} = \begin{cases} (a_0, a_1, \dots, b_i - \varepsilon_i(q-z), \dots, a_{m-1}, a_{\delta_0} - b_0 - 1, \dots, a_{\delta_{t-1}} - b_{t-1} - 1)_q & \text{if } a_{\delta_i} \notin X_{b_i, z}, \forall i \in [0, t) \\ * & \text{otherwise} \end{cases} \quad (22)$$

where  $X_{b_i, z} = \{b_i, b_i - 1, \dots, b_i - (z - 1)\}$ . Here the operations are performed modulo  $q$ .

Based on the array generated by Construction 1, the proof of Theorem 2 can be obtained and is included in Appendix A.

**Example 4:** Assume that  $m = 2$ ,  $q = 3$  and  $t = 1$ . When  $z = 2$ ,  $\lfloor \frac{q-1}{q-z} \rfloor = 2$  holds. Then (21) can be written as follows.

$$\mathcal{F} = \{(a_0, a_1, \varepsilon) \mid a_0, a_1 \in \mathbb{Z}_3, \varepsilon \in [0, 1]\} \quad \text{and} \quad \mathcal{K} = \{(b_0, \delta_0) \mid b_0 \in \mathbb{Z}_3, 0 \leq \delta_0 \leq 1\}$$

By (22), the following array can be obtained.

$(a_0, a_1, \varepsilon) \setminus (b_0, \delta_0)$	(0, 0)	(1, 0)	(2, 0)	(0, 1)	(1, 1)	(2, 1)
(0, 0, 0)	*	*	(2, 0, 0)	*	*	(0, 2, 0)
(1, 0, 0)	(0, 0, 0)	*	*	*	*	(1, 2, 0)
(2, 0, 0)	*	(1, 0, 0)	*	*	*	(2, 2, 0)
(0, 1, 0)	*	*	(2, 1, 0)	(0, 0, 0)	*	*
(1, 1, 0)	(0, 1, 0)	*	*	(1, 0, 0)	*	*
(2, 1, 0)	*	(1, 1, 0)	*	(2, 0, 0)	*	*
(0, 2, 0)	*	*	(2, 2, 0)	*	(0, 1, 0)	*
(1, 2, 0)	(0, 2, 0)	*	*	*	(1, 1, 0)	*
(2, 2, 0)	*	(1, 2, 0)	*	*	(2, 1, 0)	*
(0, 0, 1)	*	*	(1, 0, 0)	*	*	(0, 1, 0)
(1, 0, 1)	(2, 0, 0)	*	*	*	*	(1, 1, 0)
(2, 0, 1)	*	(0, 0, 0)	*	*	*	(2, 1, 0)
(0, 1, 1)	*	*	(1, 1, 0)	(0, 2, 0)	*	*
(1, 1, 1)	(2, 1, 0)	*	*	(1, 2, 0)	*	*
(2, 1, 1)	*	(0, 1, 0)	*	(2, 2, 0)	*	*
(0, 2, 1)	*	*	(1, 2, 0)	*	(0, 0, 0)	*
(1, 2, 1)	(2, 2, 0)	*	*	*	(1, 0, 0)	*
(2, 2, 1)	*	(0, 2, 0)	*	*	(2, 0, 0)	*

(23)

In fact the array in (23) is exactly the PDA in (20) when we replace the vectors by the integers in  $[1, 9]$ .

### B. The second construction

Let us also consider an example. When  $m = 2$  and  $q = 3$ , from the construction in [24], the following  $(9, 9, 3, 9)$  PDA can be obtained.

$$\mathbf{P} = \left( \begin{array}{ccc|ccc|ccc} * & 11 & 3 & * & 13 & 7 & * & 1 & 10 \\ 1 & * & 12 & * & 14 & 8 & 11 & * & 2 \\ 10 & 2 & * & * & 15 & 9 & 3 & 12 & * \\ * & 14 & 6 & 1 & * & 16 & 13 & * & 4 \\ 4 & * & 15 & 2 & * & 17 & 5 & 14 & * \\ 13 & 5 & * & 3 & * & 18 & * & 6 & 15 \\ * & 17 & 9 & 10 & 4 & * & 7 & 16 & * \\ 7 & * & 18 & 11 & 5 & * & * & 8 & 17 \\ 16 & 8 & * & 12 & 6 & * & 18 & * & 9 \end{array} \right) \quad (24)$$

For the detailed constructions, the interested readers can be referred to [24]. Similar to generating the array in (20), the following (9, 18, 12, 9) PDA can be obtained.

$$\mathbf{P}' = \begin{pmatrix} * & * & 3 & * & * & 7 & * & 1 & * \\ 1 & * & * & * & * & 8 & * & * & 2 \\ * & 2 & * & * & * & 9 & 3 & * & * \\ * & * & 6 & 1 & * & * & * & * & 4 \\ 4 & * & * & 2 & * & * & 5 & * & * \\ * & 5 & * & 3 & * & * & * & 6 & * \\ * & * & 9 & * & 4 & * & 7 & * & * \\ 7 & * & * & * & 5 & * & * & 8 & * \\ * & 8 & * & * & 6 & * & * & * & 9 \\ \hline * & * & 2 & * & * & 4 & 1 & * & * \\ 3 & * & * & * & * & 5 & * & 2 & * \\ * & 1 & * & * & * & 6 & * & * & 3 \\ * & * & 5 & 7 & * & * & * & 4 & * \\ 6 & * & * & 8 & * & * & * & * & 5 \\ * & 4 & * & 9 & * & * & 6 & * & * \\ * & * & 8 & * & 1 & * & * & * & 7 \\ 9 & * & * & * & 2 & * & 8 & * & * \\ * & 7 & * & * & 3 & * & * & 9 & * \end{pmatrix} \quad (25)$$

It is worth noting that the replacing method in (25) is more complex than that in (20) since the number of columns of the array in (24) is greater than that of the array in (19). However we can still propose our second construction for any positive integers  $m$ ,  $q$  and  $z$ .

**Construction 2:** When  $t = 1$ , (21) can be written as

$$\mathcal{F} = \{(a_0, a_1, \dots, a_{m-1}, \varepsilon) \mid a_0, \dots, a_{m-1} \in \mathbb{Z}_q, \varepsilon \in [0, \lfloor \frac{q-1}{q-z} \rfloor]\} \quad \text{and} \quad \mathcal{K} = \{(b, \delta) \mid b \in \mathbb{Z}_q, 0 \leq \delta < m\}$$

for any positive integers  $q$ ,  $z$  and  $m$  with  $z < q$ . Let  $\mathcal{K}_1 = \{(b, m) \mid b \in \mathbb{Z}_q\}$ . Define an array  $\mathbf{H} = (\mathbf{P}, \mathbf{C})$  where

- $\mathbf{P} = (p_{\mathbf{a}, \mathbf{b}})$ ,  $\mathbf{a} \in \mathcal{F}$ ,  $\mathbf{b} \in \mathcal{K}$ , is the  $\lfloor \frac{q-1}{q-z} \rfloor q^m \times mq$  array generated by Construction 1.
- $\mathbf{C} = (c_{\mathbf{a}, \mathbf{b}})$ ,  $\mathbf{a} \in \mathcal{F}$ ,  $\mathbf{b} \in \mathcal{K}_1$  is a  $\lfloor \frac{q-1}{q-z} \rfloor q^m \times q$  array define in the following way.

$$c_{\mathbf{a}, \mathbf{b}} = \begin{cases} (a_0, \dots, a_{m-1}, b - (\sum_{l=0}^{m-1} a_l - \varepsilon(q-z)) - 1)_q & \text{if } \sum_{l=0}^{m-1} a_l - \varepsilon(q-z) \notin Y_{b,z} \\ * & \text{otherwise} \end{cases} \quad (26)$$

where  $Y_{b,z} = \{b, b+1, \dots, b+(z-1)\}$ . All the above operations are performed modulo  $q$ .

Based on the array generated by Construction 2, the proof of Theorem 3 can be obtained and is included in Appendix B.

**Example 5:** Suppose that  $m = 2$ ,  $q = 3$  and  $z = 2$ . Then we have  $\mathcal{K}_1 = \{(b, 2) \mid b \in \mathbb{Z}_3\}$ . By (22) and (26), the following array can be obtained.

$(a_0, a_1, \varepsilon) \setminus (b, \delta)$	(0, 0)	(1, 0)	(2, 0)	(0, 1)	(1, 1)	(2, 1)	(0, 2)	(1, 2)	(2, 2)
(0, 0, 0)	*	*	(2, 0, 0)	*	*	(0, 2, 0)	*	(0, 0, 0)	*
(1, 0, 0)	(0, 0, 0)	*	*	*	*	(1, 2, 0)	*	*	(1, 0, 0)
(2, 0, 0)	*	(1, 0, 0)	*	*	*	(2, 2, 0)	(2, 0, 0)	*	*
(0, 1, 0)	*	*	(2, 1, 0)	(0, 0, 0)	*	*	*	*	(0, 1, 0)
(1, 1, 0)	(0, 1, 0)	*	*	(1, 0, 0)	*	*	(1, 1, 0)	*	*
(2, 1, 0)	*	(1, 1, 0)	*	(2, 0, 0)	*	*	*	(2, 1, 0)	*
(0, 2, 0)	*	*	(2, 2, 0)	*	(0, 1, 0)	*	(0, 2, 0)	*	*
(1, 2, 0)	(0, 2, 0)	*	*	*	(1, 1, 0)	*	*	(1, 2, 0)	*
(2, 2, 0)	*	(1, 2, 0)	*	*	(2, 1, 0)	*	*	*	(2, 2, 0)
(0, 0, 1)	*	*	(1, 0, 0)	*	*	(0, 1, 0)	(0, 0, 0)	*	*
(1, 0, 1)	(2, 0, 0)	*	*	*	*	(1, 1, 0)	*	(1, 0, 0)	*
(2, 0, 1)	*	(0, 0, 0)	*	*	*	(2, 1, 0)	*	*	(2, 0, 0)
(0, 1, 1)	*	*	(1, 1, 0)	(0, 2, 0)	*	*	*	(0, 1, 0)	*
(1, 1, 1)	(2, 1, 0)	*	*	(1, 2, 0)	*	*	*	*	(1, 1, 0)
(2, 1, 1)	*	(0, 1, 0)	*	(2, 2, 0)	*	*	(2, 1, 0)	*	*
(0, 2, 1)	*	*	(1, 2, 0)	*	(0, 0, 0)	*	*	*	(0, 2, 0)
(1, 2, 1)	(2, 2, 0)	*	*	*	(1, 0, 0)	*	(1, 2, 0)	*	*
(2, 2, 1)	*	(0, 2, 0)	*	*	(2, 0, 0)	*	*	(2, 2, 0)	*

The array in (27) is exactly the PDA in (25) when we replace the vectors by the integers in  $[1, 9]$ .

### C. The first extended construction

It is interesting that we can obtain the following (12, 9, 6, 9) PDA  $\mathbf{P}''$  by copying or replacing three integer entries of each column of  $\mathbf{P}$  in (19) by “\*”s elaborately.

$$\mathbf{P}'' = \begin{pmatrix} * & * & 3 & * & * & 2 & * & * & 7 & * & * & 4 \\ 1 & * & * & 3 & * & * & * & * & 8 & * & * & 5 \\ * & 2 & * & * & 1 & * & * & * & 9 & * & * & 6 \\ * & * & 6 & * & * & 5 & 1 & * & * & 7 & * & * \\ 4 & * & * & 6 & * & * & 2 & * & * & 8 & * & * \\ * & 5 & * & * & 4 & * & 3 & * & * & 9 & * & * \\ * & * & 9 & * & * & 8 & * & 4 & * & * & 1 & * \\ 7 & * & * & 9 & * & * & * & 5 & * & * & 2 & * \\ * & 8 & * & * & 7 & * & * & 6 & * & * & 3 & * \end{pmatrix} \quad (28)$$

By modifying (21), the following construction can be obtained.

**Construction 3:** For any positive integers  $q, z, m$  and  $t$  with  $z < q$  and  $t < m$ , let

$$\mathcal{F} = \{(a_0, a_1, \dots, a_{m-1}) \mid a_0, \dots, a_{m-1} \in \mathbb{Z}_q\}. \quad (29)$$

and

$$\mathcal{K} = \{(b_0, b_1, \dots, b_{t-1}, \delta_0, \delta_1, \dots, \delta_{t-1}, \varepsilon_0, \dots, \varepsilon_{t-1}) \mid b_0, \dots, b_{t-1} \in \mathbb{Z}_q, \\ 0 \leq \delta_0 < \dots < \delta_{t-1} < m, \varepsilon_0, \dots, \varepsilon_{t-1} \in [0, \lfloor \frac{q-1}{q-z} \rfloor]\} \quad (30)$$

Using the same rule defined in (22), we can obtain a  $q^m \times \binom{m}{t} \lfloor \frac{q-1}{q-z} \rfloor^t q^t$  array.

**Example 6:** When  $m = 2, q = 3$  and  $z = 2$ , (29) and (30) can be written as

$$\mathcal{F} = \{(a_0, a_1) \mid a_0, a_1 \in \mathbb{Z}_3\} \quad \text{and} \quad \mathcal{K} = \{(b, \delta, \varepsilon) \mid b \in \mathbb{Z}_3, 0 \leq \delta \leq 1, \varepsilon \in [0, 1]\}.$$

By (22), the following array can be obtained. It is easy to check that this array is a (12, 9, 6, 9) PDA.

$(a_0, a_1) \setminus (b, \delta, \varepsilon)$	$(0, 0, 0)$	$(1, 0, 0)$	$(2, 0, 0)$	$(0, 0, 1)$	$(1, 0, 1)$	$(2, 0, 1)$	$(0, 1, 0)$	$(1, 1, 0)$	$(2, 1, 0)$	$(0, 1, 1)$	$(1, 1, 1)$	$(2, 1, 1)$
$(0, 0)$	*	*	$(2, 0, 0)$	*	*	$(1, 0, 0)$	*	*	$(0, 2, 0)$	*	*	$(0, 1, 0)$
$(1, 0)$	$(0, 0, 0)$	*	$(2, 0, 0)$	*	*	*	*	*	$(1, 2, 0)$	*	*	$(1, 1, 0)$
$(2, 0)$	*	$(1, 0, 0)$	*	*	$(0, 0, 0)$	*	*	*	$(2, 2, 0)$	*	*	$(2, 1, 0)$
$(0, 1)$	*	*	$(2, 1, 0)$	*	*	$(1, 1, 0)$	$(0, 0, 0)$	*	*	$(0, 2, 0)$	*	*
$(1, 1)$	$(0, 1, 0)$	*	*	$(2, 1, 0)$	*	*	$(1, 0, 0)$	*	*	$(1, 2, 0)$	*	*
$(2, 1)$	*	$(1, 1, 0)$	*	*	$(0, 1, 0)$	*	$(2, 0, 0)$	*	*	$(2, 2, 0)$	*	*
$(0, 2)$	*	*	$(2, 2, 0)$	*	*	$(1, 2, 0)$	*	$(0, 1, 0)$	*	*	$(0, 0, 0)$	*
$(1, 2)$	$(0, 2, 0)$	*	*	$(2, 2, 0)$	*	*	*	$(1, 1, 0)$	*	*	$(1, 0, 0)$	*
$(2, 2)$	*	$(1, 2, 0)$	*	*	$(0, 2, 0)$	*	*	$(2, 1, 0)$	*	*	$(2, 0, 0)$	*

The array in (31) is exactly the PDA in (28) when we replace the vectors by the integers in [1, 9]. Similar to the proof of Theorem 2, Theorem 4 can be proved.

### D. The second extended construction

Similar to generating the PDA in (28), we can obtain the following (15, 9, 6, 9) PDA  $\mathbf{P}''$  by copying or replacing three integer entries of each column of  $\mathbf{P}$  in (24) by “\*”s elaborately.

$$\mathbf{P}'' = \begin{pmatrix} * & * & 3 & * & * & 2 & * & * & 7 & * & * & 4 & * & 1 & * \\ 1 & * & * & 3 & * & * & * & * & 8 & * & * & 5 & * & * & 2 \\ * & 2 & * & * & 1 & * & * & * & 9 & * & * & 6 & 3 & * & * \\ * & * & 6 & * & * & 5 & 1 & * & * & 7 & * & * & * & * & 4 \\ 4 & * & * & 6 & * & * & 2 & * & * & 8 & * & * & 5 & * & * \\ * & 5 & * & * & 4 & * & 3 & * & * & 9 & * & * & * & 6 & * \\ * & * & 9 & * & * & 8 & * & 4 & * & * & 1 & * & 7 & * & * \\ 7 & * & * & 9 & * & * & * & 5 & * & * & 2 & * & * & 8 & * \\ * & 8 & * & * & 7 & * & * & 6 & * & * & 3 & * & * & * & 9 \end{pmatrix} \quad (32)$$

**Construction 4:** When  $t = 1$ , (29) and (30) can be written as

$$\mathcal{F} = \{(a_0, a_1, \dots, a_{m-1}) \mid a_0, \dots, a_{m-1} \in \mathbb{Z}_q\} \quad \text{and} \quad \mathcal{K} = \{(b, \delta, \varepsilon) \mid b \in \mathbb{Z}_q, 0 \leq \delta \leq m, \varepsilon \in [0, \lfloor \frac{q-1}{q-z} \rfloor]\}$$

for any positive integers  $q$ ,  $z$  and  $m$  with  $z < q$ .

- Using the same rule defined in (22), we can obtain a  $q^m \times m \lfloor \frac{q-1}{q-z} \rfloor q$  array  $\mathbf{P} = (p_{\mathbf{a}, \mathbf{b}})$ ,  $\mathbf{a} \in \mathcal{F}$ ,  $\mathbf{b} \in \mathcal{K}$ .
- Let  $\mathcal{K}_1 = \{(b, m) \mid b \in \mathbb{Z}_q\}$ . We can define a  $q^m \times q$  array  $\mathbf{C} = (c_{\mathbf{a}, \mathbf{b}})$ ,  $\mathbf{a} \in \mathcal{F}$ ,  $\mathbf{b} \in \mathcal{K}_1$  in the following way.

$$c_{\mathbf{a}, \mathbf{b}} = \begin{cases} (a_0, \dots, a_{m-1}, b - \sum_{l=0}^{m-1} a_l - 1)_q & \text{if } \sum_{l=0}^{m-1} a_l \notin Y_{b,z} \\ * & \text{otherwise} \end{cases} \quad (33)$$

where  $Y_{b,z} = \{b, b+1, \dots, b+(z-1)\}$ . All the above operations are performed modulo  $q$ .

Based on the array generated by Construction 4, the proof of Theorem 5 can be obtained and is included in Appendix C.

## V. CONCLUSION

In this paper, four classes of new schemes were constructed. Our new constructions did not only include some previously known constructions as special cases, but also had the following advantages: In the first two classes,  $F$  grew sub-exponentially with  $K$ ; And the other two classes could significantly reduce the value of  $F$  by just increasing some fixed times of  $R$  comparing with MN PDAs. In addition, an exact tradeoff between  $R$  and  $F$  was proposed.

## REFERENCES

- [1] White paper: Cisco VNI forecast and methodology, 2015-2020, Tech. Report, 2015.
- [2] K. C. Almeroth and M. H. Ammar, The use of multicast delivery to provide a scalable and interactive video-on-demand service, *IEEE Journal on Selected Areas in Communications*, vol. 14, no. 6, pp. 1110-1122, 1996.
- [3] M. M. Amiri and D. Gündüz, Fundamental limits of caching: Improved delivery rate-cache capacity trade-off, *IEEE Transactions on Communications*, vol. 65, no. 2, pp. 806-815, 2016.
- [4] M. Cheng, Q. Yan, X. Tang, and J. Jiang, Coded caching schemes with low rate and subpacketizations, arXiv:1703.01548v2 [cs.IT], Apr 2017.
- [5] H. Ghasemi and A. Ramamoorthy, Improved lower bounds for coded caching, in Proc. *IEEE International Symposium on Information Theory*, Hong Kong, Jun. 2015, pp. 1696-1700.
- [6] N. Golrezaei, A. F. Molisch, A. G. Dimakis, and G. Caire, Femtocaching and device-to-device collaboration: A new architecture for wireless video distribution, *IEEE Communications Magazine*, vol. 51, no. 4, pp. 142-149, 2013.
- [7] M. Gregori, J. Gomez-Vilardebo, J. Matamoros, and D. Gündüz, Wireless content caching for small cell and D2D networks, *IEEE Journal on Selected Areas in Communications*, Vol. 34, no. 5, pp. 1222-1234, 2016.
- [8] M. Ji, G. Caire and A. F. Molisch, Fundamental Limits of Caching in Wireless D2D Networks, *IEEE Transactions on Information Theory*, vol. 62, no. 2, pp.849-869, 2016.
- [9] M. Ji, A. M. Tulino, J. Llorca, and G. Caire, Order-optimal rate of caching and coded multicasting with random demands, in Proc. *11th International Symposium on Wireless Communications Systems*, Barcelona, Aug. 2014, pp. 922-926.
- [10] N. Karamchandani, U. Niesen, M. A. Maddah-Ali, and S. Diggavi, Hierarchical coded caching, in Proc. *IEEE International Symposium on Information Theory*, Honolulu, HI, Jun. 2014, pp. 2142-2146.
- [11] J. Li, X. Tang, U. Parampalli, A Framework of Constructions of Minimal Storage Regenerating Codes With the Optimal Access/Update Property, *IEEE Transactions on Information Theory*, vol. 61, no. 4, pp. 1920-1932, 2015.
- [12] M. A. Maddah-Ali and U. Niesen, Fundamental limits of caching, *IEEE Transactions on Information Theory*, vol. 60, no. 5, pp. 2856-2867, 2014.
- [13] M. A. Maddah-Ali, Urs Niesen, Decentralized coded caching attains order-optimal memory-rate tradeoff, *IEEE/ACM Transactions on Networking*, vol. 23, no. 4, pp.1029-1040, 2015.
- [14] R. Pedarsani, M. A. Maddah-Ali, and U. Niesen, Online coded caching, in Proc. *IEEE International Conference on Communications*, Sydney, Australia, Jun. 2014, pp. 1878-1883.
- [15] A. Sengupta, R. Tandon, and T. C. Clancy, Improved approximation of storage-rate tradeoff for caching via new outer bounds, in Proc. *IEEE International Symposium on Information Theory*, Hong Kong, Jun. 2015, pp. 1691-1695.
- [16] C. Shanguan, Y. Zhang, G. Ge, Centralized coded caching schemes: A hypergraph theoretical approach, arXiv:1608.03989 [cs.IT], Aug. 2016.
- [17] K. Shanmugam, M. Ji, A. M. Tulino, J. Llorca, and A. G. Dimakis, Finite-length analysis of caching-aided coded multicasting, *IEEE Transactions on Information Theory*, vol. 62, no. 10, pp. 5524-5537, 2016.
- [18] K. Shanmugam, A. M. Tulino, and A. G. Dimakis, Coded caching with linear subpacketization is possible using Ruzsa-Szemredi graphs, arXiv:1701.07115v1 [cs.IT], Jan 2017.
- [19] I. Tamo, Z. Wang, and J. Bruck, Zigzag codes: MDS array codes with optimal rebuilding, *IEEE Transactions on Information Theory*, vol. 59, no. 3, pp. 1597-1616, 2013.
- [20] L. Tang and A. Ramamoorthy, Coded caching with low subpacketization levels, in Proc. *IEEE International Symposium on Information Theory*, Aachen, Germany, Jun. 2017, pp. 2790-2794.

- [21] C. Tian and J. Chen, Caching and delivery via interference elimination, in Proc. *IEEE International Symposium on Information Theory*, Barcelona, July 2016, pp. 830-834.
- [22] K. Wan, D. Tuninetti and P. Piantanida, On the optimality of uncoded cache placement, in Proc. *IEEE Information Theory Workshop*, Cambridge, UK, Sept. 2016.
- [23] C. Y. Wang, S. H. Lim, M. Gastpar, A new converse bound for coded caching, in Proc. *IEEE Information Theory Workshop*, Robinson College, Oct. 2016.
- [24] Q. Yan, M. Cheng, X. Tang and Q. Chen, On the placement delivery array design in centralized coded caching scheme, *IEEE Transactions on Information Theory*, vol. 63, no. 9, pp. 5821-5833, 2017.
- [25] Q. Yan, U. Parampalli, X. Tang, and Q. Chen, Online coded caching with random access, *IEEE Communications Letters*, 2016.
- [26] Q. Yan, X. Tang, and Q. Chen, On the gap between decentralized and centralized coded caching schemes, arXiv: 1605.04626 [cs.IT], May 2016.
- [27] Q. Yan, X. Tang, Q. Chen, and M. Cheng, Placement delivery array design through strong edge coloring of bipartite graphs, *IEEE Communications Letters*, to appear.
- [28] L. Zhang, Z. Wang, M. Xiao, G. Wu, and S. Li, Decentralized caching in two-layer networks: Algorithms and limits, *IEEE 12th International Conference on Wireless and Mobile Computing, Networking and Communications*, New York, Oct. 2016.

## APPENDIX A: PROOF OF THEOREM 2

*Proof.* It is easy to check that there are exactly  $q^{m-t}(q-z)^t \lfloor \frac{q-1}{q-z} \rfloor^t$  vector entries in each column. This implies that  $Z = (q^m - q^{m-t}(q-z)^t) \lfloor \frac{q-1}{q-z} \rfloor^t$ . So C1 holds. From (22), the vector set of  $\mathbf{P}$  in Construction 1 is

$$\mathcal{S} = \{(a_0, a_1, \dots, a_{m-1}, a_m, \dots, a_{m+t-1}) \mid a_0, \dots, a_{m-1} \in \mathbb{Z}_q, a_{m+1}, \dots, a_{m+t-1} \in [0, q-z]\}.$$

So  $S = |\mathcal{S}| = q^m(q-z)^t$ . Each  $\mathbf{s} = (s_0, s_1, \dots, s_{m+t-1}) \in \mathcal{S}$  appears in the entries at row  $\mathbf{a}$  and column  $\mathbf{b}$  of  $\mathbf{P}$ , if and only if

$$\begin{aligned} \mathbf{a} &= (a_0, \dots, a_{\delta_i}, \dots, a_{m-1}, \varepsilon_0, \dots, \varepsilon_{t-1}) = (s_0, \dots, s_{\delta_i} + s_{m+i} + \varepsilon_i(q-z) + 1, \dots, s_{m-1}, \varepsilon_0, \dots, \varepsilon_{t-1})_q \quad \text{and} \\ \mathbf{b} &= (b_0, b_1, \dots, b_{t-1}, \delta_0, \delta_1, \dots, \delta_{t-1}) = (s_{\delta_0} + \varepsilon_0(q-z), s_{\delta_1} + \varepsilon_1(q-z), \dots, s_{\delta_{t-1}} + \varepsilon_{t-1}(q-z), \delta_0, \delta_1, \dots, \delta_{t-1}). \end{aligned} \quad (34)$$

Clearly, for any fixed  $\delta_0, \delta_1, \dots, \delta_{t-1}$  and  $\varepsilon_0, \dots, \varepsilon_{t-1}$ , there is exactly a unique pair of vectors  $\mathbf{a}$  and  $\mathbf{b}$ , such that  $p_{\mathbf{a}, \mathbf{b}} = \mathbf{s}$ . Based on the above observation, C2 is clear. Now, let us consider C3. Suppose there exists another two vectors  $\mathbf{a}'$  and  $\mathbf{b}'$  with  $p_{\mathbf{a}', \mathbf{b}'} = \mathbf{s}$ . Then there must exist another  $2t$  integers  $\delta'_0, \dots, \delta'_{t-1}$  and  $\varepsilon'_0, \dots, \varepsilon'_{t-1}$  satisfying

$$\begin{aligned} \mathbf{a}' &= (a'_0, \dots, a'_{\delta'_i}, \dots, a'_{m-1}, \varepsilon'_0, \dots, \varepsilon'_{t-1}) = (s_0, \dots, s_{\delta'_i} + s_{m+i} + \varepsilon'_i(q-z) + 1, \dots, s_{m-1}, \varepsilon'_0, \dots, \varepsilon'_{t-1})_q \quad \text{and} \\ \mathbf{b}' &= (b'_0, b'_1, \dots, b'_{t-1}, \delta'_0, \delta'_1, \dots, \delta'_{t-1}) = (s_{\delta'_0} + \varepsilon'_0(q-z), s_{\delta'_1} + \varepsilon'_1(q-z), \dots, s_{\delta'_{t-1}} + \varepsilon'_{t-1}(q-z), \delta'_0, \delta'_1, \dots, \delta'_{t-1}). \end{aligned} \quad (35)$$

It is sufficient to consider the following cases, where all the operations are performed modulo  $q$ .

- If  $\{\delta_0, \dots, \delta_{t-1}\} \neq \{\delta'_0, \dots, \delta'_{t-1}\}$ , there at least exist two integers  $i, i' \in [0, t)$  such that  $\delta_i \notin \{\delta'_0, \dots, \delta'_{t-1}\}$  and  $\delta'_{i'} \notin \{\delta_0, \dots, \delta_{t-1}\}$ . Now we show C3-a) and C3-b) hold respectively.

- If  $\mathbf{s}$  occurs in a row at least twice, we have  $\mathbf{a} = \mathbf{a}'$ . This implies that  $a_{\delta_i} = a'_{\delta'_i}$ . From (34) and (35), we have

$$a_{\delta_i} = s_{\delta_i} + s_{m+i} + \varepsilon_i(q-z) + 1, \quad a'_{\delta'_i} = s_{\delta'_i}.$$

This implies that  $s_{m+i} + \varepsilon_i(q-z) + 1 = 0$ . This is impossible since

$$1 \leq s_{m+i} + \varepsilon_i(q-z) + 1 < q-z + \left(\frac{q-1}{q-z} - 1\right)(q-z) + 1 = q$$

by the facts  $s_{m+i} \in [0, q-z)$  and  $\varepsilon \in [0, \lfloor \frac{q-1}{q-z} \rfloor]$ . So C3-a) holds.

- First from (34) and (35), we have

$$a_{\delta'_{i'}} = s_{\delta'_{i'}}, \quad b'_{i'} = s_{\delta'_{i'}} + \varepsilon'_{i'}(q-z), \quad a'_{\delta_i} = s_{\delta_i}, \quad b_i = s_{\delta_i} + \varepsilon_i(q-z)$$

If  $p_{\mathbf{a}, \mathbf{b}'} \neq *$ , we have  $a_{\delta'_{i'}} \in \{b'_{i'} + 1, b'_{i'} + 2, \dots, b'_{i'} + (q-z)\}$ . Then there exists an integer  $j \in [1, q-z)$  such that  $a_{\delta'_{i'}} = b'_{i'} + j$ , i.e.,

$$a_{\delta'_{i'}} = s_{\delta'_{i'}} = s_{\delta'_{i'}} + \varepsilon'_{i'}(q-z) + j.$$

This implies that  $\varepsilon'_{i'}(q-z) + j = 0$ . This is impossible since

$$1 \leq \varepsilon'_{i'}(q-z) + j < \left(\frac{q-1}{q-z} - 1\right)(q-z) + (q-z) = q-1$$

by the fact  $\varepsilon'_i \in [0, \lfloor \frac{q-1}{q-z} \rfloor]$ . Similarly we can also show that  $p_{\mathbf{a}', \mathbf{b}} = *$  too.

- If  $\{\delta_0, \dots, \delta_{t-1}\} = \{\delta'_0, \dots, \delta'_{t-1}\}$ , we have  $(\varepsilon_0, \dots, \varepsilon_{t-1}) \neq (\varepsilon'_0, \dots, \varepsilon'_{t-1})$  by (34) and (35). So there exist integer  $i$  such that  $\varepsilon_i \neq \varepsilon'_i$ . Then we have

$$b_i = s_{\delta_i} + \varepsilon_i(q-z), \quad a_{\delta_i} = s_{m+i} + b_i + 1, \quad b'_i = s_{\delta_i} + \varepsilon'_i(q-z), \quad a'_{\delta_i} = s_{m+i} + b'_i + 1. \quad (36)$$

Now we show C3-a) and C3-b) hold respectively.

- If  $\mathbf{s}$  occurs in a column at least twice, we have  $\mathbf{b} = \mathbf{b}'$ . This implies that  $b_{\delta_i} = b'_{\delta_i}$ . By the first and third items in (36),  $\varepsilon_i(q-z) = \varepsilon'_i(q-z)$  holds. Clearly this is impossible since  $\varepsilon_i, \varepsilon'_i \in [0, \lfloor \frac{q-1}{q-z} \rfloor]$ . So C3-a) holds.
- If  $p_{\mathbf{a}, \mathbf{b}'} \neq *$ , we have  $a_{\delta_i} \in \{b'_i + 1, b'_i + 2, \dots, b'_i + (q-z)\}$ . Then there exists an integer  $j \in [0, q-z)$  such that  $a_{\delta_i} = b'_i + j$ . By the first, second and third items in (36),

$$s_{m+i} + s_{\delta_i} + \varepsilon_i(q-z) + 1 = s_{\delta_i} + \varepsilon'_i(q-z) + j. \quad (37)$$

If  $\varepsilon_i < \varepsilon'_i$ , we have  $s_{m+i} + 1 = (\varepsilon'_i - \varepsilon_i)(q-z) + j$ . This is impossible since

$$1 \leq s_{m+i} + 1 \leq q-z \quad \text{and} \quad q-z+1 \leq (\varepsilon'_i - \varepsilon_i)(q-z) + j < q$$

by the facts  $0 \leq \varepsilon_i, \varepsilon'_i < \lfloor \frac{q-1}{q-z} \rfloor$  and  $1 \leq j, s_{m+i} < q-z$ . Clearly (37) does not hold either if  $\varepsilon_i > \varepsilon'_i$ . So we have  $p_{\mathbf{a}, \mathbf{b}'} = *$ . Similarly we can also show that  $p_{\mathbf{a}', \mathbf{b}} = *$  too. □

## APPENDIX B: PROOF OF THEOREM 3

*Proof.* From Construction 2, and the proof of Theorem 2, we have that  $\mathbf{P}$  is a  $(mq, \lfloor \frac{q-1}{q-z} \rfloor q^m, z \lfloor \frac{q-1}{q-z} \rfloor q^{m-1}, (q-z)q^m)$  PDA with the vector set

$$\mathcal{S} = \{(s_0, s_1, \dots, s_m) \mid s_0, s_1, \dots, s_{m-1} \in [0, q), s_m \in [0, q-z)\}.$$

From (26), it is easy to count that each column of  $\mathbf{C}$  has exactly  $z \lfloor \frac{q-1}{q-z} \rfloor q^{m-1}$  stars and the vector set of  $\mathbf{C}$  is  $\mathcal{S}$ . So C1 and C2 hold. Now it is sufficient to verify C3. For any two distinct vector entries, say  $h_{\mathbf{a}, \mathbf{b}}$  and  $h_{\mathbf{a}', \mathbf{b}'}$ , assume that  $h_{\mathbf{a}, \mathbf{b}} = h_{\mathbf{a}', \mathbf{b}'} \in \mathcal{S}$  where

$$\mathbf{a} = (a_0, a_1, \dots, a_{m-1}, \varepsilon)_q, \quad \mathbf{a}' = (a'_0, a'_1, \dots, a'_{m-1}, \varepsilon')_q, \quad \mathbf{b} = (b, \delta), \quad \mathbf{b}' = (b', \delta').$$

We only need to consider the following cases, where all the operations are performed modulo  $q$ .

- When  $\delta \in [0, m)$  and  $\delta' = m$ , we have  $h_{\mathbf{a}, \mathbf{b}} = p_{\mathbf{a}, \mathbf{b}}$  and  $h_{\mathbf{a}', \mathbf{b}'} = c_{\mathbf{a}', \mathbf{b}'}$  in distinct columns. From (22) and (26) we have

$$a_\delta - b - 1 = b' - \left( \sum_{l=0}^{m-1} a'_l - \varepsilon'(q-z) \right) - 1, \quad a'_\delta = b - \varepsilon(q-z), \quad a_l = a'_l \quad (38)$$

for any  $l \in [0, m) \setminus \{\delta\}$ . From the second equation in (38), we can show  $p_{\mathbf{a}, \mathbf{b}'}$  and  $c_{\mathbf{a}', \mathbf{b}}$  in distinct rows too. Otherwise we have  $a_l = a'_l$  for any  $l = 0, 1, \dots, m-1$  and  $\varepsilon = \varepsilon'$ . Then combining the two equations in (38) we have

$$a_\delta - b = b' - \left( \sum_{l=0}^{m-1} a'_l - \varepsilon'(q-z) \right) = -\varepsilon(q-z).$$

Since  $\sum_{l=0}^{m-1} a'_l - \varepsilon'(q-z) \notin Y_{b, z}$ , let  $\sum_{l=0}^{m-1} a'_l - \varepsilon'(q-z) = b' - j$  where  $j \in [1, q-z]$ . Then the above equation can be written as

$$b' - \left( \sum_{l=0}^{m-1} a'_l - \varepsilon'(q-z) \right) = b' - (b' - j) = -\varepsilon(q-z).$$

This implies that  $j + \varepsilon(q-z) = 0$ . This is impossible since

$$1 \leq j + \varepsilon(q-z) < (q-z) + \left( \frac{q-1}{q-z} - 1 \right) (q-z) = q-1 \quad (39)$$

by the fact  $\varepsilon \in [0, \lfloor \frac{q-1}{q-z} \rfloor]$ . So C3-a) holds. Now let us consider the entries  $c_{\mathbf{a}, \mathbf{b}'}$  and  $p_{\mathbf{a}', \mathbf{b}}$  in the following cases.

- If  $c_{\mathbf{a},\mathbf{b}'}$  is a vector,  $\sum_{l=0}^{m-1} a_l - \varepsilon(q-z) \in \{b' - (q-z), b' - (q-z-1), \dots, b' - 1\}$ . That is, there must exist an integer  $j \in \{1, 2, \dots, q-z\}$  such that  $\sum_{l=0}^{m-1} a_l - \varepsilon(q-z) = b' - j$ . From (38),

$$\begin{aligned}
\sum_{l=0}^{m-1} a_l - \varepsilon(q-z) &= \sum_{l \in [0, m) \setminus \{\delta\}} a_l + a_\delta - \varepsilon(q-z) \\
&= \sum_{l \in [0, m) \setminus \{\delta\}} a_l + (b + b' - (\sum_{l=0}^{m-1} a'_l - \varepsilon'(q-z))) - \varepsilon(q-z) \\
&= \sum_{l \in [0, m) \setminus \{\delta\}} a_l + (b + b' - \sum_{l \in [0, m) \setminus \{\delta\}} a'_l - a'_\delta + \varepsilon'(q-z)) - \varepsilon(q-z) \\
&= b + b' - a'_\delta + \varepsilon'(q-z) - \varepsilon(q-z) \\
&= b + b' - (b - \varepsilon(q-z)) + \varepsilon'(q-z) - \varepsilon(q-z) \\
&= b' + \varepsilon'(q-z).
\end{aligned}$$

So we have  $b' - j = b' + \varepsilon'(q-z)$ , i.e.,  $j + \varepsilon'(q-z) = 0$ . This is impossible by the similar proof in (39). So  $c_{\mathbf{a},\mathbf{b}'} = *$ .

- If  $p_{\mathbf{a}',\mathbf{b}}$  is a vector,  $a'_\delta \in \{b+1, b+2, \dots, b+q-z\}$ . There must exist an integer  $j \in \{1, 2, \dots, q-z\}$  such that  $a'_\delta = b+j$ . From (38), we have  $b+j = b - \varepsilon(q-z)$ , i.e.,  $j + \varepsilon(q-z) = 0$ . This is impossible by (39). So  $p_{\mathbf{a}',\mathbf{b}} = *$ .

So C3-b) holds.

- When  $\delta = \delta' = m$ , we have  $h_{\mathbf{a},\mathbf{b}} = c_{\mathbf{a},\mathbf{b}}$  and  $h_{\mathbf{a}',\mathbf{b}'} = c_{\mathbf{a}',\mathbf{b}'}$ . From (26) we have

$$b - (\sum_{l=0}^{m-1} a_l - \varepsilon(q-z)) - 1 = b' - (\sum_{l=0}^{m-1} a'_l - \varepsilon'(q-z)) - 1, \quad a_l = a'_l \quad (40)$$

for any  $l \in [0, m)$ . Clearly  $\varepsilon \neq \varepsilon'$  always holds. Otherwise we have  $\mathbf{a} = \mathbf{a}'$  and  $b = b'$ , i.e.,  $\mathbf{b} = \mathbf{b}'$ . This is impossible. If  $\mathbf{b} = \mathbf{b}'$ , we have  $\varepsilon = \varepsilon'$  since  $0 \leq \varepsilon'(q-z), \varepsilon'(q-z) < q$  by the fact  $\varepsilon, \varepsilon' \in [0, \lfloor \frac{q-1}{q-z} \rfloor]$ . So C3-a) is clear. Now let us consider C3-b). If  $c_{\mathbf{a},\mathbf{b}'}$  is a vector, there must exist an integer  $x \in \{1, 2, \dots, q-z\}$  satisfying

$$\sum_{l=0}^{m-1} a_l - \varepsilon(q-z) = b' - x.$$

Since  $c_{\mathbf{a},\mathbf{b}}$  is a vector entry, there exist an integer  $y \in \{1, 2, \dots, q-z\}$  such that

$$(\sum_{l=0}^{m-1} a_l - \varepsilon(q-z)) = b - y.$$

So we have

$$(\sum_{l=0}^{m-1} a_l - \varepsilon(q-z)) = b' - x = b - y. \quad (41)$$

Moreover, from (40) we have

$$b + \varepsilon(q-z) = b' + \varepsilon'(q-z). \quad (42)$$

Combining (41) and (42), we have  $x - y = (\varepsilon - \varepsilon')(q-z)$ . From the above discussion, we know that  $\varepsilon \neq \varepsilon'$ , i.e.,  $(\varepsilon - \varepsilon')(q-z) \neq 0$ . So we have  $x \neq y$ . Furthermore,  $(q-z)|(x-y)$  holds since  $q-z \leq |(\varepsilon - \varepsilon')(q-z)| < q-1$ . Clearly this is impossible by the fact that  $1 \leq x \neq y \leq q-z$ . So  $c_{\mathbf{a},\mathbf{b}'} = *$ . Similarly we can also show that  $c_{\mathbf{a}',\mathbf{b}} = *$ . So C3-b) holds. □



## APPENDIX C: PROOF OF THEOREM 5

*Proof.* Similar to the proof of Theorem 3, denote  $\mathbf{H} = (\mathbf{P}, \mathbf{C})$ . From Construction 4  $\mathbf{P}$  is a  $(m \lfloor \frac{q-1}{q-z} \rfloor q, q^m, zq^m, (q-z)q^m)$  PDA with the vector set

$$\mathcal{S} = \{(s_0, s_1, \dots, s_m) \mid s_0, s_1, \dots, s_{m-1} \in [0, q], s_m \in [0, q-z]\}.$$

From (33), it is easy to count that each column of  $\mathbf{C}$  has exactly  $zq^{m-1}$  stars and the vector set of  $\mathbf{C}$  is  $\mathcal{S}$ . So C1 and C2 hold. Now it is sufficient to verify C3. For any two distinct vector entries, say  $h_{\mathbf{a}, \mathbf{b}}$  and  $h_{\mathbf{a}', \mathbf{b}'}$ , assume that  $h_{\mathbf{a}, \mathbf{b}} = h_{\mathbf{a}', \mathbf{b}'} \in \mathcal{S}$  where

$$\mathbf{a} = (a_0, a_1, \dots, a_{m-1})_q, \quad \mathbf{a}' = (a'_0, a'_1, \dots, a'_{m-1})_q, \quad \mathbf{b} = (b, \delta, \varepsilon), \quad \mathbf{b}' = (b', \delta', \varepsilon').$$

We only need to consider the following cases, where all the operations are performed modulo  $q$ .

- When  $\delta \in [0, m)$  and  $\delta' = m$ , we have  $p_{\mathbf{a}, \mathbf{b}}$  and  $c_{\mathbf{a}', \mathbf{b}'}$  in distinct columns. From (22) and (33) we have

$$a_\delta - b - 1 = b' - \sum_{l=0}^{m-1} a'_l - 1, \quad a'_\delta = s_\delta = b - \varepsilon(q-z), \quad a_l = a'_l \quad (43)$$

for any  $l \in [0, m) \setminus \{\delta\}$ . From the second equation in (43), we can show  $p_{\mathbf{a}, \mathbf{b}}$  and  $c_{\mathbf{a}', \mathbf{b}'}$  in distinct rows. Otherwise  $a_\delta = a'_\delta$  holds. By (43) the following equation can be obtained.

$$a_\delta = a'_\delta = b + b' - \sum_{l=0}^{m-1} a'_l = b - \varepsilon(q-z)$$

Since  $\sum_{l=0}^{m-1} a'_l \notin Y_{b', z}$ , there exists an integer  $j \in \{1, 2, \dots, q-z\}$  such that  $\sum_{l=0}^{m-1} a'_l = b' - j$ . Combining the above formula we have

$$b + b' - (b' - j) = b - \varepsilon(q-z)$$

i.e.,  $j + \varepsilon(q-z) = 0$ . This is impossible since

$$1 \leq j + \varepsilon(q-z) < (q-z) + \left(\frac{q-1}{q-z} - 1\right)(q-z) = q-1 \quad (44)$$

by the fact  $\varepsilon \in [0, \lfloor \frac{q-1}{q-z} \rfloor]$ . So C3-a) holds. Now let us consider the entries  $p_{\mathbf{a}, \mathbf{b}'}$  and  $p_{\mathbf{a}', \mathbf{b}}$  in the following cases.

- If  $c_{\mathbf{a}, \mathbf{b}'}$  is a vector, we have  $\sum_{l=0}^{m-1} a_l \in \{b' - (q-z), b' - (q-z-1), \dots, b' - 1\}$ . There must exist an integer  $j \in \{1, 2, \dots, q-z\}$  such that  $\sum_{l=0}^{m-1} a_l = b' - j$ . From (43), we have

$$\begin{aligned} \sum_{l=0}^{m-1} a_l &= \sum_{l \in [0, m) \setminus \{\delta\}} a_l + a_\delta \\ &= \sum_{l \in [0, m) \setminus \{\delta\}} a_l + (b + b' - \sum_{l=0}^{m-1} a'_l) \\ &= \sum_{l \in [0, m) \setminus \{\delta\}} a_l + (b + b' - \sum_{l \in [0, m) \setminus \{\delta\}} a'_l - a'_\delta) \\ &= b + b' - a'_\delta \\ &= b + b' - (b - \varepsilon(q-z)) \\ &= b' + \varepsilon(q-z) \end{aligned}$$

So we have  $b' - j = b' + \varepsilon(q-z)$ , i.e.,  $j + \varepsilon(q-z) = 0$ . This is impossible by (44). So  $c_{\mathbf{a}, \mathbf{b}'} = *$ .

- If  $p_{\mathbf{a}', \mathbf{b}}$  is a vector, we have  $a'_\delta \in \{b+1, b+2, \dots, b+q-z\}$ . There must exist an integer  $j \in \{1, 2, \dots, q-z\}$  such that  $a'_\delta = b+j$ . From (43), we have  $b+j = b - \varepsilon(q-z)$ , i.e.,  $j + \varepsilon(q-z) = 0$ . This is impossible by (44). So  $p_{\mathbf{a}', \mathbf{b}} = *$ .

So C3-b) holds.

- When  $\delta = \delta' = m$ , from (33) we have

$$b - \sum_{l=0}^{m-1} a_l - 1 = b' - \sum_{l=0}^{m-1} a'_l - 1, \quad a_l = a'_l$$

for any  $l \in [0, m)$ . This implies that  $\mathbf{a} = \mathbf{a}'$  and  $\mathbf{b} = \mathbf{b}'$ , a contradiction to our hypothesis. So C3-b) holds.

□