

Enumerations of maximum partial triple systems on 16 and 17 points

Fatih Demirkale

Department of Mathematics
Yıldız Technical University
Esenler, İstanbul 34220
TURKEY
(fatihd@yildiz.edu.tr)

Diane Donovan

Centre for Discrete Mathematics and Computing
University of Queensland, St. Lucia 4072
AUSTRALIA
(dmd@maths.uq.edu.au)

Mike Grannell*

School of Mathematics and Statistics
The Open University, Walton Hall
Milton Keynes MK7 6AA
UNITED KINGDOM
(m.j.grannell@open.ac.uk)

November 16, 2021

Abstract

For $v \equiv 1$ or $3 \pmod{6}$, maximum partial triple systems on v points are Steiner triple systems, $\text{STS}(v)$ s. The 80 non-isomorphic $\text{STS}(15)$ s were first enumerated around 100 years ago, but the next case for Steiner triple systems was unresolved until around 2004 when it was established that there are precisely 11 084 874 829 non-isomorphic $\text{STS}(19)$ s. In this paper we complete enumeration of non-isomorphic maximum partial triple systems for $v \leq 19$. It is shown that there are 35 810 097 systems on 17 points and 47 744 568 on 16 points. We also establish that there are precisely 157 151 non-isomorphic pairwise balanced designs, $\text{PBD}(17, \{3, 5\})$ s, having a single block of size 5. Structural properties of these systems are determined, including their automorphism groups, and the numbers of Pasch configurations, mitres and Fano planes contained in them. The systems themselves are available from the authors.

*Corresponding author

AMS classification: 05B07.

Keywords: Maximum partial triple system; Steiner triple system; Pairwise balanced design; Pasch configuration; mitre configuration.

1 Introduction

We will be concerned with certain types of combinatorial designs, in particular partial triple systems. A *partial triple system* of order v and index λ , $\text{PTS}(v, \lambda)$, is a pair (V, \mathcal{B}) , where V is a set of v elements (the *points*) and \mathcal{B} is a collection of 3-element subsets of V (the *triples* or *blocks*) with the property that every 2-element subset of V occurs at most λ times amongst the triples of \mathcal{B} . A $\text{PTS}(v, \lambda) = (V, \mathcal{B})$ is said to be *maximal* if there is no other $\text{PTS}(v, \lambda) = (V, \mathcal{B}')$ with $\mathcal{B} \subseteq \mathcal{B}'$. Informally, a $\text{PTS}(v, \lambda) = (V, \mathcal{B})$ is maximal if no further triples can be added to \mathcal{B} without violating the condition that pairs appear at most λ times. A maximal $\text{PTS}(v, \lambda)$ is denoted as $\text{MPTS}(v, \lambda)$. A *maximum* $\text{MPTS}(v, \lambda)$, denoted as $\text{MMPTS}(v, \lambda)$, is an $\text{MPTS}(v, \lambda)$ having the greatest number of triples of any $\text{MPTS}(v, \lambda)$. Rather than calling an $\text{MMPTS}(v, \lambda)$ a “maximum maximal partial triple system”, we prefer to use the term “maximum partial triple system”. In the case $\lambda = 1$, we will denote these systems as $\text{PTS}(v)$, $\text{MPTS}(v)$ and $\text{MMPTS}(v)$ respectively, and note that \mathcal{B} is then necessarily a *set* of triples, i.e. no triple can be repeated. The *leave* of an $\text{PTS}(v)$ (or $\text{MPTS}(v)$, $\text{MMPTS}(v)$) is the set of pairs of points that do not appear in any triple of the system.

If K is a set of positive integers, then a *pairwise balanced design*, $\text{PBD}(v, K, \lambda)$, of order v and index λ , is a pair (V, \mathcal{B}) , where V is a set of v elements (the *points*) and \mathcal{B} is a collection of subsets of V (the *blocks*) with the properties that every 2-element subset of V occurs precisely λ times amongst the blocks of \mathcal{B} , and if $B \in \mathcal{B}$ then $|B| \in K$. The elements of K are called the block sizes. If $\lambda = 1$, such a system is denoted more simply as $\text{PBD}(v, K)$. If the leave of a $\text{PTS}(v)$ is combined with its set of triples, then a pairwise balanced design $\text{PBD}(v, \{2, 3\})$ is obtained. Thus there is a rough equivalence between pairwise balanced designs with block sizes 2 and 3, and partial triple systems.

A *Steiner triple system* of order v , $\text{STS}(v)$, is an $\text{MMPTS}(v)$ in which each pair of points appears in some triple, that is to say the leave is empty. It is well known that the necessary and sufficient condition for the existence of an $\text{STS}(v)$ is that $v \equiv 1$ or $3 \pmod{6}$, and these values are called *admissible*. When $v \not\equiv 1$ or $3 \pmod{6}$, an $\text{STS}(v)$ does not exist and an $\text{MMPTS}(v)$ is, in a sense, the closest we can get to a Steiner triple system. When $v \equiv 5 \pmod{6}$ an $\text{MMPTS}(v)$ necessarily has a leave comprising four pairs of the form $\{a, b\}, \{b, c\}, \{c, d\}, \{d, a\}$ (see [1, page 553]). In this case, taking the pairs as edges of a graph on the four points, the leave can be represented as a 4-cycle (a, b, c, d) . For the cases $v \equiv 0$ or $2 \pmod{6}$, an $\text{MMPTS}(v)$ corresponds to an $\text{STS}(v+1)$ in which one point a has been deleted, resulting in a leave comprising $v/2$ disjoint pairs. For the case $v \equiv 4 \pmod{6}$ a similar correspondence holds between an $\text{MMPTS}(v)$ and an $\text{MMPTS}(v+1)$ where a point a from the leave

of the latter has been deleted giving the former a leave consisting of the three intersecting pairs $\{b, c\}$, $\{c, d\}$, $\{c, e\}$ and, disjoint from these, a further $(v-4)/2$ disjoint pairs.

It should be noted that maximum partial triple systems, MMPTS(v)s, are also called *optimal* $2-(v, 3, 1)$ *packings* (see [1, Section VI.40]). For $v \equiv 0, 1, 2, 3 \pmod{6}$, they are additionally known as (v_r, b_3) *configurations*, where b is the number of triples and r is the number of triples containing each point (see [1, Section VI.7]).

Two designs, (V_1, \mathcal{B}_1) and (V_2, \mathcal{B}_2) (of the same order v) are said to be isomorphic if there is a mapping ϕ from V_1 onto V_2 that takes blocks of \mathcal{B}_1 onto those of \mathcal{B}_2 . Such a mapping ϕ is called an isomorphism. An isomorphism from a design to itself is called an automorphism, and the set of all automorphisms of a given design forms a group under composition. This group is referred to as the automorphism group of the design. In discussing enumerations, it is generally the number of isomorphism classes of each design that is sought, in which case the point set may be fixed, for example as the set of integers $\{0, 1, \dots, v-1\}$.

As regards enumeration of these designs for small orders v , the situation for Steiner triple systems (up to isomorphism) is as follows. There are unique STS(v)s for $v = 3$, $v = 7$ and $v = 9$. There are two STS(13)s and 80 STS(15)s; the latter result [19] dates from 1919 and it was certainly not trivial to obtain this in the days prior to the advent of electronic computers. It took a further 85 years before the STS(19)s were enumerated in a paper by Kaski and Östergård [13]; there are 11 084 874 829 such systems and this result was obtained by a computer search. It seems unlikely that the next case, STS(21), will be enumerated any time soon since it is estimated that the number of these is several orders of magnitude greater than in the STS(19) case. It is known that the number of STS(v)s is $v^{v^2(\frac{1}{6}-o(1))}$ as $v \rightarrow \infty$ [20] (see also [3] Section 5.2.), a result that extends to MMPTS(v)s.

In the case when $v \equiv 5 \pmod{6}$, the MMPTS(5) is unique up to isomorphism, but there are two MMPTS(11)s [2] and these are given in Table 1. There are generally two types of MMPTS(v) designs for $v \equiv 5 \pmod{6}$. The first type is when the leave $\{a, b\}$, $\{b, c\}$, $\{c, d\}$, $\{d, a\}$ is accompanied by a pair of intersecting triples of the design, $\{a, c, e\}$ and $\{b, d, e\}$. Putting the 6 pairs from these two triples together with the leave gives a copy of the complete graph K_5 , and we will refer to this as the *quintuple* case, and such designs as being of Type-Q. The second type is when the leave is accompanied by a pair of disjoint triples of the design, $\{a, c, e\}$ and $\{b, d, f\}$ ($e \neq f$), and we will refer to this as the *non-quintuple* case, and such designs as being of Type-N. In Table 1, the design Q is of Type-Q, and the design N is of Type-N. In both cases the point set is $\{0, 1, \dots, 10\}$ (with 10 denoted by A) and the leave is the 4-cycle $(0, 1, 2, 3)$. Here and elsewhere, when listing pairs and triples, set brackets and/or commas may be omitted, so that $\{a, b, c\}$ may be written as abc . A reader with some patience can construct these two designs by hand and verify that there is precisely one of each type up to isomorphism.

Given the correspondence between MMPTS(v)s and MMPTS($v+1$)s, we be-

System	triples
Q	024, 056, 078, 09A, 134, 159, 168, 17A, 257, 269, 28A, 35A, 367, 389, 458, 46A, 479.
N	024, 057, 068, 09A, 135, 148, 169, 17A, 259, 26A, 278, 34A, 367, 389, 456, 479, 58A.

Table 1: The two maximum partial triple systems of order 11.

gin by enumerating MMPTS(17)s. Our results then enable us to enumerate the MMPTS(16)s. We first show that there are precisely 35 810 097 MMPTS(17)s. We find that there are 2 350 733 MMPTS(17)s of Type-Q, and 33 459 364 MMPTS(17)s of Type-N. These results were obtained by a computer search described in Section 2. The next value of $v \equiv 5 \pmod{6}$ is $v = 23$. The number of MMPTS(23)s is likely to be further orders of magnitude greater than the number of STS(21)s and so will probably remain out of reach for a considerable period of time.

There is a general construction of MMPTS(v)s for $v \equiv 5 \pmod{6}$ from PBD($v, \{3, 5\}$)s having precisely one block of size 5; such designs (denoted as PBD($v, \{3, 5^*\}$)s) exist for all $v \equiv 5 \pmod{6}$ (see [3] Section 6). If the block of size 5 is $\{a, b, c, d, e\}$, this can be replaced with four pairs ab, bc, cd and da and two triples ace and bde . There are fifteen distinct ways to do this, and the resulting systems may or may not be isomorphic. Conversely, an MMPTS(v) of Type-Q with leave (a, b, c, d) , and triples ace and bde can be converted to a PBD($v, \{3, 5^*\}$) having precisely one block of size 5. We take advantage of this correspondence in Section 4 to enumerate PBD(17, $\{3, 5^*\}$)s.

In the analysis of MMPTS(17)s (and other designs), the numbers of Pasch and mitre configurations are determined. A *Pasch* configuration in a PTS(v), also known as a *quadrilateral*, comprises a set of four triples of the system on six points, having the form abc, ade, bdf, cef . A *mitre* configuration has five triples on seven points, having the form abc, ade, afg, bdf, ceg ; the point a that appears in three triples will be called the *root* of the mitre and the other six points will be called the *leaves*. An (r, s) -configuration in a PTS(v) is a configuration having r triples and s points. So a Pasch configuration is a $(4, 6)$ -configuration, and in fact the only one. A mitre is a $(5, 7)$ -configuration and there is one other, obtained from a Pasch configuration by adding a triple afg . It is a longstanding conjecture regarding Steiner triple systems that, given $r \geq 4$, there exists v_r such that for all admissible $v \geq v_r$, there exists an STS(v) having no $(r', r' + 2)$ -configurations for all r' with $4 \leq r' \leq r$ [6]. Such an STS(v) is said to be *r-sparse*. A 4-sparse system is therefore one without Pasch configurations, and is also known as an *anti-Pasch* or *quadrilateral-free* system. An *anti-mitre* system is one without mitres, and a 5-sparse system is one without both mitres and Pasch configurations, equivalently it is both anti-Pasch and anti-mitre. It is

known that there are anti-Pasch STS(v)s for every admissible $v \neq 7, 13$ [16, 10], and an anti-mitre STS(v) for every admissible $v \neq 9$ [22]. It was shown in [21] that 5-sparse STS(v) exist for almost all admissible v , and their existence for orders $v \equiv 3 \pmod{6}$ ($v \neq 9, 15$) was established in [23]. Further results are given in [7], but the exact set of values v for which 5-sparse STS(v)s exist remains undetermined.

Given an MMPTS(17) and one of the points of its leave, by deleting this point and the 7 triples that contain it, an MMPTS(16) is formed. Since all MMPTS(16)s may be formed in this way, it is possible to determine all such systems from the MMPTS(17)s. After applying an isomorphism check, we establish that there are 47 744 568 MMPTS(16)s. Details of these systems are given in Section 5.

2 Searching for MMPTS(17)s

2.1 Seeding the search

The point set of each design was taken to be $V = \{0, 1, \dots, 16\}$ and the leave was taken to be $\{01, 03, 12, 23\}$. With equipment available to us at the present time it was not feasible to search all possible triples on V exhaustively, and this would anyway have been a colossal waste of resources. Instead, a small number of partial designs (*seeds*) were constructed that, without loss of generality, could potentially be extended to provide collectively at least one representative of each isomorphism class of MMPTS(17)s. These seeds were formed as follows.

Consider first the non-quintuple case. Without loss of generality, the leave and some of the triples of a non-quintuple MMPTS(17) may be taken as:

$$\{0, 1\}, \{0, 3\}, \{1, 2\}, \{2, 3\}, \{0, 2, 4\}, \{1, 3, 5\}, \{0, 5, 6\}, \\ \{0, 7, 8\}, \{0, 9, 10\}, \{0, 11, 12\}, \{0, 13, 14\}, \{0, 15, 16\}, \{2, 5, 7\}.$$

Again without loss of generality, the design may be taken to contain either the triple $\{2, 6, 8\}$ or $\{2, 6, 9\}$. For the first of these possibilities, the next triple may be taken to be $\{2, 9, 11\}$, while for the second possibility the next triple may be taken as either $\{2, 8, 10\}$ or $\{2, 8, 11\}$, all without loss of generality. The design requires three further blocks containing the point 2. By considering the possibilities it will be seen that, without loss of generality, the design must be an extension of one of the five possibilities N_1, N_2, N_3, N_4 and N_5 shown in Table 2.

Next consider the quintuple case. The argument is almost identical to the non-quintuple case. The only difference is that the triple $\{1, 3, 5\}$ is replaced by the triple $\{1, 3, 4\}$. Without loss of generality, the design must be an extension of one of the five possibilities Q_1, Q_2, Q_3, Q_{2a} and Q_4 shown in Table 3. The reason that the fourth of these alternatives is denoted Q_{2a} is that there exists an isomorphism of this design and Q_2 . An isomorphism taking Q_2 to Q_{2a} is given by the permutation $(5, 13)(6, 14)(7, 15, 10)(8, 16, 9)$. Consequently, design Q_{2a} is not required as a seed in an exhaustive search.

N_1	N_2	N_3	N_4	N_5
0, 2, 4	0, 2, 4	0, 2, 4	0, 2, 4	0, 2, 4
1, 3, 5	1, 3, 5	1, 3, 5	1, 3, 5	1, 3, 5
0, 5, 6	0, 5, 6	0, 5, 6	0, 5, 6	0, 5, 6
0, 7, 8	0, 7, 8	0, 7, 8	0, 7, 8	0, 7, 8
0, 9, 10	0, 9, 10	0, 9, 10	0, 9, 10	0, 9, 10
0, 11, 12	0, 11, 12	0, 11, 12	0, 11, 12	0, 11, 12
0, 13, 14	0, 13, 14	0, 13, 14	0, 13, 14	0, 13, 14
0, 15, 16	0, 15, 16	0, 15, 16	0, 15, 16	0, 15, 16
2, 5, 7	2, 5, 7	2, 5, 7	2, 5, 7	2, 5, 7
2, 6, 8	2, 6, 8	2, 6, 9	2, 6, 9	2, 6, 9
2, 9, 11	2, 9, 11	2, 8, 10	2, 8, 11	2, 8, 11
2, 10, 12	2, 10, 13	2, 11, 13	2, 10, 12	2, 10, 13
2, 13, 15	2, 12, 15	2, 12, 15	2, 13, 15	2, 12, 15
2, 14, 16	2, 14, 16	2, 14, 16	2, 14, 16	2, 14, 16

Table 2: Seeds for the computer search in the non-quintuple case.

Q_1	Q_2	Q_3	Q_{2a}	Q_4
0, 2, 4	0, 2, 4	0, 2, 4	0, 2, 4	0, 2, 4
1, 3, 4	1, 3, 4	1, 3, 4	1, 3, 4	1, 3, 4
0, 5, 6	0, 5, 6	0, 5, 6	0, 5, 6	0, 5, 6
0, 7, 8	0, 7, 8	0, 7, 8	0, 7, 8	0, 7, 8
0, 9, 10	0, 9, 10	0, 9, 10	0, 9, 10	0, 9, 10
0, 11, 12	0, 11, 12	0, 11, 12	0, 11, 12	0, 11, 12
0, 13, 14	0, 13, 14	0, 13, 14	0, 13, 14	0, 13, 14
0, 15, 16	0, 15, 16	0, 15, 16	0, 15, 16	0, 15, 16
2, 5, 7	2, 5, 7	2, 5, 7	2, 5, 7	2, 5, 7
2, 6, 8	2, 6, 8	2, 6, 9	2, 6, 9	2, 6, 9
2, 9, 11	2, 9, 11	2, 8, 10	2, 8, 11	2, 8, 11
2, 10, 12	2, 10, 13	2, 11, 13	2, 10, 12	2, 10, 13
2, 13, 15	2, 12, 15	2, 12, 15	2, 13, 15	2, 12, 15
2, 14, 16	2, 14, 16	2, 14, 16	2, 14, 16	2, 14, 16

Table 3: Seeds for the computer search in the quintuple case.

2.2 The searching programs

Given one of the nine seeds described in the previous section, the problem is to cover all uncovered pairs (apart from those in the leave) by triples. This can be represented as a graph covering problem by taking a graph G on the points $0, 1, \dots, 16$ whose edges are the pairs that are neither in the leave nor covered by triples in the seed. A solution corresponds to covering each edge of G precisely once with triangles of edges from G , and the program must find all possible solutions. A program written by the current authors was used for this purpose. If a point is reached where an uncovered pair has no remaining triples available to cover it, the program backtracks. If it reaches a point where all pairs are covered, it records the solution and then backtracks. The principal difficulty is adjusting the records for backtracks. In each of the nine cases, the program took about 90 minutes on a desktop computer to enumerate the solutions. It was then adapted to test solutions for isomorphisms and to record solutions as described below. It is prudent to verify computational results independently and so a minor variant of Knuth's *dancing links* graph covering program [14, 15] was used to check the number of solutions from each of the 9 runs; each of these checks took about 8 minutes.

The outcomes of the nine runs are shown in Table 4. The total number

Seed	Number of designs	Seed	Number of designs
N_1	161 885 696	Q_1	134 263 296
N_2	165 881 472	Q_2	140 978 304
N_3	166 118 112	Q_3	142 761 312
N_4	170 428 416	Q_4	144 371 376
N_5	171 571 376	Total	562 374 288
Total	835 885 072		

Table 4: Search results for the numbers of MMPTS(17)s.

of systems obtained in the nine runs was 1 398 259 360. No solution for the quintuple case can be isomorphic with any solution from the non-quintuple case, but within each class there are many isomorphisms.

2.3 Isomorphism testing

The software package *nauty* [18] was used to analyse the search outputs for isomorphisms. This treats each solution as a bipartite graph obtained from the design with an edge joining point x to triple B if and only if $x \in B$. Two designs are isomorphic if and only if the two bipartite graphs are isomorphic. Nauty assigns a canonical labelling to each graph so that two graphs are isomorphic if and only if they have the same canonical labelling. The canonical labellings

were stored, as they were produced, in a sorted list. The time taken to insert a new labelling into such a list is proportional to the logarithm of the number of objects in the list. If the labelling was already present in the list, the system was immediately recognised as an isomorphic copy and so could be discarded. Nauty is a long-established package with an excellent record for reliability, so there is a high degree of confidence in the results. Nevertheless, steps were taken to check these results.

Given a solution produced by the searching program, three numbers were obtained for each point x of the design: the number $n_p(x)$ of Pasch configurations incident with x , the number of mitres $n_r(x)$ with x as a root, and the number of mitres $n_l(x)$ with x as a leaf. In the quintuple case, any isomorphism between two solutions must preserve the point sets $V_1 = \{0, 1, 2, 3\}$, $V_2 = \{4\}$, $V_3 = \{5, 6 \dots, 16\}$, and in the non-quintuple case any isomorphism must preserve the point sets $V_1 = \{0, 1, 2, 3\}$, $V_2 = \{4, 5\}$, $V_3 = \{6, 7 \dots, 16\}$. Consequently the three sets of ordered triples $\{(n_p(x), n_r(x), n_l(x)) : x \in V_i\}$ for $i = 1, 2, 3$ form an invariant in each case. Using this invariant, it was easy to determine whether or not two solutions with the same invariant were isomorphic since the number of possible isomorphisms was small in every case. Samples of the search output were taken and the results produced by consideration of this invariant were found to agree with those produced by nauty. It had been hoped that this invariant might be complete, but this is not the case: there are non-isomorphic solutions sharing the same invariant.

Table 5 shows the number of isomorphism classes in the two cases. Altogether, up to isomorphism, there are precisely 35 810 097 MMPTS(17)s.

Non-quintuple case:	33 459 364
Quintuple case:	2 350 733
Total number:	35 810 097

Table 5: The numbers of isomorphism classes.

2.4 Storing the solutions

Storing a representative of each isomorphism class in the form of 44 triples requires about 14 GB of storage space. However this can be brought down to a modest 1.5 GB as follows. Given one of these designs, each triple $\{i, j, k\}$ is ordered so that $i < j < k$. These triples are taken in ascending order by their first entry, and then by their second. The list of 44 third entries is recorded with numbers represented by characters. The numbers 0-9 are represented as characters 0-9 and the numbers 10-16 are represented as characters a-g. In this way a design can be represented as a character string of length 44. In principle, not all 44 triples are needed because some triples are fixed in all 9 runs, but incorporating this saving isn't worth the extra effort. Compressing

the resulting files in .rar format reduces the required storage space to 13 MB for the quintuple case, and 179 MB for the non-quintuple case. The files are available as MMPTS(17)_Type-Q.rar and MMPTS(17)_Type-N.rar from [5].

3 Properties of the MMPTS(17)s

The number of Pasch configurations, mitres, and STS(7) subsystems (Fano planes) in each design was determined by two independently written programs.

In the quintuple case, 78 designs have no Pasch configurations, and 2 designs have the maximum number of 44 Pasch configurations. In the non-quintuple case, 843 designs have no Pasch configurations, and 1 design has the maximum number of 47 Pasch configurations. Hence there are anti-Pasch (4-sparse) MMPTS(17)s of both types. Table 6 gives the numbers of Type-Q and Type-N MMPTS(17)s having P Pasch configurations for $0 \leq P \leq 47$.

P	Type-Q	Type-N	P	Type-Q	Type-N	P	Type-Q	Type-N
0	78	843	16	106 560	1 487 105	32	105	868
1	487	7 531	17	76 724	1 072 999	33	35	578
2	2 415	37 136	18	52 071	745 422	34	39	303
3	8 161	124 920	19	34 839	501 176	35	42	177
4	21 834	326 465	20	21 773	325 369	36	9	133
5	46 990	696 112	21	13 707	206 993	37	4	87
6	85 332	1 258 629	22	8 038	128 418	38	11	32
7	133 629	1 962 982	23	4 957	78 709	39	9	34
8	185 730	2 697 978	24	2 782	48 182	40	6	6
9	227 426	3 292 317	25	1 919	29 380	41	1	17
10	255 310	3 642 666	26	1 088	17 815	42	0	14
11	261 623	3 697 358	27	676	11 220	43	4	1
12	248 943	3 477 394	28	423	6 664	44	2	1
13	220 388	3 053 390	29	271	4 032	45	0	4
14	182 791	2 528 980	30	208	2 456	46	0	1
15	143 166	1 984 888	31	127	1 578	47	0	1

Table 6: Pasch configuration frequencies in MMPTS(17)s

In the quintuple case, 4 designs have the minimum number of 5 mitres, and 3 designs have the maximum number of 48 mitres. In the non-quintuple case, 1 design has the minimum number of 3 mitres, and 1 design has the maximum number of 47 mitres. Note there are no anti-mitre and, consequently, no 5-sparse MMPTS(17)s. Table 7 gives the numbers of Type-Q and Type-N MMPTS(17)s having M mitres for $0 \leq M \leq 48$.

Table 8 gives the numbers of Type-Q and Type-N MMPTS(17)s having S STS(7) subsystems (also known as Fano planes) for $0 \leq S \leq 4$ (the maximum number in any MMPTS(17)).

The automorphism groups of the designs were also determined. The numbers of each order were checked by two independently written programs. Table 9 gives full details. The results in Table 9 can be also used to determine the total number of MMPTS(17)s on a specified point set from the orbit-stabiliser

M	Type-Q	Type-N
0	0	0
1	0	0
2	0	0
3	0	1
4	0	1
5	4	13
6	13	55
7	21	195
8	69	695
9	224	2368
10	533	6855
11	1367	18010
12	3406	43678
13	7139	98058
14	14637	199350
15	26544	373500

M	Type-Q	Type-N
16	44626	643122
17	69604	1020033
18	100089	1493054
19	134564	2026513
20	167649	2560458
21	197183	3014058
22	217341	3317425
23	225116	3422818
24	220267	3300754
25	205106	2992985
26	180070	2547896
27	150370	2042250
28	119906	1536133
29	89588	1086488
30	64205	720441
31	43397	449377

M	Type-Q	Type-N
32	27912	263661
33	17221	143731
34	10441	74012
35	5749	35097
36	3131	15668
37	1640	6630
38	852	2498
39	376	985
40	201	320
41	73	101
42	56	46
43	12	19
44	24	10
45	3	1
46	1	0
47	0	1
48	3	0

Table 7: Mitre frequencies in MMPTS(17)s

S	Type-Q	Type-N
0	2335059	33198472
1	15499	258750
2	125	2104
3	46	38
4	4	0

Table 8: STS(7) frequencies in MMPTS(17)s.

theorem as

$$\sum_D \frac{17!}{|Aut(D)|},$$

where $Aut(D)$ denotes the automorphism group of the design D , and D runs through the representatives of the isomorphism classes of the MMPTS(17)s. The number of Type-Q is 834 485 388 804 587 520 000 and the number of Type-N is 11 900 365 174 781 411 328 000. Hence the total number of MMPTS(17)s is 12 734 850 563 585 998 848 000.

Automorphisms		Type-Q	Type-N
order	group	number of systems	number of systems
12	dihedral D_{12}	3	0
8	dihedral D_8	20	0
8	$\mathbb{Z}_4 \times \mathbb{Z}_2$	4	0
6	dihedral $D_6 (= S_3)$	27	0
6	\mathbb{Z}_6	3	0
4	Klein	269	0
4	\mathbb{Z}_4	74	0
3	\mathbb{Z}_3	108	0
2	\mathbf{Z}_2	8 470	3 992
1	trivial	2 341 755	33 455 372

Table 9: Automorphisms of MMPTS(17)s.

4 The PBD(17, {3, 5*})s

The PBD(17, {3, 5*})s were enumerated by two methods. The first was to take the 2 350 733 non-isomorphic MMPTS(17)s of Type-Q, delete the blocks 024 and 134 (giving a leave forming the complete graph K_5 on the set $\{0, 1, 2, 3, 4\}$), and then test the resulting systems for isomorphisms. The second was to start again from Table 3, delete the blocks 024, 134 from each seed, generate systems from the seeds, and then test for isomorphisms. Isomorphisms were identified using nauty and checked with our own independent program. Both approaches gave the number of non-isomorphic PBD(17, {3, 5*})s as 157 151. Thus the vast majority of these designs must each generate 15 non-isomorphic MMPTS(17)s when two intersecting triples are formed from the leave K_5 in all possible ways. The non-isomorphic PBD(17, {3, 5*})s are available in compact form as strings of length 42 from [5] in the file PBD(17,{3,5}).rar (1 MB).

Analysis of these systems shows that there are 39 systems with no Pasch configurations, and 1 system with the maximum number of 34 Pasch configurations. Table 10 gives the number n_p of systems having P Pasch configurations for $0 \leq P \leq 34$.

P	n_p
0	39
1	259
2	911
3	2701
4	5440
5	9324
6	13619
7	17149
8	19363

P	n_p
9	19285
10	17904
11	14844
12	11935
13	8479
14	5958
15	3982
16	2468
17	1467

P	n_p
18	855
19	472
20	270
21	176
22	88
23	61
24	35
25	23
26	9

P	n_p
27	8
28	12
29	1
30	5
31	5
32	1
33	2
34	1

Table 10: Pasch configuration frequencies in PBD(17, {3, 5*})s.

The minimum number of mitres in a PBD(17, {3, 5*}) is 5 (10 systems have this number), and 2 systems have the maximum number of 36 mitres. Table 11 gives the number n_m of systems having M mitres for $5 \leq M \leq 36$.

M	n_m
5	10
6	19
7	52
8	153
9	430
10	1001
11	2211
12	4026

M	n_m
13	6640
14	10146
15	13561
16	16615
17	18614
18	18821
19	17540
20	15142

M	n_m
21	11618
22	8449
23	5448
24	3306
25	1809
26	892
27	386
28	171

M	n_m
29	54
30	22
31	2
32	9
34	1
35	1
36	2

Table 11: Mitre frequencies in PBD(17, {3, 5*})s.

Each PBD(17, {3, 5*}) has either $S = 0$ or $S = 1$ STS(7) subsystems (Fano planes). Table 12 gives the number n_s of systems having S STS(7) subsystems.

S	n_s
0	156713
1	438

Table 12: STS(7) frequencies in PBD(17, {3, 5*})s.

Table 13 tabulates the automorphism groups of the PBD(17, {3, 5*})s. Again, the numbers of each order were checked by two independently written programs. The groups of order 24 were identified using the GAP system [8]. The symbol \times denotes a semidirect product. The total number of PBD(17, {3, 5*})s on a specified point set can be obtained from the orbit-stabiliser theorem, and is 55 632 359 253 639 168 000.

Automorphisms		number of PBD(17, {3, 5*})s
order	group	
24	S_4	2
24	$\mathbb{Z}_4 \times S_3$	1
24	$(\mathbb{Z}_6 \times \mathbb{Z}_2) \rtimes \mathbb{Z}_2$	1
12	\mathbb{Z}_{12}	2
12	dihedral D_{12}	2
12	$\mathbb{Z}_3 \rtimes \mathbb{Z}_4$	2
8	dihedral D_8	5
8	$\mathbb{Z}_4 \times \mathbb{Z}_2$	3
6	\mathbb{Z}_6	15
6	dihedral $D_6 (= S_3)$	18
4	\mathbb{Z}_4	19
4	Klein	63
3	\mathbf{Z}_3	64
2	\mathbf{Z}_2	1 190
1	trivial	155 764

Table 13: Automorphisms of PBD(17, {3, 5*})s.

5 The MMPTS(16)s

As explained in the Introduction, the MMPTS(16)s can be obtained by deleting one of the points 0, 1, 2 or 3 of each of the MMPTS(17)s described in Section 2. This results in $4 \times 35\,810\,097 = 143\,240\,388$ systems. But amongst these there are isomorphisms, and nauty was used to eliminate isomorphic copies. It was found that there are 47 744 568 non-isomorphic solutions. The dancing links program was used to check this number with an independent search. Of these solutions, 3 133 686 originate from MMPTS(17)s of Type-Q, and the remaining 44 610 882 are from systems of Type-N. If an MMPTS(16) originates from an MMPTS(17) of Type-Q, then adding a new point to each of the seven non-intersecting pairs and to one of the three intersecting pairs of its leave gives a Type-Q MMPTS(17). Hence no MMPTS(16) originating from a quintuple type MMPTS(17) can be isomorphic to one originating from a non-quintuple type MMPTS(17). A MMPTS(16) derived from a quintuple type MMPTS(17) will be described as Type-Q, and a MMPTS(16) derived from a non-quintuple type MMPTS(17) will be described as Type-N. The designs are available in files MMPTS(16)_Type-Q.rar (17 MB) and MMPTS(16)_Type-N.rar (208 MB) from [5]. The designs are mapped so that the leave is $\{01, 02, 03, 45, 67, \dots, ef\}$, with characters a to f representing points 10 to 15, and the designs are specified in compact form as strings of length 37.

For MMPTS(16)s of Type-Q, 1 346 designs have no Pasch configurations, and 1 design has the maximum number of 43 Pasch configurations. For MMPTS(16)s of type-N, 15 877 designs have no Pasch configurations, and 1 design has the

maximum number of 44 Pasch configurations. Hence there are anti-Pasch (4-sparse) MMPTS(16)s of both types. Table 14 gives the numbers of Type-Q and Type-N MMPTS(16)s having P Pasch configurations for $0 \leq P \leq 44$.

P	Type-Q	Type-N	P	Type-Q	Type-N	P	Type-Q	Type-N
0	1 346	15 877	15	36 534	565 757	30	29	314
1	9 571	125 270	16	22 010	344 867	31	18	163
2	38 020	502 936	17	12 897	207 237	32	14	125
3	100 154	1 347 901	18	7 562	123 640	33	10	76
4	198 137	2 704 815	19	4 448	73 743	34	2	38
5	309 441	4 278 201	20	2 606	43 954	35	5	36
6	395 447	5 549 182	21	1 661	26 351	36	1	17
7	432 141	6 121 166	22	1 013	15 917	37	1	12
8	414 833	5 905 615	23	653	9 497	38	2	10
9	358 284	5 122 025	24	355	5 761	39	2	3
10	282 328	4 069 661	25	264	3 480	40	1	3
11	206 951	3 016 887	26	148	2 096	41	1	5
12	142 368	2 113 200	27	120	1 327	42	0	1
13	94 556	1 407 703	28	85	755	43	1	0
14	59 617	904 747	29	49	510	44	0	1

Table 14: Pasch configuration frequencies in MMPTS(16)s

For MMPTS(16)s of Type-Q, 2 designs have no mitres, and 1 design has the maximum number of 48 mitres. For MMPTS(16)s of type-N, 5 designs have no mitres, and 1 design has the maximum number of 45 mitres. Hence there are anti-mitre MMPTS(16)s of both types. Table 15 gives the numbers of Type-Q and Type-N MMPTS(16)s having M mitres for $0 \leq M \leq 48$. The 7 designs with no mitres all have Pasch configurations, so there are no 5-sparse MMPTS(16)s.

M	Type-Q	Type-N	M	Type-Q	Type-N	M	Type-Q	Type-N
0	2	5	16	335 963	4 967 389	32	153	469
1	3	14	17	289 977	4 289 496	33	41	310
2	26	119	18	231 425	3 427 280	34	77	153
3	116	879	19	172 907	2 532 692	35	4	89
4	526	5 005	20	121 074	1 746 307	36	35	87
5	2 165	21 590	21	79 603	1 122 077	37	8	16
6	6 559	73 898	22	50 082	677 356	38	10	23
7	17 563	206 848	23	29 615	385 028	39	1	15
8	39 376	488 364	24	16 792	205 365	40	9	3
9	77 123	988 266	25	9 284	103 465	41	0	6
10	131 409	1 741 921	26	4 974	50 430	42	3	14
11	198 113	2 700 157	27	2 591	23 424	43	0	1
12	267 765	3 725 024	28	1 548	10 649	44	1	0
13	324 803	4 620 691	29	695	4 811	45	0	1
14	359 205	5 184 315	30	441	2 148	46	0	0
15	361 423	5 303 746	31	195	936	47	0	0
						48	1	0

Table 15: Mitre frequencies in MMPTS(16)s

Table 16 gives the numbers of Type-Q and Type-N MMPTS(16)s having S

STS(7) subsystems (Fano planes) for $0 \leq S \leq 4$ (the maximum number in any MMPTS(16)).

S	Type-Q	Type-N
0	3 120 879	44 379 670
1	12 703	229 796
2	80	1 392
3	22	24
4	2	0

Table 16: STS(7) frequencies in MMPTS(16)s.

Table 17 tabulates the automorphism groups of the MMPTS(16)s. Again, the numbers of each order were checked by two independently written programs. The total number of MMPTS(16)s on a specified point set can be obtained from the orbit-stabiliser theorem, and this gives the number of Type-Q systems as 65 449 834 416 046 080 000 and the number of Type-N systems as 933 361 974 492 659 712 000. Thus the overall total number of MMPTS(16)s is 998 811 808 908 705 792 000.

Automorphisms		Type-Q	Type-N
order	group	number of systems	number of systems
12	dihedral D_{12}	9	0
12	alternating A_4	2	0
12	$\mathbb{Z}_6 \times \mathbb{Z}_2$	1	0
6	dihedral $D_6 (= S_3)$	33	6
6	\mathbb{Z}_6	28	0
4	Klein	470	0
4	\mathbb{Z}_4	32	0
3	\mathbb{Z}_3	280	504
2	\mathbf{Z}_2	9 802	1 434
1	trivial	3 123 029	44 608 938

Table 17: Automorphisms of MMPTS(16)s.

6 Concluding remarks

Details of MMPTS(v)s for $v \leq 11$ are given in [2]. In this Section we give some details of the systems of orders 12 and 14 that have been enumerated elsewhere. As recorded in [9], there are precisely five non-isomorphic MMPTS(12)s arising from the two non-isomorphic STS(13)s, but these are not explicitly listed. Table 18 gives them here in compact form with the numbers of Pasch configurations and mitres that they contain, and their automorphism groups. In each case the

leave is $\{01, 23, \dots, ab\}$ with a, b representing points 10, 11. Since no STS(13) has an STS(7) subsystem, neither do these MMPTS(12)s. The total number of MMPTS(12)s on a specified point set is 1 197 504 000.

MMPTS(12)	#Pasch	#mitre	Automorphism group
468ab798abb8ab7aa99b	4	6	\mathbb{Z}_2
468ab79a6b9baba889ba	5	3	\mathbb{Z}_2
468ab78ab96ba9abb89a	5	3	dihedral $D_6 (= S_3)$
468ab94ba87baa9b8ba9	4	5	trivial
468aba498b79bb9ab8aa	7	0	\mathbb{Z}_3

Table 18: The non-isomorphic MMPTS(12)s.

There are 787 non-isomorphic MMPTS(14)s, and these are enumerated in [11] where they are recorded as $(14_6, 28_3)$ configurations. Here we give counts for Pasch configurations, mitres and STS(7) subsystems, and details of the automorphism groups. The designs are available in the file MMPTS(14).txt (23KB) from [5]. They are mapped so that the leave is $\{01, 23, \dots, cd\}$ with characters a to d representing points 10 to 13, and they are specified in compact form as strings of length 28.

There are 6 MMPTS(14)s with no Pasch configurations, and 1 MMPTS(14) has the maximum number of 63 Pasch configurations. Table 19 gives the number n_p of MMPTS(14)s having P Pasch configurations for $0 \leq P \leq 63$.

There are 15 MMPTS(14)s with no mitres, and 1 MMPTS(14) has the maximum number of 22 mitres. Table 20 gives the number n_m of MMPTS(14)s having M mitres for $0 \leq M \leq 22$. The 15 designs with no mitres all have Pasch configurations, so there are no 5-sparse MMPTS(14)s.

Table 21 gives the number n_s of MMPTS(14)s having S STS(7) subsystems (Fano planes) for $0 \leq S \leq 8$ (the maximum number in any MMPTS(14)).

Table 22 tabulates the automorphism groups of the MMPTS(14)s. Again, the numbers of each order were checked by two independently written programs.

P	n_p	P	n_p	P	n_p	P	n_p
0	6	9	36	19	12	29	2
1	7	10	26	20	7	31	7
2	34	11	30	21	5	33	1
3	94	12	13	22	4	35	1
4	94	13	29	23	10	39	1
5	117	14	23	25	2	43	1
6	72	15	9	26	1	47	1
7	62	16	3	27	2	63	1
8	56	17	17	28	1		

Table 19: Pasch configuration frequencies in MMPTS(14)s

M	n_m
0	15
2	5
3	8
4	24
5	12

M	n_m
6	44
7	35
8	73
9	87
10	90

M	n_m
11	96
12	100
13	71
14	51
15	40

M	n_m
16	23
17	7
18	1
19	3
20	1
22	1

Table 20: Mitre frequencies in MMPTS(14)s

S	n_s
0	730
1	43
2	11
4	2
8	1

Table 21: STS(7) frequencies in MMPTS(14)s.

The groups of orders 24 and above were identified using the GAP system [8]. The symbol \rtimes denotes a semidirect product. The system with the automorphism group of order 1344 is the system obtained from the projective STS(15) that has an automorphism group A_8 , the alternating group of order 20160, and whose automorphism partition consists of a single part containing all 15 points; this is STS(15) number 1 in the listing of [17]. The total number of MMPTS(14)s on a specified point set can be obtained from the orbit-stabiliser theorem, and is 60 281 712 691 200.

Collecting together previous results and the results of this paper, Table 23 gives the numbers of non-isomorphic MMPTS(v)s for $v \leq 19$. The entry for $v = 18$ comes from a personal communication [12] cited in [11] and given also in [1, page 354].

Automorphisms		number of MMPTS(14)s
order	group	
1344	$\mathbb{Z}_2^3 \rtimes PSL(3, 2)$	1
192	$((\mathbb{Z}_2 \times D_8) \rtimes \mathbb{Z}_2) \rtimes \mathbb{Z}_3 \rtimes \mathbb{Z}_2$	1
96	$(\mathbb{Z}_2^2 \times A_4) \rtimes \mathbb{Z}_2$	1
32	$(\mathbb{Z}_2^4) \rtimes \mathbb{Z}_2$	3
24	symmetric S_4	3
24	$A_4 \times \mathbb{Z}_2$	2
21	$\mathbb{Z}_7 \rtimes \mathbb{Z}_3$	2
16	\mathbb{Z}_2^4	1
12	alternating A_4	4
8	dihedral D_8	10
6	dihedral $D_6 (= S_3)$	1
6	\mathbb{Z}_6	1
4	Klein	18
4	\mathbb{Z}_4	13
3	\mathbb{Z}_3	37
2	\mathbb{Z}_2	40
1	trivial	649

Table 22: Automorphisms of MMPTS(14)s.

v	The number of non-isomorphic MMPTS(v)s
≤ 9	1
10	2
11	2
12	5
13	2
14	787
15	80
16	47 744 568
17	35 810 097
18	210 611 385 743
19	11 084 874 829

Table 23: The number of non-isomorphic MMPTS(v)s.

References

- [1] C. J. Colbourn and J. H. Dinitz. The CRC Handbook of Combinatorial Designs (2nd edition), CRC Press, 2007.
- [2] C. J. Colbourn and A. Rosa. Maximal partial Steiner triple systems of order $v \leq 11$. *Ars Combin.* 20, (1985) 5–28.
- [3] C. J. Colbourn and A. Rosa. Triple Systems. Oxford University Press, 1999.
- [4] C. J. Colbourn, E. Mendelsohn, A. Rosa and J. Siran. Anti-mitre Steiner triple systems. *Graphs Combin.* 10 (1994) 215–224.
- [5] F. Demirkale, D. M. Donovan and M. J. Grannell. Maximum partial triple systems on 14, 16 and 17 points. <http://www.yildiz.edu.tr/~fatihd/MMPTS/>
- [6] P. Erdős. Problems and results in combinatorial analysis. *Colloquio Internazionale sulle Teorie Combinatorie (Rome, 1973)*, Tomo II, *Atti dei Convegni Lincei*, No. 17, (1976) 3–17.
- [7] Y. Fujiwara. Infinite classes of anti-mitre and 5-sparse Steiner triple systems. *J. Combin. Des.* 14(3) (2006) 237-250.
- [8] GAP - Groups, Algorithms, Programming - a System for Computational Discrete Algebra. <https://www.gap-system.org/>
- [9] M. J. Grannell, T. S. Griggs, K. A. S. Quinn and R. G. Stanton. A census of minimal pair-coverings with restricted largest block length. *Ars Combin* 52 (1999) 71–96.
- [10] M. J. Grannell, T. S. Griggs and C. A. Whitehead. The resolution of the anti-Pasch conjecture. *J. Combin. Des.* 8(4) (2000) 300–309.
- [11] H. Gropp. Existence and enumeration of configurations. *Bayreuth. Math. Schr.* 74 (2005) 123–129.
- [12] P. Kaski. Personal communication to Harald Gropp. April 2005.
- [13] P. Kaski and P. Östergård. The Steiner triple systems of order 19. *Math. Comp.* 73 (2004) no.248 2075–2092.
- [14] D. E. Knuth. Dancing links, *in* *Millennial Perspectives in Computer Science: Proceedings of the 1999 Oxford-Microsoft Symposium in Honour of Sir Tony Hoare*. J. Davies, B. Roscoe, and J. Woodcock, editors. Palgrave, 2000, 187–214.
- [15] D. E. Knuth. Selected papers on fun & games. *CSLI Lecture Notes 192*, CSLI Publications, Stanford, CA, 2011.

- [16] A. C. H. Ling, C. J. Colbourn, M. J. Grannell and T. S. Griggs. Construction techniques for anti-Pasch Steiner triple systems. *J. London Math. Soc.* (2), 61 (2000) 641–657.
- [17] R. A. Mathon, K. T. Phelps and A. Rosa. Small Steiner triple systems and their properties. *Ars Combin.* 15 (1983) 3–110.
- [18] B. D. McKay and A. Piperno. Nauty and Traces users guide (Version 2.5). Computer Science Department, Australian National University, Canberra, Australia (2013).
- [19] H. S. White, F. N. Cole and L. D. Cummings. Complete classification of the triad systems on fifteen elements. *Memoirs Nat. Acad. Sci. U.S.A.* 14 (1919) 1–89.
- [20] R. M. Wilson. Nonisomorphic Steiner triple systems. *Math. Zeitschr.* 135 (1973/4) 303–313.
- [21] A. J. Wolfe, 5-Sparse Steiner triple systems of order n exist for almost all admissible n . *Electron. J. Combin.* 12 (2005) #R68, 42 pp. (electronic).
- [22] A. J. Wolfe. The resolution of the anti-mitre Steiner triple system conjecture. *J. Combin. Des.* 14(3) (2006) 229–236.
- [23] A. J. Wolfe, The existence of 5-sparse Steiner triple systems of order $n \equiv 3 \pmod{6}$, $n \notin \{9, 15\}$. *J. Combin. Theory, Ser. A* 115(8) (2008) 1487–1503.