

Spanning closed walks with bounded maximum degrees of graphs on surfaces

MORTEZA HASANVAND

Department of Mathematical Sciences, Sharif University of Technology, Tehran, Iran

hasanvand@alum.sharif.edu

Abstract

Gao and Richter (1994) showed that every 3-connected graph which embeds on the plane or the projective plane has a spanning closed walk meeting each vertex at most 2 times. Brunet, Ellingham, Gao, Metzlar, and Richter (1995) extended this result to the torus and Klein bottle. Sanders and Zhao (2001) obtained a sharp result for higher surfaces by proving that every 3-connected graph embeddable on a surface with Euler characteristic $\chi \leq -46$ admits a spanning closed walk meeting each vertex at most $\lceil \frac{6-2\chi}{3} \rceil$ times. In this paper, we develop these results to the remaining surfaces with Euler characteristic $\chi \leq 0$.

Keywords:

Graphs on a surface; spanning walk; spanning trail; connectivity.

1 Introduction

In this article, all graphs have no loop, but multiple edges are allowed and a simple graph is a graph without multiple edges. Let G be a graph. The vertex set, the edge set, and the number of components of G are denoted by $V(G)$, $E(G)$, and $\omega(G)$, respectively. For a vertex set S of G , we denote by $e_G(S)$ the number of edges of G with both ends in S . Also, S is called an *independent set*, if there is no edge of G connecting vertices in S . A *minor* of G refers to a graph R which can be obtained from G by contracting some vertex-disjoint connected subgraphs of G . A *k-walk* (*k-trail*) in a graph refers to a spanning closed walk (trail) meeting each vertex at most k times. In this paper, we assume that all walks use each edge at most two times. Note that 1-walks and

1-trails of a graph are equivalent to Hamilton cycles. A graph G is called m -tree-connected, if it has m edge-disjoint spanning trees. It was known that the vertex set of any graph G can be expressed uniquely (up to order) as a disjoint union of vertex sets of some induced m -tree-connected subgraphs [2]. These subgraphs are called the m -tree-connected components of G . For a graph G , we define $\Omega(G) = |P| - \frac{1}{2}e_G(P)$, in which P is the unique partition of $V(G)$ obtained from the 2-tree-connected components of G and $e_G(P)$ denotes the number of edges of G joining different parts of P .

In 1956 Tutte [18] showed that every 4-connected plane graph admits a Hamilton cycle. Later, Gao and Richter (1994) [4] proved that every 3-connected graph which embeds on the plane or the projective plane admits a 2-walk, which was conjectured by Jackson and Wormald (1990) [10]. In 1995 Brunet, Ellingham, Gao, Metzlar, and Richter [1] extended this result to the torus and Klein bottle, and also proposed the following conjecture. This conjecture is verified in [3, 15] with linear bounds on k_χ (but not sharp).

Conjecture 1.1.([1]) *For every integer χ , there is a positive integer k_χ such that every 3-connected graph which embeds on a surface with Euler characteristic χ admits a k_χ -walk.*

In 2001 Sander and Zhao [16] obtained a sharp bound on k_χ for surfaces with small enough Euler characteristic and established the following theorem. In Section 3, we develop this result to the surfaces with Euler characteristic $\chi \leq 0$. Our proof is based on a recent result in [9] and inspired by some methods that introduced in [13, 16].

Theorem 1.2.([16]) *Every 3-connected graph G embeddable on a surface with Euler characteristic $\chi \leq -46$ has a $\lceil \frac{6-2\chi}{3} \rceil$ -walk.*

In 1994 Gao and Wormald [5] investigated trails in triangulations and derived the following theorem. They also deduced that every 5-connected triangulation in the double torus with representativity at least 6 admits a 4-trail. In Section 4, we conjecture that for each χ , there is a positive integer k_χ such that every 4-connected graph which embeds on a surface with Euler characteristic χ admits a k_χ -trail, and verify it for 5-connected graphs. As a consequence, we deduce that every 5-connected graph which embeds on the double torus (not necessarily triangulation) admits a 3-trail.

Theorem 1.3.([5]) *All triangulations in the projective plane, the torus and the Klein bottle have 4-trails.*

2 Minor minimal 3-connected graphs having no k -walks

The following theorem is inspired by Theorem 5 in [13] and provides a common improvement for Lemmas 3.1 and 3.2 in [16]. We will apply it in the next section.

Theorem 2.1. *Let G be a 3-connected graph and let k be an integer with $k \geq 3$. If G has no k -walks, then G contains a minor 3-connected bipartite graph R with the bipartition (X, Y) with the following properties:*

1. R has no k -walks.
2. $X = \{v \in V(G) : d_R(v) \geq k + 1\}$.
3. $Y = \{v \in V(G) : d_R(v) = 3\}$.

Let G be a 3-connected graph. For an edge $e \in E(G)$, define G/e to be the graph obtained from G by contracting e . An edge e is said to be *contractible* if G/e is still 3-connected. For proving Theorem 2.1, we require the following two lemmas, which the first one is well-known.

Lemma 2.2.(See Halin [7]) *Let G be a 3-connected graph except for K_4 . Then every vertex of degree 3 in G is incident with a contractible edge.*

Lemma 2.3.(Halin [7, 8]) *Let G be a minimally 3-connected graph and define $V_3(G) = \{v \in V(G) : d_G(v) = 3\}$. Then the following statements hold:*

- (i) $V_3(G) \neq \emptyset$.
- (ii) *Every edge connecting two vertices in $V(G) \setminus V_3(G)$ is contractible.*
- (iii) *The graph obtained by contracting any edge connecting two vertices in $V(G) \setminus V_3(G)$ is also minimally 3-connected.*
- (iv) *Every cycle of G contains at least two vertices of $V_3(G)$.*

Proof of Theorem 2.1. Let G be a counterexample with the minimum $|V(G)|$. We may assume that G is minimally 3-connected and also $|V(G)| \geq 5$. We here prove the following claim which was essentially shown by Sanders and Zhao in [16, Lemma 3.2].

Claim 1. $\{v \in V(G) : 3 \leq d_G(v) \leq k\}$ is independent.

Suppose otherwise that there is an edge $xy \in E(G)$ such that $d_G(x) \leq k$ and $d_G(y) \leq k$. If all edges incident with x or y are not contractible, then by Lemma 2.2, we must have $d_G(x) > 3$ and $d_G(y) > 3$, which contradicts Lemma 2.3 (ii). Thus we may assume that there is a contractible edge incident with y , say yz (possibly $z = x$). By the minimality of G , the graph G/yz has a k -walk W . Define H to be the graph with the vertex set $V(G)$ having the same edges of W by considering multiplicity of each edge. First, assume $d_H(y) > 0$. If both of $d_H(y)$ and $d_H(z)$ are odd, define H' to be the graph obtained from H by adding a copy of yz ; otherwise, define H' to be the graph obtained from H by adding two copies of yz . Next, assume $d_H(y) = 0$. In this case, define H' to be the graph obtained from H by adding two copies of xy . It is not difficult to check that H' is an Eulerian graph with $\Delta(H') \leq 2k$. Hence G admits a k -walk, which is a contradiction.

We now prove the next claim.

Claim 2. $\{v \in V(G) : d_G(v) \geq 4\}$ is independent.

Let $P = x_0x_1 \dots x_l$ be a maximal path in the subgraph of G induced by $\{v \in V(G) : d_G(v) \geq 4\}$. By Lemma 2.3 (iv), this subgraph is a forest. Also, x_0 and x_l have degree one in it and $x_ix_j \notin E(G)$, for any $0 \leq i < j \leq l$ and $j \neq i + 1$. By applying Lemma 2.3 (iii) repeatedly, one can conclude that G/P is 3-connected. By the minimality of G , the graph G/P has a k -walk W . Define H to be the graph with the vertex set $V(G)$ having the same edges of W by considering multiplicity of each edge. Since x_0 and x_l have degree one in P , there are two edges $x_0y_1, x_ly_2 \in E(G)$ such that $y_1, y_2 \in Y$. Note that $\sum_{0 \leq i \leq l} d_H(x_i) \leq 2k$. Let x_j be a vertex of $V(P)$ with the maximum $d_H(x_j)$. If $d_H(x_j) \leq 2k - 3$, define H' to be the graph obtained from H by adding some of the edges of a copy of P and adding another new copy of P such that H' forms an Eulerian graph. Otherwise, if $d_H(x_j) \geq 2k - 2$, define H' to be the graph obtained from H by adding a copy of the paths $y_1x_0Px_{i-1}$ (if $i \neq 1$) and $x_{i+1}Px_ly_2$ (if $i \neq l$) and adding some of the edges of y_1Py_2 such that H' forms an Eulerian graph. According to the construction, it is not difficult to check $\Delta(H') \leq 2k + 1$. Hence G admits a k -walk, which is a contradiction again.

By the above-mentioned claims, G is a bipartite graph with the bipartition (X, Y) in which $X = \{v \in V(G) : d_R(v) \geq k + 1\}$ and $Y = \{v \in V(G) : d_R(v) = 3\}$. By taking $R = G$, we derive that G is not a counterexample and so the proof is completed. \square

3 3-connected graphs on surfaces

We shall below develop Theorem 1.2 as mentioned in the Abstract. For this purpose, we recall the following recent result from [9] that grants the existence of walks with bounded maximum degrees on specified independent vertex set.

Lemma 3.1.([9]) *Let G be a connected graph with the independent set $X \subseteq V(G)$ and let k be a positive integer. Then G contains an spanning closed walk meeting each $v \in X$ at most k times, if for every $S \subseteq X$, at least one of the following conditions holds:*

1. $\omega(G \setminus S) \leq (k - \frac{1}{2})|S| + 1$.
2. G contains an spanning closed walk meeting each $v \in S$ at most k times.

Now, we are in a position to prove the main result of this paper.

Theorem 3.2. *Every 3-connected graph G embeddable on a surface with Euler characteristic $\chi \leq 0$ admits a $\lceil \frac{6-2\chi}{3} \rceil$ -walk.*

Proof. We may assume that $\chi < 0$, as the assertion was already proved in [1] for the surfaces with Euler characteristic zero (namely the torus and Klein bottle). By Theorem 2.1, we may also assume that G is a bipartite graph with the bipartition (X, Y) such that $X = \{v \in V(G) : d_G(v) \geq k + 1\}$ and $Y = \{v \in V(G) : d_G(v) = 3\}$, where $k = \lceil \frac{6-2\chi}{3} \rceil$. Let $S \subseteq X$ be a vertex cut of G so that $|S| \geq 3$. Define H to be the bipartite simple graph obtained from G by contracting any component of $G \setminus S$ such that one partite set is S . Note that H can be embedded on the surface as G is embedded. Since G is 3-connected, the minimum degree of H is at least 3. Since H is triangle-free, by Euler's formula, it is easy to check that $|E(H)| \leq 2|V(G)| - 2\chi$. Thus

$$3\omega(G \setminus S) \leq |E(H)| \leq 2|V(H)| - 2\chi = 2(|S| + \omega(G \setminus S)) - 2\chi,$$

and so

$$\omega(G \setminus S) \leq 2|S| - 2\chi \leq (2 + \frac{-2\chi - 3/2}{3})|S| + 3/2 \leq (\lceil \frac{6-2\chi}{3} \rceil - 1/2)|S| + 3/2.$$

If the equalities hold, then we must have $|S| = 3$ and $\omega(G \setminus S) = 6 - 2\chi$, and also every component of $G \setminus S$ has exactly 3 neighbours in S . In this case, we shall show that G has a spanning closed walk meeting each $v \in S$ at most k times. Since G is 3-connected, there is a cycle C of G containing all

three vertices of S . Since S is independent, the cycle C contains at least one vertex of exactly three components of $G \setminus S$. For the remaining $3 - 2\chi$ components of $G \setminus S$, join every of them to C such that each vertex in S has degree at most $\lceil (3 - 2\chi)/3 \rceil + 2$. Finally, add the edges of any component of $G \setminus S$ to this new graph. Obviously, the resulting spanning subgraph of G has a spanning closed walk meeting each $v \in S$ at most k times and so does G . Therefore, by Lemma 3.1, the graph G have a spanning closed walk meeting each $v \in X$ at most k times. Since $k \geq 3$ and every vertex of Y in G has degree 3, this walk forms a k -walk for G . Hence the theorem holds. \square

4 Graphs with higher connectivity

As we already mentioned, Tutte [18] proved that every 4-connected planar graph admits a Hamilton cycle. Thomas and Yu [17] extended this result to the projective plane and Grünbaum [6] and independently Nash-Williams [12] conjectured that Tutte's result could be developed to the torus. Motivated by Conjecture 1.1 and these results, we now propose the following conjecture. By considering the complete bipartite graph $K_{4, -\chi+4}$ (which can be embedded on a surface with Euler characteristic χ), if this conjecture would be true, then we must have $k_\chi \geq \lceil \frac{4-\chi}{4} \rceil$.

Conjecture 4.1. *For every integer χ , there is a positive integer k_χ such that every 4-connected graph which embeds on a surface with Euler characteristic χ admits a k_χ -trail.*

We shall below prove Conjecture 4.1 for 5-connected graphs. For this purpose, we require the following recent result which gives a sufficient condition for the existence of k -trails.

Lemma 4.2.([9]) *Let G be a graph and let k be a positive integer. If for all $S \subseteq V(G)$,*

$$\Omega(G \setminus S) \leq (k - \frac{1}{2})|S| + 1 - \frac{1}{2}e_G(S),$$

then G admits a k -trail.

Now, we are ready to prove the main result of this section.

Theorem 4.3. *Every 5-connected graph G which embeds on a surface with Euler characteristic $\chi \leq 0$ admits a $\lceil \frac{6-3\chi}{4} \rceil$ -trail.*

Proof. Let $S \subseteq V(G)$ with $\Omega(G \setminus S) > 1$. For each vertex v , we have $\Omega(G - v) = 1$, because the graph $G - v$ is 2-tree-connected [11, 19]. Thus $|S| \geq 2$. Define H to be the simple graph

obtained from G by contracting the 2-tree-connected components of $G \setminus S$. Set $S' = V(H) \setminus S$. Note that $|V(H)| = |S| + |S'| \geq 3$ and H can be embedded on the surface as G is embedded. By Euler's formula, it is easy to check that $|E(H)| \leq 3|V(H)| - 3\chi$. Note also that there is no pair of edges of G joining two different 2-tree-connected components of $G \setminus S$. Since G is 5-connected, the minimum degree of H is at least 5. Thus

$$5|S'| \leq |E(H)| - e_H(S) + e_H(S'),$$

and so

$$5|S'| - e_H(S') \leq |E(H)| - e_H(S) \leq 3|V(H)| - 3\chi - e_H(S) = 3(|S'| + |S|) - 3\chi - e_G(S).$$

Therefore,

$$\Omega(G \setminus S) = |S'| - \frac{1}{2}e_H(S') \leq \frac{3}{2}|S| + \frac{-3\chi}{2} - \frac{1}{2}e_G(S).$$

and so

$$\Omega(G \setminus S) \leq \left(\frac{3}{2} + \frac{-3\chi/2 - 1}{2}\right)|S| + 1 - \frac{1}{2}e_G(S) \leq \left(\lceil \frac{6-3\chi}{4} \rceil - \frac{1}{2}\right)|S| + 1 - \frac{1}{2}e_G(S).$$

Hence the theorem follows from Lemma 4.2 with $k = \lceil \frac{6-3\chi}{4} \rceil$. \square

The following result improves Corollary 2 in [5], which says that every 5-connected triangulation in the double torus with representativity at least 6 admits a 4-trail. Note that the Euler characteristic of the double torus is -2 .

Corollary 4.4. *Every 5-connected graph G which embeds on the double torus admits a 3-trail.*

References

- [1] R. Brunet, M. N. Ellingham, Z. Gao, A. Metzlar, and R. B. Richter, Spanning planar subgraphs of graphs in the torus and Klein bottle, *J. Combin. Theory Ser. B*, 65 (1995), pp. 7–22.
- [2] P.A. Catlin, Double cycle covers and the Petersen graph, *J. Graph Theory* 13 (1989), 465–483.
- [3] M. N. Ellingham, Spanning paths, cycles, trees and walks for graphs on surfaces, *Congr. Numer.*, 115 (1996), pp. 55–90.
- [4] Z. Gao and R. B. Richter, 2-walks in circuit graphs, *J. Combin. Theory Ser. B*, 62 (1994), pp. 259–267.

- [5] Z. Gao and N. C. Wormald, Spanning Eulerian subgraphs of bounded degree in triangulations, *Graphs Combin.*, 10 (1994), pp. 123–131.
- [6] B. Grünbaum, Polytopes, graphs, and complexes, *Bull. Amer. Math. Soc.*, 76 (1970), pp. 1131–1201.
- [7] R. Halin, A theorem on n -connected graphs, *J. Combinatorial Theory*, 7 (1969), pp. 150–154.
- [8] R. Halin, Untersuchungen über minimale n -fach zusammenhängende Graphen, *Math. Ann.*, 182 (1969), pp. 175–188.
- [9] M. Hasanvand, Spanning trees and spanning Eulerian subgraphs with small degrees. II, <https://arxiv.org/abs/1702.06203>
- [10] B. Jackson and N. C. Wormald, k -walks of graphs, *Australas. J. Combin.*, 2 (1990), pp. 135–146.
- [11] C. St. J. A. Nash-Williams, Edge-disjoint spanning trees of finite graphs, *J. London Math. Soc.*, 36 (1961), pp. 445–450.
- [12] C. St. J. A. Nash-Williams, Unexplored and semi-explored territories in graph theory, (1973), pp. 149–186.
- [13] K. Ota and K. Ozeki, Spanning trees in 3-connected $K_{3,t}$ -minor-free graphs, *J. Combin. Theory Ser. B*, 102 (2012), pp. 1179–1188.
- [14] K. Ozeki, Spanning trees with bounded maximum degrees of graphs on surfaces, *SIAM J. Discrete Math.* 27 (2013), pp. 422–435.
- [15] D. P. Sanders and Y. Zhao, On 2-connected spanning subgraphs with low maximum degree, *J. Combin. Theory Ser. B*, 74 (1998), pp. 64–86.
- [16] D. P. Sanders and Y. Zhao, On spanning trees and walks of low maximum degree, *J. Graph Theory*, 36 (2001), pp. 67–74.
- [17] R. Thomas and X. Yu, 4-connected projective-planar graphs are Hamiltonian, *J. Combin. Theory Ser. B*, 62 (1994), pp. 114–132.
- [18] W. T. Tutte, A theorem on planar graphs, *Trans. Amer. Math. Soc.*, 82 (1956), pp. 99–116.
- [19] W. T. Tutte, On the problem of decomposing a graph into n connected factors, *J. London Math. Soc.*, 36 (1961), pp. 221–230.