

ON A CONJECTURE OF KARASEV

SEUNGHUN LEE AND KANGMIN YOO

ABSTRACT. Karasev conjectured that for any set of $3k$ lines in general position in the plane, which is partitioned into 3 color classes of equal size k , the set can be partitioned into k colorful 3-subsets such that all the triangles formed by the subsets have a point in common. Although the general conjecture is false, we show that Karasev's conjecture is true for lines in convex position. We also discuss possible generalizations of this result.

1. INTRODUCTION

One of the classical results in discrete geometry is Tverberg's Theorem [10] which asserts that any set of $(d+1)(r-1)+1$ points in \mathbb{R}^d can be partitioned into r disjoint subsets whose convex hulls have a point in common. Tverberg's theorem has many variants as we discuss in this section.

1.1. Dual Tverberg theorems. We say that a collection of hyperplanes in \mathbb{R}^d is in *general position* if the intersection of any d of the hyperplanes is a single point, but the intersection of any $d+1$ of them is empty. Note that, for any $d+1$ hyperplanes in general position in \mathbb{R}^d , there is a unique bounded d -dimensional simplex formed by the hyperplanes. There are dual versions of Tverberg's theorem, which consider a set of hyperplanes in general position and bounded simplices formed by its subsets of size $d+1$. For a set of lines, Roudneff [8] proved the following result.

Theorem 1.1. *For a set of $3r$ lines in general position in the plane, the set can be partitioned into r subsets of size 3 so that all the triangles formed by the subsets have a point in common.*

And Karasev [7] proved a generalization of this result.

Theorem 1.2. *Let r be a prime power. For a set of $(d+1)r$ hyperplanes in general position in \mathbb{R}^d , the set can be partitioned into r subsets of size $d+1$ so that all the d -dimensional simplices formed by the subsets have a point in common.*

1.2. Colored Tverberg theorems. For a set P which is partitioned into color classes, a subset Q of P is called *colorful* if it contains exactly one from each color class. A research direction which has received a significant amount of attention in recent years is to establish a colored version of Tverberg's theorem.

Theorem 1.3 (The colored Tverberg theorem). *For any d and r with $d \geq 1$ and $r \geq 2$, there exists a positive integer t which satisfies the following: For a set of $(d+1)t$ points in \mathbb{R}^d which is partitioned into $d+1$ color classes of size t , there exist r disjoint colorful subsets whose convex hulls have a point in common.*

This research was supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education (NRF-2016R1D1A1B03930998).

This theorem was originally a conjecture of Bárány, Füredi and Lovász [1]. Let $t(d, r)$ be the smallest integer for which the conclusion of Theorem 1.3 holds. A particular case $t(2, 3) \leq 7$ was given in the same paper by Bárány, Füredi and Lovász [1]. Later, Bárány and Larman [2] proved $t(2, r) = r$ and conjectured $t(d, r) = r$ for every dimension d . Using topological methods, the general case was first proved by Živaljević and Vrećica [12], who also showed that $t(d, r) \leq 2r - 1$ whenever r is a prime number. The same bound was extended to all r which are prime powers [13]. Later Blagojević, Matschke and Ziegler [4] obtained the optimal bound $t(d, r) = r$ whenever $r + 1$ is a prime number.

1.3. The dual colored Tverberg theorem and our main result. Karasev [7] also proved a dual version of the colored Tverberg theorem.

Theorem 1.4 (The dual colored Tverberg theorem). *For any d and r with $d \geq 1$ and $r \geq 2$, there exists a positive integer k which satisfies the following: For a set of $(d+1)k$ hyperplanes in \mathbb{R}^d in general position, which is partitioned into $d + 1$ color classes of size k , there exist r disjoint colorful subsets such that all the d -dimensional simplices formed by the subsets have a point in common.*


Similarly with $t(d, r)$, let $k(d, r)$ be the smallest integer for which the conclusion of Theorem 1.4 holds. In [7], Karasev proved $k(d, r) \leq 2r - 1$ when r is a prime power. Especially for the 2-dimensional case, he conjectured that the equality $k(2, r) = r$ holds, which is an analogue of $t(2, r) = r$ for the colored Tverberg theorem. However, the following example shows that $k(2, r) \neq r$ whenever $r = 2t$ for some odd number t (The counterexample showing $k(2, 2) \neq 2$ was also discovered independently by Liping Yuan [11]).

Example 1.5. Consider the set \mathcal{L} with 6 lines shown in Figure 1. Note that the set \mathcal{L} is in general position and each line in \mathcal{L} is colored by either RED, BLUE, or GREEN. There are 4 ways to partition \mathcal{L} into 2 colorful subsets, and in every case the subsets form two disjoint triangles. Therefore, the set \mathcal{L} is a counterexample to the conjecture.

Now we make an arbitrarily large counterexample. For a fixed odd number t , we make t copies for each line in \mathcal{L} . The copied lines are perturbed a little from the original lines so that the copies form a new set $\mathcal{L}^{(t)}$ of lines in general position, and every selection of copies, one for each original line, gives a set of lines whose arrangement is isomorphic to that of \mathcal{L} . Also, each copy is painted by the same color with its original line.

To show that $\mathcal{L}^{(t)}$ is a counterexample, it is sufficient to show that in any partition of $\mathcal{L}^{(t)}$ into colorful subsets, there are two parts L_1 and L_2 such that every line l in \mathcal{L} has a copy in $L_1 \cup L_2$, which makes $L_1 \cup L_2$ isomorphic to \mathcal{L} as a line arrangement. Combinatorially, this is same as asking whether it is possible to choose $2t$ facets from the regular octahedron, possibly multiple times, so that there are no pairs of opposite facets among selected ones and each vertex is covered by the facets exactly t times. The only possible way to do this is first taking 4 facets so that any two of them intersect exactly at a vertex, and choosing each of them exactly $t/2$ times. And this is impossible when t is odd.

Even though the general conjecture is false, we can show that Karasev's conjecture is true when we add an additional condition on the arrangement of lines. We say that lines l_1, \dots, l_n in the plane \mathbb{R}^2 are *in convex position* if they are in general position and the complement $\mathbb{R}^2 \setminus (\bigcup_{i=1}^n l_i)$ has a connected component whose boundary meets every line. And here is our main result.

FIGURE 1. A counterexample to the conjecture $k(2, r) = r$.

Theorem 1.6. For any set of $3k$ lines in \mathbb{R}^2 in convex position which is partitioned into 3 color classes of size k , the set can be partitioned into k colorful subsets such that all the triangles formed by the subsets have a point in common.

This note is organized as follows. In Section 2, we reduce Theorem 1.6 to its dual version Theorem 2.1 below and make some observations for the proof of Theorem 2.1. In Section 3, we prove Theorem 2.1. Finally in Section 4, we discuss possible generalizations of Theorem 1.6.

2. PRELIMINARIES

In this section, we reduce Theorem 1.6 to its dual version Theorem 2.1 below and make some observations which are useful in the proof of Theorem 2.1. Before that, let us introduce some conventions we use in the remainder of this note.

By a k -set (or a k -subset), we mean a set (or a subset, respectively) of size k . For a set P which is partitioned into color classes, a subset Q of P is called *colorful* if it contains exactly one from each color class. And when we consider a partition of a set, we naturally identify it with the family of all parts from the partition.

2.1. Reduction to a dual version on a circle. In this subsection, we state a dual version of Theorem 1.6 regarding points on a circle, and show how it implies Theorem 1.6.




FIGURE 2. Two types of 3-sets on a circle.

A 3-set of points on a circle without antipodal pairs is said to be *bounding* if its convex hull contains the center of the circle, and *unbounding* otherwise. For any unbounding 3-sets, there is a unique point in the set which is contained in the shorter arc connecting the other two points, and we call this a *middle point* of the 3-set (In Figure 2-(B), the middle point is labelled by m).

Let P be a set of $3k$ points on a circle S partitioned into 3 color classes of equal size, and let \mathcal{P} be a partition of P into disjoint 3-subsets. The partition \mathcal{P} is said to be *colorful* if all parts of the partition \mathcal{P} are colorful. If a point p is a middle point of some unbounding 3-set in \mathcal{P} , then we say that the point p is a *middle point of the partition* \mathcal{P} . Finally, we say that the middle points of a partition \mathcal{P} are *consecutive* if

- there is a semicircle in S containing all middle points of \mathcal{P} , and
- the shortest closed arc in S containing all middle points has no other points from P .

For convenience, we also say that the middle points of \mathcal{P} are consecutive even in the case when there are no middle points of \mathcal{P} .

Now, we state the dual version of Theorem 1.6.

Theorem 2.1. *Let P be a set of $3k$ points on a circle without antipodal pairs, which is partitioned into 3 color classes of size k . Then there exists a partition of P into k disjoint colorful 3-sets whose middle points are consecutive.*

Before proving this theorem in the next section, we first show that Theorem 2.1 implies Theorem 1.6.

Proof of Theorem 1.6. Let \mathcal{L} be a given set of $3k$ lines in convex position. We choose a sufficiently small circle S so that it is contained in a connected component of the complement $\mathbb{R}^2 \setminus (\bigcup_{l \in \mathcal{L}} l)$ which meets every line in \mathcal{L} . For each line l in \mathcal{L} , let $p(l)$ be the intersection point of S with a ray from the center of S , which perpendicularly intersects l . And paint the point $p(l)$ with the color of l . If we let $P = \{p(l) : l \in \mathcal{L}\}$, then P has no antipodal pairs and is partitioned into 3 color classes.

Let \mathcal{P} be a partition of P given by Theorem 2.1. With respect to the partition \mathcal{P} , we make each line l in \mathcal{F} correspond to a halfspace $H(l)$ in the following way:

- If $p(l)$ is a middle point of \mathcal{P} , then let $H(l)$ be the closed halfplane bounded by l which does not contain the center of S .
- Otherwise, let $H(l)$ be the closed halfplane bounded by l which contains the center.

Then for each part $\{p(l_1), p(l_2), p(l_3)\}$ in the partition \mathcal{P} , the 3-set $\{l_1, l_2, l_3\}$ is colorful and the triangle formed by $\{l_1, l_2, l_3\}$ is equal to the intersection of halfplanes $H(l_1) \cap H(l_2) \cap H(l_3)$.

So, it is sufficient to show that the intersection $\bigcap_{l \in \mathcal{L}} H(l)$ is nonempty. If there is at most one middle point of \mathcal{P} , then this is obvious. So we assume that there are at least two middle points of \mathcal{P} . Let γ be the shortest closed arc in S containing all the middle points of \mathcal{P} , and let l_a and l_b be the members in \mathcal{L} such that $p(l_a)$ and $p(l_b)$ are the boundary points of γ . And let q be the intersection point of l_a and l_b . Since the arc γ is contained in a semicircle, for each $l \in \mathcal{L}$, the halfplane $H(l)$ contains the common point q . \square

2.2. A colorful partition with respect to a circular ordering. In this subsection, we consider some observations which guarantee the existence of a colorful partition with respect to a circular ordering. We first recall a concept, which was introduced and investigated by Bárány, Holmsen and Karasev [3].

Definition 2.2. The **geometric join** of m point sets C_1, \dots, C_m in the Euclidean space \mathbb{R}^d , denoted by $\mathcal{G}(C_1, \dots, C_m)$, is the set of all convex combinations $\sum_{j=1}^m t_j p_j \in \mathbb{R}^d$ where $p_j \in C_j$, $t_j \geq 0$ and $\sum_{j=1}^m t_j = 1$.

In particular, in the plane, it was shown that the geometric join of 3 point sets is starshaped [5, 9]. Using this, we can prove following proposition.

Proposition 2.3. *Let P be a set of points on a unit circle S centered at the origin O , partitioned into 3 nonempty color classes C_1, C_2 and C_3 . If all colorful 3-subsets of P are unbounding, then there is a point q on S such that the closed line segment connecting q and O is disjoint from $\mathcal{G}(C_1, C_2, C_3)$.*

Proof. Since $\mathcal{G}(C_1, C_2, C_3)$ is starshaped, we can find a point p in $\mathcal{G}(C_1, C_2, C_3)$ so that the point p can see every point in $\mathcal{G}(C_1, C_2, C_3)$. When t is a nonnegative scalar, $t \cdot (-p)$ is not contained in $\mathcal{G}(C_1, C_2, C_3)$. Otherwise the origin O must be contained in $\mathcal{G}(C_1, C_2, C_3)$, which implies there is a colorful bounding 3-set in P leading to a contradiction. Therefore, we can choose the point $-p/||p||$ as q on S . \square

The following proposition gives a combinatorial characterization to have a certain colorful partition.

Proposition 2.4. *Let P be a set of $3k$ elements, which is partitioned into 3 subsets C_1, C_2 and C_3 of size k . Suppose that there is another partition of P into 3 subsets A_1, A_2 and A_3 of size k . Then, we can find a partition of P into k 3-sets T_1, \dots, T_k such that $|T_i \cap A_j| = |T_i \cap C_j| = 1$ for all $i \in \{1, \dots, k\}$ and $j \in \{1, 2, 3\}$.*

Proof. We use induction on k . Since it is obvious when $k = 1$, suppose that $k > 1$. Define a bipartite graph G with $V(G) = \{A_1, A_2, A_3\} \cup \{C_1, C_2, C_3\}$ and $E(G) = \{A_i C_j : A_i \cap C_j \neq \emptyset\}$. By the Pigeonhole principle, union of any t of C_1, C_2, C_3 intersects with at least t of A_1, A_2, A_3 for each $t \in \{1, 2, 3\}$. Thus there exists a perfect matching M in G by Hall's theorem [6]. Choose $x_i \in A_i \cap C_j$ for each $A_i C_j \in M$, then let $T = \{x_1, x_2, x_3\}$. Then we have $|T \cap A_j| = |T \cap C_j| = 1$ for every $j \in \{1, 2, 3\}$. By adding T to the partition for $P \setminus T$ obtained by the induction hypothesis, we can get the desired partition. \square

Now, let P be a set of $3k$ points on a unit circle S centered at the origin O , partitioned into 3 color classes C_1, C_2 and C_3 of size k . We say that a set of n points on a circle has a *clockwise ordering* p_1, \dots, p_n from the point q if, as we go along the circle S clockwise from




FIGURE 3. An example for Observation 2.5 and 2.6 when $k = 2$. The geometric join of the color classes is described by its radial projection to the circle as a thick arc.

the point q , p_i is the i th point we meet during the tour. For two non-antipodal points p and q on a circle S , the arc moving from p to q clockwise is denoted by $\gamma_S(p, q)$.

Observation 2.5. *If there are no colorful bounding 3-subsets of P , then we can find a colorful partition of P with consecutive middle points.*

Proof. Since there are only unbounding 3-sets among colorful subsets of P , we can apply Proposition 2.3 to P in order to get the point q in the conclusion of Proposition 2.3. Let p_1, \dots, p_{3k} be the clockwise ordering from q . And let $A_j = \{p_i : (j-1)k+1 \leq i \leq jk\}$ for $j \in \{1, 2, 3\}$. By using Proposition 2.4 on two partitions $\{A_1, A_2, A_3\}$ and $\{C_1, C_2, C_3\}$, we can find a colorful partition \mathcal{P} into 3-subsets such that each part uses exactly one from each A_j . Note that each part must be unbounding by the assumption. And the assumption that the line segment connecting O and p does not have intersection with $\mathcal{G}(C_1, C_2, C_3)$ concludes that only points in A_2 can be used as a middle point of the partition. This readily implies that the middle points of \mathcal{P} , which are the points in A_2 , are consecutive. \square

We need one last observation. As in the proof of Observation 2.5, we assume that points in P has a clockwise ordering p_1, p_2, \dots, p_{3k} from some point, and let $A_j = \{p_i : (j-1)k+1 \leq i \leq jk\}$ for $j \in \{1, 2, 3\}$.

Observation 2.6. *Suppose that each of the arcs $\gamma_S(p_1, p_{2k})$ and $\gamma_S(p_{k+1}, p_{3k})$ are strictly contained in a semicircle (but not necessarily the same semicircle). Then there exists a colorful partition \mathcal{P} of the set P such that each part of \mathcal{P} contains exactly one from each A_j , and the set of middle points of \mathcal{P} is contained in A_2 .*

Proof. By using Proposition 2.4 on two partitions $\{A_1, A_2, A_3\}$ and $\{C_1, C_2, C_3\}$, we can find a colorful partition into 3-subsets such that each part uses exactly one from each A_j .

Suppose that one of the parts T is unbounding. Let us denote the unique point in $T \cap A_j$ by x_j . The point x_3 cannot be a middle point of T , since $\gamma_S(x_1, x_2)$ does not contain x_3 but it is contained in $\gamma_S(p_1, p_{2k})$ which implies that $\gamma_S(x_1, x_2)$ is the shorter arc connecting x_1 and x_2 . Similarly x_1 cannot be a middle point of T , so x_2 is a middle point of T . \square

3. PROOF OF THEOREM 2.1

In this section, we prove Theorem 2.1. Throughout this section, let S be the unit circle centered at the origin O , and P be a set of $3k$ points on S without antipodal pairs of points. Assume that P is partitioned into 3 color classes C_1, C_2 and C_3 of size k .

Recall that we need to find a colorful partition of P whose middle points are consecutive. If there exists a colorful partition of P which contains at most one colorful unbounding 3-set, then we are done. So we assume that any colorful partition of P contains at least two colorful unbounding 3-sets.

If the middle points of a colorful partition \mathcal{P} of P are contained in a semicircle, then the shortest closed arc containing all the middle points of \mathcal{P} is well-defined. We denote this arc by $\gamma(\mathcal{P})$. In particular, the boundary points of $\gamma(\mathcal{P})$ are middle points of two distinct unbounding 3-sets in \mathcal{P} . Then the rest of proof of Theorem 2.1 can be drawn by the following lemma.

Lemma 3.1. *Let \mathcal{P} be a colorful partition of P which satisfies the following conditions:*

- (i) \mathcal{P} contains the minimum number of colorful unbounding 3-sets,
- (ii) The middle points of \mathcal{P} are contained in a semicircle.
- (iii) If a point in P is contained in $\gamma(\mathcal{P})$, then it is either a middle point of \mathcal{P} or a point from a colorful bounding 3-set in \mathcal{P} .

If $\gamma(\mathcal{P})$ contains a point from a colorful bounding 3-set in \mathcal{P} , then there exists another colorful partition \mathcal{P}' of P satisfying (i), (ii) and (iii), such that $\gamma(\mathcal{P}')$ is strictly contained in $\gamma(\mathcal{P})$.

Proof of Theorem 2.1. First, we show that there exists at least one colorful partition of P satisfying (i), (ii) and (iii). Let \mathcal{P} be a colorful partition of P which satisfies (i). Let \mathcal{P}_U be the subfamily of \mathcal{P} which consists of all unbounding 3-sets in \mathcal{P} , and let X be the union of all 3-sets in \mathcal{P}_U . In particular, $\mathcal{P}_U \subseteq \mathcal{P}$ and $X \subseteq P$. Note that there is no colorful bounding 3-subset of X by (i). So we can use Observation 2.5 to obtain a colorful partition \mathcal{P}'_U of X with consecutive middle points. In particular, the middle points of \mathcal{P}'_U are contained in a semicircle. If we replace \mathcal{P}_U with \mathcal{P}'_U in \mathcal{P} , then we obtain a new colorful partition \mathcal{P}' of the whole set P , which satisfies (i), (ii) and (iii).

Now let \mathcal{F} be the collection of all colorful partitions of P satisfying (i), (ii) and (iii). Choose $\mathcal{P}_0 \in \mathcal{F}$ so that $|\gamma(\mathcal{P}_0) \cap P|$ is minimum possible among all partitions in \mathcal{F} , i.e. $|\gamma(\mathcal{P}_0) \cap P| = \min\{|\gamma(\mathcal{P}) \cap P| : \mathcal{P} \in \mathcal{F}\}$. We claim that the middle points of \mathcal{P}_0 are consecutive. Suppose otherwise. Then there is a point $p \in P$ contained in $\gamma(\mathcal{P}_0)$ which is not a middle point of \mathcal{P}_0 . Thus p is a point from a colorful bounding 3-set in \mathcal{P}_0 by (iii). By Lemma 3.1, there exists another partition $\mathcal{P}_1 \in \mathcal{F}$ with $\gamma(\mathcal{P}_1) \subsetneq \gamma(\mathcal{P}_0)$. Since $\gamma(\mathcal{P}_1)$ misses at least one boundary point of $\gamma(\mathcal{P}_0)$, which is a point in P , we have $|\gamma(\mathcal{P}_1) \cap P| < |\gamma(\mathcal{P}_0) \cap P|$. This contradicts the choice of \mathcal{P}_0 . \square

Let \mathcal{P} be a colorful partition of P which satisfies (i), (ii) and (iii), and $\gamma(\mathcal{P})$ contains a point from a colorful bounding 3-set in \mathcal{P} . In what follows, we choose certain three members in \mathcal{P} and repartition their union (9-point set, three points for each color) into another appropriate 3-sets, so that if we replace the old ones with the new ones, then we get a desired partition \mathcal{P}' in Lemma 3.1. More precisely, let B be a bounding 3-set in \mathcal{P} which has a point contained in $\gamma(\mathcal{P})$. Let U_1 and U_2 be the unbounding 3-sets in \mathcal{P} whose middle points are the boundary points of $\gamma(\mathcal{P})$. And let $Q = B \cup U_1 \cup U_2$. Under these assumptions, the following claim implies Lemma 3.1.

Claim. *There exists a new colorful partition of Q into one bounding and two unbounding 3-sets, whose set of all middle points is contained in $\gamma(\mathcal{P})$ and contains all points in $B \cap \gamma(\mathcal{P})$.*




FIGURE 4. A possible case when $|\gamma \cap B| = 2$. In (a), the thick triangle represents B' . In (b), two thick triangles represent the partition given by Observation 2.6. Here, b_1 and b_2 become the new middle points.




FIGURE 5. A possible case when $|\gamma \cap B| = 1$. In (a), the thick triangle represents B' . In (b), two thick triangles represent the partition given by Observation 2.6. Here, m_1 and b_1 become the new middle points.

Proof of Lemma 3.1. Let $\{B', U'_1, U'_2\}$ be a partition given by the claim, where B' is a bounding, and U'_1 and U'_2 are unbounding 3-sets. By replacing $\{B, U_1, U_2\}$ with $\{B', U'_1, U'_2\}$, we obtain from \mathcal{P} another colorful partition \mathcal{P}' of the whole set P . Clearly, \mathcal{P}' satisfies (i) and (ii). Since the two middle points of $\{B', U'_1, U'_2\}$ are contained in $\gamma(\mathcal{P})$, we have $\gamma(\mathcal{P}') \subseteq \gamma(\mathcal{P})$. Note that at least one of them is a point in $B \cap \gamma(\mathcal{P})$. So $\gamma(\mathcal{P}')$ misses at least one boundary point of $\gamma(\mathcal{P})$, which is the middle point of either U_1 or U_2 . Thus $\gamma(\mathcal{P}') \subsetneq \gamma(\mathcal{P})$.

Now we show \mathcal{P}' satisfies (iii). Choose an arbitrary point $p \in P \cap \gamma(\mathcal{P}')$. If p is in $(P \setminus Q) \cap \gamma(\mathcal{P}')$, then it is contained in a 3-set T with $T \in \mathcal{P} \cap \mathcal{P}'$. Since \mathcal{P} satisfies (iii) and

$p \in \gamma(\mathcal{P}') \subsetneq \gamma(\mathcal{P})$, T must be a bounding 3-set or an unbounding 3-set whose middle point is p .

Now suppose that p is in $Q \cap \gamma(\mathcal{P}')$. Note that $B \cap \gamma(\mathcal{P})$ consists of either one or two points in B since $\gamma(\mathcal{P})$ cannot contain B by (ii) of \mathcal{P} . Also, we have $Q \cap \gamma(\mathcal{P}) = (B \cap \gamma(\mathcal{P})) \cup \{m_1, m_2\}$ where m_1 and m_2 are two boundary points of $\gamma(\mathcal{P})$, by (iii) of \mathcal{P} . For example, see Figure 4a and 5a (in the figures, B , U_1 and U_2 are drawn by dotted triangles.)

If there are two points in $B \cap \gamma(\mathcal{P})$, then they are exactly the two middle points of $\{B', U'_1, U'_2\}$. So neither m_1 nor m_2 is contained in $\gamma(\mathcal{P}')$, that is, $Q \cap \gamma(\mathcal{P}') = B \cap \gamma(\mathcal{P})$. Thus p must be a middle point of either U'_1 or U'_2 .

If there is only one point in $B \cap \gamma(\mathcal{P})$, then one of the two middle points of $\{B', U'_1, U'_2\}$ is the point in $B \cap \gamma(\mathcal{P})$, and the other one is m_1 or m_2 , say m_1 . Since $\gamma(\mathcal{P}')$ does not contain the other point m_2 , the points in $Q \cap \gamma(\mathcal{P}')$ are exactly the two middle points of $\{B', U'_1, U'_2\}$. Therefore, \mathcal{P}' satisfies (iii). \square

Before we prove the claim, let us outline the proof. First, We choose a suitable colorful bounding 3-set in Q which becomes B' in the proof of Lemma 3.1. After we choose B' , we show that the remaining 6-point set $Q \setminus B'$ can be partitioned into two colorful unbounding 3-sets, which becomes U'_1 and U'_2 . Here we apply Observation 2.6. See Figure 4 and 5.

Let us note that the condition (i) of \mathcal{P} directly implies the following.

(*) *Any colorful partition of Q contains at most one colorful bounding 3-set.*

Also, as shown in the proof of Lemma 3.1, the condition (ii) of \mathcal{P} implies that $B \cap \gamma(\mathcal{P})$ consists of either one or two points. Recall that $\gamma_S(p, q)$ denotes the arc moving from p to q clockwise.

Now we start to prove the claim. Denote $\gamma = \gamma(\mathcal{P})$. Let $B = \{b_1, b_2, b_3\}$ with $b_1 \in \gamma$. Let $U_1 = \{l_1, m_1, r_1\}$ and $U_2 = \{l_2, m_2, r_2\}$, where m_i denotes the middle point of U_i for $i \in \{1, 2\}$. To fix the relative positions of the points in U_1 and U_2 , we assume that γ equals $\gamma_S(m_1, m_2)$ and that the shorter arc connecting l_i and r_i , which thus contains m_i , equals $\gamma_S(l_i, r_i)$ for $i \in \{1, 2\}$. Since U_1 and U_2 are unbounding 3-sets and \mathcal{P} satisfies (iii), it follows that $l_1, l_2 \in \gamma_S(-m_2, m_1)$ and $r_1, r_2 \in \gamma_S(m_2, -m_1)$. The situation is described in Figure 4 and 5.

Now we consider the two cases according to the number of points in $B \cap \gamma$. For simplicity, we assume that $b_i \in C_i$ for $i \in \{1, 2, 3\}$.

First, consider the case when there are two points in $B \cap \gamma$. In this case, we assume $b_2 \in \gamma$ and $b_3 \notin \gamma$. Since b_1 and b_2 are in $\gamma_S(m_1, m_2)$, b_3 must be in $\gamma_S(-m_1, -m_2)$. See Figure 4. Suppose that there exists a colorful bounding 3-set B' in Q which consists of b_3 and exactly one point in each of $\{m_1, l_1, l_2\}$ and $\{m_2, r_1, r_2\}$. If we denote the clockwise ordering of $Q \setminus B'$ from b_3 by p_1, \dots, p_6 , then $\{p_3, p_4\}$ equals $\{b_1, b_2\}$. Note that this ordering satisfies the condition in Observation 2.6. By (*), $Q \setminus B'$ contains no colorful bounding 3-set. So Observation 2.6 gives a partition of $Q \setminus B'$ into two colorful unbounding 3-sets U'_1 and U'_2 , whose middle points are exactly b_1 and b_2 . Therefore, we have a desired partition $\{B', U'_1, U'_2\}$ of Q .

Now we show that such B' always exists. Note that b_3 forms bounding 3-sets with the pairs $\{m_1, r_1\}$, $\{l_1, r_1\}$, $\{l_2, m_2\}$, $\{l_2, r_2\}$, $\{m_1, m_2\}$ and $\{m_1, r_2\}$. If any one of them is colorful, then we are done. So suppose otherwise. The first four pairs imply that $r_1, l_2 \in C_3$, so that $m_1, m_2, r_2 \notin C_3$. But then either $\{b_3, m_1, m_2\}$ or $\{b_3, m_1, r_2\}$ is colorful, which is a contradiction. This concludes the first case.

Second, consider the case when there is only one point in $B \cap \gamma$. In this case, we assume $b_2 \in \gamma_S(m_2, -b_1)$ and $b_3 \in \gamma_S(-b_1, m_1)$. See Figure 5. As before, we need to find a suitable colorful bounding 3-set in Q . Precisely, we need one of either of the following two types.

- (1) A colorful bounding 3-set which consists of exactly one point from $\{l_1, l_2, b_3\}$ and two points in $\{m_2, r_1, r_2, b_2\}$. It should contain b_2 if b_2 lies in $\gamma_S(-m_1, -b_1)$.
- (2) A colorful bounding 3-set which consists of exactly one point from $\{r_1, r_2, b_2\}$ and two points in $\{m_1, l_1, l_2, b_3\}$. It should contain b_3 if b_3 lies in $\gamma_S(-b_1, -m_2)$.

Suppose that there exists a colorful bounding 3-set B' in Q which is of either type (1) or type (2). Denote the clockwise ordering of $Q \setminus B'$ from $-b_1$ by p_1, \dots, p_6 . If B' is of type (1), then $\{p_3, p_4\}$ equals $\{m_1, b_1\}$. If B' is of type (2), then $\{p_3, p_4\}$ equals $\{b_1, m_2\}$. In each case, the ordering satisfies the condition of Observation 2.6. Combined with (*), Observation 2.6 gives a partition of $Q \setminus B'$ into two colorful unbounding 3-sets U'_1 and U'_2 , whose middle points are exactly p_3 and p_4 . Therefore, we have a desired partition $\{B', U'_1, U'_2\}$.

Finally, we show that such B' of either type (1) or (2) always exists. By a way of contradiction, suppose that there is no B' of any type. We divide cases according to the position of b_2 and b_3 .

- Suppose that neither b_2 nor b_3 is in $\gamma_S(-m_1, -m_2)$. Note that $\{b_2, b_3, m_2\}$ is a bounding 3-set. If it is colorful, then it is of type (1), which is forbidden. This implies that m_2 should not be in C_1 , so we have either $r_2 \in C_1$ or $l_2 \in C_1$.

If $r_2 \in C_1$, then $\{b_2, b_3, r_2\}$ is colorful. If $\gamma_S(b_3, r_2)$ is contained in a semicircle, then $\{b_2, b_3, r_2\}$ is a bounding 3-set, so that it is of type (1). Thus $\gamma_S(r_2, b_3)$ must be contained in a semicircle. Since $-m_2$ and $-l_2$ are contained in $\gamma_S(r_2, b_3)$, $\{b_3, m_2, r_2\}$ and $\{b_3, l_2, r_2\}$ are bounding 3-sets. Note that either one of them must be colorful. If $\{b_3, m_2, r_2\}$ is colorful, then it is of type (1). If $\{b_3, l_2, r_2\}$ is colorful, then it is of type (2). This leads to a contradiction. The case when $l_2 \in C_1$ can be treated in the same way.

- Suppose that at least one of b_2 and b_3 is in $\gamma_S(-m_1, -m_2)$. Without loss of generality, we assume that b_2 is in $\gamma_S(-m_1, -m_2)$. More precisely, b_2 is in $\gamma_S(-m_1, -b_1)$. Note that $\{b_2, l_2, m_2\}$ and $\{b_2, l_2, r_2\}$ are bounding 3-sets. If any one of them is colorful, then it is of type (1), which is forbidden. So both of them must not be colorful. This implies $l_2 \in C_2$.

Now suppose that b_3 is also in $\gamma_S(-m_1, -m_2)$, so it is in $\gamma_S(-b_1, -m_2)$. By a symmetric argument to the previous one, we have $r_1 \in C_3$. Note that $\{b_2, l_1, r_1\}$ is a bounding 3-set. If it is colorful, then it is of type (1). So l_1 must be in C_2 or C_3 . Since l_1 and r_1 are in different colors, we have $l_1 \in C_2$. Similarly, if $\{b_3, l_2, r_2\}$, being a bounding 3-set, is colorful, then it is of type (2). So r_2 must be in C_2 or C_3 . Since $l_2 \in C_2$, we have $r_2 \in C_3$. Consequently, m_1 and m_2 are in C_1 . But then we have two disjoint colorful bounding 3-sets $\{b_2, m_1, r_1\}$ and $\{b_3, l_2, m_2\}$, which contradicts (*).

Now suppose that b_3 is not in $\gamma_S(-b_1, -m_2)$. Then b_3 is in $\gamma_S(-m_2, m_1)$. Note that $\{b_2, b_3, m_2\}$ is a bounding 3-set. If it is colorful, then it is of type (1). So m_2 must be in C_2 or C_3 . Since $l_2 \in C_2$, we have $m_2 \in C_3$, so $r_2 \in C_1$. With this coloring, both $\{b_2, b_3, r_2\}$ and $\{l_2, r_2, b_3\}$ are colorful. Note that those two 3-sets have common points r_2 and b_3 . If $-r_2$ is in $\gamma_S(-m_2, b_3)$, then $\{b_2, b_3, r_2\}$ is a bounding 3-set, so it is of type (1). So $-r_2$ must be in $\gamma_S(b_3, m_1)$. But then $\{b_3, l_2, r_2\}$ is a bounding 3-set, so it is of type (2). This contradiction concludes the proof.

4. FINAL REMARKS

In this section, we discuss whether it is possible to generalize Theorem 1.6 in two different aspects. In Section 4.1, we show that there is a set of colored planes in convex position in \mathbb{R}^3 such that in every colorful partition of the set, simplices determined by parts of the partition have an empty intersection. Hence a direct generalization of Theorem 1.6 for hyperplanes in convex position in \mathbb{R}^d seems impossible. And in Section 4.2, we see an example of lines in the plane showing that being in convex position is not a necessary condition for the conclusion of Theorem 1.6.

4.1. An example in 3-dimension. Similar with lines in the plane, we say that hyperplanes h_1, \dots, h_n in general position in \mathbb{R}^d are *in convex position* if there is a connected component of the complement $\mathbb{R}^d \setminus (\bigcup_{i=1}^n h_i)$ whose boundary meets every hyperplane. Especially, planes in general position in \mathbb{R}^3 and tangent to an open hemisphere are in convex position, since the connected component containing the origin has a boundary which meets every plane at its tangent point.

We choose the following 8 points

$$\begin{aligned} p_1 &= (1/3, 2/3, 2/3) & p_2 &= (7/9, -4/9, 4/9) & p_3 &= (6/7, 2/7, 3/7) \\ p_4 &= (17/19, 6/19, 6/19) & p_5 &= (1/3, -2/3, 2/3) & p_6 &= (2/3, -2/3, 1/3) \\ p_7 &= (6/7, 3/7, 2/7) & p_8 &= (-2/7, 3/7, 6/7) \end{aligned}$$

from the open hemisphere $S = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 + z^2 = 1 \text{ and } z > 0\}$. For $1 \leq i \leq 8$, let h_i be a plane tangent to S at p_i , and define $\mathcal{H} = \{h_i : 1 \leq i \leq 8\}$. It is straightforward but tedious to check that \mathcal{H} is in general position. Make a partition of \mathcal{H} into color classes

$$C_1 = \{h_1, h_5\}, C_2 = \{h_2, h_6\}, C_3 = \{h_3, h_7\} \text{ and } C_4 = \{h_4, h_8\}.$$

We claim that for every colorful partition of \mathcal{H} , the simplices determined by parts of the partition have an empty intersection. Let h_i^+ (or h_i^-) be the closed halfspace bounded by h_i and not containing (or, containing, respectively) the origin. For every quadruple (i_1, i_2, i_3, i_4) of distinct elements from $\{1, \dots, 8\}$, and every sign vector w in $\{+, -\}^8$, let $w|_{(i_1, i_2, i_3, i_4)}$ be the closed region $\bigcap_{j=1}^4 h_{i_j}^{w(i_j)}$ where $w(i)$ is the i th coordinate of w . Let $v(i_1, i_2, i_3, i_4) \in \{+, -\}^8$ be the unique sign vector such that $v(i_1, i_2, i_3, i_4)|_{(i_1, i_2, i_3, i_4)}$ is the bounded simplex determined by planes $h_{i_1}, h_{i_2}, h_{i_3}$ and h_{i_4} , and $v(i) = 0$ for every $i \notin \{i_1, \dots, i_4\}$. Note that $-v(i_1, i_2, i_3, i_4)|_{(i_1, i_2, i_3, i_4)}$ is an emptyset.

Let us consider a partition $\{h_{i_1}, h_{i_2}, h_{i_3}, h_{i_4}\} \cup \{h_{i_5}, h_{i_6}, h_{i_7}, h_{i_8}\}$ of \mathcal{H} . If there is a quadruple (j_1, j_2, j_3, j_4) of distinct elements such that the vector $v(i_1, i_2, i_3, i_4) + v(i_5, i_6, i_7, i_8)$ in positions j_1, j_2, j_3, j_4 has the same entries as $-v(j_1, j_2, j_3, j_4)$ for a certain choice of positions j_1, j_2, j_3, j_4 , then we can see that the intersection of two simplices determined by the subsets must be empty, since the intersection is contained in $-v(j_1, j_2, j_3, j_4)|_{(j_1, j_2, j_3, j_4)}$ which is empty. Therefore, it is sufficient to find such a quadruple for every colorful partition. Which quadruple (j_1, j_2, j_3, j_4) can be chosen for each colorful partition is summarized at Table 1, with corresponding sign vectors.

4.2. Being in convex position is not necessary. In this subsection, we show that being in convex position is not necessary for the conclusion of Theorem 1.6. More precisely, we give an example of 6 lines which is not in convex position such that for every coloring using 3

$(i_1, i_2, i_3, i_4, i_5, i_6, i_7, i_8)$	$v(i_1, i_2, i_3, i_4) + v(i_5, i_6, i_7, i_8)$	(j_1, j_2, j_3, j_4)
(1, 2, 3, 4, 5, 6, 7, 8)	(-, -, +, -, +, -, +, -)	(1, 2, 5, 7)
(1, 2, 3, 8, 5, 6, 7, 4)	(+, +, -, +, -, -, -, -)	(1, 2, 3, 4)
(1, 2, 7, 4, 5, 6, 3, 8)	(+, -, +, +, +, -, -, -)	(2, 4, 5, 8)
(1, 2, 7, 8, 5, 6, 3, 4)	(+, +, +, -, -, +, -, -)	(1, 4, 5, 6)
(1, 6, 3, 4, 5, 2, 7, 8)	(-, +, +, -, -, -, -, +)	(1, 2, 5, 8)
(1, 6, 3, 8, 5, 2, 7, 4)	(-, -, +, +, +, -, -, +)	(1, 2, 3, 8)
(1, 6, 7, 4, 5, 2, 3, 8)	(-, -, +, +, +, -, -, -)	(1, 2, 4, 5)
(1, 6, 7, 8, 5, 2, 3, 4)	(+, +, +, -, -, +, -, -)	(1, 4, 5, 6)

TABLE 1. For all colorful partitions, the sign vector $-v(j_1, j_2, j_3, j_4)$ guarantees that simplices formed by the parts have empty intersection.




FIGURE 6. An example not in convex position, which has a colorful partition with intersecting triangles for every possible coloring.

color classes of equal size, we can always find a partition into colorful parts which determine intersecting triangles. This suggests finding a broader class of line arrangements where the conclusion of Theorem 1.6 holds.

See Figure 6. There are 6 lines l_1, \dots, l_6 not in convex position, and the 5 lines l_1, \dots, l_5 bounds the shaded region. We color each line with colors RED, BLUE and GREEN so that each color class has size 2. Without loss of generality, we assume that l_6 is painted by BLUE.

First, we suppose that l_2 and l_6 are in different colors. We assume that l_2 is colored by RED. If l_3 is colored by GREEN, then we can choose a partition $\{l_2, l_3, l_6\} \cup \{l_1, l_4, l_5\}$. If l_4 is colored by GREEN, then we can choose a partition $\{l_2, l_4, l_6\} \cup \{l_1, l_3, l_5\}$. So, we only need to consider the case when both l_1 and l_5 are colored by GREEN. But then, we can take a partition $\{l_2, l_5, l_6\} \cup \{l_1, l_3, l_4\}$.

Next, suppose l_2 and l_6 are in the same color BLUE. If l_1 and l_5 have different colors, then we can take a partition $\{l_1, l_2, l_5\} \cup \{l_3, l_4, l_6\}$. In this partition, we have a common intersection between triangles formed by 3-sets $\{l_1, l_2, l_5\}$ and $\{l_3, l_4, l_6\}$ in the region bounded by lines l_3, l_4 and l_5 . So we can assume that l_1 and l_5 have the same color. Then, we take a partition into $\{l_1, l_3, l_6\} \cup \{l_2, l_4, l_5\}$. In this case again, we have a common intersection in the region bounded by lines l_3, l_4 and l_5 .

ACKNOWLEDGEMENTS

We are grateful to Andreas Holmsen, Roman Karasev, Liping Yuan, Pablo Soberón, Minki Kim and Hong Chang Ji for their excellent comments.

REFERENCES

- [1] I. Bárány, Z. Füredi and L. Lovász: On the number of halving planes, *Combinatorica* **10(2)** (1990), 175–183.
- [2] I. Bárány and D.G. Larman: A colored version of Tverberg’s theorem, *J. London Math. Soc.* **s2-45(2)** (1992), 314–320.
- [3] I. Bárány, A. Holmsen and R. Karasev: Topology of geometric joins, *Discrete Comput. Geom.* **53** (2015), 402–413.
- [4] P.V.M. Blagojević, B. Matschke and G.M. Ziegler: Optimal bounds for the colored Tverberg problem, *J. Eur. Math. Soc.* **17** (2015), 739–754.
- [5] J.-D. Boissonnat, O. Devillers and F.P. Preparata: Computing the union of 3-colored triangles, *Intern. J. Comput. Geom. Appl.* **1** (1991), 187–196.
- [6] P. Hall: On Representatives of Subsets, *J. London Math. Soc.* **s1-10(1)** (1935), 26–30.
- [7] R.N. Karasev: Dual central point theorems and their generalizations, *Sbornik: Mathematics* **199(10)** (2008), 1459–1479.
- [8] J.-P. Roudneff: Tverberg-type theorems for pseudoconfigurations of points in the plane, *European J. Combin.* **9(2)** (1988), 189–198.
- [9] A. Schulz and C. Tóth: The union of colorful simplices spanned by a colored point set, *Comput. Geom. Theory Appl.* **46** (2011), 574–590.
- [10] H. Tverberg: A generalization of Radon’s theorem, *J. London Math. Soc.*, **s1-41(1)** (1966), 123–128.
- [11] Liping Yuan. Personal communication.
- [12] R.T. Živaljević and S.T. Vrećica: The colored Tverberg’s problem and complexes of injective functions, *J. Comb. Theory, Ser. A* **61(2)** (1992), 309–318.
- [13] R.T. Živaljević: User’s guide to equivariant methods in combinatorics, *II. Publ. Inst. Math.* **64(78)** (1998), 107–132.

S. LEE, DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, DAEJEON, SOUTH KOREA
E-mail address: prosolver@kaist.ac.kr

K. YOO, DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, DAEJEON, SOUTH KOREA
E-mail address: dearmymind@kaist.ac.kr