

Regular Bipartite Lattices with Large Values of $\theta_{2,2,2}/C_4$

Paul Federbush
 Department of Mathematics
 University of Michigan
 Ann Arbor, MI, 48109-1043

December 26, 2021

Abstract

The quantities C_4 and $\theta_{2,2,2}$ are as defined by Wanless, C_4 just the number of 4-loops of a graph. The construction of this paper provides a counterexample to a conjecture of Butera, Pernici, and the author about the monomer-dimer entropy, λ , of a regular bipartite lattice. The lattice we construct is not a lattice graph in its most common definition.

We let G_1 and G_2 be the graphs illustrated in Figure 1. Then given a graph G , we let $C_4 = C_4(G)$ be the number of subgraphs of G isomorphic to G_1 , and $\theta_{2,2,2}(G)$ be the number of subgraphs of G isomorphic to G_2 . This notation follows Section 2 of [1].

We propose the following definition for a lattice in the statistical mechanics setting: A connected infinite graph is a lattice if there is a finitely generated Abelian group of isomorphic maps on G for which the set of equivalence classes of vertices is finite. Two vertices are in the same equivalence class if there is a translation (a member of the Abelian group) that carries one into the other. The lattice is i -dimensional if exactly i of a set of generators of the group are infinite order.

The lattice (infinite graph) we construct in this paper has the following properties:

- 1) It is embedded in R^3 . Edges embed as straight line segments joining vertices.
- 2) It is invariant under translations by an integer in the three coordinate directions.
- 3) We consider a tessellation of R^3 by unit cubes whose vertices have integer coordinates. Each vertex of the graph is in the interior of one of these cubes. Each cube contains a finite number of vertices.
- 4) Each edge either lies in the interior of some cube, or connects two vertices in nearest neighbor cubes.




Figure 1:

This lattice does not have the property of arising from a tessellation, and thus is not a lattice graph as most commonly understood, though I believe it fits the idea of being a lattice for many researchers.

For a lattice the quantities C_4 and $\theta_{2,2,2}$ may both be infinite, but the ratio $\theta_{2,2,2}/C_4$ makes sense. Taking a sequence of finite graphs that approach the lattice in the sense of Benjamini-Schramm [2] the ratios $\theta_{2,2,2}/C_4$ converge to a limit we take to be the value for the lattice.

Given any number κ we find a three dimensional regular bipartite lattice for which

$$\theta_{2,2,2}/C_4 > \kappa \tag{1}$$

$$C_6 = 0. \tag{2}$$

Here C_6 is the number of 6-loops as C_4 is the number of 4-loops.

In [3] Friedland and I introduced an expansion for the monomer-dimer entropy of a regular lattice,

$$\lambda(p) = \lambda_B(p) + \sum_k d_k p^k \tag{3}$$

where λ_B is the monomer-dimer entropy for the Bethe lattice. In [4] Butera, Pernici and I conjectured that for a regular bipartite lattice all the $d_k \geq 0$. Actually a slightly weaker conjecture was stated in [4], see [5] Section 3. In [6] Pernici proved in the bipartite case d_2, d_3, d_4, d_5 were all nonnegative using a formalism of Wanless, [1]. For d_6 he finds the expression

$$d_6 = \frac{5\bar{C}_4}{d^6} + \frac{\bar{C}_6}{2d^6} - 2\frac{\bar{\theta}_{2,2,2}}{d^6} \tag{4}$$

where we have d-regularity and the bars indicate an average over the number of vertices, again as may be taken as a Benjamini-Schramm limit. We find a three dimensional regular bipartite lattice for each κ such that

$$\theta_{2,2,2}/C_4 > \kappa$$

and also

$$C_6 = 0.$$

Thus there are bipartite regular lattices for which $d_6 < 0$. Pernici erred in stating otherwise.

However the conjecture that all the $d_i \geq 0$ should not be lightly dismissed. For hyper-cubic lattices of all dimensions it is shown in [4] that the first 20 such coefficients satisfy this condition! It seems quite possible there is a class of 'nice' lattices that satisfy the conjecture, possibly vertex-transitive lattices.

We start the construction of our lattice with a 'root unit graph', a finite graph. To this we apply a number of 2-lifts to end with a 'full unit graph'. The full unit graphs are attached to yield the three dimensional lattice. We use 2-lifts similarly to as in Section 4 of [7], derived from [8]. The construction requires that $d \geq 5$.

The Root Unit Graph

This graph has black vertices:

$$c_i, \quad i = 1, \dots, d$$

and white vertices:

$$t, b, lx, ly, lz, rx, ry, rz, \\ f_i, \quad i = 1, \dots, d-5$$

The edges are:

$$\begin{aligned}
(t, c_i) & \quad i = 1, \dots, d \\
(b, c_i) & \quad i = 1, \dots, d \\
(c_i, f_j) & \quad i = 1, \dots, d \quad j = 1, \dots, d - 5 \\
& \quad (lx, c_1), (ly, c_1), (lz, c_1) \\
(rx, c_i) & \quad i = 2, \dots, d \\
(ry, c_i) & \quad i = 2, \dots, d \\
(rz, c_i) & \quad i = 2, \dots, d
\end{aligned}$$

It has a subgraph, the ‘root central subgraph’. The root central subgraph has black vertices:

$$c_i, \quad i = 1, \dots, d$$

and white vertices:

$$t, b$$

Its edges are:

$$\begin{aligned}
(t, c_i) & \quad i = 1, \dots, d \\
(b, c_i) & \quad i = 1, \dots, d
\end{aligned}$$

The quantities $\theta_{2,2,2}$ and C_4 for the root central subgraph are easily computed to be

$$C_4 = \frac{d(d-1)}{2} \tag{5}$$

$$\theta_{2,2,2} = \frac{d(d-1)(d-2)}{6} \tag{6}$$

Exploitation of the root central subgraph is the key idea in this paper.

Thank Heaven for 2-lifts

The full unit graph is constructed from the root unit graph through a sequence of s 2-lifts. Each edge in the full unit graph projects in a natural sense onto an edge of the root unit graph. 2^s edges of the full unit graph project onto each edge of the root unit graph (likewise 2^s vertices project onto each vertex). The edges that project onto edges of the root central subgraph have all been chosen parallel, none crossed (this is also in a natural sense). Thus there are 2^s disjoint subgraphs of the full unit graph isomorphic to the root central subgraph. After s 2-lifts any vertex may be assigned a binary integer of length s , that we call its ‘level’. If a zero appears in the i^{th} spot then at the i^{th} 2-lift the lower vertex was chosen, if a one appears the upper vertex chosen. (One arrives at the vertex by following a sequence of s splittings.)

The sequence of 2-lifts are chosen so that for the full unit graph there are no 6-loops, and 4-loops only in the 2^s subgraphs isomorphic to the root central subgraph.

Connecting the Graph

We associate a full unit graph to each of the aforementioned unit cubes that tessellate R^3 . Corresponding to each nearest neighbor pair of cubes we create a connection. Let cubes a and b

be nearest neighbors, and suppose cube b is obtained from cube a by the translation $x \rightarrow x + 1$. Then we connect the full unit graph associated to a to the full unit graph associated to b by identifying each of the 2^s vertices that project onto vertex rx of the graph a with one of the 2^s vertices that project onto lx of the graph b in a level preserving way. Nearest neighbor cubes lined up in the other two directions are treated analogously. The lattice or graph thus constructed will have $C_6 = 0$ and $\frac{\theta_{2,2,2}}{C_4} = \frac{g-2}{3}$.

Locating the Graph

It remains to associate to each vertex of the lattice a point in R^3 . To do this it is clearly sufficient to locate each vertex of a single full unit graph included in the construction. We let G be the full unit graph associated to the unit cube $[0, 1] \times [0, 1] \times [0, 1]$, and g its root unit graph. We divide the vertices of g into five disjoint sets S_1, S_2, S_3, S_4 and S_5 with $S_1 = \{rx\}$, $S_2 = \{ry\}$, $S_3 = \{rz\}$ and $S_4 = \{lx, ly, lz\}$. We let \tilde{S}_i be the vertices in G that project to vertices in S_i . A 'try' is an assignment of the vertices in \tilde{S}_5 into a set of disjoint points in $(1/3, 2/3) \times (1/3, 2/3) \times (1/3, 2/3)$, an assignment of \tilde{S}_1 into a set of disjoint points in $(2/3, 1) \times (1/3, 2/3) \times (1/3, 2/3)$, an assignment of \tilde{S}_2 into a set of disjoint points in $(1/3, 2/3) \times (2/3, 1) \times (1/3, 2/3)$, and an assignment of \tilde{S}_3 into a set of disjoint points in $(1/3, 2/3) \times (1/3, 2/3) \times (2/3, 1)$. Note that the assignments for \tilde{S}_1, \tilde{S}_2 , and \tilde{S}_3 determine an assignment for \tilde{S}_4 by the way connections were determined. For example, if a vertex in \tilde{S}_1 is assigned a point \mathbf{w} in R^3 , then the vertex of the same level that projects onto lx is assigned the point $\mathbf{w} - (1, 0, 0)$. A try is a "good try" if the edges of our full unit graph map into line segments that may intersect only in a vertex. But almost all tries are good tries, we pick a good try to fix the assignment. The resulting embedding of the lattice satisfies all our conditions.

Acknowledgement We would like to thank Sasha Barvinok for a lecture on 2-lifts and John Stembridge for putting up with my continual questions.

References

- [1] Wanless, I. M., *Counting matchings and tree-like walks in regular graphs*, Combinatorics, Probability and Computing **19** (2010) 463-480.
- [2] Csikvari, P., Abert, M., Hubai, T., *Matching measure, Benjamini-Schramm convergence and the monomer-dimer free energy*, Journal of Statistical Physics **161** (2015), 16-34.
- [3] Federbush, P., Friedland, S., *An asymptotic expansion and recursive inequalities for the monomer-dimer problem*, J. Stat. Phys. **143** (2011) 306-325.
- [4] Butera, P., Federbush, P., Pernici, M., *Higher-order expansions for the entropy of a dimer or a monomer-dimer system on d-dimensional lattices*, Phys. Rev. E **87**, 062113 (2013).
- [5] Csikvari, P. *Matchings in vertex-transitive bipartite graphs*, accepted to Israel Journal of Mathematics.
- [6] Pernici, M., *1/n expansion for the number of matchings on regular graphs and monomer-dimer entropy*, J. Stat. Phys. **168** (2017) 666.
- [7] Csikvari, P., *Lower matching conjecture, and a new proof of Schrijver's and Gurvits's theorems*, accepted to the Journal of the European Mathematical Society.
- [8] Linial, N., *Lifts of graphs*, (talk slides on the web).