

Distinguishing critical graphs

Saeid Alikhani * Samaneh Soltani

March 14, 2022

Department of Mathematics, Yazd University, 89195-741, Yazd, Iran
alikhani@yazd.ac.ir, s.soltani1979@gmail.com

Abstract

The distinguishing number $D(G)$ of a graph G is the least integer d such that G has a vertex labeling with d labels that is preserved only by a trivial automorphism. We say that a graph G is d -distinguishing critical, if $D(G) = d$ and $D(H) \neq D(G)$, for every proper induced subgraph H of G . This generalizes the usual definition of a d -chromatic critical graph. While the investigation of d -critical graphs is a well established part of coloring theory, not much is known about d -distinguishing critical graphs. In this paper we determine all d -distinguishing critical graphs for $d = 1, 2, 3$ and observe that all of these kind of graphs are k -regular graph for some $k \leq d$. Also, we show that the disconnected d -distinguishing critical graph with c connected components such that $c \geq \frac{d}{2}$, is a regular graph.

Keywords: distinguishing, critical, graph.

AMS Subj. Class.: 05C15, 05E18

1 Introduction

Let $G = (V, E)$ be a simple graph. We use the standard graph notation. In particular, $\text{Aut}(G)$ denotes the automorphism group of G . For simple connected graph G , and $v \in V$, the *neighborhood* of a vertex v is the set $N_G(v) = \{u \in V(G) : uv \in E(G)\}$. The *degree* of a vertex v in a graph G , denoted by $\deg_G(v)$, is the number of edges of G incident with v . In particular, $\deg_G(v)$ is the number of neighbours of v in G . We denote by $\delta(G)$ and $\Delta(G)$ the minimum and maximum degrees of the vertices of G . A graph G is k -regular if $\deg_G(v) = k$ for all $v \in V$. For $u, v \in V$ the *distance* of u and v , denoted by $d_G(u, v)$ or $d(u, v)$, is the length of a shortest path between u and v . The *eccentricity* of a vertex v , $\text{ecc}(v)$, is $\max\{d(u, v) : u \in V\}$. The maximum and minimum eccentricity of vertices of G are called *diameter* and *radius* of G and are denoted by $\text{diam}(G)$ and $\text{rad}(G)$, respectively. *Center* of G is the subgraph induced by vertices with eccentricity $\text{rad}(G)$. A graph is called *self-centered* if it is equal to its center,

*Corresponding author

or equivalently, its diameter equals its radius. A graph G is called *k-self-centered* if $\text{diam}(G) = \text{rad}(G) = k$. The terminology *k-equi-eccentric* graph is also used by some authors. For studies on these graphs see [4]. A *clique* of a graph is a set of mutually adjacent vertices, and that the maximum size of a clique of G , the clique number of G , is denoted $\omega(G)$. The complementary notion of a clique is an independent set, a set of vertices no two of which are adjacent. An independent set in a graph is maximum if the graph contains no larger stable set. The cardinality of a maximum independent set in G is called the *independence number* of G , denoted by $\alpha(G)$. Clearly, a subset S of V is an independent set in G if and only if S is a clique in \overline{G} , the complement of G . In particular, $\omega(\overline{G}) = \alpha(G)$.

A labeling of G , $\phi: V \rightarrow \{1, 2, \dots, r\}$, is said to be *r-distinguishing*, if no non-trivial automorphism of G preserves all of the vertex labels. The point of the labels on the vertices is to destroy the symmetries of the graph, that is, to make the automorphism group of the labeled graph trivial. Formally, ϕ is *r-distinguishing* if for every non-trivial $\sigma \in \text{Aut}(G)$, there exists x in $V = V(G)$ such that $\phi(x) \neq \phi(\sigma(x))$. We will often refer to a labeling as a coloring, but there is no assumption that adjacent vertices get different colors. Of course the goal is to minimize the number of colors used. Consequently the *distinguishing number* of G is defined by

$$D(G) = \min\{r \mid G \text{ has a labeling that is } r\text{-distinguishing}\}.$$

This number has introduced in [1]. If a graph has no nontrivial automorphisms, its distinguishing number is 1. In other words, $D(G) = 1$ for the asymmetric graphs. The other extreme, $D(G) = |V(G)|$, occurs if and only if $G = K_n$. The distinguishing number of some examples of graphs was obtained in [1]. For instance, $D(P_n) = 2$ for every $n \geq 3$, and $D(C_n) = 3$ for $n = 3, 4, 5$, $D(C_n) = 2$ for $n \geq 6$. Also, $D(K_{n,m}) = n$ where $n > m \geq 1$, and $D(K_{n,n}) = n + 1$ for any $n \geq 3$. Albertson and Collins in [1] showed that $D(G) = D(\overline{G})$, where \overline{G} is the complement graph of G .

A graph G is (colour) critical if $\chi(H) < \chi(G)$ for every proper subgraph H of G , and G is said to be *k-critical* if G is critical and $\chi(G) = k$. The investigation of *k-critical* graphs is a well established part of coloring theory. Actually, criticality is a general concept in graph theory and can be defined with respect to various graph parameters or graph properties. The importance of the notion of criticality lies in the fact that problems for graphs may often be reduced to problems for critical graphs, and the structure of the latter is more restricted. Critical graphs with respect to the chromatic number were first defined and used by Dirac [7] in 1951 ([11]). Stiebitz, Tuza and Voigt introduced and discussed the list critical graphs in [11].

In this paper we introduce critical graphs with respect to the distinguishing number and we discuss some basic properties of *d-distinguishing* critical graphs. More precisely, we characterize connected *d-distinguishing* critical graphs for $d = 1, 2$ in Section 2. We state some necessary condition for 3-distinguishing critical graphs in this section, too. In Section 3, we study disconnected *d-distinguishing* critical graphs and show that there are exactly five 3-distinguishing critical graphs. Also we obtain disconnected *d-*

distinguishing graphs for $d = 5, 6$. Finally we propose two conjectures and a problem in the last section.

2 Characterization of d -distinguishing critical graphs

Criticality is a general concept in graph theory and can be defined with respect to various graph parameters or graph properties. Motivated by chromatic critical graphs and list critical graphs we state the following definition.

Definition 2.1 *A graph G is d -distinguishing critical, if $D(G) = d$ and $D(H) \neq D(G)$, for every proper induced subgraph H of G .*

Let us first discuss some elementary facts about distinguishing critical graphs by the following example.

Example 2.2

- *The complete graph K_n (and also its complement) is an n -distinguishing critical graph for every $n \geq 3$.*
- *The complete bipartite graph $K_{n,n}$ (and also its complement) is a $(n+1)$ -distinguishing critical graph for every $n \geq 3$.*
- *The cycles graphs C_3, C_4 and C_5 are 3-distinguishing critical graphs.*

Since the distinguishing number of each graph and its complement is equal, so we have the following proposition:

Proposition 2.3 *A graph G is d -distinguishing critical, if and only if \overline{G} is a d -distinguishing critical graph.*

First, we determine d -distinguishing critical graphs for all $d \leq 2$. An undirected graph G on at least two vertices is *minimal asymmetric* if G is asymmetric and no proper induced subgraph of G on at least two vertices is asymmetric. Pascal Schweitzer and Patrick Schweitzer in [10] showed that there are exactly 18 finite minimal asymmetric undirected graphs up to isomorphism. These 18 graphs depicted in Figure 1.

Theorem 2.4

- (i) *There is no 1-distinguishing critical graph of order $n \geq 2$.*
- (ii) *The only 2-distinguishing critical graphs are K_2 and $\overline{K_2}$.*

Proof.

- (i) Since K_1 is a proper induced subgraph of G with the distinguishing number 1, so the result follows.




Figure 1: The 18 minimal asymmetric graphs. For each graph the triple $(n, m, co - G)$, describes the number of vertices, edges and the name of the complement graph, respectively. The graphs are ordered first by number of vertices and second by number of edges.

- (ii) Let G be a 2-distinguishing critical graph of order $n \geq 3$. Then there exist at least two distinct vertices of G , say v and w . If H is the proper induced subgraph of G generated by v and w , then $D(H) = 2$, and hence G is not a 2-distinguishing critical graph, which is a contradiction. \square

In sequel, we want to characterize the d -distinguishing critical graphs with $d \geq 3$. We need the following theorem.

Theorem 2.5 [6, 8] *If G is a connected graph with maximum degree Δ , then $D(G) \leq \Delta + 1$. Furthermore, equality holds if and only if G is a K_n , $K_{n,n}$ or C_5 .*

Theorem 2.6 *Let G be a d -distinguishing critical graph with $d \geq 3$ and order n . Then the following properties are satisfied:*

- (i) *If G is connected and $\Delta(G) = d - 1$, then G is K_d , $K_{d-1,d-1}$ or C_5 .*
(ii) *If G is connected and $G \neq K_d, K_{d-1,d-1}, C_5$, then $\Delta(G) \geq d$.*

- (iii) If G and \overline{G} are connected and $G \neq C_5$, then $\delta(G) \leq n - d - 1$.
- (iv) The graph G is a $K_{1,d}$ -free graph, i.e., G does not have the star graph $K_{1,d}$ as its induced subgraph.
- (v) If $G \neq K_d$, then the clique number of G is $\omega(G) \leq d - 1$.
- (vi) If $G \neq \overline{K_d}$, then the independence number of G is $\alpha(G) \leq d - 1$.

Proof. The cases (i) and (ii) follow directly from Theorem 2.5. It can be concluded from case (ii) that $n - 1 - \delta(G) = \Delta(\overline{G}) \geq d$, and this prove the case (iii). To prove the case (iv), it is sufficient to know that $D(K_{1,d}) = d$. For (v), if $\omega(G) \geq d$, then G has a proper induced subgraph K_d with $D(K_d) = d$, which is a contradiction. Finally, if $\alpha(G) \geq d$, then the complement graph $\overline{K_d}$ is a proper induced subgraph of G with the distinguishing number d , which is a contradiction. \square

Theorem 2.7

- (i) Let G be a connected d -distinguishing critical graph for some $d \geq 2$. If G is a triangle free graph, then $G = K_2, K_{d-1, d-1}$, or C_5 .
- (ii) Let G and \overline{G} be two connected graphs. If G is a claw free graph such that $G \neq C_5$, then G is not a d -distinguishing critical graph for any $d \geq 2$.

Proof.

- (i) Let G be a connected d -distinguishing critical graph for some $d \geq 2$, such that $G \neq K_2, K_{d-1, d-1}, C_5$. We show that G has a triangle. Let v be a vertex of G with maximum degree Δ , and $N_G(v) = \{v_1, \dots, v_\Delta\}$. By Theorem 2.6, we have $\Delta \geq d$ and $\alpha(G) \leq d - 1$, so there exist at least two adjacent vertices in $N_G(v)$, say v_1 and v_2 . Hence, the graph generated by vertices v, v_1, v_2 makes a triangle graph as the induced subgraph of G .
- (ii) By contradiction suppose that G is a d -distinguishing critical graph for some $d \geq 2$. So \overline{G} is a d -distinguishing critical graph, by Proposition 2.3. Since G is claw free, so \overline{G} is a triangle free graph. Now by Part (i), and connectivity of G and \overline{G} we conclude that $G = C_5$, which is a contradiction. \square

3 Disconnected distinguishing critical graphs

A vertex v in a graph G of order n , is called an a_1 -vertex if $\deg_G(v) = n - 1$ or there is a vertex $u \in N_G(v)$ such that $N_G(u) \cup N_G(v) = V(G)$; otherwise, u is an a_2 -vertex, see [9]. Malaravan in [9], showed that if $\text{diam}(G) \geq 4$, then every vertex of G is an a_2 -vertex.

Theorem 3.1 [9] *Let G be a graph with no isolates. Then, each vertex of G is an a_2 -vertex in G if and only if \overline{G} is a 2-self-centered graph.*

Theorem 3.2 *If G is a disconnected d -distinguishing critical graph with no isolated vertex, then \overline{G} is a connected 2-self-centered graph.*

Proof. Since G is a disconnected graph, so $\text{diam}(G) \geq 4$, and hence every vertex of G is an a_2 -vertex. Now, we have the result by Theorem 3.1. \square

Let (G, ϕ) denote the labeled version of G under the labeling ϕ . Given two distinguishing k -labelings ϕ and ϕ' of G , we say that ϕ and ϕ' are *equivalent* if there is some automorphism of G that maps (G, ϕ) to (G, ϕ') . We need $D(G, k)$, the number of inequivalent k -distinguishing labelings of G , which was first considered by Arvind and Devanur [3] and Cheng [5] to determine d -distinguishing critical graph with $d \geq 3$.

Lemma 3.3 [3] *Let H be a connected graph. If G consists of $c(G)$ copies of H , i.e., $G = c(G)H$, then $D(G) = \min\{k : D(H, k) \geq c(G)\}$.*

Lemma 3.4 *Let G be a disconnected d -distinguishing critical graph ($d \geq 3$) with $c(G)$ connected components.*

- (i) *The connected components of G are pairwise isomorphic.*
- (ii) *If the connected components of G are isomorphic to graph H , then*
 - 1) $D(nH) < d$ for every $1 \leq n < c(G)$, and
 - 2) $c(G) = D(H, d - 1) + 1$, and $D(H, d) \geq c(G)$, and
 - 3) $c(G) > D(H, D(H))$.

Proof. (i) Let G_{i1}, \dots, G_{is_i} be the connected components of G such that $G_{i1} \cong \dots \cong G_{is_i}$ for every $1 \leq i \leq k$, and also $G_{i1} \not\cong G_{j1}$ for $i \neq j$. Set $H_i = G_{i1} \cup \dots \cup G_{is_i}$. It is clear that $D(G) = \max\{D(H_i)\}_{i=1}^k$. Without loss of generality we can assume that $D(H_1) = \max\{D(H_i)\}_{i=1}^k$. If $k \geq 2$, then the distinguishing number of the proper induced subgraph H_1 of G is $D(G)$, which is a contradiction. Therefore $k = 1$, i.e., all connected components of G are isomorphic to each other.

(ii-1) By Lemma 3.3, we have $D(nH) \leq d$ for every $1 \leq n < c(G)$. If there exist n such that $D(nH) = d$, then the graph $K := nH$ is a proper induced subgraph of G with $D(K) = D(G)$, which is a contradiction.

(ii-2) By part (1), and Lemma 3.3, we have $D((c(G) - 1)H) \leq d - 1$ and $D(H, d - 1) < c(G)$. Thus $D(H, d - 1) \geq c(G) - 1$ and $D(H, d - 1) < c(G)$. Therefore, $c(G) = D(H, d - 1) + 1$.

(ii-3) By contradiction suppose that $c(G) \leq D(H, D(H))$. So $d = D(G) = D(H)$, by Lemma 3.3. Since $c(G) \geq 2$, so H is a proper induced subgraph of G with $D(H) = D(G)$, which is a contradiction. \square

Before we prove the next result we need some additional information about the distinguishing number of complete multipartite graphs. Let $K_{a_1^{j_1}, a_2^{j_2}, \dots, a_r^{j_r}}$ denote the complete multipartite graph that has j_i partite sets of size a_i for $i = 1, 2, \dots, r$ and $a_1 > a_2 > \dots > a_r$.

Theorem 3.5 [6] *Let $K_{a_1^{j_1}, a_2^{j_2}, \dots, a_r^{j_r}}$ denote the complete multipartite graph that has j_i partite sets of size a_i for $i = 1, 2, \dots, r$, and $a_1 > a_2 > \dots > a_r$. Then*

$$D(K_{a_1^{j_1}, a_2^{j_2}, \dots, a_r^{j_r}}) = \min \left\{ p : \binom{p}{a_i} \geq j_i \text{ for all } i \right\}.$$

Since the distinguishing number of a graph and its complement is equal, so if G consists of $c(G)$ copies of complete graph K_s , then $D(c(G)K_s) = \min\{p : \binom{p}{s} \geq c(G)\}$.

Theorem 3.6 *Let G be a disconnected d -distinguishing critical graph ($d \geq 3$) with $c(G)$ isomorphic connected components H , and the independence number $\alpha(G)$.*

- (i) *If $c(G) \geq \frac{d}{2}$, then $G = c(G)K_s$, where $c(G) = \binom{d-1}{s} + 1$ and $\binom{d}{s} \geq c(G)$.*
- (ii) *If $\alpha(G) = c(G)$, or $\alpha(G)$ is a prime number, then $G = c(G)K_s$, where $c(G) = \binom{d-1}{s} + 1$ and $\binom{d}{s} \geq c(G)$.*

Proof.

- (i) We consider the two following cases:

Case 1) If $c(G) = d$, then by choosing one vertex from each component of G we can make the induced subgraph $\overline{K_d}$ of G . Since G is a d -distinguishing critical graph, so $G = \overline{K_d}$.

Case 2) If $c(G) \leq d - 1$, then $G \neq \overline{K_d}$, and so $\alpha(G) \leq d - 1$, by part (vi) of Theorem 2.5. Since $\alpha(G) = c(G)\alpha(H)$ and $d \leq 2c(G)$, so we obtain $\alpha(H) = 1$, and hence H is a complete graph, say K_s . Thus $G = c(G)K_s$. Now by $D(G) = d$ and $D(c(G)K_s) = \min\{p : \binom{p}{s} \geq c(G)\}$ (by Theorem 3.5), we have

$$d = \min \left\{ p : \binom{p}{s} \geq c(G) \right\}. \quad (1)$$

On the other hand the distinguishing number of the proper induced subgraph $H = (c(G) - 1)K_s$ is $D(H) \leq d - 1$, by Lemma 3.4 (i). In fact $D(H) = d - 1$, since otherwise $D(H) \leq d - 2$, and so $D(H \cup K_s) = D(G) \leq d - 1$, which is a contradiction. Thus

$$d - 1 = \min \left\{ p : \binom{p}{s} \geq c(G) - 1 \right\}. \quad (2)$$

Now using Equations (1) and (2), we conclude that $\binom{d-1}{s} = c(G) - 1$ and $\binom{d}{s} \geq c(G)$.

- (ii) If $\alpha(G) = c(G)$, or $\alpha(G)$ is a prime number, then we obtain $\alpha(H) = 1$, from $\alpha(G) = c(G)\alpha(H)$. Hence H is a complete graph, say K_s . The remaining proof is the same as the proof of case (2) of Part (i). \square

To determine exactly 3-distinguishing critical graphs we need the following lemma.

Lemma 3.7 *If G is a 3-distinguishing critical graph, then $\Delta(G) \leq 2$.*

Proof. By contradiction, suppose that G has a vertex v of degree at least three. Let w, y, z be three distinct adjacent vertices to v . If at least the two vertices of w, y, z are adjacent, say w, y , then the proper induced subgraph H of G generated by v, w, y is a triangle with the distinguishing number 3, which is a contradiction. If non of vertices w, y, z are adjacent, then the proper induced subgraph H of G generated by v, w, y, z is the star graph $K_{1,3}$ with the distinguishing number 3, which is a contradiction. \square

Theorem 3.8 *There are exactly five 3-distinguishing critical graphs, $C_3, C_4, C_5, \overline{C_3}, \overline{C_4}$.*

Proof. By Theorem 2.5 and Lemma 3.7 it is easy to see that the graphs C_3, C_4, C_5 are all connected 3-distinguishing critical graph. Also the complements of graphs C_3, C_4, C_5 are the only disconnected 3-distinguishing critical graphs, by Theorem 3.6 (i). \square

Theorem 3.9 *If G is a disconnected d -distinguishing critical graph ($d \geq 5$) with $c(G)$ isomorphic connected components H , such that $c(G) < \frac{d}{2}$, then $d = D(H)+1$, $D(H) \geq 4$ and $D(H, D(H)) < \frac{D(H)-1}{2}$.*

Proof. By part (ii-1) of Lemma 3.4, we can suppose that $d = D(H) + i$ for some $i \geq 1$. By Lemma 3.3, we have $c = D(H, D(H)+i-1)+1$. Since there exist $\binom{D(H)+i-1}{D(H)}$ distinct sets with $D(H)$ labels, so $\binom{D(H)+i-1}{D(H)} \leq D(H, D(H) + i - 1)$. Now since $c(G) < \frac{d}{2}$, so we can conclude that

$$\binom{D(H) + i - 1}{D(H)} < \frac{D(H) + i - 2}{2}. \quad (3)$$

It can be seen that if $i \geq 2$, then by Equation (3) we have $D(H) \leq 0$, which is a contradiction. Thus $i = 1$, and so we obtain $D(H) \geq 4$ from Equation (3). Also $c(G) = D(H, D(H)) + 1$. The last inequality follows directly from $c(G) < \frac{d}{2}$. \square

Corollary 3.10 *If G is a disconnected d -distinguishing critical graph ($d \geq 5$) with $c(G)$ isomorphic connected components K_s (the complete graph of order s), such that $c(G) < \frac{d}{2}$, then $s = d - 1$, and $G = 2K_{d-1}$.*

Proof. It is sufficient to know that $D(K_s, D(K_s)) = D(K_s, s) = 1$. Now the result follows from Theorem 3.9. \square

Lemma 3.11 *Let G be a disconnected d -distinguishing critical graph ($d \geq 5$) with $c(G)$ isomorphic connected components H such that H is not a complete graph, and also $c(G) < \frac{d}{2}$. Let ϕ be an arbitrary distinguishing labeling of H with $D(H)$ labels and $X_i = \{x \in V(H) : \phi(x) = i\}$ where $i = 1, 2, \dots, D(H)$. If the degrees sequence of vertices in X_i is denoted by the sequence S_i , then for every $i, j \in \{1, 2, \dots, D(H)\}$ we have $S_i = S_j$.*

Proof. Let n_i be the number of vertices of degree p in X_i . By contradiction assume that there exists a degree p for which $B_p \neq \emptyset$ where $A_p = \{X_k : n_k = n_1\}$ and $B_p = \{X_k : n_k \neq n_1\}$. We consider the two following cases:

Case 1) If $|A_p| < \frac{D(H)-1}{2}$, then $|B_p| \geq \frac{D(H)-1}{2}$. For every $X_k \in B_p$, we define the distinguishing labeling φ_k of H as follows:

$$\varphi_k(x) = \begin{cases} \phi(x) & x \notin X_1, X_k, \\ 1 & x \in X_k, \\ k & x \in X_1. \end{cases}$$

Since the number of vertices of degree p in X_k is not equal with the number of vertices of degree p in X_1 , we conclude that the distinguishing labelings φ_k, X_k , are nonisomorphic distinguishing labeling to each other and also to ϕ . Since $|B| \geq \frac{D(H)-1}{2}$, so $D(H, D(H)) \geq \frac{D(H)-1}{2}$, which is a contradiction by Theorem 3.9.

Case 2) If $|A_p| \geq \frac{D(H)-1}{2}$. By a similar reasoning as Case 1, we obtain a contradiction. \square

Theorem 3.12 *Let G be a disconnected d -distinguishing critical graph ($d \geq 5$) with $c(G)$ isomorphic connected components H such that H is not a complete graph, and also $c(G) < \frac{d}{2}$. If $\{p_1, \dots, p_s\}$ is the set of degrees of vertices of G , then the number of vertices of degree p_i in G is at least $2c(G)(d-1)$, for any $1 \leq i \leq s$.*

Proof. Let ϕ be a distinguishing labeling of H with $D(H)$ labels and $X_i = \{x \in V(H) : \phi(x) = i\}$ where $i = 1, 2, \dots, D(H)$. It is clear that $\{p_1, \dots, p_s\}$ is the set of degrees of vertices of H , too. By Lemma 3.11, we know that the number of vertices of degree p_i in each of sets $X_1, \dots, X_{D(H)}$ are equal, for any $1 \leq i \leq s$. Let t_i is the number of vertices of degree p_i in each of sets $X_1, \dots, X_{D(H)}$. First, we prove that $t_i \geq 2$ for any $1 \leq i \leq s$. By contradiction, we assume that there exists $p \in \{p_1, \dots, p_s\}$ such that the number of vertices of degree p in each of sets $X_1, \dots, X_{D(H)}$ is 1. So we have exactly $D(H)$ vertices of degree p in graph H , say $v_1, \dots, v_{D(H)}$ such that $v_i \in X_i$ where $1 \leq i \leq D(H)$. We claim that the restriction of the automorphism group of H to the set $\{v_1, \dots, v_{D(H)}\}$ is isomorphic to the permutation group $\mathbb{S}_{D(H)}$.

To prove our claim note that for every $a, b \in \{1, \dots, D(H)\}$, we have a labeling $\phi_{a,b}$ of $V(H)$ with $X_i^{a,b} = \{x \in V(H) : \phi_{a,b}(x) = i\}$ such that $X_a^{a,b} = X_a - \{v_a\}$, $X_b^{a,b} = X_b \cup \{v_a\}$, and $X_i^{a,b} = X_i$ for every $i \in \{1, \dots, D(H)\} \setminus \{a, b\}$. Since the number of vertices of degree p in $X_a^{a,b}$ and $X_b^{a,b}$ are distinct, so the labeling $\phi_{a,b}$ is not distinguishing, by Lemma 3.11. Hence, there exists a nonidentity automorphism $f_{a,b}$ of H such that $f_{a,b}$ preserves the labeling $\phi_{a,b}$. Thus $f_{a,b}(v_a) = v_b$ and $f_{a,b}(v_b) = v_a$, since otherwise $f_{a,b}(v_b) = v_b$ and $f_{a,b}(v_a) = v_a$, and so $f_{a,b}$ preserves the distinguishing labeling ϕ , and hence $f_{a,b}$ should be the identity automorphism, which is a contradiction. Therefore $f_{a,b}(v_a) = v_b$, $f_{a,b}(v_b) = v_a$, and $f_{a,b}(v_i) = v_i$ for every $i \in \{1, \dots, D(H)\} \setminus \{a, b\}$. Thus for any $a, b \in \{1, \dots, D(H)\}$, there exists an automorphism $f_{a,b}$ of H such that the restriction of $f_{a,b}$ to $v_1, \dots, v_{D(H)}$ is isomorphic to the transposition (a, b) in $\mathbb{S}_{D(H)}$.

Hence, we can conclude that the restriction of automorphism group of H to the set $\{v_1, \dots, v_{D(H)}\}$ is isomorphic to the permutation group $\mathbb{S}_{D(H)}$, and this proves the claim.

Now since H is a connected graph, so by claim the set of vertices $\{v_1, \dots, v_{D(H)}\}$ generates the induced subgraph $K_{D(H)}$ of H . Since $c(G) \geq 2$, so we can make the induced subgraph $K = K_{D(H)} \cup K_{D(H)}$ of G (here \cup is the disjoint union symbol) with $D(K) = D(H) + 1$. On the other hand, $d = D(H) + 1$ by Theorem 3.9, and so $D(K) = d$. Since H is not a complete graph, so K is a proper induced subgraph of G with $D(K) = D(G)$, which is contradiction to that G is a d -distinguishing critical graph. Therefore $t_i \geq 2$, for any $1 \leq i \leq s$.

To complete the proof, it is sufficient to know that the number of vertices of degree p_i in G is $c(G)D(H)t_i$ for any $1 \leq i \leq s$, and also $D(H) = d - 1$. \square

Theorem 3.13 *If G is a disconnected d -distinguishing critical graph ($d \geq 5$) with $c(G)$ isomorphic connected components H , such that $c(G) < \frac{d}{2}$, then $D(H) \mid |V(H)|$.*

Proof. Let ϕ be an arbitrary distinguishing labeling of H with $D(H)$ labels and $X_i = \{x \in V(H) : \phi(x) = i\}$ where $i = 1, 2, \dots, D(H)$. By Lemma 3.11, we have $D(H) \mid |X_i| = |V(H)|$, for every $i = 1, 2, \dots, D(H)$, and so $D(H) \mid |V(H)|$. \square

Theorem 3.14 *Let G be a disconnected d -distinguishing critical graph ($d \geq 5$) with $c(G)$ isomorphic connected components H such that $c(G) < \frac{d}{2}$. If $\overline{G} \neq K_{d-1, d-1}, K_d$, then $c(G) \leq \frac{d-1}{3}$, and so $D(H, D(H)) \leq \frac{D(H)-3}{3}$. Also, $d \geq 7$ and $D(H) \geq 6$.*

Proof. Since \overline{G} is a connected d -distinguishing critical graph ($d \geq 5$), so \overline{G} has a triangle as a induced subgraph, by Theorem 2.7. Thus G has a claw graph as a induced subgraph. Since G is disjoint union of connected graphs H , so each of connected components H has a claw. Hence, $\alpha(H) \geq 3$, and using $\alpha(G) = \alpha(H)c(G)$ and $\alpha(G) \leq d - 1$ by Theorem 2.6, we get $3c(G) \leq d - 1$, and so $D(H, D(H)) \leq \frac{D(H)-3}{3}$, because of $c(G) = D(H, D(H)) + 1$ and $d = D(H) + 1$ by Lemma 3.4. Finally, by $3c(G) \leq d - 1$, $c(G) \geq 2$ and $d = D(H) + 1$, we get $d \geq 7$ and $D(H) \geq 6$. \square

Here, we can determine exactly all disconnected d -distinguishing critical graphs for $d = 5, 6$.

Corollary 3.15

- (i) *The only disconnected 5-distinguishing critical graphs are $\overline{K_5}$ and $\overline{K_{4,4}}$.*
- (ii) *The only disconnected 6-distinguishing critical graphs are $\overline{K_6}$ and $\overline{K_{5,5}}$.*

Proof. It follows directly from Theorem 3.14. \square

Now we close this section by determining the distinguishing critical trees. To do this, we need the following theorem:

Theorem 3.16 [6] *If T is a tree of order $n \geq 3$, then $D(T) \leq \Delta(T)$. Furthermore, equality is achieved if and only if T is a symmetric tree or a path of odd length.*

Theorem 3.17 *The only distinguishing critical tree is K_2 .*

Proof. Let T be a d -distinguishing critical tree of order $n \geq 3$. By Theorem 3.16, we have $d \leq \Delta(T)$. Let v be the vertex of T with maximum degree $\Delta(T)$. Since $\Delta \geq d$ and T does not have any cycle, so we can make the star graph $K_{1,d}$ as a proper induced subgraph of G . Since $D(K_{1,d}) = d$ and $T \neq K_{1,d}$ (T is a d -distinguishing critical tree), so we have a contradiction. \square

4 Conclusion

In this paper we characterized the d -distinguishing critical graphs. In particular, we could determine exactly all d -distinguishing critical graphs for $d = 1, 2, 3$. It can be seen that all such d -distinguishing critical graphs are k -regular graph for some $k \leq d$. Also, we showed that the disconnected d -distinguishing critical graph with c connected components such that $c \geq \frac{d}{2}$, is a regular graph. We propose the following conjecture:

Conjecture 4.1

- (i) *If G is a d -distinguishing critical graph, then G is a k -regular graph for some $k \leq d$.*
- (ii) *If G is a disconnected d -distinguishing critical graph, then each components of G is a complete graph.*

We can extend the definition of d -distinguishing critical graph to a strong version. A graph G is called *strong d -distinguishing critical*, if $D(G) = d$ and $D(G - v) \neq D(G)$ for every vertex v of G , where $G - v$ denotes the graph obtained from G by removal of a vertex v and all edges incident to v . It is immediate that any d -distinguishing critical graph is a strong d -distinguishing critical graph, but the converse is not true; the minimal asymmetric graphs are strong 1-distinguishing critical graphs which are not 1-distinguishing critical graphs. This leads us to ask the following.

Problem 4.2 *Characterize the strong d -distinguishing critical graphs.*

References

- [1] M.O. Albertson and K.L. Collins, *Symmetry breaking in graphs*, Electron. J. Combin. 3 (1996), #R18.
- [2] M.O. Albertson, *Distinguishing Cartesian product of graphs*, Electron. J. Combin. 12 (2005), #N17.

- [3] V. Arvind and N. Devanur, *Symmetry breaking in trees and planar graphs by vertex coloring*, In Proceedings of the Nordic Combinatorial Conference, 2004.
- [4] F. Buckley, *Self-centered graphs*, in: Graph theory and its applications: East 359 and West (Jinan, 1986), Vol. 576, Ann. New York Acad. Sci., New York 360 Acad. Sci., New York, 1989, pp. 71-78.
- [5] C. Cheng, *On computing the distinguishing numbers of trees and forests*, Electron. J. Combin. 13 (2006), #R11.
- [6] K.L. Collins and A.N. Trenk, *The distinguishing chromatic number*, Electron. J. Combin. 13 (1) (2006), #R16.
- [7] G.A. Dirac, *Note on the colouring of graphs*, Math. Z. 54 (1951), 347-353.
- [8] S. Klavžar, T.-L. Wong, X. Zhu, *Distinguishing labelings of group action on vector spaces and graphs*, J. Algebra. 303 (2006) 626-641.
- [9] A.C. Malaravan, *A study on center of a graph complement*, Discrete Math. Algorithm. Appl. 7 (04) (2015), 1550046.
- [10] P. Schweitzer, P. Schweitzer, *Minimal Asymmetric Graphs*, arXiv preprint arXiv:1605.01320 (2016).
- [11] M. Stiebitz, Z. Tuza, and M. Voigt, *On list critical graphs*, Discrete Math. 309 (2009), 4931-4941.