

RELATIVE DING AND K -STABILITY OF TORIC FANO MANIFOLDS IN LOW DIMENSIONS

YASUFUMI NITTA, SHUNSUKE SAITO, AND NAOTO YOTSUTANI*

Abstract. The purpose of this paper is to clarify all of the uniformly relatively Ding stable toric Fano threefolds and fourfolds as well as unstable ones. The key player in our classification result is the Mabuchi constants, which can be calculated by combinatorial data of the associated moment polytopes due to the work of Yao [33]. In Tables 1–3, we give the list of uniform relative Ding stability of all toric Fano manifolds in dimension up to four with the values of the Mabuchi constants.

As an application of our main theorem (Theorem 1.1), we clarify the difference between relative K -stability and relative Ding stability by considering some specific toric Fano manifolds (Corollaries 1.6 and 1.9). In the proof of Corollary 1.9, we used Bott tower structure of relatively Ding unstable toric Fano manifolds.

1. INTRODUCTION

In [27], Mabuchi generalized the notion of Kähler-Einstein metrics to Fano manifolds with non-vanishing Futaki invariant as follows: let X be an n -dimensional Fano manifold defined over \mathbf{C} with complex structure J . Denoting the identity component of the automorphism group of (X, J) by $\text{Aut}^0(X)$, we fix a maximal algebraic torus T of $\text{Aut}^0(X)$. Let $S := \text{Hom}(\mathbf{C}^*, T) \otimes_{\mathbf{Z}} S^1$ be the maximal compact torus of T and $\mathcal{H}(X)^S$ the space of all S -invariant Kähler forms representing $c_1(X)$. Let F_ω be the Ricci potential of ω , that is,

$$\text{Ric}(\omega) - \omega = \frac{\sqrt{-1}}{2\pi} \partial\bar{\partial}F_\omega, \quad \text{and} \quad \int_X (1 - e^{F_\omega}) \frac{\omega^n}{n!} = 0.$$

A Kähler form $\omega \in \mathcal{H}(X)^S$ is said to be *Mabuchi soliton*, if the Riemannian gradient

$$\text{grad}_\omega(1 - e^{F_\omega})$$

with respect to ω is real holomorphic. For each $\omega \in \mathcal{H}(X)^S$, let $\mathfrak{s}_0(X, \omega)$ be the subspace of $C^\infty(X, \mathbf{R})^S$ consisting of all $\phi \in C^\infty(X, \mathbf{R})^S$ so that the Hamiltonian real vector field $2\pi J \text{grad}_\omega \phi$ lies in $\text{Lie}(S)$, where $\text{Lie}(S)$ is the Lie algebra of the compact torus S and $\int_X \phi \omega^n / n! = 0$. Let $\text{pr}_\omega : C^\infty(X, \mathbf{R})^S \rightarrow \mathfrak{s}_0(X, \omega)$ be the L^2 -orthogonal projection with respect to ω . Mabuchi also introduced the constant

$$(1.1) \quad M_X := \max_X \text{pr}_\omega(1 - e^{F_\omega})$$

and showed that this is independent of the choice of ω . In particular, the existence of Mabuchi solitons implies $M_X < 1$. Throughout the paper, we call M_X the *Mabuchi constant* of X .

Recently, Mabuchi solitons were rediscovered by Yao [33] following Donaldson's infinite dimensional moment map picture. He showed that Mabuchi soliton is a critical point

Key words and phrases. Relative stability, Fano manifolds, toric variety.

* Corresponding author.

of the functional

$$\mathcal{H}(X)^S \ni \omega \mapsto \int_X (1 - e^{F_\omega})^2 \frac{\omega^n}{n!} \in \mathbf{R},$$

which can be interpreted as the norm square of Donaldson's new moment map given in [17]. For a general (not necessarily toric) Fano manifold, the corresponding version of Yau-Tian-Donaldson conjecture regarding Mabuchi soliton has been studied by many researchers. Han-Li [22] proved that *uniform relative Ding stability* and $M_X < 1$ imply the existence of Mabuchi solitons building upon the work of variational approach [7]. In [34], Yao showed that additional assumption $M_X < 1$ can be derived from uniform relative Ding stability. We also mention that Han-Li's work [22] deals with more general g -solitons which include Mabuchi solitons and Kähler-Ricci solitons on (possibly singular) Fano varieties. In the toric case, Yao proved equivalences of $M_X < 1$, the existence of Mabuchi solitons, and uniform relative Ding stability for toric Fano manifolds (or more generally toric Fano orbifolds).

The notion of uniform relative Ding stability for toric Fano manifolds was introduced by Yao [33] as a generalization of Ding stability ([8] and [18]) and uniform Ding stability ([7] and [9]). Yao's work is the toric reduction of relative Ding stability of general Fano manifolds. One of the advantage of our toric setting is that everything can be described in terms of convex functions on the associated moment polytopes. In fact, Delcroix [15] and Li-Li [26] showed that there is a one-to-one correspondence between T -equivariant test configurations of a polarized toric manifold and rational piecewise affine convex functions on the associated moment polytope. See Sections 2.3 for more details.

Let $v := \text{grad}_\omega \text{pr}_\omega(1 - e^{F_\omega})$ be the extremal Kähler vector field of $(X, -K_X)$ with respect to the maximal algebraic torus $T = (\mathbf{C}^*)^n$ defined in Futaki-Mabuchi [19], and $\langle \cdot, \cdot \rangle$ the bilinear pairing for \mathbf{C}^* -actions introduced by Székelyhidi [31]. Uniform relative Ding stability requires the existence of a constant $\delta > 0$ such that

$$D_v^{NA}(\mathcal{X}, \mathcal{L}) := D^{NA}(\mathcal{X}, \mathcal{L}) + \langle (\mathcal{X}, \mathcal{L}), v \rangle \geq \delta J_T^{NA}(\mathcal{X}, \mathcal{L})$$

holds for any T -equivariant test configuration $(\mathcal{X}, \mathcal{L})$ of $(X, -K_X)$, where D^{NA} and J_T^{NA} denote the *non-Archimedean Ding functional* (introduced in [8] and [18]) and the *non-Archimedean reduced J-functional* with respect to T (introduced in [24]) respectively. Later, this condition for (not necessarily toric) Fano manifolds was studied in [34].

In this paper, we focus on (uniform) relative Ding stability and relative K -stability of toric Fano manifolds. See Section 2.3 for definitions of relative Ding/ K -stability. More precisely, we firstly determine which toric Fano manifolds in dimension three and four are uniformly relatively Ding stable by direct computations of the Mabuchi constants. We secondly find explicit examples of toric Fano manifolds which are relatively K -stable but which are relatively Ding unstable (see Corollaries 1.6 and 1.9). For the proof of Theorem 1.1, we refer the reader to see Tables 2–3.

Theorem 1.1. (1) *In dimension three, 12 toric Fano manifolds out of 18 are uniformly relatively Ding stable and the rest are relatively Ding unstable.*
 (2) *In dimension four, 49 toric Fano manifolds out of 124 are uniformly relatively Ding stable and the rest are relatively Ding unstable.*

Note that the uniform relative Ding stability of all Gorenstein toric del Pezzo surfaces was verified in [33, Example 5.14] (see also Table 1 in Section 3.2 for smooth surfaces). We simultaneously complete the classification of toric Fano threefolds and fourfolds admitting Mabuchi solitons by Yao's result. The key theorem to achieve this is the following. (We refer the reader to Section 2 for notations used in Proposition 1.2.)

Proposition 1.2 (Yao [33]). *Let P be a Fano polytope. Then we have the following:*

- (a) *The following conditions are equivalent.*
 - (1) X_P is uniformly relatively Ding stable.
 - (2) There exists $\delta > 0$ such that $D_v^{NA}(f) \geq \delta \int_P f(x) dx$ for any normalized rational PL convex function f on P .
 - (3) $M_{X_P} = \max_P \theta_P < 1$, where θ_P is the potential function defined in Section 2.3.
- (b) *The following conditions are equivalent.*
 - (1) X_P is relatively Ding stable.
 - (2) X_P is relatively Ding semistable.
 - (3) $M_{X_P} \leq 1$.
- (c) *The following conditions are equivalent.*
 - (1) X_P is relatively Ding unstable.
 - (2) $M_{X_P} > 1$.

For reader's convenience, we shall give a proof of Proposition 1.2 in Section 2.3. Theorem 1.1 and Proposition 1.2 offer concrete examples of Fano manifolds which admit Mabuchi solitons.

Corollary 1.3. (1) *In dimension three, 12 toric Fano manifolds admit Mabuchi solitons and the others do not.*

(2) *In dimension four, 49 toric Fano manifolds admit Mabuchi solitons and the others do not.*

With remarkable progress made in research in Ding stability, it is known that uniform Ding stability is equivalent to Ding stability for general Fano manifolds. Keeping this in our mind, we see that our computation also reveals the following corollary.

Corollary 1.4. *Up to dimension four, there are no toric Fano manifolds satisfying $M_X = 1$. Hence for each toric Fano n -fold with $n \leq 4$, relative Ding stability is equivalent to uniform relative Ding stability.*

In [33, Example 5.14], Yao presented examples of Gorenstein toric del Pezzo surfaces with $M_X = 1$. However, any example of Fano manifolds with $M_X = 1$ has not yet known. Hence it is natural to expect that there are no Fano manifolds with the Mabuchi constant being precisely 1. If this is the case, uniform relative Ding stability is equivalent to relative Ding stability for arbitrary toric Fano manifold.

Finally, we clarify the difference between relative K -stability and relative Ding stability with specific examples. Toric reduction of relative K -stability was studied by Zhou and the third author in [35], where they established instability criterion of relative K -stability in terms of the associated moment polytope. More precisely, for a given Fano polytope P , we consider $P^- = \{x \in P \mid 1 - \theta_P(x) \leq 0\}$ where $\theta_P(x) = \sum \theta_i x_i + c$ is the potential function of P (see, Section 2.3 for the definition). If $\text{Int}(P^-) \neq \emptyset$ and the inequality

$$(1.2) \quad 1 - c < \frac{\int_{P^-} (1 - \theta_P)^2 dx}{\text{vol}(P^-)}$$

holds, then the corresponding toric Fano manifold $(X_P, -K_{X_P})$ is relatively K -unstable (see [35, Theorem 1.4 (2)]). Although the instability criterion (1.2) is correct, there is a technical mistake in [35] when they apply (1.2) into six toric Fano threefolds $\mathcal{B}_1, \mathcal{C}_2, \mathcal{D}_1, \mathcal{D}_2, \mathcal{E}_1$ and \mathcal{E}_2 . Here we used same notation in Section 3.3, Table 2. In fact, the first and the third authors recomputed the values of the right hand side in (1.2), and they found that

the above six toric Fano threefolds ($\mathcal{B}_1, \mathcal{C}_2, \mathcal{D}_1, \mathcal{D}_2, \mathcal{E}_1$ and \mathcal{E}_2) do *not* satisfy the instability condition (1.2) by using the computational package [10] (see Section 3.3 for the detail).

However, we can confirm relative K -stability for some of toric Fano manifolds by the existence theorem of extremal Kähler metrics due to Guan [20, Theorem 2], Hwang [25, Corollary 1.2], Apostolov-Calderbank-Gauduchon-Tønnesen-Friedman [1, Proposition 11] as follows.

Proposition 1.5 (Guan, Hwang, ACGT-F). *(1) For the toric Fano threefold $\mathbf{P}_{\mathbf{P}^2}(\mathcal{O} \oplus \mathcal{O}(2))$ which is labeled \mathcal{B}_1 in Table 2, it admits an extremal Kähler metric in every Kähler classes. (2) For the toric Fano fourfold $\mathbf{P}_{\mathbf{P}^3}(\mathcal{O} \oplus \mathcal{O}(k))$ for $k = 3$ (resp. $k = 2$) which is labeled B_1 (resp. B_2) in Table 3, it admits an extremal Kähler metric in every Kähler classes.*

In [36], Zhou-Zhu proved that relative K -stability is a necessary condition for the existence of extremal Kähler metrics on polarized toric manifolds. Combining this, Proposition 1.5 and Theorem 1.1, we have the following.

Corollary 1.6. *(1) The toric Fano threefold \mathcal{B}_1 is relatively K -stable, but it is relatively Ding unstable. (2) Moreover, toric Fano fourfolds B_1 and B_2 are relatively K -stable, but they are relatively Ding unstable.*

After we posted the third version of this paper on arXiv, the third author found that we may adapt the *Bott Manifold structure* for discovering more examples of toric Fano manifolds which clarify the difference between relative Ding stability and relative K -stability as follows.

Recall that a Kähler manifold is called *Calabi dream manifold* if it admits an extremal Kähler metric in each Kähler class [11, Definition 1.6]. Hence the Hirzebruch surfaces, toric Fano manifolds appeared in Proposition 1.5 are examples of Calabi dream manifolds. The key to make them Calabi dream manifolds is *admissible construction* that has been so successful in producing many explicit examples of extremal Kähler metrics [1]. This construction was applied to *Bott manifolds* and many interesting explicit examples of extremal Kähler metrics on certain stage n Bott manifolds were discovered [2]. See, Section 3.5 for the definition of the *stage k Bott manifold of the Bott tower of height n* . We remark that any Bott manifold is toric since it can be constructed as a quotient of n copies of $\mathbf{C}_*^2 := \mathbf{C}^2 \setminus \{0\}$ by an algebraic torus $T = (\mathbf{C}^*)^n$. The number of holomorphically nontrivial \mathbf{P}^1 -bundles in the Bott tower is called the *twist*. In [2, Proposition 5.4], it has been proved that the following types of Bott manifolds with twist one admit an extremal Kähler metric in every Kähler class.

Proposition 1.7 (BCT-F). *For $\mathbf{k} = (k_1, k_2, \dots, k_{n-1}) \in \mathbb{Z}^{n-1}$ with $\prod_{i=1}^{n-1} k_i \neq 0$, let $X_n(\mathbf{k})$ be the stage n Bott manifold of the Bott tower of height n*

$$(1.3) \quad \mathbf{P}_{(\mathbf{P}^1)^{n-1}}(\mathcal{O} \oplus \mathcal{O}(k_1, k_2, \dots, k_{n-1})) \rightarrow (\mathbf{P}^1)^{n-1}$$

which is the \mathbf{P}^1 -bundle over the product of projective lines. Then $X_n(\mathbf{k})$ is a Calabi dream manifold.

For our purpose, we shall consider *Fano Bott manifold* which is a special class of toric Fano manifold. According to the classification results of the Bott manifolds [12, 23, 13], there are five classes of Fano Bott manifolds at stage three, and 13 classes of Fano Bott manifolds at stage four. More precisely, we have the following.

Proposition 1.8. *Fano Bott manifolds at stage three are exactly type C in Table 2, and Fano Bott manifolds at stage four are exactly type L in Table 3.*

See [12, Remark 4.5], [23, Section 5.3.2], and see also [2, Proposition 3.21] for the proof. Among these 18 classes of Fano Bott manifolds at stage n ($n \leq 4$), there is only one Fano Bott manifold with twist one which is relatively Ding unstable, that is,

Corollary 1.9. *The toric Fano fourfold $\mathbf{P}_{(\mathbf{P}^1)^3}(\mathcal{O} \oplus \mathcal{O}(1, 1, 1))$ which is labeled L_1 in Table 3, is a relatively Ding unstable Calabi dream manifold.*

We refer the reader to Section 3.5 for further details.

Acknowledgements. This work was supported by first author's JSPS KAKENHI Grant Number JP21K03234, second author's JP20K14321, and third author's JP18K13406. The third author would like to thank Professors, Vestislav Apostolov, C.W. Tønnesen-Friedman, Mikiya Masuda, Yi Yao for helpful comments. Finally, we are grateful to the referees for comments which improved the presentation of our manuscript.

2. PRELIMINARIES

2.1. Polarized toric varieties. We first recall the construction of polarized toric varieties, as can be found in [14]. Let $P \subset \mathbf{R}^n$ be an n -dimensional integral convex polytope. Define

$$C(P) := \{(x, r) \in \mathbf{R}^n \times \mathbf{R} \mid r > 0, r^{-1}x \in P\} \cup \{0\}$$

to be the cone over $P \times \{1\}$ with the vertex at the origin. Then the semigroup $S_P := C(P) \cap \mathbf{Z}^{n+1}$ is finitely generated by Gordan's lemma. Let $\mathbf{C}[S_P]$ denote its semigroup algebra. The character corresponding to $(m, k) \in S_P$ is χ^{mt^k} and $\mathbf{C}[S_P]$ is graded by height, i.e., $\deg(\chi^{mt^k}) = k$. Consequently, we obtain the graded \mathbf{C} -algebra

$$\mathbf{C}[S_P] = \bigoplus_{k=0}^{+\infty} R_k, \quad R_k := \{f \in \mathbf{C}[S_P] \mid \deg f = k\}$$

from the polytope P . We define the polarized toric variety (X_P, L_P) by

$$(X_P, L_P) := (\text{Proj } \mathbf{C}[S_P], \mathcal{O}_{X_P}(1)).$$

It is well-known that X_P is a smooth projective variety if and only if P is Delzant.

An alternative way to construct toric varieties is to take a GIT quotient of affine n -space by a linear torus action. Then it turns out that every n -dimensional projective toric variety can be written as a quotient of \mathbf{C}^n by some torus action (cf. Section 3.5).

2.2. Fano polytopes. Let $P \subset \mathbf{R}^n$ be an n -dimensional integral Delzant polytope containing the origin in its interior. The dual polytope P^\vee of P is defined by

$$P^\vee = \{y \in \mathbf{R}^n \mid \langle x, y \rangle \geq -1 \text{ for all } x \in P\},$$

which is an n -dimensional rational polytope containing the origin in its interior. P is said to be *reflexive* if P^\vee is again an integral polytope. We call P a *Fano polytope* if it is a reflexive Delzant polytope. Remark that we considered the Euclidean space \mathbf{R}^n including P as $M_{\mathbf{R}} = \text{Hom}(T, \mathbf{C}^*) \otimes \mathbf{R}$, and its dual \mathbf{R}^n containing the extremal Kähler vector field as $N_{\mathbf{R}} = \text{Hom}(\mathbf{C}^*, T) \otimes \mathbf{R}$, the latter identified with the real part of $\text{Lie}(T)$. For example, the standard n -simplex is a Fano polytope. It is well-known that there is a bijective correspondence between isomorphism classes of Fano polytopes and isomorphism classes of toric Fano manifolds with the anticanonical polarization.

We will use the notation of the Batyrev classification (3-dimensional case) [4] (see also [32] for another geometrical description of toric Fano threefolds), and the Batyrev-Sato classification (4-dimensional case) of Fano polytopes ([5] and [30]).

2.3. Relative Ding stability. We recall the notion of uniform relative Ding stability of toric Fano manifolds. Suppose that P is an n -dimensional Fano polytope in \mathbf{R}^n . A function $f: P \rightarrow \mathbf{R}$ is called *rational piecewise affine convex* or *rational PL convex* if f is a convex function of the form

$$f(x) = \max\{\ell_1(x), \dots, \ell_m(x)\}$$

with each ℓ_k a rational affine function. Moreover, f is said to be *normalized* if it satisfies

$$\min_P f = f(0) = 0.$$

Note that each rational PL convex function can be assumed to be normalized after adding some rational affine function.

Here and hereafter, we denote the maximal algebraic torus $(\mathbf{C}^*)^n$ by T . The importance of such functions comes from the fact that each rational PL convex function f corresponds to a T -equivariant test configuration of $(X_P, -K_{X_P})$ as follows. We refer the reader to [16, Section 4.2] for more detailed description to explain why all T -equivariant test configurations can be induced by a rational PL convex function.

For a fixed integer L with $L > \max_P f$, we obtain an $(n + 1)$ -dimensional rational convex polytope

$$(2.1) \quad \mathcal{P} := \{(x, y) \in \mathbf{R}^n \times \mathbf{R} \mid x \in P, 0 \leq y \leq L - f(x)\}$$

and some multiple of (2.1) gives an $(n + 1)$ -dimensional polarized toric variety $(\mathcal{X}_f, \mathcal{L}_f)$ as in Section 2.1. This can be regarded as a one parameter degeneration of $(X_P, -K_{X_P})$ into a (possibly non-normal) toric variety defined by the ‘‘roof’’ of $r\mathcal{P}$. Specifically, it has the structure of a $T \times \mathbf{C}^*$ -equivariant flat projective toric morphism $\pi: \mathcal{X}_f \rightarrow \mathbf{P}^1$ and then the test configuration is $\mathcal{X}_f \setminus \pi^{-1}(\infty) \rightarrow \mathbf{P}^1 \setminus \{\infty\}$. Note that the test configuration $(\mathcal{X}_\ell, \mathcal{L}_\ell)$ for affine ℓ is *product* in the sense that $\mathcal{X}_\ell \cong X_P \times \mathbf{P}^1$, T -equivariantly (see [16]). Conversely, for each T -equivariant test configuration $(\mathcal{X}, \mathcal{L})$, there exists a rational piecewise affine convex function f such that $(\mathcal{X}, \mathcal{L}) = (\mathcal{X}_f, \mathcal{L}_f)$ (cf. [15] and [26]). Moreover, the non-Archimedean K -energy can be written as

$$M^{NA}(\mathcal{X}_f, \mathcal{L}_f) = \frac{1}{\text{vol}(P)} \left(\int_{\partial P} f(x) d\sigma - n \int_P f(x) dx \right)$$

where σ denotes the $(n - 1)$ -dimensional Hausdorff measure on the boundary ∂P .

Let v be the extremal Kähler vector field of the corresponding toric Fano manifold $(X_P, -K_{X_P})$ with respect to the maximal algebraic torus T . Let θ_P be the potential function of the extremal Kähler vector field v [19], which is affine linear on P , and normalized by $\int_P \theta_P dx = 0$. For a convex function f , we define the functional

$$\mathcal{L}_P(f) = \int_{\partial P} f(x) d\sigma - \int_P (n + \theta_P) f(x) dx.$$

Then the potential function θ_P is uniquely determined by the $n + 1$ -equations

$$(2.2) \quad \mathcal{L}_P(1) = 0, \quad \mathcal{L}_P(x_i) = 0 \quad \text{for} \quad i = 1, \dots, n.$$

According to [35, Definition 1.1], $(X_P, -K_{X_P})$ is said to be *relatively K -stable* if $\mathcal{L}_P(f) \geq 0$ for any rational PL convex function f and the equality holds if and only if f is affine linear. In [33], Yao observed that the Mabuchi constant defined in (1.1) is given by

$$M_{X_P} = \max_P \theta_P$$

by the Atiyah-Guillemin-Sternberg convexity theorem ([3] and [21]).

Let $(\mathcal{X}_f, \mathcal{L}_f)$ be a T -equivariant test configuration arising from a rational PL convex function f on P . The *non-Archimedean relative Ding functional* of $(\mathcal{X}_f, \mathcal{L}_f)$ was computed in [33, Theorem 4.3] as

$$\begin{aligned} D_v^{NA}(f) &:= D_v^{NA}(\mathcal{X}_f, \mathcal{L}_f) = -f(0) + \frac{1}{\text{vol}(P)} \int_P f(x)(1 - \theta_P(x)) dx \\ &= \frac{1}{\text{vol}(P)} \int_P (f(x) - f(0))(1 - \theta_P(x)) dx. \end{aligned}$$

Observe that the linear function θ_P is uniquely determined by the conditions $D^{NA}(\ell) = 0$ for all affine ℓ . The *non-Archimedean reduced J -functional* of $(\mathcal{X}_f, \mathcal{L}_f)$, first introduced in [24], is expressed by

$$J_T^{NA}(f) := J_T^{NA}(\mathcal{X}_f, \mathcal{L}_f) = \inf_{\ell} \left(\frac{1}{\text{vol}(P)} \int_P (f(x) + \ell(x)) dx - \min_P (f + \ell) \right),$$

where ℓ runs over all the affine functions on \mathbf{R}^n . It is clear that the functional is invariant under the (torus) action $f \mapsto f + \ell$ for affine ℓ . We also know that $J_T^{NA}(f) = 0$ is equivalent to f being affine.

Using these quantities, we can define (uniform) relative Ding stability of toric Fano manifolds as follows.

Definition 2.1. *Let X be a toric Fano manifold and P the corresponding Fano polytope.*

- (1) *X is relatively Ding semistable if $D_v^{NA}(f) \geq 0$ for any rational PL convex function f .*
- (2) *X is relatively Ding stable if X is relatively Ding semistable, and $D_v^{NA}(f) = 0$ if and only if f is affine.*
- (3) *X is relatively Ding unstable if X is not relatively Ding semistable.*
- (4) *X is uniformly relatively Ding stable if there exists $\delta > 0$ such that $D_v^{NA}(f) \geq \delta J_T^{NA}(f)$ for any rational PL convex function f .*

Since the triviality of non-Archimedean reduced J -functional characterizes affine functions, uniform relative Ding stability is a priori stronger than relative Ding stability.

It was proved in [33] that (uniform) relative Ding stability can be rephrased as the condition on the Mabuchi constant as stated in Proposition 1.2. We shall give a proof of it for reader's convenience.

Proof of Proposition 1.2. (a) The equivalence between (1) and (2) is coming from the equivalence of non-Archimedean reduced J -functional and L^1 -norm on *normalized* convex functions (see [29, Proposition 5.4.1 (3)]). First we show (2) \Rightarrow (3). Let $\ell := (1 - \theta_P)/\text{vol}(P)$. Note that $D_v^{NA}(\ell) = 0$ implies

$$\ell(0) = \int_P \ell(x)^2 dx.$$

As a first step, we prove $\ell \geq 0$ on P by contradiction. Assume $\{\ell < 0\}$ is non-empty. Then

$$\int_{\{\ell < 0\}} \ell(x)^2 dx > 0.$$

On the other hand, for a rational PL convex function $\ell^+ := \max\{0, \ell\}$, we get

$$0 \leq D_v^{NA}(\ell^+) = -\ell(0) + \int_{\{\ell \geq 0\}} \ell(x)^2 dx = - \int_{\{\ell < 0\}} \ell(x)^2 dx < 0,$$

which yields a contradiction. Now assume $\min_P \ell = (1 - \max_P \theta_P) / \text{vol}(P) = 0$. Then for each $i \in \mathbf{N}$, $\{\ell < 1/i\}$ is non-empty and hence we can define

$$g_i(x) := \max\left\{0, \frac{1}{i} - \ell(x)\right\}, \quad f_i(x) := \frac{g_i(x)}{\int_P g_i dx}.$$

Each f_i is a non-negative PL convex function on P . Since $\ell(0) = 1/\text{vol}(P) > 0$, we have $1/i - \ell(0) < 0$ for sufficiently large i and hence $f_i(0) = 0$. This implies that f_i is normalized. Furthermore, the rationality of g_i shows that of f_i . Since $\{f_i > 0\} = \{\ell < 1/i\}$, we have

$$\delta = \delta \int_P f_i(x) dx \leq D_v^{NA}(f_i) = \int_{\{f_i > 0\}} f_i(x)\ell(x) dx \leq \frac{1}{i},$$

which yields a contradiction. Now we show (3) \Rightarrow (2). Pick a normalized rational PL convex function f . Then a simple computation gives

$$D_v^{NA}(f) = \frac{1}{\text{vol}(P)} \int_P f(x)(1 - \theta_P(x)) dx \geq \frac{1 - \max_P \theta_P}{\text{vol}(P)} \int_P f(x) dx.$$

(b) The proof of (a) shows (1) \Rightarrow (2) \Rightarrow (3). To prove (3) \Rightarrow (1), apply Jensen's inequality. \square

3. LISTS OF UNIFORM RELATIVE DING STABILITY OF TORIC FANO MANIFOLDS

3.1. How to determine relative Ding stability. As it was seen in Proposition 1.2, we can check uniform relative Ding stability by computing the Mabuchi constants. In this subsection, we illustrate how to compute the potential function θ_P and the Mabuchi constant M_{X_P} for a given Fano polytope P .

Let P be an n -dimensional Fano polytope in \mathbf{R}^n and v the extremal Kähler vector field of $(X_P, -K_{X_P})$. Suppose that θ_P is of the form

$$\theta_P(x) = \sum_{i=1}^n \theta_i \left(x_i - \frac{b_i}{b_0} \right),$$

with

$$b_0 := \text{vol}(P), \quad b_i := \int_P x_i dx \quad \text{for } i = 1, \dots, n.$$

The coefficients $\theta_1, \dots, \theta_n$ are uniquely determined by (2.2), that is, the n -equations

$$(3.1) \quad \sum_{j=1}^n (b_0 c_{ij} - b_i b_j) \theta_j + b_0 b_i = 0 \quad \text{for } i = 1, \dots, n,$$

where

$$c_{ij} := \int_P x_i x_j dx \quad \text{for } i, j = 1, \dots, n.$$

In [28, Table 1, 2, 3], Nakagawa listed the vector $(\theta_1, \dots, \theta_n)$ for all Fano polytopes in dimension up to four. Instead of applying (3.1), one can also use his list to compute θ_{X_P} and M_{X_P} from the polytope data. We used the software packages Normaliz [10] and Maxima¹ for our computations. See [35, Section 5.1] for more details.

Example 1. Let P be the reflexive polytope in \mathbf{R}^4 whose vertices are given by

$$\left\{ \begin{array}{l} (-1, -1, -1, -1), (-1, -1, -1, 1), (-1, -1, 0, -1), (-1, -1, 2, 1) \\ (-1, 1, -1, -1), (-1, 1, 0, -1), (-1, 3, -1, 1), (-1, 3, 2, 1) \\ (1, -1, -1, -1), (1, -1, 0, -1), (3, -1, -1, 1), (3, -1, 2, 1) \end{array} \right\}.$$

The dual polytope P^\vee of P is given by

$$P^\vee = \text{conv} \left(\left\{ \begin{array}{l} (1, 0, 0, 0), (0, 1, 0, 0), (0, 0, 1, 0), (0, 0, 0, 1) \\ (0, 0, 0, -1), (-1, -1, 0, 1), (0, 0, -1, 1) \end{array} \right\} \right),$$

and the associated toric Fano fourfold is $X_P = \mathbf{P}_{\mathbf{P}^1 \times \mathbf{P}^2}(\mathcal{O} \oplus \mathcal{O}(1, 1))$. Following the computation in (3.1), we find $(\theta_1, \theta_2, \theta_3, \theta_4) = (0, 0, 0, 2790/1973)$ which is consistent with [28, Table 3, D -6]. Since $b_0 = \text{vol}(P) = 62/3$ and $b_4 = \int_P x_4 dx = 36/5$, we obtain

$$\theta_P(x) = \frac{1}{1973}(2790x_4 - 972).$$

For the vertex $p = (3, -1, 2, 1)$ of P , one can see that

$$M_{X_P} = \max_P \theta_P = \theta_P(p) = \frac{1818}{1973} < 1,$$

which shows that $X_P = \mathbf{P}_{\mathbf{P}^1 \times \mathbf{P}^2}(\mathcal{O} \oplus \mathcal{O}(1, 1))$ is uniformly relatively Ding stable.

3.2. Toric del Pezzo surfaces. Although the result of the surface case was already obtained in [33, Example 16], we list it here for reader's convenience. In dimension two, there are five isomorphism classes of Fano polygons and all of them are uniformly relatively Ding stable. In Table 1, dP_{9-i} corresponds to a (smooth) del Pezzo surface with degree $(9 - i)$, that is, a smooth projective surface given by the blowing-up of \mathbf{P}^2 at i points.

TABLE 1. Relative Ding stability of smooth toric del Pezzo surfaces

No.	Notation	Relative Ding stability	M_X
1	\mathbf{P}^2	uniformly stable	0
2	$\mathbf{P}^1 \times \mathbf{P}^1$	uniformly stable	0
3	dP_8	uniformly stable	5/11
4	dP_7	uniformly stable	304/409
5	dP_6	uniformly stable	0

3.3. Toric Fano threefolds. In dimension three, in turn, there are 18 isomorphism classes of Fano polytopes. In this subsection, we verify relative Ding stability of all toric Fano threefolds and prove that there is an example of toric Fano threefolds which is relatively K -stable, but it is relatively Ding unstable.

¹Maxima is available from <http://maxima.sourceforge.net/>.

3.3.1. *Relative Ding stability of toric Fano threefolds.* Among all 18 classes of toric Fano threefolds, if a given toric Fano threefold X is one of $\mathcal{B}_1, \mathcal{C}_2, \mathcal{D}_1, \mathcal{D}_2, \mathcal{E}_1$ and \mathcal{E}_2 , then X is relatively Ding unstable. On the other hand, if X is one of $\mathbf{P}^3, \mathcal{B}_2, \mathcal{B}_3, \mathcal{B}_4, \mathcal{C}_1, \mathcal{C}_3, \mathcal{C}_4, \mathcal{C}_5, \mathcal{E}_3, \mathcal{E}_4, \mathcal{F}_1$ and \mathcal{F}_2 , then X is relatively Ding stable. Remark that these 12 relatively Ding stable ones are actually *uniformly* relatively Ding stable (see [33, Example 5.15]). We list all results in Table 2.

3.3.2. *Relative K -stability of toric Fano threefolds.* This subsection aims to show that the toric Fano threefold $X = \mathbf{P}_{\mathbf{P}^2}(\mathcal{O} \oplus \mathcal{O}(2))$, which is labelled \mathcal{B}_1 in Table 2, is relatively K -stable. Comparing this and the result in Table 2, we see the following (see, Corollary 1.6 (1)).

Proposition 3.1. *The toric Fano threefold \mathcal{B}_1 is relatively K -stable, but it is relatively Ding unstable.*

The rest of this subsection is devoted to prove Proposition 3.1 where we use the existence theorem of an extremal Kähler metric in each Kähler class of \mathcal{B}_1 (Proposition 1.5). We also clarify which part is a problematic in [35] when the third author apply the instability criterion (1.2) into \mathcal{B}_1 . This effects the “only if” part of Theorem 1.5 and Corollary 1.6 in [35], and provides inconclusive results for relative K -stability of $\mathcal{C}_2, \mathcal{D}_1, \mathcal{D}_2, \mathcal{E}_1$ and \mathcal{E}_2 .

Let P be the Fano polytope corresponding to \mathcal{B}_1 . Let us define $P^- = \{x \in P \mid 1 - \theta_P \leq 0\}$, where $\theta_P = -\frac{620}{349}x_3 - \frac{240}{349}$ is the potential function of P . It was computed in [35, Proposition 5.1 (b)] that the right-hand side value of (1.2) is given by

$$(3.2) \quad P^- = \text{conv} \left(\left\{ (4, -1, -1), \left(\frac{39}{10}, -1, -\frac{19}{20}\right), \left(-1, -1, -\frac{19}{20}\right), \right. \right. \\ \left. \left. \left(-1, \frac{39}{10}, -\frac{19}{20}\right), (-1, -1, -1), (-1, 4, -1) \right\} \right), \quad \text{vol}(P^-) = \frac{7351}{12000}$$

and

$$(3.3) \quad \int_{P^-} (1 - \theta_P)^2 dx = \frac{1475918766336271}{1461612000}$$

respectively. Although the values of (3.2) are correct, the integral value of (3.3) is incorrect because the third author forgot to put the normalize condition into the input data of P^- when he run the software [10] at the time. More precisely, the first and third authors recomputed the value of $\int_{P^-} (1 - \theta_P)^2 dx$ and confirmed that the correct value is

$$\int_{P^-} (1 - \theta_P)^2 dx = \frac{23785711}{14616120000}.$$

Since the left-hand side value of (1.2) is $1 - c = \frac{589}{349}$, we see that the toric Fano threefold \mathcal{B}_1 is inconclusive.

However, if we put emphasis on the geometric description of \mathcal{B}_1 as a projective bundle $\mathbf{P}_{\mathbf{P}^2}(\mathcal{O} \oplus \mathcal{O}(2))$, we may conclude that it is relatively K -stable as follows (cf. [33, Section 5.2.4]).

In fact, since the toric Fano manifold \mathcal{B}_1 can be expressed as a \mathbf{P}^1 -bundle over the base space \mathbf{P}^2 , we may apply “Calabi ansatz” for constructing extremal Kähler metrics which were accomplished by many researchers [20, 25, 1]. Consequently, \mathcal{B}_1 admits an extremal Kähler metric in every Kähler class (see Proposition 1.5 (1)). Hence the assertion is verified by [36].

Remark 3.2. The first and third authors also confirmed that other five classes of toric Fano threefolds $\mathcal{C}_2, \mathcal{D}_1, \mathcal{D}_2, \mathcal{E}_1$ and \mathcal{E}_2 do *not* satisfy the instability condition (1.2) by

TABLE 2. Relative Ding stability of toric Fano threefolds

No.	Notation	Relative Ding stability	M_X
1	\mathbf{P}^3	uniformly stable	0
2	\mathcal{B}_1	unstable	380/349
3	\mathcal{B}_2	uniformly stable	55/97
4	\mathcal{B}_3	uniformly stable	35/43
5	\mathcal{B}_4	uniformly stable	0
6	\mathcal{C}_1	uniformly stable	60/73
7	\mathcal{C}_2	unstable	38/33
8	\mathcal{C}_3	uniformly stable	0
9	\mathcal{C}_4	uniformly stable	5/11
10	\mathcal{C}_5	uniformly stable	0
11	\mathcal{D}_1	unstable	711861/467581
12	\mathcal{D}_2	unstable	694595/650251
13	\mathcal{E}_1	unstable	27195/19651
14	\mathcal{E}_2	unstable	2936215/2735927
15	\mathcal{E}_3	uniformly stable	304/409
16	\mathcal{E}_4	uniformly stable	185791/394975
17	\mathcal{F}_1	uniformly stable	0
18	\mathcal{F}_2	uniformly stable	31/67

direct computations as above. Thus the relative K -stability of these five classes are still inconclusive.

3.4. Relative Ding stability and relative K -stability of toric Fano fourfolds. Now we verify relative Ding stability of toric Fano fourfolds. Among all 124 isomorphism classes of toric Fano fourfolds, we see that there are 49 classes of toric Fano fourfolds which are uniformly relatively Ding stable, while the remaining 75 classes are relatively Ding unstable.

In dimension four, there are at least three examples of toric Fano manifolds which are relatively K -stable, but they are relatively Ding unstable. The geometric description of these manifolds are $X = \mathbf{P}_{\mathbf{P}^3}(\mathcal{O} \oplus \mathcal{O}(k))$ which are labeled B_1 and B_2 for $k = 3$ and 2 , respectively. The proof is similar to it of Proposition 3.1, and details are left it to the reader.

3.5. Bott structure of certain toric Fano manifolds. In this final section, we prove Corollary 1.9. We firstly recall toric description of Bott manifolds. For further details, see [2, 12, 13, 23].

For a given $n \in \mathbf{N}$ with $n \geq 2$, we inductively construct complex manifolds X_k for $k = 1, 2, \dots, n$ as follows. Let X_0 be a point $\text{Spec } \mathbf{C}$, and let X_1 be the projective line. For $k \geq 2$, we assume that X_{k-1} is already determined and choose a holomorphic line bundle \mathcal{L}_k on X_{k-1} . Then we define X_k as the total space of the \mathbf{P}^1 -bundle over X_{k-1} and denote it by

$$X_k := \mathbf{P}_{X_{k-1}}(\mathcal{O}_{X_{k-1}} \oplus \mathcal{L}_k) \xrightarrow{\pi_k} X_{k-1}.$$

Through this construction, we call X_k the *stage k Bott manifold of the Bott tower of height n* :

$$X_n \xrightarrow{\pi_n} X_{n-1} \xrightarrow{\pi_{n-1}} \dots \xrightarrow{\pi_2} X_1 \xrightarrow{\pi_1} \text{Spec } \mathbf{C}.$$

On the other hand, a stage n Bott manifold X_n can be written as a quotient of $(\mathbf{C}_*^2)^n$ by an algebraic torus $(\mathbf{C}^*)^n$, where $(\mathbf{C}_*^2)^n$ is n copies of $\mathbf{C}_*^2 := \mathbf{C}^2 \setminus \{0\}$. In order to see this, we denote an algebraic torus $(\mathbf{C}^*)^n$ by T . Let A be a lower triangular unipotent matrix of the form

$$(3.4) \quad A = \begin{pmatrix} 1 & 0 & \cdots & 0 & 0 \\ a_2^1 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ a_{n-1}^1 & a_{n-1}^2 & \cdots & 1 & 0 \\ a_n^1 & a_n^2 & \cdots & a_n^{n-1} & 1 \end{pmatrix}$$

with $a_j^i \in \mathbf{Z}$. Let us consider the induced action λ of $\mathbf{t} = (t_1, \dots, t_n) \in T$ on $(\mathbf{z}; \mathbf{w}) = ((z_1, \dots, z_n); (w_1, \dots, w_n)) \in (\mathbf{C}_*^2)^n$ which is given by

$$(3.5) \quad \begin{array}{ccc} \lambda : T \times (\mathbf{C}_*^2)^n & \longrightarrow & (\mathbf{C}_*^2)^n \\ \downarrow & & \downarrow \\ (\mathbf{t}, (\mathbf{z}; \mathbf{w})) & \longmapsto & \left((t_1 z_1, \dots, t_n z_n); \left(\left(\prod_{i=1}^n t_i^{a_i^1} \right) w_1, \dots, \left(\prod_{i=1}^n t_i^{a_i^n} \right) w_n \right) \right). \end{array}$$

For the unit sphere S^3 in \mathbf{C}_*^2 , there is the induced action of $(\mathbf{R}_{>0})^n$ on $(\mathbf{C}_*^2)^n$ by (3.5), which is transverse to $(S^3)^n$. Hence the orbits of this action are bijectively correspond to orbits of the induced free $(S^1)^n$ -action on $(S^3)^n$. Consequently, the geometric quotient $(\mathbf{C}_*^2)^n/T$ is a complex n -dimensional compact manifold which will be denoted by $X_n(A)$.

As a special case, for $\mathbf{k} = (k_1, \dots, k_{n-1}) \in \mathbf{Z}^{n-1}$, we consider the case where $a_n^i = k_i$ and $a_j^i = 0$ otherwise. Then a lower triangular matrix in (3.4) is written as

$$A = \begin{pmatrix} 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \\ k_1 & k_2 & \cdots & k_{n-1} & 1 \end{pmatrix}.$$

The associated Bott tower $X_n(A)$ is the sequence of \mathbf{P}^1 -bundles

$$(3.6) \quad \mathbf{P}_{(\mathbf{P}^1)^{n-1}}(\mathcal{O} \oplus \mathcal{O}(k_1, \dots, k_{n-1})) \xrightarrow{\pi_n} (\mathbf{P}^1)^{n-1} \xrightarrow{\pi_{n-1}} \cdots \xrightarrow{\pi_2} \mathbf{P}^1 \xrightarrow{\pi_1} \text{Spec } \mathbf{C}$$

which has twist one unless $\mathbf{k} = (0, \dots, 0)$. Since any stage n Bott manifold in (3.6) with $\prod_{i=1}^{n-1} k_i \neq 0$ admits an extremal Kähler metric in each Kähler class by Proposition 1.7, our problem can be reduced to find examples of relatively Ding unstable Fano Bott manifolds in the form of (3.6). In particular, they must be twist one. There are six candidates of such examples in Tables 2–3, that is, \mathcal{C}_2 , L_1 , L_2 , L_3 , L_4 , and L_6 . Their twists are computable by primitive relations of the vertices of P^\vee (see [5, Proposition 3.3.6]). Alternatively, one can use the *signed rooted forest* associated with a Fano Bott manifold for computing twist (see [13, Section 2]). In fact, the height of signed rooted forest is equal to the twist of a Fano Bott manifold, and all signed rooted forests of stage four Fano Bott manifolds are illustrated in [13, Figure 2]. Consequently, we have the following list of twist for each relatively Ding unstable Fano manifold.

Notation	\mathcal{C}_2	L_1	L_2	L_3	L_4	L_6
twist	2	1	2	2	3	2
Number in [13, Figure 2]	–	(12)	(10)	(8)	(7)	(3)

The assertion is verified.

Remark 3.3. Note that there exists *some* Kähler classes of the toric Fano threefold \mathcal{C}_2 that admit extremal Kähler metrics by Theorem 1 (2) in [2]. However, it is difficult to determine *which* Kähler class may admit extremal Kähler metrics in general. For the moment, we do not know if every Kähler class of \mathcal{C}_2 has an extremal Kähler metric, or even if its first Chern class has an extremal Kähler metric.

Hence the following problem is still open.

Problem 3.4. *Does the toric Fano threefold \mathcal{C}_2 admit an extremal Kähler metric in its first Chern class? Moreover, is it a Calabi Dream manifold?*

If the answer is affirmative, then any stage three Fano Bott manifold is a Calabi dream manifold.

TABLE 3. Relative Ding stability of toric Fano fourfolds

No.	Notation	Relative Ding stability	M_X
1	P^4	uniformly stable	0
2	B_1	unstable	186633/108133
3	B_2	unstable	1827/1577
4	B_3	uniformly stable	62937/101357
5	B_4	uniformly stable	0
6	B_5	unstable	81/71
7	C_1	unstable	5168/2803
8	C_2	unstable	5056/4961
9	C_3	uniformly stable	1876/2921
10	C_4	uniformly stable	0
11	E_1	unstable	3117506664/1528096589
12	E_2	unstable	1840195984/1038057499
13	E_3	unstable	8725985172/6541476757
14	D_1	unstable	11387/8057
15	D_2	unstable	2344247/1129107
16	D_3	unstable	23933079/13920179
17	D_4	unstable	10601177/5141177
18	D_5	unstable	380/349
19	D_6	uniformly stable	1818/1973
20	D_7	unstable	748/523
21	D_8	unstable	3165248067/2130145727
22	D_9	unstable	1380/1177
23	D_{10}	unstable	18353/11683
24	D_{11}	unstable	1727/1007
25	D_{12}	uniformly stable	55/97
26	D_{13}	uniformly stable	0
27	D_{14}	uniformly stable	35/43
28	D_{15}	uniformly stable	5/11
29	D_{16}	uniformly stable	1046933/1061921
30	D_{17}	uniformly stable	59/134
31	D_{18}	uniformly stable	3927/5177
32	D_{19}	uniformly stable	238/1613
33	G_1	unstable	29663642268/16641653953
34	G_2	unstable	372321328/163900203
35	G_3	unstable	2130146436/1907713037
36	G_4	unstable	91293727706236/48057952407691
37	G_5	unstable	7423814976/5976434111

No.	Notation	Relative Ding stability	M_X
38	G_6	unstable	6431616/4388521
39	H_1	unstable	391552/172227
40	H_2	unstable	23712381468/12261327193
41	H_3	unstable	1551503556/999563887
42	H_4	unstable	173016/111181
43	H_5	unstable	9770225016/8410595791
44	H_6	unstable	7892183028/6776874793
45	H_7	unstable	2510023476/2191715291
46	H_8	uniformly stable	304/409
47	H_9	uniformly stable	3339308412/5523943807
48	H_{10}	uniformly stable	33344/116609
49	L_1	unstable	1827/1577
50	L_2	unstable	4491311/2637399
51	L_3	unstable	29551791/20356871
52	L_4	unstable	12523717673/6291985913
53	L_5	uniformly stable	60/73
54	L_6	unstable	38/33
55	L_7	uniformly stable	10/11
56	L_8	uniformly stable	0
57	L_9	uniformly stable	5/11
58	L_{10}	uniformly stable	11999/15439
59	L_{11}	uniformly stable	0
60	L_{12}	uniformly stable	98007/118907
61	L_{13}	uniformly stable	93/203
62	I_1	unstable	708660642/298795537
63	I_2	unstable	3024970758/1752843533
64	I_3	unstable	1265748565401/557417025511
65	I_4	unstable	94549671/42931741
66	I_5	unstable	11713596999245802/6984760752795427
67	I_6	unstable	448243263/230998253
68	I_7	unstable	711861/467581
69	I_8	unstable	3126600189529/1577014856729
70	I_9	unstable	51542252/31553027
71	I_{10}	unstable	1376202783/1008086248
72	I_{11}	unstable	13278385/7524593
73	I_{12}	uniformly stable	3762595665/10127095471
74	I_{13}	unstable	694595/650251
75	I_{14}	uniformly stable	211431316083/218756440813
76	I_{15}	unstable	949610187/739116617
77	M_1	unstable	4575996/2853121
78	M_2	unstable	9794436236/4944865931
79	M_3	unstable	25645181180008/16440690503203
80	M_4	unstable	19189680326248/13060549547443

No.	Notation	Relative Ding stability	M_X
81	M_5	unstable	12265145105056/5748810067571
82	J_1	unstable	326672481/271746296
83	J_2	uniformly stable	141120/450983
84	Q_1	unstable	510608/266403
85	Q_2	unstable	518137667/237174992
86	Q_3	unstable	90760711/67077371
87	Q_4	unstable	121849781/73313456
88	Q_5	unstable	977249482633/573603226348
89	Q_6	unstable	27195/19651
90	Q_7	unstable	138419/99094
91	Q_8	unstable	2936215/2735927
92	Q_9	uniformly stable	173974704/197727059
93	Q_{10}	unstable	5389/4499
94	Q_{11}	uniformly stable	304/409
95	Q_{12}	unstable	203968842/196251577
96	Q_{13}	uniformly stable	31595933/35875483
97	Q_{14}	unstable	3046408136973/2450627729348
98	Q_{15}	uniformly stable	185791/394975
99	Q_{16}	uniformly stable	80506889/111477439
100	Q_{17}	uniformly stable	17172/162187
101	K_1	unstable	18300/16571
102	K_2	unstable	400/341
103	K_3	uniformly stable	20550/35413
104	K_4	uniformly stable	0
105	R_1	unstable	567397500/324135871
106	R_2	unstable	2758057258564/2028544442569
107	R_3	uniformly stable	277721596/369479321
108		unstable	182110284/153124261
109	U_1	uniformly stable	2850/3389
110	U_2	unstable	1095285/924382
111	U_3	uniformly stable	1275/1424
112	U_4	uniformly stable	5/11
113	U_5	uniformly stable	0
114	U_6	uniformly stable	31/67
115	U_7	uniformly stable	119/258
116	U_8	uniformly stable	0
117	\tilde{V}^4	unstable	737568/395263
118	V^4	uniformly stable	0
119	$dP_7 \times dP_7$	unstable	608/409
120	$dP_7 \times dP_6$	uniformly stable	304/409
121	$dP_6 \times dP_6$	uniformly stable	0
122	Z_1	uniformly stable	1013328/1289443
123	Z_2	unstable	3418840588/2592694163
124	W	uniformly stable	0

REFERENCES

- [1] V. Apostolov, D.M.J. Calderbank, P. Gauduchon, and C.W. Tønnesen-Friedman, Hamiltonian 2-forms in Kähler geometry. III. Extremal metrics and stability. *Invent. Math.* **173** (2008), 547–601.
- [2] C.P. Boyer, D.M.J. Calderbank, and C.W. Tønnesen-Friedman, The Kähler geometry of Bott manifolds. *Adv. Math.* **350** (2019), 1–62.
- [3] M. F. Atiyah, Convexity and commuting hamiltonians. *Bull. London Math. Soc.* **14** (1982), 1–15.
- [4] V. V. Batyrev, Troidal Fano 3-folds. *Izv. Akad. Nauk SSSR.* **45** (1981), 704–717.
- [5] V. V. Batyrev, On the classification of toric Fano 4-folds. *J. Math. Sci.* **94** (1999), 1021–1050.
- [6] V. V. Batyrev and E. Selivanova, Einstein-Kähler metrics on symmetric toric Fano manifolds. *J. Reine Angew. Math.* **512** (1999), 225–236.
- [7] R. J. Berman, S. Boucksom, and M. Jonsson, A variational approach to the Yau-Tian-Donaldson conjecture. *J. Amer. Math.* **34** (2021) 605–652.
- [8] R. J. Berman, K -polystability of \mathbf{Q} -Fano varieties admitting Kähler-Einstein metrics. *Invent. Math.* **203** (2016), 973–1025.
- [9] S. Boucksom, T. Hisamoto, and M. Jonsson, Uniform K -stability, Duistermaat-Heckman measures and singularity of pairs. *Ann. Inst. Fourier (Grenoble)* **67** (2017), 743–841.
- [10] W. Bruns, B. Ichim, T. Römer and C. Söger, Normaliz. Algorithms for rational cones and affine monoids. Available from <http://www.math.uos.de/normaliz>.
- [11] X. Chen and J. Cheng, On the constant scalar curvature Kähler metrics (II) – Existence results. arXiv:1801.00656
- [12] Y. Cho, E. Lee, M. Masuda, and S. Park, Unique toric structure on a Fano Bott manifold. arXiv:2005.02740
- [13] Y. Cho, E. Lee, M. Masuda, and S. Park, On the enumeration of Fano Bott manifolds. arXiv:2106.12788
- [14] D. A. Cox, J. B. Little, and H. K. Schenck, *Toric varieties*. Graduate Studies in Mathematics **124**. American Mathematical Society, Providence, RI, (2011). xxiv+841 pp.
- [15] T. Delcroix, Uniform K -stability of polarized spherical varieties. arXiv:2009.06463v2.
- [16] S. K. Donaldson, Scalar curvature and stability of toric varieties. *J. Differential Geom.* **62** (2002), 289–349.
- [17] S. K. Donaldson, The Ding Functional, Berndtsson Convexity and Moment Maps. in *Geometry, Analysis and Probability, Progress in Math.* **310** (2017), 57–67.
- [18] K. Fujita, Optimal bounds for the volumes of Kähler-Einstein Fano manifolds. *Amer. J. Math.* **140** (2018), 391–414.
- [19] A. Futaki and T. Mabuchi, Bilinear forms and extremal Kähler vector fields associated with Kähler classes. *Math. Ann.* **301** (1995), 199–210.
- [20] D. Guan, Existence of extremal metrics on compact almost homogeneous Kähler manifolds with two ends. *Trans. Amer. Math. Soc.* **347** (1995), 2255–2262.
- [21] V. Guillemin and S. Sternberg, Convexity properties of the moment mapping I and II. *Invent. Math.* **67** (1982), 491–513; *Invent. Math.* **77** (1984), 533–546.
- [22] J. Han and C. Li, On the Yau-Tian-Donaldson conjecture for generalized Kähler-Ricci soliton equations. To appear in *Commun. Pur. Appl. Math.* arXiv:2006.00903
- [23] A. Higashitani, K. Kurimoto, and M. Masuda, Cohomological rigidity for toric Fano manifolds of small dimensions or large Picard numbers. arXiv:2005.13795
- [24] T. Hisamoto, Stability and coercivity for toric polarizations. arXiv:1610.07998v2.
- [25] A.D. Hwang, On existence of Kähler metrics with constant scalar curvature. *Osaka J. Math.* **31** (1994), 561–595.
- [26] Y. Li and Z. Li, Equivariant \mathbb{R} -test configurations of polarized spherical varieties. arXiv:2206.04880v3.
- [27] T. Mabuchi, Kähler-Einstein metrics for manifolds with nonvanishing Futaki character. *Tohoku Math. J.* **53** (2001), 171–182.
- [28] Y. Nakagawa, Combinatorial formulae for Futaki characters and generalized killing forms of toric Fano orbifolds. *The Third Pacific Rim Geometry Conference (Seoul, 1996)*, 223–260, Monogr. Geom. Topology **25**, Int. Press, Cambridge, MA, 1998.
- [29] Y. Nitta and S. Saito, A uniform version of the Yau-Tian-Donaldson correspondence for extremal Kähler metrics on polarized toric manifolds. arXiv:2110.10386.
- [30] H. Sato, Toward the classification of higher-dimensional toric Fano varieties. *Tohoku Math. J.* **52** (2000) 383–413.

- [31] G. Székelyhidi, Extremal metrics and K-stability. *Bull. Lond. Math. Soc.* **39** (2007), 76–84.
- [32] K. Watanabe and M. Watanabe, The classification of Fano 3-folds with torus embeddings. *Tokyo. J. Math.* **5** (1982), 37–48.
- [33] Y. Yao, Mabuchi solitons and relative Ding stability of toric Fano varieties. *Int. Math. Res. Not.*, published electronically 22 September 2021, <https://doi.org/10.1093/imrn/rnab226>
- [34] Y. Yao, Relative Ding stability and an obstruction to the existence of Mabuchi solitons. *J. Geom. Anal.* **32** (2022), Article No. 105, 51 pp.
- [35] N. Yotsutani and B. Zhou, Relative algebro-geometric stabilities of toric manifolds. *Tohoku Math. J.* **71** (2019), 495–524.
- [36] B. Zhou and X. Zhu, K -stability on toric manifolds. *Proc. Amer. Math. Soc.* **136** (2008), 3301–3307.

APPENDIX A. COMPUTATIONAL DATA

A.1. 2-dimensional Fano polytopes.

A.1.1. \mathbf{P}^2 .

$$\begin{aligned} b_0 &= \text{vol}(P) = \frac{9}{2}, & b_1 &= \int_P x_1 dx_1 dx_2 = 0, \\ b_2 &= \int_P x_2 dx_1 dx_2 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 = \frac{9}{4}, \\ c_{12} &= \int_P x_1 x_2 dx_1 dx_2 = -\frac{9}{8}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 = \frac{9}{4}, \end{aligned}$$

$$\begin{aligned} \theta_P(x_1, x_2) &= 0, \\ M_X &= \max_P \theta_P = 0 < 1. \end{aligned}$$

A.1.2. $\mathbf{P}^1 \times \mathbf{P}^1$.

$$\begin{aligned} b_0 &= \text{vol}(P) = 4, & b_1 &= \int_P x_1 dx_1 dx_2 = 0, \\ b_2 &= \int_P x_2 dx_1 dx_2 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 = \frac{4}{3}, \\ c_{12} &= \int_P x_1 x_2 dx_1 dx_2 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 = \frac{4}{3}, \end{aligned}$$

$$\begin{aligned} \theta_P(x_1, x_2) &= 0, \\ M_X &= \max_P \theta_P = 0 < 1. \end{aligned}$$

A.1.3. dP_8 .

$$\begin{aligned} b_0 &= \text{vol}(P) = 4, & b_1 &= \int_P x_1 dx_1 dx_2 = -\frac{2}{3}, \\ b_2 &= \int_P x_2 dx_1 dx_2 = -\frac{1}{3}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 = \frac{4}{3}, \\ c_{12} &= \int_P x_1 x_2 dx_1 dx_2 = \frac{2}{3}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 = 2, \end{aligned}$$

$$\begin{aligned} \theta_P(x_1, x_2) &= -\frac{6x_1 + 1}{11}, \\ M_X &= \max_P \theta_P = \frac{5}{11} < 1. \end{aligned}$$

A.1.4. dP_7 .

$$\begin{aligned}
b_0 = \text{vol}(P) &= \frac{7}{2}, & b_1 &= \int_P x_1 dx_1 dx_2 = -\frac{1}{3}, \\
b_2 &= \int_P x_2 dx_1 dx_2 = \frac{1}{3}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 = \frac{13}{12}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 = \frac{5}{24}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 = \frac{13}{12},
\end{aligned}$$

$$\theta_P(x_1, x_2) = \frac{-168x_1 + 168x_2 - 32}{409},$$

$$M_X = \max_P \theta_P = \frac{304}{409} < 1.$$

A.1.5. dP_6 .

$$\begin{aligned}
b_0 = \text{vol}(P) &= 3, & b_1 &= \int_P x_1 dx_1 dx_2 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 = \frac{5}{6}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 = -\frac{5}{12}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 = \frac{5}{6},
\end{aligned}$$

$$\theta_P(x_1, x_2) = 0,$$

$$M_X = \max_P \theta_P = 0 < 1.$$

A.2. **3-dimensional Fano polytopes.**A.2.1. \mathbf{P}^3 .

$$\begin{aligned}
b_0 = \text{vol}(P) &= \frac{32}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = 0, \\
c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = \frac{32}{5}, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = -\frac{32}{15}, \\
c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = -\frac{32}{15}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{32}{5}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = -\frac{32}{15}, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = \frac{32}{5},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = 0,$$

$$M_X = \max_P \theta_P = 0 < 1.$$

A.2.2. \mathcal{B}_1 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{31}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = \frac{8}{3}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = \frac{8}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = -4, \\
c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = \frac{311}{30}, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = -\frac{53}{20}, \\
c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = -\frac{38}{15}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{311}{30}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = -\frac{38}{15}, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = \frac{19}{5},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = -\frac{620x_3 + 240}{349},$$

$$M_X = \max_P \theta_P = \frac{380}{349} > 1.$$

A.2.3. \mathcal{B}_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{28}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = \frac{2}{3}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = \frac{2}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = -2, \\
c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = \frac{88}{15}, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = -\frac{12}{5}, \\
c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = -\frac{16}{15}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{88}{15}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = -\frac{16}{15}, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = \frac{16}{5},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = -\frac{70x_3 + 15}{97},$$

$$M_X = \max_P \theta_P = \frac{55}{97} < 1.$$

A.2.4. \mathcal{B}_3 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 9, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = -\frac{9}{8}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = -\frac{9}{8}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = \frac{9}{8}, \\
c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = \frac{81}{20}, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = -\frac{27}{20}, \\
c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = -\frac{27}{20}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{81}{20}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = -\frac{27}{20}, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = \frac{111}{20},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = -\frac{20x_1 + 20x_2 + 5}{43},$$

$$M_X = \max_P \theta_P = \frac{35}{43} < 1.$$

A.2.5. \mathcal{B}_4 .

$$\begin{aligned} b_0 &= \text{vol}(P) = 9, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = 0, \\ b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = 0, \\ c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = \frac{9}{2}, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = -\frac{9}{4}, \\ c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{9}{2}, \\ c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = 0, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = 3, \end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = 0,$$

$$M_X = \max_P \theta_P = 0 < 1.$$

A.2.6. \mathcal{C}_1 .

$$\begin{aligned} b_0 &= \text{vol}(P) = \frac{26}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = \frac{4}{3}, \\ b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = \frac{4}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = -\frac{8}{3}, \\ c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = \frac{24}{5}, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = \frac{23}{30}, \\ c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = -\frac{23}{15}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{24}{5}, \\ c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = -\frac{23}{15}, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = \frac{46}{15}, \end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = -\frac{260x_3 + 80}{219},$$

$$M_X = \max_P \theta_P = \frac{60}{73} < 1.$$

A.2.7. \mathcal{C}_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{25}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = -\frac{2}{3}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = \frac{4}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = -2, \\
c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = \frac{37}{10}, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = -\frac{89}{60}, \\
c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = -\frac{11}{15}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{161}{30}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = -\frac{16}{15}, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = \frac{43}{15},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = -\frac{7600x_1 + 17750x_3 + 4868}{17787},$$

$$M_X = \max_P \theta_P = \frac{38}{33} > 1.$$

A.2.8. \mathcal{C}_3 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 8, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = 0, \\
c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = \frac{8}{3}, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = 0, \\
c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{8}{3}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = 0, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = \frac{8}{3},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = 0,$$

$$M_X = \max_P \theta_P = 0 < 1.$$

A.2.9. \mathcal{C}_4 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 8, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = \frac{2}{3}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = -\frac{4}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = 0, \\
c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = 4, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = -\frac{4}{3}, \\
c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{8}{3}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = 0, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = \frac{8}{3},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = -\frac{6x_2 + 1}{11},$$

$$M_X = \max_P \theta_P = \frac{5}{11} < 1.$$

A.2.10. \mathcal{C}_5 .

$$\begin{aligned} b_0 = \text{vol}(P) &= \frac{22}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = 0, \\ b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = 0, \\ c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = \frac{16}{5}, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = -\frac{17}{30}, \\ c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = -\frac{17}{15}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{16}{5}, \\ c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = \frac{17}{15}, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = \frac{34}{15}, \end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = 0,$$

$$M_X = \max_P \theta_P = 0 < 1.$$

A.2.11. \mathcal{D}_1 .

$$\begin{aligned} b_0 = \text{vol}(P) &= \frac{25}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = \frac{11}{8}, \\ b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = -\frac{21}{8}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = \frac{5}{8}, \\ c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = \frac{319}{60}, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = -\frac{13}{20}, \\ c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = -\frac{7}{3}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{11}{4}, \\ c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = -\frac{21}{20}, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = \frac{109}{20}, \end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = \frac{99600x_1 - 627000x_2 - 213939}{467581},$$

$$M_X = \max_P \theta_P = \frac{711861}{467581} > 1.$$

A.2.12. \mathcal{D}_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{23}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = \frac{11}{12}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = -\frac{7}{8}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = -\frac{11}{24}, \\
c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = \frac{19}{5}, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = \frac{17}{30}, \\
c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = -\frac{19}{10}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{47}{20}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = -\frac{17}{60}, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = \frac{211}{60},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = \frac{219420x_1 - 318320x_2 - 62565}{650251},$$

$$M_X = \max_P \theta_P = \frac{694595}{650251} > 1.$$

A.2.13. \mathcal{E}_1 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{23}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = -\frac{37}{24}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = -\frac{37}{24}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = \frac{37}{24}, \\
c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = \frac{151}{60}, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = -\frac{7}{60}, \\
c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = -\frac{6}{5}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{151}{60}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = -\frac{6}{5}, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = \frac{311}{60},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = -\frac{17020x_1 + 17020x_2 + 6845}{19651},$$

$$M_X = \max_P \theta_P = \frac{27195}{19651} > 1.$$

A.2.14. \mathcal{E}_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{22}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = -\frac{7}{4}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = -\frac{11}{24}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = \frac{7}{8}, \\
c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = \frac{71}{30}, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = -\frac{4}{15}, \\
c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = -\frac{71}{60}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{139}{60}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = \frac{2}{15}, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = \frac{229}{60},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = -\frac{2646160x_1 + 982960x_2 + 692905}{2735927},$$

$$M_X = \max_P \theta_P = \frac{2936215}{2735927} > 1.$$

A.2.15. \mathcal{E}_3 .

$$\begin{aligned} b_0 = \text{vol}(P) &= 7, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = -\frac{2}{3}, \\ b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = -\frac{2}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = 0, \\ c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = \frac{13}{6}, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = -\frac{5}{12}, \\ c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{13}{6}, \\ c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = 0, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = 7/3 \end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = -\frac{168x_1 + 168x_2 + 32}{409},$$

$$M_X = \max_P \theta_P = \frac{304}{409} < 1.$$

A.2.16. \mathcal{E}_4 .

$$\begin{aligned} b_0 = \text{vol}(P) &= \frac{20}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = -\frac{7}{8}, \\ b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = \frac{5}{12}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = \frac{5}{24}, \\ c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = \frac{121}{60}, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = -\frac{17}{30}, \\ c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = -\frac{17}{60}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{59}{30}, \\ c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = \frac{59}{60}, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = \frac{181}{60}, \end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = -\frac{171040x_1 - 39680x_2 + 24929}{394975},$$

$$M_X = \max_P \theta_P = \frac{185791}{394975} < 1.$$

A.2.17. \mathcal{F}_1 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 6, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = 0, \\
c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = \frac{5}{3}, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = \frac{5}{6}, \\
c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{5}{3}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = 0, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = 2,
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = 0,$$

$$M_X = \max_P \theta_P = 0 < 1.$$

A.2.18. \mathcal{F}_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 6, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 = \frac{5}{12}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 = \frac{5}{6}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 = \frac{5}{12}, \\
c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 = \frac{5}{3}, & c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 = \frac{5}{6}, \\
c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 = \frac{5}{12}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 = \frac{5}{3}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 = \frac{5}{6}, & c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 = \frac{17}{6},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3) = \frac{36x_2 - 5}{67},$$

$$M_X = \max_P \theta_P = \frac{31}{67} < 1.$$

A.3. 4-dimensional Fano polytopes.

A.3.1. \mathbf{P}^4 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{625}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{625}{36}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{625}{144}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{625}{144}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{625}{144}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{625}{36}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{625}{144}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{625}{144}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{625}{36}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{625}{144}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{625}{36},
\end{aligned}$$

$$\begin{aligned}
\theta_P(x_1, x_2, x_3, x_4) &= 0, \\
M_X &= \max_P \theta_P = 0 < 1.
\end{aligned}$$

A.3.2. B_1 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{100}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{89}{20}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{89}{20}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{89}{20}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{89}{20}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{3757}{90}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{229}{18}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{229}{18}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{229}{18}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{3757}{90}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{229}{18}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{229}{18}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{3757}{90}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{229}{18}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{3757}{90},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{400500x_1 + 400500x_2 + 400500x_3 + 400500x_4 - 213867}{108133},$$

$$M_X = \max_P \theta_P = \frac{186633}{108133} > 1.$$

A.3.3. B_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{80}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{28}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{28}{5}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{1208}{45}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{32}{3}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{728}{45}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{728}{45}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1208}{45}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{728}{45}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{728}{45}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1456}{45}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{1456}{45},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{3150x_1 + 3150x_2 + 3150x_3 + 3150x_4 - 1323}{1577},$$

$$M_X = \max_P \theta_P = \frac{1827}{1577} > 1.$$

A.3.4. B_3 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{68}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{27}{20}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{27}{20}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{27}{20}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{27}{20}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{1441}{90}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{421}{90}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{421}{90}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{421}{90}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1441}{90}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{421}{90}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{421}{90}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1441}{90}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{421}{90}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{1441}{90},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{82620x_1 + 82620x_2 + 82620x_3 + 82620x_4 - 19683}{101357},$$

$$M_X = \max_P \theta_P = \frac{62937}{101357} < 1.$$

A.3.5. B_4 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{64}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{64}{5}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{64}{15}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{64}{15}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{64}{5}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{64}{15}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{64}{5}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{64}{9},
\end{aligned}$$

$$\begin{aligned}
\theta_P(x_1, x_2, x_3, x_4) &= 0, \\
M_X &= \max_P \theta_P = 0 < 1.
\end{aligned}$$

A.3.6. B_5 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{64}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = -\frac{32}{15}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{32}{15}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{16}{5}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{16}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{512}{45}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{128}{45}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{128}{45}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{128}{45}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{512}{45}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{128}{45}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{128}{45}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{224}{15}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{32}{5}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{224}{15},
\end{aligned}$$

$$\begin{aligned}
\theta_P(x_1, x_2, x_3, x_4) &= \frac{30x_3 + 30x_4 - 9}{71}, \\
M_X &= \max_P \theta_P = \frac{81}{71} > 1.
\end{aligned}$$

A.3.7. C_1 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{99}{4}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{153}{10}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{153}{20}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{51}{5}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{51}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{837}{40}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{837}{80}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{279}{20}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{279}{20}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{81}{8}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{279}{40}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{279}{40}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1461}{40}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{69}{16}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{1461}{40},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{3740x_1 - 2312}{2803},$$

$$M_X = \max_P \theta_P = \frac{5168}{2803} > 1.$$

A.3.8. C_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{171}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{36}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{18}{5}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{12}{5}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{12}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{567}{40}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{567}{80}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{189}{40}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{189}{40}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{189}{20}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{189}{80}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{189}{80}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{309}{20}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{429}{80}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{309}{20},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{3040x_1 - 1024}{4961},$$

$$M_X = \max_P \theta_P = \frac{5056}{4961} > 1.$$

A.3.9. C_3 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{171}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{63}{20}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{63}{20}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{21}{10}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{21}{10}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{243}{20}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{621}{80}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{117}{80}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{117}{80}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{243}{20}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{117}{80}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{117}{80}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{57}{4}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{453}{80}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{57}{4},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{21280x_1 + 21280x_2 - 6272}{23368},$$

$$M_X = \max_P \theta_P = \frac{1876}{2921} < 1.$$

A.3.10. C_4 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{81}{4}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{81}{8}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{81}{16}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{81}{8}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{81}{8}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{81}{16}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{81}{8},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = 0,$$

$$M_X = \max_P \theta_P = 0 < 1.$$

A.3.11. E_1 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{605}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 12, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{67}{10}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{173}{30}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{173}{30}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{587}{60}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{367}{80}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{719}{144}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{719}{144}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{623}{24}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{1007}{180}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{1007}{180}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{4709}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{3809}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{4709}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{4447297800x_1 + 166293600x_2 - 2161259136}{1528096589},$$

$$M_X = \max_P \theta_P = \frac{3117506664}{1528096589} > 1.$$

A.3.12. E_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{163}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{34}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 3, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{19}{15}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{19}{15}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{83}{12}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{247}{240}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{157}{80}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{157}{80}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{883}{60}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{73}{16}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{73}{16}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{883}{60}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{983}{240}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{883}{60},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{1517634320x_1 + 215186080x_2 - 538182656}{1038057499},$$

$$M_X = \max_P \theta_P = \frac{1840195984}{1038057499} > 1.$$

A.3.13. E_3 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{431}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{11}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{47}{20}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{47}{60}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{47}{60}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{329}{60}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{347}{240}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{347}{720}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{347}{720}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1319}{120}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{1319}{360}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{1319}{360}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{361}{36}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{2291}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{361}{36},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{3423372660x_1 + 1994323200x_2 - 680357088}{6541476757},$$

$$M_X = \max_P \theta_P = \frac{8725985172}{6541476757} > 1.$$

A.3.14. D_1 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{74}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{118}{15}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{118}{15}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{59}{10}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{59}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{403}{15}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{203}{30}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{10}{3}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{20}{3}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{403}{15}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{10}{3}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{20}{3}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{697}{45}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = 5, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = 10,
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{21830x_4 - 10443}{8057},$$

$$M_X = \max_P \theta_P = \frac{11387}{8057} > 1.$$

A.3.15. D_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 24, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{157}{15}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{157}{15}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{157}{10}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{43}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{1631}{45}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{391}{90}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{124}{9}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{298}{45}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1631}{45}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{124}{9}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{298}{45}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{62}{3}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{149}{15}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{80}{9},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{1461480x_3 + 588060x_4 - 1166773}{1129107},$$

$$M_X = \max_P \theta_P = \frac{2344247}{1129107} > 1.$$

A.3.16. D_3 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{70}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{83}{12}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{83}{12}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{219}{20}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{49}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{2297}{90}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{262}{45}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{751}{90}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{83}{15}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{2297}{90}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{751}{90}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{83}{15}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1571}{90}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{341}{45}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{406}{45},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{6623400x_3 + 23783700x_4 - 13097421}{13920179},$$

$$M_X = \max_P \theta_P = \frac{23933079}{13920179} > 1.$$

A.3.17. D_4 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{70}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{157}{10}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{161}{60}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{157}{20}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{158}{15}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{182}{5}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -7, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{91}{5}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{56}{5}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{571}{30}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{7}{2}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{18}{5}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{2263}{90}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{28}{5}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{46}{5},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{1389500x_1 + 10897600x_4 - 5854423}{5141177},$$

$$M_X = \max_P \theta_P = \frac{10601177}{5141177} > 1.$$

A.3.18. D_5 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{62}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{16}{3}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{16}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 8, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{53}{10}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{76}{15}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{311}{15}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{76}{15}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{38}{5}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{62}{9},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{620x_3 - 240}{349},$$

$$M_X = \max_P \theta_P = \frac{380}{349} > 1.$$

A.3.19. D_6 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{62}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{12}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{12}{5}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{18}{5}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{36}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = 14, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{86}{15}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{19}{15}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{38}{15}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = 14, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{19}{15}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{38}{15}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{532}{45}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{19}{5}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{38}{5},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{2790x_4 - 972}{1973},$$

$$M_X = \max_P \theta_P = \frac{1818}{1973} < 1.$$

A.3.20. D_7 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{81}{4}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{99}{10}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{99}{20}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{99}{20}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{99}{20}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{153}{10}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{153}{20}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{153}{20}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{153}{20}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{171}{20}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{153}{40}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{153}{40}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{303}{20}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{153}{40}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{303}{20},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{495x_1 - 242}{523},$$

$$M_X = \max_P \theta_P = \frac{748}{523} > 1.$$

A.3.21. D_8 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 20, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{161}{60}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{161}{60}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{153}{40}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{27}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{269}{18}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{469}{90}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{407}{90}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{89}{45}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{269}{18}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{407}{90}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{89}{45}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{407}{30}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{89}{15}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{316}{45},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{783290760x_1 + 783290760x_2 - 1168558320x_3}{2130145727} + \frac{2808344100x_4 - 811654353}{2130145727},$$

$$M_X = \max_P \theta_P = \frac{3165248067}{2130145727} > 1.$$

A.3.22. D_9 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{58}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{8}{3}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{8}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 4, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 4, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{661}{45}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{449}{90}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{106}{45}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{106}{45}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{661}{45}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{106}{45}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{106}{45}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{298}{45}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{4}{9}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{298}{45},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{870x_3 + 870x_4 - 360}{1177},$$

$$M_X = \max_P \theta_P = \frac{1380}{1177} > 1.$$

A.3.23. D_{10} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{58}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{36}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{16}{15}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{18}{5}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{76}{15}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{82}{5}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -6, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{41}{5}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{22}{5}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{158}{15}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -3, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{6}{5}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{644}{45}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{11}{5}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{34}{5},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{18125x_1 + 34945x_4 - 15908}{46732},$$

$$M_X = \max_P \theta_P = \frac{18353}{11683} > 1.$$

A.3.24. D_{11} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{153}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{36}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{18}{5}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{39}{5}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{39}{10}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{261}{20}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{261}{40}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{189}{20}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{189}{40}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{333}{40}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{189}{40}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{189}{80}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{321}{20}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{321}{40}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{573}{40},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{465x_1 + 375x_3 - 328}{1007},$$

$$M_X = \max_P \theta_P = \frac{1727}{1007} > 1.$$

A.3.25. D_{12} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{56}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{4}{3}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{4}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 4, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{176}{15}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{24}{5}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{32}{15}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{176}{15}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{32}{15}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{32}{5}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = -\frac{56}{9},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{70x_3 - 15}{97},$$

$$M_X = \max_P \theta_P = \frac{55}{97} < 1.$$

A.3.26. D_{13} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 18, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = 9, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{9}{2}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = 3, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = 6, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = 6,
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = 0,$$

$$M_X = \max_P \theta_P = 0 < 1.$$

A.3.27. D_{14} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 18, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{9}{2}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{9}{4}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{9}{4}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{54}{5}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{27}{5}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{27}{5}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{81}{10}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{27}{10}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = 6, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{111}{10},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{20x_1 - 5}{43},$$

$$M_X = \max_P \theta_P = \frac{35}{43} < 1.$$

A.3.28. D_{15} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 18, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 3, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{3}{2}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = 9, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{9}{2}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = 9, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = 6, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = 3, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = 9,
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{6x_3 - 1}{11},$$

$$M_X = \max_P \theta_P = \frac{5}{11} < 1.$$

A.3.29. D_{16} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 18, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{79}{60}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{79}{60}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{107}{20}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{7}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{1033}{90}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{206}{45}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{35}{18}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{17}{45}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1033}{90}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{35}{18}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{17}{45}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{947}{90}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{211}{45}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{262}{45},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{796536x_3 - 335484x_4 - 210655}{1061921},$$

$$M_X = \max_P \theta_P = \frac{1046933}{1061921} < 1.$$

A.3.30. D_{17} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{135}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{9}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{9}{5}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{9}{10}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{9}{10}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{171}{20}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{99}{20}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{171}{40}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{99}{40}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{171}{20}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{99}{40}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{171}{40}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{363}{40}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{99}{80}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{363}{40},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{75x_1 + 75x_2 - 16}{134},$$

$$M_X = \max_P \theta_P = \frac{59}{134} < 1.$$

A.3.31. D_{18} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = -\frac{99}{80}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{14}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{14}{5}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{21}{10}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{21}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{73}{5}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{23}{6}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{26}{15}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{52}{15}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{73}{5}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{26}{15}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{52}{15}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{53}{9}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{13}{5}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{26}{5},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{-5250x_4 - 1323}{5177},$$

$$M_X = \max_P \theta_P = \frac{3927}{5177} < 1.$$

A.3.32. D_{19} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = -\frac{99}{80}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{4}{15}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{4}{15}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{2}{5}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{4}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{142}{15}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{58}{15}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{13}{15}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{26}{15}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{142}{15}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{13}{15}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{26}{15}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{316}{45}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{13}{5}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{26}{5},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = -\frac{250x_4 + 12}{1613},$$

$$M_X = \max_P \theta_P = \frac{238}{1613} < 1.$$

A.3.33. G_1 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{529}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{93}{10}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{35}{6}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{71}{30}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{629}{60}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{1517}{180}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{2803}{720}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{77}{120}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{5837}{720}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{3149}{180}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{989}{180}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{4781}{720}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{4187}{360}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{745}{144}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{5147}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{37598378760x_1 - 8882851620x_3 - 16817588112}{16641653953},$$

$$M_X = \max_P \theta_P = \frac{29663642268}{16641653953} > 1.$$

A.3.34. G_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{75}{4}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{89}{10}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{89}{20}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{197}{15}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{7}{3}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{517}{40}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{517}{80}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{595}{36}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{839}{180}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{911}{120}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{595}{72}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{839}{360}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{12301}{360}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{401}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{1727}{120},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{1410100500x_1 + 366948000x_3 - 926354392}{1802902233},$$

$$M_X = \max_P \theta_P = \frac{372321328}{163900203} > 1.$$

A.3.35. G_3 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{433}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{89}{30}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{133}{60}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{133}{30}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{37}{10}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{217}{36}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{19}{720}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{19}{360}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{2189}{720}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1787}{180}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{877}{240}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{265}{144}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{877}{120}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{265}{72}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{2333}{180},
\end{aligned}$$

$$\begin{aligned}
\theta_P(x_1, x_2, x_3, x_4) &= \frac{28885436928x_1 - 35779946976x_3 - 13541869440}{45785112888}, \\
M_X &= \max_P \theta_P = \frac{2130146436}{1907713037} > 1.
\end{aligned}$$

A.3.36. G_4 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{139}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = -\frac{17}{20}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{301}{60}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{127}{30}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{1}{30}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{147}{20}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{785}{144}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{2333}{720}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{161}{240}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{383}{36}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{53}{80}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{181}{80}, \\
c_{33} &= \int_P x_3 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{2261}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{1039}{240}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{583}{60},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{10356297206700x_1 + 33083176976940x_2 + 16155069457920x_3}{48057952407691} - \frac{12981537414704}{48057952407691},$$

$$M_X = \max_P \theta_P = \frac{91293727706236}{48057952407691} > 1.$$

A.3.37. G_5 .

$$\begin{aligned} b_0 = \text{vol}(P) &= \frac{203}{12}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = -\frac{16}{5}, \\ b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{8}{5}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{28}{15}, \\ b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{14}{15}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{767}{120}, \\ c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{767}{240}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{68}{45}, \\ c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{34}{45}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{3181}{360}, \\ c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{34}{45}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{17}{45}, \\ c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1039}{120}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{1039}{240}, \\ c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{569}{72}, \end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{-4015697280x_1 + 2205554400x_3 - 1002991104}{5976434111},$$

$$M_X = \max_P \theta_P = \frac{7423814976}{5976434111} > 1.$$

A.3.38. G_6 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{401}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{12}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{6}{5}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{24}{5}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{6}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{997}{120}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{997}{240}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{2437}{360}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{277}{360}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1357}{180}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{2437}{720}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{277}{720}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{2437}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{2437}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{1357}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{1732320x_3 - 497664}{4388521},$$

$$M_X = \max_P \theta_P = \frac{6431616}{4388521} > 1.$$

A.3.39. H_1 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{93}{4}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{153}{40}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{161}{10}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{161}{15}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{161}{15}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{3043}{360}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{2449}{240}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{2449}{360}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{2449}{360}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{2449}{120}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{2449}{180}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{2449}{180}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{12947}{360}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{3151}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{12947}{360},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{299460x_2 - 207368}{172227},$$

$$M_X = \max_P \theta_P = \frac{391552}{172227} > 1.$$

A.3.40. H_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{505}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{163}{20}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{247}{20}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{41}{6}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{41}{6}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{1391}{180}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{6139}{720}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{3901}{720}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{3901}{720}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{2963}{180}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{1999}{240}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1999}{240}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{4361}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{3773}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{4361}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{12590952900x_1 + 11321625300x_2 - 11521822032}{12261327193},$$

$$M_X = \max_P \theta_P = \frac{23712381468}{12261327193} > 1.$$

A.3.41. H_3 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{239}{12}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{42}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{179}{20}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{28}{5}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{28}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{881}{120}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{1697}{240}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{881}{180}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{881}{180}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{4771}{360}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{1697}{360}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1697}{360}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{7363}{360}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{3839}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{7363}{360},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{1983987756x_1 + 260718408x_2 - 953921016}{999563887},$$

$$M_X = \max_P \theta_P = \frac{1551503556}{999563887} > 1.$$

A.3.42. H_4 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{149}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{89}{20}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{89}{10}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{89}{30}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{89}{30}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{1133}{180}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{311}{48}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{311}{144}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{311}{144}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{311}{24}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{311}{72}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{311}{72}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{2599}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{3643}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{2599}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{795660x_2 - 380208}{778267},$$

$$M_X = \max_P \theta_P = \frac{173016}{111181} > 1.$$

A.3.43. H_5 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{415}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{97}{20}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{29}{5}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{97}{60}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{97}{60}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{139}{24}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{1259}{240}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{139}{72}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{139}{72}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{367}{36}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{1259}{720}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1259}{720}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{2021}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{3347}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{2021}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{6818815200x_1 + 2922039900x_2 - 2892669984}{8410595791},$$

$$M_X = \max_P \theta_P = \frac{9770225016}{8410595791} > 1.$$

A.3.44. H_6 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{409}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{27}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{61}{20}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{31}{12}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{31}{12}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{1013}{180}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{3049}{720}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{337}{144}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{337}{144}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{2941}{260}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{1}{10}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1}{10}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{2411}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{3173}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{2411}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{11025012180x_1 - 2015224800x_2 - 3132829152}{6776874793},$$

$$M_X = \max_P \theta_P = \frac{7892183028}{6776874793} > 1.$$

A.3.45. H_7 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{191}{12}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{61}{20}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{31}{20}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{46}{15}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{46}{15}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{335}{72}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{1859}{720}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{497}{360}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{497}{360}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1963}{360}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{689}{360}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{689}{360}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1031}{72}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{2783}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{1031}{72},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{3404937900x_1 - 2489673540x_2 - 894914424}{2191715291},$$

$$M_X = \max_P \theta_P = \frac{2510023476}{2191715291} > 1.$$

A.3.46. H_8 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{63}{4}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{3}{2}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 3, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{33}{8}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{63}{16}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{63}{8}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{63}{8}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{63}{16}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{63}{8},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{168x_2 - 32}{409},$$

$$M_X = \max_P \theta_P = \frac{304}{409} < 1.$$

A.3.47. H_9 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{367}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{11}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{11}{20}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{11}{20}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{11}{20}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{827}{180}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{2267}{720}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{347}{720}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{347}{720}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1133}{180}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{151}{144}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{151}{144}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1613}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{535}{144}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{1613}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = -\frac{-3841851420x_1 + 1395187200x_2 + 502543008}{5523943807},$$

$$M_X = \max_P \theta_P = \frac{3339308412}{5523943807} < 1.$$

A.3.48. H_{10} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{117}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = -\frac{4}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{117}{8}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{117}{8}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{117}{8}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{151}{36}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{619}{240}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{619}{720}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{619}{720}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{619}{120}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{619}{360}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{619}{360}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1567}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{503}{144}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{1567}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{-37440x_2 - 4096}{116609},$$

$$M_X = \max_P \theta_P = \frac{33344}{116609} < 1.$$

A.3.49. L_1 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 20, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{42}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{21}{5}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{21}{5}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{21}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{116}{15}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{58}{15}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{58}{15}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{58}{15}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{542}{45}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{29}{15}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{29}{15}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{542}{45}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{29}{15}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{542}{45},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{3150x_1 - 1323}{1577},$$

$$M_X = \max_P \theta_P = \frac{1827}{1577} > 1.$$

A.3.50. L_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{58}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{92}{15}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{43}{10}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{313}{60}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{313}{60}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{314}{45}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{2}{45}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{52}{15}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{52}{15}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{362}{45}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -4, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -4, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{149}{10}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{56}{15}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{149}{10},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{3878228x_1 - 2370692x_2 - 1757609}{2637399},$$

$$M_X = \max_P \theta_P = \frac{4491311}{2637399} > 1.$$

A.3.51. L_3 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{56}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{103}{15}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{103}{30}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{112}{15}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{56}{15}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{104}{15}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{52}{15}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{248}{45}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{124}{45}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{487}{45}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{124}{45}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{62}{45}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{193}{15}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{193}{30}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = 13,
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{26763240x_1 + 7896000x_3 - 13003449}{20356871},$$

$$M_X = \max_P \theta_P = \frac{29551791}{20356871} > 1.$$

A.3.52. L_4 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 18, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{76}{15}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{169}{60}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{241}{30}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{241}{60}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{286}{45}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{34}{45}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{22}{5}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{11}{5}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{139}{18}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{73}{15}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{73}{30}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{79}{5}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{79}{10}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{251}{18},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{6334289640x_1 - 2670575400x_2 + 2239779600x_3}{6291985913} - \frac{3200486167}{6291985913},$$

$$M_X = \max_P \theta_P = \frac{12523717673}{6291985913} > 1.$$

A.3.53. L_5 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{52}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{16}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{8}{3}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{8}{3}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{52}{9}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = 0, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{92}{15}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{46}{15}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{46}{15}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{48}{5}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{23}{15}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{48}{5},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{-260x_2 - 80}{219},$$

$$M_X = \max_P \theta_P = \frac{60}{73} < 1.$$

A.3.54. L_6 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = -\frac{99}{80}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -4, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{16}{3}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{8}{3}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{50}{9}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = 0, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{86}{15}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{64}{15}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{32}{15}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{51}{5}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{51}{10}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{161}{15},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{-10150x_2 + 7600x_3 - 4868}{17787},$$

$$M_X = \max_P \theta_P = \frac{38}{33} > 1.$$

A.3.55. L_7 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 16, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{8}{3}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{4}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{8}{3}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{4}{3}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{16}{3}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{8}{3}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{4}{9}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{2}{9}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = 8, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{2}{9}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1}{9}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{16}{3}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{8}{3}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = 8,
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{6x_1 + 6x_3 - 2}{11},$$

$$M_X = \max_P \theta_P = \frac{10}{11} < 1.$$

A.3.56. L_8 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 16, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{16}{3}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = 0, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{16}{3}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{16}{3}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{16}{3},
\end{aligned}$$

$$\begin{aligned}
\theta_P(x_1, x_2, x_3, x_4) &= 0, \\
M_X &= \max_P \theta_P = 0 < 1.
\end{aligned}$$

A.3.57. L_9 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 16, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{8}{3}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{4}{3}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{16}{3}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = 0, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{16}{3}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{16}{3}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{8}{3}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = 8,
\end{aligned}$$

$$\begin{aligned}
\theta_P(x_1, x_2, x_3, x_4) &= \frac{6x_3 - 1}{11}, \\
M_X &= \max_P \theta_P = \frac{5}{11} < 1.
\end{aligned}$$

A.3.58. L_{10} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 16, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{71}{15}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{71}{30}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{71}{60}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{71}{60}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{256}{45}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{128}{45}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{64}{45}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{64}{45}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{122}{15}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{119}{45}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{61}{15}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{53}{6}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{119}{90}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{53}{6},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{17040x_1 - 5041}{15439},$$

$$M_X = \max_P \theta_P = \frac{11999}{15439} < 1.$$

A.3.59. L_{11} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{44}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{44}{9}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = 0, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{68}{15}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{34}{15}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{34}{15}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{32}{5}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{17}{15}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{32}{5},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = 0,$$

$$M_X = \max_P \theta_P = 0 < 1.$$

A.3.60. L_{12} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{44}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = -\frac{17}{15}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{17}{30}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{16}{5}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{8}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{68}{15}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{34}{15}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{136}{45}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{68}{45}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{271}{45}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{68}{45}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{34}{45}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{113}{15}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{113}{30}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{127}{15},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{5940x_1 + 58080x_3 - 12213}{118907},$$

$$M_X = \max_P \theta_P = \frac{98007}{118907} < 1.$$

A.3.61. L_{13} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{44}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{34}{15}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{17}{15}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{17}{15}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{17}{15}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{68}{15}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{34}{15}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{34}{15}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{34}{15}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{322}{45}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{17}{15}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{17}{15}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{322}{45}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{17}{15}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{254}{45},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{110x_1 - 17}{203},$$

$$M_X = \max_P \theta_P = \frac{93}{203} < 1.$$

A.3.62. I_1 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{62}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{173}{20}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{521}{60}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{521}{30}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{1}{30}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{1393}{180}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{443}{72}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{443}{36}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{571}{360}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{2197}{90}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{88}{5}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{953}{360}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{176}{5}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{953}{180}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{563}{45},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{455864610x_1 + 140393265x_3 - 308777028}{298795537},$$

$$M_X = \max_P \theta_P = \frac{708660642}{298795537} > 1.$$

A.3.63. I_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{463}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 8, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 4, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{57}{5}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{23}{10}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{141}{20}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{141}{40}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{1627}{180}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{911}{360}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{4121}{360}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{1627}{360}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{911}{720}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{4363}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{1109}{360}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{1607}{120},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{3407814270x_1 + 302232510x_3 - 1591773552}{1752843533},$$

$$M_X = \max_P \theta_P = \frac{3024970758}{1752843533} > 1.$$

A.3.64. I_3 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{221}{12}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{127}{20}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{101}{12}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{61}{6}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{23}{10}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{286}{45}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{679}{120}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{2771}{260}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{259}{120}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1583}{120}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{1193}{120}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1069}{240}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1663}{72}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{131}{48}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{139}{10},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{1119451198840x_1 + 553671270165x_2 + 212169307415x_3}{1114834051022} - \frac{756143145443}{1114834051022},$$

$$M_X = \max_P \theta_P = \frac{1265748565401}{557417025511} > 1.$$

A.3.65. I_4 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{433}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{59}{10}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{121}{30}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{419}{30}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{13}{12}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{557}{90}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{349}{180}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{151}{15}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{52}{45}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{5023}{360}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{3533}{360}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{2137}{720}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1069}{36}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{1721}{360}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{3607}{360},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{29597715x_1 + 23135190x_3 - 27588804}{42931741},$$

$$M_X = \max_P \theta_P = \frac{94549671}{42931741} > 1.$$

A.3.66. I_5 .

$$\begin{aligned}
b_0 = \text{vol}(P) &= \frac{415}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{63}{10}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{67}{20}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{403}{69}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{67}{10}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{427}{72}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{551}{240}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{2263}{360}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{551}{120}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1577}{120}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{43}{90}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{463}{60}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{2881}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{43}{45}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{463}{30},
\end{aligned}$$

$$\begin{aligned}
\theta_P(x_1, x_2, x_3, x_4) &= \frac{10897419402618300x_1 + 101247812541450x_3}{6984760752795427} \\
&\quad + \frac{1845227650200450x_4 - 4719505353973848}{6984760752795427},
\end{aligned}$$

$$M_X = \max_P \theta_P = \frac{11713596999245802}{6984760752795427} > 1.$$

A.3.67. I_6 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{137}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{53}{15}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{43}{10}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{43}{5}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{38}{15}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{253}{45}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{239}{90}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{239}{45}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{7}{45}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{4933}{360}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{923}{120}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{1813}{720}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{923}{60}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{1813}{360}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{2693}{360},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{48154815x_1 + 164153400x_3 - 92371752}{230998253},$$

$$M_X = \max_P \theta_P = \frac{448243263}{230998253} > 1.$$

A.3.68. I_7 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = -\frac{99}{80}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{21}{4}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 8, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{5}{4}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{11}{2}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = 0, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{34}{5}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{21}{10}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{50}{9}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{281}{15}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{77}{30}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{109}{10},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{527400x_1 + 99600x_3 - 213939}{467581},$$

$$M_X = \max_P \theta_P = \frac{711861}{467581} > 1.$$

A.3.69. I_8 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 16, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = -\frac{53}{20}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{88}{15}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{163}{60}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1}{4}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{443}{90}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{1547}{360}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{431}{360}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{103}{360}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{3611}{360}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{581}{360}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1483}{720}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{3743}{360}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{2731}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{829}{90},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{-354416417280x_1 + 991006584000x_2 + 428321791680x_3}{1577014856729} - \frac{494794770791}{1577014856729},$$

$$M_X = \max_P \theta_P = \frac{3126600189529}{1577014856729} > 1.$$

A.3.70. I_9 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{65}{4}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{71}{60}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{169}{60}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{409}{60}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{169}{60}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{919}{180}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{43}{120}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{79}{18}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{43}{120}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{637}{60}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{37}{8}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{299}{120}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{491}{36}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{37}{8}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{247}{36},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{-11160375x_1 + 23983575x_3 - 9248098}{31553027},$$

$$M_X = \max_P \theta_P = \frac{51542252}{31553027} > 1.$$

A.3.71. I_{10} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{389}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{41}{10}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{41}{20}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{161}{30}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{19}{30}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{313}{60}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{313}{120}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{202}{45}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{131}{360}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1021}{120}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{101}{45}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{131}{720}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{106}{9}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{164}{45}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{551}{72},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{762471120x_1 + 302223825x_3 - 292939812}{1008086248},$$

$$M_X = \max_P \theta_P = \frac{1376202783}{1008086248} > 1.$$

A.3.72. I_{11} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 16, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{7}{4}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{11}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{59}{12}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1}{4}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{443}{90}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{2}{9}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{341}{90}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{61}{90}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{248}{45}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{116}{45}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{71}{45}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1169}{90}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{149}{45}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{829}{90},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{1527360x_1 + 5306880x_2 + 2119680x_3 - 2034575}{7524593},$$

$$M_X = \max_P \theta_P = \frac{13278385}{7524593} > 1.$$

A.3.73. I_{12} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{347}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = -\frac{3}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{3}{10}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{3}{5}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{6}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{251}{60}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{251}{120}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{119}{45}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{277}{360}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{2309}{360}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{119}{90}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{277}{720}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{373}{45}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{127}{45}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{2309}{360},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = -\frac{1780035048x_1 - 859184145x_3}{10127095471} - \frac{1344488976x_4 + 221112504}{10127095471},$$

$$M_X = \max_P \theta_P = \frac{3762595665}{10127095471} < 1.$$

A.3.74. I_{13} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{46}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{7}{4}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{43}{12}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{11}{12}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{47}{10}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = 0, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{107}{30}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{17}{30}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{46}{9}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{301}{30}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{97}{30}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{211}{30},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{98900x_1 + 219420x_3 - 62565}{650251},$$

$$M_X = \max_P \theta_P = \frac{694595}{650251} > 1.$$

A.3.75. I_{14} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{119}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{61}{30}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{13}{10}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{59}{30}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{13}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{1619}{360}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{427}{720}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{142}{45}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{427}{360}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{3043}{360}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{1003}{720}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{473}{120}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{2963}{360}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{1003}{360}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{473}{60},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{21660926805x_1 + 90160555275x_3}{218756440813} + \frac{112125015975x_4 - 34479642672}{218756440813},$$

$$M_X = \max_P \theta_P = \frac{211431316083}{218756440813} < 1.$$

A.3.76. I_{15} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{337}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{5}{2}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{5}{4}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{23}{5}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{79}{20}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{79}{40}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{821}{180}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{601}{360}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1859}{360}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{821}{360}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{601}{720}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{2381}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{739}{360}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{1009}{120},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{329975235x_1 + 184724865x_3 - 119264508}{739116617},$$

$$M_X = \max_P \theta_P = \frac{949610187}{739116617} > 1.$$

A.3.77. M_1 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{385}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = -\frac{79}{30}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{79}{20}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{79}{30}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{79}{30}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{193}{36}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{1949}{720}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{19}{720}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{19}{720}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{2411}{180}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{1949}{720}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1949}{720}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{193}{36}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{19}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{193}{36},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{-1824900x_1 + 1824900x_3 + 1824900x_4 - 898704}{2853121},$$

$$M_X = \max_P \theta_P = \frac{4575996}{2853121} > 1.$$

A.3.78. M_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{139}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{161}{30}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{239}{60}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{20}{3}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{20}{3}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{1109}{180}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{481}{240}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{481}{240}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{481}{240}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{2503}{180}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{1223}{240}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1223}{240}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{2101}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{173}{48}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{2101}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{2555701260x_1 + 2483476860x_3}{4944865931} + \frac{2483476860x_4 - 2695172464}{4944865931},$$

$$M_X = \max_P \theta_P = \frac{9794436236}{4944865931} > 1.$$

A.3.79. M_3 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{123}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{89}{30}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{13}{10}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{7}{60}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{109}{20}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{185}{36}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{19}{720}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{1549}{720}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{2189}{720}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1433}{180}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{413}{720}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1733}{720}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1279}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{-2741}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{2099}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{4151273350320x_1 + 4524155739540x_3}{16440690503203} + \frac{10150485305940x_4 - 4433392348592}{16440690503203},$$

$$M_X = \max_P \theta_P = \frac{25645181180008}{16440690503203} > 1.$$

A.3.80. M_4 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{123}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{41}{30}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{157}{60}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{119}{30}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{79}{30}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{893}{180}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{253}{720}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{553}{144}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{301}{720}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1879}{180}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{1813}{720}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1789}{720}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1553}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{179}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{185}{36},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{-2793968096940x_1 + 8560548364140x_3}{13060549547443} + \frac{8232843774060x_4 - 3370292079152}{13060549547443},$$

$$M_X = \max_P \theta_P = \frac{19189680326248}{13060549547443} > 1.$$

A.3.81. M_5 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{91}{6}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{347}{60}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{3}{2}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{169}{30}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 3, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{577}{45}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{2287}{720}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{141}{16}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{2287}{360}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{2399}{360}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{2423}{720}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{313}{80}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1039}{90}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{2423}{360}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{313}{40},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{2003379647360x_1 + 2700009652340x_3}{5748810067571} + \frac{-1155042818020x_4 - 1538316805364}{5748810067571},$$

$$M_X = \max_P \theta_P = \frac{12265145105056}{5748810067571} > 1.$$

A.3.82. J_1 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{46}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = -\frac{139}{30}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{71}{20}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{37}{30}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{37}{30}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{871}{180}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{2173}{720}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{431}{720}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{431}{720}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1417}{120}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{313}{180}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{313}{180}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{2827}{360}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{3581}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{2827}{360},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{-353629140x_1 + 95218620x_3 + 95218620x_4 - 122175279}{271746296},$$

$$M_X = \max_P \theta_P = \frac{326672481}{271746296} > 1.$$

A.3.83. J_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{163}{12}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{2}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{2}{3}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{2}{3}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{137}{36}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{137}{144}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{137}{144}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{137}{144}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{2381}{360}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{34}{45}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{34}{45}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{2381}{360}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{2989}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{2381}{360},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{-78240x_1 + 156480x_3 + 156480x_4 - 15360}{450983},$$

$$M_X = \max_P \theta_P = \frac{141120}{450983} < 1.$$

A.3.84. Q_1 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{221}{12}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{329}{60}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{329}{30}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{329}{60}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{329}{60}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{233}{36}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{437}{60}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{437}{120}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{437}{120}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{437}{30}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{437}{60}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{437}{60}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{293}{20}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{437}{120}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{293}{20},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{363545x_2 - 216482}{266403},$$

$$M_X = \max_P \theta_P = \frac{510608}{266403} > 1.$$

A.3.85. Q_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{135}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{257}{60}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{257}{30}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{124}{15}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{62}{15}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{2071}{360}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{145}{24}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{109}{24}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{109}{48}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{145}{12}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{109}{12}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{109}{24}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{311}{20}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{311}{40}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{4883}{360},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{211857525x_2 + 80463375x_3 - 146967508}{237174992},$$

$$M_X = \max_P \theta_P = \frac{518137667}{237174992} > 1.$$

A.3.86. Q_3 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{197}{12}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{88}{15}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{391}{60}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{44}{15}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{44}{15}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{173}{30}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{65}{12}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{173}{60}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{173}{60}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1861}{180}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{65}{24}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{65}{24}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{187}{20}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{173}{120}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{187}{20},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{269739295x_1 + 61718130x_2 - 120893422}{201232113},$$

$$M_X = \max_P \theta_P = \frac{90760711}{67077371} > 1.$$

A.3.87. Q_4 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{135}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{28}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{257}{30}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{257}{60}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{14}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{119}{20}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{1123}{180}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{1123}{360}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{119}{40}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{145}{12}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{145}{24}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1123}{360}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1501}{120}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1123}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{379}{40},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{51832575x_1 + 58443525x_2 - 46869844}{73313456},$$

$$M_X = \max_P \theta_P = \frac{121849781}{73313456} > 1.$$

A.3.88. Q_5 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{373}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{281}{60}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{82}{15}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{167}{30}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{167}{60}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{21}{4}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{577}{120}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{49}{12}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{49}{24}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{671}{72}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{473}{120}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{473}{240}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{199}{20}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{199}{40}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{3727}{360},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{394596985345x_1 + 192548075640x_2 + 234003144575x_3 - 270449943142}{573603226348} - \frac{270449943142}{573603226348},$$

$$M_X = \max_P \theta_P = \frac{977249482633}{573603226348} > 1.$$

A.3.89. Q_6 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{46}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{37}{12}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{37}{6}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{37}{12}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{151}{30}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{24}{5}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{12}{5}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{48}{5}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{24}{5}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{46}{9}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{311}{30},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{17020x_2 - 6845}{19651},$$

$$M_X = \max_P \theta_P = \frac{27195}{19651} > 1.$$

A.3.90. Q_7 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{121}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{97}{30}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{97}{15}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{97}{60}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{97}{60}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{59}{12}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{226}{45}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{59}{24}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{19}{360}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{452}{45}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{113}{45}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{113}{45}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{931}{120}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{19}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{931}{120},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{176055x_2 - 75272}{198188},$$

$$M_X = \max_P \theta_P = \frac{138419}{99094} > 1.$$

A.3.91. Q_8 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{44}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{7}{2}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{53}{12}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{7}{4}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{71}{15}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{21}{5}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{71}{30}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{83}{10}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{21}{10}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{44}{9}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{229}{30},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{1663200x_1 + 982960x_2 - 692905}{2735927},$$

$$M_X = \max_P \theta_P = \frac{2936215}{2735927} > 1.$$

A.3.92. Q_9 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{341}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{113}{30}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{71}{30}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{113}{60}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{7}{10}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{91}{20}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{152}{45}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{911}{40}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{211}{360}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{59}{9}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{76}{45}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{143}{90}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{901}{120}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{211}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{479}{72},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{461229780x_1 - 53988825x_2 - 113280372}{395454118},$$

$$M_X = \max_P \theta_P = \frac{173974704}{197727059} < 1.$$

A.3.93. Q_{10} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 14, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{4}{3}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{8}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{7}{3}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{7}{6}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{13}{3}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{7}{2}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{2}{9}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1}{9}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = 7, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{4}{9}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{2}{9}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{14}{3}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{7}{3}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = 7,
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{1848x_2 + 2454x_3 - 761}{4499},$$

$$M_X = \max_P \theta_P = \frac{5389}{4499} > 1.$$

A.3.94. Q_{11} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 14, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{4}{3}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{8}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{13}{3}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{7}{2}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = 7, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{14}{3}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{14}{3},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{168x_2 - 32}{409},$$

$$M_X = \max_P \theta_P = \frac{304}{409} < 1.$$

A.3.95. Q_{12} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{331}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{16}{5}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{113}{30}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{113}{60}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{17}{60}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{203}{45}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{169}{45}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{169}{90}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{17}{45}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{427}{60}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{427}{120}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{121}{72}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{329}{40}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{121}{144}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{2081}{360},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{115654710x_1 + 66679950x_2 - 45045768}{196251577},$$

$$M_X = \max_P \theta_P = \frac{203968842}{196251577} > 1.$$

A.3.96. Q_{13} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{55}{4}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{139}{60}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{29}{60}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{7}{5}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{7}{5}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{733}{180}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{227}{90}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{31}{40}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{31}{40}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{185}{36}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{157}{120}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{157}{120}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1243}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{25}{24}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{1243}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{38978775x_1 - 23199825x_2 - 7382842}{35875483},$$

$$M_X = \max_P \theta_P = \frac{31595933}{35875483} < 1.$$

A.3.97. Q_{14} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{325}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{61}{30}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{13}{60}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{103}{30}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{103}{60}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{1459}{360}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{122}{45}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{427}{360}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{427}{720}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1951}{360}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{119}{72}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{119}{144}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{437}{60}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{437}{120}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{2911}{360},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{1233812146125x_1 - 151909753125x_2 + 1142554261875x_3 - 472512532902}{2450627729348} - \frac{472512532902}{2450627729348},$$

$$M_X = \max_P \theta_P = \frac{3046408136973}{2450627729348} > 1.$$

A.3.98. Q_{15} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{40}{3}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{7}{4}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{11}{12}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{5}{12}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{121}{30}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{29}{10}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{17}{30}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{57}{10}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{7}{5}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{40}{9}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{181}{30},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{210720x_1 - 39680x_2 - 24929}{394975},$$

$$M_X = \max_P \theta_P = \frac{185791}{394975} < 1.$$

A.3.99. Q_{16} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{155}{12}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{17}{15}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{139}{60}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{17}{30}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{17}{30}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{37}{10}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{179}{80}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{37}{20}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{37}{20}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1127}{180}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{179}{120}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{179}{120}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{71}{12}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{37}{40}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{929}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{1544575x_1 + 43444950x_2 - 7927586}{111477439},$$

$$M_X = \max_P \theta_P = \frac{80506889}{111477439} < 1.$$

A.3.100. Q_{17} .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{299}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = -\frac{4}{15}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{8}{15}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{2}{15}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{2}{15}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{211}{60}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{109}{45}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{197}{360}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{211}{120}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{218}{45}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{109}{90}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{109}{90}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1949}{360}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{197}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{1949}{360},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{-17940x_2 - 768}{162187},$$

$$M_X = \max_P \theta_P = \frac{17172}{162187} < 1.$$

A.3.101. K_1 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{91}{6}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 5, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{5}{2}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{10}{3}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{10}{3}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{281}{60}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{281}{120}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{281}{90}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{281}{90}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{253}{60}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{281}{180}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{281}{180}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{2527}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{1403}{360}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{2527}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{27300x_1 - 9000}{16571},$$

$$M_X = \max_P \theta_P = \frac{18300}{16571} > 1.$$

A.3.102. K_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{59}{4}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{15}{4}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{15}{4}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{5}{2}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{5}{2}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{13}{3}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{281}{120}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{89}{40}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{89}{40}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{13}{3}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{89}{40}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{89}{40}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{121}{10}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{153}{40}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{121}{10},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{2950x_1 + 2950x_2 - 1500}{3751},$$

$$M_X = \max_P \theta_P = \frac{400}{341} > 1.$$

A.3.103. K_3 .

$$\begin{aligned}
b_0 = \text{vol}(P) &= \frac{167}{12}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{5}{2}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{5}{4}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{5}{6}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{5}{6}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{239}{60}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{239}{120}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{239}{180}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{239}{180}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{58}{15}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{239}{180}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{239}{180}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{761}{90}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{1283}{360}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{761}{90},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{25050x_1 - 4500}{35413},$$

$$M_X = \max_P \theta_P = \frac{20550}{35413} < 1.$$

A.3.104. K_4 .

$$\begin{aligned}
b_0 = \text{vol}(P) &= \frac{27}{2}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{15}{4}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{15}{8}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{15}{4}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{27}{4}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{27}{8}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{27}{4},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = 0,$$

$$M_X = \max_P \theta_P = 0 < 1.$$

A.3.105. R_1 .

$$b_0 = \text{vol}(P) = \frac{83}{6},$$

$$b_1 = \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 1,$$

$$b_2 = \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 1,$$

$$b_3 = \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{35}{8},$$

$$b_4 = \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{23}{12},$$

$$c_{11} = \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{2551}{360},$$

$$c_{12} = \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{3149}{720},$$

$$c_{13} = \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{1709}{720},$$

$$c_{14} = \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{61}{360},$$

$$c_{22} = \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{2551}{360},$$

$$c_{23} = \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{1709}{720},$$

$$c_{24} = \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{61}{360},$$

$$c_{33} = \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{2033}{180},$$

$$c_{34} = \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{2357}{720},$$

$$c_{44} = \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{521}{120},$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{-54894540x_1 - 54894540x_2 + 238278060x_3 - 92542140}{324135871},$$

$$M_X = \max_P \theta_P = \frac{567397500}{324135871} > 1.$$

A.3.106. R_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{107}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{31}{20}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{31}{20}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{269}{60}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{157}{60}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{1471}{180}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{2989}{720}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{79}{30}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{5}{16}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{1471}{180}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{79}{30}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{5}{16}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{2929}{360}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{2483}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{731}{180},
\end{aligned}$$

$$\begin{aligned}
\theta_P(x_1, x_2, x_3, x_4) &= \frac{451036894560x_1 + 451036894560x_2 + 701882402340x_3}{2028544442569} \\
&\quad + \frac{-984670344060x_4 - 532451679296}{2028544442569},
\end{aligned}$$

$$M_X = \max_P \theta_P = \frac{2758057258564}{2028544442569} > 1.$$

A.3.107. R_3 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{305}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{23}{15}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{23}{15}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{23}{10}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{23}{60}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{95}{12}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{943}{240}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{463}{240}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{247}{240}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{95}{12}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{463}{240}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{247}{240}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{17}{4}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{47}{240}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{659}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{91335300x_1 + 91335300x_2 + 142965700x_3 - 47914704}{369479321},$$

$$M_X = \max_P \theta_P = \frac{277721596}{369479321} < 1.$$

A.3.108.

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{155}{12}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{19}{12}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{7}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{3}{2}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{19}{12}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{1417}{360}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{2093}{720}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{253}{720}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{1}{9}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{2749}{360}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{1741}{720}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{71}{90}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{997}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{253}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{1417}{360},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{74703180x_1 - 57721380x_3 - 74703180x_4 - 25017456}{153124261},$$

$$M_X = \max_P \theta_P = \frac{182110284}{153124261} > 1.$$

A.3.109. U_1 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{77}{6}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{10}{3}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{5}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{5}{3}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{5}{3}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{19}{5}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{19}{10}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{19}{10}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{19}{10}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{107}{30}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{19}{20}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{19}{20}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{599}{90}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{19}{20}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{599}{90},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{3850x_1 - 1000}{3389},$$

$$M_X = \max_P \theta_P = \frac{2850}{3389} < 1.$$

A.3.110. U_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{149}{12}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{5}{2}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{5}{4}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{7}{6}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{11}{6}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{107}{30}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{107}{60}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{41}{45}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{239}{180}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{69}{20}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{41}{90}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{239}{180}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{238}{45}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{197}{90}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{1387}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{907410x_1 - 409005x_3 - 221130}{924382},$$

$$M_X = \max_P \theta_P = \frac{1095285}{924382} > 1.$$

A.3.111. U_3 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{149}{12}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{5}{2}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{5}{2}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{5}{4}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{5}{4}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{107}{30}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{19}{10}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{107}{60}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{19}{20}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{107}{30}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{19}{20}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{107}{60}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{1109}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{19}{40}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{1109}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{11175x_1 + 11175x_2 - 4500}{19936},$$

$$M_X = \max_P \theta_P = \frac{1275}{1424} < 1.$$

A.3.112. U_4 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 12, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = -2, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 1, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{10}{3}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{5}{3}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{10}{3}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = 4, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -2, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = 6,
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{-6x_3 - 1}{11},$$

$$M_X = \max_P \theta_P = \frac{5}{11} < 1.$$

A.3.113. U_5 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 12, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{10}{3}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{5}{3}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{10}{3}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = 4, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = 4,
\end{aligned}$$

$$\begin{aligned}
\theta_P(x_1, x_2, x_3, x_4) &= 0, \\
M_X &= \max_P \theta_P = 0 < 1.
\end{aligned}$$

A.3.114. U_6 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = 12, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{5}{3}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{5}{6}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{5}{6}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{10}{3}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{5}{3}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{5}{3}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{10}{3}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{5}{6}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{17}{3}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = 4,
\end{aligned}$$

$$\begin{aligned}
\theta_P(x_1, x_2, x_3, x_4) &= \frac{36x_1 - 5}{67}, \\
M_X &= \max_P \theta_P = \frac{31}{67} < 1.
\end{aligned}$$

A.3.115. U_7 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{139}{12}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{5}{6}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{5}{3}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{5}{12}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{5}{12}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{31}{10}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{5}{3}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{31}{20}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{43}{60}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{10}{3}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{5}{6}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{5}{6}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{931}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{43}{120}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{931}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{139x_2 - 20}{258},$$

$$M_X = \max_P \theta_P = \frac{119}{258} < 1.$$

A.3.116. U_8 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{67}{6}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{43}{15}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = \frac{43}{30}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{43}{30}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{43}{30}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{31}{10}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{43}{60}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{43}{60}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{421}{90}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{43}{60}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{421}{90},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = 0,$$

$$M_X = \max_P \theta_P = 0 < 1.$$

A.3.117. \tilde{V}^4 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{307}{24}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{13}{10}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{13}{10}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{13}{10}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{13}{10}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{743}{180}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{263}{720}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{263}{720}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{263}{720}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{743}{180}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{263}{720}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{263}{720}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{743}{180}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{263}{720}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{743}{180},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{1436760x_1 + 1436760x_2 + 1436760x_3}{2766841} + \frac{1436760x_4 - 584064}{2766841},$$

$$M_X = \max_P \theta_P = \frac{737568}{395263} > 1.$$

A.3.118. V^4 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{115}{12}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{263}{90}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{263}{360}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{263}{360}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{263}{360}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{263}{90}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{263}{360}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{263}{360}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{263}{90}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{263}{360}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{263}{90},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = 0,$$

$$M_X = \max_P \theta_P = 0 < 1.$$

A.3.119. $dP_7 \times dP_7$.

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{49}{4}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = -\frac{7}{6}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{7}{6}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{7}{6}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{7}{6}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{91}{24}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{35}{48}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{1}{9}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1}{9}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{91}{24}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{1}{9}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{1}{9}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{91}{24}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{35}{48}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{91}{24},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = -\frac{168x_1 + 168x_2 + 168x_3 + 168x_4 + 64}{409},$$

$$M_X = \max_P \theta_P = \frac{608}{409} > 1.$$

A.3.120. $dP_7 \times dP_6$.

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{21}{2}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = -1, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -1, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = -1, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{13}{4}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{5}{8}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{13}{4}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{35}{12}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{35}{24}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{35}{12},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = -\frac{168x_1 + 168x_2 + 32}{409},$$

$$M_X = \max_P \theta_P = \frac{304}{409} < 1.$$

A.3.121. $dP_6 \times dP_6$.

$$\begin{aligned}
b_0 &= \text{vol}(P) = 9, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{5}{2}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{5}{4}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{5}{2}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = 0, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = 0, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{5}{2}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{5}{4}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{5}{2},
\end{aligned}$$

$$\begin{aligned}
\theta_P(x_1, x_2, x_3, x_4) &= 0, \\
M_X &= \max_P \theta_P = 0 < 1.
\end{aligned}$$

A.3.122. Z_1 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{161}{12}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = \frac{31}{30}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{31}{15}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{31}{15}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{31}{15}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{421}{72}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{217}{80}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{145}{36}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{1069}{720}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{217}{40}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{1831}{720}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{1831}{720}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{145}{18}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{1069}{360}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{145}{18},
\end{aligned}$$

$$\begin{aligned}
\theta_P(x_1, x_2, x_3, x_4) &= \frac{598920x_3 + 598920x_4 - 184512}{1289443}, \\
M_X &= \max_P \theta_P = \frac{1013328}{1289443} < 1.
\end{aligned}$$

A.3.123. Z_2 .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{109}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = -\frac{19}{12}, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = \frac{19}{6}, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = \frac{9}{10}, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = \frac{9}{10}, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{46}{9}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{303}{80}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{1351}{720}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{61}{40}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{303}{40}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{253}{720}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{253}{720}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{2327}{360}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{257}{72}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{2327}{360},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = \frac{1384138680x_2 + 1120256220x_4 + 1120256220x_3}{2592694163} - \frac{469692992}{2592694163},$$

$$M_X = \max_P \theta_P = \frac{3418840588}{2592694163} > 1.$$

A.3.124. W .

$$\begin{aligned}
b_0 &= \text{vol}(P) = \frac{83}{8}, & b_1 &= \int_P x_1 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_2 &= \int_P x_2 dx_1 dx_2 dx_3 dx_4 = 0, & b_3 &= \int_P x_3 dx_1 dx_2 dx_3 dx_4 = 0, \\
b_4 &= \int_P x_4 dx_1 dx_2 dx_3 dx_4 = 0, & c_{11} &= \int_P x_1^2 dx_1 dx_2 dx_3 dx_4 = \frac{463}{120}, \\
c_{12} &= \int_P x_1 x_2 dx_1 dx_2 dx_3 dx_4 = -\frac{463}{240}, & c_{13} &= \int_P x_1 x_3 dx_1 dx_2 dx_3 dx_4 = -\frac{601}{360}, \\
c_{14} &= \int_P x_1 x_4 dx_1 dx_2 dx_3 dx_4 = \frac{601}{720}, & c_{22} &= \int_P x_2^2 dx_1 dx_2 dx_3 dx_4 = \frac{463}{120}, \\
c_{23} &= \int_P x_2 x_3 dx_1 dx_2 dx_3 dx_4 = \frac{601}{720}, & c_{24} &= \int_P x_2 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{601}{360}, \\
c_{33} &= \int_P x_3^2 dx_1 dx_2 dx_3 dx_4 = \frac{463}{120}, & c_{34} &= \int_P x_3 x_4 dx_1 dx_2 dx_3 dx_4 = -\frac{463}{240}, \\
c_{44} &= \int_P x_4^2 dx_1 dx_2 dx_3 dx_4 = \frac{463}{120},
\end{aligned}$$

$$\theta_P(x_1, x_2, x_3, x_4) = 0,$$

$$M_X = \max_P \theta_P = 0 < 1.$$

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE DIVISION II, TOKYO UNIVERSITY OF SCIENCE, 1-3 KAGURAZAKA, SHINJUKU-KU, TOKYO 162-8601, JAPAN

Email address: nitta@rs.tus.ac.jp

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE DIVISION I, TOKYO UNIVERSITY OF SCIENCE, 1-3 KAGURAZAKA, SHINJUKU-KU, TOKYO 162-8601, JAPAN

Email address: saito@rs.tus.ac.jp

FACULTY OF EDUCATION, KAGAWA UNIVERSITY, 1-1 SAIWAI-CHO, TAKAMATSU 760-8521, JAPAN

Email address: yotsutani.naoto@kagawa-u.ac.jp