

A SHORT NOTE ON UNDIRECTED FITCH GRAPHS

MANUELA GEISS, MARC HELLMUTH, YANGJING LONG, AND PETER F. STADLER

ABSTRACT. The symmetric version of Fitch’s xenology relation coincides with class of complete multipartite graph and thus cannot convey any non-trivial phylogenetic information.

Fitch graphs [4] form a class of directed graphs that is derived from rooted, $\{0, 1\}$ -edge-labeled trees T in the following manner: The vertices of the Fitch graph are the leaves of T . Two distinct leaves x and y of T are connected by an arc (x, y) from x to y if and only if there is at least one edge with label 1 on the (unique) path in T that connects the least common ancestor $\text{lca}(x, y)$ of x and y with y . Fitch graphs model “xenology”, an important binary relation among genes introduced by Walter M. Fitch [2]. Interpreting T as a phylogenetic tree and 1-edges as horizontal gene transfer events, the arc (x, y) in the Fitch graph encodes the fact that y is xenologous with respect to x . The directed Fitch graphs are the topic of Ref. [4], which among other results proves a characterization in terms of eight forbidden induced subgraphs.

It is natural to consider the symmetrized version of this relationship, i.e., to interpret $\{x, y\}$ as a xenologous pair whenever the evolutionary history separated x and y by at least one horizontal transfer event. In mathematical terms, this idea is captured by

Definition 1. *Let T be a rooted tree with leaf set X and let $\lambda : E(T) \rightarrow \{0, 1\}$. Then the undirected Fitch graph F explained by (T, λ) has vertex set X and edges $\{x, y\} \in E(F)$ if and only if the (unique) path from x to y in T contains at least one edge e with $\lambda(e) = 1$. A graph F is an undirected Fitch graph if and only if it is explained in this manner by some edge-labeled rooted tree (T, λ) .*

Undirected Fitch graphs are closely related to their directed counterparts. Since the path \wp connecting two leaves x and y is unique and contains their last common ancestor $\text{lca}(x, y)$, there is a 1-edge along \wp if and only if there is 1-edge on the path between x and $\text{lca}(x, y)$ or between $\text{lca}(x, y)$ and y . The undirected Fitch graph is therefore the underlying undirected graph of the directed Fitch graph, i.e., it is obtained from the directed version by ignoring the direction of the arcs.

The undirected Fitch graphs form a heritable family, i.e., if F is a an undirected Fitch graph, so are all its induced subgraphs. This is an immediate consequence of the fact that directed Fitch graphs are a heritable family of digraphs [4]. The fact can also be obtained directly by considering the restriction of T to a subset of leaves. This obviously does not affect the paths or their labeling between the remaining vertices.

Clearly F does not depend on which of the non-leaf vertices in T is the root. Furthermore, a vertex v with only two neighbors and its two incident edges e' and e'' can be replaced by a single edge e . The new edge is labeled $\lambda(e) = 0$ if both $\lambda(e') = \lambda(e'') = 0$, and $\lambda(e) = 1$ otherwise. These operations do not affect the undirected Fitch graph. Hence, we can replace the rooted tree T by an unrooted tree in Def. 1 and assume that all non-leaf edges have at least degree 3. To avoid trivial cases we assume throughout that T has at least two leaves and hence a Fitch graph has at least two vertices.

Lemma 1. *If G is an undirected Fitch graph, then F does not contain $K_1 \cup K_2$ as an induced subgraph. In particular every undirected Fitch graph is a complete multipartite graph.*

Date: December 8, 2021.

1991 Mathematics Subject Classification. 05C75, 05C05, 92B10.

Key words and phrases. Labeled trees, forbidden subgraphs, phylogenetics, xenology, Fitch graph.

This work is supported in part by the BMBF-funded project “Center for RNA-Bioinformatics” (031A538A, de.NBI/RBC).




FIGURE 1. The complete multipartite graph $K_{3,2,1,1}$ is the Fitch graph explained by the tree $T[3,2,1,1]$ with edge labeling λ^* shown with bold numbers **0** and **1**.

Proof. There is a single unrooted tree with three leaves, namely the star S_3 , which admits four non-isomorphic $\{0,1\}$ -edge labelings defined by the number N of 1-edges. The undirected Fitch graphs F_N are easily obtained. In the absence of 1-edges, $F_0 = \overline{K_3}$ is edge-less. For $N = 2$ and $N = 3$ there is 1-edge along the path between any two leaves, i.e., $F_2 = F_3 = K_3$. For $N = 1$ one leaf is connected to the other two by a path in S_3 with an 1-edge; the path between the latter two leaves consists of two 0-edges, hence $F_1 = P_3$, the path of length two. Hence, only three of the four possible undirected graphs on three vertices can be realized, while $K_1 \cup K_2$ is not an undirected Fitch graph. By heredity, $K_1 \cup K_2$ is therefore a forbidden induced subgraph for the class of undirected Fitch graphs. Finally, it is well known that the class of graphs that do not contain $K_1 \cup K_2$ as an induced subgraph are exactly the complete multipartite graphs, see e.g. [6]. \square

In order to show that forbidding $K_1 \cup K_2$ is also sufficient, we explicitly construct the edge-labeled trees necessary to explain complete multipartite graphs. We start by recalling that each complete multipartite graph K_{n_1, \dots, n_k} is determined by its independent sets V_1, \dots, V_k with $|V_i| = n_i$ for $1 \leq i \leq k$. By definition, $\{x, y\} \in E(K_{n_1, \dots, n_k})$ if and only if $x \in V_i$ and $y \in V_j$ with $i \neq j$. In particular, therefore, K_{n_1, \dots, n_k} with at least two vertices is connected if and only if $k \geq 2$. The complete 1-partite graphs are the edge-less graphs $\overline{K_n}$.

Since $K_1 \cup K_2$ is an induced subgraph of the path on four vertices P_4 , any graph G that does not contain $K_1 \cup K_2$ as an induced subgraph must be P_4 -free, i.e., a cograph [1]. Cographs are associated with vertex-labeled trees known as cotrees, which in turn are a special case of modular decomposition trees [3]. The cotrees of connected multipartite graphs have a particularly simple shape, illustrated without the vertex labels in Fig. 1. The cotree has a root labeled “1” and all inner vertices labeled “0”. Here we do not need the connection between cographs and their cotrees, however. Therefore we introduce these trees together with an edge-labeling that is useful for our purposes in the following

Definition 2. For $k = 1$, $T[n]$ is the star graph S_n with n leaves. For $k \geq 2$, the tree $T[n_1, \dots, n_k]$ has a root r with k children c_i , $1 \leq i \leq k$. The vertex c_i is a leaf if $|V_i| = n_i = 1$ and has exactly n_i children that are leaves if $|V_i| = n_i \geq 2$. For $k = 1$ all edges e of $T[n]$ are labeled $\lambda^*(e) = 0$. For $k \geq 2$ we set $\lambda^*({r, c_i}) = 1$ for $1 \leq i \leq k$ and $\lambda^*(e) = 0$ for all edges not incident to the root.

Now we can prove our main result:

Theorem 1. A graph G is an undirected Fitch graph if and only if it is a complete multipartite graph. In particular, K_{n_1, \dots, n_k} is explained by $(T[n_1, \dots, n_k], \lambda^*)$.

Proof. Lemma 1 implies that an undirected Fitch graph is a complete multipartite graph. To show the converse, we fix an arbitrary complete multipartite graph $G = K_{n_1, \dots, n_k}$ and find an edge-labeled rooted tree (T, λ^*) that explains G .

For $k = 1$ it is trivial that $(T[n], \lambda^*)$ explains $\overline{K_n}$.

For $k \geq 2$ consider the tree $T[n_1, \dots, n_k]$ with edge labeling λ^* and let F be the corresponding Fitch graph. The leaf-set of $T[n_1, \dots, n_k]$ partitioned into exactly k subsets L_1, \dots, L_k defined by (a) singletons adjacent to the root and (b) subsets comprising at least two leaves adjacent to the same child c_i of the root. Furthermore, we can order the leaf sets so that $|L_i| = n_i$. By construction, all vertices within a leaf set L_i are connected by a path that does not run through the root and thus, contains only 0-edges, if $|L_i| > 1$ and no edge, otherwise. The L_i are independent sets in F .




FIGURE 2. The non-isomorphic trees (T, λ^*) , (T', λ^*) , $(T[2, 2, 1], \lambda^*)$, and $(T^*[2, 2, 1], \lambda^*)$ all explain the same complete multipartite graph $K_{2,2,1}$. Three of these trees have the smallest possible number (7) of vertices and thus are minimal. These can be obtained from the tree $(T[2, 2, 1], \lambda^*)$ specified in Definition 1 by contracting of one of its inner 1-edge and possibly re-routing the resulting tree.

On the other hand any two leaves $x \in L_i$ and $y \in L_j$ with $i \neq j$ are connected only by path through the root, which contains two 1-edges. Thus x and y are connected by an edge in F . Thus F is a complete multipartite graph of the form $K_{|L_1|, \dots, |L_k|} = K_{n_1, \dots, n_k}$. Since K_{n_1, \dots, n_k} is explained by $(T[n_1, \dots, n_k], \lambda^*)$ for all $n_i \geq 1$ and all $k \geq 2$, and K_n is explained by $(T[n], \lambda^*)$, we conclude that every complete multipartite graph is a Fitch graph. \square

Complete multipartite graphs $G = (V, E)$ obviously can be recognized in $O(|V|^2)$ time e.g., by checking that its complement is a disjoint union of complete graphs), and even in $O(|V| + |E|)$ time by explicitly constructing its modular decomposition tree [5]. Given the tree $T[n_1, \dots, n_k]$, the canonical edge labeling λ^* is then assigned in $O(|V|)$ time.

A tree (T, λ) that explains a Fitch graph F is *minimum* if it has the smallest number of vertices among all trees that explain F . In this case, (T, λ) is also *least-resolved*, i.e., the contraction of any edge in (T, λ) results in a tree that does not explain F . Not surprisingly, the tree $T[n_1, \dots, n_k]$ is almost minimum in most, and minimum in some cases: Since the vertices of the Fitch graph must correspond to leaves of the tree, $T[n_1, \dots, n_k]$ is necessarily minimum whenever it is a star, i.e., for $T[n]$ and $T[1, \dots, 1]$. In all other cases, its only potentially “superfluous” part is its root. Indeed, exactly one of the edges connecting the root with a non-leaf neighbor can be contracted without changing the corresponding Fitch graph. It is clear that this graph is minimal: The leaf sets L_i must be leaves of an induced subtree without an intervening 1-edge. Having all vertices of L_i adjacent to the same vertex is obviously the minimal choice. Since the L_i must be separated from all other leaves by a 1-edge, at least one neighbor of c_i must be a 1-edge. Removing all leaves incident to a 0-edge results in a tree with at least k vertices that must contain at least $k - 1$ 1-edges, since every path between leaves in this tree must contain a 1-edge. The contraction of exactly one of the k 1-edges incident to the root r in $T[n_1, \dots, n_k]$ indeed already yields a minimal tree. In general, the minimal trees are not unique, see Fig. 2.

The practical implication of Thm. 1 in the context of phylogenetic combinatorics is that the mutual xenology relation cannot convey any interesting phylogenetic information: Since the undirected Fitch graphs are exactly the complete multipartite graphs, which in turn are completely defined by their independent sets, the only insight we can gain by considering mutual xenology is the identification of the maximal subsets of taxa that have not experienced any horizontal transfer events among them.

REFERENCES

[1] D. G. Corneil, H. Lerchs, and L. Steward Burlingham. Complement reducible graphs. *Discr. Appl. Math.*, 3:163–174, 1981.
 [2] W M Fitch. Homology a personal view on some of the problems. *Trends Genet.*, 16:227–231, 2000.
 [3] T Gallai. Transitiv orientierbare Graphen. *Acta Math. Acad. Sci. Hungar.*, 18:1967, 25-66.
 [4] Manuela Geiß, John Anders, Peter Stadler, Nicolas Wieseke, and Marc Hellmuth. Reconstructing gene trees from Fitch’s xenology relation. 2017. arXiv:1711.02152.
 [5] Ross M. McConnell and Jeremy P. Spinrad. Modular decomposition and transitive orientation. *Discrete Math*, 201:189–241, 1999.
 [6] Igor E Zverovich. Near-complete multipartite graphs and forbidden induced subgraphs. *Discrete Math*, 207:257–262, 1999.

BIOINFORMATICS GROUP, DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITÄT LEIPZIG, HÄRTELSTRASSE 16-18, D-04107 LEIPZIG, GERMANY

E-mail address: manuela@bioinf.uni-leipzig.de

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF GREIFSWALD, WALTHER-RATHENAU-STRASSE 47, D-17487 GREIFSWALD, GERMANY; CENTER FOR BIOINFORMATICS, SAARLAND UNIVERSITY, BUILDING E 2.1, P.O. Box 151150, D-66041 SAARBRÜCKEN, GERMANY

E-mail address: mhellmuth@mailbox.org

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF GREIFSWALD, WALTHER-RATHENAU-STRASSE 47, D-17487 GREIFSWALD, GERMANY

E-mail address: yjlong@sjtu.edu.cn

BIOINFORMATICS GROUP, DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITÄT LEIPZIG, HÄRTELSTRASSE 16-18, D-04107 LEIPZIG, GERMANY; INTERDISCIPLINARY CENTER FOR BIOINFORMATICS, THE GERMAN CENTRE FOR INTEGRATIVE BIODIVERSITY RESEARCH (iDiv) HALLE-JENA-LEIPZIG THE COMPETENCE CENTER FOR SCALABLE DATA SERVICES AND SOLUTIONS DRESDEN-LEIPZIG THE LEIPZIG RESEARCH CENTER FOR CIVILIZATION DISEASES, AND THE CENTRE FOR BIOTECHNOLOGY AND BIOMEDICINE AT LEIPZIG UNIVERSITY; THE MAX PLANCK INSTITUTE FOR MATHEMATICS IN THE SCIENCES, LEIPZIG, GERMANY; THE INSTITUTE FOR THEORETICAL CHEMISTRY, UNIVERSITY OF VIENNA, VIENNA, AUSTRIA; THE CENTER OF NONCODING RNA IN HEALTH AND TECHNOLOGY (RTH) AT THE UNIVERSITY OF COPENHAGEN; AND THE SANTA FE INSTITUTE, SANTA FE, NM

E-mail address: studla@bioinf.uni-leipzig.de