

# On the size-Ramsey number of tight paths

Linyuan Lu <sup>\*</sup>      Zhiyu Wang <sup>†</sup>

March 14, 2022

## Abstract

For any  $r \geq 2$  and  $k \geq 3$ , the  $r$ -color size-Ramsey number  $\hat{R}(\mathcal{G}, r)$  of a  $k$ -uniform hypergraph  $\mathcal{G}$  is the smallest integer  $m$  such that there exists a  $k$ -uniform hypergraph  $\mathcal{H}$  on  $m$  edges such that any coloring of the edges of  $\mathcal{H}$  with  $r$  colors yields a monochromatic copy of  $\mathcal{G}$ . Let  $\mathcal{P}_{n,k-1}^{(k)}$  denote the  $k$ -uniform tight path on  $n$  vertices. Dudek, Fleur, Mubayi and Ródl showed that the size-Ramsey number of tight paths  $\hat{R}(\mathcal{P}_{n,k-1}^{(k)}, 2) = O(n^{k-1-\alpha}(\log n)^{1+\alpha})$  where  $\alpha = \frac{k-2}{\binom{k-1}{2}+1}$ . In this paper, we improve their bound by showing that  $\hat{R}(\mathcal{P}_{n,k-1}^{(k)}, r) = O(r^k(n \log n)^{k/2})$  for all  $k \geq 3$  and  $r \geq 2$ .

## 1 Introduction

Given two simple graphs  $G$  and  $H$  and a positive integer  $r$ , say that  $H \rightarrow (G)_r$  if every  $r$ -edge-coloring of  $H$  results in a monochromatic copy of  $G$  in  $H$ . In this notation, the Ramsey number  $R(G)$  of  $G$  is the minimum  $n$  such that  $K_n \rightarrow (G)_2$ . The *size-Ramsey number*  $\hat{R}(G, r)$  of  $G$  is defined as the minimum number of edges in a graph  $H$  such that  $H \rightarrow (G)_r$ , i.e.

$$\hat{R}(G, r) = \min\{|E(H)| : H \rightarrow (G)_r\}.$$

When  $r = 2$ , we ignore  $r$  and simply use  $\hat{R}(G)$ .

Size-Ramsey number was first studied by Erdős, Faudree, Rousseau and Schelp [8] in 1978. By the definition of  $R(G)$ , we have

$$\hat{R}(G) \leq \binom{R(G)}{2}.$$

Chvátal (see, e.g. [8]) showed that this bound is tight for complete graphs, i.e.  $\hat{R}(K_n) = \binom{R(K_n)}{2}$ . Answering a question of Erdős [9], Beck [3] showed by a probabilistic construction that

$$\hat{R}(P_n) = O(n).$$

---

<sup>\*</sup>University of South Carolina, Columbia, SC 29208, (lu@math.sc.edu). This author was supported in part by NSF grant DMS-1600811.

<sup>†</sup>University of South Carolina, Columbia, SC 29208, (zhiyuw@math.sc.edu).

Alon and Chung [1] gave an explicit construction of a graph  $G$  with  $O(n)$  edges such that  $G \rightarrow P_n$ . Recently, Dudek and Prałat [6] provided a simple alternative proof for this result (See also [10]). The best upper bound  $\hat{R}(P_n) \leq 74n$  is due to Dudek and Prałat [7] by considering a random 27-regular graph of a proper order.

Dudek, Fleur, Mubayi, and Ródl [11] first initiated the study of size-Ramsey number in hypergraphs. A  $k$ -uniform hypergraph  $\mathcal{G}$  on a vertex set  $V(\mathcal{G})$  is a family of  $k$ -element subsets (called edges) of  $V(\mathcal{G})$ . We use  $E(\mathcal{G})$  to denote the edge set. Given  $k$ -uniform hypergraphs  $\mathcal{G}$  and  $\mathcal{H}$ , we say that  $\mathcal{H} \rightarrow (\mathcal{G})_r$  if every  $r$ -edge-coloring of  $\mathcal{H}$  results in a monochromatic copy of  $\mathcal{G}$  in  $\mathcal{H}$ . Define the *size-Ramsey number*  $\hat{R}(\mathcal{G}, r)$  of a  $k$ -uniform hypergraph  $\mathcal{G}$  as

$$\hat{R}(\mathcal{G}, r) = \min\{|E(\mathcal{H})| : \mathcal{H} \rightarrow (\mathcal{G})_r\}.$$

When  $r = 2$ , we simply use  $\hat{R}(\mathcal{G})$  for the ease of reference.

Given integers  $1 \leq l < k$  and  $n \equiv l \pmod{k-l}$ , an  $l$ -path  $\mathcal{P}_{n,l}^{(k)}$  is a  $k$ -uniform hypergraph with vertex set  $[n]$  and edge set  $\{e_1, \dots, e_m\}$ , where  $e_i = \{(i-1)(k-l) + 1, (i-1)(k-l) + 2, \dots, (i-1)(k-l) + k\}$  and  $m = \frac{n-l}{k-l}$ , i.e. the edges are intervals of length  $k$  in  $[n]$  and consecutive edges intersect in exactly  $l$  vertices. A  $\mathcal{P}_{n,1}^{(k)}$  is commonly referred as a *loose* path and a  $\mathcal{P}_{n,k-1}^{(k)}$  is called a *tight* path.

Dudek, Fleur, Mubayi and Ródl [11] showed that when  $l \leq \frac{k}{2}$ , the size-Ramsey number of a path  $\mathcal{P}_{n,l}^{(k)}$  can be easily reduced to the graph case. In particular, they showed that if  $1 \leq l \leq \frac{k}{2}$ , then

$$\hat{R}(\mathcal{P}_{n,l}^{(k)}) \leq \hat{R}(P_n) = O(n).$$

For tight paths, they showed in the same paper that for fixed  $k \geq 3$ ,

$$\hat{R}(\mathcal{P}_{n,k-1}^{(k)}) = O(n^{k-1-\alpha}(\log n)^{1+\alpha}),$$

where  $\alpha = (k-2)/\binom{k-2}{2} + 1$ . Observe that  $\hat{R}(\mathcal{P}_{n,l}^{(k)}) \leq \hat{R}(\mathcal{P}_{n,k-1}^{(k)})$ . Thus any upper bound on the size-Ramsey number of tight paths is also an upper bound for other  $l$ -path  $\mathcal{P}_{n,l}^{(k)}$ .

Motivated by their approach, we use a different probabilistic construction and improve the upper bound to  $O((n \log n)^{k/2})$ . In particular, we show the following result on the multi-color size-Ramsey number of tight paths in hypergraphs:

**Theorem 1.** *For any fixed  $k \geq 3$ , any  $r \geq 2$ , and sufficiently large  $n$ , we have*

$$\hat{R}(\mathcal{P}_{n,k-1}^{(k)}, r) = O\left(r^k (n \log n)^{\frac{k}{2}}\right).$$

## 2 Proof of Theorem 1

The approach of our proof is inspired by Dudek, Fleur, Mubayi and Rödl's approach in their proof of Theorem 2.8 in [11]. In their proof, they constructed their hypergraph by setting edges to be the  $k$ -cliques of an Erdős-Rényi random graph. Then they use a greedy algorithm to show that the number of edges of each color is smaller than  $\frac{1}{r}$  fraction of the total number of edges, which gives a contradiction. Motivated by their approach, we use the same greedy algorithm but a different probabilistic construction of the hypergraph. Instead of using  $k$ -cliques of an Erdős-Rényi random graph as edges, we use  $k$ -cycles of a random  $C_k$ -colorable graph (which will be defined later) as edges.

Throughout the paper, we will use the following version of Chernoff inequalities for the binomial random variables  $X \sim \text{Bin}(n, p)$  (for details, see, e.g. [4]):

$$\Pr(X \leq E(X) - \lambda) \leq \exp\left(-\frac{\lambda^2}{2E(X)}\right). \quad (1)$$

$$\Pr(X \geq E(X) + \lambda) \leq \exp\left(-\frac{\lambda^2}{2(E(X) + \lambda/3)}\right). \quad (2)$$

We follow a similar notation as [11]. A graph  $G$  is  $C_k$ -colorable if there is a graph homomorphism  $\pi$  mapping  $G$  to the cycle  $C_k$ . That is,  $V(G)$  can be partitioned into  $k$ -parts  $V_1 \cup V_2 \cup \dots \cup V_k$  so that  $E(G) \subseteq \bigcup_{i=1}^k E(V_i, V_{i+1})$  with  $V_{k+1} = V_1$  and  $E(V_i, V_{i+1})$  denoting the set of edges between a vertex in  $V_i$  and a vertex in  $V_{i+1}$ . For such a graph  $G$ , we say a  $k$ -cycle  $C$  in  $G$  is *proper* if it intersects each  $V_i$  by exactly one vertex. For  $1 \leq l \leq k-1$ , we say a path  $P_l$  of  $l$  vertices in  $G$  is *proper* if it intersects each  $V_i$  by at most one vertex. Let  $\mathcal{T}_{k-1}(G)$  denote the set of all proper  $(k-1)$ -paths in  $G$ . Let  $\mathcal{B} \subseteq \mathcal{T}_{k-1}(G)$  be a family of pairwise vertex-disjoint proper  $(k-1)$ -paths. Let  $t_{\mathcal{B}}$  be the total number of proper  $k$ -cycles in  $G$  that extend some  $B \in \mathcal{B}$ . For  $A \subseteq V$ , define  $y_{A, \mathcal{B}}$  as the number of proper  $k$ -cycles in  $G$  that extend a proper  $(k-1)$ -path  $B \in \mathcal{B}$  with a vertex  $v \in A \cup \bigcup_{B \in \mathcal{B}} V(B)$ . Given  $C \subseteq V(G)$ , we use  $z_C$  to denote the number of proper  $k$ -cycles in  $G$  that intersect  $C$ . We use  $t_k$  to denote the total number of proper  $k$ -cycles in  $G$ .

We say an event in a probability space holds *a.a.s.* (aka, *asymptotically almost surely*) if the probability that it holds tends to 1 as  $n$  goes to infinity. Finally, we use  $\log n$  to denote natural logarithms.

**Proposition 1.** *For every  $r \geq 2$ ,  $k \geq 3$ , and sufficiently large  $n$ , there exists a  $C_k$ -colorable graph  $G = (V, E)$  satisfying the following:*

- (i) *For every  $\mathcal{B}$  consisting of  $n$  pairwise vertex-disjoint proper  $(k-1)$ -paths, and every  $A \subseteq V \setminus \bigcup_{B \in \mathcal{B}} V(B)$  with  $|A| \leq n$ , we have*

$$y_{A, \mathcal{B}} < \frac{1}{2kr} t_{\mathcal{B}}.$$

(ii) For every  $C \subseteq V$  with  $|C| \leq (k-1)n$ , we have

$$z_C < \frac{t_k}{2r}.$$

(iii) The total number of proper  $k$ -cycles satisfies

$$t_k = O(r^k(n \log n)^{k/2}).$$

*Proof.* Set  $c = 16k^2r$  and  $p = \frac{\sqrt{\log n}}{\sqrt{n}}$ . Consider the following random  $C_k$ -colorable graph  $G$ . Let  $V(G) = V_1 \cup V_2 \cup \dots \cup V_k$  be the disjoint union of  $k$  sets. Each  $V_i$  (for  $1 \leq i \leq k$ ) has the same size  $cn$ . For any pair of vertices  $\{u, v\}$  in two consecutive parts, i.e., there is an  $i \in [k]$ , such that  $u \in V_i$  and  $v \in V_{i+1}$  (with the convention  $V_{k+1} = V_1$ ), add  $uv$  as an edge of  $G$  with probability  $p$  independently. There is no edge inside each  $V_i$  or between two non-consecutive parts.

We will show that this random  $C_k$ -colorable graph  $G$  satisfies a.a.s. (i)–(iii).

First we show that  $G$  a.a.s. satisfies (i). For a fixed family  $\mathcal{B}$  of  $n$  pairwise vertex-disjoint proper  $(k-1)$ -paths, we would like to give a lower bound of  $t_{\mathcal{B}}$ . For each proper  $(k-1)$ -path  $B \in \mathcal{B}$ , there are  $cn$  vertices that can extend  $B$  into a proper  $k$ -cycle, each with probability  $p^2$  independently. Thus, we have  $t_{\mathcal{B}} \sim \text{Bin}(cn^2, p^2)$  with

$$E[t_{\mathcal{B}}] = cn^2p^2 = cn \log n = 16k^2rn \log n.$$

Applying Chernoff inequality, we have

$$\begin{aligned} \Pr \left( t_{\mathcal{B}} \leq \frac{E[t_{\mathcal{B}}]}{2} \right) &\leq \exp \left( -\frac{1}{8} E[t_{\mathcal{B}}] \right) \\ &= \exp \left( -2k^2rn \log n \right). \end{aligned}$$

Now for fixed  $A \subseteq V \setminus \bigcup_{B \in \mathcal{B}} V(B)$ , we estimate the upper bound of  $y_{A, \mathcal{B}}$ . Without loss of generality, we can assume that  $|A| = n$ . We have  $y_{A, \mathcal{B}} \leq Y \sim \text{Bin}(2n^2, p^2)$ , thus

$$E[Y] = 2n^2p^2 = 2n \log n.$$

Thus if we apply the Chernoff bound (2) with  $\lambda = (2k-1)E[Y]$ , then

$$\begin{aligned} \Pr \left( Y \geq \frac{1}{4kr} E[t_{\mathcal{B}}] \right) &= \Pr (Y \geq 2kE[Y]) \\ &= \Pr (Y \geq E[Y] + \lambda) \\ &\leq \exp \left( -\frac{\lambda^2}{2(E[Y] + \lambda/3)} \right) \\ &\leq \exp \left( -\frac{3(2k-1)^2}{2k+2} n \log n \right). \end{aligned}$$

The number of possible choices of  $\mathcal{B}$  is upper bounded by  $\left(\binom{cn}{n} \cdot n!\right)^k$ . The number of possible choices of  $A$  and  $\mathcal{B}$  is upper bounded by  $\left(\binom{cn}{n, \lceil n/k \rceil} \cdot n!\right)^k \leq \left(\binom{cn}{n, n} \cdot n!\right)^k$ . Stirling approximation of binomial coefficient gives us that

$$\begin{aligned}\log \left( \binom{cn}{n} \cdot n! \right)^k &= (1 + o(1)) (kn \log n), \\ \log \left( \binom{cn}{n, n} \cdot n! \right)^k &= (1 + o(1)) (kn \log n).\end{aligned}$$

Therefore by the union bound, we have

$$\begin{aligned}Pr \left( \bigcup_{\mathcal{B}} \{t_{\mathcal{B}} \leq \frac{E[t_{\mathcal{B}}]}{2}\} \right) &\leq \left( \binom{cn}{n} \cdot n! \right)^k Pr \left( t_{\mathcal{B}} \leq \frac{E[t_{\mathcal{B}}]}{2} \right) \\ &\leq \exp \left( (1 + o(1)) kn \log n - 2k^2 rn \log n \right) \\ &= o(1).\end{aligned}$$

Similarly, we have

$$\begin{aligned}Pr \left( \bigcup_{A, \mathcal{B}} \{y_{A, \mathcal{B}} \geq \frac{1}{4kr} E[t_{\mathcal{B}}]\} \right) &\leq \left( \binom{cn}{n, n} \cdot n! \right)^k Pr \left( Y \geq \frac{1}{4kr} E[t_{\mathcal{B}}] \right) \\ &\leq \exp \left( (1 + o(1)) kn \log n - \frac{3(2k-1)^2}{2k+2} n \log n \right) \\ &= o(1).\end{aligned}$$

In the last step, we observe  $\frac{3(2k-1)^2}{2k+2} > k$  for all  $k \geq 3$ .

Therefore, combining previous inequalities, it follows that for all  $A, \mathcal{B}$  satisfying the condition in (i), we have, a.a.s.,

$$y_{A, \mathcal{B}} < \frac{1}{4kr} E[t_{\mathcal{B}}] \leq \frac{1}{2kr} t_{\mathcal{B}}.$$

This finishes the proof of (i).

Now we will prove that  $G$  satisfies (ii) and (iii) a.a.s.

We will use the Kim-Vu inequality [12] stated as below:

*Let  $H$  be a (weighted) hypergraph with  $V(H) = [n]$ . Edge  $e$  has some weight  $w(e)$ . Suppose  $\{t_i : i \in [n]\}$  is a set of Bernoulli independent random variables with probability  $p$  of being 1. Consider the polynomial*

$$Y_H = \sum_{e \in E(H)} w(e) \prod_{s \in e} t_s.$$

*Furthermore, for a subset  $A$  of  $V(H)$ , define*

$$Y_{H_A} = \sum_{e, A \subseteq e} w(e) \prod_{i \in e \setminus A} t_i.$$

If we define  $E_i(H) = \max_{A \subset V(H), |A|=i} E(Y_{H_A})$ ,  $E(H) = \max_{i \geq 0} E_i(H)$  and  $E'(H) = \max_{i \geq 1} E_i(H)$ , then

$$\Pr \left( |Y_H - E_0(H)| > a_k (E(H)E'(H))^{1/2} \lambda^k \right) = O(\exp(-\lambda + (k-1) \log n)) \quad (3)$$

for any positive number  $\lambda > 1$  and  $a_k = 8^k (k!)^{1/2}$ .

In our context, for a fixed  $v \in V(G)$ , let  $H$  be the  $k$ -uniform hypergraph constructed by the proper  $k$ -cycles of  $G$  containing  $v$ . The edge set of  $H$  is the collection of all  $k$ -tuples  $\{vv_1, v_1v_2, \dots, v_{k-2}v_{k-1}, v_{k-1}v\}$  such that  $vv_1v_2 \dots v_{k-1}v$  is a proper  $k$ -cycle in  $G$  and all edges have weight 1.

Fix  $v \in V(G)$ . we let  $X_v$  denote the number of proper  $k$ -cycles in  $G$  that contain  $v$ . Then it's not hard to see that

$$\begin{aligned} E_0(X_v) &= E(X_v) = (cn)^{k-1} p^k = c^{k-1} n^{\frac{k-2}{2}} (\log n)^{\frac{k}{2}}. \\ E'(X_v) &= (cn)^{k-2} p^{k-1} = c^{k-2} n^{\frac{k-3}{2}} (\log n)^{\frac{k-1}{2}}. \end{aligned}$$

Applying Kim-Vu inequality with  $\lambda = 2(k-1) \log n$ , we get that for each  $v \in V(G)$ ,

$$\Pr \left( |X_v - E_0(X_v)| > a_k (E(X_v)E'(X_v))^{1/2} \lambda^k \right) = O(\exp(-(k-1) \log n)).$$

Observe that  $a_k (E(X_v)E'(X_v))^{1/2} \lambda^k = o(E_0(X_v))$ . Applying union bound for all  $v \in V(G)$ , we obtain that a.a.s that

$$X_v = (1 \pm o(1))(cn)^{k-1} p^k = (1 \pm o(1))c^{k-1} n^{\frac{k}{2}-1} (\log n)^{\frac{k}{2}}.$$

Recall that  $t_k$  denotes the total number of proper  $k$ -cycles in  $G$  and  $z_C$  denotes the number of proper  $k$ -cycles in  $G$  that intersect  $C$ . Suppose  $|C| \leq (k-1)n$ . Then

$$z_C \leq (1 + o(1))(k-1)nc^{k-1} n^{\frac{k}{2}-1} (\log n)^{\frac{k}{2}} = (1 + o(1))(k-1)c^{k-1} (n \log n)^{\frac{k}{2}}.$$

Note that  $t_k = \frac{1}{k} \sum_{v \in V(G)} X_v$ . Thus

$$\begin{aligned} t_k &\geq \frac{1}{k} (1 - o(1))kcn \cdot c^{k-1} n^{\frac{k}{2}-1} (\log n)^{\frac{k}{2}} \\ &\geq (1 - o(1))c^k (n \log n)^{\frac{k}{2}}. \end{aligned}$$

Since  $c = 16k^2r$ , we have that for  $n$  sufficiently large,

$$z_C < \frac{t_k}{2r}.$$

Moreover, similar to the above calculation, we have that a.a.s.,

$$t_k \leq (1 + o(1))c^k (n \log n)^{\frac{k}{2}} = O(r^k (n \log n)^{\frac{k}{2}}).$$

□

Now we will prove the main result. We use the same greedy algorithm approach by Dudek, Fleur, Mubayi and Ródl in [11].

*Proof of Theorem 1:* We show that there exists a  $k$ -uniform hypergraph  $\mathcal{H}$  with  $|E(\mathcal{H})| = O(r^k n^{\frac{k}{2}} (\log n)^{\frac{k}{2}})$  such that any  $r$ -coloring of the edges of  $\mathcal{H}$  yields a monochromatic copy of  $\mathcal{P}_{n,k-1}^{(k)}$ .

Let  $G$  be the graph constructed from Proposition 1 for  $n$  sufficiently large. Let  $\mathcal{H}$  be a  $k$ -uniform hypergraph such that  $V(\mathcal{H}) = V(G)$  and  $E(\mathcal{H})$  be the collection of all proper  $k$ -cycles in  $G$ .

Take an arbitrary  $r$ -coloring of the edges  $\mathcal{H}_0 = \mathcal{H}$  and assume that there is no monochromatic  $\mathcal{P}_{n,k-1}^{(k)}$ . Without loss of generality, suppose the color class with the most number of edges is blue. We will consider the following greedy algorithm:

- (1) Let  $\mathcal{B} = \emptyset$  be a *trash set* of proper  $(k-1)$ -paths in  $G$ . Let  $A$  be a blue tight path in  $\mathcal{H}$  that we will iteratively modify. Throughout the process, let  $U = V(\mathcal{H}) \setminus (V(A) \cup \bigcup_{B \in \mathcal{B}} V(B))$  be the set of *unused* vertices. If at any point  $|\mathcal{B}| = n$ , terminate.
- (2) If possible, choose a blue edge  $v_1 v_2 \cdots v_{k-1} v_k$  from  $U$  and put these vertices into  $A$  and set the pointer to  $v_k$ . Otherwise, if not possible, terminate.
- (3) Suppose the pointer is at  $v_i$  and  $v_{i-k+2}, \dots, v_{i-1}, v_i$  are the last  $k-1$  vertices of the constructed blue path  $A$ . There are two cases:

Case 1: If there exists a vertex  $u \in U$  such that  $v_{i-k+2}, \dots, v_{i-1}, v_i, u$  form a blue edge in  $\mathcal{H}$ , then we *extend*  $P$ , i.e. add  $v_{i+1} = u$  into  $A$ . Set the pointer to  $v_{i+1}$  and restart Step (3).

Case 2: Otherwise, remove the last  $k-1$  vertices from  $A$  and set  $\mathcal{B} = \mathcal{B} \cup \{v_{i-k+2}, \dots, v_{i-1}, v_i\}$ . Set the pointer to  $v_{i-k+1}$ . Now if  $|A| < k$ , then set  $A = \emptyset$  and go to Step (2). Otherwise, restart Step (3).

Note that this procedure will terminate under two circumstances: either  $|\mathcal{B}| = n$  or there is no blue edge in  $U$ .

Let us first consider the case when  $|\mathcal{B}| = n$ , i.e. there are  $n$  pairwise vertex-disjoint proper  $(k-1)$ -paths in  $\mathcal{B}$ . Moreover,  $|A| \leq n$  since there is no blue path of  $n$  vertices. Applying Proposition 1 with sets  $A$  and  $\mathcal{B}$ , we obtain that

$$y_{A,\mathcal{B}} < \frac{1}{2kr} t_{\mathcal{B}}.$$

Observe that every edge of  $\mathcal{H}$  that extends some  $B \in \mathcal{B}$  with a vertex from  $V(\mathcal{H}_0) \setminus \left( V(A) \cup \bigcup_{B \in \mathcal{B}_m} B \right)$  must be non-blue. Therefore, the number of blue edges of  $\mathcal{H}$  that contain some  $B \in \mathcal{B}$  as subgraph is at most  $y_{A,\mathcal{B}}$ .

Consider  $A, \mathcal{B}$  as  $A_0, \mathcal{B}_0$  respectively. Now remove all the blue edges from  $\mathcal{H}_0$  that contain some  $B \in \mathcal{B}_0$  as subgraph and denote the resulting hypergraph

as  $\mathcal{H}_1$ . Perform the greedy procedure again on  $\mathcal{H}_1$ . This will generate a new  $A_1$  and  $\mathcal{B}_1$ . Applying Proposition 1 again, we have  $y_{A_1, \mathcal{B}_1} \leq \frac{1}{2kr} t_{\mathcal{B}_1}$ . Keep repeating the procedure until it is no longer possible. Observe that  $\mathcal{B}_i \cap \mathcal{B}_j = \emptyset$  for  $i \neq j$ .

When the above procedure can not be repeated anymore, we are in the case that  $|\mathcal{B}_m| < n$  for some positive integer  $m$  and there are no more blue edges in  $V(\mathcal{H}) \setminus \bigcup_{B \in \mathcal{B}_m} B$ . In this case,  $A_m = \emptyset$  and all the blue edges remaining in  $\mathcal{H}_m$  have to intersect the set  $C = \bigcup_{B \in \mathcal{B}_m} B$ . By Proposition 1, it follows that

$$z_C < \frac{1}{2r} t_k.$$

Let  $e_b(\mathcal{H})$  denote the total number of blue edges in  $\mathcal{H}$ . We have

$$\begin{aligned} e_b(\mathcal{H}) &\leq \sum_{i=0}^{m-1} y_{A_i, \mathcal{B}_i} + z_C \\ &< \sum_{i=0}^{m-1} \frac{1}{2kr} t_{\mathcal{B}_i} + \frac{1}{2r} t_k. \end{aligned}$$

Note that every proper  $k$ -cycle can extend exactly  $k$  proper  $(k-1)$ -paths. We have  $\sum_{i=0}^{m-1} t_{\mathcal{B}_i} \leq k t_k$ . Thus,

$$\begin{aligned} e_b(\mathcal{H}) &< \frac{1}{2kr} \sum_{i=0}^{m-1} t_{\mathcal{B}_i} + \frac{1}{2r} t_k \\ &\leq \frac{1}{2r} t_k + \frac{1}{2r} t_k \\ &= \frac{1}{r} |E(\mathcal{H})|. \end{aligned}$$

The conclusion is that the number of blue edges in  $\mathcal{H}$  is strictly smaller than  $\frac{1}{r}$  of the total number of edges in  $\mathcal{H}$ , which contradicts that blue is the color class with the most number of edges of  $\mathcal{H}$ .  $\square$

## References

- [1] N. Alon and F. R. K. Chung, *Explicit construction of linear sized tolerant networks*, Discrete Math. **72** (1988), no. 1-3, 15–19.
- [2] Noga Alon and Joel H. Spencer, *The Probabilistic Method*. Wiley Series in Discrete Mathematics and Optimization. Wiley, 2016.
- [3] J. Beck, *On size Ramsey number of paths, trees, and circuits. I*, Journal of Graph Theory **7** (1983), no. 1, 115–129.



- [4] Herman Chernoff, *A note on an inequality involving the normal distribution*, Ann. Probab., **9** (1981), no. 3, 533–535.
- [5] D. Clemens, M. Jenssen, Y. Kohayakawa, N. Morrison, G.O. Mota, D. Reding, B. Roberts, *The size-Ramsey number of power of paths*, arXiv:1707.04297.
- [6] A. Dudek and P. Prałat, *An alternative proof of the linearity of the size-Ramsey number of paths*, Combin. Probab. Comput. **24** (2015), no. 3, 551–555.
- [7] A. Dudek and P. Prałat, *On some multicolour Ramsey properties of random graphs*, SIAM J. Discrete Math., **31** (2017), no. 3, 2079–2092.
- [8] P. Erdős, R. Faudree, C. Rousseau, and R. Schelp, *The size Ramsey number*, Periodica Mathematica Hungarica **9** (1978), no. 1-2, 145–161.
- [9] P. Erdős, *On the combinatorial problems which I would most like to see solved*, Combinatorica **1** (1981), no. 1, 25–42.
- [10] S. Letzter, *Path Ramsey number for random graphs*, Combin. Probab. Comput. **25** (2016), no. 4, 612–622.
- [11] A. Dudek, S. L. Fleur, D. Mubayi, and V. Rödl, *On the size-Ramsey number of hypergraphs*, J. Graph Theory **86** (2017), no. 1, 104–121.
- [12] J. H. Kim and V.H. Vu, *Concentration of multivariate polynomials and its applications* (2000), Combinatorica **20**, pp. 417–434.
- [13] M. Krivelevich, *Long cycles in locally expanding graphs, with applications*, arXiv: 1609.06851.