

The minimum forcing number of perfect matchings in the hypercube

Ajit A. Diwan

Department of Computer Science and Engineering,
Indian Institute of Technology Bombay, Mumbai 400076, India.

January 19, 2022

Abstract Let M be a perfect matching in a graph. A subset S of M is said to be a forcing set of M , if M is the only perfect matching in the graph that contains S . The minimum size of a forcing set of M is called the forcing number of M . Pachter and Kim [Discrete Math. 190 (1998) 287–294] conjectured that the forcing number of every perfect matching in the n -dimensional hypercube is at least 2^{n-2} , for all $n \geq 2$. Riddle [Discrete Math. 245 (2002) 283-292] proved this for even n . We show that the conjecture holds for all $n \geq 2$. The proof is based on simple linear algebra.

1 Introduction

Let M be a perfect matching in a graph G . A subset S of edges in M is said to be a *forcing set* of M , if M is the only perfect matching in G that contains S . The minimum size of a forcing set of M is called the forcing number of M .

Forcing sets and forcing numbers of perfect matchings were first studied by Harary et al. [3], and have applications in chemistry [2]. Pachter and Kim [4] found tight upper and lower bounds on the forcing number of perfect matchings in a 2-dimensional square grid. They also conjectured that every perfect matching in the n -dimensional hypercube has forcing number at least 2^{n-2} , for all $n \geq 2$. This was proved by Riddle [5] for even n . Adams et al. [1] showed that for all $n \geq 5$ and integers i in the interval $[2^{n-2}, 2^{n-2} + 2^{n-5}]$,

there exists a perfect matching in the n -dimensional hypercube with forcing number i .

In this note, we complete the proof of Pachter and Kim's conjecture, and show that it holds for all $n \geq 2$. Our proof is much simpler than the one in [5], and uses only basic linear algebra. We actually prove a slightly stronger statement. If S is a forcing set of a perfect matching M in a graph, then the graph obtained by deleting all endvertices of edges in S has a unique perfect matching $M \setminus S$. Thus an upper bound on the order of an induced subgraph with a unique perfect matching directly gives a lower bound on the forcing number of every perfect matching in the graph. We show that any induced subgraph of the n -dimensional hypercube, with a unique perfect matching, has at most 2^{n-1} vertices, for $n \geq 2$. This implies the lower bound on the forcing number. The technique used is quite general, and can be applied to any bipartite graph.

2 Hypercube

We describe our technique in general, and then apply it to the specific case of the hypercube. Let G be a bipartite graph, with a bipartition X, Y of the vertex set. Let $X = \{x_1, \dots, x_n\}$ and $Y = \{y_1, \dots, y_m\}$. The weighted bipartite adjacency matrix of G is an $n \times m$ matrix W , with $W_{ij} = 0$ if x_i is not adjacent to y_j , and $W_{ij} = w$ otherwise. Here, w is an indeterminate used to indicate the presence of an edge in the graph.

Let H be an induced subgraph of G with a unique perfect matching. The rows and columns of W that correspond to vertices in H form a square sub-matrix W_H of W . Since H has a unique perfect matching, the determinant of W_H must be $\pm w^{|H|/2}$, since only one term in the expansion of the determinant is non-zero. This implies that if we replace each occurrence of w in W by an arbitrary *non-zero* element from some field \mathcal{F} , the resulting matrix must have rank at least $|H|/2$ over \mathcal{F} . Thus, if we choose a field \mathcal{F} , and appropriate non-zero values from the field as the weights of the edges, then twice the rank of the resulting bipartite adjacency matrix is an upper bound on the order of any induced subgraph with a unique perfect matching. We now apply this technique to the hypercube.

The vertex set of the n -dimensional hypercube Q_n is $\{0, 1\}^n$, the set of all sequences of length n in which each element is either 0 or 1. Two sequences are adjacent if they differ in exactly one position. Let E_n denote the subset of

these sequences with an even number of 1 elements, and O_n the subset with an odd number of 1 elements. The hypercube is a bipartite graph with E_n, O_n the two parts of the bipartition. Construct the weighted bipartite adjacency matrix W_n of Q_n with rows indexed by elements of E_n in lexicographic order, and columns indexed by elements of O_n in lexicographic order. Then W_n is a symmetric matrix with a simple recursive structure. W_1 is the 1×1 matrix with w as the only entry and

$$W_{n+1} = \left[\begin{array}{c|c} W_n & wI_{2^{n-1}} \\ \hline wI_{2^{n-1}} & W_n \end{array} \right].$$

Here, I_k denotes the $k \times k$ identity matrix.

We use the recursive structure of W_n to assign non-zero values from the field Z_3 of integers modulo 3, to the w elements in W_n , so that the rank of the resulting matrix is 2^{n-2} over Z_3 .

Lemma 1 *There exists a nonsingular matrix A_n , obtained by assigning non-zero values from the field Z_3 to the w elements in W_n , such that A_n^{-1} can also be obtained by assigning non-zero values to the w elements in W_n .*

Proof: We construct the matrix inductively. For $n = 1$, let A_1 be the 1×1 matrix containing the entry 1. Clearly, $A_1^{-1} = A_1$ satisfies the required property. Let

$$A_{n+1} = \left[\begin{array}{c|c} 2A_n & I_{2^{n-1}} \\ \hline I_{2^{n-1}} & A_n^{-1} \end{array} \right].$$

Then it is easy to verify that

$$A_{n+1}^{-1} = \left[\begin{array}{c|c} A_n^{-1} & 2I_{2^{n-1}} \\ \hline 2I_{2^{n-1}} & 2A_n \end{array} \right].$$

By induction, and using the recursive structure of W_{n+1} , we conclude that both A_{n+1} and A_{n+1}^{-1} can be obtained from W_{n+1} by assigning non-zero values to the w elements in W_{n+1} . \square

Theorem 1 *Any induced subgraph of Q_n that has a unique perfect matching contains at most 2^{n-1} vertices, for all $n \geq 2$.*

Proof: Let A_{n-1} be the matrix over Z_3 satisfying the properties in Lemma 1. Let

$$B_n = \left[\begin{array}{c|c} A_{n-1} & I_{2^{n-2}} \\ \hline I_{2^{n-2}} & A_{n-1}^{-1} \end{array} \right].$$

Now, it is easy to see that B_n has rank 2^{n-2} over Z_3 , and is obtained from W_n by replacing all occurrences of w in W_n by non-zero values from Z_3 . \square

Note that for even n , we get a simpler proof by considering the field Z_2 . In this case, if we replace all occurrences of w in W_n by 1, we get a matrix of rank 2^{n-2} over Z_2 . However, for odd n , this gives a nonsingular matrix which is its own inverse. Riddle's proof for even n is purely combinatorial. It would be interesting to see if the proof for odd n can also be made purely combinatorial. It would also be interesting to see if this technique can be used to find tight bounds on the forcing numbers of perfect matchings in other bipartite graphs.

References

- [1] P. Adams, M. Mahdian, E. S. Mahmoodian, On the forced matching number of bipartite graphs, *Discrete Math.* 281 (2004) 1–12.
- [2] Z. Che, Z. Chen, Forcing on perfect matchings—a survey, *MATCH Commun. Math. Comput. Chem.* 66 (2011) 93–136.
- [3] F. Harary, D. J. Klein, T. P. Živković, Graphical properties of polyhexes: perfect matching vector and forcing, *J. Math. Chem.* 6 (1991) 295–306.
- [4] L. Pachter, P. Kim, Forcing matchings on square grids, *Discrete Math.* 190 (1998) 287–294.
- [5] M. E. Riddle, The minimum forcing number for the torus and hypercube, *Discrete Math.* 245 (2002) 283–292.