

A Generalization of Schur's Theorem

Jon Henry Sanders

JHS Consulting jon_sanders@partech.com

arXiv:1712.03620v1 [math.CO] 11 Dec 2017

Chapter 1

1.1 A GENERALIZATION OF SCHUR'S THEOREM

In 1916 Schur [1] proved the following theorem:

Theorem 1. Given a number¹ there exists a number $J(t)$ such that if the set $\{1, \dots, J(t)\}$ is divided into t classes A_1, \dots, A_t then there exists numbers a, b and a number $\ell, 1 \leq \ell \leq t$, such that $a, b, a + b \in A_\ell$. We prove the following generalization:

Theorem 2. Let $P(t, n)$ be the statement, "Given numbers t, n there exists a number $J(t, n)$ such that if the set $S = \{1, \dots, J(t, n)\}$ is divided into t classes A_1, \dots, A_t there exists a sequence of (not necessarily distinct) numbers a_1, \dots, a_n and a number $\ell, 1 \leq \ell \leq t$, such that $\sum_{i \in I} a_i \in A_\ell$ for all $I \subseteq \{1, \dots, n\}$." Then $P(t, n)$ is true for all numbers t, n .

We first prove the following corollary:

Corollary 1.1. The a_i may be assumed distinct in the above theorem.

Proof. Let $J'(t, n) = J(t, n^4)$. We show if the set $S = \{1, \dots, J'(t, n)\}$ is divided into t classes there are n distinct numbers in S with the desired property. For let a_1, \dots, a_{n^4} be a sequence of n^4 not necessarily distinct

¹The word "number" will mean "positive integer" throughout this section.

numbers of S with the desired property. Then either there are at least n^2 distinct numbers among them or there is a number k such that $a_1 = k$ for at least n^2 distinct numbers among them or there is a number k such that $a_i = k$ for at least n^2 distinct values of i . In the first case we are through. In the second define $b_i = ik$, $i = 1, \dots, n$. Then $\sum_{i \in I} b_i, I \subseteq \{1, \dots, n\}$ is equal to ℓk , $1 \leq \ell \leq \frac{n(n-1)}{2} \leq n^2 = \sum_{i \in I'} a_i, I' \subseteq \{1, \dots, n^4\}$ where I' is a set of ℓ (distinct) indices which satisfy $i \in I'$ implies $a_i = k$. Thus the b_1 have the desired property since the a_1 do and furthermore they are distinct. \square

We need to make one remark before proceeding. Let $P(t, n, k)$ be the statement derived from $P(t, n)$ by substituting the set $S' = \{k, 2k, \dots, J(t, n)k\}$ for the set $S = \{1, 2, \dots, J(t, n)\}$.

Remark. For any given number k , $P(t, n)$ implies² $P(t, n, k)$.

Proof. Let A_1, \dots, A_t be a division of S' into t sets. Divide S into t sets by the rule for all $a \in S, a \in A'_j$ iff $ka \in A_j$. If $P(t, n)$ is true there exists a sequence b_1, \dots, b_n such that $\sum_{i \in I} b_i \in A'_\ell$ for some fixed ℓ , $1 \leq \ell \leq t$, for all $I \subseteq \{1, \dots, n\}$. But then the sequence $a_i = b_i k$, $i = 1, \dots, n$ and the number ℓ , $1 \leq \ell \leq t$, have the desired property for S' so $P(t, n, k)$ is true. \square

We now prove an ‘‘Iterated Ramsey Theorem’’ needed for the proof of Theorem 2. Let $R(k, r, t)$ be Ramsey’s function, i.e., let $R(k, r, t)$ be the smallest number such that when the r -subsets of any set S of order $R(k, r, t)$ are divided into t classes A_1, \dots, A_t there always exists a k -subset K of S and a number ℓ , $1 \leq \ell \leq t$ such that all the r -subsets of K are contained in A_ℓ . We prove

²Actually, $P(t, n)$ and $P(t, n, k)$ are equivalent.

Lemma 1. Given numbers k, r, t , $k \geq r$ there exists a number $N(k, r, t)$ such that if all the non-empty subsets of a set S of order $n \geq N(k, r, t)$ are divided into t classes A_1, \dots, A_t then there exists a k -subset $K \subseteq S$ and a sequence A_{i_1}, \dots, A_{i_r} , $1 \leq i_j \leq t$, $j = 1, \dots, r$, such that all the j -subsets of K are contained in A_{i_j} for $j = 1, \dots, r$.

Proof. The proof is by induction on r . If $r = 1$ we take $N(k, l, t) > kt$. Suppose the theorem is true for $r = r_o - 1$ and arbitrary k, t . We wish to prove it true for $r = r_o$ and arbitrary values of k, t , say k_o, t_o . Take $N(k_o, r_o, t_o) = N(R(k_o, r_o - 1, t_o), r_o, t_o)$. If the order of S is larger than $N(k_o, r_o, t_o)$ then by the induction hypothesis there exists a subset K' of S of order $R(k_o, r_o, t_o)$ and a sequence $i_1, \dots, i_{r_o - 1}$ such that each j -subset of K' is contained in A_{i_j} , $j = 1, \dots, r_o - 1$. But by Ramsey's Theorem there exists a k_o -subset K'' of K' such that all the r_o -subsets of K'' are in some A' . Then K'' is the desired subset of S and $A_{i_1}, \dots, A_{i_{r_o - 1}}, A'$ is the required sequence of classes. \square

We are now ready to prove Theorem 2.

Proof. The proof is by induction on n . Let $P'(n)$ be the statement " $P(t, n)$ is true for all possible values of t ." Then $P'(2)$ is Theorem 1. Assume $P'(n_o - 1)$ is true. We prove $P'(n_o)$ is true. Take

$$J(n_o, t) = N(4J(n_o - 1, t), 4J(n_o - 1, t), t).$$

Let the set $S = \{1, \dots, J(n_o, t)\}$ be divided into t classes A_1, \dots, A_t . If $B = \{a_1, a_2, \dots, a_k\}$ is any subset of S , $a_1 < a_2 < \dots < a_k$ define $f(B)$ by $f(B) = \sum_{j=1}^k (-1)^j a_j$ if k is even and by $f(B) = \sum_{j=1}^k (-1)^{j+1} a_j$ if k is odd. Divide all the non-empty subsets of S into t classes A'_1, \dots, A'_t by the rule

$B \in A'_i$ iff $f(B) \in A_i$. (Note that this is well defined since $f(B) \in S$ if $\theta \neq B \subseteq S$). By Lemma 1³, there exists a subset K of S of order $4J(n_o^{-1}, t) := N$ and a sequence of classes $A'_{j_1}, \dots, A'_{j_N}$ such that all the i -subsets of K are contained in $A'_{j_i}, i = 1, \dots, N$. Now divide the set $S' = \{4, 8, \dots, 4J(n_o-1, t)\}$ into t classes A''_1, \dots, A''_t by the rule for all $a \in S', a \in A''_j$ iff all the a -subsets of K are contained in A'_j . Then by the induction hypothesis and the preceding remark we can find a sequence a_1, \dots, a_{n_o-1} , say $a_1 \leq a_2 \leq \dots \leq a_{n_o-1}$, and a number $\ell, 1 \leq \ell \leq t$, such that $\sum_{i \in I} a_i \in A''_\ell$ for all $I \subseteq \{1, \dots, n_o-1\}$. Since $\sum_{i=I} a_i \leq \text{order of } K^4$ we may choose a strictly increasing sequence $b_1, b_2, \dots, b_{\theta_{n_o-1}}$ of elements of K , where $\theta_{n_o-1} = \sum_{i=1}^{n_o-1} a_i$. In general define $\theta_i = \sum_{j=1}^i a_j, i = 1, \dots, n_o-1$ and define some subsets of K as follows (see Fig. 1).

$$B_1 = \{b_1, \dots, b_{a_1}\},$$

$$B_i = \{b_{\theta_{i-1}+1}, \dots, b_{\theta_i}\}, \quad i = 2, \dots, n_o-1,$$

$$P_1 = \left\{ b_{\frac{a_1}{2}-1}, \dots, b_{a_1-1} \right\},$$

$$P_2 = \left\{ b_{a_1+2}, \dots, b_{\frac{3a_1}{2}+2} \right\},$$

and $B_{n_o} = P_1 \cup P_2$.

³with $k = r = N$.

⁴Since $\sum_{i=1}^{n_o-1} a_i \in A''_\ell \subseteq S', \sum_{i=1}^{n_o-1} a_i \leq \max S' = 4J(n_o-1, t) = \text{order of } K$.

$$\begin{array}{ccc}
\overbrace{b_1, \dots, b_{\frac{a_1}{2}-1}, \dots, b_{a_1-1}, b_{a_1}}^{B_1} & \overbrace{b_{a_1+1}, b_{a_1+2}, \dots, b_{\frac{3a_1}{2}+2}, \dots, b_{a_1+a_2}}^{B_2} & \overbrace{b_{a_1+a_2+1}, \dots}^{B_3} \\
\underbrace{\hspace{10em}}_{P_1} & \underbrace{\hspace{10em}}_{P_2} &
\end{array}$$

Figure 1.

We claim that $f(B_i), i = 1, \dots, n_o$ and A_ℓ are the required numbers and class, i.e.,

$$\sum_{i \in I} f(B_i) \in A_\ell, \quad I \subseteq \{1, \dots, n_o\}. \quad T$$

We first state two facts about the function f .

$$F_1 \quad f(A \cup B) = f(A) + f(B) \text{ if } A \text{ and } B \text{ are of even order and } A < B^5.$$

$$F_2 \quad f(A) = f(A - P) - f(P) \text{ when } P \text{ is a consecutive}^6 \text{ subset of } A \text{ of even order such that all the elements of } A \text{ larger than the largest element of } P \text{ are odd in number.}$$

We first show T assuming either $n_o \notin I$ or $n_o \in I$ but $1, 2 \notin I$. By F_1 we have $\sum_{i \in I} f(B_i) = f(\bigcup_{i \in I} B_i)$. But $\bigcup_{i \in I} B_i$ is of order $\sum_{i \in I} a_i$ if $n_o \notin I$ and of order $\sum_{i \in I - \{n_o\}} a_i + a_1$ if $n_o \in I, 1, 2 \notin I$ since the B_i are disjoint in either case and $|B_{n_o}| = a_1$ and $|B_i| = a_i, i = 1, \dots, n_o - 1$. Thus in either case $\bigcup_{i \in I} B_i \in A'_\ell$ (since $|\bigcup_{i \in I} B_i| \in A''_\ell$) so $\sum_{i \in I} f(B_i) = f(\bigcup_{i \in I} B_i) \subseteq A_\ell$. Next we consider the case $n_o, 1 \in I, 2 \notin I$. We claim $f(B_{n_o}) + f(B_1) = f((B_1 - P_1) \cup P_2)$. [This follows

⁵ $A < B$ denotes here that the largest element of A is less than the smallest element of B .

⁶ By a consecutive subset we mean that $a \in A, x < a < y, x, y \in P$ implies $a \in P$.

from the fact that $\frac{a_1}{2}$ is even so $f(B_{n_o}) = f(P_1 \cup P_2) = f(P_1) + f(P_2)$ (by F_1 since $|P_1| = |P_2| = \frac{a_1}{2}$ and $P_1 < P_2$). Then by F_2 , $f(B_1) + f(P_1) = f(B_1 - P_1)$ and by F_1 , $f(B_1 - P_1) + f(P_2) = f((B_1 - P_1) \cup P_2)$.] But $(B_1 - P_1) \cup P_2$ is of order a_1 and $(B_1 - P_1) \cup P_2 < B_i, i = 3, \dots, n_o - 1$,

so by F_1 , $\sum_{i \in I} f(B_i) = f(B_{n_o}) + f(B_1) + \sum_{i \in I - \{n_o, 1\}} f(B_i) = f((B_1 - P_1) \cup P_2) + \sum_{i \in I - \{n_o, 1\}} f(B_i) = f((B_1 - P_1) \cup P_2 \cup \bigcup_{i \in I - \{n_o, 1\}} B_i)$. But $(B_1 - P_1) \cup P_2 \cup \bigcup_{i \in I - \{n_o, 1\}} B_i$ is of order $\sum_{i \in I - \{n_o\}} a_i$. Thus $((B_1 - P_1) \cup P_2) \cup \bigcup_{i \in I - \{n_o, 1\}} B_i \in A_\ell$, so $\sum_{i \in I} f(B_i) = f(\bigcup_{i=1-\{1, n_o\}} B_i \cup (B_1 - P_1) \cup P_2) \in A_\ell$.

Similar arguments hold in the cases $n_o, 2 \in I, 1 \notin I$ and $n_o, 2, 1 \in I$ noting $f(B_{n_o}) + f(B_2) = f((B_2 - P_2) \cup B_1)$ and $f(B_{n_o}) + f(B_2) + f(B_1) = f((B_1 \cup B_2) - B_{n_o})$. \square

1.2 FURTHER GENERALIZATIONS

It is natural to ask whether either Lemma 1 or Theorem 2 generalize in the following way:

Lemma 1'. Let all the non-empty finite subsets of a countably infinite set S be divided into t classes A_1, \dots, A_t . Then there exists a countably infinite subset $K \subseteq S$ and an infinite sequence $i_1, i_2, \dots, i \leq i_j \leq t, i = 1, 2, \dots$, such that all the j -subsets of K are contained in $A_{i_j}, j = 1, 2, \dots$.

Theorem 2'. Let the natural numbers be divided into t classes A_1, \dots, A_t . Then there exists an infinite sequence a_1, a_2, \dots , and a number $\ell, 1 \leq \ell \leq t$, such that $\sum_{i \in I} a_i \in A_\ell$, for all (non-empty) finite sets I of natural numbers.

A counter example to Lemma 1' is as follows. Divide all the finite (non-empty) subsets S of the natural numbers N into two classes A_1 and A_2 by

the rule:

$$\begin{aligned} S \in A_1 & \quad \text{iff} \quad |S| \in S, \\ S \in A_2 & \quad \text{iff} \quad |S| \notin S. \end{aligned}$$

Then K contains k -subsets which contain k and k -subsets which do not, i.e., subsets which belong to A_1 and ones which belong to A_2 .

We note that if Theorem 2' were true then using an argument similar to that of Corollary 1.1 we could assume the a_1 are distinct, however, at present the validity of Theorem 2' is not known.

BIBLIOGRAPHY

1. I. Schur, Über die Kongruenz $x^m + y^m = z^m \pmod{p}$, Jahresbericht der Deut. Mathematiker-Vereinigung, Band 25 (1916) pp. 114-117.