

ON THE SPACE OF 2-LINKAGES

GUANTAO CHEN, HEIN VAN DER HOLST, SERGUEI NORINE, AND ROBIN THOMAS

ABSTRACT. Let $G = (V, E)$ be a finite undirected graph. If P is an oriented path from $r_1 \in V$ to $r_2 \in V$, we define $\partial(P) = r_2 - r_1$. If $R, S \subseteq V$, we denote by $P(G; R, S)$ the span of the set of all $\partial P \otimes \partial Q$ with P and Q disjoint oriented paths of G connecting vertices in R and S , respectively. By $L(R, S)$, we denote the submodule of $\mathbb{Z}\langle R \rangle \otimes \mathbb{Z}\langle S \rangle$ consisting all $\sum_{r \in R, s \in S} c(r, s)r \otimes s$ such that $c(r, r) = 0$ for all $r \in R \cap S$, $\sum_{r \in R} c(r, s) = 0$ for all $s \in S$, and $\sum_{s \in S} c(r, s) = 0$ for all $r \in R$.

In this paper, we provide, when G is sufficiently connected, characterizations when $P(G; R, S)$ is a proper subset of $L(R, S)$.

1. INTRODUCTION

Let R be a finite set. By $\mathbb{Z}\langle R \rangle$, we denote the free \mathbb{Z} -module generated by R ; that is, $\mathbb{Z}\langle R \rangle$ is the module of all $\sum_{r \in R} c(r)r$ with $c(r) \in \mathbb{Z}$. By $\mathbb{Z}\langle R \rangle^{\otimes 2}$, we denote the tensor product of $\mathbb{Z}\langle R \rangle$ and $\mathbb{Z}\langle R \rangle$; that is, $\mathbb{Z}\langle R \rangle^{\otimes 2}$ is the module of all $\sum_{r, s \in R} c(r \otimes s)r \otimes s$, where $c(r \otimes s) \in \mathbb{Z}$. Observe that we can identify elements $c \in \mathbb{Z}\langle R \rangle^{\otimes 2}$ with linear forms $c : \mathbb{Z}\langle R \rangle \otimes \mathbb{Z}\langle R \rangle \rightarrow \mathbb{Z}$.

Suppose that R is a finite set with a cyclic ordering r_1, r_2, \dots, r_n . So the successor of r_i in this cyclic ordering is r_{i+1} if $i \neq n$, and r_1 if $i = n$. If $u, v \in R$ are distinct elements, we define

$$\text{link}(u \otimes v; r_1, r_2, \dots, r_n) = \begin{cases} 1 & \text{if } v \text{ comes after } u \text{ in the sequence } r_1, \dots, r_n \\ -1 & \text{if } v \text{ comes before } u \text{ in the sequence } r_1, \dots, r_n. \end{cases}$$

Let $D(R) = \{\sum_{r, s \in R} c(r \otimes s)r \otimes s \in \mathbb{Z}\langle R \rangle^{\otimes 2} \mid c(r \otimes r) = 0 \text{ for all } r \in R\}$. We extend $\text{link}(\cdot; r_1, r_2, \dots, r_n)$ linearly to $D(R)$. It is easy to see that, for all $c \in D(R)$,

$$\text{link}(c; r_1, r_2, \dots, r_n) = \text{link}(c; r_n, r_1, \dots, r_{n-1})$$

if and only if, for all $r \in R$,

$$\sum_{s \in R} c(r \otimes s) = 0 \text{ and } \sum_{s \in R} c(s \otimes r) = 0.$$

We denote by $L(R)$ the submodule of $D(R)$ of all $c \in D(R)$ such that, for all $r \in R$,

$$\sum_{s \in R} c(r \otimes s) = 0 \text{ and } \sum_{s \in R} c(s \otimes r) = 0.$$

Then, if $c \in L(R)$, $\text{link}(c; r_1, r_2, \dots, r_n)$ is independent of where we start the cyclic ordering of r_1, r_2, \dots, r_n .

Let $R = \{r_1, r_2, \dots, r_n\}$ with a cyclic ordering r_1, r_2, \dots, r_n . If $u_1, u_2, v_1, v_2 \in R$, then

$$c(u_1 u_2; v_1 v_2) := (u_1 - u_2) \otimes (v_1 - v_2) \in L(R).$$

Suppose u_1, u_2, v_1, v_2 are distinct. If u_1, u_2, v_1, v_2 occur in this order in r_1, r_2, \dots, r_n , then

$$\text{link}(c(u_1 u_2; v_1 v_2); r_1, r_2, \dots, r_n) = 0,$$

while, if u_1, v_1, u_2, v_2 occur in this order in r_1, r_2, \dots, r_n , then

$$\text{link}(c(u_1 u_2; v_1 v_2); r_1, r_2, \dots, r_n) = 2,$$

and if u_2, v_1, u_1, v_2 occur in this order in r_1, r_2, \dots, r_n , then

$$\text{link}(c(u_1 u_2; v_1 v_2); r_1, r_2, \dots, r_n) = -2.$$

If $u_1, u_2, u_3 \in R$, then

$$c(u_1, u_2, u_3) := u_1 \otimes u_2 - u_1 \otimes u_3 + u_2 \otimes u_3 - u_2 \otimes u_1 + u_3 \otimes u_1 - u_3 \otimes u_2 \in L(R).$$

Suppose u_1, u_2, u_3 are distinct. If u_1, u_2, u_3 occur in this order in r_1, r_2, \dots, r_n , then

$$\text{link}(c(u_1, u_2, u_3); r_1, r_2, \dots, r_n) = 2,$$

while, if u_3, u_2, u_1 occur in this order in R , then

$$\text{link}(c(u_1, u_2, u_3); r_1, r_2, \dots, r_n) = -2.$$

If R, S are finite sets, we denote by $L(R, S)$ the submodule of $L(R \cup S)$ of all $c \in L(R \cup S)$ with $c(r, s) = 0$ if $r \notin R$ or $s \notin S$. If $u_1, u_2 \in R$ and $s_1, s_2 \in S$ are distinct, then $c(u_1 u_2; v_1 v_2) \in L(R, S)$. Observe that $c(u_1, u_2, u_3) \in L(R, S)$ only if $u_1, u_2, u_3 \in R \cap S$.

Let $G = (V, E)$ be a finite graph. If P is an oriented path in G from u_2 to u_1 , then

$$\partial(P) := u_1 - u_2 \in \mathbb{Z}\langle V \rangle.$$

If $R_1, R_2 \subseteq V$, an $(R_1; R_2)$ -linkage in G is a pair of disjoint paths (P_1, P_2) in G such that, for $i = 1, 2$, P_i connects vertices in R_i . Let u_1, u_2, v_1, v_2 be vertices in V . A $(u_1 u_2; v_1 v_2)$ -linkage in G is a pair of disjoint paths (P_1, P_2) in G such that P_1 connects u_1 and u_2 , and P_2 connects v_1 and v_2 . If $R_1, R_2 \subseteq V$, we denote by $P(G; R_1, R_2)$ the span of all $\partial(P_1) \otimes \partial(P_2) \in \mathbb{Z}\langle R_1 \rangle \otimes \mathbb{Z}\langle R_2 \rangle$ where (P_1, P_2) are $(R_1; R_2)$ -linkages in G . Here we give P_1 and P_2 arbitrary orientations. Clearly, $P(G; R_1, R_2) \subseteq L(R_1, R_2)$.

A *separation* of a graph G is a pair (G_1, G_2) of subgraphs with $G_1 \cup G_2 = G$ and $E(G_1) \cap E(G_2) = \emptyset$. The *width* of a separation (G_1, G_2) is $k := V(G_1 \cap G_2)$. We call a separation of width k a k -separation.

Let $R, S \subseteq V$. We call a 2-separation (G_1, G_2) of G a *sided (R, S) -separation* if $|V(G_2) - V(G_1)| > 0$, $(V(G_1) - V(G_2)) \cap S = \emptyset$, and $G_1 - V(G_2)$ contains at least two vertices u_1, u_2 from R and there are $(u_1 w_1; u_2 w_2)$ - and $(u_1 w_2; u_2 w_1)$ -linkages in G_1 , where $\{w_1, w_2\} = V(G_1 \cap G_2)$. A graph G is *(R, S) -connected* if G is 2-connected and G has no sided (R, S) - and no sided (S, R) -separations.

Let $G = (V, E)$ be a graph. If $X \subseteq V$, we denote by $N_G(X)$ the set of vertices in $V - X$ that are adjacent to a vertex in X . If it is clear what graph we mean, we also write $N(X)$ for $N_G(X)$. We call a family $\mathcal{A} = \{A_1, \dots, A_k\}$ of pairwise disjoint subsets of V *separated* if for $1 \leq i \neq j \leq k$, $N(A_i) \cap A_j = \emptyset$. If \mathcal{A} is a separated family of subsets of V , we denote by $p(G, \mathcal{A})$ the graph obtained from G by deleting A_i for each i and adding new edges joining every pair of distinct vertices in $N(A_i)$.

A *nearly planar* graph is a pair (G, \mathcal{A}) , where $G = (V, E)$ is a graph and $\mathcal{A} = \{A_1, \dots, A_k\}$ is a separated family of subsets of V such that for each A_i , $|N(A_i)| \leq 3$ and $p(G, \mathcal{A})$ can be drawn in a closed disc D with no pair of edges crossing and for each A_i with $|N(A_i)| = 3$, $N(A_i)$ induces a facial triangle in $p(G, \mathcal{A})$. If, in addition, r_1, \dots, r_n are vertices in G such that $r_i \notin A_j$ for any $A_j \in \mathcal{A}$ and r_1, \dots, r_n occur on the boundary of D in the specified order, then we say that $(G, \mathcal{A}, r_1, \dots, r_n)$ is *nearly planar*. If there is no need to specify the family \mathcal{A} , we also write (G, r_1, \dots, r_n) for $(G, \mathcal{A}, r_1, \dots, r_n)$.

Let t_1, \dots, t_n be a cyclic ordering and suppose $R, S \subseteq \{t_1, \dots, t_n\}$ such that $R \cup S = \{t_1, \dots, t_n\}$. We say that R and S *cross* if $R = S$ and $|R| = 3$, or there exist vertices $u_1, u_2 \in R$ and vertices $v_1, v_2 \in S$ such that u_1, v_1, u_2, v_2 are distinct and occurring in this order in the cyclic ordering t_1, \dots, t_n .

In this paper, we prove the following theorem.

Theorem 1.1. *Let $G = (V, E)$ be a graph, let $R, S \subseteq V$ and suppose G is (R, S) -connected. Then the following are equivalent:*

- (1) $L(R, S) - P(G; R, S) \neq \emptyset$;
- (2) *there exists a cyclic ordering t_1, \dots, t_n of the vertices in $R \cup S$ such that R and S cross and (G, t_1, \dots, t_n) is nearly planar; and*
- (3) *there exists a cyclic ordering t_1, \dots, t_n of the vertices in $R \cup S$ such that R and S cross and, for all $z \in L(R, S)$, $\text{link}(z; t_1, \dots, t_n) = 0$ if and only if $z \in P(G; R, S)$.*

2. ELEMENTARY TENSORS

An *elementary* (R, S) -tensor is a tensor of the form $c(u_1u_2; v_1v_2)$ with $u_1, u_2 \in R$ and $v_1, v_2 \in S$ and u_1, u_2, v_1, v_2 distinct, or a tensor of the form $c(u_1, u_2, u_3)$ with $u_1, u_2, u_3 \in R \cap S$ and u_1, u_2, u_3 distinct. In this section, we show that each element in $L(R, S)$ can be written as a sum of elementary (R, S) -tensors. The proof follows the proof of Theorem 7 in [2].

Lemma 2.1. *Let R, S be finite sets. Then $L(R, S)$ is generated by elementary (R, S) -tensors.*

Proof. Order the elements in R as r_1, \dots, r_k such that we start with the elements that also belong to S . Similarly, we order the elements in S as s_1, \dots, s_m such that we start with the elements that also belong to R . For every $c = \sum_{r \in R, s \in S} c(r \otimes s)r \otimes s \in L(R, S)$, we define the height of c as the minimum of $i + j$ over $c(r_i \otimes s_j) \neq 0$. Suppose for a contradiction that there exists a $c \in L(R, S)$ that is not a sum of elementary (R, S) -tensors. Choose such a c whose height is maximal. Then the height of c is not greater than $k + m$.

Let i and j with $c(r_i \otimes s_j) \neq 0$ and $i + j$ minimum. Let $r_{i'} \in R$ be an element unequal to r_i such that $c(r_{i'} \otimes s_j) \neq 0$, and let $s_{j'} \in S$ be an element unequal to s_j such that $c(r_i \otimes s_{j'}) \neq 0$. Then $r_i \neq s_j, r_i \neq s_{j'}$, and $r_{i'} \neq s_j$.

Suppose that $r_{i'} \neq s_{j'}$. Then $c' = c - c(r_i \otimes s_j)c(r_{i'}r_{i'}; s_j s_{j'}) \in L(R, S)$ is not a sum of elementary (R, S) -tensors. As the height of c' is larger than the height of c , we obtain a contradiction.

Therefore $r_{i'} = s_{j'}$. Then, by choice of the ordering the elements r_1, \dots, r_k and s_1, \dots, s_m and by the minimality of $i + j$, $r_i \in S$ and $s_j \in R$. So each of $r_i, s_j, r_{i'} = s_{j'}$ belongs to both R and S . Then $c' = c - c(r_i \otimes s_j)c(r_i, s_j, r_{i'}) \in L(R, S)$ is not a sum of elementary (R, S) -tensors. As the height of c' is larger than the height of c , we obtain a contradiction. \square

3. SUMS OF LINKAGES

Let $G = (V, E)$ be a graph and let u_1, u_2, v_1, v_2 be distinct vertices of G . In this section, we show that the existence of certain linkages in G implies that $c(u_1u_2; v_1v_2) \in P(G; R, S)$.

Lemma 3.1. *Let $G = (V, E)$ be a graph and let $R, S \subseteq V$. Let $u_1, u_2, u \in R$ and $v_1, v_2 \in S$. If G contains a $(u_1u; v_1v_2)$ - and a $(uu_2; v_2v_1)$ -linkage, then $c(u_1u_2; v_1v_2) \in P(G; R, S)$.*

Proof. Since G has a $(u_1u; v_1v_2)$ -linkage, $c(uu_1; v_2v_1) \in P(G; R, S)$. Since G has a $(uu_2; v_2v_1)$ -linkage, $c(uu_2; v_1v_2) \in P(G; R, S)$. Then $c(u_1u_2; v_1v_2) = c(u_1u; v_1v_2) + c(uu_2; v_1v_2) \in P(G; R, S)$. \square

Lemma 3.2. *Let $G = (V, E)$ be a graph and let $R, S \subseteq V$. Let $u_1, u_2, u \in R$ and $v_1, v_2, v \in S$. If G contains a $(u_1u; v_1v_2)$ -, a $(uu_2; vv_1)$ -, and a $(u_2u; v_2v)$ -linkage, then $c(u_1u_2; v_1v_2) \in P(G; R, S)$.*

Proof. Since G contains a $(u_1u; v_1v_2)$ -linkage, $c(uu_1; v_2v_1) \in P(G; R, S)$. Since G contains a $(uu_2; vv_1)$ -linkage, $c(u_2u; v_1v) \in P(G; R, S)$. Since G contains a $(u_2u; v_2v)$ -linkage, $c(uu_2; vv_2) \in P(G; R, S)$. Then $c(u_1u_2; v_1v_2) = c(uu_1; v_2v_1) + c(u_2u; vv_1) + c(uu_2; vv_2) \in P(G; R, S)$. \square

Lemma 3.3. *Let $G = (V, E)$ be a graph and let $R, S \subseteq V$. Let $u_1, u_2, u \in R$ and $v_1, v_2, v', v'' \in S$. If G contains a $(u_2u_1; v'v_1)$ -, a $(uu_2; v'v_2)$ -, a $(uu_1; v'v_1)$ -, a $(uu_1; v''v_2)$ -, and a $(uu_1; v''v_1)$ -linkage, then $c(u_1u_2; v_1v_2) \in P(G; R, S)$.*

Proof. Since G contains a $(u_1u_2; v'v_1)$ -linkage, $c(u_1u_2; v_1v') \in P(G; R, S)$. Since G contains a $(uu_2; v'v_2)$ -linkage, $c(u_2u; v_2v') \in P(G; R, S)$. Since G contains a $(uu_1; v'v_1)$ -linkage, $c(uu_1; v_1v') \in P(G; R, S)$. Since G contains a $(uu_1; v''v_2)$ -linkage, $c(uu_1; v_2v'') \in P(G; R, S)$. Since G contains a $(uu_1; v''v_1)$ -linkage, $c(u_1u; v_1v'') \in P(G; R, S)$. Then $c(u_1u_2; v_1v_2) = c(u_1u_2; v_1v') + c(u_2u; v_2v') + c(uu_1; v_1v') + c(uu_1; v_2v'') + c(u_1u; v_1v'') \in P(G; R, S)$. \square

Lemma 3.4. *Let $G = (V, E)$ be a graph and let $R, S \subseteq V$. Let $u_1, u_2 \in R \cap S$, let $v_1, v_2 \in S$, and let $u \in R$. If G contains a $(u_1u; v_1u_2)$ -, a $(uu_2; u_1v_1)$ -, a $(uu_1; u_2v_2)$ -, and a $(uu_2; u_1v_2)$ -linkage, then $c(u_1u_2; v_1v_2) \in P(G; R, S)$.*

Proof. Since G contains a $(u_1u; v_1u_2)$ -linkage, $c(uu_1; vu_2) \in P(G; R, S)$. Since G contains a $(uu_2; u_1v_1)$ -linkage, $c(uu_2; v_1u_1) \in P(G; R, S)$. Since G contains a $(uu_1; u_2v_2)$ -linkage, $c(uu_1; v_2u_2) \in P(G; R, S)$. Since G contains a $(uu_2; u_1v_2)$ -linkage, $c(uu_2; v_2u_1) \in P(G; R, S)$. Then $c(u_1u_2; v_1v_2) = c(u_1u; vu_2) + c(uu_2; v_1u_1) - c(uu_1; v_2u_2) + c(uu_2; v_2u_1) \in P(G; R, S)$. \square

4. LINKAGES IN NEARLY PLANAR GRAPHS

In this section, we prove the following theorem.

Theorem 4.1. *Let G be a 2-connected graph and let $R, S \subseteq V(G)$. If $L(R, S) - P(G; R, S) \neq \emptyset$, then either G has a sided (R, S) -separation or a side (S, R) -separation, or there exists a cyclic ordering w_1, \dots, w_n of the vertices in $R \cup S$ such that (G, w_1, \dots, w_n) is nearly planar and R and S cross.*

If P is a path in a graph $G = (V, E)$ connecting u and v , we also write that P is a $u \rightarrow v$ path in G . If $S \subseteq V$ and P is a path in G connecting u and a vertex in S , we also write that P is a $u \rightarrow S$ path in G .

Let (G, \mathcal{A}) be a nearly planar graph. We call an $A \in \mathcal{A}$ *minimal* if there is no collection \mathcal{H} of pairwise disjoint subsets of A such that $\mathcal{H} \neq \{A\}$ and $(G[A \cup N(A)], \mathcal{H}, a_1, \dots, a_m)$ is nearly planar, where $\{a_1, \dots, a_m\} = N(A)$. If every element of \mathcal{A} is minimal, we say that \mathcal{A} is *minimal*.

Lemma 4.2. *Let $(G, \mathcal{A}; r_1, \dots, r_n)$ be a nearly planar graph. Let $A \in \mathcal{A}$ with $|N(A)| = 3$ and A minimal, let $\{u_1, u_2, u_3\} = N(A)$, and let $u \in A$. Then $G[A \cup N(A)]$ contains a $(u_1u_2; u_3u)$ -, a $(u_1u_3; u_2u)$ -, and a $(u_1u; u_2u_3)$ -linkage.*

Lemma 4.3. *Let (G, \mathcal{A}) be a nearly planar graph, where \mathcal{A} is minimal. Let $H = p(G, \mathcal{A})$ and let u_1, u_2, v_1, v_2 be vertices of H . If H contains a $(u_1u_2; v_1v_2)$ -linkage, then G contains a $(u_1u_2; v_1v_2)$ -linkage.*

Proof. Let (P, Q) be a $(u_1u_2; v_1v_2)$ -linkage in H . Replace each edge e of P and Q whose ends u_e and v_e belong to $N(A)$ for some $A \in \mathcal{A}$ by a $u_e \rightarrow v_e$ path T in $G[A \cup N(A)]$. Let P' and Q' be the resulting paths. Then (P', Q') is a $(u_1u_2; v_1v_2)$ -linkage in G . \square

Lemma 4.4. *Let (G, \mathcal{A}) be a nearly planar graph, where \mathcal{A} is minimal. Let $H = p(G, \mathcal{A})$. Let u_1, u_2, v_1, v_2 be vertices of G , with $u_1, u_2, v_1 \in V(H)$ and $v_2 \notin V(H)$. Let P be a $u_1 \rightarrow u_2$ path in H , and let $A \in \mathcal{A}$ with $v_2 \in A$. Let $r \in N(A) - V(P)$ and suppose Q is a $v_1 \rightarrow r$ path disjoint from P . Then G has a $(u_1u_2; v_1v_2)$ -linkage.*

Proof. Suppose e is an edge of P whose ends u_e and v_e belongs to $N(A)$. Since \mathcal{A} is minimal, A is minimal, and so $G[A \cup N(A)]$ contains a $(u_e v_e; ur)$ -linkage (T_1, T_2) . Replace the edge e in P by T_1 . Replace any other edge f of P whose ends s_f and t_f belong to $N(A')$ for some $A' \in \mathcal{A}$ with $A' \neq A$ by a $s_f \rightarrow t_f$ path T in $G[A' \cup N(A')]$. Let P' be the resulting path. Replace each edge f of Q whose ends s_f and t_f belong to $N(A)$ for some $A \in \mathcal{A}$ by a $s_f \rightarrow t_f$ path T in $G[A \cup N(A)]$. Let Q' be the resulting path. Then $(P', Q' \cup T_2)$ is a $(u_1u_2; v_1v_2)$ -linkage in G . \square

Let C be an oriented cycle of a graph G . If $u, v \in V(C)$, we denote by $C[u, v]$ the path in C when traversing C from u to v in forward direction. By $C(u, v)$ we denote $C[u, v] - \{u, v\}$.

Lemma 4.5. *Let (G, \mathcal{A}) be a nearly planar graph, where G is 2-connected and \mathcal{A} is minimal, and let $R, S \subseteq V(G)$. Let $H = p(G, \mathcal{A})$ and let C be an oriented cycle of H . Suppose that u_1, u_2, v_1, v_2 are distinct vertices with $u_1, u_2 \in R$ and $v_1, v_2 \in S$ such that u_1, v_1, u_2, v_2 occurs in this order on C . Let B be a component of $G - V(C)$ that has a neighbor in $C(v_1, v_2)$ and a neighbor in $C(v_2, v_1)$. If B contains a vertex of R , then $c(u_1u_2; v_1v_2) \in P(G; R, S)$.*

Proof. Let $u \in R$ belong to B . Suppose first that $u \notin A$ for all $A \in \mathcal{A}$. Let P'_1 be a $u_1 \rightarrow u$ path in $H - V(C[v_1, v_2])$, and let P'_2 be a $u \rightarrow u_2$ path in $H - V(C[v_2, v_1])$. Then $(P'_1, C[v_1, v_2])$ and $(P'_2, C[v_2, v_1])$

are (u_1u, v_1v_2) - and $(uu_2; v_2v_1)$ -linkages in H , respectively. By Lemma 4.3, G contains a (u_1u, v_1v_2) - and a $(uu_2; v_2v_1)$ -linkage. By Lemma 3.1, $c(u_1u_2; v_1v_2) \in P(G; R, S)$.

Suppose next that $u \in A$ for some $A \in \mathcal{A}$. Since B has a neighbor in $C[v_2, v_1]$ and a neighbor in $C[v_1, v_2]$, $N(A) - V(C[v_2, v_1]) \neq \emptyset$ and $N(A) - V(C[v_1, v_2]) \neq \emptyset$. Let $r \in N(A) - V(C[v_2, v_1])$. Since B has a neighbor in $C[v_2, v_1]$ and a neighbor in $C[v_1, v_2]$, there exists an $r \rightarrow u_2$ disjoint from $C[v_2, v_1]$. Hence H contains a (v_1v_2, ru_2) -linkage. By Lemma 4.4, G contains a (v_1v_2, uu_2) -linkage in G . In the same way, G contains a (v_1v_2, uu_1) -linkage. By Lemma 3.1, $c(u_1u_2; v_1v_2) \in P(G; R, S)$. \square

Lemma 4.6. *Let $(G, \mathcal{A}, t_1, \dots, t_n)$ be a nearly planar graph, where G is 2-connected and \mathcal{A} is minimal, and let $R, S \subseteq V(G)$ such that $\{t_1, \dots, t_n\} \subseteq R \cup S$. Suppose u_1, u_2, v_1, v_2 are distinct vertices with $u_1, u_2 \in R$ and $v_1, v_2 \in S$ such that u_1, v_1, u_2, v_2 occur in this order in the cyclic ordering t_1, \dots, t_n . Let $(H, t_1, \dots, t_n) = p(G, \mathcal{A}, t_1, \dots, t_n)$ and let C be the cycle bounding the infinite face of a plane drawing of H with the vertices t_1, \dots, t_n on C . Let B be a component of $G - V(C)$ such that $N(B) \subseteq C[v_2, v_1]$, with at least one neighbor of B in $C(v_2, v_1)$. Then at least one of the following holds:*

- (1) $c(u_1u_2; v_1v_2) \in P(G; R, S)$,
- (2) G has a sided (R, S) -separation or a sided (S, R) -separation,
- (3) there exists a cyclic ordering w_1, \dots, w_{n+1} with $\{t_1, \dots, t_n\} \subset \{w_1, \dots, w_{n+1}\} \subseteq R \cup S$ such that t_1, \dots, t_n occur in this ordering in w_1, \dots, w_{n+1} and (G, w_1, \dots, w_{n+1}) is nearly planar, or
- (4) no vertex of B belongs to $R \cup S$.

Proof. For each component K of $G - V(C)$ such that $N(K) \subseteq C[v_2, v_1]$ and at least one neighbor belongs to $C(v_2, v_1)$, let $r_i(K)$, $i = 1, 2$, be the neighbor of K in $C[v_2, v_1]$ that is closer to v_i . Choose a component B of $G - V(C)$ such that the length $C[r_2(B), r_1(B)]$ is minimum. For $i = 1, 2$, let $r_i = r_i(B)$. We may assume that a vertex $u \in R \cup S$ belongs to B . Let \mathcal{A}' be the family of all sets $A \in \mathcal{A}$ such that $A \subseteq V(B)$ and let $B' = p(B, \mathcal{A}')$.

Suppose first that $C(r_2, r_1)$ does not contain a vertex of $R \cup S$. For $i = 1, 2$, let z_i be the vertex of $\{t_1, \dots, t_n\}$ in $C[r_1, r_2]$ closer to r_i . Then $C(z_1, z_2)$ has no vertices of $\{t_1, \dots, t_n\}$. Let the cyclic ordering w_1, \dots, w_{n+1} be obtained from t_1, \dots, t_n by adding u between z_1 and z_2 . Then (G, w_1, \dots, w_{n+1}) is nearly planar, $\{t_1, \dots, t_n\} \subset \{w_1, \dots, w_{n+1}\} \subseteq R \cup S$, and t_1, \dots, t_n occur in this ordering in w_1, \dots, w_{n+1} .

We may therefore assume that $C(r_2, r_1)$ contains at least one vertex of $R \cup S$.

Suppose that $u \in R$ and $C(r_2, r_1) \cap S \neq \emptyset$, or that $u \in S$ and $C(r_2, r_1) \cap R \neq \emptyset$. By symmetry, we may assume that the former holds; let v be a vertex in $C(r_2, r_1) \cap S$.

Suppose first that $u \notin A$ for every $A \in \mathcal{A}$. Let P_0 be a $u \rightarrow u_1$ path in $H - V(C[v_1, v_2])$ and let $Q_0 = C[v_1, v_2]$. Then (P_0, Q_0) is a $(uu_1; v_1v_2)$ -linkage in H . By Lemma 4.3, G contains a $(uu_1; v_1v_2)$ -linkage. Let P be an $r_2 \rightarrow u$ path in B' that is internally disjoint from $C[r_2, r_1]$ and let P_1 be the concatenation of P and $C[u_2, r_2]$. Let $Q_1 = C[v_1, v]$. Then (P_1, Q_1) is a $(uu_2; v_1v)$ -linkage in H . By Lemma 4.3, G contains a $(uu_2; v_1v)$ -linkage. In the same way, we define the $(uu_2; v_2v)$ -linkage (P_2, Q_2) . By Lemma 3.2, $c(u_1u_2; v_1v_2) \in P(G; R, S)$.

We may therefore assume that $u \in A$ for some $A \in \mathcal{A}$. Suppose that $N(A) - V(C) = \emptyset$. Then $N(A) \subseteq V(C[v_2, v_1])$ and $r_1, r_2 \in N(A)$. Since $v \in C(r_1, r_1)$, $|N(A)| = 3$. Let $r \in N(A) - \{r_1, r_2\}$ and let P be a $r \rightarrow u_1$ path in $C(v_2, v_1)$. Then $(P, C[v_1, v_2])$ is a $(ru_1; v_1v_2)$ -linkage in H . By Lemma 4.4, there exists a $(uu_1; v_1v_2)$ -linkage in G . Since $(C[r_1, u_2], C[v_2, v])$ is a $(r_1u_2; vv_2)$ -linkage in H , G contains a $(uu_2; v_2v)$ -linkage, by Lemma 4.4. Since $(C[u_2, r_2], C[v, v_1])$ is a $(u_2r_2; vv_1)$ -linkage in H , G contains a $(u_2u; vv_1)$ -linkage. By Lemma 3.2, $c(u_1u_2; v_1v_2) \in P(G; R, S)$.

Suppose next that $N(A) - V(C) \neq \emptyset$. Let $r \in N(A) - V(C)$. Let P be a $r \rightarrow r_1$ path in B' and let P_0 be the concatenation of P and $C[r_1, u_1]$. Then $(P_0, C[v_1, v_2])$ is a $(ru_1; v_1v_2)$ -linkage in H . By Lemma 4.4, G contains a $(uu_1; v_1v_2)$ -linkage. Let P' be a $r \rightarrow r_1$ path in B' and let P_1 be the concatenation of P' and $C[r_1, u_2]$. Then $(P_1, C[v_2, v])$ is a $(ru_2; vv_2)$ -linkage in H . By Lemma 4.4, G contains a $(uu_2; vv_2)$ -linkage. Let P'' be a $r \rightarrow r_2$ path and let P_2 be the concatenation of P'' and $C[u_2, r_2]$.

Then $(P_2, C[v, v_1])$ is a $(ru_2; vv_1)$ -linkage in H . By Lemma 4.4, G contains a $(uu_2; vv_1)$ -linkage. By Lemma 3.2, $c(u_1u_2; v_1v_2) \in P(G; R, S)$.

Suppose next that B and $C(r_2, r_1)$ contain no vertex of R , but that $C(r_2, r_1)$ contains a vertex of S , or that B and $C(r_2, r_1)$ contain no vertex of S , but that $C(r_2, r_1)$ contains a vertex of R . We may assume that B and $C(r_2, r_1)$ contain no vertex of R , but that $C(r_2, r_1)$ contains a vertex $v \in S$. Since B contains a vertex $R \cup S$, B contains a vertex $u \in S$. Let V_1 be the set of vertices w in G such that there exists a $w \rightarrow \{u, v\}$ path that is internally disjoint from $\{r_1, r_2\}$. Let $V_2 = V(G) - (V_1 - \{r_1, r_2\})$. Let $G_1 = G[V_1]$ and let $G_2 = G[V_2]$. Then (G_1, G_2) is a 2-separation. It remains to show that G_1 contains a (r_1u, r_2v) - and a $(r_2u; r_1v)$ -linkage.

Suppose $u \notin A$ for all $A \in \mathcal{A}$. Since $C[r_2, r_1] \cup B'$ contains a $(ur_1; vr_2)$ -linkage, G_1 contains a $(ur_1; vr_2)$ -linkage. In the same way, G_1 contains a $(ur_2; vr_1)$ -linkage. Suppose next that $u \in A$ for some $A \in \mathcal{A}$. If $N(A) - V(C) \neq \emptyset$, let $r \in N(A) - V(C)$. Since $C[r_2, r_1] \cup B'$ contains a $(rr_1; vr_2)$ -linkage, G_1 contains a $(ur_1; vr_2)$ -linkage, by Lemma 4.4. In the same way, G_1 contains a $(ur_2; vr_1)$ -linkage. Suppose next that $N(A) - V(C) = \emptyset$. Let $r_1, r_2 \in N(A)$. Let $r_3 \in N(A) - \{r_1, r_2\}$. By Lemma 4.2, $G[A \cup N(A)]$ contains a $(ur_2; r_3r_1)$ -linkage and a $(ur_1; r_3r_2)$ -linkage. Hence G_1 contains a $(ur_2; vr_1)$ - and a $(ur_1; vr_2)$ -linkage. \square

Lemma 4.7. *Let (G, \mathcal{A}) be a nearly planar graph, where G is 2-connected, and let $R, S \subseteq V(G)$. Let $H = p(G, \mathcal{A})$ and let C be an oriented cycle of H . Suppose u_1, v_1, u_2, v_2 are distinct vertices with $u_1, u_2 \in R$ and $v_1, v_2 \in S$ such that u_1, v_1, u_2, v_2 occurs in this order on C . Let B be a component of $G - V(C)$ such that $N(B) = \{v_1, v_2\}$. If B contains a vertex of R and both $C(v_2, v_1)$ and $C(v_1, v_2)$ contain a vertex of S , then $c(u_1u_2; v_1v_2) \in P(G; R, S)$.*

Proof. Let $u \in V(B) \cap R$, and let $v', v'' \in S$ be vertices in $C(v_2, v_1)$ and $C(v_1, v_2)$, respectively.

We first assume that $u_1 \notin S$ or $u_2 \notin S$. By symmetry, we may assume that $u_1 \notin S$. By symmetry, we may assume that v' is on $C(u_1, v_1)$. Let $P_1 = C[u_2, u_1]$ and let $Q_1 = C[v', v_1]$. Hence G contains a $(u_2u_1; v'v_1)$ -linkage. Let S_2 be a $u \rightarrow v_1$ path in $B \setminus \{v_2\}$, and let P_2 be the concatenation of $C[v_1, u_2]$ and S_2 . Let $Q_2 = C[v_2, v']$. Hence G contains a $(uu_2; v'v_2)$ -linkage. Let S_3 be a $u \rightarrow v_2$ path in $B \setminus \{v_1\}$, and let P_3 be the concatenation of $C[v_2, u_1]$ and S_3 . Let $Q_3 = C[v', v_1]$. Hence G contains a $(uu_1; v'v_1)$ -linkage. Let S_4 be a $u \rightarrow v_1$ path in $B \setminus \{v_2\}$, and let P_4 be the concatenation of S_4 and $C[u_1, v_1]$. Let $Q_4 = C[v'', v_2]$. Hence G contains a $(uu_1; v''v_2)$ -linkage. Let S_5 be a $u \rightarrow v_2$ path in $B \setminus \{v_1\}$, and let P_5 be the concatenation of S_5 and $C[u_1, v_2]$. Let $Q_5 = C[v_1, v'']$. Hence G contains a $(uu_1; v''v_1)$ -linkage. By Lemma 3.3, $c(u_1u_2; v_1v_2) \in P(G; R, S)$.

Next we assume that $u_1, u_2 \in S$. Let P_1 be a $u_1 \rightarrow u$ path in $C[u_1, v_2] \cup (B \setminus \{v_1\})$ and let $Q_1 = C[v_1, u_2]$. Hence G contains a $(u_1u; v_1u_2)$ -linkage. Let P_2 be a $u_2 \rightarrow u$ path in $C[u_2, v_2] \cup (B \setminus \{v_1\})$ and let $Q_2 = C[u_1, v_1]$. Hence G contains a $(uu_2; u_1v_1)$ -linkage. Let P_3 be a $u_1 \rightarrow u$ path in $C[u_1, v_1] \cup (B \setminus \{v_2\})$ and let $Q_3 = C[u_2, v_2]$. Hence G contains a $(uu_1; u_2v_2)$ -linkage. Let P_4 be a $u_2 \rightarrow u$ path in $C[v_1, u_2] \cup (B \setminus \{v_2\})$ and let $Q_4 = C[v_2, u_1]$. Hence G contains a $(uu_2; u_1v_2)$ -linkage. By Lemma 3.4, $c(u_1u_2; v_1v_2) \in P(G; R, S)$. \square

Lemma 4.8. *Let (G, \mathcal{A}) be a nearly planar graph, where G is 2-connected, and let $R, S \subseteq V(G)$. Let $H = p(G, \mathcal{A})$ and let C be an oriented cycle bounding the infinite face of a plane drawing of H . Suppose u_1, v_1, u_2, v_2 are distinct vertices with $u_1, u_2 \in R$ and $v_1, v_2 \in S$ such that u_1, v_1, u_2, v_2 occurs in this order on C . Let B be a component of $G - V(C)$ such that $N(B) = \{v_1, v_2\}$. Then at least one of the following holds:*

- (i) $c(u_1u_2; v_1v_2) \in P(G; R, S)$,
- (ii) G has a sided (R, S) -separation or a sided (S, R) -separation, or
- (iii) no vertex of B belongs to $R \cup S$.

Proof. Suppose first that B contains a vertex of S . Then, by Lemma 4.5, $c(u_1u_2; v_1v_2) \in P(G; R, S)$.

Suppose next that B contains a vertex of R . If both $C(v_1, v_2)$ and $C(v_2, v_1)$ contain a vertex of S , then, by Lemma 4.7, $c(u_1u_2; v_1v_2) \in P(G; R, S)$. Suppose $C(v_1, v_2)$ or $C(v_2, v_1)$ contains no vertex of S ; say $C(v_2, v_1)$ contains no vertex of S . Let G_1 be the subgraph of G consisting of all vertices and edges reachable by a path from $\{u_1\} \cup V(B)$ that has no vertex in $\{v_1, v_2\}$, and let $G_2 := G[(V(G) \setminus V(G_1)) \cup \{v_1, v_2\}]$. Since G_1 contains a $(uv_1; u_1v_2)$ - and a $(uv_2; u_1v_1)$ -linkage, (G_1, G_2) is a sided (R, S) -separation of G . The case where $C(v_1, v_2)$ contains no vertex of S is similar. \square

Lemma 4.9. *Let $(G, \mathcal{A}, t_1, \dots, t_n)$ be a nearly planar graph, where G is 2-connected and \mathcal{A} is minimal, and let $R, S \subseteq V(G)$ such that $\{t_1, \dots, t_n\} \subseteq R \cup S$. Suppose that t_1, \dots, t_n contains distinct vertices u_1, v_1, u_2, v_2 in this order, where $u_1, u_2 \in R$ and $v_1, v_2 \in S$. Let $(H, t_1, \dots, t_n) = p(G, \mathcal{A}, t_1, \dots, t_n)$ and let C be the cycle bounding the infinite face of a plane drawing of H with t_1, \dots, t_n on C . If $\{t_1, \dots, t_n\} \subset R \cup S$, then at least one of the following holds:*

- (i) $c(u_1u_2; v_1v_2) \in P(G; R, S)$,
- (ii) there exists a sided (R, S) -separation or a side (S, R) -separation, or
- (iii) there exists a cyclic ordering w_1, \dots, w_{n+1} with $\{t_1, \dots, t_n\} \subset \{w_1, \dots, w_{n+1}\} \subseteq R \cup S$ such that t_1, \dots, t_n occur in this ordering in w_1, \dots, w_{n+1} and (G, w_1, \dots, w_{n+1}) is nearly planar.

Proof. Suppose $\{t_1, \dots, t_n\} \subset R \cup S$. If no component B of $G - V(C)$ contains a vertex of $R \cup S$, then $R \cup S \subseteq V(C)$. We may assume that a component B of $G - V(C)$ contains a vertex u of R ; the case where a component B of $G - V(C)$ contains a vertex v in S is similar. If a neighbor of B belongs to $C(v_1, v_2)$ and a neighbor of B belongs to $C(v_2, v_1)$, then, by Lemma 4.5, $c(u_1u_2; v_1v_2) \in P(G; R, S)$.

We may therefore assume that $N(B) \subseteq C[v_2, v_1]$. Suppose $N(B) = \{v_1, v_2\}$. By Lemma 4.8, $c(u_1u_2; v_1v_2) \in P(G; R, S)$ or G has a sided (R, S) -separation or a sided (S, R) -separation.

We may therefore assume that a neighbor of B belongs to $C(v_2, v_1)$. By Lemma 4.6, $c(u_1u_2; v_1v_2) \in P(G; R, S)$, G has a sided (R, S) -separation or a sided (S, R) -separation, or there exists a cyclic ordering w_1, \dots, w_{n+1} with $\{t_1, \dots, t_n\} \subset \{w_1, \dots, w_{n+1}\} \subseteq R \cup S$ such that (G, w_1, \dots, w_{n+1}) is nearly planar and t_1, \dots, t_n occur in this ordering in w_1, \dots, w_{n+1} . \square

Lemma 4.10. *Let G be a 2-connected graph and let $R, S \subseteq V$, where $R \cup S$ has at least four vertices. Let $u_1, u_2, u_3 \in R \cap S$ be distinct vertices. Suppose $c(u_1, u_2, u_3) \notin P(G; R, S)$. Then there exist distinct vertices $u_4, u_5 \in R$ and distinct vertices $v_1, v_2 \in S$ such that $c(u_4u_5; v_1v_2) \notin P(G; R, S)$ and $c(u_1, u_2, u_3) + c(u_4u_5; v_1v_2) \in P(G; R, S)$.*

Proof. Since $R \cup S$ has at least four vertices, there exists a vertex $v \in R \cup S$ distinct from u_1, u_2, u_3 . By symmetry, we may assume that $v \in R$. Since G is 2-connected, there exist two vertex-disjoint paths P_1, Q_1 from $\{v, u_1\}$ to $\{u_2, u_3\}$ and two vertex-disjoint paths P_2, Q_2 from $\{v, u_2\}$ to $\{u_1, u_3\}$. Let v and $z_2 \in \{u_2, u_3\}$ be the ends of P_1 , and let z_1 and z_3 be the ends of Q_1 . Let v and z_1 be the ends of P_2 , and let z_2 and z_3 be the ends of Q_2 . Then $\{z_1, z_2, z_3\} = \{u_1, u_2, u_3\}$, and

$$c(vz_2; z_1z_3) + c(vz_1; z_3z_2) = c(z_1, z_2, z_3) + c(vz_3; z_1z_2).$$

Since either $c(z_1, z_2, z_3) = c(u_1, u_2, u_3)$ or $c(z_1, z_2, z_3) = -c(u_1, u_2, u_3)$, $c(z_1, z_2, z_3) \notin P(G; R, S)$. Since $c(vz_2; z_1z_3), c(vz_1; z_3z_2) \in P(G; R, S)$, we obtain that $c(vz_3; z_1z_2) \notin P(G; R, S)$. \square

In the proof of the Theorem 4.1, we use the following theorem; see [1].

Theorem 4.11. *Let $G = (V, E)$ be a graph and let s_1, s_2, t_1, t_2 be vertices of G . Then G contains no $(s_1t_1; s_2t_2)$ -linkage if and only if $(G; s_1, s_2, t_1, t_2)$ is nearly planar.*

Proof of Theorem 4.1. Since $L(R, S) - P(G; R, S) \neq \emptyset$, there exist vertices $u_1, u_2 \in R$ and vertices $v_1, v_2 \in S$ such that $c(u_1u_2; v_1v_2) \in L(R, S) - P(G; R, S)$, or there exist vertices $u_1, u_2, u_3 \in R \cap S$ such that $c(u_1, u_2, u_3) \in L(R, S) - P(G; R, S)$, by Lemma 2.1.

If $c(u_1, u_2, u_3) \in L(R, S) - P(G; R, S)$ and $|R \cup S| = 3$, then $(G; u_1, u_2, u_3)$ is nearly planar and R and S cross. If $c(u_1, u_2, u_3) \in L(R, S) - P(G; R, S)$ and $|R \cup S| > 3$, then there exist vertices $u_4, u_5 \in R$ and vertices $v_1, v_2 \in S$ such

that $c(u_4u_5; v_1v_2) \in L(R, S) - P(G; R, S)$, by Lemma 4.10. Hence we may assume that $c(u_1u_2; v_1v_2) \notin P(G; R, S)$. By Theorem 4.11, $(G; u_1, v_1, u_2, v_2)$ is nearly planar. By Lemma 4.9, either G has a sided (R, S) -separation or a side (S, R) -separation, or there exists a cyclic ordering w_1, \dots, w_n of the vertices in $R \cup S$ such that u_1, v_1, u_2, v_2 occur in this order in w_1, \dots, w_n and (G, w_1, \dots, w_n) is nearly planar. Also R and S cross. \square

5. PROOF OF THEOREM 1.1

Lemma 5.1. *Let $(G, \mathcal{A}, r_1, \dots, r_n)$ be a 2-connected nearly planar graph. If u_1, u_2, v_1, v_2 are distinct vertices occurring in this order in r_1, \dots, r_n , then there exists a $(u_1u_2; v_1v_2)$ -linkage in G .*

Lemma 5.2. *Let $(G, \mathcal{A}, t_1, \dots, t_n)$ be a 2-connected nearly planar graph, and let $R, S \subseteq \{t_1, \dots, t_n\}$. Suppose $u_1, v_1, u_2, v_2 \in R \cap S$. If u_1, v_1, u_2, v_2 are distinct vertices occurring in this order in t_1, \dots, t_n , then $c(u_1u_2; v_1v_2) + c(v_1v_2; u_1u_2) \in P(G; R, S)$.*

Proof. Since $(G, \mathcal{A}, t_1, \dots, t_n)$ is 2-connected, G contains a $(u_1v_1; u_2v_2)$ -linkage, a $(v_1u_2; v_2u_1)$ -linkage, a $(u_2v_2; u_1v_1)$ -linkage, and a $(v_2u_1; v_1u_2)$ -linkage. Hence $c(v_1u_1; v_2u_2) \in P(G; R, S)$, $c(u_2v_1; u_1v_2) \in P(G; R, S)$, $c(v_2u_2; v_1u_1) \in P(G; R, S)$, and $c(u_1v_2; u_2v_1) \in P(G; R, S)$. Then

$$c(u_1u_2; v_1v_2) + c(v_1v_2; u_1u_2) = c(v_1u_2; v_2u_1) - c(u_2v_1; u_1v_2) + c(v_2u_2; v_1u_1) - c(u_1v_2; u_2v_1),$$

and hence $c(u_1u_2; v_1v_2) + c(v_1v_2; u_1u_2) \in P(G; R, S)$. \square

Lemma 5.3. *Let $(G, \mathcal{A}, t_1, \dots, t_n)$ be a 2-connected nearly planar graph, and let $R, S \subseteq \{t_1, \dots, t_n\}$. Suppose that $u_1, u_2, \bar{u}_1, \bar{u}_2 \in R$ and $v_1, v_2, \bar{v}_1, \bar{v}_2 \in S$. If u_1, v_1, u_2, v_2 are distinct vertices, occurring in this order in t_1, \dots, t_n , and $\bar{u}_1, \bar{v}_1, \bar{u}_2, \bar{v}_2$ are distinct vertices, occurring in this order in t_1, \dots, t_n , then*

$$c(u_1u_2; v_1v_2) - c(\bar{u}_1\bar{u}_2; \bar{v}_1\bar{v}_2) \in P(G; R, S).$$

Proof. Suppose for a contradiction that the statement of the lemma is false. Choose the sequences u_1, v_1, u_2, v_2 and $\bar{u}_1, \bar{v}_1, \bar{u}_2, \bar{v}_2$ such that $c(u_1u_2; v_1v_2) - c(\bar{u}_1\bar{u}_2; \bar{v}_1\bar{v}_2) \notin P(G; R, S)$.

If $\{u_1, u_2\} = \{\bar{v}_1, \bar{v}_2\}$ and $\{v_1, v_2\} = \{\bar{u}_1, \bar{u}_2\}$, then, by Lemma 5.2, $c(u_1u_2; v_1v_2) - c(\bar{u}_1\bar{u}_2; \bar{v}_1\bar{v}_2) \in P(G; R, S)$. We may therefore assume that $\{u_1, u_2\} \neq \{\bar{v}_1, \bar{v}_2\}$ or $\{v_1, v_2\} \neq \{\bar{u}_1, \bar{u}_2\}$. By symmetry, we may assume that $u_1 \notin \{\bar{v}_1, \bar{v}_2\}$. Furthermore, by symmetry we may also assume that u_1 is a vertex in between \bar{v}_2 and \bar{v}_1 in the cyclic ordering t_1, \dots, t_n . Since $(G, \mathcal{A}, t_1, \dots, t_n)$ is 2-connected, G contains a $(u_1\bar{u}_1; \bar{v}_1\bar{v}_2)$ -linkage. Hence

$$c(u_1\bar{u}_1; \bar{v}_1\bar{v}_2) \in P(G; R, S),$$

and therefore

$$c(u_1u_2; v_1v_2) - c(\bar{u}_1\bar{u}_2; \bar{v}_1\bar{v}_2) - c(u_1\bar{u}_1; \bar{v}_1\bar{v}_2) \notin P(G; R, S).$$

Since

$$c(u_1u_2; v_1v_2) - c(u_1\bar{u}_2; \bar{v}_1\bar{v}_2) = c(u_1u_2; v_1v_2) - c(\bar{u}_1\bar{u}_2; \bar{v}_1\bar{v}_2) - c(u_1\bar{u}_1; \bar{v}_1\bar{v}_2),$$

we obtain

$$c(u_1u_2; v_1v_2) - c(u_1\bar{u}_2; \bar{v}_1\bar{v}_2) \notin P(G; R, S).$$

We may therefore assume that $u_1 = \bar{u}_1$. Since u_1, v_1, u_2, v_2 and $u_1, \bar{v}_1, \bar{u}_2, \bar{v}_2$ occur in the same ordering in t_1, \dots, t_n , either v_1 is between u_1 and \bar{u}_2 in the cyclic ordering t_1, \dots, t_n or v_2 is between \bar{u}_2 and u_1 in the cyclic ordering t_1, \dots, t_n . We may assume that v_1 is between u_1 and \bar{u}_2 in the cyclic ordering t_1, \dots, t_n . Since $(G, \mathcal{A}, t_1, \dots, t_n)$ is 2-connected, G contains a $(u_1\bar{u}_2; v_1\bar{v}_1)$ -linkage. Hence

$$c(u_1\bar{u}_2; v_1\bar{v}_1) \in P(G; R, S),$$

and therefore

$$c(u_1u_2; v_1v_2) - c(u_1\bar{u}_2; \bar{v}_1\bar{v}_2) - c(u_1\bar{u}_2; v_1\bar{v}_1) \notin P(G; R, S).$$

Since

$$c(u_1u_2; v_1v_2) - c(u_1\bar{u}_2; v_1\bar{v}_2) = c(u_1u_2; v_1v_2) - c(u_1\bar{u}_2; \bar{v}_1\bar{v}_2) - c(u_1\bar{u}_2; v_1\bar{v}_1),$$

we obtain

$$c(u_1 u_2; v_1 v_2) - c(u_1 \bar{u}_2; v_1 \bar{v}_2) \notin P(G; R, S).$$

We may therefore assume that $\bar{v}_1 = v_1$. In the same manner, $u_2 = \bar{u}_2$ and $v_2 = \bar{v}_2$. This is a contradiction. \square

Proof of Theorem 1.1. (1) \implies (2) Suppose $L(R, S) - P(G; R, S) \neq \emptyset$. Since G is (R, S) -connected, there exists a cyclic ordering t_1, \dots, t_n of the vertices of $R \cup S$ such that (G, t_1, \dots, t_n) is nearly planar and R and S cross, by Theorem 4.1.

(2) \implies (3) Suppose (G, t_1, \dots, t_n) is nearly planar and R and S cross. Suppose first that $R = S$ and $|R| = 3$. Then $P(G; R, S) = \{0\}$ and $L(R, S)$ is generated by $c(u_1, u_2, u_3)$, where u_1, u_2, u_3 are the vertices in R . Since $\text{link}(c(u_1, u_2, u_3); t_1, \dots, t_n) \in \{-2, 2\}$, we see that in this case, for all $z \in L(R, S)$, $\text{link}(z; t_1, \dots, t_n) = 0$ if and only if $z \in P(G; R, S)$. Suppose next that $|R \cup S| > 3$. Let $c \in L(R, S)$ with $\text{link}(c; t_1, \dots, t_n) = 0$. By Lemma 2.1, $c = \sum_{i=1}^m c_i$, where each c_i is an elementary (R, S) -tensor. Observe that $\text{link}(c; t_1, \dots, t_n) = \sum_{i=1}^m \text{link}(c_i; t_1, \dots, t_n)$ and, for each elementary (R, S) -tensor d , $\text{link}(d; t_1, \dots, t_n) \in \{-2, 0, 2\}$. If $\text{link}(c_j; t_1, \dots, t_n) \neq 0$ for some $j \in \{1, \dots, m\}$, then there exists an c_k such that $\text{link}(c_k; t_1, \dots, t_n) = -\text{link}(c_j; t_1, \dots, t_n)$. Then $\text{link}(c_j - c_k; t_1, \dots, t_n) = 0$. By Lemmas 5.3 and 4.10, $c_j - c_k \in P(G; R, S)$. Hence $c \in P(G; R, S)$.

Conversely, if $c \in P(G; R, S)$, then $c = \sum_i c_i$, where each c_i is of the form $c(u_1 u_2; v_1 v_2)$ with u_1, u_2, v_1, v_2 distinct, and $u_1, u_2 \in R$ and $v_1, v_2 \in S$, such that there exists an $(u_1 u_2; v_1 v_2)$ -linkage in G . Since $\text{link}(c(u_1 u_2; v_1 v_2); t_1, \dots, t_n) = 0$ if there exists an $(u_1 u_2; v_1 v_2)$ -linkage in G , $\text{link}(c; t_1, \dots, t_n) = 0$.

(3) \implies (1) Suppose that there exists a cyclic ordering t_1, \dots, t_n of the vertices in $R \cup S$ such that R and S cross and, for all $z \in L(R, S)$, $\text{link}(z; t_1, \dots, t_n) = 0$ if and only if $z \in P(G; R, S)$. Since R and S cross, either $R = S$ and $|R| = 3$, or there exist vertices $u_1, u_2 \in R$ and $v_1, v_2 \in S$ such that u_1, v_1, u_2, v_2 are distinct and occur in this order in the cyclic ordering t_1, \dots, t_n . If $R = S$ and $|R| = 3$, let u_1, u_2, u_3 be the vertices of R . Then $c(u_1, u_2, u_3) \notin P(G; R, S)$. Suppose next that there exist vertices $u_1, u_2 \in R$ and $v_1, v_2 \in S$ such that u_1, v_1, u_2, v_2 are distinct, occurring in this order in the cyclic ordering t_1, \dots, t_n . Since $\text{link}(c(u_1 u_2; v_1 v_2); t_1, \dots, t_n) \neq 0$, $c(u_1 u_2; v_1 v_2) \notin P(G; R, S)$. \square

REFERENCES

- [1] P. D. Seymour. Disjoint paths in graphs. *Discrete Math.*, 29(3):293–309, 1980.
- [2] Hein van der Holst. A polynomial-time algorithm to find a linkless embedding of a graph. *J. Combin. Theory Ser. B*, 99(2):512–530, 2009.

DEPARTMENT OF MATHEMATICS AND STATISTICS, GEORGIA STATE UNIVERSITY, ATLANTA, GEORGIA , USA

DEPARTMENT OF MATHEMATICS AND STATISTICS, GEORGIA STATE UNIVERSITY, ATLANTA, GEORGIA , USA

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCGILL UNIVERSITY, MONTREAL, QC, H3A 2K6, CANADA

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GEORGIA 30332-0160, USA